Main Manuscript for

Locomotor savings relies on attentional control of walking in older, but not younger adults.

Shuqi Liu[1,2], Andrea L. Rosso[3], Emma M. Baillargeon[4], Andrea M Weinstein[5], Gelsy Torres-Oviedo[1,2]

[1]Dept. of Bioengineering; [2]Center for the Neural Basis of Cognition; [3]Dept. of Epidemiology; [4]Dept. of Physical Therapy & Human Movement Sciences, Northwestern University, Chicago, IL, USA; [5]Dept. of Psychiatry, University of Pittsburgh, Pittsburgh, PA, USA.

Corresponding Author: Dr. Gelsy Torres-Oviedo

Corresponding Author's Email: (gelsyto@pitt.edu)

Author Contributions: S.L., A.R., A.W., and G.T. conceptualized and designed research; A.R., E.B., A.W., and G.T. advised on data analysis and interpretation and reviewed and edited the manuscript; G.T. acquired funding; S.L. performed research, analyzed data, and wrote the paper.

Competing Interest Statement: The authors declare no competing interest.

Keywords: multiday savings, functional near-infrared spectroscopy, dual-task walking, split-belt treadmill, aging

Preprint server: BioRxiv

Abstract

The ability to recall learned movements and rapidly adapt to environmental changes, known as locomotor savings, is crucial for mobility in community-dwelling older adults. However, the influence of aging on locomotor savings and the underlying mechanisms remains poorly understood. Attentional compensation is a particularly relevant mechanism because the control of automatic motor behaviors like walking tend to recruit more attentional/executive resources with aging. We hypothesize that locomotor savings is diminished with age and relies on attentional rather than automatic control of walking. To test this, we compared savings of a novel walking pattern learned on a split-belt treadmill, where each leg moves at a different speed, across multiple days in 21 older and 21 younger adults. Attentional control of walking was assessed by overground dual-task walking while prefrontal cortex (PFC) activity was recorded using functional near-infrared spectroscopy (fNIRS). We found that older adults exhibited less locomotor savings than younger adults after practice. Older adults also relied more on attentional resources during dual-task walking. Importantly, greater locomotor savings was associated with higher attentional control of walking in older adults, suggesting that the use of attentional resources during challenging walking facilitates the recall of previously learned movements. These results indicate that cognitive compensation strategies utilizing attentional resources are important neural mechanisms modulating locomotor savings. Understanding the role of cognitive compensation in locomotor savings may inform rehabilitation design to enhance mobility in older adults ensuring movement corrections practiced in clinical settings are saved for long-term benefit in daily life.

Introduction

Mobility in the community occurs within a dynamic environment and often requires adapting walking patterns according to the situations at hand and to recall previously adapted movements (1, 2), known as locomotor savings (3–6). For example, navigating icy surfaces initially requires reactive motor commands; however, subsequent encounters allow for the anticipatory retrieval of learned patterns, ensuring stability. Locomotor savings over multiple days is especially important in community mobility, where individuals must repeatedly draw on previously learned walking patterns to handle recurring challenges (1, 2, 7, 8), such as crossing busy intersections or navigating construction zones.

Inability to recall previously learned walking patterns when needed, i.e. poor locomotor savings, could contribute to restrictions in community-mobility, the ability to move outside one's home (9), which are common (10, 11) in older ages. However, the effect of aging on multi-day locomotor savings remains unclear. Prior studies (12–14) suggested that older adults exhibited impaired locomotor savings when tested across days, but these findings are inconclusive due to uncontrolled intervals between testing days and the absence of practice to induce robust locomotor savings. In this study, we systematically test whether older adults can demonstrate locomotor savings across multiple days following a structured practice and quantify the extent of locomotor savings relative to younger adults.

Furthermore, we are interested in elucidating the neural mechanisms of locomotor savings, particularly the role of attentional compensation. Attention will be defined as the information processing capacity of the prefrontal cortex (PFC) here (15). With older age, walking control shifts from predominantly automatic to compensatory strategies, relying more heavily on attentional and executive resources (16). This shift in control is often assessed by dual-task walking (walking while performing an attention demanding task) with recording of prefrontal cortex (PFC) activity. In this context, decreased task performance and/or increased PFC activity indicate that walking is competing with the secondary task for attentional resources, which is interpreted as reduced automaticity and increased attentional control of walking (17–21). Consistently, reaching motor savings in younger adults was related to brain activation in the cortico-striatal loops (22–24) but in older adults was linked to more frontal brain activities (25). Locomotor savings in younger adults

54 are typically linked to subcortical circuits, such as the cerebellar-thalamic-cortical and cerebellar-
55 basal ganglia circuits (26), and these structures often undergo age-related atrophy (27, 28).
56 Therefore, we hypothesize attentional compensation during walking will be relevant to locomotor
57 savings in older adults.
58
In summary, this study investigates age-related differences in locomotor savings across multiple
days and evaluates the contribution of attentional compensation. We hypothesize that with aging,
locomotor savings is diminished and relies on attentional rather than automatic control of walking.
To test this, we assessed locomotor savings and attentional control of walking in older and
younger adults. Locomotor savings was quantified by retrieval of a novel walking pattern acquired
through repeated exposure to a split-belt treadmill, where each leg moves at a different speed.
Attentional control of walking was evaluated using a well-established overground dual-task
walking paradigm (20, 29), with neural compensation from the PFC measured by functional near-
infrared spectroscopy (fNIRS). Finally, to explore the clinical relevance of these findings in older
adults, we tested the relationship between locomotor savings, dynamic balance and mobility as
measured by the Functional Gait Assessment (FGA) (30), and cognitive abilities evaluated by
neuropsychological assessment in older adults.
71
72
73 **Results**
74
We recruited 21 older (>= 65 years old) and 21 younger (19-40 years old) adult participants in this
study. Demographics and self-selected overground walking speeds are shown in Table 1. Leg
dominance was determined by self-reported leg used to kick a soccer ball. Self-selected
overground walking speed was calculated as the average speed walking back and forth on a
straight 9.2m walkway for 150 strides (Figure 1A baseline overground). Older adults were
screened for no signs of cognitive impairments as indicated by a Montreal Cognitive Assessment
(MoCA) score greater than or equal to 24. The full inclusion criteria for both age groups are
included in the Methods. The Functional Gait Assessment (FGA) was administered to older adults
to evaluate dynamic balance.
84
Older adults showed less locomotor savings than younger adults
Locomotor savings was evaluated by the change in step length asymmetry (StepAsym, Fig 1B)
before and after completing a split-belt treadmill walking practice session (Figure 1A). StepAsym
is defined as the normalized difference in bilateral step lengths (see methods, Equation 1), which
is a robust and clinically relevant measure conventionally used to characterize gait changes (31).
During the practice session, participants experienced multiple bouts of alternating split (legs
moving at different speeds) and tied belt (regular treadmill) walking (Figure 1A), which has been
shown to accelerate locomotor savings in young adults (2, 32).
93
Figure 2A illustrates the time course of StepAsym during adaptation (split-belt) relative to baseline
(tied-belt) in the pre- and post-practice sessions. Locomotor savings was observed by reduced
gait perturbation (closer to zero StepAsym) in the post-practice (open circles) compared to the
pre-practice session (filled circles, Figure 2A). We quantified locomotor savings with a savings
index, defined as the change in StepAsym during early split-belt walking from pre- to post-
practice as a proportion of the pre-practice StepAsym (see methods, Equation 2). We found a
significantly smaller locomotor savings index in older adults than younger adults (two-sample t-
test, $t = -3.96$, $p < 0.001$, Cohen's $d = -1.22$; Figure 2B), suggesting that locomotor savings is
reduced with age.
103
Despite differences in locomotor savings, older and younger adults adapted to similar steady
state walking behavior in both sessions (Figure 2C). StepAsym during the pre- and post-practice
steady state periods (last 40 strides of adaptation) showed no significant main effect of age group
($F_{(1,40)} = 0.22$, $p = 0.64$), session ($F_{(1,40)} = 0.13$, $p = 0.72$), nor their interactions ($F_{(1,40)} = 0.05$, $p =$

0.82). These results suggest that older adults maintained the ability to adapt walking patterns with
sustained exposures to a novel environment, but they saved less across days.
110
Locomotor savings was correlated with higher attentional control of gait in older adults
To determine the role of attentional control of walking in locomotor savings, we evaluated the
attentional control of gait using a well-established dual-task walking paradigm (20, 29).
Participants walked at their comfortable pace around an oval while reciting every two (easy dual-
task) or three letters of the alphabet (hard dual-task), with concurrent prefrontal cortical (PFC)
activity recordings from a functional near-infrared spectroscopy (fNIRS) headband (Figure 1C).
Similar results were observed in both tasks, and the following section will focus on the hard dual-
task (see supplementary Fig. S1, Table S2, S3, for the easy dual-task). Attentional control of
walking was quantified with an attentional gait index (33), which combines the PFC activation and
motor and cognitive performance change from single- to dual-task walking (Equation 3 in
methods). Higher values on the attentional gait index reflect greater reliance on attentional
resources and reduced automatic control of gait (16, 18–21, 33). This indirect assessment was
done as a first step to evaluate attentional control during walking due to technical challenges in
acquiring reliable fNIRS-based PFC estimations during a dynamic adaptation task likes locomotor
savings (see limitations).
126
We found that older adults have greater attentional control during walking than younger adults.
This is demonstrated by the significantly higher attentional gait index in older than younger adults
(z = 3.20, p = 0.001; Figure 3A). In the exploratory analysis, we found that this age difference in
attentional gait index was driven by elevated PFC activation (z = 3.52, p < 0.001; supplementary
Table S1), and a trending effect from greater motor performance cost (i.e., more reduction in
motor performance from single- to dual-task; trending interaction, $F_{(1,40)}$ = 3.13, p =0.08;
supplementary Table S1) in older than younger adults.
134
More importantly, locomotor savings was positively correlated with attentional gait index in older
($\rho = 0.62, p = 0.003$; Figure 3B), but not in younger adults ($\rho = -0.05, p = 0.82$; Figure 3B). The
association in older adults remained robust after controlling for age, sex, and self-selected
overground walking speed (partial correlation $\rho = 0.69, p = 0.001$). To identify the factors driving
the correlation between attentional gait index and locomotor savings in older adults, we
performed exploratory analyses on the subcomponents of attentional gait index (PFC activity and
performance), which are metrics that are conventionally reported in the literature (18–21, 34). We
found that locomotor savings was correlated with motor performance cost ($\rho = -0.45$, p = 0.04),
but not with cognitive performance cost ($\rho = -0.20, p = 0.39$) or PFC activity ($\rho = 0.28, p = 0.22$).
Notably, the association between locomotor savings and motor performance cost was not
influenced by single-task motor performance, as single-task motor performance was neither
associated with locomotor savings ($\rho = 0.04, p = 0.87$) nor motor performance cost ($\rho <$
$0.001, p = 1.00$). The results indicate that attentional control of walking, as measured by
attentional gait index, contributes to locomotor savings in older adults, and this relationship was
mainly driven by motor performance cost.
150
151 Taken together, we found that older adults rely more on attentional control during walking than
152 younger adults, and those with greater attentional control during walking exhibit greater locomotor
153 savings. This relationship was not present in younger adults, suggesting an age-specific
154 compensatory mechanism.
155
Locomotor savings was weakly correlated with functional gait assessment
To explore if locomotor savings is relevant for functional mobility, we tested its relationship with
the functional gait assessment (FGA), which evaluates an individual's ability to perform multiple
motor tasks while walking (35). We found a trending correlation where higher FGA score was
associated with more locomotor savings ($\rho = 0.38, p = 0.09$, supplementary Figure S2). Of note,
18 out of 21 older adults in our sample scored above the fall-risk cutoff of 22/30 on the FGA (35–

37), suggesting we have a high functioning cohort (mean = 24.71±2.43, range = [18, 29]). Taken
together, our results suggest that locomotor savings may only weakly reflect functional mobility in
older adults without significant mobility or balance limitations.
165
Locomotor savings in older adults was not correlated with cognitive abilities
Lastly, prior literature suggests cognitive processes are relevant for motor adaptation and savings
in both reaching (6, 38) and walking (4). Therefore, we explored the relationship between
locomotor savings and cognitive function using a comprehensive neuropsychological battery that
assesses general mental status and major domains of cognitive functioning, including language,
visuospatial ability, attention, memory, and executive functions (39–41). Among the domains
tested, we observed only a trending relationship between the Repeatable Battery for the
Neuropsychological Status (RBANS) (42) visuospatial index and locomotor savings where older
adults with higher scores exhibited greater locomotor savings ($\rho = 0.40, p = 0.07$). Given the
exploratory nature of these analyses and the sample size of our data these results should be
interpreted with caution. Full neuropsychological test scores and correlation results are provided
in supplementary Table S4.
178
179
180 **Discussion**
181
We found that older adults can exhibit multi-day locomotor savings following practice, but to a
less extent than younger adults. We also demonstrated that older adults have increased
attentional control of walking, and higher attentional control of walking is linked to greater
locomotor savings in older adults. We argue that with aging, different aspects of motor behaviors
(such as locomotor savings and dual-task walking) become linked due to increased reliance on
attentional compensations following age-related decline in the shared neural substrates. Future
work is needed to test how this relationship changes with different cognitive abilities such as in
older adults with cognitive impairment.
190
191 **Older adults can save and recall learned locomotor patterns with practice**
192 In this study, we demonstrated that older adults exhibited locomotor savings to a lesser extent
193 than younger adults. Mixed findings have been reported about the influence of aging on savings
194 in reaching (25, 43, 44) and walking (13, 14, 45–47) where older adults show either similar or
195 reduced savings depends on the task, protocol, and assessment intervals.
196
The main factor that could explain these discrepancies is our use of a structured practice session,
which is known to favor locomotor savings (2, 4, 32, 48). We used a practice protocol that
included abbreviated bouts of novel walking interleaved with regular walking on the practice day
(Figure 1A). These transitions provided repeated experiences of movement errors, a critical driver
for savings in younger adults (5, 48, 49). We extend previous research by demonstrating that
multiple movement errors also effectively induce locomotor savings in older adults. Unlike prior
multi-day studies (12, 14, 25) our structured practice likely enhanced memory consolidation,
allowing older adults to show locomotor savings.
205
The second factor distinguishing our study from previous is the controlled passage of time
between learning and re-learning (13, 50, 51). Previous work has reported that older adults forget
a newly learned walking pattern more and showed less retention than younger adults (13, 51, 52),
suggesting faster decays of recent motor memories in older than younger adults. In models
comprising fast and slow learning components, savings is typically attributed to the slow
component (3, 53, 54), and older adults often show reduced retention of the slow component
(55). Taken together, while older adults may demonstrate savings within a single session where
memory decay is incomplete (43–46), over longer intervals, such as the six-week in the previous
work, older adults' memory of the locomotor patterns may decay more until no savings remain
(12, 13). In our study, intervals between sessions were carefully controlled, with an average

interval of 3.98 days between initial exposure (pre-practice session) and the locomotor savings
assessment (post-practice session). The controlled time interval and the inclusion of a practice
session may have allowed us to detect the age-related difference in locomotor savings across
days.
220
221　In sum, we demonstrate that both older and younger adults can exhibit multi-day locomotor
222　savings after practice and controlled short time interval. Nonetheless, even under these controlled
223　conditions, older adults showed less locomotor savings than younger adults, suggesting age-
224　related limitations in long-term motor memory savings.
225
Older adults retain the flexibility to adapt in a novel environment
Despite differences in locomotor savings, older adults reached the same steady state behavior as
younger adults at the end of the adaptation. This indicates that although older adults were more
perturbed during early adaptation, they were able to adapt and achieve similar gait symmetry as
younger adults after a sustained exposure to the split-belt treadmill environment. This is
consistent with prior studies in walking (45, 46, 56–58) and reaching (43).
232
The hierarchical computational model of locomotor adaptation and savings (59) provides a
framework for interpreting these findings. The model comprises three key components: (1) a
reinforcement learner that improves performance, (2) a feedback controller that stabilizes the
body, and (3) a memory module that stores previously learned walking strategies. The age-
related decline in locomotor savings may emerge from deficits in one or more of these
components.
239
Dopaminergic reinforcement learning system is relevant for savings in reaching (48, 60, 61) and
dopaminergic neurotransmission has been associated with walking adaptation in aging (62, 63).
Although older adults eventually achieved similar asymmetry to younger adults, they took longer
to counteract the initial perturbation (13, 46, 56, 58, 64). Deficits in the dopaminergic
reinforcement learner can contribute to the slower learning within the session and reduced
savings across sessions.
246
Alternatively, a weakened feedback controller could contribute to poor movement adjustments
when encountering a novel environment, resulting in more perturbed movement during early split-
walking in older adults across all sessions (Figure 2A, blue circles for older adults were lower
than red circles for younger adults in both sessions). This aligns with balance (65–67) and
walking (64) literature showing reduced reactive feedback correction in older adults. A less
efficient feedback controller would result in larger errors during both learning and relearning
(savings).
254
Lastly, our results could indicate a faulty memory module in older adults that limit the update and
retrieval of the corresponding motor memory when re-encountering the split environment (59),
resulting in poor locomotor savings (Figure 2B). This is consistent with the fast and slow learning
components model, which attributed savings to a biased slow learning component during re-
learning (3, 53, 54). Older adults exhibited reduced retention in this slow component (51, 55),
which could result in a smaller bias during re-learning and therefore diminished locomotor savings
observed in this study.
262
263　Note that these hypotheses are not mutually exclusive. Future computational work is needed to
264　distinguish the specific components underlying the age-related differences in locomotor savings.
265
Attentional compensations can be a strategy for locomotor savings
We observed that higher attentional control of walking is correlated with greater locomotor
savings in older adults. This suggests that cognitive compensation, typically indicative of reduced
automaticity (16, 18–21), can be advantageous for complex walking behaviors like locomotor

savings (4, 68). This relationship may stem from shared neural correlates such as the subcortical-
frontal (e.g., basal ganglia and cerebellum to primary motor cortex) networks (45, 69, 70). These
structures are susceptible to age-related changes in white matter hyperintensity volume and
functional connectivity (71–73). Consequently, older adults may increasingly recruit PFC-
mediated attentional and executive resources as a compensatory mechanism for challenging
walking (16). Supporting this view, executive cortical-striatal network resting-state connectivity
has been linked to mild Parkinsonian signs (72, 74). In healthy aging, older adults with higher
attentional control of gait in dual-task walking may be more inclined to use cognitive strategies in
other challenging motor tasks such as locomotor adaptation, and this strategy may facilitate
locomotor savings.
280
We were unable to measure PFC activity concurrently with locomotor savings due to technical
limitations and we inferred the role of attentional control in locomotor savings using attentional
control of walking measured by behavior and PFC activation in dual-task walking. Future multi-
modal imaging studies are necessary to provide direct evidence for the neural correlates of
locomotor savings. Furthermore, the relationship between locomotor savings and attentional
control of walking was most robustly observed in the attentional gait index, highlighting the
importance to integrate performance when interpreting fNIRS-based PFC activities, as shown in
our previous work (33).
289
Attentional gait index revealed reduced automaticity and inefficient use of attentional
resources during challenging walking in older adults
We found that older adults exhibited increased attentional control of walking, as indicated by
higher attentional gait index and higher PFC activation compared to younger adults across tasks.
These findings align with prior work showing reduced gait automaticity in aging (16, 21) where
older adults showed elevated PFC activation or greater dual-task performance cost (16, 18–21).
Notably, our older adult cohort recruited more PFC yet performed worse than younger adults,
suggesting the extra PFC recruitment was inefficient. Taken together, our results indicate that
older adults had reduced gait automaticity, increased reliance on attentional resources, but
limited benefit from this compensatory recruitment during dual-task walking.
300
Other metrics of functional mobility are needed to determine the relationship between
mobility and locomotor savings
Our exploratory analysis revealed only a weak correlation between locomotor savings and
dynamic balance measured by the Functional Gait Assessment (FGA). However, this does not
mean locomotor savings is irrelevant for mobility. The lack of association may be due to a ceiling
effect in our cohort's FGA scores. Prior research has suggested that higher levels of self-reported
recent physical activity was related to better locomotor adaptations (57), and gait adaptability is
similar between physical activity matched older and younger adults (75). This warrants future
research to use more comprehensive mobility metrics, such as accelerometry or GPS-based
objective measures of real-world physical activities and community mobility (76) and examine
their associations with locomotor savings.
312
Cognitive assessment results were not directly associated with locomotor savings
Most neuropsychological test results in our battery were not correlated with locomotor savings.
While cognitive processes are relevant for savings in younger adults, such as explicit re-aiming
(6), explicit recall of perturbation size (4), and visual field dependence (77), evidence in older
populations remains sparse. One study showed no relation between adaptation rate and a variety
of neuropsychological test results in older adults (78), while Sasikumar and collegues (79)
reported that better overall cognitive scores were related to adaptation and aftereffects in patients
with Parkinson's disease. Taken together, these findings suggest that neuropsychological test
scores may only predict locomotor adaptation and savings in more impaired populations (e.g.,
Parkinson's disease patients), but not in older adults with little mobility or cognitive limitations like

in the current sample. Alternatively, the relationship between cognitive processes and locomotor
savings may only emerge under conditions that pose sufficient challenge to the nervous system
in both motor and cognitive domains simultaneously, such as in dual-task walking, rather than
through static neuropsychological assessments.
327
Limitations
The main limitation of this study is that attentional compensation was not measured concurrently
with locomotor savings. Reliable fNIRS-based PFC estimations require multiple repetitions of a
task to ensure a sufficient signal-to-noise ratio. However, since repeated exposures to split-belt
perturbations inherently alter adaptation behavior, each encounter represents a dynamic, non-
stationary state rather than a simple repetition. To address this challenge, future studies could
develop new fNIRS analysis supporting continuous measurement of a dynamic condition, use
indirect measures of attention (e.g., arousal), or incorporate imaging modalities with better
temporal resolution (80), such as electroencephalogram (EEG).
337
Clinical Implications
The relationship we identified between locomotor savings and attentional control of walking in
older adults is of clinical relevance because it reflects cognitive compensation strategies in
walking, which is a strong indicator of dementia risks (81). It is important to investigate how this
relationship changes when cognitive compensation is compromised due to neurodegeneration,
such as in older adults with cognitive impairment. Understanding these mechanisms will also help
identify older adults who are less likely to retain clinical gait interventions due to underlying
cognitive decline.
346
Conclusions
In conclusion, we found that practice effectively induced multi-day locomotor saving in older
adults, though to a lesser extent than in younger adults. We also found a moderate relationship
between attentional control of walking and locomotor savings in older adults, but not in younger
adults. This suggests that locomotor savings relies more on higher-order processes such as
attention in older ages, and the attentional compensation might support older adults to save and
recall from prior experiences. Future work is needed to directly examine the neural substrates
underlying this relationship and determine if the relationship persists when compensatory neural
resources are diminished, such as in older adults with cognitive impairment.
356
357
Materials and Methods
359
Participants
Older (>= 65 years old) and younger (19-40 years old) adult participants were recruited through
flyers posted in the community and through research registries including the Pitt+Me Research
Recruitment Program and the Pittsburgh Pepper Community Research Connection. All
participants could walk unassisted, did not take medications that change cognitive functions, and
had no major neurological conditions, severe visual impairment, or unmanaged orthopedic
conditions that would interfere with walking. Older adults were further screened for signs of
cognitive impairments as indicated by a MoCA score less than 24. A total of 25 older adult
participants were recruited; three did not meet the MoCA inclusion criteria and one dropped out,
resulting in 21 older adult participants for all data analysis. Sex matched younger adults were
recruited with one drop out, resulting in 21 participants for data analysis. Participant
demographics are included in Table 1. The Institutional Review Board at the University of
Pittsburgh approved the study. All participants gave written informed consent.
373
General Experimental Protocol
All participants performed dual-task walking while wearing a functional near-infrared spectroscopy
(fNIRS) headband to evaluate attentional control of walking (Fig. 1C) before exposure to the split-

belt treadmill. Then, participants completed three sessions of split-belt treadmill walking on
separate days, pre-practice, practice, and post-practice (Fig. 1A). Each session was one or two
days apart (between pre-practice and practice sessions: 1.67 ± 2.34 days (mean $\pm$ standard
deviation), between practice and post-practice: 1.31 ± 0.56 days). The evaluation for attentional
control of walking was completed on the same day before the first split-belt exposure for all
younger adults, resulting in three total experimental sessions in younger adults. Older adults
visited the lab one more time one- or two-days prior to the first split-belt treadmill walking session,
during which they completed a comprehensive neuropsychological exam, attentional control of
walking assessment, and Functional Gait Assessment (FGA). This extra visit was added to avoid
overburdening the participants at any single session. Importantly, in both groups, attentional
control of walking was always evaluated before any treadmill walking to characterize participants'
baseline abilities.

Protocol to Evaluate Locomotor Savings
Locomotor savings of a newly learned walking pattern was evaluated across multiple exposures
to a split-belt treadmill where the two legs moved at different speeds. The protocol consists of
three sessions (Fig. 1A): pre-practice, practice and post-practice. All sessions consist of baseline,
adaptation, and washout conditions. Detailed treadmill speed profile for the whole protocol can be
found in supplementary material Figure S3.

Baseline was collected during both overground and treadmill walking. During overground
baseline, participants walked back and forth on a 9.2m walkway at a self-selected speed for 150
strides excluding turns. A stride was defined as two consecutive heel-strikes of the same leg and
heel strikes were identified in the kinematic data as the times of the ankle marker's greatest
forward excursion (13, 82, 83). During treadmill baseline, participants walked on an instrumented
treadmill (Bertec, Columbus, OH, United States) with both legs moving at the same speed (tied-
belt) at 0.5m/s, 1.0m/s, and 0.75 m/s for 150 strides each. Then, during adaptation, participants
walked on the treadmill with the dominant leg moving at 1m/s and non-dominant leg moving at
0.5m/s (split-belt). Participants experienced this novel situation in a single block of 900 strides in
split-belt in "pre" and "post-practice" sessions (Fig. 1A). In contrast, during the "practice" session,
participants experienced five bouts of 250 strides with the treadmill belts split interleaved with four
bouts of 50 strides with the treadmill belts tied at 0.75m/s (Fig. 1A). Next, during washout,
participants walked 450 strides at their self-selected speed overground and then 150 strides with
tied-belt treadmill at 0.75m/s. The overground and tied-belt walking were included to wash out the
adaptation effect from the session.

After treadmill washout condition, participants experienced two short perturbations in opposite
directions (supplementary material Figure S3). Overground washout and short perturbations were
performed for a different aim in the larger study to evaluate adaptation in muscle activities (84,
85) and were not used in the analyses presented in this paper. We included the information here
to promote reproducibility. However, we believe these protocol design choices do not impact our
overall conclusions.

Participants took sit-down resting breaks during regular intervals to limit fatigue (dashed lines in
Fig. 1C). The breaks were administered at the same location with the same durations for all
participants (break duration for older adults: 5.35 ± 0.58 mins (mean $\pm$ standard deviation), younger
adults: 5.16 ± 0.41 mins). Participants were instructed to not move their feet during the breaks.

Locomotor Savings Data Collection
Kinematic data during split-belt walking was collected with Vicon Motion Systems (Oxford, United
Kingdom) at 100Hz. We placed passive reflective markers bilaterally over the hip (greater
trochanter) and ankle (lateral malleolus). Gaps in the raw kinematic data due to marker
occlusions were filled first with a quintic spline interpolation (Woltring; Vicon Nexus Software,

Oxford, United Kingdom) and remaining gaps were filled manually using Vicon's spline, pattern,
and cyclic fill tools when appropriate as done in previous studies (12).
432
Locomotor Savings Outcome Measures
Behaviors on the treadmill were quantified by step length asymmetry (StepAsym, Figure 1B),
defined as the normalized difference in bilateral step lengths (Equation 1). The normalization
allows fair comparison among individuals with different stride length. A zero value for StepAsym
indicated that both steps were the same length, i.e., symmetric walking. Step length is quantified
by the distance between the ankle markers at heel strike, specifically fast step length was
measured during fast leg heel strike and vice versa for the slow leg.
440

$$StepAsym = \frac{StepLength_{fast} - StepLength_{slow}}{FStepLength_{fast} + StepLength_{slow}} \ (Equation\ 1)$$

441
Locomotor savings was quantified by a locomotor savings index, defined as the difference in
StepAsym during early adaptation (first 30 strides in adaptation) between the pre- and post-
practice sessions, normalized by pre-practice (Equation 2). This normalization allowed us to
quantify locomotor savings relative to individual's baseline ability to counteract the split-belt
perturbations. The average of the first 30 strides was chosen as a model-free approximation for
the adaptation rate without assuming any specific patterns of the stride-dependent relationship (2,
26). Higher values on the locomotor savings index indicate less perturbed movements when re-
exposed to split-belt walking during the post-practice compared to pre-practice session, which
means more locomotor savings.
451

$$Locomotor\ Savings\ Index = \frac{StepAsym_{Post} - StepAsym_{pre}}{|StepAsym_{pre}|} \ (Equation\ 2)$$

452
453 StepAsym during pre-practice was always negative for all participants. So, absolute value was
454 taken on the denominator such that the sign of the locomotor savings index is determined by the
455 differences between post and pre-practice StepAsym and a positive locomotor savings index
456 represents improved StepAsym (closer to zero gait) in post- relative to pre-practice.
457
Protocol to Evaluate Attentional Control of Walking
Attentional control of walking was evaluated using a well-established dual-task walking paradigm
(20, 29). Participants walked at their comfortable pace along on oval path with a ~5.85-meter
straight walkway on each side, while reciting every two (e.g., A, C, E, etc.) or three letters (e.g., A,
D, G, etc.) of the alphabet and wearing a functional near-infrared spectroscopy (fNIRS) headband
to measure prefrontal cortical (PFC) activity (Fig. 1C). Reciting every two or three letters while
walking was considered the easy or hard dual-task conditions, respectively. Two levels of
difficulties were included to allow calculation of an attentional gait index as shown in our previous
work (33). Participants also performed a motor single-task, walking without any secondary task,
and two cognitive single-tasks, standing while reciting every two (easy) or three (hard) letters of
the alphabet (Fig. 1C). The single-task conditions allowed the quantification of performance
change from single- to dual-task walking.
470
We used a block design with six trials. Each trial includes all three single-tasks and two dual-
tasks in a pseudo random order. The task order within a trial was kept the same across all
participants, but the order of the trials was randomized. Each task was performed for 20 seconds,
and a standing rest preceded every task. During the standing rest, participants were instructed to
stand while counting silently from one to allow the hemodynamic response to return to rest level
while limiting distraction or practice of the cognitive task during rest (86).
477
Instructions for each condition were given by a computerized voice using a customized MATLAB
script that automatically controls task timing. Before starting the session, we confirmed that all

participants could hear the instructions clearly. Participants were instructed to walk at their
comfortable pace while doing the cognitive task with no specific instruction about which task to
prioritize. A practice trial was performed up to three times to ensure the participant understood
the task instructions (repeated 1.02±0.15 (mean±standard deviation) times).
484
Attentional Control of Walking Data Collection and Data Processing
Motor performance was quantified by gait speed (speed, m/s) using motion data from Vicon
Motion Systems (Oxford, United Kingdom). Motor performance during dual-task was calculated
as the total Euclidian distance in the x-y (transverse) plane traveled by the average position of the
hip markers (greater trochanter) divided by the time the marker was moving. The hip markers'
motion data was collected using the same system and processed with the same pipeline as the
locomotor savings data (see *Locomotor Savings Data Collection* section above).
492
493 Cognitive performance was measured by rate of correct letters of alphabet generated (rate, letter
494 /s). An experimenter transcribed the alphabet generated by the participant during the
495 experimental session and the rate of correct letters of alphabet generated was calculated by a
496 customized MATLAB script during post-processing.
497
PFC activity was measured from an eight-channel continuous wave functional near-infrared
spectroscopy (fNIRS) headband (Octamon, Artinis Medical Systems, Netherlands). The fNIRS
headband consisted of two detectors and eight sources with a source and detector pair distance
of 35mm. The fNIRS headband covered both the left and right PFC regions, specifically
Brodmann areas 9, 44, 45, and 46 (34). The center of the fNIRS headband was aligned with the
center of the participant's nose, and the bottom of the headband was just above the eyebrow
(87). The headband was not removed between trials to maintain stable placement.
505
fNIRS quantifies changes in the blood oxygenation based on distinct light absorption properties of
the oxygenated (Hbo) and deoxygenated (Hbr) blood, which provides a proxy for cortical
activities. Near-infrared light transmitted at 850 nm and 760 nm was used to detect changes in
Hbo and Hbr. Data were sampled at 10 Hz and collected by the OxySoft software (Artinis
Medical Systems, Netherlands). No short separation channel was available in the equipment, and
physiological and extracortical noises were addressed statistically (see below) using the NIRS
Brain AnalyzeIR toolbox (88) in MATLAB 2021a (Mathworks, Natick, Massachusetts). Hbo
typically has a stronger signal-to-noise ratio (69) and therefore was used for the analysis
presented in the main paper. The same analysis can be performed with Hbr (supplementary
Table S5, S6).
516
fNIRS data was processed with the NIRS Brain AnalyzeIR toolbox (88) following the procedures
described in our previous work (33). Data for a given task were excluded from processing and
analysis (0.01% of the tasks) if there were experimental errors or if the participant clearly violated
the protocol (e.g., walking during a standing task). The fNIRS recording for each trial was first
trimmed to keep only 2 s of the data before and after the first and last task of a trial to reduce
global baseline noise. Channels with a variance less than 10^{-9} in a 5-s moving window, likely due
to equipment malfunction, were identified as flat (0.01% of the data) and removed from analysis.
Light intensity was converted to optical density and then converted to Hbo measurements using
the modified Beer–Lambert law with a partial path length factor of 0.1 (34). The time series data
for each source–detector pair was used to fit a general linear model (GLM). The design matrix of
the GLM was the convolution of stimulus timing, duration, and a canonical hemodynamic
response function (34). To minimize motion and physiological artifacts, the model was solved with
an autoregressive pre-whitening iteratively reweighted least square approach (88). In brief, the
autoregressive filter is a statistical method to alleviate physiological noise and motion artifacts.
The iterative reweighted least square approach further downweighs large motion artifacts (87–
89). A Student's t-test was then performed on the regression coefficients, and the t-score
represents the changes in Hbo in each task compared to the standing rest while counting silently

immediately before. Higher values represent more neural activity in the experimental task relative
to standing rest. Notice that this analysis approach also alleviates the need to adjust for age-
dependent differential pathlength factor because the mean and variance from the GLM will scale
accordingly. The results across six trials and eight channels covering the whole PFC were
combined using a weighted average that accounts for the covariance using the toolbox.
539
540 ***Attentional Control of Walking Outcome Measure***
541 Attentional control of walking was measured by attentional gait index that combines PFC
542 activations and dual-task performance (Equation 3, ref Liu 2024).

$$\text{Attentional Gait Index} = \text{Gain} * PFC_{Activation} \qquad (\text{Eq. 3})$$

544 $$\text{where } \text{Gain} = f(\Delta Motor + \Delta Cognitive) = e^{-\alpha * (\Delta Motor + \Delta Cognitive)},$$

546 $$\Delta Motor = \frac{MotorPerformance_{dual-task} - MotorPerformance_{single-task}}{MotorPerformance_{single-task}},$$

$$\Delta Cognitive = \frac{CognitivePerformance_{dual-task} - CognitivePerformance_{single-task}}{CognitivePerformance_{single-task}}$$

The optimization of the hyperparameter α and index calculation followed the same procedure as
described in our previous work (33). The definition of ΔMotor and ΔCognitive is similar to dual-
task cost in existing literature (90, 91) and the normalization relative to single-task allowed fair
comparisons among individuals with different baseline motor and cognitive performances.

The attentional gait index is a monotonic axis that quantifies the need for attentional control
during challenging walking, which has been shown to be more informative than PFC or dual-task
performance alone (33). Higher attentional gait index values represent greater PFC recruitment
and poorer performance in dual-task relative to single-task walking, which are indicative of
greater reliance on attentional resources and reduced automatic control of gait (16, 18–21, 33).

Self-selected overground walking speed
Self-selected overground walking speed was calculated as the average speed walking back and
forth on a straight 9.2m walkway excluding turns for 150 strides during the baseline overground
shown in Figure 1A.

Functional Gait Assessment Protocol
The Functional Gait Assessment (FGA) was scored in older adults on an ordinal scale for 10
items with a max score of 30 and higher score indicates better balance function (30). The 10
items are: gait on level surface, change in gait speed, gait with horizontal and vertical head turns,
gait with 180° pivot turn, stepping over obstacles, gait with narrow base of support, gait with eyes
closed, backwards gait and stairs (30). Scoring of each of item is done on a 4-point ordinal scale
ranging from 0 indicating severe impairment to 3 indicating no impairment. The assessment was
always administered by the same experimenter (SL) to reduce inter rater bias.

Neuropsychological Assessment Battery
Older adults completed a comprehensive neuropsychological assessment in their first visit to the
lab. The battery was designed and reviewed by a clinical neuropsychologist (AW) and the tests
were administered and scored by trained psychometrists. General cognitive status was measured
with the Montral Cognitive Assessment (MoCA). Verbal ability/literacy was measured with the
Wide Range Achievement Test Fifth Edition (WRAT5) (92). Overall cognition was measured with
the Repeatable Battery for the Assessment of Neuropsychological Status (RBANS) (42), which
includes tests that factor onto five cognitive domain index scores: Attention (Digit Span, Coding),
Language (Picture Naming, Semantic Fluency),Visuospatial (Figure Copy, Line Orientation),
Immediate Memory (List Learning, Story Learning), and Delayed Memory (List Recall, Story

Recall, Figure Recall). A supplemental language test, the action verbal fluency test (93), was
administered because the task is sensitive to basal ganglia function and may be relevant in
locomotor savings (94). Executive function was measured with subtests of the Delis-Kaplan
Executive Functioning System: Color Word Interference, Verbal Fluency, and Trail Making Test.
Since these subtests were co-developed and co-normed, we combined them into a composite
executive function score by averaging scaled scores from Color-Word Interference condition 3,
Color-Word test 4, Trail Making Test condition 4, and Verbal Fluency conditions 1, 2, and 3 total
correct. Similarly, cognitive switching was quantified by a composite score averaging Color-Word
Interference test condition 3, Trail Making Test condition 4, and Verbal Fluency condition 3
category switching scaled scores from D-KEFS (95).
594
Power Analysis
This study aims to evaluate the correlation between attentional control of walking and locomotor
savings in older adults. A sample of 21 older adults achieves 80% power to detect a moderate
(96) correlation (r = 0.57) at α = 0.05. To compare across age, a sex-matched sample of 21
younger adults was also recruited. This total sample size provides 80% power to detect a large
(97) between-group difference in locomotor savings (Cohen's d = 0.89) at α = 0.05. All power
analyses were conducted in R (version 4.5.1) using the pwr package.
602
Statistical analysis
In the primary analysis, t-test with equal variance was used to compare locomotor savings index
between older and younger adults after checking that normality (Shapiro-Wilk test, p = 0.24 and p
= 0.58 for older and younger adults respectively) and homoscedasticity were satisfied (Levene's
absolute test, p = 0.18). A Spearman's correlation was performed for each age group to quantify
the relationship between locomotor savings and attentional control of walking (measured by the
attentional gait index), because we were interested in quantifying the monotonic relationship and
the attentional gait index was not always normal.
611
In the exploratory analysis, due to violations of normality, non-parametric tests were performed.
To compare age-related differences in attentional gait index and PFC activation, Wilcoxon's rank
sum test was performed. To compare difference in motor or cognitive performance in single- and
dual-task walking as well as differences in steady state split-belt treadmill walking, aligned rank
tests were performed (98–100) with age as the between-subjects factor and task (single- or dual-
task for motor and cognitive performance) or session (for steady state walking) as the within-
subjects factor. When a significant correlation was found in the primary analysis, we performed
exploratory Spearman's correlations to identify if the relationship was driven by any
subcomponents of the attentional gait index. Spearman's correlation was also performed to
explore the relationship between locomotor savings, functional gait assessment, and
neuropsychological test results.
623
624 A significance level of alpha = 0.05 was used for all analysis. Aligned rank test was performed in
625 R (version 4.5.1) and all other tests was performed in MATLAB 2021a (Mathworks, Natick,
626 Massachusetts).
627
628

629 **Acknowledgments**
630
631 This study was supported by National Institutes of Health 1R01AG089175-01, National Science
632 foundation Career Award 2419849, and Pittsburgh Pepper Center P30AG024827. SL is
633 supported by the National Institutes of Health R90DA060340-02S1. AW is supported by the
634 National Institutes of Health (K23AG076663). EB is supported by the National Institutes of Health
635 (T32AG021885).
636

The authors thank Devin Stafford, Elan Albalak, Emma Stein, Shaoyi Li, and Shaina Patel for their tremendous help during data collection. The authors thank Dr. Dulce M. Mariscal, Dr. Krista Fjeld, Marcela Gonzalez-Rubio, and Jiwon Choi for their insightful feedback on the manuscripts and help during data collection. The authors thank Nathan Brantly and Adwoa A. Awuah for their feedback on the manuscripts. The authors thank Dr. Ted Huppert for his feedback on the fNIRS statistical analysis.

References

1. C. McCrum, K. Karamanidis, P. Willems, W. Zijlstra, K. Meijer, Retention, savings and interlimb transfer of reactive gait adaptations in humans following unexpected perturbations. *Commun. Biol.* **1** (2018).

2. K. A. Day, K. A. Leech, R. T. Roemmich, A. J. Bastian, Accelerating locomotor savings in learning: Compressing four training days to one. *J. Neurophysiol.* **119**, 2100–2113 (2018).

3. F. Mawase, L. Shmuelof, S. Bar-Haim, A. Karniel, Savings in locomotor adaptation explained by changes in learning parameters following initial adaptation. *J. Neurophysiol.* **111**, 1444–1454 (2014).

4. R. T. Roemmich, A. J. Bastian, Two ways to save a newly learned motor pattern. *J. Neurophysiol.* **113**, 3519–3530 (2015).

5. D. J. Herzfeld, P. A. Vaswani, M. K. Marko, R. Shadmehr, A memory of errors in sensorimotor learning. *Science (80-.).* **345**, 1349–1353 (2014).

6. J. R. Morehead, S. E. Qasim, M. J. Crossley, R. Ivry, Savings upon re-aiming in visuomotor adaptation. *J. Neurosci.* **35**, 14386–14396 (2015).

7. C. McCrum, M. H. G. Gerards, K. Karamanidis, W. Zijlstra, K. Meijer, A systematic review of gait perturbation paradigms for improving reactive stepping responses and falls risk among healthy older adults. *Eur. Rev. Aging Phys. Act.* (2017). Available at: https://link.springer.com/article/10.1186/s11556-017-0173-7 [Accessed 9 January 2026].

8. M. H. G. Gerards, C. McCrum, A. Mansfield, K. Meijer, Perturbation-based balance training for falls reduction among older adults: Current evidence and implications for clinical practice. *Geriatr. Gerontol. Int.* (2017). Available at: https://onlinelibrary.wiley.com/doi/full/10.1111/ggi.13082 [Accessed 9 January 2026].

9. S. C. Webber, M. M. Porter, V. H. Menec, Mobility in older adults: a comprehensive framework. *Gerontologist* **50**, 443–50 (2010).

10. M. Rantakokko, *et al.*, Changes in life-space mobility and quality of life among community-dwelling older people: a 2-year follow-up study. *Qual. Life Res.* **25**, 1189–1197 (2016).

11. A. L. Rosso, J. A. Taylor, L. P. Tabb, Y. L. Michael, Mobility, disability, and social engagement in older adults. *J. Aging Health* **25**, 617–37 (2013).

12. C. J. Sombric, G. Torres-Oviedo, Cognitive and Motor Perseveration Are Associated in Older Adults. *Front. Aging Neurosci.* **13**, 1–16 (2021).

13. C. J. Sombric, H. M. Harker, P. J. Sparto, G. Torres-Oviedo, Explicit action switching interferes with the context-specificity of motor memories in older adults. *Front. Aging Neurosci.* **9** (2017).

14. A. Bakkum, S. M. Gunn, D. S. Marigold, How aging affects visuomotor adaptation and
retention in a precision walking paradigm. *Sci. Rep.* **11**, 1–8 (2021).

15. M. Woollacott, A. Shumway-cook, Attention and the control of posture and gait : a review
of an emerging area of research. **16**, 1–14 (2002).

16. D. J. Clark, Automaticity of walking: Functional significance, mechanisms, measurement
and rehabilitation strategies. *Front. Hum. Neurosci.* **9**, 1–13 (2015).

17. S. B. Festini, L. Zahodne, P. A. Reuter-Lorenz, Theoretical Perspectives on Age
Differences in Brain Activation: HAROLD, PASA, CRUNCH—How Do They STAC Up?
Oxford Res. Encycl. Psychol. 1–24 (2018).
https://doi.org/10.1093/acrefore/9780190236557.013.400.

18. C. Udina, *et al.*, Functional Near-Infrared Spectroscopy to Study Cerebral Hemodynamics
in Older Adults During Cognitive and Motor Tasks: A Review. *Front. Aging Neurosci.* **11**,
1–19 (2020).

19. A. L. Rosso, *et al.*, Neuroimaging of an attention demanding dual-task during dynamic
postural control. *Gait Posture* **57**, 193–198 (2017).

20. R. Holtzer, *et al.*, Online fronto-cortical control of simple and attention-demanding
locomotion in humans. *Neuroimage* **112**, 152–159 (2015).

21. P. H. S. Pelicioni, M. Tijsma, S. R. Lord, J. Menant, Prefrontal cortical activation measured
by fNIRS during walking: Effects of age, disease and secondary task. *PeerJ* **7**, e6833
(2019).

22. M. F. L. Ruitenberg, *et al.*, Neural correlates of multi-day learning and savings in
sensorimotor adaptation. *Sci. Rep.* **8**, 1–14 (2018).

23. K. Cassady, *et al.*, Neural predictors of sensorimotor adaptation rate and savings. *Hum.*
Brain Mapp. **39**, 1516–1531 (2018).

24. R. D. Seidler, B. S. Gluskin, B. Greeley, Right prefrontal cortex transcranial direct current
stimulation enhances multi-day savings in sensorimotor adaptation. *J. Neurophysiol.* **117**,
429–435 (2017).

25. E. M. Reuter, L. A. Leow, T. J. Carroll, Task Feedback Processing Differs Between Young
and Older Adults in Visuomotor Rotation Learning Despite Similar Initial Adaptation and
Savings. *Neuroscience* **451**, 79–98 (2020).

26. F. Mawase, S. Bar-Haim, L. Shmuelof, Formation of long-term locomotor memories is
associated with functional connectivity changes in the cerebellar-thalamic-cortical network.
J. Neurosci. **37**, 349–361 (2017).

27. K. B. Walhovd, *et al.*, Consistent neuroanatomical age-related volume differences across
multiple samples. *Neurobiol. Aging* **32**, 916–932 (2011).

28. R. Beurskens, O. Bock, Age-Related Deficits of Dual-Task Walking: A Review. *Neural*
Plast. **2012**, 131608 (2012).

29. R. Holtzer, *et al.*, fNIRS study of walking and walking while talking in young and old
individuals. *Journals Gerontol. - Ser. A Biol. Sci. Med. Sci.* **66 A**, 879–887 (2011).

30. D. M. Wrisley, G. F. Marchetti, D. K. Kuharsky, S. L. Whitney, Reliability, internal

consistency, and validity of data obtained with the functional gait assessment. *Phys. Ther.*
84, 906–18 (2004).

31. D. S. Reisman, H. McLean, J. Keller, K. A. Danks, A. J. Bastian, Repeated Split-belt
Treadmill Training Improves Poststroke Step Length Asymmetry. *Neurorehabil. Neural*
Repair **27**, 460–468 (2013).

32. L. A. Malone, E. V. L. Vasudevan, A. J. Bastian, Motor adaptation training for faster
relearning. *J. Neurosci.* **31**, 15136–15143 (2011).

33. S. Liu, *et al.*, Novel attentional gait index reveals a cognitive ability-related decline in gait
automaticity during dual-task walking. *Front. Aging Neurosci.* **15**, 2023.07.31.551290
(2024).

34. C. W. Hoppes, *et al.*, Changes in Cortical Activation during Dual-Task Walking in
Individuals with and without Visual Vertigo. *J. Neurol. Phys. Ther.* **44**, 156–163 (2020).

35. D. M. Wrisley, N. A. Kumar, Functional Gait Assessment: Concurrent, Discriminative, and
Predictive Validity in Community-Dwelling Older Adults. *Phys. Ther.* **90**, 761–773 (2010).

36. Functional Gait Assessment | RehabMeasures Database. Available at:
https://www.sralab.org/rehabilitation-measures/functional-gait-assessment [Accessed 9
January 2026].

37. M. Beninato, A. Fernandes, L. S. Plummer, Minimal Clinically Important Difference of the
Functional Gait Assessment in Older Adults. *Phys. Ther.* **94**, 1594–1603 (2014).

38. S. Coltman, P. Gribble, J. Cashaback, Both fast and slow learning processes contribute to
savings following sensorimotor adaptation. *J. Neurophysiol.* **121** (2019).

39. M. D. Lezak, D. B. Howieson, E. D. Bigler, D. Tranel, *Neuropsychological Assessment*,
Fifth (Oxford University Press, 2012).

40. E. Bolceková, *et al.*, Cognitive impairment in cerebellar lesions: a logit model based on
neuropsychological testing. *Cerebellum & Ataxias* **4**, 13 (2017).

41. F. Hoche, X. Guell, M. G. Vangel, J. C. Sherman, J. D. Schmahmann, The cerebellar
cognitive affective/Schmahmann syndrome scale. *Brain* **141**, 248–270 (2018).

42. C. Randolph, M. C. Tierney, E. Mohr, T. N. Chase, The Repeatable Battery for the
Assessment of Neuropsychological Status (RBANS): Preliminary clinical validity. *J. Clin.*
Exp. Neuropsychol. **20**, 310–319 (1998).

43. H. A. Clayton, S. Abbas, B. M. Hart, D. Y. P. Henriques, Visuomotor adaptation across the
lifespan. *PLoS One* **19**, e0306276 (2024).

44. R. Binyamin-Netser, N. Goldhamer, I. Avni, A. Ressel Zviely, L. Shmuelof, Cognitive
Impairments After Stroke Do Not Attenuate Explicit Visuomotor Adaptation in Reaching
and Savings With the Unaffected Arm. *Neurorehabil. Neural Repair* **37**, 444 (2023).

45. S. Bohm, L. Mademli, F. Mersmann, A. Arampatzis, Predictive and Reactive Locomotor
Adaptability in Healthy Elderly: A Systematic Review and Meta-Analysis. *Sport. Med.* **45**,
1759–1777 (2015).

46. R. T. Roemmich, *et al.*, Locomotor adaptation and locomotor adaptive learning in
Parkinson's disease and normal aging. *Clin. Neurophysiol.* **125**, 313–319 (2014).

47. C. Rossi, R. T. Roemmich, N. Schweighofer, A. J. Bastian, K. A. Leech, Younger and Late
Middle-Aged Adults Exhibit Different Patterns of Cognitive-Motor Interference During
Locomotor Adaptation, With No Disruption of Savings. *Front. Aging Neurosci.* **13**, 1–18
(2021).

48. L. A. Leow, A. de Rugy, W. Marinovic, S. Riek, T. J. Carroll, Savings for visuomotor
adaptation require prior history of error, not prior repetition of successful actions. *J.*
Neurophysiol. **116**, 1603–1614 (2016).

49. L. A. Leow, W. Marinovic, A. de Rugy, T. J. Carroll, Task Errors Drive Memories That
Improve Sensorimotor Adaptation. *J. Neurosci.* **40**, 3075–3088 (2020).

50. R. Shadmehr, T. Brashers-Krug, Functional Stages in the Formation of Human Long-Term
Motor Memory. *J. Neurosci.* **17**, 409–419 (1997).

51. L. A. Malone, A. J. Bastian, Age-related forgetting in locomotor adaptation. *Neurobiol.*
Learn. Mem. **128**, 1–6 (2016).

52. C. Krishnan, E. P. Washabaugh, C. E. Reid, M. M. Althoen, R. Ranganathan, Learning
new gait patterns: Age-related differences in skill acquisition and interlimb transfer. *Exp.*
Gerontol. **111**, 45–52 (2018).

53. M. A. Smith, A. Ghazizadeh, R. Shadmehr, Interacting Adaptive Processes with Different
Timescales Underlie Short-Term Motor Learning. *PLoS Biol.* **4**, e179 (2006).

54. W. M. Joiner, M. A. Smith, Long-term retention explained by a model of short-term
learning in the adaptive control of reaching. *J. Neurophysiol.* **100**, 2948–2955 (2008).

55. K. M. Trewartha, X. A. Garcia, X. D. M. Wolpert, X. J. R. Flanagan, Fast But Fleeting□:
Adaptive Motor Learning Processes Associated with Aging and Cognitive Decline. **34**,
13411–13421 (2014).

56. S. B. Swart, A. R. den Otter, C. J. C. Lamoth, Natural ageing primarily affects the initial
response to a sustained walking perturbation but not the ability to adapt over time. *Front.*
Physiol. **14**, 1065974 (2023).

57. S. Hiew, L. Eibeck, C. Nguemeni, D. Zeller, The Influence of Age and Physical Activity on
Locomotor Adaptation. *Brain Sci. 2023, Vol. 13, Page 1266* **13**, 1266 (2023).

58. C. McCrum, *et al.*, Locomotor stability and adaptation during perturbed walking across the
adult female lifespan. *J. Biomech.* **49**, 1244–1247 (2016).

59. N. Seethapathi, B. C. Clark, M. Srinivasan, Exploration-based learning of a stabilizing
controller predicts locomotor adaptation. *Nat. Commun.* **15**, 9498 (2024).

60. V. S. Huang, A. Haith, P. Mazzoni, J. W. Krakauer, Rethinking motor learning and savings
in adaptation paradigms: model-free memory for successful actions combines with internal
models. *Neuron* **70**, 787 (2011).

61. L. A. Leow, A. de-Rugy, A. M. Loftus, G. Hammond, Different mechanisms contributing to
savings and anterograde interference are impaired in Parkinson's disease. *Front. Hum.*
Neurosci. **7**, 39031 (2013).

62. L. M. Chahine, *et al.*, Nigrostriatal dopaminergic neurotransmission and resilience to
peripheral systemic risk factors for gait slowing upon transition to uneven surfaces in older
adult. *Aging Brain* **8**, 100143 (2025).

63.   A. L. Rosso, *et al.*, Nigrostriatal dopaminergic integrity in relation to prefrontal cortex
802   activity and gait performance during dual-task walking in older adults. *Journals Gerontol. -*
803   *Ser. A Biol. Sci. Med. Sci.* **80** (2025).

804   64.   S. Bierbaum, A. Peper, K. Karamanidis, A. Arampatzis, Adaptive feedback potential in
805   dynamic stability during disturbed walking in the elderly. *J. Biomech.* **44**, 1921–1926
806   (2011).

807   65.   J. Falk, *et al.*, Increased co-contraction reaction during a surface perturbation is
808   associated with unsuccessful postural control among older adults. *BMC Geriatr.* **22**, 438-
809   (2022).

810   66.   R. A. J. Koster, *et al.*, Balance training in older adults enhances feedback control after
811   perturbations. *PeerJ* **12**, e18588 (2024).

812   67.   T. H. Haid, A. C. M. Doix, B. M. Nigg, P. A. Federolf, Age effects in postural control
813   analyzed via a principal component analysis of kinematic data and interpreted in relation
814   to predictions of the optimal feedback control theory. *Front. Aging Neurosci.* **10**, 310279
815   (2018).

816   68.   K. A. Leech, R. T. Roemmich, Independent voluntary correction and savings in locomotor
817   learning. *J. Exp. Biol.* **221** (2018).

818   69.   J. C. Menant, *et al.*, A consensus guide to using functional near-infrared spectroscopy in
819   posture and gait research. *Gait Posture* **82**, 254–265 (2020).

820   70.   E. D. Thompson, D. S. Reisman, Split-Belt Adaptation and Savings in People with
821   Parkinson Disease. *J. Neurol. Phys. Ther.* **46**, 293–301 (2022).

822   71.   T. Ghanavati, *et al.*, Deep white matter hyperintensities, microstructural integrity and dual
823   task walking in older people. *Brain Imaging Behav.* **12**, 1488–1496 (2018).

824   72.   H. T. Karim, *et al.*, Resting state connectivity within the basal ganglia and gait speed in
825   older adults with cerebral small vessel disease and locomotor risk factors. *NeuroImage*
826   *Clin.* **28**, 102401 (2020).

827   73.   A. L. Rosso, *et al.*, Contributors to Poor Mobility in Older Adults: Integrating White Matter
828   Hyperintensities and Conditions Affecting Other Systems. *J. Gerontol. A. Biol. Sci. Med.*
829   *Sci.* **72**, 1246–1251 (2017).

830   74.   J. B. Hengenius, N. I. Bohnen, A. Rosso, T. J. Huppert, C. Rosano, Cortico-striatal
831   functional connectivity and cerebral small vessel disease: Contribution to mild
832   Parkinsonian signs. *J. Neuroimaging* **32**, 352–362 (2022).

833   75.   S. W. Ducharme, J. A. Kent, R. E. A. van Emmerik, Comparable Stride Time Fractal
834   Dynamics and Gait Adaptability in Active Young and Older Adults Under Normal and
835   Asymmetric Walking. *Front. Physiol.* **10**, 478311 (2019).

836   76.   P. M. Dunlap, *et al.*, Global Positioning System Indicators of Community Mobility and
837   Future Health Outcomes Among Older Adults. *Journals Gerontol. Ser. A* **79**, 1–8 (2024).

838   77.   M. F. L. Ruitenberg, *et al.*, Multi-day Adaptation and Savings in Manual and Locomotor
839   Tasks. *J. Mot. Behav.* **50**, 517–527 (2018).

840   78.   J. A. Roper, *et al.*, Higher relative effort of the knee relates to faster adaptation in older
841   adults at risk for mobility disability. *Exp. Gerontol.* **144**, 111192 (2021).

79. S. Sasikumar, G. Sorrento, A. E. Lang, A. P. Strafella, A. Fasano, Cognition affects gait
adaptation after split-belt treadmill training in Parkinson's disease. *Neurobiol. Dis.* **181**,
106109 (2023).

80. N. A. Jacobsen, D. P. Ferris, Exploring Electrocortical Signatures of Gait Adaptation:
Differential Neural Dynamics in Slow and Fast Gait Adapters. *eNeuro* **11** (2024).

81. Q. Tian, *et al.*, Dual cognitive and mobility impairments and future dementia - Setting a
research agenda. *Alzheimer's Dement.* **19**, 1579–1586 (2023).

82. G. Torres-Oviedo, A. J. Bastian, Seeing Is Believing: Effects of Visual Contextual Cues on
Learning and Transfer of Locomotor Adaptation. *J. Neurosci.* **30**, 17015–17022 (2010).

83. G. Torres-Oviedo, A. J. Bastian, Natural error patterns enable transfer of motor learning to
novel contexts. *J. Neurophysiol.* **107**, 346–356 (2012).

84. P. A. Iturralde, G. Torres-Oviedo, Corrective Muscle Activity Reveals Subject-Specific
Sensorimotor Recalibration. *eneuro* **6**, ENEURO.0358-18.2019 (2019).

85. D. de Kam, P. A. Iturralde, G. Torres-Oviedo, Cerebral Contribution to the Execution, But
Not Recalibration, of Motor Commands in a Novel Walking Environment. *eneuro* **7**,
ENEURO.0493-19.2020 (2020).

86. F. Herold, *et al.*, Functional near-infrared spectroscopy in movement science: a systematic
review on cortical activity in postural and walking tasks. *Neurophotonics* **4**, 041403 (2017).

87. K. Bohlke, *et al.*, Exercise interventions, postural control, and prefrontal cortex activation
in older adults. *Brain Cogn.* **171**, 106063 (2023).

88. H. Santosa, X. Zhai, F. Fishburn, T. Huppert, The NIRS Brain AnalyzIR toolbox.
Algorithms **11** (2018).

89. J. W. Barker, A. Aarabi, T. J. Huppert, Autoregressive model based algorithm for
correcting motion and serially correlated errors in fNIRS. *Biomed. Opt. Express* **4**, 1366
(2013).

90. F. O. Brauner, *et al.*, The Performance Index Identifies Changes Across the Dual Task
Timed Up and Go Test Phases and Impacts Task-Cost Estimation in the Oldest-Old.
Front. Hum. Neurosci. **15**, 1–12 (2021).

91. J. K. Longhurst, *et al.*, A Novel Way of Measuring Dual-Task Interference: The Reliability
and Construct Validity of the Dual-Task Effect Battery in Neurodegenerative Disease.
Neurorehabil. Neural Repair **36**, 346–359 (2022).

92. G. S. Wilkinson, G. J. Robertson, *(WRAT5) Wide Range Achievement Test*, Fifth
(Pearson Assessment, 2017).

93. S. P. Woods, *et al.*, Action (verb) fluency: Test-retest reliability, normative standards, and
construct validity. *J. Int. Neuropsychol. Soc.* **11**, 408–415 (2005).

94. D. C. Hinton, D. M. Conradsson, C. Paquette, Understanding Human Neural Control of
Short-term Gait Adaptation to the Split-belt Treadmill. *Neuroscience* **451**, 36–50 (2020).

95. D. C. Delis, E. Kaplan, J. H. Kramer, *Delis-Kaplan Executive Function System* (Pearson
Assessment, 2001).

96.   H. Akoglu, User's guide to correlation coefficients. *Turkish J. Emerg. Med.* **18**, 91–93
(2018).

97.   J. Cohen, *Statistical Power Analysis for the Behavioral Sciences* (Routledge, 2013).

98.   M. Kay, L. Elkin, J. J. Higgins, J. O. Wobbrock, ARTool: Aligned Rank Transform for
Nonparametric Factorial ANOVAs. Available at: https://github.com/mjskay/ARTool.

99.   J. O. Wobbrock, L. Findlater, D. Gergle, J. J. Higgins, The aligned rank transform for
nonparametric factorial analyses using only anova procedures in *Proceedings of the
SIGCHI Conference on Human Factors in Computing Systems*, (ACM, 2011), pp. 143–
146.

100. L. A. Elkin, M. Kay, J. J. Higgins, J. O. Wobbrock, An Aligned Rank Transform Procedure
for Multifactor Contrast Tests in *The 34th Annual ACM Symposium on User Interface
Software and Technology*, (ACM, 2021), pp. 754–768.

Figures and Tables
895

896
Figure 1. Experimental protocols and key outcome measures. (A) Schematics of the split-belt
treadmill walking paradigm to evaluate locomotor savings. Participants completed three sessions
on separate days: pre-practice, practice, and post-practice with each session 1 or 2 days apart. In
the pre-practice session, participants learned a new walking pattern on a split-belt treadmill.
Locomotor savings quantifies the ability to recall this newly learned pattern after practice and was
measured by the change in step length asymmetry (StepAsym, B) in the post- compared to pre-
practice session. Patched blocks represent self-paced overground walking, white blocks
represent tied-belt treadmill walking, grey blocks represent split-belt walking, and dashed grey
vertical lines represent breaks. (B) Illustrations of StepAsym parameter calculation where zero
represents symmetric walking. Step length (SL) is defined as the distance between two ankles
during heel strike. (C) Schematics of the dual-task walking paradigm with prefrontal cortical (PFC)
recordings from functional near infrared spectroscopy (fNIRS) to evaluate attentional control of
walking. Both motor (walking speed) and cognitive (correct number of alphabet letters generated
per second) performances were measured. Participants performed motor (walking) and cognitive
(reciting every two or three letters of the alphabet) tasks either separately (single-task) or
simultaneously (dual-task) in randomized order for 6 repetitions.
913
914

Figure 2. Step length asymmetry (StepAsym) time course (A) with inserts comparing locomotor savings (B) and steady state walking (C) between older and younger adults. (A) Group averaged time course of StepAsym during the adaptation condition (split-belt) in pre- (closed circles) and post-practice (open circles) sessions. All data is baseline (tied-belt) removed. Closer to zero values represent more symmetric (i.e., less perturbed) walking. Dot represents the average of 5 consecutive strides, and shaded area represents the standard error of each group. Both groups showed typical adaptation trajectories in both sessions (closer to zero StepAsym towards the end of adaptation), but the difference in early adaptation (grey shaded area) between pre- and post-practice sessions is larger in younger adults (red) compared to older adults (blue). (B) Locomotor savings. Bar plot represents the mean per group ± standard deviation. Dot represents the locomotor saving index value, calculated by equation shown in the figure. Older adults showed significantly less locomotor savings than younger adults. (C) StepAsym during steady state walking. Bar plot shows the mean per group ± standard deviation, and each dot corresponds to the average StepAsym during steady state walking (last 40 strides of adaptation) per participant. All participants reached the same steady state after adaptation in both sessions (as shown by the non-significant age and session effect).

Figure 3. Attentional control of gait (measured by the Attentional Gait Index) comparisons between older and younger adults (A) and associations with locomotor savings (B). (A) Older adults showed significantly higher Attentional Gait Index values than younger adults. Bar plot represents the group mean per group ± standard deviation. Dot represents Attentional Gait Index (a metric combining prefrontal cortical activations and performance change from single- to dual-task walking) of each participant. (B) Scatter plots of Attentional Gait Index vs. locomotor savings. For visualization purpose, a linear regression line was displayed when we found a significant correlation. A significant positive correlation was observed in older adults.

944 **Table 1. Participant characteristics**.

Variables	Older Adults Mean (SD) or n (%)	Younger Adults Mean (SD) or n (%)
Sample size	21	21
Age (years)	71.85 ± 3.14	25.05 ± 5.20
Female, n (%)	11 (52.38%)	11 (52.38%)
Race, n (%)		
White	19 (90.48%)	16 (76.19%)
Black or African American	1 (4.76%)	1 (4.76%)
More than one race	1 (4.76%)	0
Asian	0	4 (19.05%)
Highest level of education, n (%)		
Secondary level education / GED / Equivalent	2 (9.52%)	0
Some college	4 (19.05%)	8 (38.10%)
Bachelor's degree or higher	15 (71.43%)	13 (61.90%)
Right leg dominant, n (%)	21 (100%)	20 (95.24%)
Self-selected overground walking speed (m/s)	0.96 ± 0.17	1.01 ± 0.14
Montreal Cognitive Assessment (MoCA) score (max 30)	27.05 ± 1.47	not administered
Functional Gait Assessment (FGA) score (max 30)	24.71 ± 2.43	not administered