Water Stress Induces Diel Patterns of Root Growth

Richard Nair[1, 2], Martin Strube[2], Javier Pacheco-Labrador[3], Marion Schrumpf Schrump[2,4], Mirco Migliavacca[5]

1. Discipline of Botany, School of Natural Sciences, Trinity College Dublin, Dublin Ireland

   Department of Biogeochemical Integration, Max Planck Institute for Biogeochemistry, 07745 Jena, Germany

2. Environmental Remote Sensing and Spectroscopy Laboratory (SpecLab), Spanish National Research Council, 28037 Madrid, Spain Marions new affiliation

3. European Commission, Joint Research Centre (JRC), Ispra, Italy

ABSTRACT

Growth and death of vegetation control both the amount of carbon held in living biomass and much of the flow between the atmosphere and soils, which is the largest terrestrial carbon reservoir. However, long-term field datasets on belowground growth dynamics are rare. Knowledge is lacking on how root growth patterns relate to the environment, fluctuating resources availability such as carbon or water, and internal plant physiological control at both seasonal and diel scales. We imaged roots using automated minirhizotrons and interpreting root length density (RLD) automatically twice a day for nine months in a temperate grassland. From winter to autumn, we always observed some root growth in individual images, but with decreasing net RLD (indicative of net turnover) becoming predominant

across all imaging locations later in the season. Roots grew day-round except during a period of about

one month in spring when root growth was only in daytime. This was a period without rain, during

which soil moisture depleted rapidly, but not at its driest point. We discuss potential causes for both

overall and short-term growth patterns, particularly the change in drivers of growth from internal

storage to new photosynthates, or limits imposed by water availability - both soil moisture and rainfall -

under shifting environmental conditions.

INTRODUCTION
29

High-frequency measurement of ecosystem carbon (C) and water fluxes is now routine (Baldocchi, -

2020), revealing patterns in land-atmosphere exchange, often due to combined variation in meteorology

and partially coupled plant growth and turnover (e.g. Y. Liu & Wu, 2020; C. Wu et al., 2012). However,

validation from whole-plant dynamics in-field is often opaque. While remote sensing provides above-

ground information, at least 1/3 of plant biomass (Jackson *et al.*, 1996; McCormack *et al.*, 2015) and half

of total productivity (Gherardi & Sala, 2020) is allocated belowground to 'cryptic' roots, mycorrhizal

fungi, and exudates. When studied, root phenology is commonly asynchronous with leaves (Abramoff &

Finzi, 2015), but measurements of roots at high frequency are also far less frequent. Overall,

observational data on root growth dynamics on diel to seasonal scales is sparse compared to above-

ground observations.

At seasonal scale, precipitation, temperature, and light control carbon and water availability, and soil

nutrient turnover (e.g. Graham et al., 2003; Huete et al., 2006; Z. Wu et al., 2021), and ultimately plant

phenology (e.g. Luo et al., 2020; Zelikova et al., 2015). Meteorological conditions and light availability

also oscillate between day and night, influencing resource supply above-ground. Circadian rhythms help

plants buffer these conditions, but activity still has diel patterns (Rozema *et al.*, 1987; Walter *et al.*,

2009; Poiré *et al.*, 2010). Cell turgor limits initial leaf expansion (Pantin *et al.*, 2011), often favouring

nighttime (Rozema *et al.*, 1987; Poiré *et al.*, 2010), and similar patterns underly radial tree stem growth

(Zweifel *et al.*, 2021). Above-ground, turgor limits imposed by hydraulic conductance are a key growth

driver.

Roots are located at opposite ends of the vascular system to leaves, so growth may relate differently to

water and carbon (Ruts *et al.*, 2012). They inhabit a dense, environmentally buffered, heterogeneous

environment (Morris & Nair, 2025) and have different meristem mechanisms to shoots (Stahl & Simon,

2010). Daytime (Halter *et al.*, 1996) and nighttime (Head, 1965) tree root growth has been reported in

walk-in rhizotrons, but generally, roots tend to grow at night or around dawn (Ruts *et al.*, 2012) and

respond to environmental fluctuations (Bengough *et al.*, 2011; Heinze *et al.*, 2023; Ceolin *et al.*, 2025).

These data usually come from cuttings, isolated plants, short observations, or artificial conditions

(Walter *et al.*, 2002, 2009, (Yazdanbakhsh & Fisahn, 2009), whereas high frequency field measurements

are limited (c.f. Belaud et al., 2024; Kuwabe & and Ohashi, 2023).

Here we used automated minirhizotron root cameras (Nair *et al.*, 2022) to examine sub-daily fine root

growth for 9 months in a temperate grassland. Over 250,000 images were analysed using a

convolutional neural network (Smith *et al.*, 2022) to track root length density (RLD) and rooting depth.

We aimed to answer the following questions:

Q1: **Under which environmental conditions do roots grow and die?** We predicted that we would

observe the highest growth rates during warm periods when soil moisture was not limiting. We

expected to see accelerating growth due to the cumulative effect of increased green biomass through

spring, leading to increased C available to grow roots. We expected a seasonal cycle with net turnover

later in the year.

Q2: **Is there a diel pattern of root growth, and if so, does this shift seasonally?** Following the evidence

above, we expected night-time growth (if turgor limited) or no pattern (if it didn't), without priors for

seasonal changes beyond soil water availability.

RESULTS

Net RLD growth was immediate when measurements began in February and continued until June, when

it started to decline. Although the instruments matched each other in terms of gross patterns (Figure

S3), the absolute values of RLD observed at each instrument microsite differed (Figure 1a),. The mean

rooting depth (MRD) deepened with a deepening soil moisture profile going into summer (Figure 1b,

Figure 1d). The number of images with growing roots declined throughout the experiment (Figure S4).

Minor recovery of MRD in September was driven by new roots in shallow soil layers (Figure S5).

We calculated daytime growth as the 3-day smoothed difference between the previous night and the

current day, and vice-versa, for nighttime growth. Across the entire 40 cm depth, roots grew evenly

between day and night until mid-April, when growth was predominantly during the day (Figure 1c). This

returned to no difference by late May. Day-and-night growth was significantly different (sign test, $P <$

0.05) through most of May. No sustained or significant desynchrony was observed afterwards.

In April, there was a small but notable increase in rooting depth, which preceded decreasing soil

moisture. Desynchrony between day and night occurred during this dry period, which was notable

because was the longest precipitation gap in the entire timeseries (Figure 1c, Figure 1e).

DISCUSSION

We investigated the timing of root growth in the field to identify 1) what environmental conditions roots

grow and die, and 2) if there are diel patterns. We assumed that expansive growth is maximised at

night/early morning, because during daytime, transpiration depletes water necessary for cell expansion

(Walter et al., 2002). In lab-grown Arabidopsis, nighttime root elongation is under circadian control

(Yazdanbakhsh et al., 2011) but imprints via hydrological conditioning (Caldeira et al., 2014). We found

continuous root growth and only found diel desynchrony under late spring water deficits (Figure 1c).

Here, more roots were produced during day than at night.

Growth could be limited by demand for resources, *source control* or other limits to use of these

resources, *sink control* (Körner, 2015). Desynchrony could be due to 1) interruption or variation of C

supply (source control) or 2) external temperature constraints, 3) a shift in internal partitioning or 4)

interruption in water available for turgor. Explanations 2, 3, and 4 are sink controls. We discount shifts in

endogeny because diurnal shifts only occurred at a single period, and our measurements were

community scale. Overall, we interpret the interplay between carbon and water stress determining

short-term desynchrony, as follows:

Did carbon fixation (source control) control growth?

Net RLD increased steadily until May (Figure 1a), and some roots grew throughout the year even when

net RLD declined (S3). Early in the growing season, C fixation was likely low, yet the trajectory of long-

term RLD increase February-May was almost linear, suggesting no C limitation. Roots may respond

slower to diurnal or weather-related shifts in C fixation than leaves due to C transport in the phloem,

which operates via pressure-driven bulk flow and osmotic gradient regulation, (Münch, 1930;

Mencuccini & Höltta, 2010; Jensen et al., 2016). Transport time constraints are consistent with observed

lags between photosynthesis and soil respiration in grassland ecosystems (Bahn et al., 2009). In such

short-stature systems, leaf to root tip distance is short. Phloem sugar concentrations vary diurnally

(Jensen et al., 2016), but with no consistent diurnal RLD change for most of our grassland timeseries,

either pressure flow equilibrates supply and/or photosynthates were always sufficient. Either way,

diurnal growth did not appear to be limited by C supply.

112   <u>Did temperature control growth?</u>

Aboveground growth in temperate grasses becomes negligible below 5°C (Peacock, 1975; Wingler &

Hennessy, 2016), but photosynthesis continues at lower temperatures (Körner, 2008) and leaf expansion

is phytohormone regulated (Wingler & Hennessy, 2016). Soils are buffered from temperature and may

be warmer than air; this was observed in February (Figure 1e), which may have allowed root growth

despite stalled leaf production. RLD increased as soon as observations began (Figure 1a), initially mostly

close to the surface (Figure 1b), followed by deepening roots as the season progressed. Summer mean

daily air temperatures, rarely exceeding 25°C (optimal for temperate plants (Hatfield & Prueger, 2015)),

were steadily increasing during the desynchrony period, suggesting temperature did not limit growth or

drive the pattern.

122   <u>Did we miss carbon assignment to exudates or mycorrhizal fungi?</u>

Diel asynchrony in root observations could potentially be due to variations in exudate or mycorrhizal

components (i.e. total C remains the same, relative assignment different). Although automated

minirhizotrons have observed mycorrhizal fungi (Allen & Kitajima, 2013; Defrenne et al., 2021), our

systems were not designed for fungal hyphal resolution. Minirhizotron resolution usually trades off

against field of view; high 'zoom' increases imaging time. Relevant fungal structures are further probably

limited to rhizomorphs in ectomycorrhizal system (grasslands are dominated by arbuscular mycorrhizal

fungi (Brundrett, 1991)) with less obvious effect on root phenotype. In any case, total C allocation to

mycorrhizal fungi is around 1-13 % (Hawkins et al., 2023). In grasslands, most photo-assimilates also

remain in plants in the short term (Ostle *et al.*, 2007). From lab experiments, exudate patterns are also

probably not diurnal (Badri *et al.*, 2010), and we could not find diel patterns in mycorrhizal exchange.

Complete belowground ^{13}C traces through to efflux CO_2 also do not reveal diurnal patterns (Liu *et al.*,

2021). Thus, there are not likely alternate cryptic belowground sinks, which vary on diel or seasonal

scales.

Water and Turgor limited growth

Overall, seasonal patterns observed in this study matched root observations in similar ecosystems

(Edwards *et al.*, 2004; Steinaker & Wilson, 2008). Desynchrony (Figure 1c) occurred prior to a RLD

decline over summer (Figure 1a). During this period, soil moisture (Figure 1e) and MRD (Figure 1b)

decreased. Thus, the ecosystem was in transition from abundant to scarce soil water availability. Pasture

grasses are susceptible to hydraulic failure and follow embolism avoidance strategies (Jacob *et al.*,

2022). Leaf expansion is tightly coupled to soil water potential, but root growth is less so (Majerus,

1975), perhaps because of root proximity to the source of water . Because root extension may supply

water, but leaves extends the transpiration stream, root growth may continue while leaf growth stalls.

Photosynthesis likely persisted under mild soil drought conditions, so there was fresh C available.

Sentinel-2 Normalized Difference Vegetation Index (NDVI) remained stable during this desynchrony

(Figure S6). Thus, daytime root growth could reflect diverted C from non-growing leaves, while

nighttime was limited by water being used to refill depleted xylems. The plants may no longer have been

able to reach turgor for root cell growth as the xylem refilled. Supporting this, MRD deepened before

desynchrony, ahead of soil moisture decline, perhaps searching for additional water driven by leaf stress

(Figure 1b, Figure 1d). The absence of rainfall during this period (Figure 1d) may have further shifted

functional activity, with root building influenced by increasing leaf dryness and xylem vulnerability, even

before soil moisture levels were fully depleted.

Wider Implications

Growth timing matters for carbon budgets. In soil, respiration can be partitioned into root and heterotroph sources (Högberg *et al.*, 2009; Taylor *et al.*, 2015; Savage *et al.*, 2018; Sun *et al.*, 2019), or into growth, maintenance, and redox balancing components at plant level (De Vries *et al.*, 1974; Johnson, 1990; Hirano *et al.*, 2023). However, such partitioning requires assumptions, since data from undisturbed systems is scarce (Hirano *et al.*, 2023). Respiration temperature sensitivity is usually modelled using Q_{10} enzyme kinetics, applied in ecosystem models (Meyer *et al.*, 2018), partitioning of eddy covariance data (e.g. Reichstein *et al.*, 2005), and interpreting processes such as acclimation and adaptation (King *et al.*, 2006) plus other physiological insights. However, growth and maintenance have a different response to temperature, being driven by substrate sink/source limits and rates of reaction, respectively (De Vries *et al.*, 1974). If their timing diverges, daily or long term average Q_{10} may not reflect underlying processes.

Other evidence supports this view. Root temperature does not affect growth rates or respiration coefficients (Szaniawski & Kielkiewicz, 1982) but does affect total respiration via substrate supply (Atkin *et al.*, 2000; Thurgood *et al.*, 2014). Ecosystem respiration mostly comes from belowground (Raich & Schlesinger, 1992), with roots contributing 42 ± 18% (Jian et al., 2022). Root respiration has the highest Q_{10} of numerous factors contributing to soil CO_2 fluxes (Boone *et al.*, 1998), so our finding that root growth timing can switch between continuous and diel patterns under stress indicates such ecosystem generalizations may not hold consistently.

Similarly, models used to predict climate change are parameterised at half-hour scales for fluxes but fit to biomass over months to years. Fine roots are often simply prescribed via fixed ratios, thresholds, or optimality assumptions (e.g., Clark et al., 2011, Thum et al., 2019), without firm validation. Here we show diel desynchrony under environmental stress. Further work could link to above-ground dynamics

177 and be used for model improvement, allowing teasing out source-sink controls and improving predictive

178 capacity of coupled biogeochemical cycles. Understanding overall ecosystem activity may depend on

179 what roots are doing and when they are doing it.

MATERIALS AND METHODS

Field Installation

The study was conducted at a grassland on the floodplain of the river Saale in Germany 50°55◻N,

11°35◻E. The site is a loamy Eutric fluvisol and hosts the Jena biodiversity experiment (Roscher *et al.*,

2004; Weisser *et al.*, 2017). We set up automated minirhizotron instruments in two ancillary plots

without manipulation. The plots were unmanaged except for two cuts during the growing season when

the above ground biomass was manually clipped above the instruments corresponding to mowing of the

wider experiment.

We used an minirhizotron instrument design to capture images autonomously (Nair *et al.*, 2022). In

brief, this system took a series of images around a 1 m (length) x 10 cm (external diameter) observatory

made of plexiglass. Roots grew around the observatory and were visible in the images. Image resolution

was 25 μm per pixel, which covered the root diameters expected at the site. We installed observatories

in September 2021 at 40 degrees to the horizontal, for a vertical span of ~ 40 cm and a horizontal span

of ~ 70 cm, following best practice (Box *et al.*, 1989; Johnson *et al.*, 2001; Strand *et al.*, 2018; Freschet *et*

al., 2021) to ensure a tight fit between soil and tube. The length of 'rest' time before observations varies

in the literature; here it was five months. Issues like soil-tube gaps and compaction may influence root

behaviour (Bengough *et al.*, 2006) and even follow hydrological changes in soil properties. As our rest

period was over winter, it is likely that there was minimal root growth but time for soil expansion in the

wetter part of the year.

We deployed two automatic minirhizotrons on 1 Feb 2022 and added a further two in Jul 2022. The

instruments were in two clusters but because of the short depth of field of minirhizotron imagery

(Johnson *et al.*, 2001), we treated each minirhizotron as independent. The instruments ran on mains

power and sampled at 01:00, 09:00, 13:00, and 21:00 UTC. At each sampling interval, the instrument

imaged the entire observatory surface in 112 images over approximately 35 minutes. Lights, provided by

a ring of LEDs angled away from the field of view, were switched on at image capture and off

immediately afterwards. Images were saved in a .jpeg format and retrieved via exchange of an SD card

on site. The instruments ran on mains power except on a few occasions where they failed to start; these

missing data were gap filled as detailed later. Here we show data until 29 September 2022. This period

covered the main growing season, with a gradual progression from winter conditions (air temperature

min -7°C) through summer (air max. of 39 °C) and early autumn. Soil temperature was more buffered

and at 8 cm depth reached minima of 1.6 °C. Even at the beginning of this dataset (the first 6 weeks are

covered coarsely in Nair *et al.*, 2022), root growth could be observed in sequential images. We could

observe the disappearance of roots from images, indicating turnover or herbivory. Over the period of

data collection there were periods with clear artefacts in the images, such as condensation and soil

fauna, which we were mostly successfully able to train the Convolutional Neural Network (CNN) ignore

in segmentations.

Image Processing

Field minirhizotron imagery is still difficult to process automatically and while much progress has been

made in using neural networks for root images in general this is largely in simple agricultural or

mesocosm settings and designed on sparse timepoints (e.g. Gillert *et al.*, 2021; Smith *et al.*, 2022)) which

are effective for sparse timepoints but had potentially high noise between sequential images. We

smoothed the data as described later to offset this. Further development of sequential models could

improve the interpretation of dense timeseries.

We processed every image collected using the same workflow (Figure 2), processing 268972 images over

2480 individual sampling cycles over four instruments). The size of this dataset meant we had to limit

training data to a small fraction of the total image set, despite using a corrective annotation method,

rootpainter (Smith *et al.*, 2022) that allows a large throughput in training. Workflows based on

rootpainter have been successfully used to interpret root images in both mesocosm (Alonso-Crespo *et*

al., 2021) and field sites (Bauer *et al.*, 2021; Nair *et al.*, 2022). At this site, a cross-validated R^2 of 88 %

against was achieved image-level manual markup in the first six weeks of the experiment (Nair *et al.*,

2022). Our philosophy for the interpretation of automatic instruments is to maximise *overall* consistency

through time and at the scale of whole images than strict pixel-correct segmentation. Hence a declining

value on our graphs is not disappearance of individual roots, but declining segmented area across all

images.

We excluded images on the top or bottom plane of the minirhizotron and those immediately at the soil

surface, leaving 76 images per instrument per observation. We trained the model specifically on this site

using a dataset consisting of 25 randomly selected images per instrument-week of the experiment,

which were used to train in a random order. This dataset was fully generated after data collection was

complete (i.e., not the same model as we used in Nair *et al.*, (2022)) so there was not a bias in training

245 the model from images in the sequence they were collected nor bias towards earlier images in the

246 training. We trained for 350 random images at which point the f1 score indicated that the model was

247 not improving. We repeated the complete training on an independent random set of 350 images and

248 compared segmentations on 500 unseen images to validate that our training was reproducible (Figure

249 S1). Overall, the two segmentations had a correlation coefficient r = 0.96.

250 Following segmentation, we processed binary pixel maps of the images to extract basic trait data, mostly

251 in Ppython 3 using the skikit-image package. We followed the same protocol as (Nair *et al.*, 2022) with

252 the addition of an additional property, mean rooting depth (MRD). This was done as follows: first, we

253 removed non-contiguous segmented areas of less than 0.5 mm^2 to limit the influence of time-variable

254 segmentation artefacts. We skeletonized the remaining segmented area to a single pixel-wide line,

255 preserving original connectivity using medial axis segmentation and pruned all skeleton segments with a

256 length of less than 10 pixels. Short segments were predominantly artefacts of the segmentation

257 procedure and contributed little individually to the total root length. We counted all pixels in the

258 remaining skeleton; we consider this root length. It contains a small artefact due to not adjusting for

259 diagonal connections, but because roots are mostly linear in the images - unlike scanned root samples

260 from soil core extractions - we expected this effect to be minimal. RLD (mm^2 cm^{-2}) is root length / image

261 area. We calculated the *mean rooting* depth per instrument:timestep in R. As the minirhizotron

262 observations were not evenly distributed in the vertical profile we fit a loess regression to generate the

263 best fit smooth curve over the depth profile and predicted a RLD value at even 0.5 mm steps between 0

264 and 50 cm depth. We then proceeded through the predicted root length to find the depth at which 50%

265 of the total root length form the model had been reached, which we treated as MRD (cm).

Gap Filling

The image archive had a few gaps as on rare occasions the instruments did not start when scheduled. There were also instances when image artefacts occurred in the colour balance or brightness of the images due to data transfer from the camera, or power issues with the Light Emitted Diode (LED) lighting. We filtered out these occurrences automatically based on image colour statistics as these issues could affect the segmentation result. We also gap filled data points which were more than 25 % larger or smaller than the mean of the two observations before and after it; this accounted for short term anomalies such as soil animals which may obscure roots or be segmented as roots in a single image in a time series. Overall these data gaps affected 4% of images across the whole dataset. Having removed these data, we then gap filled the data for each trait via linear interpolation for a maximum gap size of 10 sequential observations. In practice, this meant an observation (real or gap filled) was available for every potential data point. Finally, in creating the RLD and MRD curves shown, we smoothed the data with a 3-day rolling average.

Meteorological and Remote Sensing Data

We used data from the MPI BGC Jena 'Saaleaue' meteorological station, located approximately 150 m from the root observations. The station provided us with air temperature, soil moisture (SM) content, and precipitation at the site level not paired to any instrument. For air temperature and moisture, we took the average of all sensors between 5 and 40 cm (the maximum depth of the minirhizotron). For precipitation, we averaged three rain gauges. The weather station data was available on 30 minute resolution, and further averaged to a daily averages to pair with daily averages growth rates from the minirhizotrons and calculated a further lagged 5 day (air temperature) and 14 day average (all meteorological variables except Rain and SM, which we left out because of their tendency to have strong effects on short timescales).

We lacked a dedicated paired above-ground canopy dynamics assessment paired to the minirhizotrons

instruments. To assess whether the canopy was growing, we used Sentinel 2 imagery of the Jena

Experiment site at 10 m spatial resolution, and computed Normalised Difference vegetation Index

(NDVI) product with a cut-out of the Jena Experiment site for gross patterns of biomass above ground in

our herbaceous system. There were 49 overpasses in the period of the experiment, of which 30 were

obscured by clouds. With these removed from the dataset, we fit a LOESS regression across the

remaining data for a smooth to produce a smooth curve that was used to interpret changes in the more

frequent belowground data (Figure S6).

Sub-daily Patterns

Minirhizotron data may be subject to variation on sub-daily timescales which are not growth patterns

but real sub-daily patterns of cell turgor (Huck *et al.*, 1970) or artefacts in neural network segmentation

due to root or soil appearance (Nair *et al.*, 2022). We wanted to discount these as much as possible to

examine actual sub-daily growth. To do this, we started on the raw data (without the smoothing

previously mentioned), observing that the biggest differences were between observations at midday

and midnight – midday tended to have slightly higher RLD than midnight (Figure S2). This cycle was

almost at inflection at 7 and 19 UTC without a significant difference in observed mean RLD between

instruments when grouped by calendar week (Supplementary Table S2.1)). After we found the

desynchronous behaviour, we re-ran this analysis, splitting the dataset into three groups: between the

periods before, during, and after the desynchrony, and found significant effects of hour only in this

period. Details in this post-hoc analysis are in Supplementary Table S2.2.

We hence treated the period between 7 and 19 UTC as 'day' and from 19 and to 7 as 'night'. We

smoothed the date across the time axis by applying a 3-days rolling average between day and night

periods and calculated daytime growth as the difference between the previous NIGHT and the current

DAY, and vice versa for nighttime growth.

To examine the trends in timing of growth over time we aggregated the DAY/NIGHT data arbitrarily into

calendar week bins, each containing 7 observations of DAY and 7 of night from each instrument. We

treated these as independent. We tested if day and night were significantly different for every

sequential day-night and night-day pair of image sets within each calendar week using sign tests, a non-

parametric method to test the median of two distributions.

ACKNOWLEDGEMENTS

The authors are grateful for the Jena Experiment for hosting this study.

We also acknowledge technical help in minirhizotron development and maintenance from Agnes

Fastnacht, Kerstin Hippler, Nadine Hempel and Tarek El-Madany.

AUTHOR CONTRIBUTION

RN: Richard Nair, MS1: Martin Strube, JPL: Javier Pacheco-Labrador, MS2: Marion Schrumpf, MM: Mirco Migliavacca

Conceptualization: RN, MM. Methodology: RN, MS1, MS2, JPL. Investigation: RN. Data Curation, RN,

Formal Analysis, RN, Software: RN. Writing – original draft: RN, Writing – review & editing: MS2, MM,

JPL

DATA AVAILABILITY

REFERENCES

Abramoff R, Finzi AC. **2015**. Are above- and below-ground phenology in sync? *New Phytologist* **205**: 1054–1061.

Allen MF, Kitajima K. **2013**. In Situ High-Frequency Observations of Mycorrhizas. *New Phytologist* **200**: 222–228.

Alonso-Crespo IM, Weidlich EWA, Temperton VM, Delory BM. **2021**. Assembly history modulates vertical root distribution in a grassland experiment. : 2021.08.24.457510.

Atkin OK, Edwards EJ, Loveys BR. **2000**. Response of root respiration to changes in temperature and its relevance to global warming. *New Phytologist* **147**: 141–154.

Badri DV, Loyola-Vargas VM, Broeckling CD, Vivanco JM. **2010**. Root secretion of phytochemicals in Arabidopsis is predominantly not influenced by diurnal rhythms. *Molecular Plant* **3**: 491–498.

Bahn M, Schmitt M, Siegwolf R, Richter A, Brüggemann N. **2009**. Does photosynthesis affect grassland
soil-respired CO2 and its carbon isotope composition on a diurnal timescale? *New Phytologist* **182**: 451–
460.

Baldocchi DD. **2020**. How eddy covariance flux measurements have contributed to our understanding of
Global Change Biology. *Global Change Biology* **26**: 242–260.

Bauer FM, Lärm L, Morandage S, Lobet G, Vanderborght J, Vereecken H, Schnepf A. **2021**. *Combining*
deep learning and automated feature extraction to analyze minirhizotron images: development and
validation of a new pipeline.

Belaud E, Jourdan C, Barry-Etienne D, Marsden C, Robin A, Taschen E, Hedde M. **2024**. In situ soil
imaging, a tool for monitoring the hourly to monthly temporal dynamics of soil biota. *Biology and*
Fertility of Soils **60**: 1055–1071.

Bengough AG, Bransby MF, Hans J, McKenna SJ, Roberts TJ, Valentine TA. **2006**. Root responses to soil
physical conditions; growth dynamics from field to cell. *Journal of Experimental Botany* **57**: 437–447.

Bengough AG, McKenzie BM, Hallett PD, Valentine TA. **2011**. Root elongation, water stress, and
mechanical impedance: a review of limiting stresses and beneficial root tip traits. *Journal of*
Experimental Botany **62**: 59–68.

Boone RD, Boone RD, Canary JD, Canary JD. **1998**. Roots Exert a Strong Influence on the Temperature
Sensitivity of Soil Respiration. *Nature* **396**: 570–572.

Box JE, Smucker AJM, Ritchie JT. **1989**. Minirhizotron Installation Techniques for Investigating Root
Responses to Drought and Oxygen Stresses. *Soil Science Society of America Journal* **53**: 115–118.

Brundrett M. **1991**. Mycorrhizas in Natural Ecosystems. In: Begon M, Fitter AH, Macfadyen A, eds.
Advances in Ecological Research. Academic Press, 171–313.

Caldeira CF, Jeanguenin L, Chaumont F, Tardieu F. **2014**. Circadian rhythms of hydraulic conductance
and growth are enhanced by drought and improve plant performance. *Nature Communications* **5**: 5365.

Ceolin S, Schymanski SJ, van Dusschoten D, Koller R, Klaus J. **2025**. Root growth dynamics and
allocation as a response to rapid and local changes in soil moisture. *Biogeosciences* **22**: 691–703.

De Vries FWTP, Brunsting AHM, Van Laar HH. **1974**. Products, requirements and efficiency of
biosynthesis a quantitative approach. *Journal of Theoretical Biology* **45**: 339–377.

Defrenne CE, Childs J, Fernandez CW, Taggart M, Nettles WR, Allen MF, Hanson PJ, Iversen CM. **2021**.
High-resolution minirhizotrons advance our understanding of root-fungal dynamics in an experimentally
warmed peatland. *Plants, People, Planet* **3**: 640–652.

Edwards EJ, D.G. B, A. ML, H. FA. **2004**. Root Production Is Determined by Radiation Flux in a Temperate
Grassland Community. *Global Change Biology* **10**: 209–227.

Freschet GT, Pagès L, Iversen CM, Comas LH, Rewald B, Roumet C, Klimešová J, Zadworny M, Poorter
H, Postma JA, *et al.* **2021**. A starting guide to root ecology: strengthening ecological concepts and

standardising root classification, sampling, processing and trait measurements. *New Phytologist* **232**:
973–1122.

Gherardi LA, Sala OE. **2020**. Global patterns and climatic controls of belowground net carbon fixation.
Proceedings of the National Academy of Sciences **117**: 20038–20043.

Gillert A, Peters B, Freiherr von Lukas, U, Kreyling, J. **2021**. Identification and Measurement of
Individual Roots in Minirhizotron Images of Dense Root Systems. In: Proceedings of the IEEE/CVF
International Conference on Computer Vision (ICCV) Workshops. 1323–1331.

Graham EA, Mulkey SS, Kitajima K, Phillips NG, Wright SJ. **2003**. Cloud cover limits net CO2 uptake and
growth of a rainforest tree during tropical rainy seasons. *Proceedings of the National Academy of*
Sciences **100**: 572–576.

Halter R, Sands R, Nambiar EKS, Ashton DH. **1996**. Elongation of Eucalyptus roots during day and night.
Tree Physiology **16**: 877–881.

Hatfield JL, Prueger JH. **2015**. Temperature extremes: Effect on plant growth and development.
Weather and Climate Extremes **10**: 4–10.

Hawkins H-J, Cargill RIM, Van Nuland ME, Hagen SC, Field KJ, Sheldrake M, Soudzilovskaia NA, Kiers
ET. **2023**. Mycorrhizal mycelium as a global carbon pool. *Current Biology* **33**: R560–R573.

Head GC. **1965**. Studies of Diurnal Changes in Cherry Root Growth and Nutational Movements of Apple
Root Tips by Time-lapse Cinematography. *Annals of Botany* **29**: 219–224.

Heinze J, Werger L, Ogden M, Heinken T, Hoefgen R, Weber E. **2023**. Short wind pulses consistently
change the morphology of roots, but not of shoots, across young plants of different growth forms. *Stress*
Biology **3**: 43.

Hirano T, Cui R, Sun L, Teramoto M, Liang N. **2023**. Partitioning of root respiration into growth,
maintenance, and ion uptake components in a young larch-dominated forest. *Plant and Soil* **482**: 57–72.

Högberg P, Bhupinderpal-Singh, Löfvenius MO, Nordgren A. **2009**. Partitioning of soil respiration into its
autotrophic and heterotrophic components by means of tree-girdling in old boreal spruce forest. *Forest*
Ecology and Management **257**: 1764–1767.

Huck MG, Klepper B, Taylor HM. **1970**. Diurnal Variations in Root Diameter. *Plant Physiology* **45**: 529–
530.

Huete AR, Didan K, Shimabukuro YE, Ratana P, Saleska SR, Hutyra LR, Yang W, Nemani RR, Myneni R.
2006. Amazon rainforests green-up with sunlight in dry season. *Geophysical Research Letters* **33**.

Jackson RB, Canadell J, Ehleringer JR, Mooney HA, Sala OE, Schulze ED. **1996**. A Global Analysis of Root
Distributions for Terrestrial Biomes. *Oecologia* **108**: 389–411.

Jacob V, Choat B, Churchill AC, Zhang H, Barton CVM, Krishnananthaselvan A, Post AK, Power SA,
Medlyn BE, Tissue DT. **2022**. High safety margins to drought-induced hydraulic failure found in five
pasture grasses. *Plant, Cell & Environment* **45**: 1631–1646.

Jensen KH, Berg-Sørensen K, Bruus H, Holbrook NM, Liesche J, Schulz A, Zwieniecki MA, Bohr T. 2016.
Sap flow and sugar transport in plants. *Reviews of Modern Physics* **88**: 035007.

Jian J, Frissell M, Hao D, Tang X, Berryman E, Bond-Lamberty B. 2022. The global contribution of roots
to total soil respiration. *Global Ecology and Biogeography* **31**: 685–699.

Johnson IR. 1990. Plant respiration in relation to growth, maintenance, ion uptake and nitrogen
assimilation. *Plant, Cell & Environment* **13**: 319–328.

Johnson MG, Tingey DT, Phillips DL, Storm MJ. 2001. Advancing fine root research with minirhizotrons.
Environmental and Experimental Botany **45**: 263–289.

King AW, Gunderson CA, Post WM, Weston DJ, Wullschleger SD. 2006. Plant Respiration in a Warmer
World. *Science* **312**: 536–537.

Körner C. 2008. Winter crop growth at low temperature may hold the answer for alpine treeline
formation. *Plant Ecology & Diversity* **1**: 3–11.

Körner C. 2015. Paradigm shift in plant growth control. *Current Opinion in Plant Biology* **25**: 107–114.

Kuwabe N, and Ohashi M. 2023. Evaluation of spatial and temporal variation in fine root dynamics in a
temperate mixed forest using a scanner method. *Journal of Forest Research* **28**: 186–193.

Liu Y, Wu C. 2020. Understanding the role of phenology and summer physiology in controlling net
ecosystem production: a multiscale comparison of satellite, PhenoCam and eddy covariance data.
Environmental Research Letters **15**: 104086.

Liu M, Xu X, Nannipieri P, Kuzyakov Y, Gunina A. 2021. Diurnal dynamics can modify plant–microbial
competition for N uptake via C allocation. *Biology and Fertility of Soils* **57**: 949–958.

Luo Y, El-Madany T, Ma X, Nair R, Jung M, Weber U, Filippa G, Bucher SF, Moreno G, Cremonese E, *et
***al*. 2020**. Nutrients and water availability constrain the seasonality of vegetation activity in a
Mediterranean ecosystem. *Global Change Biology* **26**: 4379–4400.

Majerus ME. 1975. Response of Root and Shoot Growth of Three Grass Species to Decreases in Soil
Water Potential. *Journal of Range Management* **28**: 473.

McCormack ML, Dickie IA, Eissenstat DM, Fahey TJ, Fernandez CW, Guo D, Helmisaari HS, Hobbie EA,
Iversen CM, Jackson RB, *et al*. 2015. Redefining Fine Roots Improves Understanding of Below-Ground
Contributions to Terrestrial Biosphere Processes. *New Phytologist* **207**: 505–518.

Mencuccini M, Hölttä T. 2010. The significance of phloem transport for the speed with which canopy
photosynthesis and belowground respiration are linked. *New Phytologist* **185**: 189–203.

Meyer N, Welp G, Amelung W. 2018. The Temperature Sensitivity (Q10) of Soil Respiration: Controlling
Factors and Spatial Prediction at Regional Scale Based on Environmental Soil Classes. *Global*
Biogeochemical Cycles **32**: 306–323.

Morris K, Nair R. **2025**. Below the leaves: Integrating above- and below-ground phenology for
earth-system predictability. *Functional Ecology*: 1365-2435.70057.

Münch E. **1930**. *Die Stoffbewegungen in der Pflanze*. Gustav Fischer.

Nair R, Strube M, Hertel M, Kolle O, Rolo V, Migliavacca M. **2022**. *HIGH FREQUENCY ROOT DYNAMICS:*
SAMPLING AND INTERPRETATION USING REPLICATED ROBOTIC MINIRHIZOTRONS. Ecology.

Ostle N, Briones MJI, Ineson P, Cole L, Staddon P, Sleep D. **2007**. Isotopic detection of recent
photosynthate carbon flow into grassland rhizosphere fauna. *Soil Biology and Biochemistry* **39**: 768–777.

Pantin F, Simonneau T, Rolland G, Dauzat M, Muller B. **2011**. Control of Leaf Expansion: A
Developmental Switch from Metabolics to Hydraulics. *Plant Physiology* **156**: 803–815.

Peacock JM. **1975**. Temperature and Leaf Growth in Lolium perenne. III. Factors Affecting Seasonal
Differences. *Journal of Applied Ecology* **12**: 685–697.

Poiré R, Wiese-Klinkenberg A, Parent B, Mielewczik M, Schurr U, Tardieu F, Walter A. **2010**. Diel time-
courses of leaf growth in monocot and dicot species: endogenous rhythms and temperature effects.
Journal of Experimental Botany **61**: 1751–1759.

Raich JW, Schlesinger WH. **1992**. The global carbon dioxide flux in soil respiration and its relationship to
vegetation and climate. *Tellus B: Chemical and Physical Meteorology* **44**: 81–99.

Reichstein M, Falge E, Baldocchi D, Papale D, Aubinet M, Berbigier P, Bernhofer C, Buchmann N,
Gilmanov T, Granier A, et al. **2005**. On the separation of net ecosystem exchange into assimilation and
ecosystem respiration: review and improved algorithm. *Global Change Biology* **11**: 1424–1439.

Roscher C, Schumacher J, Baade J, Wilcke W, Gleixner G, Weisser WW, Schmid B, Schulze E-D. **2004**.
The role of biodiversity for element cycling and trophic interactions: an experimental approach in a
grassland community. *Basic and Applied Ecology* **5**: 107–121.

Rozema J, Arp W, Digglen JV, Kok E, Letschert J. **1987**. An Ecophysiological Comparison of
Measurements of the Diurnal Rhythm of the Leaf Elongation and Changes of the Leaf Thickness of Salt-
resistant Dicotyledonae and Monocotyledonae. *Journal of Experimental Botany* **38**: 442–453.

Ruts T, Matsubara S, Wiese-Klinkenberg A, Walter A. **2012**. Diel patterns of leaf and root growth:
endogenous rhythmicity or environmental response? *Journal of Experimental Botany* **63**: 3339–3351.

Savage KE, Davidson EA, Abramoff RZ, Finzi AC, Giasson M-A. **2018**. Partitioning soil respiration:
quantifying the artifacts of the trenching method. *Biogeochemistry* **140**: 53–63.

Smith AG, Han E, Petersen J, Olsen NAF, Giese C, Athmann M, Dresbøll DB, Thorup-Kristensen K. **2022**.
RootPainter: deep learning segmentation of biological images with corrective annotation. *New*
Phytologist **236**: 774–791.

Stahl Y, Simon R. **2010**. Plant primary meristems: shared functions and regulatory mechanisms. *Current*
Opinion in Plant Biology **13**: 53–58.

Steinaker DF, Wilson SD. **2008**. Phenology of Fine Roots and Leaves in Forest and Grassland. *Journal of*
Ecology **96**: 1222–1229.

Strand AE, Pritchard SG, Mccormack ML, Davis MAMA, Strand AE, Pritchard SG, Mccormack ML, Davis
MAMA, Oren R. **2018**. Irreconcilable Differences⬚: Fine-Root Life Spans and Soil Carbon Persistance.
Science **319**: 456–458.

Sun L, Hirano T, Yazaki T, Teramoto M, Liang N. **2019**. Fine root dynamics and partitioning of root
respiration into growth and maintenance components in cool temperate deciduous and evergreen
forests. *Plant and Soil*.

Szaniawski RK, Kielkiewicz M. **1982**. Maintenance and growth respiration in shoots and roots of
sunflower plants grown at different root temperatures. *Physiologia Plantarum* **54**: 500–504.

Taylor AJ, Lai CT, Hopkins FM, Wharton S, Bible K, Xu X, Phillips C, Bush S, Ehleringer JR. **2015**.
Radiocarbon-Based Partitioning of Soil Respiration in an Old-Growth Coniferous Forest. *Ecosystems* **18**:
459–470.

Thurgood A, Singh B, Jones E, Barbour MM. **2014**. Temperature sensitivity of soil and root respiration in
contrasting soils. *Plant and Soil* **382**: 253–267.

Walter A, Silk WK, Schurr U. **2009**. Environmental Effects on Spatial and Temporal Patterns of Leaf and
Root Growth. *Annual Review of Plant Biology* **60**: 279–304.

Walter A, Spies H, Terjung S, Küsters R, Kirchgeßner N, Schurr U. **2002**. Spatio-temporal dynamics of
expansion growth in roots: automatic quantification of diurnal course and temperature response by
digital image sequence processing. *Journal of Experimental Botany* **53**: 689–698.

Weisser WW, Roscher C, Meyer ST, Ebeling A, Luo G, Allan E, Beßler H, Barnard RL, Buchmann N,
Buscot F, *et al.* 2017. Biodiversity effects on ecosystem functioning in a 15-year grassland experiment:
Patterns, mechanisms, and open questions. *Basic and Applied Ecology* **23**: 1–73.

Wingler A, Hennessy D. **2016**. Limitation of Grassland Productivity by Low Temperature and Seasonality
of Growth. *Frontiers in Plant Science* **7**.

Wu Z, Chen S, De Boeck HJ, Stenseth NC, Tang J, Vitasse Y, Wang S, Zohner C, Fu YH. **2021**.
Atmospheric brightening counteracts warming-induced delays in autumn phenology of temperate trees
in Europe. *Global Ecology and Biogeography* **30**: 2477–2487.

Wu C, Gonsamo A, Chen JM, Kurz WA, Price DT, Lafleur PM, Jassal RS, Dragoni D, Bohrer G, Gough CM,
***et al.* 2012**. Interannual and spatial impacts of phenological transitions, growing season length, and
spring and autumn temperatures on carbon sequestration: A North America flux data synthesis. *Global*
and Planetary Change **92–93**: 179–190.

Yazdanbakhsh N, Fisahn J. **2009**. High Throughput Phenotyping of Root Growth Dynamics, Lateral Root
Formation, Root Architecture and Root Hair Development Enabled by PlaRoM. *Functional Plant Biology*
36: 938–946.

Yazdanbakhsh N, Sulpice R, Graf A, Stitt M, Fisahn J. **2011**. Circadian control of root elongation and C
partitioning in Arabidopsis thaliana. *Plant, Cell & Environment* **34**: 877–894.

Zelikova TJ, Williams DG, Hoenigman R, Blumenthal DM, Morgan JA, Pendall E. **2015**. Seasonality of
soil moisture mediates responses of ecosystem phenology to elevated CO2 and warming in a semi-arid
grassland. *Journal of Ecology* **103**: 1119–1130.

Zweifel R, Sterck F, Braun S, Buchmann N, Eugster W, Gessler A, Häni M, Peters RL, Walthert L,
Wilhelm M, *et al.* 2021. Why trees grow at night. *New Phytologist* **231**: 2174–2185.

Figure Captions

Figure 1. Root dynamics during the observation period. a) Mean root length density (RLD) dynamics between two instruments with two added in July (three day rolling mean, shapes indicate different instruments). Shaded area indicates the minimum and maximum observation within each five-day window. b) Mean rooting depth (MRD), the mean depth of roots observed in the minirhizotron. c) Sub-daily RLD growth. Throughout the observations, roots both grew (+ve growth) and turned over (-ve growth). Stars indicate periods with significantly different night-time growth than day-time growth (sign test). d) soil moisture as a % at measured 8 cm. e) rain, soil temperature (blue), and air temperature (red). The grey lines in the top three figures indicate (dashed) installation of new instruments, and (solid) first harvest and second mowing of above ground biomass removal in the management cycle.

Figure 2. Summary of our workflow. Black boxes indicate data, grey boxes automated processing, and blue boxes automated processing where parameter choice was meaningful.