Rhythmic Nuclear Import Mediated by Importins Regulates the *Neurospora*

Circadian Clock.

Ziyan Wang[a], Bin Wang[a], Bradley M. Bartholomai[a], Jennifer J. Loros[b], Jay C. Dunlap[a+]

[a] Department of Molecular and Systems Biology, [b] Department of Biochemistry and Cell Biology,

Geisel School of Medicine at Dartmouth, Hanover, NH, USA

[+]Corresponding author: J. C. Dunlap, jay.c.dunlap@dartmouth.edu

ABSTRACT

Circadian clocks in eukaryotes rely on precisely regulated negative feedback loops to generate
daily rhythms. However, the delay mechanisms that extend this structurally simple feedback
loop to ~24 hours are not yet fully understood. In the filamentous fungal model organism
Neurospora crassa, the negative arm complex, centered by FREQUENCY (FRQ), must enter
the nucleus to repress the White Collar Complex (WCC) and close the feedback loop, but the
mechanisms and dynamics of its nuclear transport have remained unresolved. Using long-term
live-cell imaging and fluorescence recovery after photobleaching (FRAP), we demonstrate that
FRQ nuclear import is an active circadian-regulated process that is fastest early in the
subjective day and progressively decreases as nuclear FRQ approaches peak levels,
corresponding to altered direct binding between FRQ and Importin α. We further establish that
Importin α is required for the spatial regulation of FRQ and WC-1 and the correct timing of
Neurospora circadian clock. Analysis of the three *Neurospora* Importin β homologs reveals that
each of them contributes differently to the circadian clock through pathways beyond FRQ or
WCC nuclear import. More specifically, we find a genetic interaction between Impβ3 and the
phosphatase PP-4. Together, these findings indicate that nuclear import is a selective, dynamic,
and rate-limiting regulatory step in the fungal circadian clock and reveal both conserved and
fungal-specific mechanisms by which importins tune circadian timing.

INTRODUCTION

Circadian clocks are self-sustained timekeeping systems that allow organisms to anticipate and adapt to daily changes in the environment. These oscillators are based on conserved transcription-translation feedback loops (TTFLs) in which the Positive Arm (typically transcription factors) drives expression of the Negative Arm that, once translated and imported into nuclei, inhibit its own activators (Dunlap, 1999; Hardin, 2011; Partch et al., 2014). A key feature of this design is that multiple built-in delays extend the simple feedback loop to maintain a ~24-hour periodicity that approximates environmental changes.

Precisely timed nuclear entry of Negative Arm components can act as a rate-limiting step that introduces phase delay between transcriptional activation and repression. Importins (karyopherins) mediate the nucleocytoplasmic transport of most proteins through the nuclear pore complex (NPC) (Chook and Blobel 2001; Pemberton and Paschal 2005). There are two major nuclear import pathways: the classical Importin α (Impα)-dependent pathway, in which Impα, as an adaptor, binds cargos with canonical nuclear localization signals (NLSs) and delivers them to Importin β (Impβ); and the Impα-independent pathway, in which Impβ directly recognizes non-classical NLSs. In *Drosophila*, the clock protein dTIM binds directly to Impα1 and co-transports dPER, and perturbations of Impα1 alter circadian period and rhythmicity (Jang et al. 2015; Lin et al. 2023). In mammals, the extent to which nuclear import of PERs and CRYs requires Impα remains debatable and may rely more directly on Impβ family members. In human cells, KPNB1 (Impβ1) mediates nuclear import of some PER/CRY complexes independently of Impα (KPNA1/2), and TNPO1 (Impβ2) selectively regulates PER1 localization and period length (Lee et al. 2015; Korge et al. 2018; Lee et al. 2019). However, chemical inhibition of the classical Impα/β1 pathway also alters period length (Öllinger et al. 2014), and in mouse, nuclear import of mCRY2 is Impα-dependent (Sakakida et al. 2005). These findings suggest that controlled nuclear entry of the Negative Arm is a conserved step important for circadian timing, but the exact pathways and molecular players differ: Impα is essential in *Drosophila*, while in mammals PER and CRY can sometimes bypass Impα and depend on specific Impβs. In contrast to animal circadian systems, little is known about how the Positive and Negative Arm components of fungal clocks such as *Neurospora* enter the nucleus.

The filamentous fungus *Neurospora crassa* has been a powerful model for studying circadian clocks. The Positive Arm of its oscillator is the White Collar Complex (WCC, a heterodimer of WC-1 and WC-2), which activates transcription of the *frequency* (*frq* gene) whose product serves to nucleate the Negative Arm complex of FRQ, FRH, and CK1. This complex then feeds back to repress WCC activity (Crosthwaite et al. 1997; Dunlap 1999; Dunlap and Loros 2017; Costa et al. 2025). FRQ is highly concentrated in the nucleus, which makes sense because its nuclear entry is required for inhibitory function on WCC and is essential for closing the negative feedback loop (Wang et al. 2025). Thus, FRQ nuclear transport is a central regulatory point in the *Neurospora* clock.

Despite its importance, the mechanism of FRQ nuclear transport remains unclear. An NLS within FRQ was reported more than two decades ago and suggested to be essential for both nuclear localization and clock function (Luo et al. 1998), but this has not been thoroughly tested with modern tools, and the transport machinery has also not been defined. Unlike animals,

which express three classes of Importin α subunits (e.g. six in human genome) with overlapping
and specialized functions, plants and fungi only have one class of Importin α that resembles
animal α1(Goldfarb et al. 2004; Mason et al. 2009; Bernardes et al. 2020). Within the fungal
kingdom, nuclear import pathways have only been systematically studied in yeast and
Aspergillus nidulans. A few related studies in *Neurospora* focused on the conserved structure
and epigenetic function of Impα (Takeda et al. 2013; Bernardes et al. 2015; Klocko et al. 2015).
Through identification of homologs using NCBI orthologs based on the Karyopherin-β family in
Saccharomyces cerevisiae and *Aspergillus nidulans* (Ström and Weis 2001; Mans et al. 2004;
Markina-Iñarrairaegui et al. 2011), we identified a single Impα, three Impβs and nine other
potential Karyopherins including Exportin-1 in *Neurospora crassa* (Supplementary Table 1).
Mass spectrometry studies suggested FRQ may weakly interact with Impα and/or Impβ3 (Baker
2010; Pelham et al. 2023), but biochemical, functional and biological validation is lacking. How
these Importins contribute to FRQ nuclear entry or to subcellular localization of other clock
proteins is unknown. In part this is due to technical challenges which have further limited
progress. The insufficient brightness of FRQ-GFP fusion protein prevented reliable imaging, and
an "anchor-away" assay (Haruki et al. 2008), which used FRQ fused to FRB artificially tethered
to FKBP-tagged histone H1 to assay nuclear import, introduced non-physiological import routes
and disrupted normal clock function (Diernfellner et al., 2009). As a result, the dynamics and
regulation of FRQ nuclear transport in *Neurospora* remain poorly characterized.

In this study, we used live-cell imaging combined with FRAP to directly measure FRQ nuclear
import in *Neurospora*. We found that FRQ is imported slowly, requiring tens of minutes for
recovery, and that import is strongly regulated across the circadian cycle, being highest early in
the subjective day and reduced near nuclear peak levels. These dynamics are comparable to
those of mammalian PER proteins, suggesting that slow nuclear import is a conserved delay
mechanism in circadian clocks. We further show that FRQ import requires importin α but not the
previously proposed NLS, and that import of WC-1, but not WC-2, also depends on importin α.
Finally, we find that the three importin βs have distinct contributions to circadian regulation,
including a potential link between Impβ3 and PPH-4. Together, these results reveal nuclear
import as a selective and regulated process that is essential for circadian timing in *Neurospora*.

RESULTS

Nuclear import rate of FRQ is circadianly regulated.

Precisely timed nuclear localization of FRQ is essential for the normal operation of the circadian clock. To understand how and when FRQ enters nuclei, we optimized FRAP experiments for *Neurospora* building on the continuous live-cell imaging approach reported separately (Wang et al. 2025). An imaging strain blind to blue light carrying endogenously mNeonGreen-tagged FRQ (strain 1952-2; *his-3::frq$_{cbox}$-luc; ras-1^{bd}; wc-1$^{\Delta LOV}$, for, frq$^{mNeonGreen\_Flag}$*) retains an intact circadian clock (Fig. S1A-C). The application of microfluidic devices allows convenient medium exchange, enabling application of the microtubule inhibitor benomyl at specific times. Benomyl slows nuclear movement by disrupting active transport along microtubules and by reducing growth, which decreases passive movement driven by bulk cytoplasmic flow and thereby facilitating imaging. Importantly, benomyl does not affect circadian functions (Fig. S1D). For FRAP experiments, a thick optical section (around 2 µm, corresponding to the diameter of *Neurospora* nuclei) was chosen to minimize artifacts from vertical random movement of nuclei. Cultures in microfluidic devices (Wang et al. 2025) were entrained with 12h:12h temperature cycles mimicking natural conditions, and FRAP was then performed every 30 minutes during the subjective day on the same culture (Fig.1A). After bleaching entire target nuclei, the subsequent recovery of fluorescence reflects import of unbleached cytoplasmic FRQ^{mNG} at that time.

The nuclear fluorescence from FRQ^{mNeonGreen} recovered most rapidly early in the subjective day, indicating a higher import rate, while the recovery was slower later in the subjective day (Fig. 1B-C, Supplementary Video 1). This temporal difference is consistent with the progressive nuclear accumulation of FRQ during the subjective day that we observed previously (Wang et al. 2025). Near subjective dawn, the recovered signal could even exceed the pre-bleaching level (e.g. 129 ± 29% at 23 minutes post-bleaching at CT3). By contrast, at subjective dusk, when nuclear FRQ levels reach their peak, nuclear import was remarkably reduced (Fig. 1C-D).

Due to the slow FRQ import rate we observed, most recovery curves failed to reach a stable plateau within the 25-minute observation window. To ensure consistent analyses, we used linear regression rather than the commonly used exponential fit to calculate nuclear import rates. To compensate for high variability in nuclear movements and possible artifacts, multiple control nuclei within each culture were averaged and used to correct for unavoidable photobleaching caused by continuously imaging. The nuclear import rate shows a significant negative correlation with time during the subjective day (Fig. 1D, Fig. S1E). Consistent results were observed when using a single corresponding nucleus from the same experiment, without averaging (Fig. S1F). In conclusion, nuclear import rates of FRQ are circadian regulated with a peak in the subjective morning.

FRQ is transported into nuclei through the Impα import pathway.

Because previous mass spectrometry had identified Impα as a possible weak interactor of FRQ (Baker et al. 2009; Baker 2010), we hypothesized that the time-dependent regulation of FRQ nuclear import is mediated by its interaction with importin α (Impα). To verify this interaction, we endogenously tagged Impα with the V5 epitope tag at its C-terminus. Co-immunoprecipitation with V5 beads successfully enriched Impα^{V5}, and FRQ co-precipitated with Impα^{V5} at early time

points, corresponding to times showing rapid nuclear import (Fig. 2A, Fig, S2A-B). However, at
later time points when FRQ is more abundant, binding between FRQ and Impα^{V5} is barely
detectable, corresponding to times of slower import. The change in binding strength is
consistent with this being the cause for precisely controlled nuclear import.

Impα is expected to act through nuclear localization signals (NLS) (Conti et al. 1998; Takeda et
al. 2013). Several putative NLS have been suggested on FRQ, with one of them being essential
to circadian clock (Luo et al. 1998). We reconstituted Neurospora strains bearing FRQ at its
endogenous locus with the FRQ-NLS deleted, with or without the mNeonGreen tag to allow for
fluorescent imaging, to directly assess the contribution of the predicted NLS to FRQ nuclear
entry. The reconstituted strains recapitulate the arrhythmic phenotype as expected (Fig. S2C).
To delineate nuclear boundaries, two nuclear envelope proteins, Nup159 (NCU08604) in the
cytoplasmic side of nuclear pore complexes and Nup211 (NCU04059) in the nuclear basket
complexes, were tagged at their endogenous loci using different fluorescent tags to serve as
markers for the outer and inner sides of the nuclear membrane, respectively (Beck and Hurt
2017; Bartholomai 2022). Both proteins form fragmented ring-like structures that largely align
with each other at the nuclear periphery, with the two layers clearly distinguishable when
imaged on SoRa super-resolution microscope (Fig. S2D). To our surprise, although rhythmicity
was plainly diminished (Fig. S2C), FRQ still localized to nuclei and nuclear FRQ levels remained
constantly high when the proposed NLS is absent (Fig. 2B-C, Fig, S2E). This nuclear
localization is not an artifact caused by mNeonGreen tagging, as other tagged *Neurospora*
proteins, like the septin CDC-11, localized to cytoplasm as expected (Fig. S2F). Nuclear FRQ
still forms highly dynamic nuclear bodies in the *frq^{ΔNLS}* strain, with subnuclear heterogeneity
levels indistinguishable from that of wildtype FRQ (Fig. S2G, Supplementary Video 2)

Impα plays an important role in *Neurospora* development and growth (Klocko et al. 2015) and
complete depletion of Impα is lethal. To assess the functional role of Impα, we used an RNAi
approach to knock down *impa* under the control of a quinic acid (QA)-inducible promoter (Giles
et al. 1985; Shi et al. 2010) (Fig. 3A). Addition of QA leads to severe growth defects (Fig. 3B),
and short induction (20 hours) is sufficient to significantly reduce both *impa* transcript and Impα
protein levels, while not causing macroscopical sickness or death (Fig. 3C-D). Due to leakiness
of the *qa-2* promoter (Larrondo et al. 2009), some reduction of Impα was also observed in the
inducible RNAi strains even without induction (Fig. 3C-D, Fig, S3A). The progressive decrease
in Impα levels, from the no-dsRNA control strain (*impa^{V5}*) to inducible RNAi strains without QA,
and finally to the inducible RNAi strains with QA, provided a platform to assay dose-dependent
circadian phenotypes caused by Impα deficiency.

We examined circadian rhythms in Impα knockdown strains using the luciferase reporter driven
by the *frq$_{cbox}$* promoter described above. Control cultures exhibited ~22 h rhythms, whereas
progressive depletion of Impα shortened the period and stronger knockdown abolished
rhythmicity, resulting in arrhythmic or inviable cultures (Fig. 3E, Fig. S3B). Indeed, cultures
displayed sensitivity to extremely low levels of inhibition, comparing reductions seen in Fig. 3C-
D with rhythm phenotypes in Fig. 3E. Another striking effect caused by Impα deficiency is that
all tested core clock proteins, as well as non-clock-related protein Tubulin, decrease in bulk
tissues collected from liquid cultures (Fig. 3D). In contrast, the most abundant proteins that

dominate unspecific protein measurements remain unchanged. This may reflect a reduction in
metabolically active tissue due to sickness in Impα-deficient culture, while the structural
components in older tissue remain largely intact.

Live-cell fluorescent imaging allows dissection of local changes in metabolically active tissue,
such as the actively growing tip regions. These results are consistent with biochemical
analyses, showing that with Impα knocked down, FRQ nuclear import was clearly impaired (Fig.
4A). In the control strains, addition of QA (which also provides an extra nutrient source) boosts
nuclear FRQ levels (Fig. 4A-C, Fig. S3C). In contrast, in the RNAi strains, QA addition caused
reduced or unchanged nuclear FRQ, compared to the increase in the control strain, despite total
FRQ levels in the tip region remaining relatively constant (Fig. 4A-C, Fig. S3C). This indicates
that without sufficient Impα, FRQ accumulates in the cytoplasm and fails to efficiently enter
nuclei to shut down its own synthesis. Even in the absence of QA, the leaky knockdown of Impα
resulted in a mild reduction in nuclear FRQ compared to controls (Fig. 2B). Together, these
observations support the model that FRQ nuclear import is actively mediated by the Impα
pathway and is regulated across the circadian cycle to ensure appropriate timing and rates of
nuclear entry, both of which are essential for maintaining circadian stability.

WC-1 but not WC-2 also depends on Impα for nuclear import.

The most obvious change in clock protein levels is that WC-1 becomes undetectable by western
blot with severe deficiency of Impα (Fig. 3D), even though the expression level of *wc-1*
transcripts is unchanged (Fig. 3C). This divergence indicates that the defect occurs at the post-
transcriptional level, either through impaired translation or reduced WC-1 protein stability.
Interestingly, WC-1 co-immunoprecipitates with Impα in a phase-dependent manner (Fig. 2A),
similar to FRQ. Because WC-1 is predominantly found in the nucleus, this suggests a model in
which WC-1 is unstable in the cytoplasm. When nuclear import of WC-1 is blocked by deficiency
of Impα, newly synthesized WC-1 is trapped in cytoplasm, being unstable, and is rapidly
degraded (Fig. 4D).

Although rhythmic turnover of WCC is not essential for circadian clock regulation, the presence
of nuclear WC-1 is essential. While WC-1 is below the detection limit of western blotting in Impα
knockdown strains (Fig. S3A), a trace amount of WC-1 may still be able to support circadian
function (Lee et al. 2003; Káldi et al. 2006). Consistent with this, the expression of its direct
target *frq* is unaffected (Fig. 3C). The photoreceptor function of WC-1 is intact as well; both *wc-1*
and *frq* transcript levels increase drastically after a 15-minute light pulse (Fig. 4E-F).
Surprisingly, *impα* itself also turns out to be light-inducible (Fig. 4G), although it was not
previously identified as a light responsive gene by RNA-seq (Chen et al. 2009; Smith et al.
2010; Dekhang et al. 2017). Even though the residual WC-1 retains functions when Impα is
reduced, its overall efficiency is compromised, as the fold increase of gene expression in
response to light is smaller compared to no QA (Fig. S3D). In this context, it may be that less
FRQ is needed to fully inhibit the limited WCC activity during circadian cycles, which may
explain the observed period shortening despite slow FRQ nuclear import.

WC-2 also co-precipitated with Impα in a phase-dependent manner (Fig. 2A). However, both
nuclear and total levels of WC-2 in tip regions increase significantly as the level of Impα

decreases (Fig. 4A, H-I, Fig. S3E). Overall, the nuclear import of WC-2 does not appear to rely
on the Impα pathway. The observed co-immunoprecipitation is likely mediated through WC-2's
association with WC-1, and may not reflect a direct or biologically relevant interaction.

Deficiency of importin βs have divergent effects on circadian clock.

The genes encoding the three identified Impβ subunits in *Neurospora crassa* are NCU02011,
NCU03690, and NCU02357 (Supplementary Table 1) (Galagan et al. 2003). Homologs of two of
the three Impβs, Impβ1 (NCU02011; mammalian KPNB1) and Impβ2 (NCU03690; mammalian
TNPO1), have been associated with nuclear import of Negative Arm components PER1 and
PER2 in mammalian systems (Lee et al. 2015; Korge et al. 2018; Lee et al. 2019). Impβ3
(NCU02357; mammalian IPO5) has not been linked to circadian regulation in any system, but
was identified as a potential FRQ interactor at DD24 by mass spectrometry (Pelham et al.
2023). We therefore investigated how deficiency of these three Impβs affect circadian clock
using different strategies.

impβ1 is an essential gene - neither homozygous knockouts nor inducible *impβ1* strains (with
impβ1 expression driven by either *qa-2* or *tcu-1* promoters) are viable. Moreover, RNAi targeting
of two distinct regions of the *impβ1* gene, using the same strategy applied to *impα*, failed to
cause a strong growth defect and at best only reduced the expression level by around 40% (Fig.
S4A). As an alternative approach to achieve deficiency of Impβ1, we tested two Impβ1
inhibitors, Importazole and Ivermectin (Soderholm et al. 2011; Wagstaff et al. 2011). Both drugs
cause growth defects, with Importazole having less severe effects in a dose-dependent manner
(Fig. S4B). Based on the growth tests, 40µM drugs were used for fluorescent imaging
experiments. Importazole inhibits FRQ nuclear localization (Fig. 5A-C), suggesting a potential
role of Impβ1 in nuclear import, possibly together with Impα. Ivermectin treatment instead leads
to an overall increase in hyphal fluorescence without changing in nuclear signal (Fig. S4C-E),
which could result from cytoplasmic accumulation of FRQ or increased autofluorescence due to
growth-associated stress.

A full knockout of *impβ2* causes minor growth defects (Fig. S4F), shortens circadian period by
around 2.5 hours, and slightly reduces WCC activity leading to loss of persistence of the rhythm
(Fig. 5D-E). Consistent with this, knockdown or knockout of TNPO1 in mammalian cells also
shortens circadian period (Korge et al. 2018), suggesting a similar regulatory mechanism
between the two circadian systems. However, nuclear localization of neither FRQ nor WC-2 is
impaired in the *Δimpβ* knockout strains (Fig. 5F-G). FRAP experiments at CT4 showed no
difference in FRQ nuclear import rate either (Fig. 5H). There are also no obvious changes in
core clock protein expression levels in the *Δimpβ* knockout strains (Fig. S4G).

impβ3 is also essential, but viable conditional knockdowns were achieved by insertion of the
inducible *qa-2* promoter directly upstream of the ORF (Fig. 6A). A VHF tag was fused to its C
terminus to facilitate protein detection. In the absence of QA, the expression level of Impβ3 is
significantly decreased (Fig. 6B, Fig. S5A), resulting in a substantial circadian period length
shortening to around 15 hours. A relatively small amount of Impβ3 is sufficient to sustain a
normal clock, as even a low concentration (10^{-4}M) of QA restores the period to wild-type length,
and higher QA concentrations have no additional effect (Fig. 6C, Fig. S5B-C). Expression of

core clock proteins remains normal in Impβ3-deficient samples (Fig. 6B, Fig. S5A). To examine
subcellular localization of core clock components, endogenous fluorescent-tagged *frq* or *wc-2*
alleles were introduced into the inducible strain, which exhibited dramatically shortened periods
as expected (Fig. S5D). Despite the period effects, loss of Impβ3 resulted in no reduction in
nuclear localization of either FRQ or WC-2 (Fig. 6D-E).

Period shortening, particularly the dramatic period shortening seen with reduced Impβ3, is less
often seen with clock mutants. This prompted us to further investigate the possible mechanism
by which Impβ3 regulates the circadian clock by looking for possible interactions with protein
phosphatase 4 (PP4) whose loss also results in period shortening. PP4 has previously been
shown to regulate the nuclear enrichment of the WCC, and loss of *pph-4* also results in a
shortened circadian period, similarly to Impβ3 knockdown (Fig. S5E) (Cha et al. 2008; Dasgupta
2015). Interestingly, the period phenotypes caused *Δpph4* and *impβ3* reduction are not additive;
that is, knockdown on *impβ3* is epistatic to *pph4*. When Impβ3 is present (with QA), *Δpph-4*
caused the expected period shortening (Fig. S5E). However, using the same pair of strains,
combining *Δpph-4* with Impβ3 reduction (no QA) does not further shorten the circadian period
although rhythm persistence is reduced (Fig. 6F, Fig. S5E). This non-additive effect suggests
that Impβ3 and PP4 act within the same or overlapping pathways to regulate circadian
rhythmicity. Impβ3 may facilitate the proper subcellular localization of PP4 or its associated co-
enzymes, thereby enabling PP4 to precisely regulate phosphorylations of core clock proteins.
These findings point to functional crosstalk between nucleocytoplasmic transportation and
phosphatase-mediated regulation in maintaining the accuracy of circadian clock.

DISCUSSION

Our findings establish nuclear import as a key regulatory step in the *Neurospora* circadian clock. Using the live-cell imaging platform (Wang et al. 2025), we are able to measure in vivo subcellular dynamics in *N. crassa* under native, biologically relevant conditions. Through FRAP, we observed that FRQ is imported into nuclei slowly during the subjective day, requiring >20 min to recover fluorescence (Fig. 1). This disagrees with an earlier report of rapid nucleocytoplasmic shuttling of FRQ based on an anchor-away strategy that fused FRQ to FRB and tethered it to FKBP-tagged histone H1 (Diernfellner et al., 2009). While powerful, that system relies on rapamycin-induced heterodimerization of FRB-FKBP, which introduces an artificial import route that could diminish intrinsic clock function, raising the possibility that the observed fast shuttling is a result of methodological differences rather than the intrinsic behavior of FRQ in a free-running circadian cycle.

FRAP analyses revealed that FRQ nuclear import is not constant but varies across different stages of FRQ nuclear accumulation of the circadian cycle, being most active early in the subjective day and strongly weakened near subjective dusk (Fig. 1). This temporal regulation ensures that nuclear FRQ accumulation occurs with proper timing and rate, contributing to period stability. A previous report using FRAP to estimate nuclear import of mammalian Negative Arm proteins tested only extreme time points - when nuclear levels were at their peak or trough (Smyllie et al. 2016). As these conditions represent equilibrium states with little net nucleocytoplasmic exchange, it may be reasonable that no difference in import rate was detected (Smyllie et al. 2016). In contrast, by assaying import continuously across the subjective day, we observed a gradual decrease in nuclear import as FRQ approached its peak nuclear accumulation. The timescale of FRQ import we observed by FRAP (tens of minutes for recovery) is comparable to that of PER1 and PER2, functional homologs of FRQ in mammalian systems, whether assayed through endogenous tagging or transient expression (Öllinger et al. 2014; Smyllie et al. 2016; Korge et al. 2018). This similarity across fungal and mammalian systems suggests that slow nuclear import is a conserved feature of Negative Arm proteins and may represent an important delay step required to sustain a ~24-hour feedback loop.

Impα has been shown to regulate the subcellular localization of the Negative Arm in both mammalian systems and *Drosophila* (Umemura et al. 2014; Jang et al. 2015). The dependence of FRQ nuclear import on Impα, supported by both biochemical Co-IP and fluorescent imaging after RNAi knockdown experiments, places Impα at the core of this regulatory process (Fig. 2-4). Impα facilitates nuclear import by directly binding to the NLS on a target protein (Conti et al. 1998). Surprisingly, a domain previously shown to be essential for circadian rhythmicity suggested to function as a NLS on FRQ (Luo et al. 1998), is not required for nuclear localization of FRQ (Fig. 2) (Schunke et al. 2025). These strains exhibit persistently high WCC activity, suggesting that the loss of rhythmicity arises from a failure of FRQ to inhibit WCC despite its nuclear presence. Deletion of this motif has also been proposed to disrupt FRQ dimerization (Lauinger et al. 2014), which is essential for circadian clock and could be the cause of arrhythmicity (Cheng et al. 2001). A very recent work overexpressing fluorescent tagged FRQ fragments in mammalian cell culture support our conclusion and suggest the possible presence of multiple redundant NLSs on FRQ (Schunke et al. 2025). Similar redundancy has also been

reported for the Positive Arm WCC, whose nuclear localization doesn't rely on a single NLS
(Schwerdtfeger and Linden 2000; Cheng et al. 2003; Wang et al. 2016). Together, these
observations suggest that nuclear import in *Neurospora* is governed by a more complex and
possibly redundant regulatory mechanism than seen in other eukaryotes (Sakakida et al. 2005;
Kwon et al. 2006; Saez et al. 2011), consistent with the unique challenges posed by its syncytial
organization.

In addition to FRQ, our results demonstrate that WC-1 nuclear import is also strongly dependent
on Impα (Fig. 3-4). In Impα-deficient strains, WC-1 protein became undetectable despite normal
transcript levels, consistent with post-transcriptional destabilization. It may be that WC-1 is
unstable in the cytoplasm and is rapidly degraded when nuclear import is blocked, thereby
explaining the dramatic loss of WC-1 protein in Impα-deficient strains. By contrast, WC-2
nuclear import does not require Impα and its nuclear localization even increases with reduced
Impα. Although the overall amount of nuclear WCC does not show strong oscillations (Denault
et al. 2001), WCC has been reported to undergo rapid nucleocytoplasmic shuttling (Schafmeier
et al. 2008). NCU01249 encodes the sole Impα subunit in *Neurospora*, which must control
nuclear import of numerous cargo proteins and has been linked to additional cellular functions
beyond nuclear import (Klocko et al. 2015). Therefore, the effects of Impα deficiency on both
growth and circadian rhythmicity is likely mediated through multiple pathways beyond altered
localization of core clock proteins.

Our analysis of Impβ proteins further underscores the complexity of importin-mediated
regulation (Fig. 5-6). Impβ1 is essential and difficult to genetically manipulate, but chemical
inhibition suggests a role in FRQ nuclear localization, possibly in cooperation with Impα. Impβ2
deletion shortened the circadian period by ~2.5 h, paralleling results from knockdowns of its
mammalian homolog TNPO1 (Korge et al. 2018). However, nuclear localization of FRQ and
WC-2 was unaffected, suggesting that Impβ2 influences the clock through other cargoes or
regulatory pathways. Impβ3 is also important for precise circadian control: reduced expression
caused dramatic period shortening without altering nuclear localization of FRQ or WC-2, again
suggesting other cargoes or regulatory pathways. Genetic interaction experiments with *pph-4*
indicate that Impβ3 and PPH4 act within the same regulatory pathway, with Impβ3 potentially
functioning upstream to facilitate PPH4 activity or localization. This unexpected connection
highlights the interplay between nuclear transport and phosphatase-mediated regulation of
WCC.

Together, these results demonstrate that nuclear import is a dynamic and selective process that
plays a critical role in shaping circadian timing in *Neurospora*. Our findings parallel observations
in other eukaryotic systems (Sakakida et al. 2005; Saez et al. 2011; Öllinger et al. 2014;
Umemura et al. 2014; Jang et al. 2015; Lee et al. 2015; Korge et al. 2018; Korge et al. 2018),
underscoring the evolutionary conservation of importin-mediated circadian regulation, while also
revealing unique fungal-specific mechanisms.

MATERIALS AND METHODS

Strain construction:

Plasmids were constructed using pRS426 as backbone. Gene fragments were amplified from genomic DNA of *N. crassa* 74-OR23-1VA (FGSC, 2489) using Phusion Flash High-Fidelity PCR Master Mix (Thermo Scientific # F548S). Other fragments were either ordered as a gBlock or amplified from previously plasmids from the lab. Fragments were then assembled together using NEBuilder® HiFi DNA Assembly Master Mix (New England Biolabs Cat. #E2621) according to the manufacturer's instructions. All plasmids and their sequence data have been deposited at Addgene.

Plasmid	Addgene ID	Corresponding allele	Source or reference
frq[mNG_Flag]_hph	245515	*frq^{mNeonGreen_Flag}::hph+*	(Wang et al. 2025)
wc-2[mAp]_hph	245516	*wc-2^{mApple}::hph+*	
frq[ΔNLS, mNG_V5]_nat-1	251399	*frq^{ΔNLS, mNeonGreen_V5}::nat-1+*	
nup159[mApple]_bar	251400	*nup159^{mApple}::bar+*	
nup211[mTagBFP2]_hph	251401	*nup211^{mTagBFP2}::hph+*	
csr-1_qa-2_dsimpα_bar	251402	*csr-1::qa_dsimpα*	This Study
wc-1[ΔLOV, mAp]_bar	251403	*wc-1^{ΔLOV, mAp}::bar+*	
qa-2_[VHF]impβ3_bar	251404	*qa-2_^{VHF}impβ3::bar+*	
qa-2_[VHF]impβ3_nat-1	251405	*qa-2_^{VHF}impβ3::nat-1+*	

frq^{ΔNLS, mNeonGreen_V5}::nat-1+ contains the full *frq* gene with NLS1 (Luo et al. 1998) deleted, fused at its C-terminus to codon-optimized mNeonGreen and a V5 tag, followed by the nourseothricin resistance marker (He et al. 2020), designed to target the endogenous *frq* locus. The transformation cassette was amplified using Phusion Flash High-Fidelity PCR Master Mix (Thermo Scientific # F548S) and then transformed into strain 576-4 (*Δmus-52::hph+; Δfrq::hph+*).

nup159 (NCU08604) and *nup211* (NCU04059) were endogenously tagged at their C-termini using similar strategy as described previously (Bartholomai 2022) using different fluorescent tags and drug selection markers. *impα* (NCU01249) was endogenously V5-tagged at its C-terminus using a hygromycin resistant cassette for selection, resulting in the *impα^{V5}::hph+* allele.

The expression cassette producing double-stranded RNA targeting exon 3 of *importin α* (NCU01249) under control of the QA-inducible promoter was constructed as shown in Fig. 3A and inserted at the *csr-1* locus.

The *qa-2* promoter, along with a V5_6xHis_3xFlag (VHF) tag, was inserted at the N-terminus of *importin β3* (NCU03257) coding sequence using standard lab protocols to regulate its expression (Aronson et al. 1994). Two series of strains with different drug selection markers were constructed to accompany with other alleles during sexual crosses.

Transformations of *Neurospora crassa* were performed following lab protocol as described
previously (Bartholomai 2022), after which sexual crosses were carried out using standard
Neurospora methods (http://www.fgsc.net/Neurospora/NeurosporaProtocolGuide.htm) to
generate homokaryotic progeny with the desired combination of alleles.

FRAP:
5 independent runs were performed for FRAP experiments using strain 1946-1 (*csr-1::son-*
1^{mTagBFP2}; ras-1^{bd}; wc-1^{ΔLOV}::bar+, for, frq^{mNeonGreen_Flag}::hph+) in the customized microfluidic
devices as described previously (Wang et al. 2025) with Bird medium (Metzenberg 2004)
containing 0.2% glucose, 0.2% poly-acrylic acid, and 1mM sodium formate. The cultures were
entrained by temperature steps before resetting to CT0. 2.5 µg/ml benomyl was added to the
medium at the same time of resetting. For each independent run, one FRAP experiment was
performed on a new region within the same culture every half an hour from CT2 to CT10. Single
focal plane images were acquired on LSM 880 with the pinhole size adjusted to an optical
section of 2.8 µm. For each recording, 40 to 50 frames were collected at 30 seconds interval.
After the first two frames, 4-7 nuclei were photobleached with a 488 nm laser to around 30% of
the initial fluorescence intensity. SON-1^{mTagBFP2} (DAPI channel) was recorded at the same time
as nuclear marker.

Measurements of F_ROI (manually bleached nuclei), P_ROI (untreated nuclei) and B_ROI
(outside of the cell) were taken in ImageJ. Rolling averages of 3 time points (n-1, n, n+1) were
calculated and used for further analysis. Fluorescence recordings were corrected for
background. The fitted values based on an averaged slope (photobleaching rate) calculated
from the linear regressions of *P_ROI - B_ROI* within the same run were used to correct for
photobleaching during recording:

$$418 \qquad F\_ROI_{corrected} = \frac{(F\_ROI - B\_ROI)}{(\widehat{P\_ROI - B\_ROI})}$$

419 *F_ROI_{corrected}* was then normalized to pre-beach intensities (set as 100%) and grouped into 1-
420 hour bins, i.e. all *F_ROI_{corrected}* from HT3 and HT3.5 were considered as group HT3, for Fig. 1D.

421 A linear regression was calculated for each nucleus (*F_ROI_{corrected}*). Then, the slopes of these
422 fitted curves were grouped into 1-hour bins for Fig. 1E. A small number of nuclei were excluded
423 because of poor goodness of fit (r^2).

Time	CT2	CT3	CT4	CT5	CT6	CT7	CT8	CT9	CT10
N of included nuclei ($r^2 \geqslant 0.5$)	4	22	26	32	35	37	32	24	18
N of excluded nuclei ($r^2 < 0.5$)	0	1	1	1	0	1	1	2	0

424 For Fig. S1E, a similar analysis was performed using linear regression of a single example
425 *P_ROI - B_ROI* within the same FRAP experiment (same recording/region) to correct for
426 photobleaching, rather than using the average in one run. Recordings without a suitable P_ROI
427 were excluded from the analysis.

Time	CT2	CT3	CT4	CT5	CT6	CT7	CT8	CT9	CT10
N of included nuclei ($r^2 \geqslant 0.5$)	4	21	20	25	26	36	28	24	16
N of excluded nuclei ($r^2 < 0.5$)	0	0	0	0	1	2	1	2	2

Luciferase assay: All luciferase assays were performed and analyzed in the same way as
described previously (Kelliher et al. 2023). Unless otherwise specified, all entrainments and
recordings were performed at 25°C. For the benomyl test in Fig. S1D, 50 µl per well of water or
2.5 µg/ml benomyl was injected at indicated time using 1ml syringes. All other drugs, as well as
the inducer quinic acid, were added directly to the medium before inoculation at the indicated
concentrations.

Immunoprecipitation and Western blot: Co-IP experiments were performed following lab
protocol with slight modifications (Chen et al. 2010). Protein samples were collected, and Input
samples were saved, using the same method described in Wang et al., (2025). For V5 IP, V5
beads were prepared by preincubating 2µl V5 antibody (Invitrogen #46-0705) with 30µl
Dynabeads™ Protein G (Invitrogen #1003D) per sample at 4°C for 1 hour. After determining
protein concentrations in the protein samples by Bradford assay, 2mg protein per sample in a
1ml volume was incubated with the V5 beads at 4°C for 2 hours or overnight. The bound
complex was then eluted with 100µl 2x Sample Buffer (Diluted from NuPAGE™ LDS Sample
Buffer (4X), Invitrogen #NP0007) following two washes with Wash Buffer (50mM Hepes-KOH,
pH 7.4, 150mM NaCl, 0.4% NP-40) and two washes with no-NP Wash Buffer (50mM Hepes-
KOH, pH 7.4, 150mM NaCl). For detection, 30µg Input and 10µl IP samples were loaded onto
NuPAGE™ 3-8% Tris-Acetate Mini Protein Gels (ThermoFisher Scientific, Catalog
#EA03785BOX). Western blotting was then performed as described in Wang et al., (2025).

For the *qa_dsimpα* samples, cultures were initiated in 10cm dishes with LCM (no QA). Five
~1cm diameter circles were cut and transferred to 275ml flasks containing 50ml LCM, with or
without 10^{-2}M QA (as indicated in figures), and cultured for 21 hours before harvest. For all other
samples, strains were maintained in the same LCM medium (with or without 10^{-2}M QA)
throughout the culture. All cultures were performed in light at 25°C unless stated otherwise.

Fluorescent imaging: Samples were grown at 25 °C in the light on agarose pads (1× Vogel's
Salts, 2% sucrose, 0.5% arginine, 50 ng/mL biotin, 1% agarose) overnight, unless otherwise
specified, and imaged using the 'inverted agar block' method (Hickey et al. 2004) on one of two
Nikon Eclipse Ti-E stand-based spinning disk confocal microscope systems described in
Chapter 4. 10 µm z-stacks (35 slices at 300 nm step size) were collected in corresponding
channels. Tip regions were cropped in ImageJ and then 3D reconstituted in Arivis Vision 4D for
segmentation and quantification. Mean fluorescent signals from both the total hyphae and
individual nuclei were measured and background signals, defined as the mean intensity from a
manually generated object outside the hyphae in each z-stack, were subtracted. The coefficient
of variation (CV) for each nucleus was calculated as the standard deviation of its fluorescence

signal (SD) divided by its background subtracted mean fluorescent signal

($MEAN_{background\ substracted}$):

$$CV = \frac{SD}{MEAN_{background\ substracted}}$$

For Fig. 5E, samples were germinated on agarose pads without drug in light and then
transferred to darkness for 12 hours. At DD12, a small piece of agarose at the growth front was
cut, and its top surface was immersed in water or a 40μM drug solution. After 2 hours of drug
and light exposure, samples were imaged using the same methods. Quinic acid was added to
the agarose pad prior to inoculation in corresponding experiments.

Race tubes / growth tests: Race tube assays were performed as described previously
(Aronson et al. 1994). or growth tests with Impβ1 inhibitors, race tube medium was
supplemented with the indicated concentrations of drug. Conidia of strain 661-4a (*his-3::frq_cbox-*
luc) were inoculated at the edge of 35-mm dishes and incubated overnight at 25 °C in constant
light before being scanned.

RT-qPCR: Samples were cultured and harvested as described for Western blotting. RNA was
extracted from ~200μg ground powder per sample using Direct-zol RNA Miniprep Kits
(ZYMO Research, R2050). For cDNA synthesis, 3μg extracted RNA reverse-transcribed using
SuperScript™ IV First-Strand Synthesis System (Invitrogen, 18091050) with the provided Oligo
d(T)$_{20}$ primer. The resulting cDNA was diluted to ~7.5 ng/μl, and 2 μl was used for RT-qPCR with
iTaq Universal SYBR Green Supermix (Bio-Rad, #1725121) measured with an Applied
Biosystems™ StepOne™ Real-Time PCR System. The primers used for RT-qPCR are listed
below:

Target	Primers	Product Length	Source
crp-43 NCU08964	Forward: CTGTCCGTACTCGTGACTCC Reverse: ACCATCGATGAGGAGCTTGC	173bp	(Hurley et al. 2015)
Impα NCU01249	Forward: GCATGCTCAGGAATGCTACC Reverse: TCGTCGTCGAGCGAGTAGAC	178bp	This study
wc-1 NCU02356	Forward: TCAGCAGCATCAGTTCAACC Reverse: GTTGATGTTCGCCCATCTCT	116bp	(Chen et al. 2009)
frq NCU02265	Forward: TGGCTCGGATAAGAATGGTC Reverse: ATGAAAGGTGTCCGAAGGTG	103bp	(Chen et al. 2009)

	Forward: GGATCCTAACCGCAGCGAAG Reverse: GATCATGGCGGTGATCCAGT	179bp	This study
Impβ1 NCU02011			

RT-qPCR data were analyzed using the $\Delta\Delta C_t$ method, with *crp-43* (NCU08964) (Hurley et al.
2015) as the internal control. All reactions were performed in triplicate for each biological sample
in one biological replicate, and results are presented as mean ± standard deviation from 3
independent biological replicates.

For light pulse experiments, samples were initiated in light, the cut circles were transferred to
flasks in the dark and cultured for 21 h. Cultures were then either harvested in dark (DD21) or
exposed to light for 15 minutes before harvest (15min LP).

Statistical analysis and data visualization:

Statistical analysis was performed using Prism 10 (GraphPad) with either one-way ANOVA
followed by multiple comparisons or Welch's t test. The number of replicates (n) used for each
experiment are indicated in the relevant figure legends. All schematic diagrams were created in
BioRender.

DATA AVAILABILITY STATEMENT

The authors affirm that all data necessary for confirming the conclusions of the article are
present within the article, figures, and tables. Strains and plasmids are available upon request.

ACKNOWLEDGMENTS

This study was supported by a National Institutes of Health (NIH) grant awarded to Jay C.
Dunlap (R35GM118021). We acknowledge use of stocks from the Fungal Genetics Stock
Center (FGSC.net), NIAID-supported informatic resources at FungiDB (Contract
HHSN75N93019C00077), and Molecular Biology facilities supported by the Dartmouth CQB
COBRE (NIH NIGMS grant P20GM130454), and the Dartmouth BioMT (NIH NIGMS grant P20-
GM113132). Imaging was performed in the Life Sciences Light Microscopy Facility
(RRID:SCR_027143) at Dartmouth College.

REFERENCES

Aronson BD, Johnson KA, Loros JJ, Dunlap JC. 1994. Negative Feedback Defining a Circadian Clock: Autoregulation of the Clock Gene frequency. Science. 263(5153):1578–1584. doi:10.1126/science.8128244.

Baker CL. 2010. Post-translational regulation of the Neurospora crassa circadian system [Ph.D.]. Dartmouth College. [accessed 2025 Sept 14]. https://www.proquest.com/docview/219994612/abstract/9232CEC881334441PQ/1.

Baker CL, Kettenbach AN, Loros JJ, Gerber SA, Dunlap JC. 2009. Quantitative Proteomics Reveals a Dynamic Interactome and Phase-Specific Phosphorylation in the Neurospora Circadian Clock. Mol Cell. 34(3):354–363. doi:10.1016/j.molcel.2009.04.023.

Bartholomai BM. 2022. Cell Biological Dissection of the Neurospora Circadian Oscillator: Tools, Approaches, and Fundamental Observations [Ph.D.]. [United States -- New Hampshire]: Dartmouth College. https://www.proquest.com/dissertations-theses/cell-biological-dissection-neurospora-circadian/docview/2632997349/se-2?accountid=10422.

Beck M, Hurt E. 2017. The nuclear pore complex: understanding its function through structural insight. Nat Rev Mol Cell Biol. 18(2):73–89. doi:10.1038/nrm.2016.147.

Bernardes NE, Fukuda CA, da Silva TD, de Oliveira HC, de Barros AC, Dreyer TR, Bertolini MC, Fontes MRM. 2020. Comparative study of the interactions between fungal transcription factor nuclear localization sequences with mammalian and fungal importin-alpha. Sci Rep. 10(1):1458. doi:10.1038/s41598-020-58316-9.

Bernardes NE, Takeda AAS, Dreyer TR, Freitas FZ, Bertolini MC, Fontes MRM. 2015. Structure of Importin-α from a Filamentous Fungus in Complex with a Classical Nuclear Localization Signal. PLOS ONE. 10(6):e0128687. doi:10.1371/journal.pone.0128687.

Cha J, Chang S, Huang G, Cheng P, Liu Y. 2008. Control of WHITE COLLAR localization by phosphorylation is a critical step in the circadian negative feedback process. EMBO J. 27(24):3246–3255. doi:10.1038/emboj.2008.245.

Chen C-H, DeMay BS, Gladfelter AS, Dunlap JC, Loros JJ. 2010. Physical interaction between VIVID and white collar complex regulates photoadaptation in Neurospora. Proc Natl Acad Sci. 107(38):16715–16720. doi:10.1073/pnas.1011190107.

Chen C-H, Ringelberg CS, Gross RH, Dunlap JC, Loros JJ. 2009. Genome-wide analysis of light-inducible responses reveals hierarchical light signalling in Neurospora. EMBO J. 28(8). doi:10.1038/emboj.2009.54.

Cheng P, Yang Y, Heintzen C, Liu Y. 2001. Coiled-coil domain-mediated FRQ–FRQ interaction is essential for its circadian clock function in Neurospora. EMBO J. 20(1–2):101–108. doi:10.1093/emboj/20.1.101.

Cheng P, Yang Y, Wang L, He Q, Liu Y. 2003. WHITE COLLAR-1, a Multifunctional
NeurosporaProtein Involved in the Circadian Feedback Loops, Light Sensing, and
Transcription Repression of wc-2 *. J Biol Chem. 278(6):3801–3808.
doi:10.1074/jbc.M209592200.

Chook Y, Blobel G. 2001. Karyopherins and nuclear import. Curr Opin Struct Biol.
11(6):703–715. doi:10.1016/S0959-440X(01)00264-0.

Conti E, Uy M, Leighton L, Blobel G, Kuriyan J. 1998. Crystallographic Analysis of the
Recognition of a Nuclear Localization Signal by the Nuclear Import Factor Karyopherin α.
Cell. 94(2):193–204. doi:10.1016/S0092-8674(00)81419-1.

Costa JI, Muñoz-Guzmán F, Larrondo LF. 2025. Clock Regulation and Fungal Physiology:
Molecular Mechanisms Underpinning the Timely Control of Just About Everything.
doi:10.1146/annurev-micro-060424-051158. [accessed 2025 Aug 9].
https://www.annualreviews.org/content/journals/10.1146/annurev-micro-060424-051158.

Crosthwaite SK, Dunlap JC, Loros JJ. 1997. Neurospora wc-1 and wc-2: Transcription,
Photoresponses, and the Origins of Circadian Rhythmicity. Science. 276(5313):763–769.
doi:10.1126/science.276.5313.763.

Dasgupta A. 2015. Biological Significance of VIVID Photocycle Length and the Control of
Gene Expression by the Neurospora Clock [Ph.D.]. Dartmouth College. [accessed 2025
Sept 15].
https://www.proquest.com/docview/1758936730/abstract/29396917764B407CPQ/1.

Dekhang R, Wu C, Smith KM, Lamb TM, Peterson M, Bredeweg EL, Ibarra O, Emerson JM,
Karunarathna N, Lyubetskaya A, et al. 2017. The Neurospora Transcription Factor ADV-1
Transduces Light Signals and Temporal Information to Control Rhythmic Expression of
Genes Involved in Cell Fusion. G3 GenesGenomesGenetics. 7(1):129–142.
doi:10.1534/g3.116.034298.

Denault DL, Loros JJ, Dunlap JC. 2001. WC-2 mediates WC-1–FRQ interaction within the
PAS protein-linked circadian feedback loop of Neurospora. EMBO J. 20(1–2):109–117.
doi:10.1093/emboj/20.1.109.

Dunlap JC. 1999. Molecular Bases for Circadian Clocks. Cell. 96(2):271–290.
doi:10.1016/S0092-8674(00)80566-8.

Dunlap JC, Loros JJ. 2017. Making Time: Conservation of Biological Clocks from Fungi to
Animals. Microbiol Spectr. 5(3):10.1128/microbiolspec.funk-0039–2016.
doi:10.1128/microbiolspec.funk-0039-2016.

Galagan JE, Calvo SE, Borkovich KA, Selker EU, Read NO, Jaffe D, FitzHugh W, Ma LJ,
Smirnov S, Purcell S, et al. 2003. The genome sequence of the filamentous fungus
Neurospora crassa. Nat 2003 4226934. 422(6934):859–868. doi:10.1038/nature01554.

Giles NH, Case ME, Baum J, Geever R, Huiet L, Patel V, Tyler B. 1985. Gene organization and
regulation in the qa (quinic acid) gene cluster of Neurospora crassa. Microbiol Rev.
49(3):338–358. doi:10.1128/mr.49.3.338-358.1985.

Goldfarb DS, Corbett AH, Mason DA, Harreman MT, Adam SA. 2004. Importin α: a
multipurpose nuclear-transport receptor. Trends Cell Biol. 14(9):505–514.
doi:10.1016/j.tcb.2004.07.016.

Haruki H, Nishikawa J, Laemmli UK. 2008. The Anchor-Away Technique: Rapid, Conditional
Establishment of Yeast Mutant Phenotypes. Mol Cell. 31(6):925–932.
doi:10.1016/j.molcel.2008.07.020.

He L, Guo W, Li J, Meng Y, Wang Y, Lou H, He Q. 2020. Two dominant selectable markers for
genetic manipulation in Neurospora crassa. Curr Genet. 66(4):835–847.
doi:10.1007/s00294-020-01063-1.

Hickey PC, Swift SR, Roca MG, Read ND. 2004. Live-cell Imaging of Filamentous Fungi
Using Vital Fluorescent Dyes and Confocal Microscopy. In: Methods in Microbiology. Vol.
34. Academic Press. (Microbial Imaging). p. 63–87. [accessed 2025 July 5].
https://www.sciencedirect.com/science/article/pii/S0580951704340031.

Hurley JH, Dasgupta A, Andrews P, Crowell AM, Ringelberg C, Loros JJ, Dunlap JC. 2015. A
Tool Set for the Genome-Wide Analysis of Neurospora crassa by RT-PCR. G3-GENES
GENOMES Genet. 5(10):2043–2049. doi:10.1534/g3.115.019141.

Jang AR, Moravcevic K, Saez L, Young MW, Sehgal A. 2015. Drosophila TIM Binds Importin
α1, and Acts as an Adapter to Transport PER to the Nucleus. PLOS Genet. 11(2):e1004974.
doi:10.1371/JOURNAL.PGEN.1004974.

Káldi K, González BH, Brunner M. 2006. Transcriptional regulation of the Neurospora
circadian clock gene wc-1 affects the phase of circadian output. EMBO Rep. 7(2):199–204.
doi:10.1038/sj.embor.7400595.

Kelliher CM, Stevenson E-L, Loros JJ, Dunlap JC. 2023. Nutritional compensation of the
circadian clock is a conserved process influenced by gene expression regulation and
mRNA stability. PLOS Biol. 21(1):e3001961. doi:10.1371/journal.pbio.3001961.

Klocko AD, Rountree MR, Grisafi PL, Hays SM, Adhvaryu KK, Selker EU. 2015. Neurospora
Importin α Is Required for Normal Heterochromatic Formation and DNA Methylation. PLOS
Genet. 11(3):e1005083. doi:10.1371/journal.pgen.1005083.

Korge S, Maier B, Brüning F, Ehrhardt L, Korte T, Mann M, Herrmann A, Robles MS, Kramer A.
2018. The non-classical nuclear import carrier Transportin 1 modulates circadian rhythms
through its effect on PER1 nuclear localization. PLoS Genet. 14:e1007189.
doi:10.1371/journal.pgen.1007189.

Kwon I, Lee J, Chang SH, Jung NC, Lee BJ, Son GH, Kim K, Lee KH. 2006. BMAL1 Shuttling
Controls Transactivation and Degradation of the CLOCK/BMAL1 Heterodimer. Mol Cell Biol.
26(19):7318–7330. doi:10.1128/MCB.00337-06.

Larrondo LF, Colot HV, Baker CL, Loros JJ, Dunlap JC. 2009. Fungal Functional Genomics:
Tunable Knockout-Knock-in Expression and Tagging Strategies. Eukaryot Cell. 8(5):800–
804. doi:10.1128/ec.00072-09.

Lauinger L, Diernfellner A, Falk S, Brunner M. 2014. The RNA helicase FRH is an ATP-
dependent regulator of CK1a in the circadian clock of Neurospora crassa. Nat Commun.
5(1):3598. doi:10.1038/ncomms4598.

Lee K, Dunlap JC, Loros JJ. 2003. Roles for WHITE COLLAR-1 in Circadian and General
Photoperception in *Neurospora crassa*. Genetics. 163(1). doi:10.1093/genetics/163.1.103.

Lee Y, Jang AR, Francey LJ, Sehgal A, Hogenesch JB. 2015. KPNB1 mediates PER/CRY
nuclear translocation and circadian clock function. Kramer A, editor. eLife. 4:e08647.
doi:10.7554/eLife.08647.

Lee Y, Shen Y, Francey LJ, Ramanathan C, Sehgal A, Liu AC, Hogenesch JB. 2019. the nRon
complex controls circadian clock function through regulated peR and cRY nuclear
translocation. Sci Rep 2019 91. 9(1):1–12. doi:10.1038/s41598-019-48341-8.

Lin C, Feng S, DeOliveira CC, Crane BR. 2023. Cryptochrome–Timeless structure reveals
circadian clock timing mechanisms. Nature. 617(7959):194–199. doi:10.1038/s41586-023-
06009-4.

Luo C, Loros JJ, Dunlap JC. 1998. Nuclear localization is required for function of the
essential clock protein FRQ. EMBO J. 17(5):1228–1235. doi:10.1093/emboj/17.5.1228.

Mans B, Anantharaman V, Aravind L, Koonin EV. 2004. Comparative Genomics, Evolution
and Origins of the Nuclear Envelope and Nuclear Pore Complex. Cell Cycle. 3(12):1625–
1650. doi:10.4161/cc.3.12.1316.

Markina-Iñarrairaegui A, Etxebeste O, Herrero-García E, Araújo-Bazán L, Fernández-
Martínez J, Flores JA, Osmani SA, Espeso EA. 2011. Nuclear transporters in a
multinucleated organism: functional and localization analyses in Aspergillus nidulans. Mol
Biol Cell. 22(20):3874–3886. doi:10.1091/mbc.e11-03-0262.

Mason DA, Stage DE, Goldfarb DS. 2009. Evolution of the Metazoan-Specific Importin α
Gene Family. J Mol Evol. 68(4):351–365. doi:10.1007/s00239-009-9215-8.

Metzenberg R. 2004. Bird Medium: an alternative to Vogel Medium. Fungal Genet Rep.
51(1):19–20. doi:10.4148/1941-4765.1138.

Öllinger R, Korge S, Korte T, Koller B, Herrmann A, Kramer A. 2014. Dynamics of the
circadian clock protein PERIOD2 in living cells. J Cell Sci. 127(19):4322–4328.
doi:10.1242/jcs.156612.

Pelham JF, Mosier AE, Altshuler SC, Rhodes ML, Kirchhoff CL, Fall WB, Mann C, Baik LS,
Chiu JC, Hurley JM. 2023. Conformational changes in the negative arm of the circadian
clock correlate with dynamic interactomes involved in post-transcriptional regulation. Cell
Rep. 42(4). doi:10.1016/j.celrep.2023.112376. [accessed 2025 July 7].
https://www.cell.com/cell-reports/abstract/S2211-1247(23)00387-X.

Pemberton LF, Paschal BM. 2005. Mechanisms of Receptor-Mediated Nuclear Import and
Nuclear Export. Traffic. 6(3):187–198. doi:10.1111/j.1600-0854.2005.00270.x.

Saez L, Derasmo M, Meyer P, Stieglitz J, Young MW. 2011. A Key Temporal Delay in the
Circadian Cycle of Drosophila Is Mediated by a Nuclear Localization Signal in the Timeless
Protein. Genetics. 188(3):591–600. doi:10.1534/genetics.111.127225.

Sakakida Y, Miyamoto Y, Nagoshi E, Akashi M, Nakamura TJ, Mamine T, Kasahara M, Minami
Y, Yoneda Y, Takumi T. 2005. Importin α/β mediates nuclear transport of a mammalian
circadian clock component, mCRY2, together with mPER2, through a bipartite nuclear
localization signal. J Biol Chem. 280(14):13272–13278. doi:10.1074/jbc.M413236200.

Schafmeier T, Diernfellner A, Schäfer A, Dintsis O, Neiss A, Brunner M. 2008. Circadian
activity and abundance rhythms of the Neurospora clock transcription factor WCC
associated with rapid nucleo–cytoplasmic shuttling. Genes Dev. 22(24):3397–3402.
doi:10.1101/gad.507408.

Schunke C, Ruppert B, Schultz S, Diernfellner ACR, Brunner M. 2025. CK1a-Mediated Two-
Step Subunit Remodeling of the Circadian FRQ-FRH Complex. :2025.05.13.653691.
doi:10.1101/2025.05.13.653691. [accessed 2025 Aug 4].
https://www.biorxiv.org/content/10.1101/2025.05.13.653691v1.

Schwerdtfeger C, Linden H. 2000. Localization and light-dependent phosphorylation of
white collar 1 and 2, the two central components of blue light signaling in Neurospora
crassa. Eur J Biochem. 267(2):414–422. doi:10.1046/j.1432-1327.2000.01016.x.

Shi M, Collett M, Loros JJ, Dunlap JC. 2010. FRQ-Interacting RNA Helicase Mediates
Negative and Positive Feedback in the Neurospora Circadian Clock. Genetics. 184(2):351–
361. doi:10.1534/genetics.109.111393.

Smith KM, Sancar G, Dekhang R, Sullivan CM, Li S, Tag AG, Sancar C, Bredeweg EL, Priest
HD, McCormick RF, et al. 2010. Transcription Factors in Light and Circadian Clock Signaling
Networks Revealed by Genomewide Mapping of Direct Targets for Neurospora White Collar
Complex. Eukaryot Cell. 9(10):1549–1556. doi:10.1128/ec.00154-10.

Smyllie NJ, Pilorz V, Boyd J, Meng Q-J, Saer B, Chesham JE, Maywood ES, Krogager TP,
Spiller DG, Boot-Handford R, et al. 2016. Visualizing and Quantifying Intracellular Behavior
and Abundance of the Core Circadian Clock Protein PERIOD2. Curr Biol. 26(14):1880–1886.
doi:10.1016/j.cub.2016.05.018.

Soderholm JF, Bird SL, Kalab P, Sampathkumar Y, Hasegawa K, Uehara-Bingen M, Weis K,
Heald R. 2011. Importazole, a Small Molecule Inhibitor of the Transport Receptor Importin-
β. ACS Chem Biol. 6(7):700–708. doi:10.1021/cb2000296.

Ström A-C, Weis K. 2001. Importin-beta-like nuclear transport receptors. Genome Biol.
2(6):reviews3008.1. doi:10.1186/gb-2001-2-6-reviews3008.

Takeda AAS, Freitas FZ, Magro AJ, Bernardes NE, Fernandes CAH, Goncalves RD, Bertolini
MC, Fontes MRM. 2013. Biophysical Characterization of the Recombinant Importin-α from
Neurospora crassa. http://www.eurekaselect.com. [accessed 2025 Sept 16].
https://www.eurekaselect.com/article/47441.

Umemura Y, Koike N, Matsumoto T, Yoo S-H, Chen Z, Yasuhara N, Takahashi JS, Yagita K.
2014. Transcriptional program of Kpna2/Importin-α2 regulates cellular differentiation-
coupled circadian clock development in mammalian cells. Proc Natl Acad Sci.
111(47):E5039–E5048. doi:10.1073/pnas.1419272111.

Wagstaff KM, Rawlinson SM, Hearps AC, Jans DA. 2011. An AlphaScreen®-Based Assay for
High-Throughput Screening for Specific Inhibitors of Nuclear Import. J Biomol Screen.
16(2):192–200. doi:10.1177/1087057110390360.

Wang B, Zhou X, Loros JJ, Dunlap JC. 2016. Alternative Use of DNA Binding Domains by the
Neurospora White Collar Complex Dictates Circadian Regulation and Light Responses. Mol
Cell Biol. 36(5):781–793. doi:10.1128/MCB.00841-15.

Wang Z, Bartholomai BM, Wang B, Ritz D, Schultz D, Loros JJ, Dunlap JC. 2025. Core Clock
Protein Subcellular Dynamics Coordinate Local and Global Circadian Control in
Syncytia. :2025.09.30.679467. doi:10.1101/2025.09.30.679467. [accessed 2025 Nov 17].
https://www.biorxiv.org/content/10.1101/2025.09.30.679467v1.

FIGURE LEGENDS

Figure 1 | Nuclear import of FRQ is circadian regulated.

A) Schematic of the experimental design. FRAP experiments were performed during the subjective day following entrainment by temperature cycles. Whole nuclei were bleached, and the recovery of their fluorescent signal reflects nuclear import of cytoplasmic FRQ. B) Example timelapses from subjective morning (CT4, top) and afternoon (CT8, bottom), showing different recovery rates`. White arrowheads indicate example target nuclei. Scale bar = 5 µm. C) Average fluorescent recovery curves for each hour of the subjective day. D) Nuclear signal recovery speed, calculated as the slope of linear regression of each target nucleus's recovery curve. Data were analyzed by Pearson correlation, with the correlation coefficient (r) and p-value indicated in the plot.

Figure 2 | Core clock proteins directly bind Importinα while bypassing the classic NLS for nuclear import.

A) Co-immunoprecipitation between endogenous Importinα and core clock proteins. Cell lysates were immunoprecipitated with anti-V5 antibody. The interaction between FRQ and FRH is included as a positive control. B) Example central focal images showing FRQ or FRQ$^{\Delta NLS}$ subcellular localization during free-run circadian cycles or in constant light. FRQ is shown in green, the inner nuclear envelope marker NUP211 in blue, and the outer nuclear envelope marker NUP159 in magenta. All targets are endogenously tagged with monomeric fluorescent proteins. Scale bar = 10 µm. C) Quantification of nuclear FRQ or FRQ$^{\Delta NLS}$ signal from 3D renderings of z-stacks (Mean ± SD). Each dot represents the mean fluorescent signal of one nucleus. N = 7-11 hyphal tips per condition, with the ungrouped data of each tip plotted in Fig. S2E.

Figure 3 | Impα is required for normal circadian rhythmicity.

A) Schematic of the siRNA knockdown design. Expression of dsDNA targeting the last exon of *impα* is driven by the *qa-2* inducible promoter. With the addition of the inducer quinic acid (QA), dsRNA is expressed and subsequently processed to inhibit *impα* expression. B) Growth of *qa_dsimpα* strains in race tubes with or without 10^{-2} M QA. C) Expression level of *impα* (top), *wc-1* (middle), or *frq* (bottom) in control (*impα^{V5}*) or *qa_dsimpα* strains with or without 10^{-2} M QA quantified by RT-qPCR. Error bars represent the standard deviation from biological triplicates. D) Protein levels in control (*impα^{V5}*) or *qa_dsimpα* strains with or without 10^{-2} M QA. Example targets, including core clock proteins, were detected by immunoblotting (left), and total protein levels were assessed by Coomassie staining (right). * indicates unspecific bands in immunoblots. E) Luciferase assays of control (*impα^{V5}*) or *qa_dsimpα* strains with varying QA concentrations (top) and corresponding period lengths (bottom). n=4 replicates per condition. Error bars show one standard deviation. Periods were analyzed by one-way ANOVA with multiple comparisons.

Figure 4 | Nuclear localization of FRQ and WC-1, but not WC-2, depends on Impα.

A) Maximum projection images of example z-stacks showing FRQ and WC-2 subcellular localization in constant light with or without 10^{-2} M QA. Scale bar = 10 μm. B-C) Quantification of B) nuclear FRQ or C) total FRQ signal in tip regions from 3D renderings of z-stacks. One control strain (1984-4, $frq^{mNeonGreen}::hph+$ $nup159^{mTagBFP2}::nat-1+$) and 2 $csr-1::qa\_dsimpα$ $nup159^{BFP}$ $wc-2^{mApple}$; frq^{mNG} strains were analyzed. 2 (controls) or 3 ($dsimpα$ strains) biological replicates were quantified, with individual replicates plotted in Fig. S3B. N = 5-12 hyphal tips per condition per replicate. Each dot represents the mean fluorescent signal of one nucleus (nuclear signal) or one hypha (total signal). D) Schematic of the potential mechanism by which WCC amount and localization are affected when Impα is deficient. E-G) Expression level of E) $wc-1$, F) frq, or G) $impα$ in a $qa\_dsimpα$ strain at DD21 with or without 10^{-2} M QA, before or after 15-minute light pulse, quantified by RT-qPCR. Error bars represent the standard deviation from biological triplicates. H-I) Quantification of H) nuclear WC-2 or I) total WC-2 signal in tip regions from 3D renderings of z-stacks. One control strain (1996-2, $wc-2^{mApple}::hph+$, $his-3::cbox-luc$) and 2 $csr-1::qa\_dsimpa$ $nup159^{BFP}$ $wc-2^{mApple}$; frq^{mNG} strains were analyzed. Same methods as Fig. 4B-C were used.

Figure 5 | Deficiency of Importinβs affects FRQ localization or circadian clock.

A) Maximum projection images of example z-stacks showing FRQ subcellular localization with or without Importazole, an Importinβ1 inhibitor. Samples were treated with drug (or water) and light for 2 hours starting at DD12. N = 5-6 hyphal tips. Each dot represents the mean fluorescent signal of one nucleus (nuclear signal) or one hypha (total signal). Scale bar = 10 μm. B-C) Quantification of B) nuclear FRQ or C) total FRQ signal in tip regions from 3D renderings of z-stacks (Mean ± SD). Each dot represents the mean fluorescent signal of one nucleus (nuclear signal) or one hypha (total signal). D) Luciferase assays of control strain ($csr-1::frq_{cbox}-luc$) or three $Δimpβ2$ strains (left) and their period lengths (right). n=4 replicates. The shade shows one standard deviation. Periods were analyzed by one-way ANOVA with multiple comparisons. D) Luciferase assays showing the free-run clock of $Δimpβ2$ FRQ imaging strain used in E) compared to its control strain (2066-4) without $Δimpβ2$. Each curve represents the average of three biological triplicates each with 4 technical replicates, with the shade showing one standard deviation. F-G) Maximum projection images of example z-stacks showing F) FRQ or G) WC-1 subcellular localization ((left) and quantification of nuclear signal (right) in constant light, with or without Impβ2. Scale bar = 10 μm. N = 3-6 hyphal tips. Each dot represents the mean fluorescent signal of one nucleus. H) Nuclear FRQ signal recovery speed at CT4 with or without Impβ2, calculated as the slope of linear regression of each target nucleus's recovery curve.

Figure 6 | Shortened period in Impβ3-deficient strains is independent of core clock protein levels or localization.

A) Schematic of the inducible $impβ3$ design. Expression of $impβ3$ is driven by the $qa-2$ inducible promoter. Without the addition of the inducer quinic acid (QA), expression of $impβ3$ is reduced. B) Core clock protein levels in control strain or inducible $impβ3$ strains with or without 10^{-2} M QA. The result shown is representative of biological triplicates; additional replicates are shown in Fig. S5D. C) Luciferase assays of the inducible $impβ3$ strain with varying QA concentrations

(left) and their period lengths (right). n=4 replicates. The shade shows one standard deviation.
Periods were analyzed by one-way ANOVA with multiple comparisons. D-E) Maximum
projection images of example z-stacks showing D) FRQ or E) WC-2 subcellular localization
((left) and quantification of nuclear signal (right) in constant light, with or without 10^{-2} M QA.
Scale bar = 10 µm. N = 4-7 hyphal tips. Each dot represents the mean fluorescent signal of one
nucleus. Two sibling strains were quantified for WC-2. F) Luciferase assays of strains with both
pph4 deletion and *impβ3* deficiency compared to strains with *impβ3 deficiency* alone. n=4
replicates. The shade shows one standard deviation.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6