Location and mapping of the human rostromedial tegmental nucleus and associated midbrain inhibitory nuclei regulating dopamine neurons

Anastasia Filimontseva[1,2], ORCID 0000-0002-3316-9035 email afil9180@uni.sydney.edu.au

YuHong Fu[1,2], ORCID 0000-0003-4539-2039 email yuhong.fu@sydney.edu.au

Glenda M Halliday[1,2,#] ORCID 0000-0003-0422-8398 email glenda.halliday@sydney.edu.au

[1]The University of Sydney Brain and Mind Centre & Faculty of Medicine and Health School of Medical Sciences, Sydney, NSW 2050, Australia

[2]Aligning Science Across Parkinson's (ASAP) Collaborative Research Network, Chevy Chase, MD, 20815

[#]Corresponding author: Professor Glenda Halliday, University of Sydney Brain and Mind Centre, 97 Church Street, Camperdown, NSW 2050 AUSTRALIA, Email: glenda.halliday@sydney.edu.au, Tel: +61 2 9351 0888

Running title: The human rostromedial tegmental nucleus

Number of text words: 3,551

Number of figures: 5 plus 3 supplementary figures and 2 supplementary tables

Acknowledgements

The authors acknowledge: Heidi Cartwright for assistance with the figures; human brain tissue sample procurement by the Sydney Brain Bank, Sydney (enabled by funding from Neuroscience Research Australia and is a special gift in memory of Jim Raftos from the Shaw family) and Flinders Medical Centre and Institute of Medical and Veterinary Sciences, Adelaide; the use of Microscopy Australia infrastructure at the Sydney Microscopy and Microanalysis facility, University of Sydney (enabled by the National Collaborative Research Infrastructure Strategy);

28   and funding support by Aligning Science Across Parkinson's [020505] through the Michael J.

29   Fox Foundation for Parkinson's Research and the National Health and Medical Research Council

30   of Australia to GMH [Investigator Grant 1176607]. For open access, the authors have applied a

31   CC BY public copyright license to all Author Accepted Manuscripts arising from this

32   submission.

Abstract

Background: Animal experiments reveal distinct GABAergic cell clusters within the

dopaminergic midbrain regions in the rostromedial tegmental nucleus (RMTg) and retrorubral

fields (RRF) that have yet to be clearly defined in humans. These neurons send prominent

inhibitory projections to dopaminergic neurons in the substantia nigra and ventral tegmental area

that impact motor, reward and threat processing.

Objective: To identify GABergic RMTg and RRF cell clusters in the human brain.

Methods: Ten µm formalin-fixed paraffin-embedded transverse human brainstem sections from

ten control cases were obtained from the Sydney Brain Bank and immunohistochemistry for

GABA and tyrosine hydroxylase was used to determine the location and cell size of RMTg and

RRF GABAergic neurons. These cell populations were further mapped in transverse 50µm thick

cresyl violet stained serial midbrain sections (every 750µm) from previously published controls

(Halliday et al. 1990a).

Results: GABAergic neurons were cytoarchitecturally distinct, with the largest GABAergic

neurons in the RRF, followed by RMTg neurons which were larger than GABAergic neurons in

the interpeduncular nucleus (Kruskal-Wallis test, p<0.0001). RMTg and RRF GABAergic

neurons first appear in caudal transverse midbrain sections approximately 38mm above the obex.

RMTg moves rostrally and medially from underneath the decussation of the superior cerebellar

peduncle to just lateral to the interpeduncular nucleus. The RRF cluster also moves rostrally and

medially to the parabrachial pigmented nucleus (PBP) just under the red nucleus.

Conclusion: The GABAergic neurons in RMTg and RRF/PBP that modulate dopamine neuronal

excitability have distinct morphologies in humans. Identifying these inhibitory neurons is key to

evaluating their role in neurodegenerative diseases.

Key words: dopaminergic neurons, GABAergic neurons, parabrachial pigmented nucleus,

rostromedial tegmental nucleus, retrorubral fields, three-dimensional reconstruction

Introduction

The majority of somatodendritic inputs to dopamine (DA) neurons in the substantia nigra pars

compacta (SNpc) are GABAergic (~40-70% of neurons)(Bolam and Smith 1990; Henny et al.

2012). These inhibitory afferents arise from a number of sources, some extrinsic to the SNpc

(e.g. from the periaqueductal gray, dorsal raphe, lateral habenula, and basal

ganglia)(Omelchenko and Sesack 2010; Beier et al. 2015; Nieh et al. 2015; Hjelmstad et al.

2013; Lerner et al. 2015; Evans et al. 2020; Vounatsos and Gittis 2025b), while others more

recently defined are interspersed within the midbrain DA regions.

The main midbrain DA regions are the SNpc, retrorubral fields (RRF) and ventral tegmental area

(VTA), with the VTA cytoarchiteturally subdivided into the more lateral parabrachial pigmented

nucleus (PBP), medial paranigral (PN) and parainterfascicular (PIF) nuclei, and midline

interfascicular (IF), central linear (CL) and rostral linear (RL) nuclei (Fu et al. 2012). Within

these DA regions, the RRF has the highest proportion of GABA to DA neurons compared to

other midbrain DA clusters (Nair-Roberts et al. 2008; Vounatsos and Gittis 2025b; Kelly et al.

2022). In contrast only around 29% of SNpc neurons are GABAergic, which is the lowest

proportion among the main DA regions, and the VTA has a moderate population of GABAergic

neurons (Nair-Roberts et al. 2008). The highest densities of GABAergic neurons in the VTA are

in the caudal PN and PIF subnuclei (Nair-Roberts et al. 2008), although the rostromedial

tegmental nucleus (RMTg) has not been identified in humans but in other species it is a

GABAergic nucleus in 'the tail of the VTA' (Barrot et al. 2012).

Many GABAergic neurons intermixed within these DA regions are local interneurons. In

contrast, GABAergic neurons broadly innervating midbrain DA neurons concentrate only in the

RMTg and RRF (Barrot et al. 2012; Smith et al. 2019; Vounatsos and Gittis 2025a). The RMTg

was initially defined in rodents with retrograde injections into the SNpc and VTA which reliably

identified this concentration of GABAergic neurons (Smith et al. 2019; Jhou et al. 2009a;

Lammel et al. 2012). In addition to its direct inhibition of the motor nigrostriatal DA pathway
(Bourdy et al. 2014), the RMTg in general encodes negative prediction errors and plays a critical
role in behavioural inhibition in general, as well as punishment learning through mesolimbic DA
pathways (Jhou 2021). As discussed above, of the midbrain DA regions the RRF has the highest
concentration of GABAergic neurons which densely innervate the SNpc, although their function
is less well defined (Vounatsos and Gittis 2025a). In gerbils, the RRF GABAergic pathway is
involved in male sex behaviour (Simmons et al. 2011) with RRF projections in general involved
in reward processing, motivation and threat evaluation (Moaddab and McDannald 2021;
Vounatsos and Gittis 2025a).

Due to species differences in midbrain anatomy and neural identification using developmental
markers (e.g. some transcription factors for RMTg neurons are only expressed in embryonic
tissue and not in adult human brain (Allen Institute for Brain Science 2010)), the human RMTg
has yet to be defined and GABAergic clusters in the human RRF have largely been ignored. The
fact that there are multiple GABAergic populations in close proximity within these anatomical
regions (Margolis et al. 2012) further complicates the identification of these important GABergic
neuronal nuclei in the human midbrain. However, the comparative identification of these
inhibitory neurons is important to determine as these midbrain nuclei provide broad inhibition to
midbrain DA neurons with any deficits likely to have significant impact.

For the first time we mapped the RMTg and a large RRF GABAergic neuronal cluster (LatC) in
the human brainstem using immunohistochemistry and three-dimensional reconstructions in
serial sections showing that these GABAergic cell clusters have distinct morphological features
and anatomical locations. Overall, we present detailed anatomical maps of these prominent
human midbrain GABAergic nuclei that innervate midbrain DA neurons.

Materials and Methods

Human brain tissue

For immunohistochemical analysis, formalin-fixed paraffin-embedded (FFPE) cross-sectional
midbrain tissue blocks from healthy controls without neurological or neuropathological diseases
were obtained from the Sydney Brain Bank (n=10, for details see Supplementary Table 1). Serial
section reconstruction using thick cross-sectional Nissl-stained midbrain sections from control

cases (n=4, for details see Supplementary Table 1) previously processed and published (Halliday

et al. 1990b) were also included. These midbrain sections were cut at 50μm on a freezing

microtome, collected at 750μm intervals, and stained with 0.5% cresyl violet (HB5608 Hello

Bio), as previously described (Halliday et al. 1990b). The study was approved by the University

of Sydney Human Research Ethics Committee (2021/845).

Immunohistochemistry

FFPE tissue blocks were sectioned at 6μm and dried on microscope slides at 37°C for 48 hours.

Sections were deparaffinized by incubation at 60°C for 1 hour and then submerged in D-

Limonene (Hurst Scientific, HISTO-5LCTN) for 2 x 7 minutes, followed by rehydration in

decreasing ethanol concentrations (100% ethanol for 2 x 3 minutes, 95% ethanol for 3 minutes,

70% ethanol for 3 minutes) and distilled H2O for 3 minutes. Sections underwent heat-induced

antigen retrieval (HIAR) in commercially available antigen retrieval solution (Invitrogen, 00-

4956-58) diluted to 1x using a programmable antigen retrieval cooker (Aptum Biologics Ltd,

UK, Retriever 2100). Peak temperature reached ~121°C and afterwards sections were gradually

cooled for two hours. Sections were washed with tris-buffered saline (TBS, Sigma-Aldrich,

94158-10TAB), and endogenous alkaline phosphatase (AP) and hydrogen peroxidase (HRP)

were quenched with BLOXALL blocking solution (SP-6000-100) for 30 minutes. Afterwards

sections were blocked for 1 hour at room temperature with in-house made normal horse serum

blocking solution (2% horse serum, 1% BSA in 0.01M TBS). Sections were incubated with a

primary cocktail of antibodies to tyrosine hydroxylase (TH, rabbit polyclonal IgG, AB152

Sigma-Aldrich diluted 1:200) and GAD67 (mouse monoclonal IgG$_3$, sc-28376 Santa-Cruz

Biotechnology diluted 1:50) for 4 nights at 4°C. Secondary antibody incubation for 90 minutes at

room temperature used a cocktail of ImmPRESS™-alkaline phosphatase (AP) horse anti-mouse

IgG polymer detection kit (MP-5402 Vector Laboratories) and ImmPRESS™-horseradish

peroxidase (HRP) horse anti-rabbit IgG polymer detection kit (MP-7401 Vector Laboratories).

Antibody detection was visulaised using SignalStain® Vibrant Red AP Substrate Kit (76713 Cell

Signaling) and ImmPACT® DAB EqV HRP Substrate Kit (SK-4103 Vector Laboratories).

Sections then underwent standard dehydration in increasing graded ethanol concentrations and

cleared in xylene, prior to mounting with Eukitt® quick-hardening mounting medium (03989

Sigma-Aldrich). In separate control experiments where either the primary or secondary

antibodies were omitted,was no AP or HRP was detectable.

Image acquisition

An automated slide scanner (SLIDEVIEW VS200, Tokyo, Japan) was used to scan the sections
using pre-set focusing and exposure parameters. The method for brightfield image acquisition
has previously been described (Filimontseva et al. 2025) with slight amendments to optimize the
exposure settings for 1) the GABA and TH stained sections, and 2) for serial Nissl stained
sections.

Anatomical mapping

Midbrain DA regions were cytoarchitecturally mapped on digitized images in Adobe Photoshop
(ver. 25.9.1, Adobe Systems Inc., San Jose, CA, USA) and Illustrator (ver. 28.5, Adobe Systems
Inc., San Jose, CA, USA) based on the human brainstem atlas and previously published
cytoarchitecture studies (Paxinos et al. 2020; McRitchie et al. 1995; McRitchie et al. 1997). For
the GABA and TH stained sections, TH staining was used to determine the DA brain regions and
for the serial Nissl stained sections, neuromelanin pigment was used to determine the DA brain
regions. To determine cytoarchitectural landmarks for the RMTg, published anatomical maps
from rodents and non-human primates were used (Supplementary Figures 1-3) and the most
frequent landmarks determined (Supplementary Table 2). GABA and TH stained thin sections
were used to identify other relevant GABAergic neuronal clusters. Both thick and thin sections
were used to reconstruct proposed three-simensional models of the DA and GABA regions of
interest. The midline and aqueduct were used to align every other 50µm, taken at 750µm
intervals. Landmarks were outlined and relevant clusters colour coded in Adobe Photoshop.
Stacked Nissl and GABA stained sections were compiled in Adobe Illustrator, and mapped
structures compared.

Quantification

A semi-automated method was used to perform morphological analysis of identified GABAergic
neuronal clusters. Images of Nissl stained and GABA stained sections were loaded into QuPath
(Bankhead et al. 2017) and the system trained to identify neurons in the region of interest based
on stain intensity. Successful training was used to create a pixel classifier and identified neurons
were filtered by size and shape. Due to technical challenges of successful GABA staining on
human FFPE tissue, six cases with conspicuous GABA neuronal staining out of the total ten
cases used for mapping were selected for this analysis. The morphological features of Nissl

stained neurons in the regions of interest were analysed between 33mm to 38mm Obex at every

1.6mm interval.

Statistical analysis

Statistical analyses were performed in GraphPad Prism (version 10.0.3, GraphPad Software Inc.,

California, United Stated). Kruskal-Wallis non-parametric test was used to compare cell

circularity of neurons in the GABAergic cell cluster of the RMTg, RRF and interpeduncular

nucleus in Nissl and GABA stained sections. A two-way ANOVA was run to analyse effect of

anatomical region and tissue stain on neurnal size, followed by post-hoc Tukey tests. The

proportion and density of GABA to Nissl neurons in the GABAergic cell clusters was also

analysed with two-way ANOVA tests. For all analysis the statistical significance was set to p-

value < 0.05.

Results

Comparative landmarks to identify the human RMTg

We identified comparative cytoarchitectural landmarks for the human RMTg based on animal

mapping literature (Supplementary Table 2). These important landmarks from this literature

oriented our anatomical search for the RMTg in human midbrain tissue sections (Supplementary

Table 2). The RMTg in other species first appears at the level of caudal interpeduncular nucleus,

posterior to PN (Supplementary Figure 1)(Jhou et al. 2009b; Kaufling et al. 2009). As the

superior cerebellar peduncle appears, the RMTg shifts dorsally (Supplementary Figure 2). The

RMTg is elongated at this level where it shifts laterally below the RRF.

Morphological identification of distinct inhibitory cell clusters in the human midbrain DA

regions

Prominent distinct inhibitory neuronal clusters were identified in the RRF (an obvious novel

GABAergic neuronal cell cluster we call LatC) and RMTg and were compared morphologically

to GABAergic neurons in the interpeduncular nucleus (Figure 1). GABAergic neurons in these

cell clusters appeared to be morphologically different, with the RRF LatC GABAergic neurons

larger, rounder and more densely labelled (Figure 1f) compared to the small-medium sized

RMTg GABAergic neurons that stained lighter and were randomly oriented (Figure 1e). The size

of GABAergic neurons in these cell clusters were compared to those in the interpeduncular

nucleus (Figure 1a-f) revealing a significant difference between anatomical region [F (2,10796)

= 166.9, p<0.0001]. Post-hoc Tukey tests revealed that RRF LatC GABAergic neurons were

significantly larger than RMTg GABAergic neurons, with both being significantly larger than

GABAergic neurons in the interpeduncular nucleus (Figure 1g). Analysis of neuronal size in

Nissl versus GABA stained sections (Figure 1a-f) revealed a significant main effect of the stain

used on neuronal size [F (1,10796) = 21.41, p<0.0001]. Further, there was a significant

difference in neuronal size between anatomical region depending on the stain used [F (2,10796)

= 31.60, p<0.0001]. GABAergic neurons in the interpeduncular nucleus were smaller on average

than Nissl stained interpeduncular neurons, whereas the reverse was true for the RMTg and RRF

LatC GABAergic neurons (Figure 1g). Cell circularity measurements confirmed that RRF LatC

neurons were on average rounder than RMTg neurons [Kruskal-Wallis test, ****

p<0.0001](Figure 1h), although not so much for the GABAergic neurons in these regions (Figure

1i). GABAergic neurons in the RRF LatC were rounder than those in the interpeduncular nucleus

[Kruskal-Wallis test, * p<0.05](Figure 1i). The size and morphology of these concentrated

midbrain GABAergic cell clusters distinguishes these different populations of midbrain

GABAergic neurons.

Figure 1. *Morphological identification of distinct inhibitory clusters in the human brainstem.*
Representative images from Nissl (CV) stained serial sections (a-c) and GABA stained sections (d-f) of specific
neuronal clusters. Interpeduncular nucleus (IPN) is evident in a) and d), and the rostromedial tegmental nucleus
(RMTg) is presented in b) and e). A novel, lateral cluster (LatC) of GABAergic neurons in the RRF is presented in
c) and f). Measures of neuronal area (g) and cell circularity (h and i) in both tissue preparations validate the
identification of these distinct inhibitory neuronal cell clusters. For area analysis, a two-way ANOVA revealed that
anatomic region [F (2,10796) = 166.9, p<0.0001] and stain type significantly affected neuronal size [F (1,10796) =
21.41, p<0.0001] in g). Error bars display SEM. Post-hoc Tukey tests showed that LatC neurons were the largest,
**** p<0.0001. IPN neurons were the smallest but significantly differed in size between CV and GABA stained
tissue [F (2,10796) = 31.60, p<0.0001 in g]. GABAergic neurons in IPN were slightly smaller than Nissl stained
neurons, with the reverse occurring in the RMTg and RRF. Cell circularity analyses also identified differences in
these neuronal cell clusters in both tissue preparations. Violin plots revealed that LatC (magenta) neurons were
significantly rounder than RMTg (blue) and IPN (yellow) in CV stained sections [Kruskal-Wallis test, ****
p<0.0001 in (h)]. GABA staining revealed significant differences only between LatC and IPN circularity [Kruskal-
Wallis test, * p<0.05 in (i)].

Chemoarchitectural mapping of GABAergic cell clusters in midbrain DA regions

Chemoarchitectural mapping of GABAergic cell clusters in midbrain DA regions

Midbrain DA regions surrounding the GABAergic RMTg and RRF LatC were

cytoarchitecturally delineated and mapped using the GABA and TH colabelled sections (Figure

2). The RMTg first appeared caudally in the midbrain approximately 33-34mm above the Obex

where it is at its largest in cross section mostly due to an absence of surrounding DA neurons and

subregions (Figure 2). It lies lateral to the interpeduncular nucleus, dorsal to the pontine nuclei

and ventromedial to the superior cerebellar peduncle (Figure 2). At these caudal levels, there is

an obvious RRF LatC GABAergic cell cluster located some distance from the RMTg at the

medial edge of the RRF on the ventrolateral edge of the superior cerebellar peduncle (Figure 2).

In more rostral midbrain sections, the RMTg GABAergic cell cluster maintains its position but is

reduced in size due to its location within the DA neurons of the VTA PN subregion and

ascending DA fibres (Figure 2). The RMTg is discontinued rostrally when the dense compact

SNpc occurs in cross sections (around 37mm above Obex)(Figure 2). The RRF LatC cell cluster

maintains its position laterally until the VTA PBP subcluster begins to expand laterally in more

rostral sections. At these levels (35-36mm above the Obex) this GABAergic cell cluster bridges

between the medial RRF and lateral PBP (Figure 2). With the discontinuation of the RRF

rostrally, the LatC moves ventromedially to embed into the PBP on the ventral aspect of the

superior cerebellar peduncle and continues to move ventromedially to be adjacent to the RMTg

within the VTA DA regions (Figure 2).

Figure 2. *Chemoarchitectural mapping of GABAergic cell clusters in midbrain DA regions.* The main
DA regions and landmarks were mapped in a) after immunohistochemical staining on 6μm FFPE sections. Tyrosine
hydroxylase (TH)+ neurons were found in the dorsal, ventral and medial substantia nigra pars compacta (SNpc) tiers
(SNd, SNv and SNM), retrorubral field (RRF), parabrachial pigmented nucleus (PBP), paranigral nucleus (PN),
ventral tegmental area (VTA), caudal linear nucleus (CLi) and the interfascicular nucleus (IF). Other relevant
landmarks include the pedunculopontine tegmental nucleus (PPTg), medial lemniscus (ml), superior cerebellar
peduncle (scp), central tegmental tract (ctg), mammillotegmental tract (mtg), the medial longitudinal fasciculus
(mlf), the interpeduncular nucleus (IPN), pontine nuclei (pn) and cerebellar peduncle (cp). Main TH+ SNpc
subclusters are annotated in dark grey, whereas VTA subclusters and other relevant landmarks are shaded in light
grey. Micrographs of GABA and TH immunohistochemically stained sections (GABA labeling with alkaline
phosphatase in pink; TH labeled with diaminobenzidine in brown) show neurons in the RMTg and RRF/PBP LatC
in b) and c) respectively. The nuclei of interest were mapped along a 4mm distance in the transverse plane. Scale
bars equal 5mm in a) and 100μm in b) and c).

Cytoarchitectural identification of the RMTg and RRF LatC in midbrain DA regions

The RMTg and RRF LatC were cytoarchitecturally identified and mapped using serial thick

Nissl stained cross-sectional midbrain sections to observe cell clusters more readily through a

greater number of midbrain levels (Figure 3). Midbrain DA regions were identified using

neuromelanin pigment. The RRF/PBP LatC was observed within the same anatomical location in

these midbrain DA subregions, that is, in the medial edge of the RRF shifting in rostral sections

to the dorsolateral PBP subregion (Figure 3).

The cytoarchitecture of the RMTg is highlighted in more detail as this subregion has not

previously been identified cytoarchitecturally in humans. In Nissl stained midbrain sections,

RMTg neurons seem smaller, more round and overall denser compared to their neighboring DA

cell regions (Figure 3 and Supplementary Figure 3). Caudally the RMTg cell cluster can be

identified lateral to the interpeduncular nucleus, and in more rostral sections collocates dorsal to

the medial SNpc cell cluster alongside/overlapping with the VTA PN DA subregion (Figure 3).

Figure 3. *Cytoarchitectural identification of the RMTg and RRF/PBP lateral GABAergic cell cluster*
(LatC) in the human midbrain. a) Schematic mapping of 50μm-thick Nissl (CV) stained sections shown
in b). Substantia nigra pars compacta (SNpc) and ventral tegmental area (VTA) subclusters, as well as
retrorubral field (RRF) and other relevant nuclei (e.g. the pedunculopontine tegmental nucleus or PPTg),
are annotated in grey. RRF, VTA, PPTg and several other nuclei are shown in light grey, whereas SNpc
subdivisions are shaded in darker grey. The aqueduct (Aq) and fiber tracts, such as medial lemniscus (ml),
superior cerebellar peduncle (scp), central tegmental tract (ctg), mammillotegmental tract (mtg) and the
medial longitudinal fasciculus (mlf), are also relevant landmarks. The rostromedial tegmental nucleus
(RMTg, blue) is located slightly lateral to the interpeduncular nucleus (IPN, yellow), whereas the
RRF/PBP lateral GABAergic cell cluster (LatC, magenta) is much more lateral. c) Higher magnification
regional micrographs of RMTg and cellular view of individual neurons in d). Scale bars equal 5mm in a),
2mm in b), 200μm in c), and 200μm in d). Abbreviations: substantianigra lateral part (SNl); medial,
dorsal and ventral tiers, SNm, SNd and SNv; parabrachial pigmented nucleus (PBP); paranigral nucleus
(PN); caudal linear nucleus (CLi); interfascicular nucleus (IF); pontine nuclei (pn) and cerebellar
peduncle (cp).
303

Integrated cyto- and chemo-architectural three-dimensional (3D) model of the GABAergic cell

clusters in the human RMTg and RRF/PBP

Three dimensional modelling revealed consistent overlap between the structural and chemical

location of the mapped GABAergic cell clusters in both tissue types (Figure 4a). Both the RMTg

and RRF/PBP LatC shift medially min the midbrain from caudal to rostral (Figure 4a). A pseudo-

3D model of the RMTg and RRF/PBP LatC was constructed across cases in the rostrocaudal

plane (Figure 4b). This model confirms the concentration of the RMTg to the lateral edge of the

interpeduncular nucleus (Figure 4b). The larger area of the LatC in the ventromedial RRF hugs

the ventrolateral edge of the superior cerebellar peduncle above and the medial lemniscus below,

then travels rostromedially to the PBP and the more ventromedial aspect of the superior cerebellar

peduncle (Figure 4b).

The composition and density of neurons in these GABAergic cell clusters was then determined

and compared. The composition of neurons was relatively uniform in the three GABAergic

neuronal regions in humans with approximately 40-60% of neurons staining with the GAD67

antibody (Figure 4c). However, the density of GABAergic neurons differs between these regions

with a higher density of GABAergic neurons in the interpeduncular nucleus compared to the

RMTg and LatC, and a lower density of non-GABAergic neurons in the LatC [$F_{(2,28)} = 3.737$,

* $p<0.0364$](Figure 4f). Overall our data show that there are discrete concentrations of

morphologically different GABAergic neurons in two midbrain DA regions in humans, the

RMTg with smaller neurons (previously identified in non-human species), and a novel more

concentrated cell cluster with larger GABAergic neurons in the RRF/PBP.

Figure 4. *Integrated cyto- and chemo-architectural three dimensional model of the GABAergic neuronal clusters in the human midbrain DA regions.* a) Schematic reconstructions of inhibitory nuclei of interest. Cross-sectional midbrain Nissl (CV) stained serial sections (left) and GABA-labeled sections (right) were collapsed and integrated across approximately 4-6mm in the rostrocaudal plane. Scale bar equals 5mm. To visualise a pseudo-3D transformation of these inhibitory midbrain nuclei, traced cytoarchitectural sections were aligned at a slight angle and the zoomed in region is shown in b). As expected, the rostromedial tegmental nucleus (RMTg, blue) closely associates with the interpeduncular nucleus (IPN), whereas the lateral cell cluster (LatC, magenta) shifts between the retrorubral field (RRF) and parabrachial pigmented nucleus (PBP) DA subregions. Within these GABAergic cell clusters, neuronal populations were further assessed in c) and d). Approximately half of IPN, RMTg

335   and LatC neurons are GABAergic in c). However, the neuronal density differs because with a significant main effect
336   of anatomical region on cell density in d) [F (2,28) = 3.737, * p=0.0364]. Error bars display SEM in c) and d).
337   Abbreviations: aqueduct (Aq); medial lemniscus (ml); central tegmental tract (ctg); mammillotegmental tract (mtg);
338   medial longitudinal fasciculus (mlf); superior cerebellar peduncle (scp); pontine nuclei (pn); cerebellar peduncle
339   (cp).

Discussion

This study identified two concentrated clusters of inhibitory neurons in the human midbrain DA

regions, with one cluster equivalent to the RMTg that has been previously identified in non-

human species (Figure 5a and RMTg information in Supplementary file) and the other a novel

dense cluster of larger GABAergic neurons we called LatC that crosses the RRF and PBP DA

subregions (Figure 5b). Morphological analyses confirmed that these two inhibitory neuronal

clusters are distinct. The RMTg has not been identified in the human midbrain, and to the best of

our knowledge the separate inhibitory neuronal population in the LatC has also not been

identified. Approximately half of the neuronal population in these cell clusters are GABAergic

with LatC having a lower density of non-GABAergic neurons compared with the RMTg. In

rodents the proportion of GABAergic neurons in the RMTg is approximately 70-90% (Morello

and Partanen 2015), although human brain tissue fixation may have contributed to an undercount

of GABAergic neurons in our study. We used a GAD67 antibody to label cytoplasmic GABA

and longer fixation can alter GAD epitopes to reduce the number of detectable GABAergic

neurons (Howat and Wilson 2014; Vincek et al. 2003). However, we used antigen retrieval

methods and the proportion of non-DA to DA neurons in the substantia nigra is known to be

higher in rodents compared with humans (Hardman et al. 2002), with our data consistent with

this orthogonal observation. Further studies are required to confirm our findings in humans and

to determine the identity of the non-GABAergic neuronal population in these midbrain nuclei.

In addition to their morphological differences, these two GABAergic nuclei have been shown to

be molecularly distinct developmentally in rodents (Lahti et al. 2016; Düdükcü et al. 2024).

Specifically, one study has shown that *Sox14* and *Zfmp2* transcription factors distinguish

developing RMTg from RRF GABAergic neurons (Lahti et al. 2016), while another study shows

that *Nkx2.2* identifies three developing GABAergic subtypes in the lateral non-SNpc regions

(RRF not identified directly) and that *Slc32a1* identified six developing GABAergic subtypes in

the SNpc and VTA regions (two GABAergic neuron subtypes and four DA+GABA neuron

subtypes) (Düdükcü et al. 2024). This suggests that there are molecular differences between
these midbrain GABAergic cell clusters which require further evaluation in humans.

External inhibitory input   **Local midbrain inhibitory input**

Figure 5. Connectivity of inhibitory nuclei to DA brain regions in humans. a) Rodent anatomical
mapping of RMTg (red outlines of dark blue neuron representative circles among other tract traced neurons in light
blue). Figure adapted and reproduced with permission from *Smith et al., 2019* (Smith et al. 2019). b) GABA and DA
clusters in the caudal human midbrain showing the lateral cluster (LatC), RMTg and interpeduncular nucleus (IPN).
c) Summarised inhibitory input from rodent and non-human primate connectivity literature. VTA receives input
from periaqueductal gray, dorsal raphe, lateral habenula and ventral pallidum. SNpc also receives projections from
the ventral pallidum, as well as other regions included in the blue box. These DA regions also receive inhibitory
projections from the RMTg, substantia nigra pars reticulata (SNpr) and RRF. Scale bars equal 5mm in b) and left
side c), and 2mm in right side c). Abbreviations: aqueduct (Aq); medial lemniscus (ml); superior cerebellar peduncle
(scp); caudal linear (CLi); parabrachial pigmented nucleus (PBP); paranigral nucleus (PN); substantia nigra pars
compacta (SNpc) lateral part, medial, dorsal and ventral tiers, SNl, SNm, SNd and SNv, respectively; central

tegmental tract (ctg); mammillotegmental tract (mtg); medial longitudinal fasciculus (mlf); cerebellar peduncle (cp);
pontine nuclei (pn).

Using anatomical landmarks, we have generated three dimensional cyto- and chemo-

architectural maps of the human midbrain to show the location of these inhibitory cell clusters

within the DA subregions. Mapping these midbrain inhibitory nuclei in humans will enable

further understanding of their molecular signatures, function and dysfunction in disease. The

recent evidence that chronic hyperactivity of DA neurons in animal models leads to DA cell

dysfunction and loss (Rademacher et al. 2025) makes identifying and evaluating these midbrain

GABAergic nuclei important.

Differences between the midbrain inhibitory circuits

As identified in the introduction, midbrain DA neurons in the SNpc and VTA receive major

GABA input from a variety of sources that can be divergent in function (Figure 5c). Midbrain

projecting external inhibitory circuits differ functionally, molecularly and structurally.

Genetically defined globus pallidus and dorsal striatum cells inhibit SNpc DA neurons but only

striatal inputs induce rebound firing (Evans et al. 2020). Aldh1a1+ SNpc neurons receive inputs

from molecularly distinct striatal clusters, which have opposite effects on locomotion (Dong et

al. 2025), potentially due to their location on distal dendrites - striosomal GABAergic axons have

a positive relationship between inhibition strength and distance away from the SNpc proximal

dendrite (Evans et al. 2020). Evidence of functional differences in the input of GABA from the

RMTg versus extrinsic sources to the VTA is that activation of RMTg GABA terminals induces

place avoidance whereas PAG GABA terminal activation induces immobility (St. Laurent et al.

2020). Further work on determining how different inhibitory circuits influence DA neuronal

function is required. Importantly, the inhibition exerted by RMTg GABAergic neurons on

midbrain DA neurons is bigger compared to the one mediated for example by the substantia

nigra pars reticulata GABA neurons (Morello and Partanen 2015), further warranting the study

of these inhibitory midbrain nuclei in humans.

The identified RMTg and RRF/PBP LatC GABAergic nuclei must be functionally different from

each other as well as external GABA inputs based on their location within different functional

brain regions (Figure 5c). The GABAergic nuclei embedded in the DA subregions likely mediate

different effects on midbrain DA neurons. Recent evidence in primates show that RRF and PBP

GABAergic neurons receive equivalent GABAergic inputs to their DA neighbours (Fudge et al. 2024). Neural circuits with such mutual inhibitory connections are able to rapidly and flexibly switch between distinct functions allowing multiple controllable functions to be managed through the same circuits (Liu et al. 2025). The demonstration of these two distinct GABAergic nuclei that receive similar afferents to their DA neighbours and innervate broadly across all DA subregions is likely to at least double the tunable flexibility of the multiple functions regulated through DA circuits. These and other inhibitory nuclei are promising anatomical correlates for unexplored biological and disease mechanisms.

Overall, this is the first study to identify RMTg and RRF/PBP LatC GABAergic nuclei in the human midbrain. The data presented show clear concentrations of large GABAergic neurons in these human midbrain regions, higher numbers than expected for the small intrinsic interneurons also found in DA subregions. Given their major input to midbrain DA neurons it will be important to understand how their interactions impacts midbrain DA motor and behavioural functions, particularly as chronic hyperactivity of DA neurons in animal models leads to DA cell dysfunction and loss (Rademacher et al. 2025).

References

Allen Institute for Brain Science (2010) Allen Human Brain Atlas. Allen Institute for Brain Science. https://human.brain-map.org. Accessed 24/10 2025

Bankhead P, Loughrey MB, Fernández JA, Dombrowski Y, McArt DG, Dunne PD, McQuaid S, Gray RT, Murray LJ, Coleman HG, James JA, Salto-Tellez M, Hamilton PW (2017) QuPath: Open source software for digital pathology image analysis. Scientific Reports 7 (1):16878. doi:10.1038/s41598-017-17204-5

Barrot M, Sesack SR, Georges F, Pistis M, Hong S, Jhou TC (2012) Braking dopamine systems: a new GABA master structure for mesolimbic and nigrostriatal functions. J Neurosci 32 (41):14094-14101. doi:10.1523/JNEUROSCI.3370-12.2012

Beier Kevin T, Steinberg Elizabeth E, DeLoach Katherine E, Xie S, Miyamichi K, Schwarz L, Gao Xiaojing J, Kremer Eric J, Malenka Robert C, Luo L (2015) Circuit Architecture of VTA Dopamine Neurons Revealed by Systematic Input-Output Mapping. Cell 162 (3):622-634. doi:10.1016/j.cell.2015.07.015

Bolam JP, Smith Y (1990) The GABA and substance P input to dopaminergic neurones in the substantia nigra of the rat. Brain Research 529 (1):57-78. doi:https://doi.org/10.1016/0006-8993(90)90811-O

Bourdy R, Sanchez-Catalan MJ, Kaufling J, Balcita-Pedicino JJ, Freund-Mercier MJ, Veinante P, Sesack SR, Georges F, Barrot M (2014) Control of the nigrostriatal dopamine neuron activity and motor function by the tail of the ventral tegmental area. Neuropsychopharmacology 39 (12):2788-2798. doi:10.1038/npp.2014.129

Dong J, Wang L, Sullivan BT, Sun L, Martinez Smith VM, Chang L, Ding J, Le W, Gerfen CR, Cai H
(2025) Molecularly distinct striatonigral neuron subtypes differentially regulate locomotion.
Nature Communications 16 (1):2710. doi:10.1038/s41467-025-58007-x
Düdükcü Ö, Raj DDA, van de Haar LL, Grossouw LM, Linders LE, Garritsen O, Adolfs Y, van
Kronenburg NCH, Broekhoven MH, Kapteijns THW, Meye FJ, Pasterkamp RJ (2024) Molecular
diversity and migration of GABAergic neurons in the developing ventral midbrain. iScience 27
(11):111239. doi:https://doi.org/10.1016/j.isci.2024.111239
Evans RC, Twedell EL, Zhu M, Ascencio J, Zhang R, Khaliq ZM (2020) Functional Dissection of Basal
Ganglia Inhibitory Inputs onto Substantia Nigra Dopaminergic Neurons. Cell Rep 32
(11):108156. doi:10.1016/j.celrep.2020.108156
Filimontseva A, Cuadros T, Chatterton Z, Burke A, Mahabadian AA, Compte J, Fu Y, Vila M, Halliday
GM (2025) Standardised TruAI Automated Quantification of Intracellular Neuromelanin
Granules in Human Brain Tissue Sections. Neuropathol Appl Neurobiol 51 (4):e70033.
doi:10.1111/nan.70033
Fu Y, Yuan Y, Halliday G, Rusznak Z, Watson C, Paxinos G (2012) A cytoarchitectonic and
chemoarchitectonic analysis of the dopamine cell groups in the substantia nigra, ventral tegmental
area, and retrorubral field in the mouse. Brain Struct Funct 217 (2):591-612. doi:10.1007/s00429-
011-0349-2
Fudge JL, Kelly EA, Love TM (2024) Amygdalo-nigral inputs target dopaminergic and GABAergic
neurons in the primate: a view from dendrites and soma. bioRxiv. doi:10.1101/2024.01.16.575910
Halliday GM, Li YW, Blumbergs PC, Joh TH, Cotton RG, Howe PR, Blessing WW, Geffen LB (1990a)
Neuropathology of immunohistochemically identified brainstem neurons in Parkinson's disease.
Ann Neurol 27 (4):373-385. doi:10.1002/ana.410270405
Halliday GM, Li YW, Blumbergs PC, Joh TH, Cotton RGH, Howe PRC, Blessing WW, Geffen LB
(1990b) Neuropathology of immunohistochemically identified brainstem neurons in Parkinson's
disease. Annals of Neurology 27 (4):373-385. doi:https://doi.org/10.1002/ana.410270405
Hardman CD, Henderson JM, Finkelstein DI, Horne MK, Paxinos G, Halliday GM (2002) Comparison of
the basal ganglia in rats, marmosets, macaques, baboons, and humans: volume and neuronal
number for the output, internal relay, and striatal modulating nuclei. J Comp Neurol 445 (3):238-
255. doi:10.1002/cne.10165
Henny P, Brown MTC, Northrop A, Faunes M, Ungless MA, Magill PJ, Bolam JP (2012) Structural
correlates of heterogeneous in vivo activity of midbrain dopaminergic neurons. Nature
Neuroscience 15 (4):613-619. doi:10.1038/nn.3048
Hjelmstad GO, Xia Y, Margolis EB, Fields HL (2013) Opioid Modulation of Ventral Pallidal Afferents to
Ventral Tegmental Area Neurons. The Journal of Neuroscience 33 (15):6454.
doi:10.1523/JNEUROSCI.0178-13.2013
Howat WJ, Wilson BA (2014) Tissue fixation and the effect of molecular fixatives on downstream
staining procedures. Methods 70 (1):12-19. doi:https://doi.org/10.1016/j.ymeth.2014.01.022
Jhou TC (2021) The rostromedial tegmental (RMTg) "brake" on dopamine and behavior: A decade of
progress but also much unfinished work. Neuropharmacology 198:108763.
doi:10.1016/j.neuropharm.2021.108763
Jhou TC, Fields HL, Baxter MG, Saper CB, Holland PC (2009a) The Rostromedial Tegmental Nucleus
(RMTg), a GABAergic Afferent to Midbrain Dopamine Neurons, Encodes Aversive Stimuli and
Inhibits Motor Responses. Neuron 61 (5):786-800.
doi:https://doi.org/10.1016/j.neuron.2009.02.001
Jhou TC, Geisler S, Marinelli M, Degarmo BA, Zahm DS (2009b) The mesopontine rostromedial
tegmental nucleus: A structure targeted by the lateral habenula that projects to the ventral
tegmental area of Tsai and substantia nigra compacta. J Comp Neurol 513 (6):566-596.
doi:10.1002/cne.21891

Kaufling J, Veinante P, Pawlowski SA, Freund-Mercier MJ, Barrot M (2009) Afferents to the GABAergic
tail of the ventral tegmental area in the rat. J Comp Neurol 513 (6):597-621.
doi:10.1002/cne.21983
Kelly EA, Contreras J, Duan A, Vassell R, Fudge JL (2022) Unbiased Stereological Estimates of
Dopaminergic and GABAergic Neurons in the A10, A9, and A8 Subregions in the Young Male
Macaque. Neuroscience 496:152-164. doi:https://doi.org/10.1016/j.neuroscience.2022.06.018
Lahti L, Haugas M, Tikker L, Airavaara M, Voutilainen MH, Anttila J, Kumar S, Inkinen C, Salminen M,
Partanen J (2016) Differentiation and molecular heterogeneity of inhibitory and excitatory
neurons associated with midbrain dopaminergic nuclei. Development 143 (3):516-529.
doi:10.1242/dev.129957
Lammel S, Lim BK, Ran C, Huang KW, Betley MJ, Tye KM, Deisseroth K, Malenka RC (2012) Input-
specific control of reward and aversion in the ventral tegmental area. Nature 491 (7423):212-217.
doi:10.1038/nature11527
Lerner Talia N, Shilyansky C, Davidson Thomas J, Evans Kathryn E, Beier Kevin T, Zalocusky Kelly A,
Crow Ailey K, Malenka Robert C, Luo L, Tomer R, Deisseroth K (2015) Intact-Brain Analyses
Reveal Distinct Information Carried by SNc Dopamine Subcircuits. Cell 162 (3):635-647.
doi:https://doi.org/10.1016/j.cell.2015.07.014
Liu B, White AJ, Lo CC (2025) Augmenting flexibility: mutual inhibition between inhibitory neurons
expands functional diversity. iScience 28 (2):111718. doi:10.1016/j.isci.2024.111718
Margolis EB, Toy B, Himmels P, Morales M, Fields HL (2012) Identification of Rat Ventral Tegmental
Area GABAergic Neurons. PLOS ONE 7 (7):e42365. doi:10.1371/journal.pone.0042365
McRitchie DA, Cartwright HR, Halliday GM (1997) Specific A10 Dopaminergic Nuclei in the Midbrain
Degenerate in Parkinson's Disease. Experimental Neurology 144 (1):202-213.
doi:https://doi.org/10.1006/exnr.1997.6418
McRitchie DA, Halliday GM, Cartwright H (1995) Quantitative analysis of the variability of substantia
nigra pigmented cell clusters in the human. Neuroscience 68 (2):539-551. doi:10.1016/0306-
4522(95)00163-D
Moaddab M, McDannald MA (2021) Retrorubral field is a hub for diverse threat and aversive outcome
signals. Curr Biol 31 (10):2099-2110 e2095. doi:10.1016/j.cub.2021.02.055
Morello F, Partanen J (2015) Diversity and development of local inhibitory and excitatory neurons
associated with dopaminergic nuclei. FEBS Letters 589 (24PartA):3693-3701.
doi:https://doi.org/10.1016/j.febslet.2015.10.001
Nair-Roberts RG, Chatelain-Badie SD, Benson E, White-Cooper H, Bolam JP, Ungless MA (2008)
Stereological estimates of dopaminergic, GABAergic and glutamatergic neurons in the ventral
tegmental area, substantia nigra and retrorubral field in the rat. Neuroscience 152 (4):1024-1031.
doi:https://doi.org/10.1016/j.neuroscience.2008.01.046
Nieh Edward H, Matthews Gillian A, Allsop Stephen A, Presbrey Kara N, Leppla Christopher A,
Wichmann R, Neve R, Wildes Craig P, Tye Kay M (2015) Decoding Neural Circuits that Control
Compulsive Sucrose Seeking. Cell 160 (3):528-541. doi:10.1016/j.cell.2015.01.003
Omelchenko N, Sesack SR (2010) Periaqueductal gray afferents synapse onto dopamine and GABA
neurons in the rat ventral tegmental area. Journal of Neuroscience Research 88 (5):981-991.
doi:https://doi.org/10.1002/jnr.22265
Paxinos G, Furlong T, Watson C (2020) Human brainstem: cytoarchitecture, chemoarchitecture,
myeloarchitecture. Academic Press,
Rademacher K, Doric Z, Haddad D, Mamaligas A, Liao SC, Creed R, Kano K, Chatterton Z, Fu Y, Garcia
JH, Vance VM, Sei YJ, Kreitzer A, Halliday G, Nelson AB, Margolis E, Nakamura K (2025)
Chronic hyperactivation of midbrain dopamine neurons causes preferential dopamine neuron
degeneration. Elife 13. doi:10.7554/eLife.98775
Simmons DA, Hoffman NW, Yahr P (2011) A forebrain-retrorubral pathway involved in male sex
behavior is GABAergic and activated with mating in gerbils. Neuroscience 175:162-168.
doi:10.1016/j.neuroscience.2010.11.048

Smith RJ, Vento PJ, Chao YS, Good CH, Jhou TC (2019) Gene expression and neurochemical characterization of the rostromedial tegmental nucleus (RMTg) in rats and mice. Brain Structure and Function 224 (1):219-238. doi:10.1007/s00429-018-1761-7

St. Laurent R, Martinez Damonte V, Tsuda AC, Kauer JA (2020) Periaqueductal Gray and Rostromedial Tegmental Inhibitory Afferents to VTA Have Distinct Synaptic Plasticity and Opiate Sensitivity. Neuron 106 (4):624-636.e624. doi:https://doi.org/10.1016/j.neuron.2020.02.029

Vincek V, Nassiri M, Nadji M, Morales AR (2003) A Tissue Fixative that Protects Macromolecules (DNA, RNA, and Protein) and Histomorphology in Clinical Samples. Laboratory Investigation 83 (10):1427-1435. doi:https://doi.org/10.1097/01.LAB.0000090154.55436.D1

Vounatsos MV, Gittis AH (2025a) GPe Projections to the Retrorubral Field Give Rise to Diverging GABAergic and DAergic Circuits. J Neurosci 45 (2). doi:10.1523/JNEUROSCI.1446-24.2024

Vounatsos MV, Gittis AH (2025b) GPe Projections to the Retrorubral Field Give Rise to Diverging GABAergic and DAergic Circuits. The Journal of Neuroscience 45 (2):e1446242024. doi:10.1523/JNEUROSCI.1446-24.2024

Statements & Declarations

Funding

This work was supported in part by Aligning Science Across Parkinson's [020505] through the Michael J. Fox Foundation for Parkinson's Research (MJFF) and also by funding from the National Health and Medical Research Council of Australia to GMH [Investigator Grant 1176607]. For open access, the authors have applied a CC BY public copyright license to all Author Accepted Manuscripts arising from this submission.

Competing interests

The authors declare no competing interests.

Author contributions

All authors contributed to the study conception and design. Material preparation and data collection were performed by Anastasia Filimontseva and analysis were performed by all authors. The first draft of the manuscript was written by Anastasia Filimontseva and all authors commented on previous versions of the manuscript. All authors read and approved the final manuscript.

Data availability

The data and analysis images and key lab materials used and generated in this study are listed in a Key Resource Table alongside their persistent identifiers at Zenodo DOI 10.5281/zenodo.17946370.

583 **Ethics approval**

584 Tissue was obtained from the Sydney Brain Bank (UNSW Sydney Human Research Ethics

585 Committee approval HC200026) and the study approved by the University of Sydney Human

586 Research Ethics Committee (2021/845).