Characterising the detectable and invisible fractions of genomic loci under balancing selection

Débora Y. C. Brandt[1,2,§], Aida M. Andrés[1,2], Tim Connallon[4,*] and Max Reuter[1,3,*,§]

[1]Research Department of Genetics, Evolution and Environment, University College London, Gower Street, London WC1E 6BT, United Kingdom

[2]UCL Genetics Institute, University College London, Gower Street, London WC1E 6BT, United Kingdom

[3]Centre of Life Origins and Evolution, University College London, Gower Street, London WC1E 6BT, United Kingdom

[4]School of Biological Sciences, Monash University, 25 Rainforest Walk, Clayton, Victoria 3800, Australia

[*]Equal contribution

[§]Corresponding authors: Débora Y. C. Brandt (d.brandt@ucl.ac.uk), and Max Reuter (m.reuter@ucl.ac.uk)

Abstract

Balancing selection refers to scenarios where selection maintains genetic polymorphisms affecting fitness and its components. Empirical studies have documented specific cases of balancing selection, yet the general prevalence of balanced polymorphisms within genomes remains a topic of longstanding debate. Although genome-wide scans for signals of balancing selection suggest that it is rare, current methods are notoriously conservative and may identify only a small and unrepresentative fraction of loci evolving under balancing selection. What, then, are the proportions of loci under balancing selection that are detectable versus invisible to genome scans? Here, we address this question using a combination of analytical modelling and population genetic simulations. Our results provide quantitative support for the intuition that a large fraction of loci under balancing selection will be invisible to standard tests for balancing selection, with the detectable loci representing a biased subset with large and symmetrical fitness effects on different fitness components. Quantifying power across parameters that vary between species also shows that the detectable and invisible fractions of genomic loci under balancing selection are likely to differ substantially between organisms. For example, a *Drosophila melanogaster*-like ratio of mutation to recombination rates (~0.1) reduces detection power by roughly a quarter of the power expected with a human-like ratio (~1). This, combined with other evidence suggesting that balancing selection could be common in *D. melanogaster*, showcases how our appreciation of the prevalence of balancing selection is limited by our ability to detect its genomic signals, especially in those species where it might be common.

Introduction

Balancing selection occurs when natural selection actively favours the maintenance of genetic polymorphism. This can arise from several specific contexts of selection, including intrinsic fitness advantages of being heterozygous (*i.e.*, 'heterozygote advantage' or fitness 'overdominance'; (Fisher, 1922)), genetic trade-offs between different fitness contexts or environments (*e.g.*, 'antagonistic pleiotropy' between fitness components: (Curtsinger et al., 1994); 'sexually antagonistic' trade-offs between sexes: (Kidwell et al., 1977)); trade-offs between temporal or spatially varying environments (Haldane & Jayakar, 1963; Levene, 1953), or negative-frequency-dependent selection, where genotypes become more favourable when rare (Fitzpatrick et al., 2007; Wright, 1942).

Loci under balancing selection are expected to be more polymorphic than loci evolving neutrally or under directional selection (though exceptions can occur when selection is weak or allele frequency equilibria under balancing selection are close to zero or one, (Robertson, 1962; Williamson et al., 2004)). When equilibrium frequencies are intermediate (*e.g.*, very roughly between 0.2 and 0.8; (Hedrick, 1999)) and balancing selection is strong relative to genetic drift, the balanced polymorphisms should persist far longer than neutrally evolving polymorphisms (Nei & Roychoudhury, 1973), potentially over timescales spanning speciation events (Klein et al., 1998). Whatever their age, balanced polymorphisms should contribute disproportionately to standing genetic variation for fitness or fitness components (*e.g.*, for survival, fertility, or longevity), with even modest numbers of such loci contributing substantially to fitness variation within populations (Charlesworth, 2015; Connallon & Czuppon, 2026; Crow, 1952, 1987; Zajitschek & Connallon, 2018).

The importance of balancing selection has, nevertheless, been a subject of longstanding debate in evolutionary biology. As an early and vociferous proponent, Dobzhansky (1955) argued that selection should often preserve genetic variation due to the greater versatility of heterozygous, relative to homozygous individuals in perpetually changing environments. Yet, by the close of the 20th Century, high levels of molecular genetic diversity were reported by studies based first on allozymes (Lewontin & Hubby, 1966; Lewontin, 1974) and subsequently DNA sequences (Kreitman, 1983). The pervasiveness of these polymorphisms made Dobzhansky's 'balance hypothesis' an implausible general explanation for molecular genomic diversity. Instead, the emerging neutral theory (Kimura, 1968; King & Jukes, 1969; Kimura, 1983) provided a simpler and more powerful explanation for the ubiquity of molecular genetic

diversity, effectively sidelining balancing selection as irrelevant for all but a few exceptional
loci within the genome (see p. 199 of Lewontin (1974); Hedrick (2012)).

More recent developments in evolutionary genetics have, however, prompted a
resurgence in enthusiasm for balancing selection. Firstly, the coupling of field studies with
genomics has identified exceptionally clear cases of balancing selection that maintains
conspicuous phenotypic polymorphisms. Beyond classic examples like sickle cell anaemia
(Allison, 1954), modern case studies include a pleiotropic trade-off between viability and
mating success that maintains horn size polymorphism in Soay sheep (mapped to the *RXFP2*
gene; Johnston et al. (2013)), a sexually antagonistic polymorphism in cuticular hydrocarbons
of *Drosophila serrata* (in the *DsFAR2-B* gene; Rusuwa et al. (2022)), alleles affecting age of
maturation in Atlantic salmon (the *VGLL3* gene; Barson et al. (2015)), temporally varying
selection that maintains genetic polymorphism for insecticide resistance in *Drosophila*
melanogaster (Karageorgi et al., 2025), and many others (reviewed in Bitarello *et al.* (2023);
Ruzicka *et al.* (2025)). Second, the advent of high-throughput sequencing techniques and
sophisticated analytical methods for detecting population genetic signatures of balancing
selection has provided new means of identifying candidate loci under long-term balancing
selection (reviewed in Bitarello et al., 2023; Fijarczyk & Babik, 2015). Applied to the human
genome, such methods have provided strong evidence for long-term balancing selection in both
African and European populations (Bitarello et al., 2018). Third, detailed lab studies of natural
genetic variation in life-history traits suggest that *Drosophila* populations harbour higher levels
of genetic variation than can be explained by models of mutation-selection balance
(Charlesworth, 2015; Charlesworth & Hughes, 2000; Sharp & Agrawal, 2018). Balancing
selection is a prime candidate to account for this apparent excess of genetic variation in life-
history traits. If the same is true in other species, balancing selection may maintain substantial
amounts of genetic and trait variation in natural populations.

While recent genome scans have identified putative balanced polymorphisms, it
remains unclear how accurately they capture the true extent of balancing selection. The
typically small fraction of loci with statistically significant signals of balancing selection (*e.g.*,
<1% of the genome in humans (Bitarello et al., 2018)) might reflect either the rarity of
balancing selection across the genome or methodological limitations to detecting it. Most
notably, classical methods cannot detect balancing selection unless intermediate-frequency
polymorphisms are maintained over exceptionally long timescales. Loci segregating for
evolutionarily 'young' balanced polymorphisms cannot easily be identified through population
genomic approaches alone (but see Isildak et al. (2021) for a potentially promising approach),

and old polymorphisms can likewise be missed due to limited data quality or other attributes
that vary among species (*e.g.*, mutation and recombination rates; effective population sizes)
and may influence the strength of balancing selection signals. The balanced polymorphisms
that are detected are, therefore, likely to represent only a minor, potentially unrepresentative,
subset of loci evolving under balancing selection. In fact, a recent genome-wide analysis
suggests there are hundreds of non-synonymous polymorphisms shared among human
populations due to recent balancing selection, even though the method can only detect an
aggregate signal across many loci rather than identify individual targets of balancing selection
(Soni et al., 2022). While it remains an open question what proportion of these loci ultimately
results in long-term stable balanced polymorphisms that are individually identifiable, this
analysis suggests that at least recent episodes of balancing selection are widespread in the
human genome.

Here, we aim to put the debate over the prevalence of balancing selection on a sounder
conceptual and quantitative footing by evaluating factors affecting opportunities for long-term
balancing selection and its detectability. It is obviously impossible to make general predictions
about the proportion of a genome that is a potential target of balancing selection. Yet it is
possible to assess—under a given model of balancing selection—how the population genetic
parameters of genetic drift and selection will affect: *i*) the long-term maintenance of
polymorphisms, and *ii*) the proportion of loci that develop signatures of balancing selection
that are clear enough to be detectable using contemporary population genomic methods.

We address these questions while considering three mechanisms of balancing selection:
overdominance, antagonistic pleiotropy and sexually antagonistic selection, each of which
have been empirically documented in natural populations (e.g. Allison, 1954; Johnston et al.,
2013; Rusuwa et al., 2022). These mechanisms represent a subset of balancing selection
mechanisms (e.g. Glémin, 2021; Prout, 2000; Ruzicka et al., 2025), yet their evolutionary
dynamics are representative of a broader array of balancing selection scenarios, which allows
our predictions to be generalized. We begin by briefly reviewing predictions from population
genetics theory, highlighting constraints associated with maintaining long-term stable
polymorphisms under each mechanism of balancing selection. We then present simulations
under each mechanism that explore the genomic signals associated with long-term balanced
polymorphisms and assess how these signals translate into statistical power to infer the
presence of balancing selection at these loci. Our analyses reveal how the maintenance of
balanced polymorphisms and their detectability depend on the properties of the genetic
polymorphism itself, as well as the genetic and demographic features of the species. They also

provide a quantitative appreciation for the extent to which loci subject to balancing selection
are detectable and suggest the biases in the specific types of balanced polymorphisms that are
most and least likely to be identified.

Results

Long-term maintenance of balanced polymorphisms: Theoretical considerations
We start by reviewing theoretical predictions regarding the conditions under which genetic
polymorphisms are stably maintained due to overdominant selection (OD), antagonistic
pleiotropy (AP) or sexually antagonistic selection (SA). Our models focus on bi-allelic, diploid
loci, and we assume that individuals mate at random and selection at each locus remains
constant over time. Violations of these assumptions tend to restrict conditions for balancing
selection (reviewed in Ruzicka et al. (2025), but see Wittmann et al. (2023) for signals of long-
term balancing selection under temporally fluctuating selection). Full details of the models are
presented in Appendix 1.

Long-term maintenance of a polymorphism under these models of selection requires
two conditions. First, the combination of allelic fitness effects must actually generate balancing
selection. Second, the resulting selection must be stronger than genetic drift for the
polymorphism to be stably maintained over time. We deal with each condition in turn.

As far as the general criteria for balancing selection are concerned, OD is the most
permissive scenario. In outbred populations, heterozygotes having higher fitness than both
homozygotes (OD) invariably leads to balancing selection (Fisher, 1922). In contrast, AP and
SA lead to balancing selection under only a subset of parameter conditions (Fig. 1). AP and
SA each involve trade-offs between the fitness effects of alleles in different contexts of
selection (between fitness components and between sexes, respectively). Considering a locus
with alleles A_1 and A_2, s_1 represents the homozygous cost associated with the A_1 allele, s_2 is the
homozygous cost associated with the A_2 allele, and dominance (h) defines the degree to which
these costs manifest in heterozygotes (Fig. 1A). For simplicity, we focus on cases in which the
dominance coefficient ranges from $h = 0.5$, implying co-dominance of the alleles, to $h = 0$,
implying complete recessivity of these costs (for generalizations beyond these conditions, see
Kidwell et al. (1977); Curtsinger et al. (1994)). Note that dominance values within the range 0
$< h < 0.5$ correspond to "dominance reversals" (Connallon & Chenoweth, 2019; Grieshop et

al., 2024; Reid, 2022), where each allele is (partially) recessive in contexts where it is harmful and (partially) dominant in contexts where it is favourable.

Balancing selection arises consistently in the SA and AP scenarios when the homozygous costs of each allele are symmetric in magnitude (s_1 is roughly equal to s_2), and the criteria for balancing selection expand with both the strength of selection and the strength of the dominance reversal (i.e., with s_1 and s_2 increasing and h decreasing; Fig. 1B). When selection is weak and there is no dominance reversal, only a small proportion of the parameter space will generate balancing selection. In this case, the proportion of random parameter combinations that leads to balancing selection is roughly $f_{bal} \approx \frac{2}{3}\bar{s}$ under AP and $f_{bal} \approx \frac{4}{3}\bar{s}$ under SA, where $\underline{s}$ represents the average fitness effect ($\bar{s} = \frac{s_{max}}{2}$ for parameters drawn from uniformly from the parameter space, with s_{max} representing the maximum selection coefficient; $s_{max} \ll 1$ when selection is consistently weak). With dominance reversals ($h < 0.5$) and weak selection, the proportions expand to approximately $f_{bal} \approx \frac{1-2h}{1-h}$ for both AP and SA, which corresponds to $f_{bal} \approx \frac{2}{3}$ when $h = 0.25$ (as in Fig. 1B) and $f_{bal} = 1$ for a complete dominance reversal ($h = 0$).

Figure 1. Conditions for balancing selection under antagonistic pleiotropy (AP) and sexually antagonistic selection (SA). **Panel A:** Visual summary of the AP and SA model parameters, with the example showing $s_1 = 0.8$, $s_2 = 0.6$, and $h = 0.5$. The two bar charts show the relative fitness of each of the three genotypes of a locus across two contexts of selection (i.e., different fitness components or sexes), where the A_1 allele is selectively disfavoured in one context and the A_2 allele is disfavoured in the other. **Panel B:** Criteria for balancing selection as a function of selection (s_1 and s_2) and dominance (h) parameters. The regions between each pair of curves (based on

colour and line type) show conditions where selection favours the maintenance of

polymorphism (i.e., where there is balancing selection). Predictions are shown when

there is no dominance between alleles (i.e., alleles are co-dominant: $h = 0.5$), and

when there is a beneficial reversal of dominance ($h = 0.25$), where negative allelic

effects are partially recessive to beneficial effects. Panel B is modified from

Connallon and Chenoweth (2019), with curves based on standard modelling results

(see Kidwell et al. 1977; Curtsinger et al. 1994).

203   The second condition for long-term stability of a balanced polymorphism is that

selection towards a polymorphic equilibrium (which we denote as $\hat{p}$) must be strong enough to

prevent fixation of one or the other allele by genetic drift. The required strength of selection

depends on the effective size of the population (N_e) and the net strength of selection to the

equilibrium (α), which differs between the mechanisms of balancing selection ($\alpha = s_1 + s_2$

under the OD model, where s_1 and s_2 represent the fitness reductions of A_1 and A_2 homozygotes

relative to the heterozygotes; $\alpha = (s_1 + s_2)(1 - 2h) + 2s_1 s_2 h^2$ and $\alpha = \frac{1}{2}(s_1 + s_2)(1 -$

$2h) + 2s_1 s_2 h^2$ under the AP and SA models, respectively, with the SA and AP parameters, s_1,

s_2, and h, as depicted in Fig. 1A; see Appendix 2 for derivations of the equations for α).

212   The stationary distribution of the allele frequencies for a locus under balancing

selection and drift (*i.e.*, the predicted long-term probability distribution of allele frequency

states) provides an intuitive measure of balancing selection's ability to maintain polymorphism

in the presence of genetic drift (Rice, 2004; Robertson, 1962). Balancing selection reliably

maintains polymorphism when the allele frequency probability densities are much greater near

the deterministic equilibrium ($\hat{p}$) than near the boundaries (*i.e.*, close to zero and one).

Heuristically, balancing selection can be regarded as effectively strong in cases where the

probability density around $\hat{p}$ is at least ten times greater than the density near the boundaries.

Under this definition, balancing selection is effectively strong when $\alpha > a_{\min}$, where:

221
$$a_{\min} = \begin{cases} \dfrac{\log(20N_e\hat{p})}{2N_e\hat{p}^2} & \text{for } \hat{p} \le 0.5 \\ \dfrac{\log(20N_e(1-\hat{p}))}{2N_e(1-\hat{p})^2} & \text{for } \hat{p} \ge 0.5 \end{cases} \tag{1}$$

Figure 2A illustrates how the effective population size and equilibrium allele frequencies under

balancing selection (N_e and $\hat{p}$) interact to determine criteria for effectively strong balancing

selection (where $\alpha > a_{\min}$). As can be seen in the figure, these conditions are most permissive

in large populations with intermediate equilibria (note that $a_{\min}$ is always minimized at $\hat{p} =$
0.5).

Because α is defined differently for each model of selection (see above), thresholds for
effectively strong balancing selection correspond to different absolute strengths of selection
under the AP, SA, and OD scenarios. For example, the minimum values of s_1 and s_2 that are
required for $\alpha > a_{\min}$ (shown in Fig. 2B) are largest for the SA model, smallest for the OD
model, and intermediate for the AP model (SA and AP cases are equal when $h = 0.5$, and AP
and OD cases are equal when $h = 0$; see Fig. 2B).

A reduction of N_e amplifies drift and restricts the parameter space of effectively strong
balancing selection (Fig. 2C), particularly so when dominance reversals are weak or absent.
For example, when $h = 0.5$ and $N_e = 2{,}000$, SA and AP alleles cannot be reliably maintained
polymorphic even by moderately strong selection (e.g., $s_1 \sim s_2 \sim 0.05$, corresponding to a 5%
difference between the homozygote genotypes in each fitness component). Larger N_e and/or
dominance reversals ($h < 0.5$) make long-term maintenance of polymorphism much more likely
(Fig. 2C). Population genetic simulations (see Methods for simulation details) confirm these
general predictions (Figs. S2, S4, S6). They further illustrate how the notion of maintaining
long-term stable polymorphisms depends on the timescale of the observation: the range of
parameter values that maintains polymorphism for $16N_e$ generations is significantly narrower
than the range that maintains polymorphism for $8N_e$ generations (Fig. S2C,D).

Figure 2. Effectively strong balancing selection. **Panel A** illustrates the thresholds
for effectively strong balancing selection relative to genetic drift ($a_{\min}$; see eq. (1)
and the surrounding text) for three population sizes (N_e) and a spectrum of allele
frequency equilibrium conditions under balancing selection ($0 < \hat{p} < 1$). Regions
above each curve denote combinations of α and $\hat{p}$ that generate effectively strong

balancing selection (i.e., where the probability density for the stationary distribution is at least a 10-fold higher near the polymorphic equilibrium, $\hat{p}$, than near the allele frequency boundaries of zero and one, which correspond to fixation). Two special cases of the stationary distribution are shown: one where selection is ineffective ($\alpha < a_{min}$, and density is concentrated near *zero*) and one where selection is effective ($\alpha > a_{min}$, and density is concentrated near the equilibrium, $\hat{p} = 0.4$). **Panel B** shows the minimum selection coefficients for each model that allow for effective balancing selection (the minimum is defined at $\hat{p} = 0.5$, where conditions for effective balancing selection are most permissive). **Panel C** shows the proportion of loci under AP balancing selection that meet the criteria for effectively strong balancing selection. Results are based on randomly drawing 10^7 sets of selection coefficients (s_1 and s_2) from independent uniform distributions between 0 and s_{max}, where $\bar{s} = \frac{1}{2}s_{max}$ is the average selection coefficient. Bar heights were calculated as the number of parameter sets that met the criterion for effectively strong balancing selection ($0 < \hat{p} < 1$ and $\alpha > a_{min}$), divided by the number of parameter sets that generated balancing selection ($0 < \hat{p} < 1$).

Empirical signatures of long-term balancing selection

The theory presented above summarizes factors that promote or constrain the maintenance of polymorphism at individual sites subject to balancing selection. Assessing the detectability of balancing selection requires additional considerations, because genomic scans for balancing selection do not focus on the balanced site itself but on signals of elevated genetic diversity at sites physically linked to the selected site. These signatures build up in the presence of long-term balancing selection, where the balanced polymorphism segregates for much longer than expected under neutrality (*i.e.*, the time to the most recent common ancestor, TMRCA, at the selected site is greater than the neutral expectation of $4N_e$ generations). Genomic regions with an old TMRCA owing to balancing selection are expected to deviate from the genomic background in two ways. First, they are expected to show an elevated density of polymorphisms (and a concomitant deficit of substitutions relative to closely related outgroups) due to the long timespans over which mutations can accumulate since the TMRCA in the region and their lower probability of going to fixation due to linkage with the balanced site (DeGiorgio et al., 2014). Second, they are expected to show an excess of intermediate-frequency alleles

when compared with the neutral expectation, as linked neutral variants tend to segregate at

frequencies similar to those of the balanced polymorphism (see e.g., Fig. 1 in Bitarello et al.

(2018)). These properties will be reflected in the local site-frequency spectrum (SFS) and

provide the signal exploited by current population genomic methods for detecting long-term

balancing selection (e.g. Bitarello et al., 2018; Cheng & DeGiorgio, 2020).

In the following section, we will explore how the presence and intensity of the signal,

and thus our statistical power to identify regions under balancing selection, depends on the

properties of the balanced polymorphism itself (*i.e.*, the nature and strength of balancing

selection, dominance, and the resulting equilibrium frequency of the balanced polymorphism),

and factors that affect population genomic data, including the effective population size, and

mutation and recombination rates, which vary among species and genomic regions.

We can get a preliminary sense of some of the key factors affecting neutral diversity

near a long-term balanced polymorphism by quantifying the expected spike in heterozygosity

near a balanced site under strong selection, which is analytically tractable. Consider a balanced

polymorphism with alleles A_1 and A_2 maintained stably at deterministic equilibrium

frequencies, $\hat{p}$ and $1 - \hat{p}$, respectively. Building on previous work (see p. 103 in Rice (2004)),

one can derive the expected width of the sequence surrounding the balanced site in which

neutral polymorphisms will have an expected level of heterozygosity that is at least K times

greater than the expected heterozygosity at unlinked neutral sites elsewhere in the genome

($4N_e\mu$, where μ is mutation rate per nucleotide site). The width of this region, in kilobases (kb),

is:

306
$$L_K \approx \frac{2\hat{p}(1 - \hat{p})}{10^3(K - 1)N_e r} \tag{2}$$

where r represents the recombination rate per nucleotide site (see Appendix 3). Inspection of

eq. (2) shows that the width of elevated diversity surrounding the balanced polymorphism

increases with the equilibrium heterozygosity of the selected site (with $2\hat{p}(1 - \hat{p})$), which is

maximized when balanced alleles are maintained at equal frequencies ($\hat{p} = 0.5$), and it

decreases with the population-scaled recombination rate per nucleotide site ($N_e r$). Thus, we

expect empirical signals of long-term balancing selection to be strongest when $\hat{p}$ is intermediate

and $N_e r$ is small, owing to small N_e and/or small r.

Power to detect balancing selection in simulated data

While these predictions lend intuition, they ignore the reality that allele frequencies at the

selected site fluctuate over time (*e.g.,* due to drift), and overlook other empirically distinctive

features of the SFS that are relevant to balancing selection scans, including the actual number of segregating linked variants and the shape of the SFS around a balanced site relative to the genome as a whole. Allele frequency fluctuations at the balanced polymorphic site are expected to reduce linked diversity relative to the more idealized case (as in eq. (2)) where the alleles are maintained at invariant equilibrium frequencies (for allele frequency fluctuations caused by temporally varying selection, see Wittmann et al. (2023)). Drift-mediated allele frequency fluctuations at the selected site should be minimal provided $N_e\alpha$ is large and the equilibrium ($\hat{p}$) is intermediate, while fluctuations increase as $N_e\alpha$ and $\hat{p}(1-\hat{p})$ decline (Charlesworth & Charlesworth, 2010, pp. 354–355; Connallon & Czuppon, 2026).

To provide an assessment of statistical power for detecting balancing selection candidates, we turned to population genetic simulations. We ran extensive simulations (see Methods for details) with population-scaled values of $N_e\mu$ and $N_e r$ chosen to roughly match those of humans and fruit flies—intensively studied species that may be representative of vertebrates and insects. We note that these simulations are not intended to provide specific power estimates for these species, since our models do not match their demographic histories (for example, the long-term N_e of human populations is smaller than 20,000 and their detailed demographic history generally increases power compared with the stable effective population size simulated here see Discussion). Rather, they are presented as toy examples that allow us to evaluate effects of relevant genomic parameters that vary among species. We first assessed power for a human-like 'reference' set of parameters ($N_e = 20,000$, $\mu = r = 10^{-8}$, age of polymorphism $T = 8N_e = 160,000$ generations), which we will later use as a point of comparison with simulations using different population parameters. We assessed the power of the NCD method (Bitarello et al., 2018) to identify balancing selection in population genetic simulations of 10Kb chromosomes with a centrally located site under balancing selection and neutral variation in the flanking regions (see Methods). The results obtained for NCD should be representative of those obtained with other contemporary methods for detecting long-term balancing selection, which use similar approaches; additional results using the BalLeRMix method (Cheng & DeGiorgio, 2020) are presented in the Supplementary Figures S11 and S12 (and see Fig. S13 for a comparison between the two methods). NCD tests for an enrichment of polymorphism in the SFS by calculating, across all SNPs in a window, the average deviation of allele frequency from an intermediate 'target frequency'. We used the equilibrium frequency of the balanced site as the target frequency for NCD, and our analyses therefore represent an idealised best-case scenario for power, where the true equilibrium frequency is known (Bitarello et al., 2018). For each set of parameters, we generated 100 replicate simulations in

which the balanced polymorphism segregated until the end of the simulation ($8N_e$ or $16N_e$

generations, thus conditioning on the maintenance of polymorphism; for results on the

probability of maintenance over these time durations, see Figs. S1–S6). From these, we

estimated power as the proportion of simulations that resulted in a statistically significant signal

of balancing selection at $P < 0.01$, measured by NCD.

We evaluated the effects of our simulation parameters on power through multiple

logistic regressions (Generalised Linear Models (GLM) with binomial error) with counts of

successful and unsuccessful detection among the 100 trials of our simulations as dependent

variables. The GLMs model power on the log-odds scale and allow us to estimate and test the

effects of the simulation parameters while taking into account the fact that power is bounded

between 0 and 1 as well as the binomial error of successful/unsuccessful detection of balancing

selection (see Methods for more details).

In our simulations, the power to detect balancing selection is relatively low overall for

our set of parameters, regardless of the exact method used for detection (i.e., NCD in Fig. 3

and Figs. S7–S9 versus BalLeRMix in Figs. S10–13). However, power varied considerably

across parameter conditions in ways that broadly align with theoretical expectations. Among

our human-like reference simulations, the variable that most strongly predicts power is the

equilibrium frequency of the balanced polymorphism ($\hat{p}$), with the likelihood of detecting

balancing selection increasing as the equilibrium approaches an intermediate frequency (i.e.,

$\hat{p} \to 0.5$; Fig. 3A). Across these reference simulations, the average power is 0.23 at $\hat{p} = 0.5$,

while power drops to 0.02 with $\hat{p} = 0.07$. Fitting a binomial GLM with equilibrium frequency

($\hat{p}$), selection strength (α), selection model and their interactions as predictors of detection

power, we found that the equilibrium frequency explains more than 60% of the variance in

power across our simulations (Tab. 1). More specifically, the odds of successfully detecting

balancing selection are estimated to increase by more than 30% with every 10% increase in the

equilibrium frequency of the minor allele at the selected site, up to $\hat{p} = 0.5$ ($b_{\text{eq. freq.}} = 2.737$ on

log-odds scale, estimated for OD).

Power also generally increases with the strength of selection (Tab. 1), but this effect

depends on interactions with other parameters. First, the positive effect of selection strength on

power is pronounced at intermediate equilibrium frequencies but declines and eventually

vanishes when equilibrium frequencies of the minor allele at the selected site are lower (Tab.

1, Fig. 3B). Second, this interaction effect differs between the three models of selection and is

strongest under SA. Under OD and AP, increasing selection strength by 0.1 increases the odds

of detecting balancing selection by 5.7% and 1.6% for equilibrium frequencies of $\hat{p} = 0.5$ and $\hat{p} = 0.25$, respectively. For SA, in contrast, the same change in selection strength increases the odds of detection by 15.7% for an equilibrium frequency of $\hat{p} = 0.5$ but decreases the odds by 40% for an equilibrium of $\hat{p} = 0.25$ (see Fig. 3B for simulation data and Fig. 4 for model predictions). Beyond these interactive effects, however, the model of selection does not impact the overall power of detecting balancing selection (Tab. 1, Fig. 3C).

Figure 3. Power as a function of equilibrium frequency ($\hat{p}$, panel A), selection strength (α, panel B), and model of selection (panel C) for the reference simulations. On panels A and C points were horizontally jittered for visualization purposes. The three panels show the same data points, with power on the vertical axis plotted against different horizontal axes of variation.

Figure 4. Power predicted by the Generalised Linear Model fitted to the data for standard simulations as a function of equilibrium frequency ($\hat{p}$, panel A) and selection strength (α, panel B). Results are shown for different parameter combinations as indicated by the right-hand legend of each panel.

Table 1. Binomial GLM of numbers of instances of successful/unsuccessful detections of balancing selection in the reference set of simulations, as a function of equilibrium frequency ($\hat{p}$), selection strength (α), selection model and their interactions. Only terms with P-value <0.05 are shown.

Predictor	Deviance	% deviance explained	P-value
$\hat{p}$	362.96	62.28%	< 0.001
α	47.66	8.2%	< 0.001
$\alpha * \hat{p}$	13.64	2.3%	< 0.001
$\hat{p} *$ selection model	13.68	2.4%	0.001
$\alpha * \hat{p} *$ selection model	33.38	5.7%	< 0.001

As current methods for inferring targets of long-term balancing selection rely largely

on a signal of excess polymorphism and intermediate frequencies in the SFS, power should

increase in conditions that generate a larger and wider signature of excess polymorphism and

decrease under those that restrict the signal. To explore these effects, we ran additional

simulations under OD while varying the recombination and mutation rates, the effective

population size, and the age of the balanced polymorphism. These simulations show that power

consistently decreases at higher recombination rates (Tab. 2), particularly so in cases where the

equilibrium frequency is intermediate (Fig. 5A; Tab. 2). The model predicts that a ten-fold

increase in recombination rate decreases the odds of detection by roughly ten-fold when the

equilibrium is intermediate ($\hat{p} = 0.5$), but only by 4.5-fold for an equilibrium of $\hat{p} = 0.25$ (Tab.

2, b values).

Conversely, power increases with a higher mutation rate per nucleotide site (Fig. 5B).

The model predicts that the odds of successfully detecting balancing selection increase by about

150% for each ten-fold increase in mutation rate (Tab. 2, b value). Note that while there is no

interaction of mutation rate with the allele frequency equilibrium (Tab. 2), there is a nominally

significant triple interaction with the equilibrium frequency and strength of selection, though

this explains a negligible proportion of variation in power (Tab. 2).

The intensity of the signal of balancing selection also depends on the timespan over
which a balanced polymorphism has been maintained, with the age of the balanced
polymorphism in our simulation data having a strong positive effect on our power to detect it
(Fig. 5C; Tab. 2). In our analyses, the allelic age also interacts significantly with the equilibrium
frequency in determining power (Tab. 2). Accordingly, for every additional 0.5 N_e generations
of allelic age in our model, the odds of successfully detecting balancing selection increase by
9.5% for a polymorphism with an equilibrium frequency of 0.5 and by 5.5% with an
equilibrium frequency of 0.25.

Finally, increasing the effective population size generally reduces the detectability of
balanced polymorphisms, though effects of N_e are complex (Fig. 5D). The binomial GLM
reveals a significant negative effect of N_e on power (Tab. 2), with a significant interaction with
the equilibrium frequency and with the interaction of equilibrium frequency with selection
strength (Tab. 2). Thus, the GLM predicts that among loci with a minor allele equilibrium
frequency close to our lower bound of $\hat{p} = 0.1$, and under weak selection ($\alpha = 0.1$), a ten-fold
increase in N_e lowers the odds of successfully detecting balancing selection by over 30%. In
contrast, for loci with an intermediate equilibrium frequency of $\hat{p} = 0.5$ and under strong
selection ($\alpha = 1.5$), the GLM predicts that the same increase in N_e would reduce the odds of
detection by only 10%.

Figure 5. Power to detect balancing selection in our reference set of simulations
(points on the left of panels A-C and on the right of panel D) compared to simulations
with increased recombination rate, mutation rate, age of the selected allele and
reduced effective population size. Balancing selection was simulated under
overdominance, and parameters in the reference set of simulations are N_e = 20,000,
recombination and mutation rates of 10^{-8} per site per generation, selection onset time
160,000 generations ago.

Table 2. Binomial GLM of numbers of instances of successful/unsuccessful detections of balancing selection (overdominance model) as a function of predictors and interactions. Only terms including the variable of interest and with P-value <0.05 are shown.

Predictors included in the model	Predictors discussed in main text	Deviance	% deviance explained	P-value	b (odds ratio)
equilibrium frequency ($\hat{p}$), selection strength (α), $\log_{10}$ of **recombination rate (r)**	r	249.89	55.3%	<0.001	-0.841 (0.43)
	r * $\hat{p}$	11.12	2.5%	0.001	-2.766 (0.06)
equilibrium frequency ($\hat{p}$), selection strength (α), $\log_{10}$ of **mutation rate (m)**	m	112.86	18.9%	<0.001	0.879 (2.41)
	m * $\hat{p}$ * α	3.98	0.7%	0.046	1.776
equilibrium frequency ($\hat{p}$), selection strength (α), **allelic age (t)**	t	557.1	51.2%	<0.001	0.002 (1.002)
	t * $\hat{p}$	69.17	6.4%	<0.001	0.015 (1.015)
	t * α	12.50	1.1%	<0.001	-0.002
equilibrium frequency ($\hat{p}$), selection strength (α), **effective population size (N_e) in thousands of generations**	N_e	82.02	23.2%	<0.001	-0.036 (0.96)
	N_e * $\hat{p}$	17.92	5.1%	<0.001	0.033 (1.03)
	N_e * $\hat{p}$ * α	6.51	1.8%	0.011	0.130 (1.14)

450
451

Discussion

We assessed the propensity of balancing selection to maintain long-term stable polymorphism and the power of current inference methods to successfully detect such loci. The main message from our analyses is that detecting balanced sites in population genomic data can be difficult, at least in the populations of constant size that we simulate here. Even where balancing selection is strong relative to drift and has maintained allelic variation at frequencies closely clustered around an equilibrium frequency of 0.5 for time spans that vastly exceed neutral coalescent times (such as $16N_e$ generations), state-of-the-art methods of inference still only rarely exceed a power of 50%. For example, a mutation under OD with relatively strong and symmetrical selection ($s_1 = s_2 = 0.1$, resulting in $\hat{p} = 0.5$) that is $8N_e$ generations old is detected as a balanced locus with a probability of only 28% in our reference simulations. These selection coefficients are of the same order of magnitude as the ones estimated for the remarkable example of net overdominance maintaining horn size polymorphism in Soay sheep (Johnston et al. (2013), see their Tab. S3) . Our results thus show that in our models, 72% of loci under similar selection strengths and equilibrium frequencies would be missed. This finding contrasts with the notion that inference power with current approaches is good (Bitarello et al., 2023), a perception that is probably due to the empirical focus on human populations, which have particular demographic histories that are generally expected to boost the power of existing methods (see below).

While power in our simulations is moderate in the best of cases, it further decreases, often substantially, under less ideal and perhaps more realistic conditions. The selection parameters have particularly pronounced effects on the long-term maintenance and eventual detectability of a balanced locus. Selection and dominance coefficients determine whether a polymorphism will be subject to balancing selection at all (as in the AP and SA cases), the equilibrium frequencies of balanced polymorphisms, and the efficacy with which selection maintains the alleles near equilibrium. Each of these factors influences the long-term stability of a balanced polymorphism. But even when a polymorphism is long-term stable, balancing selection resulting in allele frequency equilibria close to zero or one are exceedingly unlikely to be identified, not only because they are more susceptible to becoming lost by drift, but also because even those polymorphisms that persist produce weak signals of balancing selection that are very hard to identify.

Model of selection mainly affects maintenance

The models of selection differ very strongly in the strength of selection they generate and hence their propensity to maintain polymorphism in the first place. With OD the net strength of selection is a simple linear function of its underlying selection coefficients ($\alpha = s_1 + s_2$), leading to effectively strong net selection in cases where population-scaled selection parameters are large (i.e., $N_e s_1 \gg 1$ and $N_e s_2 \gg 1$). In contrast, for scenarios of AP and SA without dominance reversals, the net strength of balancing selection (when it occurs) depends on the product of the selection coefficients (i.e., $\alpha = \frac{1}{2} s_1 s_2$ when $h = 0.5$), so that the same values of s_1 and s_2 lead to much weaker balancing selection under AP and SA than under OD (see Appendix 2 for derivations of equations for α). Assuming the distribution of fitness effects is similar under the three mechanisms, then those subject to antagonistic selection should be less likely to remain polymorphic over extended periods of time and, where they do, be harder to detect by genome scans for balancing selection. While dominance reversals dampen these differences between the antagonistic selection scenarios and OD, balancing selection in the SA and AP scenarios are expected to remain weaker and less permissive than the OD case unless dominance reversals are complete.

While the specific model of balancing selection has a strong effect on the propensity to maintain balanced polymorphisms, it did not have an overall effect on detection power. However, our analyses revealed more subtle interactions between the model of selection, the net strength of selection (α), and the equilibrium frequency ($\hat{p}$). Specifically, at low equilibrium frequencies, power decreases with increasing strength of selection under SA, while selection strength has little or no effect on power under OD and AP. This effect is surprising, given that α is intended to capture the strength of selection in a way that is independent of the mechanism generating it. We note, however, that the expressions for α are approximations based on the assumption of effectively weak selection (see Appendix 1). It is therefore possible that the interaction effects we detected here were due to higher-order terms that are not included in the calculation of α.

Selection onset and effective population size

Our simulations also demonstrate that the probability of inferring balancing selection at a locus increases with the age of the polymorphism, as signatures of linked variation build up over time. Importantly, the timescale of this process increases with the effective size of the population. For the signature of selection in linked variation to be discernible, the coalescent

time at the balanced site needs to substantially exceed the average neutral coalescent time of $4N_e$ generations. This means that when N_e is larger, balanced polymorphisms need to be older, in generations, to be detectable. Likewise, for a given age in generations, polymorphisms are harder to detect in species with high N_e than in those with low N_e.

But the timescales over which signatures of selection accumulate, and how they differ between species, also matter for the presence of balancing selection more generally. Thus, the time for signatures to build up can be considerable in geological time. In humans, for example, an age of $8N_e$ generations corresponds to roughly 2My (based on a historical N_e of 10^4 and a 25-year generation time), or roughly a quarter of the time since the human–chimpanzee split. For a polymorphism to be present and detectable after this time, sizable fitness effects need to persist for this duration. This could substantially limit the prospects for the presence of long-term balancing selection in species with large N_e, since stability of the polymorphism requires a relative constancy of the environment over a long time. However, this effect is attenuated by the fact that generation times tend to correlate negatively with N_e. An analysis of the relationship between N_e and generation time (g) (Chao & Carr, 1993) suggests that, very roughly, $g \propto N_e^{-0.35}$, in which case the geological timescale (in years) that is required to generate a genomic signal of long-term balancing selection is on the order of $N_e g \propto N_e^{0.65}$. Thus, despite the inverse relationship between N_e and g, the timescale for generating a signal of balancing selection should generally be higher in lineages with larger N_e and shorter g (such as insects). As a consequence, for a given geological age of a polymorphism, the signals of balancing selection are less likely to be visible in large-N_e lineages because the criteria for "long-term balancing selection" (as defined in Bitarello et al., 2023) are more restrictive.

Interestingly, *Drosophila melanogaster* may be an exception to the broader pattern, in that it has a short generation time (~15 generations/year or 0.07 years/generation, (Turelli & Hoffmann, 1991)) but a relatively modest effective population size (i.e., historical N_e of roughly 10^6 (Arguello et al., 2019)). Based on these figures, $8N_e$ would correspond to 0.56My, much shorter than the equivalent time in humans, which makes it more likely that broader environmental conditions remain constant enough to allow for the maintenance of balanced polymorphisms in this species. In addition, extremely short generation times such as those in *Drosophila* provide further scope for balancing selection due to seasonal fluctuations in the environment (Wittmann et al., 2023), which is an unlikely avenue for balancing selection in species with long generation times.

The effective population size also has other direct and indirect effects on the presence

and detectability of balanced polymorphisms, beyond the timescales discussed above. Large

N_e leads to reduced drift and concomitantly more effective selection to maintain balanced

polymorphisms and should therefore (all else being equal), increase the proportion of balanced

loci that remains long-term polymorphic. On the other hand, N_e also has adverse effects on the

signatures at linked neutral polymorphism. Species with large N_e tend to have larger effective

recombination rates (larger $N_e r$), owing trivially to the effect of larger N_e but also due to the

fact that large-N_e species often have relatively streamlined genomes and accordingly higher

per-nucleotide recombination rates (r) (Lynch, 2006). Elevated recombination will erode the

balancing selection signal of intermediate frequencies at neutral sites that are closely linked to

selected variants and thereby make the signature sparser and inference harder. This effect can

be compensated by a higher mutation rate (μ), which favours the generation of linked

polymorphism around the balanced site. Thus, for a given N_e, the power to detect balancing

selection is determined by the ratio of mutation to recombination rate, μ/r.

Demographic history and comparison to previous power estimates

It is worth noting that while we have only simulated simple scenarios of constant population

size, more complex demographic histories are likely to substantially affect detection of

balanced polymorphisms. Changes in the effective population size over time, such as sudden

contractions (bottlenecks) and subsequent expansions, can have beneficial effects on detection.

Power can be improved because a bottleneck in the not-too-distant past will result in increased

rates of coalescence (a more recent average TMRCA) at neutral regions that are unlinked to

sites under balancing selection, and thus on average tend to cause a leftward-shifted SFS at

those regions. Peaks of intermediate frequencies and older TMRCAs at sites linked to balanced

loci stand out more clearly from such genomic backgrounds of neutral diversity. It is likely that

this phenomenon has facilitated the identification of balanced loci in humans, where migration

out of Africa has resulted in a series of bottlenecks and founder events that have systematically

shifted the neutral SFS.

The small N_e and specific demographic history of humans likely explains why previous

analyses of the power of current inference methods, including the NCD statistic used here

(Bitarello et al., 2018) or BalLeRMix (Cheng & DeGiorgio, 2020), have provided higher

estimates of power (reaching over 80% under some conditions) than the ones we report here

for our human-like 'reference' parameters. While methods like NCD or BalLeRMix are general

in scope, they have typically been benchmarked using simulated data that follow a
demographic history and a mutation-to-recombination rate ratio that is characteristic of human
populations (e.g., 2.5 in Bitarello et al. (2018) and sampled between 2.5 and 0.83 in Cheng and
DeGiorgio (2020)), and relatively old polymorphisms (*e.g.* 5 million years in Bitarello et al.
(2018)and up to 650,000 generations with $N_e = 10^4$ in Cheng and DeGiorgio (2020), which
corresponds to 16.25 million years with the generation time of 25 years used in Bitarello et al.
(2018)). Given the specific assumptions made in the original method publications and the
substantial variation in power as a function of the parameters used, we reiterate that it is
imperative to combine any application of existing methods to a careful power analysis,
parameterised as much as possible to fit the study organism and population (Bitarello et al.,
2018, 2023).

Methodological notes on NCD

We have deliberately focused our analyses on a single approach, NCD (but see Supplementary
Material for additional analyses with BalLeRMix), to evaluate how different population genetic
parameters determine the efficacy of balancing selection and the signals it generates. It is not
our aim to compare different methods, all of which essentially exploit the same signals in the
data and will hence be affected similarly by the parameters we explore. Nevertheless, we will
briefly discuss a few specificities of our application of NCD that are expected to have small
effects on the absolute power we estimate.

First, in our analyses, we use the actual equilibrium frequency, which is known in our
simulations, as the NCD 'target frequency' (i.e., the frequency value from which NCD
measures the mean squared distance of allele frequencies in a window). Doing so will have a
small positive effect on power, compared to the case of an empirical application where the
target frequency is not known, and is therefore set to an arbitrary intermediate value. Previous
analyses (Bitarello et al., 2018), however, have shown that power is minimally impacted by the
exact frequency value of the target and additional analyses performed on our simulations
confirm this (Fig. S14).

On the other hand, two other specificities of our application of NCD are likely to have
a small negative effect on power. First, we use a variant of NCD, NCD2, that takes into account
the SFS of derived alleles, including substitutions. Because we do not simulate an outgroup,
however, NCD2 is not as powerful here as the approach would be in a real-world application
where the SFS also includes fixed sites that result from substitutions inferred from outgroup

data (see Methods). Second, we use NCD with a fixed window size of 3Kb, independently of

the recombination rate. In cases where recombination rate is high and regions of elevated

polymorphism are narrow, calculating the SFS over a smaller interval can improve power,

although this is offset by the fact that the small numbers of SNPs in a narrow focal window

generate noise that negatively impacts the inference (Fig. S15). Again, real-world applications

should be preceded by power analyses that explore these effects and can help to optimise the

analysis.

General considerations on differences between species

Our analyses make it clear that the prospects of identifying targets of balancing selection are

expected to vary significantly and systematically among species. Thus, species will harbour

variable numbers of long-term balanced loci, depending on the effective sizes of their

populations. Furthermore, the effective size, combined with mutation and recombination rates

will determine how difficult these sites are to identify as targets of balancing selection. In

humans, for example, the small effective population size means that only a limited number of

balanced polymorphisms are expected to have remained polymorphic over sufficiently long

periods of time. On the other hand, the same low effective population size, together with the

high ratio of mutation to recombination rate ($\mu/r = 1.2 \times 10^{-8}/1.3 \times 10^{-8} \sim 1$ (Pritchard,

2023 Chap. 1.1)), favour detection of such loci. In contrast, *D. melanogaster*, and possibly

other insects, presents a much harder case for identification, as the recombination rate is high

relative to the mutation rate ($\mu/r \sim 3 \times 10^{-9}/3 \times 10^{-8} = 0.1$ (Wang et al., 2023)), which

our analyses suggest should reduce power by roughly four-fold compared to a species with

human-like mutation and recombination parameters (mean power across simulated parameters

in Fig. 4A, 0.14 vs. 0.03).

Of course, as discussed above, there are other important factors at work that affect the

relative frequency of detectable balanced loci in the two species, such as the time required for

long-term balancing selection, the scope for seasonal selection, effects of N_e on the

maintenance of polymorphism on the one hand and its detectability on the other, and species-

specific demographic histories. Overall, there is little doubt that the detection of balancing

selection in *Drosophila* is challenging, which suggests that there could be many targets of

balancing selection in *D. melanogaster* that are invisible to genomic scans for balancing

selection, and this may also be the case for many other species with similar population

parameters. Indeed, a genome-wide screen for balancing selection in fruit flies combining

Theta and Tajima's *D* identified roughly 100 candidate balanced loci (Croze et al., 2017),

compared to over 1000 candidates in humans (Bitarello et al., 2018). Although more

sophisticated methods may improve power in *Drosophila*, the notion that large numbers of

balanced polymorphisms might exist and be missed in genomic scans is in line with the high

levels of standing genetic variation for major fitness components in this species, which exceeds

the variation explainable under mutation-selection-drift balance (Charlesworth, 2015; Sharp &

Agrawal, 2018). The excess variance in life-history traits might be attributable to balanced

polymorphisms segregating in *D. melanogaster* populations. While it seems plausible that at

least some of these balanced polymorphisms would be detected if detection methods were more

powerful, others could remain invisible if they are relatively young (relative to $4N_e$). It is

difficult to assess these alternatives in the absence of more powerful approaches, and similar

conclusions may apply to many other species with equally unfavourable population parameters.

Methods

Population genetic simulations of balancing selection

We used simulations to assess which regions of the parameter space generate genomic

signatures of balancing selection that can be detected using current approaches. Simulations

were run in SLiM (version 4.0.1, (Haller et al., 2019)), a forward simulation framework for

population genetics. In line with the analytical derivations above, we implemented

overdominance (OD), antagonistic pleiotropy (AP) and sexually antagonistic selection (SA),

in addition to simulations under neutrality. In the simulations with selection, we simulated a

single focal site, embedded centrally in a chromosome of 10,000bp, large enough for neutral

sites in the periphery to be unaffected by selection at the central site. To implement the three

selection mechanisms, we used custom functions (see SLiM scripts in available in

https://github.com/reuterlab/bls_sim/) to run a non-Wright-Fisher model for populations of

constant size with non-overlapping generations. In each generation, the code generated

offspring by randomly drawing parents based on their fitness values, calculated according to

their genotype at the locus under selection and the selection mechanism and parameters

(selection coefficients, dominance, cf. Tab. A1 in Appendix 1) being simulated. We used the

same code for reproduction in the neutral simulations, but set the fitness of all individuals to 1.

In all simulations the default sex ratio of 1:1 was used.

At the start of each simulation, a mutation at the focal site was introduced on a single

chromosome, randomly chosen within the population. The mutation would be of type A_1 or A_2

(see Tab. A1, *e.g.*, either male- or female-beneficial for SA) with equal probability. To reduce

the number of simulation runs where the polymorphism was quickly lost due to drift when rare,

we initially subjected the mutant allele to directional positive selection with $s = 0.5$ and $h = 1/2$

until the mutant allele frequency reached 5% (or the mutant allele was lost). This selective

boost facilitates the establishment of the polymorphism, but due to its old age and short

duration should have negligible influence on the patterns of linked variation. For

polymorphisms that reached a frequency of 5%, simulations then switched from directional to

balancing selection with a specified mechanism and parameter combination. We recorded the

frequency of the mutant (derived) allele and the genealogy of all chromosomes in the

population at 16000, 32000, 160000 and 320000 generations.

To optimise the runtime of simulations, we did not include neutral mutations in any

of the SLiM simulations. Instead, we recorded the full chromosome-wide genealogies from

each simulation (using 'tree-sequence recording', (Haller et al., 2019)). We then used

'recapitation' in pyslim (version 1.0.4 (Ralph et al., 2023)) to generate the preceding

genealogical relationships of the starting chromosomes up to their most recent common

ancestor. Finally, we added neutral mutations on the complete recapitated genealogies with

msprime (version 1.2.0, (Baumdicker et al., 2022)).

Table 3. Parameter values used in the simulations. We highlight in bold the values used in our "reference" set of simulations, which approximate the parameter values estimated in the literature for humans.

Selection model	Overdominance (OD)
	Antagonistic pleiotropy (AP)
	Sexual antagonism (SA)
Selection coefficients (s)	s_{weak} = {0.01, 0.02,0.05,0.1}
	s_{strong} = {0.1,0.2,0.5,1}
Dominance coefficients (h, only defined for AP and SA)	0.25
	0.5
Time of onset of selection (T, in generations)	**8Ne**

	16Ne
(Effective) population size ($N=N_e$)	2,000 **20,000**
Recombination rate (r)	10^{-7} **10^{-8}**
Mutation rate (μ)	10^{-7} **10^{-8}**

The parameter values used in our simulations are summarised in Table 3. We explored
selection and dominance coefficients to match the analytical results. For the species-specific
parameters of effective population size (N_e), recombination rate (r) and mutation rate (μ) were
chosen to cover a broad range of relevant scenarios. Thus, we simulated population sizes of N
$= N_e = 2,000$ and $N = N_e = 20,000$ and neutral mutation at rates of $\mu = 10^{-7}$ and $\mu = 10^{-8}$ per
base per generation, such that the expected genetic diversity ($\pi = 4N_e\mu$) ranges from 8×10^{-5} to
8×10^{-3}. The highest expected genetic diversity (for $N_e = 20,000$ and $\mu = 10^{-7}$) approaches that
of *Drosophila melanogaster* ($\pi = 10^{-2}$ (Leffler et al., 2012)), while the combination of $N_e =$
$20,000$ and $\mu = 10^{-8}$ generates diversity similar to that estimated in humans ($\pi = 8 \times 10^{-4}$
(Pritchard, 2023 Chap.1.1)). We simulated recombination rates of $r=10^{-7}$ and $r = 10^{-8}$ per base
per generation to include combinations of $\mu = r =10^{-8}$, which are similar to the values estimated
in humans (Pritchard, 2023 Chap. 1.1), as well as a ratio of mutation to recombination rate
similar to that estimated in *Drosophila melanogaster* ($\mu/r = 10^{-8}/10^{-7} = 0.1$, (Wang et al., 2023))
and a scenario where mutation rate exceeds recombination rate ($\mu/r = 10^{-7}/10^{-8} = 10$), which
should facilitate the detection of signatures of selection. Finally, we used two values for the
onset of selection (or age of the polymorphism), $T = 8N_e$ and $T = 16N_e$ (16,000 and 32,000
generations for $N_e = 2,000$ and 160,000 and 320,000 generations for $N_e = 20,000$). These values
roughly correspond to the definitions of long-term and ultra long-term balancing selection
(Bitarello et al. 2023). Both significantly exceed the expected coalescence time (time to the
most recent common ancestor, TMRCA) under neutrality, $4N_e$. The probability of observing a
TMRCA larger than $8N_e$ generations under neutrality is approximately 0.054, while that of a
TMRCA larger than $16N_e$ generations is approximately 0.001 (Wakeley 2009, p.77 eqn 3.27).

To explore the effect of the strength of selection, equilibrium frequency and model of

selection on the maintenance of the selected polymorphism (Figs. S1–S6) and the inference of

balancing selection (Fig. 3), we ran simulations for a reference set with human-like parameters

(bold values in Tab. 1, N_e=20,000, $\mu = r = 10^{-8}$, $T = 8N_e$ =160,000 generations) under OD, AP

and SA across combinations of selection coefficients s_i and s_j drawn from two 4-by-4 grids

with weaker selection {0.01,0.02,0.05,0.1} and stronger selection {0.1,0.2,0.5,1.0}. To explore

the effects of mutation rate, recombination rate, selection onset time and N_e, (Fig. 5), we ran

similar sets of simulations for OD only, but changing the values of each of these parameters

relative to the reference set of simulations (i.e. $\mu = 10^{-7}$, $r = 10^{-7}$, $T = 16N_e = 320,000$

generations, $N_e = 2,000$).

In simulations to assess the maintenance of polymorphisms, we ran 100 replicate

simulations in which the mutation successfully established (*i.e.*, reached 5%) for each

parameter combination. To assess the power to detect signatures of balancing selection, we

complemented these with additional simulations to obtain sets of 100 replicate simulations for

each parameter combination in which the polymorphism was maintained.

Power of inference of balancing selection

We used the NCD (non-central deviation) method (Bitarello et al., 2018) to infer balancing

selection on the simulated data (but for complementary results using BalLeRMix (Cheng &

DeGiorgio, 2020), see Supplementary Material). Like most other current inference methods,

NCD detects regions under balancing selection due to their signal of excess intermediate-

frequency polymorphism. NCD for a genomic window with n informative sites is defined as:

739
$$NCD(tf) = \sqrt{\frac{\sum_{i=1}^{n}(p_i - tf)^2}{n}}$$

where p_i is the allele frequency at the ith informative site, and tf is the target frequency (set

as an intermediate value *e.g.*, 0.3, 0.4 or 0.5). The presence of a balanced site in the focal region

with allele frequencies close to the target frequency then results in a low value of the NCD

statistic, compared to a region under neutrality or purifying selection.

In our application of this approach, we used a refined version of the statistic, NCD2.

This statistic gains power by also considering fixed differences (relative to an outgroup) in the

site frequency spectrum (Bitarello et al., 2018), thus exploiting the signal of increased

polymorphism and lower rate of substitution in regions under balancing selection. Although

we have not simulated an outgroup, we can differentiate ancestral and derived alleles in our

simulated genotypes and then manually add an 'outgroup' individual to the VCF file from our

simulations, for which all sites are homozygous ancestral. Therefore, any sites that are fixed

for the derived allele among the simulated (ingroup) individuals will be counted as fixed

differences in the NCD2 calculation. We note that this approach adds less additional power

than a real outgroup would, as the latter would also contribute fixed differences that arise due

to substitutions in the outgroup lineage.

To assess the statistical significance of the NCD2 values, we generated a null

distribution by running 1000 neutral simulations for each combination of relevant parameters

(N_e, μ and r) and calculating NCD2 for each. We then calculated a P-value for an NCD2 value

from a simulation with selection by determining the proportion of NCD2 values from the null

distribution (with corresponding N_e, μ and r) that was lower than the NCD2 value with

selection. Power was then calculated as the proportion of the 100 simulations generated under

a particular set of parameters for which P ≤ 0.01.

Statistical analysis of power estimates

We fitted binomial Generalised Linear Models (GLMs) to our power data to directly model the

probability of detecting or failing to detect balancing selection as a function of the parameters

that were varied in our simulations. The models used a logit link function and were fitted to the

number of successful and unsuccessful instances of detection as the response variable. The

significance of model terms was determined with an analysis of deviance, a series of sequential

likelihood ratio tests based on the Chi^2 distribution. This analysis further makes it possible to

assess the importance of each model term through its deviance (the amount of variability in the

response explained by a model term) relative to the null deviance, the total amount of variability

in the response values analysed. We note that all models showed a degree of overdispersion,

as indicated by the residual deviance exceeding the number of residual degrees of freedom

several-fold. However, running alternative models that accommodate overdispersion (as a

'quasibinomial' model, combined with F-tests) produced qualitatively identical results.

These GLMs fit a linear (additive) model of detection probability (π) as a function of

the simulation parameters. The model is fitted on the log-odds scale (where odds are $\pi/(1 -$

$\pi)$) that accommodates the fact that probabilities are bounded by 0 and 1. The estimated model

parameters, back-transformed to the odds-scale, provide information on the proportional

change in the detection probability that is independent of the value of π (while the change in
the actual probability due to a change in a predictor variable changes over the range of π).
782

Acknowledgements

We would like to thank Filip Ruzicka and members of the Alliance EvolGen lab group for
helpful discussions. MR, AA and DB were supported by grants from the UK Biotechnology
and Biological Sciences Research Council, BBSRC (award BB/W007703/1 to MR and AA)
and the Leverhulme Trust (award RPG-2021-414 to MR and AA). The work was initiated and
developed during mutual visits between London and Melbourne by MR, AA and TC, made
possible by an Australia Partnering award from the BBSRC (award BB/T019921/1 to MR and
TC).

Bibliography

Allison, A. C. (1954). The distribution of the sickle-cell trait in East Africa and elsewhere, and
its apparent relationship to the incidence of subtertian malaria. *Transactions of the*
Royal Society of Tropical Medicine and Hygiene, *48*(4), 312–318.
https://doi.org/10.1016/0035-9203(54)90101-7

Arguello, J. R., Laurent, S., & Clark, A. G. (2019). Demographic History of the Human
Commensal Drosophila melanogaster. *Genome Biology and Evolution*, *11*(3), 844–854.
https://doi.org/10.1093/gbe/evz022

Barson, N. J., Aykanat, T., Hindar, K., Baranski, M., Bolstad, G. H., Fiske, P., Jacq, C., Jensen,
A. J., Johnston, S. E., Karlsson, S., Kent, M., Moen, T., Niemelä, E., Nome, T., Næsje,
T. F., Orell, P., Romakkaniemi, A., Sægrov, H., Urdal, K., … Primmer, C. R. (2015).
Sex-dependent dominance at a single locus maintains variation in age at maturity in
salmon. *Nature*, *528*(7582), 405–408. https://doi.org/10.1038/nature16062

Baumdicker, F., Bisschop, G., Goldstein, D., Gower, G., Ragsdale, A. P., Tsambos, G., Zhu,
S., Eldon, B., Ellerman, E. C., Galloway, J. G., Gladstein, A. L., Gorjanc, G., Guo, B.,

Jeffery, B., Kretzschumar, W. W., Lohse, K., Matschiner, M., Nelson, D., Pope, N. S.,

… Kelleher, J. (2022). Efficient ancestry and mutation simulation with msprime 1.0.

Genetics, *220*(3), iyab229. https://doi.org/10.1093/genetics/iyab229

Bitarello, B. D., Brandt, D. Y. C., Meyer, D., & Andrés, A. M. (2023). Inferring Balancing

Selection From Genome-Scale Data. *Genome Biology and Evolution*, *15*(3), evad032.

https://doi.org/10.1093/gbe/evad032

Bitarello, B. D., De Filippo, C., Teixeira, J. C., Schmidt, J. M., Kleinert, P., Meyer, D., &

Andrés, A. M. (2018). Signatures of Long-Term Balancing Selection in Human

Genomes. *Genome Biology and Evolution*, *10*(3), 939–955.

https://doi.org/10.1093/gbe/evy054

Chao, L., & Carr, D. E. (1993). The molecular clock and the relationship between population

size and generation time. *Evolution*, *47*(2), 688–690. https://doi.org/10.1111/j.1558-

5646.1993.tb02124.x

Charlesworth, B. (2015). Causes of natural variation in fitness: Evidence from studies of

Drosophila populations. *Proceedings of the National Academy of Sciences*, *112*(6),

1662–1669. https://doi.org/10.1073/pnas.1423275112

Charlesworth, B., & Charlesworth, D. (2010). *Elements of Evolutionary Genetics*. Roberts and

Company Publishers.

Charlesworth, B., & Hughes, K. A. (2000). The Maintenance of Genetic Variation in Life-

History Traits. In *Evolutionary Genetics: From molecules to morphology*.

Cheng, X., & DeGiorgio, M. (2020). Flexible Mixture Model Approaches That Accommodate

Footprint Size Variability for Robust Detection of Balancing Selection. *Molecular*

Biology and Evolution, *37*(11), 3267–3291. https://doi.org/10.1093/molbev/msaa134

Connallon, T., & Chenoweth, S. F. (2019). Dominance reversals and the maintenance of

genetic variation for fitness. *PLOS Biology*, *17*(1), e3000118.

https://doi.org/10.1371/journal.pbio.3000118

Connallon, T., & Clark, A. G. (2014). Balancing Selection in Species with Separate Sexes:

Insights from Fisher's Geometric Model. *Genetics*, *197*(3), 991–1006.

https://doi.org/10.1534/genetics.114.165605

Connallon, T., & Czuppon, P. (2026). Additive genetic variance for fitness maintained by

balancing selection. *Evolution, in Press*.

Crow, J. F. (1952). Dominance and Overdominance. In *Heterosis: A Record of Researches*

Directed Toward Explaining and Utilizing the Vigor of Hybrids. Iowa State College

Press.

Crow, J. F. (1987). Muller, Dobzhansky, and overdominance. *Journal of the History of Biology*,

20(3), 351–380. https://doi.org/10.1007/BF00139460

Croze, M., Wollstein, A., Božičević, V., Živković, D., Stephan, W., & Hutter, S. (2017). A

genome-wide scan for genes under balancing selection in Drosophila melanogaster.

BMC Evolutionary Biology, *17*(1), 15. https://doi.org/10.1186/s12862-016-0857-z

Curtsinger, J. W., Service, P. M., & Prout, T. (1994). Antagonistic Pleiotropy, Reversal of

Dominance, and Genetic Polymorphism. *The American Naturalist*.

https://doi.org/10.1086/285671

DeGiorgio, M., Lohmueller, K. E., & Nielsen, R. (2014). A Model-Based Approach for

Identifying Signatures of Ancient Balancing Selection in Genetic Data. *PLOS Genetics*,

10(8), e1004561. https://doi.org/10.1371/journal.pgen.1004561

Dobzhansky, T. (1955). A Review of Some Fundamental Concepts Problems of Population

Genetics. *Cold Spring Harbor Symposia on Quantitative Biology*, *20*, 1–15.

Fijarczyk, A., & Babik, W. (2015). Detecting balancing selection in genomes: Limits and

prospects. *Molecular Ecology*, *24*(14), 3529–3545. https://doi.org/10.1111/mec.13226

Fisher, R. A. (1922). On the Dominance Ratio. *Proceedings of the Royal Society of Edinburgh*,

42, 321–341. https://doi.org/10.1017/S0370164600023993

Fitzpatrick, M. J., Feder, E., Rowe, L., & Sokolowski, M. B. (2007). Maintaining a behaviour

polymorphism by frequency-dependent selection on a single gene. *Nature*, *447*(7141),

210–212. https://doi.org/10.1038/nature05764

Glémin, S. (2021). Balancing selection in self-fertilizing populations. *Evolution; International*

Journal of Organic Evolution, *75*(5), 1011–1029. https://doi.org/10.1111/evo.14194

Grieshop, K., Ho, E. K. H., & Kasimatis, K. R. (2024). Dominance reversals: The resolution

of genetic conflict and maintenance of genetic variation. *Proceedings of the Royal*

Society   B: Biological   Sciences,   *291*(2018),   20232816.

https://doi.org/10.1098/rspb.2023.2816

Haldane, J. B. S., & Jayakar, S. D. (1963). Polymorphism due to selection depending on the

composition   of   a   population.   *Journal   of   Genetics*,   *58*(3),   318–323.

https://doi.org/10.1007/BF02986301

Haller, B. C., Galloway, J., Kelleher, J., Messer, P. W., & Ralph, P. L. (2019). Tree-sequence

recording in SLiM opens new horizons for forward-time simulation of whole genomes.

Molecular   Ecology   Resources,   *19*(2),   552–566.   https://doi.org/10.1111/1755-

0998.12968

Hedrick, P. W. (1999). Antagonistic pleiotropy and genetic polymorphism: A perspective.

Heredity, *82*(2), 126–133. https://doi.org/10.1038/sj.hdy.6884400

Hedrick, P. W. (2012). What is the evidence for heterozygote advantage selection? *Trends in*

Ecology & Evolution, *27*(12), 698–704. https://doi.org/10.1016/j.tree.2012.08.012

Isildak, U., Stella, A., & Fumagalli, M. (2021). Distinguishing between recent balancing

selection and incomplete sweep using deep neural networks. *Molecular Ecology*

Resources, *21*(8), 2706–2718. https://doi.org/10.1111/1755-0998.13379

Johnston, S. E., Gratten, J., Berenos, C., Pilkington, J. G., Clutton-Brock, T. H., Pemberton, J.

M., & Slate, J. (2013). Life history trade-offs at a single locus maintain sexually

selected genetic variation. *Nature*, *502*(7469), 93–95.

https://doi.org/10.1038/nature12489

Karageorgi, M., Lyulina, A. S., Bitter, M. C., Lappo, E., Greenblum, S. I., Mouza, Z. K., Tran,

C. T., Huynh, A. V., Oken, H., Schmidt, P., & Petrov, D. A. (2025). Beneficial reversal

of dominance maintains a large-effect resistance polymorphism under fluctuating

insecticide selection. *Nature Ecology & Evolution*, *9*(11), 2145–2160.

https://doi.org/10.1038/s41559-025-02853-x

Kidwell, J. F., Clegg, M. T., Stewart, F. M., & Prout, T. (1977). Regions of stable equilibria

for models of differential selection in the two sexes under random mating. *Genetics*,

85(1), 171–183. https://doi.org/10.1093/genetics/85.1.171

Kimura, M. (1968). Evolutionary Rate at the Molecular Level. *Nature*, *217*(5129), 624–626.

https://doi.org/10.1038/217624a0

Kimura, M. (1983). *The Neutral Theory of Molecular Evolution*. Cambridge University Press.

http://dx.doi.org/10.1017/CBO9780511623486

King, J. L., & Jukes, T. H. (1969). Non-Darwinian Evolution. *Science*, *164*(3881), 788–798.

https://doi.org/10.1126/science.164.3881.788

Klein, J., Sato, A., Nagl, S., & O'hUigín, C. (1998). Molecular trans-species polymorphism.

Annual Review of Ecology, Evolution, and Systematics, *29*(Volume 29, 1998), 1–21.

https://doi.org/10.1146/annurev.ecolsys.29.1.1

Kreitman, M. (1983). Nucleotide polymorphism at the alcohol dehydrogenase locus of

Drosophila melanogaster. *Nature*, *304*(5925), 412–417.

https://doi.org/10.1038/304412a0

Leffler, E. M., Bullaughey, K., Matute, D. R., Meyer, W. K., Ségurel, L., Venkat, A.,

Andolfatto, P., & Przeworski, M. (2012). Revisiting an old riddle: What determines

genetic diversity levels within species? *PLoS Biology*, *10*(9), e1001388.

https://doi.org/10.1371/journal.pbio.1001388

Levene, H. (1953). Genetic Equilibrium When More Than One Ecological Niche is Available.

The American Naturalist. https://doi.org/10.1086/281792

Lewontin, R. C., & Hubby, J. L. (1966). A molecular approach to the study of genic

heterozygosity in natural populations. II. Amount of variation and degree of

heterozygosity in natural populations of Drosophila pseudoobscura. *Genetics*, *54*(2),

595–609. https://doi.org/10.1093/genetics/54.2.595

Lewontin, R. C. (with Internet Archive). (1974). *The genetic basis of evolutionary change*.

New York, Columbia University Press.

http://archive.org/details/geneticbasisofev00lewo

Lynch, M. (2006). The Origins of Eukaryotic Gene Structure. *Molecular Biology and*

Evolution, *23*(2), 450–468. https://doi.org/10.1093/molbev/msj050

Nei, M., & Roychoudhury, A. K. (1973). Probability of Fixation and Mean Fixation Time of

an Overdominant Mutation. *Genetics*, *74*(2), 371–380.

https://doi.org/10.1093/genetics/74.2.371

Pritchard, J. K. (2023). *An Owner's Guide to the Human Genome* (Release 1.0).

Prout, T. (2000). How well does opposing selection maintain variation? In *Evolutionary*

Genetics: From molecules to morphology (pp. 157–181). Cambridge University Press.

Ralph, P., Jeffery, B., Wong, Y., Ashander, J., Patterson, G., Kelleher, J., R, M., Kern, A.,

Tittes, S., Huang, X., & Galloway, J. (2023). *tskit-dev/pyslim: 1.0.4* [Computer

software]. Zenodo. https://doi.org/10.5281/zenodo.8205346

Reid, J. (2022). Intrinsic emergence and modulation of sex-specific dominance reversals in

threshold traits. *Evolution*, *76*, 1924–1941.

Rice, S. H. (2004). *Evolutionary Theory: Mathematical and Conceptual Foundations*. Sinauer.

Robertson, A. (1962). Selection for Heterozygotes in Small Populations. *Genetics*, *47*(9),

1291–1300. https://doi.org/10.1093/genetics/47.9.1291

Rose, M. R. (1982). Antagonistic pleiotropy, dominance, and genetic variation. *Heredity*,

48(1), 63–78. https://doi.org/10.1038/hdy.1982.7

Rusuwa, B. B., Chung, H., Allen, S. L., Frentiu, F. D., & Chenoweth, S. F. (2022). Natural

variation at a single gene generates sexual antagonism across fitness components in

Drosophila. *Current Biology*, *32*(14), 3161–3169.

https://doi.org/10.1016/j.cub.2022.05.038

Ruzicka, F., Zwoinska, M. K., Goedert, D., Kokko, H., Li Richter, X.-Y., Moodie, I. R., Nilén,

S., Olito, C., Svensson, E. I., Czuppon, P., & Connallon, T. (2025). A century of

theories of balancing selection. *Biological Reviews*. https://doi.org/10.1111/brv.70103

Sharp, N. P., & Agrawal, A. F. (2018). An experimental test of the mutation-selection balance

model for the maintenance of genetic variance in fitness components. *Proceedings.*

Biological Sciences, *285*(1890), 20181864. https://doi.org/10.1098/rspb.2018.1864

Siewert, K. M., & Voight, B. F. (2017). Detecting Long-Term Balancing Selection Using

Allele Frequency Correlation. *Molecular Biology and Evolution*, *34*(11), 2996–3005.

https://doi.org/10.1093/molbev/msx209

Soni, V., Vos, M., & Eyre-Walker, A. (2022). A new test suggests hundreds of amino acid

polymorphisms in humans are subject to balancing selection. *PLOS Biology*, *20*(6),

e3001645. https://doi.org/10.1371/journal.pbio.3001645

Turelli, M., & Hoffmann, A. A. (1991). Rapid spread of an inherited incompatibility factor in

California Drosophila. *Nature*, *353*(6343), 440–442. https://doi.org/10.1038/353440a0

Wang, Y., McNeil, P., Abdulazeez, R., Pascual, M., Johnston, S. E., Keightley, P. D., &

Obbard, D. J. (2023). Variation in mutation, recombination, and transposition rates in

Drosophila melanogaster and Drosophila simulans. *Genome Research*, *33*(4), 587–598.

https://doi.org/10.1101/gr.277383.122

Williamson, S., Fledel-Alon, A., & Bustamante, C. D. (2004). Population Genetics of

Polymorphism and Divergence for Diploid Selection Models With Arbitrary

Dominance. *Genetics*, *168*(1), 463–475. https://doi.org/10.1534/genetics.103.024745

Wittmann, M. J., Mousset, S., & Hermisson, J. (2023). Modeling the genetic footprint of

fluctuating balancing selection: From the local to the genomic scale. *Genetics*, *223*(4),

iyad022. https://doi.org/10.1093/genetics/iyad022

Wright, S. (1942). Statistical genetics and evolution. *Bulletin of the American Mathematical*

Society, *48*(4), 223–246. https://doi.org/10.1090/S0002-9904-1942-07641-5

Wright, S. (1945). The Differential Equation of the Distribution of Gene Frequencies.

Proceedings of the National Academy of Sciences, *31*(12), 382–389.

https://doi.org/10.1073/pnas.31.12.382

Zajitschek, F., & Connallon, T. (2018). Antagonistic pleiotropy in species with separate sexes,

and the maintenance of genetic variation in life-history traits and fitness. *Evolution*,

72(6), 1306–1316. https://doi.org/10.1111/evo.13493

Appendices

Appendix 1 – Conditions for internal frequency equilibria

We model balancing selection making three simplifying assumptions that should apply broadly to many (perhaps most) cases where this type of selection is present. First, we focus on populations of diploid, randomly mating (outbred) individuals (see Glémin (2021) for a comprehensive treatment of balancing selection in self-fertilizing species). Second, we consider loci with two major allele types, which we label A_1 (at a population frequency of p) and A_2 (frequency $1 - p$). Third, in scenarios involving antagonistic selection (i.e., antagonistic pleiotropy and sexual antagonism), we only consider scenarios of dominance that are amenable to balanced polymorphism, i.e. cases where each fitness component or sex either lacks dominance (fitness effects are additive in each context of selection) or there is a beneficial reversal of dominance, in which the costs of expressing the "wrong" allele for a given context of selection are partially or completely recessive (reviewed in Connallon and Chenoweth 2019; Grieshop et al. 2024). Our exposition draws upon previous studies of balancing selection theory (*e.g.*, Robertson 1962; Kidwell et al. 1977; Curtsinger et al. 1994; Connallon and Clark 2013), with Table A1 providing an outline of the fitness values associated with each genotype and for each model of selection.

Table A1. Fitness per genotype for each model of selection

	Genotype		
	A_1A_1	A_1A_2	A_2A_2
Overdominance	$1 - s_1$	1	$1 - s_2$
Antagonistic pleiotropy			
Fitness component 1	1	$1 - t_1 h$	$1 - t_1$
Fitness component 2	$1 - t_2$	$1 - t_2 h$	1
Overall Fitness	$1 - t_2$	$(1 - t_1 h)(1 - t_2 h)$	$1 - t_1$
Sexual antagonism			

Female fitness	1	$1 - s_f h_f$	$1 - s_f$
Male fitness	$1 - s_m$	$1 - s_m h_m$	1

Parameters for the three models fall within the following ranges: overdominance: $1 \geq$
$s_1, s_2 > 0$; antagonistic pleiotropy: $1 \geq t_1, t_2 > 0$; sexual antagonism: $1 \geq s_f, s_m > 0$.
Dominance coefficients are within the range $0.5 \geq h, h_f, h_m \geq 0$, which includes additive
effects ($h = 0.5$) and beneficial reversals of dominance ($0.5 > h$). Note that many of our
analyses, we focus on scenarios where fitness effects are small, *i.e.*: $1 \gg s_1, s_2 > 0$; $1 \gg$
$t_1, t_2 > 0$; and $1 \gg s_f, s_m > 0$. Note that in the Appendices, we use slightly different notation
to distinguish selection parameters of the OD, AP, and SA models (s_1/s_2, t_1/s_2, and s_f/s_m,
respectively). For simplicity, and with no loss of generality, we use a single notation, s_1/s_2, in
the main text.
999

Overdominant selection (OD)

1001　With OD, the heterozygotes for the locus (A_1A_2) have the highest fitness and A_1A_1 and A_2A_2
1002　homozygotes suffer fitness reductions of s_1 and s_2, respectively (Tab. A1). Evolutionary
1003　dynamics are captured by the following difference equation describing allele frequency change
1004　across a single generation:

$$\Delta p = \frac{(s_1 + s_2)p(1 - p)(\hat{p} - p)}{\overline{w}} \approx (s_1 + s_2)p(1 - p)(\hat{p} - p) \qquad (1)$$

1006　where $\underline{w} = p^2(1 - s_1) + 2p(1 - p) + (1 - p)^2(1 - s_2)$ denotes mean fitness and:

$$\hat{p} = \frac{s_2}{s_1 + s_2}$$

1008　is the equilibrium frequency for the A_1 allele. The final approximation in eq. (1) applies under
1009　weak selection ($0 < s_1, s_2 \ll 1$). Note that the entire range of parameter values in this model
1010　generates balancing selection (*i.e.*, $0 < \hat{p} < 1$ for the plausible parameter range, $0 < s_1, s_2 < 1$).
1011

Antagonistic pleiotropy (AP)

1013　Following previous models (e.g. Curtsinger et al., 1994; Rose, 1982), we consider a locus with
1014　antagonistic pleiotropic effects on two major fitness components. Selection is directional
1015　within each component (Tab. A1) and fitness components combine multiplicatively to
1016　determine overall fitness. The evolutionary dynamics are described by:

$$\Delta p = \frac{\left((t_1 + t_2)(1 - 2h) + 2t_1t_2h^2\right)p(1-p)(\hat{p}-p)}{\overline{w}}$$

1017

1018
$$\approx \left((t_1 + t_2)(1 - 2h) + 2t_1t_2h^2\right)p(1-p)(\hat{p}-p) \qquad (3)$$

1019   where $\overline{w} = p^2(1 - t_2) + 2p(1-p)(1 - t_1h)(1 - t_2h) + (1-p)^2(1 - t_1)$ and   the

1020   polymorphic equilibrium (whenever it is valid) is:

1021
$$\hat{p} = \frac{t_1(1 - h) - t_2h + t_1t_2h^2}{(t_1 + t_2)(1 - 2h) + 2t_1t_2h^2} \qquad (4)$$

1022   The equilibrium is valid under the following condition:

1023
$$\frac{t_2h}{1 - h + t_2h^2} < t_1 < \frac{t_2(1 - h)}{h(1 - t_2h)} \qquad (5)$$

1024   and otherwise, selection on total fitness is directional and favours fixation of either A_1 or A_2.

1025

Sexually antagonistic selection (SA)

Following Kidwell et al. (1977), sexually antagonistic alleles are under opposing directional

selection in males and females, characterised by a selection and dominance coefficient for each

sex (Tab. A1). Letting $p = (p_f + p_m)/2$ represent the overall frequency of A_1 in the population,

where p_f and p_m represent the respective frequencies in female and male gametes contributing

to fertilization, the exact allele frequency dynamics are given by $\Delta p = \frac{1}{2}(\Delta p_f + \Delta p_m)$, where:

1032
$$\Delta p_f = \frac{p_mp_f + (p - p_mp_f)(1 - s_fh_f)}{p_mp_f + 2(p - p_mp_f)(1 - s_fh_f) + (1 - p_f)(1 - p_m)(1 - s_f)} - p_f \qquad (6a)$$

1033   and

1034
$$\Delta p_m = \frac{p_mp_f(1 - s_m) + (p - p_mp_f)(1 - s_mh_m)}{p_mp_f(1 - s_m) + 2(p - p_mp_f)(1 - s_mh_m) + (1 - p_f)(1 - p_m)} - p_m \qquad (6b)$$

1035   Within the range $0 \leq h = h_f = h_m \leq 0.5$ which we consider here, and assuming modest to

1036   weak selection ($0 < s_f, s_m << 1$), the allele frequency dynamics are closely approximated by:

1037
$$\Delta p \approx \left(s(1 - 2h) + 2h^2s_fs_m\right)p(1-p)(\hat{p}-p) \qquad (7)$$

1038   (Connallon and Clark 2013), where $s = \frac{1}{2}(s_f + s_m)$, and $\hat{p}$ represents the polymorphic

1039   equilibrium when valid. With no dominance in either sex ($h = 0.5$), the equilibrium frequency

1040   (when it exists) is $\hat{p} = \frac{(s_f - s_m + s_fs_m)}{2s_fs_m}$, and for cases involving dominance reversals ($h < 0.5$), the

1041   equilibrium is $\hat{p} \approx \frac{(s_f(1-h) - s_mh)}{(s_f + s_m)(1-2h)}$. The condition favouring a polymorphic equilibrium (based

1042   on the exact eqs. (6a-6b)) is:

$$1043 \qquad \frac{s_m h_m}{1 - h_f + s_m h_m} < s_f < \frac{s_m(1 - h_m)}{h_f(1 - s_m)} \qquad (8)$$

1044   (Kidwell et al. 1977; Connallon and Chenoweth 2009).

1045   If we assume that each selection coefficient is uniformly distributed with a mean of $\bar{s} =$

1046 $\frac{1}{2} s_{\max}$ (where $s_{\max}$ represents the maximum selection coefficient, $0 < s_{\max} \leq 1$), then the

1047   proportion of parameter space that results in balancing selection under AP will be:

$$1048 \qquad f_{bal} = 1 - \frac{2h^2 s_{\max} + 2(1 - h) \log\left(\frac{1 - h}{1 - h + h^2 s_{\max}}\right)}{s_{\max}^2 h^3} = \frac{1 - 2h}{1 - h} + \frac{4h^3 \bar{s}}{3(1 - h)^2} + O(s_{\max}^2)$$

1049   while that that results in balancing selection under SA is:

$$1050 \qquad f_{bal} = 1 - \frac{2h s_{\max} + 2(1 - h) \log\left(\frac{1 - h}{1 - h + h s_{\max}}\right)}{s_{\max}^2 h} = \frac{1 - 2h}{1 - h} + \frac{4h^2 \bar{s}}{3(1 - h)^2} + O(s_{\max}^2)$$

1051

Appendix 2 - Criteria for maintaining a long-term balanced polymorphism

A long-term polymorphism requires balancing selection to be both consistent over time and

strong enough to offset the tendency of genetic drift to eliminate the polymorphism. Provided

balancing selection is temporally consistent, then its capacity to maintain polymorphism can

be quantified using standard diffusion theory (Rice, 2004; Wright, 1945). In the models

presented above, the expected change in allele frequencies at a single locus under balancing

selection (with change measured across a single generation) is:

$$1059 \qquad M = \alpha p(1 - p)(\hat{p} - p) \qquad (9)$$

where p again refers to the frequency of A_1, $p^{\wedge}$ is the equilibrium under balancing selection,

and α represents a model-specific constant that reflects the strength of balancing selection

(these can be extracted from eqs. (1), (3) and (7), which yield: $\alpha = s_1 + s_2$ for the

overdominance model, $\alpha = (t_1 + t_2)(1 - 2h) + 2t_1 t_2 h^2$ under antagonistic pleiotropy, and

$\alpha = s(1 - 2h) + 2s_f s_m h^2$ under sexual antagonism). The variance in allele frequency change

owing to drift is $V = \frac{p(1-p)}{2N_e}$, where N_e is the effective size of the population. Ignoring effects

of recurrent mutation, the stationary distribution loci subject to balancing selection (*i.e.*, the

long-term probability density function for allele frequency p) is:

$$1068 \qquad f(p) = \frac{C}{p(1 - p)} \exp\left((-2N_e \alpha(\hat{p} - p)^2)\right) \qquad (10)$$

(Wright 1945; Robertson 1962; Rice 2004) where the constant C ensures that the probability

density function integrates to one (*i.e.*, $C^{-1} = \int_0^1 [p(1 - p)]^{-1} \exp(-2N_e \alpha(\hat{p} - p)^2)\, dp$).

Balancing selection effectively maintains polymorphism when the probability density

near the equilibrium (p near $\hat{p}$) is high relative to the probability density near the boundaries of

the distribution ($p = 1/2N_e$ and $p = 1 - 1/2N_e$). A measure of these relative densities (following

Connallon & Clark, 2014) is given by:

1075
$$R \approx \frac{1}{2N_e\hat{p}(1-\hat{p})\left(exp(-2N_e\alpha\hat{p}^2) + exp(-2N_e\alpha(1-\hat{p})^2)\right)} \tag{11}$$

1076   for which balancing selection is effective when R is large ($R \gg 1$), and drift dominates when

1077 R is small ($R < 1$). Using a cut-off of $R > 10$ to represent effectively strong balancing selection,

1078   eq. (11) implies the following minimum values of α (a_{min}) are required to ensure the stability

1079   of a balanced polymorphism:

1080
$$a_{min} = \begin{cases} \dfrac{\log(20N_e\hat{p})}{2N_e\hat{p}^2} & \text{for } \hat{p} \leq 0.5 \\ \dfrac{\log\left(20N_e(1-\hat{p})\right)}{2N_e(1-\hat{p})^2} & \text{for } \hat{p} \geq 0.5 \end{cases} \tag{12}$$

From eq. (12), balancing selection should be effective at maintaining polymorphism when $\alpha >$

a_{min}, whereas drift will impede maintenance when α falls substantially below this threshold.

Figure 2 illustrates conditions for effective balancing selection across the spectrum of

polymorphic equilibrium frequency states ($0 < \hat{p} < 1$) and for three different effective

population sizes ($N_e = 1,000$; $N_e = 10,000$; $N_e = 100,000$). Two special cases of moderately

strong balancing selection in a small population ($N_e = 1,000$ and $\alpha = 0.05$) illustrate how the

probability density of the stationary distribution is concentrated near $p = \hat{p}$ when $\alpha > a_{min}$,

whereas it accumulates near $p = 0$ when $\alpha < a_{min}$ (Fig. 2B, top versus bottom panels).

1089

Appendix 3 - Linked variation

Long-term balanced polymorphisms can affect patterns of genetic diversity at linked and

otherwise neutrally evolving loci, leading to characteristic signatures of inflated heterozygosity

in genomic regions where the balanced polymorphism resides. With the exceptions of trans-

species polymorphisms, it is these signatures that are used by current methods for the inference

of balancing selection in genomic data (Bitarello et al., 2018; Cheng & DeGiorgio, 2020;

Siewert & Voight, 2017). For a balanced polymorphism with alleles A_1 and A_2 that are stably

maintained at equilibrium frequencies $\hat{p}$ and $1 - \hat{p}$ (respectively), the mean heterozygosity at

a nearby neutral site is approximately:

1099
$$\bar{H} = 4N_e\mu\left[1 + 2\hat{p}(1-\hat{p})\left(\frac{1}{2N_eR} + \frac{(1-2\hat{p})^2}{2 + 8N_eR\hat{p}(1-\hat{p})}\right)\right] \tag{13}$$

where R is the total recombinational distance between the neutral site and the site of the

balanced polymorphism, and μ is the mutation rate at the neutral site (the approximation, which

assumes that $R \gg \mu$, is obtained using mean coalescent times from p. 103 of Rice 2004).

The sum in the square brackets of eq. (13) determines the degree of inflation of

heterozygosity at the linked site relative to that of an unlinked locus, where the expected

heterozygosity would be $\bar{H} = 4N\mu$. Recombination rate and the equilibrium frequency of the

selected site also influence the width of the spike in neutral diversity that could be seen around

a long-term balanced polymorphism. Let $R = nr$ represent the cumulative recombination rate

between the selected site and a given neutral site, where n is the number of nucleotide sites

separating the two and r is the recombination rate per site (*i.e.*, the probability of a

recombination event between two directly adjacent nucleotide sites). Assuming that the

polymorphism is intermediate ($\hat{p} = 0.5$), then the window around the selected site over which

neutral diversity is elevated by at least two-fold (*i.e.*: where $\frac{\bar{H}}{4N\mu} > 2$) will be $L = \frac{1}{2N_e r}$

nucleotides, which implies that species with large effective population sizes and large per-

nucleotide recombination rates should exhibit particularly narrow windows of inflated

heterozygosity around loci maintaining long-term balanced polymorphisms.