Behavior evolves as a correlated response to selection on cuticle color in *Drosophila*

melanogaster* and *D. simulans

Sarah N. Ruckman[1*], Alexandra G. Duffy[2], Paulina Montes Mendez[1], Katelyn D. McCaffery[1],

Samuel Miller[1], Addison Crews[1], Nicholas Tan[1], Lauren A. Campbell[1], Ashley March[1],

Elizabeth B. Brown[1], and David Houle[1]

[1]Department of Biological Science, Florida State University, Tallahassee, FL, USA

[2]Department of Biological Sciences, North Carolina State University, Raleigh, NC, USA

*Email of corresponding author: sarahnruckman@gmail.com

Keywords: experimental evolution, color, behavior, pleiotropy

Abstract:

To predict adaptive evolution, we need to understand the degree to which selection on

one trait can constrain or redirect evolutionary responses in other traits. We used artificial

selection on cuticle color in *Drosophila melanogaster* and *D. simulans* to investigate whether

color and behavior evolve in tandem. We selected for light and dark thoracic colors for 16

generations in two populations per species and measured correlated responses in a suite of

behavioral traits, including aggression, basal activity, total activity, sleep, and geotaxis. Dark

selected individuals consistently showed higher aggression and basal activity than light selected

or control flies across both species and sexes, pointing to a modest but repeatable correlated

response that must be interpreted in light of low and variable aggression in our assay. In contrast,

patterns for geotaxis, sleep, and total activity evolved unpredictably, often varying across species

and populations, and showed no clear or uniform association with color. Taken together, our

results suggest that some behaviors may share a predictable relationship with color, whereas

others behave largely independently under the conditions we examined. The persistence of

correlated responses in aggression and basal activity in response to selection on color is

consistent with a conserved genetic basis, such as pleiotropy or tight linkage, although our data

do not, by themselves, distinguish among these mechanisms. Future work that identifies the loci

underlying variation in both color and behavior and tests their effects experimentally will be

important for clarifying how multivariate genetic covariance shapes the direction and pace of

adaptive evolution.

Introduction

A key goal of modern evolutionary biology is to predict if and how populations will

evolve adaptively in response to environmental change. Predicting adaptive evolution requires

identifying constraints that might limit a population's ability to respond to selection. Definitions

of adaptive constraints vary, but they all emphasize the same core concept: that populations are

not always able to respond to selection as one might predict from a simple model of adaptation

(Blows and Walsh 2009; Walsh and Blows 2009; Connallon and Hall 2018; Hansen and Pélabon

2021). One important source of long-term adaptive constraint is genetic covariance, which can

prevent traits from evolving independently or potentially slow adaptation (Houle 1991;

Connallon and Hall 2018; Hansen and Pélabon 2021).

Numerous empirical studies in both natural and experimental populations have tested

whether these genetic covariances constrain adaptation. In natural populations, selection on one

trait can cause changes in another trait, such as the linked evolution of beak morphology and

song in Darwin's finches (Podos 2001) or the covariation of body shape and armor in threespine

sticklebacks (Leinonen et al. 2011). Experimental evolution and artificial selection studies in

model organisms, such as *Drosophila melanogaster*, allow researchers to identify the nature of

underlying genomic differences associated with genetic correlations. For example, laboratory

selection for decreased development time in *D. melanogaster* led to correlated changes in body

size and lifespan, along with divergence across dozens of loci (Burke et al. 2010). Collectively,

these studies demonstrate that correlated trait responses are likely common and, thus, that our

naïve expectations about adaptation may be incorrect. Experimental studies are essential if we

are to fully understand how traits constrain adaptive evolution.

Color and behavior are two ecologically important traits that are often key targets of

selection and have been previously demonstrated to be correlated at the phenotypic level

(Ducrest et al. 2008; Roulin and Ducrest 2011; San-Jose and Roulin 2018; Ruckman et al. 2024).

One explanation for these correlations is that the biochemical pathways underlying coloration

(e.g., melanin synthesis) influence neurophysiology and behavior (Ducrest et al. 2008; San-Jose

and Roulin 2018). In invertebrates, for example, melanin coloration arises from dopamine

metabolism, and changes in gene expression within the pathway are linked to variation in color

intensity (Wittkopp and Beldade 2009; Takahashi 2013). Consequently, mutations in key

pigmentation genes, such as *ebony* and *yellow,* not only alter color but also cause predictable

behavioral defects, including abnormalities in circadian rhythms (Suh and Jackson 2007) and

mating behavior (Suh and Jackson 2007; reviewed in Takahashi 2013). However, a recent meta-

analysis across all of Animalia (Ruckman et al. 2024) revealed a consistent positive correlation

between color and behavior, which was not restricted to only melanin-based colors, challenging

the exclusivity of the melanin-pleiotropy hypothesis. Instead, the results are more consistent with

the 'badge of status' hypothesis as the most parsimonious explanation, where conspicuous colors

act as honest signals of an individual's aggressive potential (Rohwer 1975; McGraw et al. 2003;

Tibbetts and Dale 2004). Importantly, the badge of status hypothesis operates at the level of

social signaling and selection (how conspecifics respond to color), and is therefore compatible

with multiple proximate physiological mechanisms, including melanin pleiotropy and other

genetic or developmental pathways. In this view, physiological mechanisms may generate a

genetic covariance between color and aggression, while social selection on receivers' responses

maintains color as an honest badge of status. Consistent with this interpretation, Ruckman et al.

(2024) found no evidence for condition dependence, suggesting that the correlation between

color and aggression is more likely to reflect underlying genetic covariance than condition-based

signaling. While pleiotropy associated with the same genes across disparate color pathways (e.g.,

melanin and carotenoids) is unlikely, genetic covariance can also arise from linkage

disequilibrium, in which genes for coloration and aggression are inherited together because of

their close physical proximity. The conclusion from this meta-analysis, however, was inferred

from broad-scale correlational patterns, leaving the underlying genetic mechanism, pleiotropy or

linkage disequilibrium, unresolved.

Distinguishing between pleiotropy and linkage disequilibrium can be challenging when

testing for correlated responses to selection in a single species. Instead, experimental evolution

studies comparing responses among two or more related species subjected to the same selection

regime can be a powerful tool. Because linkage disequilibrium is generally not expected to be

conserved over longer evolutionary timescales (Falconer and Mackay 1996; Gaut and Long

2003; Slatkin 2008), a consistent pattern of correlated trait evolution observed repeatedly in

different taxa would provide strong evidence for pleiotropy (Lande 1979; Cheverud 1988, 1996;

Wagner and Zhang 2011). Here, we focused on two species of fruit flies, *D. melanogaster* and *D.*

simulans. These recently diverged species (estimates vary from 2-8 million years ago; Capy and

Gibert 2004; Obbard et al. 2012) are important models for experimental evolution (e.g., Burke et

al. 2010; Turner et al. 2011, 2013; Remolina et al. 2012; Barghi et al. 2017; Kelly and Hughes

2019; Bastide et al. 2022; Girardeau et al. 2025) due to their tractability for large-scale

experiments (Hoffmann 1987; Kravitz and Huber 2003; Kravitz and Fernandez 2015; Massey

and Wittkopp 2016). Furthermore, in *Drosophila*, both cuticle pigmentation (Wittkopp et al.

2011; Rajpurohit et al. 2016; Bastide et al. 2022) and behavior (Sokolowski 2001; Martin 2003;

Anholt and Mackay 2015; Kravitz and Fernandez 2015) are ecologically relevant traits that

influence fitness. By leveraging this two-species system, we used an experimental evolution

approach to test whether consistent patterns of behavioral change emerge following selection on

pigmentation and explored how those responses vary across species and sexes.

107   We hypothesized that selection for darker color would drive a correlated suite of

behavioral changes (aggression, basal activity, geotaxis, total activity, and sleep) in comparison

to light selected individuals for both *D. melanogaster* and *D. simulans*. We predicted dark

selected individuals would be more aggressive and have increased locomotor activity (Jordan et

al. 2006, 2007; Ducrest et al. 2008; Wittkopp and Beldade 2009; Takahashi 2013). Given

previous evidence of a negative correlation between locomotor activity and sleep in *D.*

melanogaster (Keene et al. 2010; Harbison et al. 2013), we predicted dark selected flies (i.e.,

more active flies) would sleep less. Finally, because activity level and geotaxis have been shown

to be genetically correlated in *D. melanogaster* (Murphey and Hall 1969; Toma et al. 2002;

Zhong et al. 2022), we predicted that dark selected flies would have a faster geotactic response.

The evolution of one or more of these behaviors in tandem with color would support our

hypothesis, providing direct experimental evidence that genetic covariance influences the

evolution of this suite of traits.

Methods

Fly maintenance

122   We housed each species in a separate incubator (Percival Model # 136VLC8Q9) on a 12-

hour light and 12-hour dark cycle. We maintained *D. melanogaster* populations (*D. mel*-1 and *D.*

mel-2) at a constant temperature of 25 °C, while *D. simulans* (*D. sim*-1 and *D. sim*-2) were

housed at a constant temperature of 24 °C (Krstevska and Hoffmann 1994; Ashburner and Roote

2007; Austin and Moehring 2013). We housed each treatment in a separate plexiglass population

cage (26.67 cm x 15.25 cm x 20.32 cm) and randomly assorted the cages into the incubator every

week. Bottles containing a standard Drosophila media composition (Ashburner and Roote 2007)

were added to each cage every week and were removed after three weeks.

Artificial Selection

To limit the effects of bottlenecks early in the experiment, we used a gradual selection

regime (Kofler and Schlötterer 2014; Vlachos and Kofler 2019). Each generation, we collected

750 virgin females and 750 males for each population (Supplemental Material for origin). We

randomly assigned them to three equal-sexed groups to serve as the parents for the three

treatments (Figure 1). In each population, we established two selection treatments (dark "D" and

light "L" selected) and one control "C" treatment. We performed selection on the cuticle

coloration of the dorsal thorax. This region of the cuticle is not sexually dimorphic (Wittkopp et

al. 2003; Massey and Wittkopp 2016). To perform selection, flies were anesthetized with CO_2.

We scored the color of the dorsal thorax by eye into four color categories, using the prominent

macrochaeta as landmarks (David et al. 1985; see Supplemental Material for details). Color

selection was conducted by a single individual and validated (n=100) for agreement with

objective color quantification measurements and repeatability (see Supplemental Material).

Visual bins differed from one another in the expected directions ($F_{3, 96}$= 138.19, $P<0.001$; Figure

S1) and were highly repeatable (ICC = 0.933; $F_{99,99.1}$ = 29.2, $p < 0.001$; Figure S2).

For D and L selection treatments, we gradually increased the coefficient of selection from

80% of 250 virgin females and 80% of 250 males to 8.4% of 250 virgin females and 8.4% of 250

males, over 16 generations. For example, for L, we selected the lightest 80% of 250 virgin

females and 80% of 250 males in the first generation of selection. By the 16th generation, we

selected only the lightest 8.4% of 250 virgin females and 8.4% of 250 males. C treatments flies

were randomly chosen with respect to thorax color in each generation, while still mimicking the

changes in population size experienced by the D and L treatments. Figure 1 illustrates this

selection scheme in detail and indicates the generations in which phenotypes other than thorax

color were assayed. No individuals used in the selection experiment were used in behavioral

tests.

Behavioral Assays

All behavioral tests were conducted on individuals 3-5 days old that were not used to

establish the next generation of flies. For all behavioral assays, flies were isolated for 24 hours

before testing. All tests were scored by observers who were blind to the hypothesis. However,

the observer was not blind to the color or sex of the individual as these features were readily

apparent.

Aggressive Behavior: Due to its previously reported association with color (Ducrest et al. 2008;

San-Jose and Roulin 2018; Ruckman et al. 2024), we were particularly interested in aggression,

and therefore, measured aggression at multiple generations (generations 5, 10, and 15; Figure 1)

using a standard protocol (Girardeau et al. 2025). To assay aggression, we paired a focal

individual with a marked opponent of the same sex, similar age, color, and size. Controlling for

sex, age, and size accounts for known behavioral covariates in *Drosophila* (Hoffmann 1991;

Chen et al. 2002; Baxter and Dukas 2017; Bath et al. 2021; Fernandez et al. 2023) Furthermore,

we paired individuals of similar color because interactions can be modulated by an opponent's

coloration, a principle established in other species (e.g., Lank et al. 1995; Sinervo and Lively

1996; Kraft et al. 2018; Sacchi et al. 2021; Scali et al. 2021).

The focal individual was assigned randomly. The opponent's left wing was marked to

distinguish it from the focal individual. Females were first aspirated into new vials without

media or water for two hours before conducting the trial to increase the likelihood of aggressive

behaviors (Ueda and Kidokoro 2002). After two hours, both females were aspirated into the

arena. For the male assays, flies were aspirated together into the arena 30 minutes after lights on

that day.

We used a standard aggression arena with a central food resource (see Supplemental

Material for details; Hutchins et al. 2024; Girardeau et al. 2025). Following a five-minute

acclimation period, we recorded the occurrence of fencing, lunging, boxing, and tussling (Chen

et al. 2002; see Supplemental Material for behavioral descriptions) every five minutes for one

hour (Altmann 1974).

Basal Activity Level: To measure basal activity level, we aspirated each individual into an open

field arena 3 hours after lights on that day (see Supplemental Material for details). After a 10-

minute acclimation period, the focal individuals were video-recorded for one minute, and we

counted the number of grid lines crossed as a measure of activity (Burnet et al. 1988; Anderson

et al. 2016).

Geotaxis Level: To measure geotaxis, we aspirated each individual into the geotaxis arena

(Taylor and Tuxworth 2019; Zhong et al. 2022; see Supplemental Material for details). After

tapping a fly to the bottom, we timed how long it took to climb a 2 cm distance. Trials were

considered censored (score of 0) after one minute if the individual did not climb the 2 cm

distance.

Total Activity Level and Sleep: We measured activity and sleep over 72 hours using the

Drosophila Activity Monitoring (DAM) system (Trikinetics, Waltham, MA). Flies were

individually housed in tubes with standard media and acclimated for 24 hours before data

collection. Total activity is measured as total beam crossings per day. Total sleep is defined as

the number of 5-minute bouts of inactivity per day (Pfeiffenberger et al. 2010a,b; see

Supplemental Material for details).

Color Quantification of Individuals Used in Behavioral Assays

Following behavioral assays, all individuals were photographed using a standardized

apparatus (WINGMACHINE; Houle et al. 2003). The mean grayscale value of the dorsal thorax

was quantified from these images using ImageJ after calibration to a color standard (Schneider et

al. 2012; see Supplemental Material for the imaging and analysis protocol).

Statistical Analyses

All statistical analyses were conducted using R version 4.5.0 (R Core Team 2025). To

test for direct and correlated responses to selection, we used a consistent generalized linear

model (GLM) framework for each trait. All models were analyzed separately for males and

females because the residual distributions differed between the sexes. For each trait, we visually

inspected residuals and fitted values to verify that model assumptions (e.g., normality,

homoscedasticity, goodness of fit) were met.

For the color and behavioral analyses, the full model for each trait related the dependent

variable (e.g., grayscale value, aggression counts) to the fixed effects of species, treatment, and

population (nested within species), including all two-way interactions. To test our prediction that

selection treatments would diverge linearly, we coded treatment as a continuous predictor with

values -1, 0, and 1 corresponding to L, C, and D lines, respectively. This parameterization

allowed us to estimate both the direction and magnitude of the evolutionary response to

selection. Models were fit by maximum likelihood, and we used F-tests for Gaussian models and

Wald χ^2 statistics for GLMs to assess the significance of main effects and interactions. Non-

significant interaction terms ($p > 0.05$) were removed before testing main effects.

We selected error distributions appropriate to each trait: a normal distribution for thorax

grayscale values and for total activity and sleep; a Poisson distribution with a log link for counts

of aggressive acts; and a negative binomial distribution for basal activity counts. For geotaxis,

we analyzed time-to-event data using a Cox proportional hazards mixed-effects model (coxme

package; Therneau 2024) with observer as a random effect and right-censoring of individuals

that did not complete the task within one minute.

When we detected a significant main effect or interaction involving treatment, we

conducted post-hoc comparisons using the emmeans package (Lenth 2025). For Gaussian and

negative binomial models, we report least-squares means and pairwise t-tests; for Poisson

models, we used asymptotic z-tests with effectively infinite degrees of freedom. When treatment-

by-population interactions were significant, we further characterized among-population

differences in response to selection using simple slopes analyses implemented with the emtrends

function in emmeans, which estimates the treatment slope within each population and tests it

using Wald X^2 statistics.

To examine the temporal dynamics of the response to selection and differences in

evolutionary trajectories among treatments, we fit models similar to those above, but also

including generation and generation^2 as predictors, along with their interactions with treatment.

For thorax color, we used a Gaussian model, and for aggression, we used a Poisson GLM with a

log link. Full model specifications, diagnostics, and trait-specific details (color, aggression, basal

activity, total activity, sleep, and geotaxis) are provided in the Supplementary Methods.

Results

Evolution of Color in Selection Treatments

To examine the temporal dynamics of the direct response to selection, we analyzed the

divergence in thorax color between D and L selected lines at generations 5, 10, and 15. Since the

full statistical model revealed complex interactions (Table S1), we visualized the response by

plotting the mean difference (contrast) in color between D and L treatments for each population

over time (Figure 2). In both sexes, the divergence between D and L lines was significant by

generation 5 and increased in a roughly linear fashion throughout the experiment (Table S2;

Figure 2). In males, the pattern was broadly consistent, though the *D. sim*-2 population tended to

exhibit less divergence than the other three populations in most generations (Figure 2A). The

response in females was remarkably consistent across all four populations (Figure 2B).

Having established that D and L lines diverged steadily in thorax color over 15

generations, we next examined the magnitude and among-population structure of color

differences at the end of the experiment (Table S3; Figure 3). There was a significant effect of

treatment by population nested in species interaction (males: $F_{2,345} = 12.43$, $p < 0.0001$; females:

$F_{2,337} = 5.20$, $p = 0.006$), indicating that the magnitude of the response to selection varied among

populations within a species. However, our analysis of rates of response was in the same

direction and was significantly negative in all cases (all slopes < -19.4, all $X^2_1 > 90.5$, all $p <$

0.0001; Table S4). Post-hoc tests further confirmed that for all populations and sexes, D

treatment flies were significantly darker than L and C treatment flies (Table S5; Figure 3), and L

treatment flies were significantly lighter than C treatment flies (Table S5; Figure 3).

260   Despite the consistent direction of the response, there were notable differences among

groups that explain the significant interaction. For instance, *D. melanogaster* males were, on

average, lighter than *D. simulans* males, a difference driven primarily by the L and C treatments

(Table S4; Figure 3A). The response in *D. mel*-1 males was particularly strong, with a selection

slope twice as steep as any other population (-46.2 vs. -21.9; Table S4). Among females, there

was a significant difference between the replicate *D. simulans* populations, with *D. sim*-

1 females tending to be darker than *D. sim*-2 females across all treatments (Tables S5 and S6;

Figure 3B).

Correlated Evolution of Aggression in Selection Treatments

269   Due to our *a priori* prediction about the correlated evolution of cuticle color and

aggression, we assayed aggression across multiple generations during the selection experiment.

To simplify interpretation (Table S7), we visualized the temporal response by plotting the

divergence in the number of behaviors (contrast) between the D and L treatments over time

(Figure 4). In males, this revealed a non-linear response that differed between the species (Figure

4A). The divergence in both species decreased slightly at generation 10 and was greatest at

generation 15 (Figure 4). Despite divergence in color at generation 5, *D. simulans* populations

did not diverge in aggressive behavior until generation 15 (Table S8; Figure 4A). However, *D.*

melanogaster populations diverged in both color and behavior by generation 5 and consistently

showed greater divergence than *D. simulans* populations (Table S8; Figure 4A). Females

exhibited a similar non-linear trajectory, with the divergence between D and L lines also peaking

at generation 15 after a minimum at generation 10 (Figure 4B). For females, however, the

variation was most pronounced between populations rather than species (Figure 4B). The *D. mel*-

1 population, in particular, exhibited a greater divergence between D and L treatments at each

generation than the other three populations (Table S8; Figure 4B).

Given these nonlinear divergence trajectories and their peak at generation 15, we next

examined the magnitude and among-population structure of aggression differences at the end of

the experiment (Table S9; Figure 5). There was a significant main effect of selection treatment

on aggression for both sexes (males: $X^2_1 = 50.3$, $p < 0.0001$; females: $X^2_1 = 17.5$, $p < 0.0001$;

Table S9). This response was consistent across all populations, as the treatment by population

interaction was not significant in initial models (males: $X^2_2 = 0.23$, $p = 0.89$; females: $X^2_2 = 0.13$,

$p = 0.94$) and was therefore dropped from the final model. As predicted, we observed a

significant positive slope for both males (slope $= 1.52$, $X^2_1 = 31.5$, $p < 0.0001$; Table S10) and

females (slope $= 1.1$, $X^2_1 = 13.3$, $p = 0.0003$; Table S10). Post-hoc tests revealed that D

individuals were significantly more aggressive than L and C treatment individuals, which did not

differ from each other (Table S11; Figure 5).

Correlated Evolution of Basal Activity Level

At generation 12, basal activity showed a significant correlated response to selection

(Table S12; Figure 6). This response was consistent across all populations, as the treatment by

population interaction in initial models was not significant (males: $X^2_2 = 1.35$, $p = 0.51$; females:

$X^2_2 = 0.10$, $p = 0.95$) and was therefore dropped from the final models. As predicted, we observed

a significant positive slope for both males (slope $= 0.51$, $X^2_1 = 34.0$, $p < 0.0001$; Table S13) and

females (slope $= 0.27$; $X^2_1 = 11.7$, $p = 0.0006$; Table S13). Post-hoc tests showed that L

individuals did not differ from C individuals (Table S14). In males, D individuals were

significantly more active than both L and C individuals, which did not differ from each other

(Figure 6A). In females, D individuals were significantly more active than L individuals, but not

C individuals (Figure 6B).

Inconsistent Correlated Responses in Other Behaviors

In contrast to aggression and basal activity, the correlated responses for geotaxis, total

activity, and sleep were inconsistent and unpredictable, with responses varying across species,

populations, and sexes (see Supplemental Material; Tables S15-S23; Figures S4-S9).

Discussion

Our experiment demonstrates that artificial selection on cuticle color in *D. melanogaster*

and *D. simulans* is accompanied by consistent correlated changes in some, but not all, behaviors.

Selection for darker coloration led to a consistent increase in aggression at generation 15 and

basal activity in both sexes of both species. In contrast, other behaviors, including geotaxis,

sleep, and total activity, evolved unpredictably, with responses varying among species,

populations, and sexes.

The consistent evolution of heightened aggression by generation 15 in dark selected flies

across both species suggests a genetic association between melanin-based color and aggressive

behavior. However, the strength of this association is modest. Absolute aggression levels in our

assay were low, and trajectories through time were noisy, which limits how strongly we can infer

a tight or deterministic link between color and aggression. Nonetheless, the fact that dark lines

consistently tended to be more aggressive than control and light lines in both species, despite low

baseline aggression levels and demographic noise, is consistent with a conserved color-

aggression association. Within D lines, males were consistently more aggressive than females on

average, suggesting that sex specific expression modulates the magnitude of the response while

preserving the direction of the association between color and aggression. Similar context

dependence in the evolution of male aggression has also been documented when aggression itself

is the direct target of selection in our *D. mel*-2 base population (Girardeau et al. 2025). In that

study, aggression was selected using three different "feedback" regimes: i) positive feedback,

where evolving males always fought other males from the same evolving population, so that

increasing aggression amplified retaliatory interactions over generations; ii) negative feedback,

where males first experienced a conspecific fight and were then tested and selected against a

standard opponent, so that evolving increases in aggression strengthened prior-experience

suppression of subsequent aggression; and iii) no feedback, where males were always tested and

selected against a standard, non-evolving opponent, eliminating indirect genetic effect-mediated

evolutionary feedbacks. These different social contexts generated distinct evolutionary

trajectories: the rate and pattern of change in total aggression differed markedly among

treatments and were often noisy rather than smoothly directional. This suggests that treatment-

specific responses are a general feature of this trait rather than unique to our selection

experiment.

Our data revealing an association with cuticle color and aggression aligns with

predictions from the melanin-pleiotropy hypothesis (Takahashi 2013; San-Jose and Roulin

2018), which posits a shared biochemical basis for the traits, and is also compatible with

scenarios in which color could function as a status-related signal, with melanin-based coloration
as the signal and aggressive propensity as the underlying quality (Rohwer 1975; Ruckman et al.
2024). However, given the low and variable aggression levels in our experiment, our data should
be viewed as suggestive rather than definitive evidence for these mechanisms. At present, they
indicate that darker cuticle coloration is associated with a greater propensity to engage in
aggressive acts under our assay conditions, but they do not by themselves establish that melanin-
based traits are universally reliable "badges of status" or that pleiotropy is the sole cause of the
observed covariance.

The correlated response in aggression was notably asymmetric in both species: D lines
became more aggressive, whereas L lines did not become less aggressive than C lines. The lack
of response in L lines is consistent with another experiment using the same assay in *D. mel*-2
(Girardeau et al. 2025). A likely explanation is that the base populations exhibited very low
aggression under our assay conditions, such that further reductions in the underlying tendency to
act aggressively would be difficult to resolve with our behavioral measure. In other words, we
may be closer to a floor than a midpoint in aggression, which would naturally make upward
responses easier to detect than downward ones.

In contrast to the linear divergence in color, the correlated response in aggression over
time was non-linear, peaking at generation 15 after a dip at generation 10. Environmental
artifacts (e.g., humidity or temperature fluctuations) are one possible explanation, but all lines
were maintained and assayed under tightly controlled conditions. Such inconsistent trajectories
are often observed for correlated traits (Falconer and Mackay 1996), whose evolution is
mediated by the interplay between indirect selection, genetic drift in finite populations (e.g., Yoo
1980; Weber 1996), and the potential for the underlying genetic covariance (the G-matrix) itself
to change over time (Steppan et al. 2002; Arnold et al. 2008). One possible future direction is to
continue selection for additional generations to reduce noise in the correlated responses. Even
with this noisy background, the repeated tendency for D lines in both species to show higher
aggression than light and control lines suggests a real association between pigmentation and
aggression rather than a purely transient effect.

The consistent correlated evolutionary response in basal activity in D flies is consistent
with a shared genetic basis between color and activity. This is in agreement with previous work
that identified *Dopa decarboxylase (Ddc)*, a gene essential for both dopamine synthesis and

melanin production, as a key locus associated with locomotor behavior in *D. melanogaster*
(Jordan et al. 2006, 2007). The finding that many genes associated with locomotor behavior are
also linked to aggression and neural development further suggests potential widespread genetic
integration among these traits (Jordan et al. 2007). Our finding that dark males were more active
than dark females in both species, and that dark lines consistently showed elevated basal activity,
indicates that these genetic connections between pigmentation and locomotor behavior are
expressed in a sex specific but directionally conserved way across taxa.

In contrast, we did not observe a consistent correlated response in total activity, which
was measured using the DAM system. This result is consistent with the findings of Dierick and
Greenspan (2006), who selected directly for high aggression and used an older, but similar,
DAM monitoring system to measure general activity. They also found no correlated increase in
total activity. Therefore, the genetic covariance between color and activity is likely context-
dependent, manifesting strongly in exploratory or basal activity but not in overall activity levels
measured in a confined tube environment.

The stochastic evolution of geotaxis, sleep, and total activity suggests that these traits
exhibit little to no conserved genetic covariance with cuticle color, although we cannot rule out
the possibility that, with continued selection across more generations, a modest association could
emerge. While our gradual selection design was intended to allow recombination to break down
transient, non-functional linkages (Kofler and Schlötterer 2014; Vlachos and Kofler 2019), the
low effective population size, particularly in the final five generations, amplifies the impact of
stochastic forces. These forces include genetic drift, but also stochasticity in the decay of any
initial linkage disequilibrium. Together, these processes are expected to cause random,
population-specific divergence in phenotypes not under strong, consistent indirect selection.
Geotaxis, sleep, and total activity in our experiment showed exactly this pattern of idiosyncratic,
population specific divergence with no consistent differences among color treatments. Their
trajectories likely reflect drift or transient linkage rather than a stable conserved genetic
association with pigmentation.

A productive direction for future research would be to determine the specific genetic
mechanisms underlying these divergent outcomes. A primary goal is to determine if the
consistent association between color, aggression, and basal activity is indeed caused by
pleiotropy. Evolve-and-resequence (E&R) studies could identify candidate genes, and

subsequent functional analyses, such as RNAi knockdowns, could then confirm their effects on

both color and behavior, clarifying the extent to which such correlated responses constrain or

facilitate evolution. Beyond pairwise relationships, it will be important to characterize

multivariate pleiotropy, in which single loci influence suites of behavioral, morphological, and

physiological traits, and to assess whether similar pleiotropic modules recur across taxa. Large-

scale mapping and association studies in multiple species, combined with comparative analyses

of genetic covariance matrices, could reveal whether the pigmentation-behavior-physiology

integration we observe is part of a broader, conserved pleiotropic architecture.

414
415
Figure 1: Selection and Behavioral Workflow. Flies were collected from wild populations to
create each population. They were then tested for aggression and split into three treatments (light
selected "L", control "C", and dark selected "D"). The L treatment is represented by pink. The C
treatment is represented by purple. The D treatment is represented by blue. Solid lines indicate
the selection and establishment of a new generation. Dashed lines and dashed boxes indicate
behavioral assays. Selection gradually increased, so fewer individuals were kept every
generation (decreasing percentage of individuals kept every generation). The L treatment had the
lightest offspring selected, while the D treatment had the darkest offspring selected each
generation. The C treatment was randomly chosen each generation. This workflow is designed
for a single population of *D. melanogaster*. This same workflow was used for each population
per species.
427
428

429
Figure 2: Difference (contrast) in mean thorax color between D and L flies within a population at
a specific generation of selection. The colors represent the different populations. The x-axis
represents the generation number in which traits were assayed, and the y-axis represents the D -
L color contrast (mean ± SE). The mean is the difference between the raw data mean of the D
and L treatments. The error bars represent the standard errors of raw data. However, as described
in the Methods, the statistical analysis was conducted using the individual fly color measure as
the dependent variable. A negative contrast indicates that D flies were darker than L flies. The
left panel shows contrast values for males of both populations of both species, while the right
panel shows the contrasts for females.
439

Figure 3: Thoracic color evolution at generation 15 for A) *D. melanogaster* and B) *D. simulans*. The x-axis indicates the different treatments (L, C, and D). The y-axis represents the average grey-scale value, with lighter values (higher on the scale) at the top and darker values (lower on the scale) at the bottom. Photos on the y-axis represent individuals before grey-scale conversion in ImageJ. The letters above and below the data indicate significance, where groups with different letters are significantly different from each other. Uppercase letters indicate the significance of *D. mel*-1 and *D. sim*-1, and lowercase letters indicate the significance of *D. mel*-2 and *D. sim*-2 populations.

453
Figure 4: Difference (contrast) in mean aggression counts between D and L flies within a
population at a specific generation of selection. The colors represent the different populations.
The x-axis represents the generation number in which traits were assayed, and the y-axis
represents the D - L color contrast (mean ± SE). The mean is the difference between the raw data
mean of the D and L treatments. The error bars represent the standard errors of the raw data.
However, as described in the Methods, the statistical analysis was conducted using the individual
fly aggression counts as the dependent variable. A positive contrast indicates that D treatment
flies were more aggressive than L flies. The left panel shows contrast values for males of both
populations of both species, while the right panel shows the contrasts for females.
463

464
465 Figure 5: Aggression at generation 15 for A) Male analysis. B) Female analysis. The x-axis
466 indicates the different treatments (L, C, and D). The y-axis represents the average aggression
467 counts (number of aggressive acts). The letters above the data indicate significance, where
468 groups with different letters are significantly different from each other.
469
470
471
472

473
474 Figure 6: Basal activity level at generation 12 for A) Male analysis. B) Female analysis. The x-
475 axis indicates the different treatments (L, C, and D). The y-axis represents the average basal
476 activity level (number of lines crossed in one minute). The letters above the data indicate
477 significance, where groups with different letters are significantly different from each other.
478
479

Acknowledgements: We thank Kimberly Hughes, Michelle Arbeitman, and Darin Roykta for their helpful comments throughout the data collection process and for valuable suggestions that improved the manuscript. We thank the editor and the two reviewers for their careful reading and constructive comments, which improved this manuscript. We thank our funding sources NSF DEB-1740466, DEB-2217558, and the Florida State University Council on Research and Creativity SEED and COFRS grants.

References:

Altmann, J. 1974. Observational study of behavior: sampling methods. Behaviour 49:227–266. Brill.

Anderson, B. B., A. Scott, and R. Dukas. 2016. Social behavior and activity are decoupled in larval and adult fruit flies. Behav. Ecol. 27:820–828.

Anholt, R. R., and T. F. Mackay. 2015. Dissecting the genetic architecture of behavior in *Drosophila melanogaster*. Curr. Opin. Behav. Sci. 2:1–7.

Arnold, S. J., R. Bürger, P. A. Hohenlohe, B. C. Ajie, and A. G. Jones. 2008. UNDERSTANDING THE EVOLUTION AND STABILITY OF THE G-MATRIX. Evolution 62:2451–2461.

Ashburner, M., and J. Roote. 2007. Maintenance of a Drosophila Laboratory: General Procedures. Cold Spring Harb. Protoc. 2007:pdb.ip35. Cold Spring Harbor Laboratory Press.

Austin, C. J., and A. J. Moehring. 2013. Optimal temperature range of a plastic species, *Drosophila simulans*. J. Anim. Ecol. 82:663–672.

Barghi, N., R. Tobler, V. Nolte, and C. Schlötterer. 2017. Drosophila simulans: A Species with Improved Resolution in Evolve and Resequence Studies. G3 GenesGenomesGenetics 7:2337–2343.

506   Bastide, H., D. Ogereau, C. Montchamp-Moreau, and P. R. Gérard. 2022. The fate of a

507   suppressed X-linked meiotic driver: experimental evolution in Drosophila simulans.

508   Chromosome Res. 30:141–150.

509   Bath, E., D. Edmunds, J. Norman, C. Atkins, L. Harper, W. G. Rostant, T. Chapman, S. Wigby,

510   and J. C. Perry. 2021. Sex ratio and the evolution of aggression in fruit flies. Proc. R.

511   Soc. B Biol. Sci. 288:20203053. Royal Society.

512   Baxter, C. M., and R. Dukas. 2017. Life history of aggression: effects of age and sexual

513   experience on male aggression towards males and females. Anim. Behav. 123:11–20.

514   Blows, M., and B. Walsh. 2009. Spherical Cows Grazing in Flatland: Constraints to Selection

515   and Adaptation. Pp. 83–101 *in* J. van der Werf, H.-U. Graser, R. Frankham, and C.

516   Gondro, eds. Adaptation and Fitness in Animal Populations: Evolutionary and Breeding

517   Perspectives on Genetic Resource Management. Springer Netherlands, Dordrecht.

518   Burke, M. K., J. P. Dunham, P. Shahrestani, K. R. Thornton, M. R. Rose, and A. D. Long. 2010.

519   Genome-wide analysis of a long-term evolution experiment with Drosophila. Nature

520   467:587–590. Nature Publishing Group.

521   Burnet, B., L. Burnet, K. Connolly, and N. Williamson. 1988. A genetic analysis of locomotor

522   activity in Drosophila melanogaster. Heredity 61:111–119. Nature Publishing Group.

523   Capy, P., and P. Gibert. 2004. *Drosophila melanogaster, Drosophila simulans*: so similar yet so

524   different. Pp. 5–16 *in* P. Capy, P. Gibert, and I. Boussy, eds. Drosophila melanogaster,

525   Drosophila simulans: So Similar, So Different. Springer Netherlands, Dordrecht.

526   Chen, S., A. Y. Lee, N. M. Bowens, R. Huber, and E. A. Kravitz. 2002. Fighting fruit flies: A

527   model system for the study of aggression. Proc. Natl. Acad. Sci. 99:5664–5668. National

528   Academy of Sciences.

Cheverud, J. M. 1988. A Comparison of Genetic and Phenotypic Correlations. Evolution

42:958–968.

Cheverud, J. M. 1996. Developmental Integration and the Evolution of Pleiotropy. Am. Zool.

36:44–50. Oxford University Press.

Connallon, T., and M. D. Hall. 2018. Genetic constraints on adaptation: a theoretical primer for

the genomics era. Ann. N. Y. Acad. Sci. 1422:65–87. Wiley-Blackwell.

David, J. R., P. Capy, V. Payant, and S. Tsakas. 1985. Thoracic trident pigmentation in

Drosophila melanogaster: Differentiation of geographical populations. Génétique

Sélection Évolution 17:211.

Dierick, H. A., and R. J. Greenspan. 2006. Molecular analysis of flies selected for aggressive

behavior. Nat. Genet. 38:1023–1031.

Ducrest, A.-L., L. Keller, and A. Roulin. 2008. Pleiotropy in the melanocortin system, coloration

and behavioural syndromes. Trends Ecol. Evol. 23:502–510.

Falconer, D. S., and T. F. C. Mackay. 1996. Introduction to quantitative genetics. Longmans

Green, Harlow, Essex, UK.

Fernandez, M. P., S. Trannoy, and S. J. Certel. 2023. Fighting Flies: Quantifying and Analyzing

Drosophila Aggression. Cold Spring Harb. Protoc. 2023:pdb.top107985. Cold Spring

Harbor Laboratory Press.

Gaut, B. S., and A. D. Long. 2003. The Lowdown on Linkage Disequilibrium. Plant Cell

15:1502–1506.

Girardeau, A. R., G. E. Enochs, and J. B. Saltz. 2025. Evolutionary feedbacks for Drosophila

aggression revealed through experimental evolution. Proc. Natl. Acad. Sci.

122:e2419068122. Proceedings of the National Academy of Sciences.

Hansen, T. F., and C. Pélabon. 2021. Evolvability: A Quantitative-Genetics Perspective. Annu.

Rev. Ecol. Evol. Syst., doi: https://doi.org/10.1146/annurev-ecolsys-011121-021241.

Harbison, S. T., L. J. McCoy, and T. F. Mackay. 2013. Genome-wide association study of sleep

in Drosophila melanogaster. BMC Genomics 14:281.

Hoffmann, A. A. 1987. A laboratory study of male territoriality in the sibling species *Drosophila*

melanogaster and *D. simulans*. Anim. Behav. 35:807–818.

Hoffmann, A. A. 1991. Heritable Variation for Territorial Success in Field-Collected Drosophila

melanogaster. Am. Nat. 138:668–679. The University of Chicago Press.

Houle, D. 1991. Genetic Covariance of Fitness Correlates: What Genetic Correlations Are Made

of and Why It Matters. Evolution 45:630–648.

Houle, D., J. Mezey, P. Galpern, and A. Carter. 2003. Automated measurement of Drosophila

wings. BMC Evol. Biol. 3:25.

Hutchins, M., T. Douglas, L. Pollack, and J. B. Saltz. 2024. Genetic Variation in Male

Aggression Is Influenced by Genotype of Prior Social Partners in Drosophila

melanogaster. Am. Nat. 203:551–561. The University of Chicago Press.

Jordan, K. W., M. A. Carbone, A. Yamamoto, T. J. Morgan, and T. F. Mackay. 2007.

Quantitative genomics of locomotor behavior in Drosophila melanogaster. Genome Biol.

8:R172.

Jordan, K. W., T. J. Morgan, and T. F. C. Mackay. 2006. Quantitative Trait Loci for Locomotor

Behavior in *Drosophila melanogaster*. Genetics 174:271–284.

Keene, A. C., E. R. Duboué, D. M. McDonald, M. Dus, G. S. B. Suh, S. Waddell, and J. Blau.

2010. Clock and cycle Limit Starvation-Induced Sleep Loss in Drosophila. Curr. Biol.

20:1209–1215. Elsevier.

Kelly, J. K., and K. A. Hughes. 2019. Pervasive Linked Selection and Intermediate-Frequency

576   Alleles Are Implicated in an Evolve-and-Resequencing Experiment of *Drosophila*

577   *simulans*. Genetics 211:943–961.

578 Kofler, R., and C. Schlötterer. 2014. A Guide for the Design of Evolve and Resequencing

579   Studies. Mol. Biol. Evol. 31:474–483.

580 Kraft, B., V. A. Lemakos, J. Travis, and K. A. Hughes. 2018. Pervasive indirect genetic effects

581   on behavioral development in polymorphic eastern mosquitofish. Behav. Ecol. 29:289–

582   300.

583 Kravitz, E. A., and M. de la P. Fernandez. 2015. Aggression in Drosophila. Behav. Neurosci.

584   129:549–563. American Psychological Association, US.

585 Kravitz, E. A., and R. Huber. 2003. Aggression in invertebrates. Curr. Opin. Neurobiol. 13:736–

586   743.

587 Krstevska, B., and A. A. Hoffmann. 1994. The effects of acclimation and rearing conditions on

588   the response of tropical and temperate populations of *Drosophila melanogaster* and *D.*

589   *simulans* to a temperature gradient (Diptera: Drosophilidae). J. Insect Behav. 7:279–288.

590 Lande, R. 1979. Quantitative Genetic Analysis of Multivariate Evolution, Applied to Brain:

591   Body Size Allometry. Evolution 33:402–416. [Society for the Study of Evolution,

592   Wiley].

593 Lank, D. B., C. M. Smith, O. Hanotte, T. Burke, and F. Cooke. 1995. Genetic polymorphism for

594   alternative mating behaviour in lekking male ruff *Philomachus pugnax*. Nature 378:59–

595   62. Nature Publishing Group.

596 Leinonen, T., J. M. Cano, and J. Merilä. 2011. Genetics of body shape and armour variation in

597   threespine sticklebacks. J. Evol. Biol. 24:206–218.

Lenth, R. 2025. emmeans: Estimated Marginal Means, aka Least-Squares Means.

Martin, J.-R. 2003. Locomotor activity: a complex behavioural trait to unravel. Behav. Processes

64:145–160.

Massey, J. H., and P. J. Wittkopp. 2016. The Genetic Basis of Pigmentation Differences Within

and Between Drosophila Species. Curr. Top. Dev. Biol. 119:27–61.

McGraw, K. J., J. Dale, and E. A. Mackillop. 2003. Social environment during molt and the

expression of melanin-based plumage pigmentation in male house sparrows (*Passer*

domesticus). Behav. Ecol. Sociobiol. 53:116–122.

Murphey, R. M., and C. F. Hall. 1969. Some correlates of negative geotaxis in *Drosophila*

melanogaster. Anim. Behav. 17:181–185.

Obbard, D. J., J. Maclennan, K.-W. Kim, A. Rambaut, P. M. O'Grady, and F. M. Jiggins. 2012.

Estimating Divergence Dates and Substitution Rates in the Drosophila Phylogeny. Mol.

Biol. Evol. 29:3459–3473.

Pfeiffenberger, C., B. C. Lear, K. P. Keegan, and R. Allada. 2010a. Locomotor Activity Level

Monitoring Using the Drosophila Activity Monitoring (DAM) System. Cold Spring Harb.

Protoc. 2010:pdb.prot5518. Cold Spring Harbor Laboratory Press.

Pfeiffenberger, C., B. C. Lear, K. P. Keegan, and R. Allada. 2010b. Processing Sleep Data

Created with the Drosophila Activity Monitoring (DAM) System. Cold Spring Harb.

Protoc. 2010:pdb.prot5520. Cold Spring Harbor Laboratory Press.

Podos, J. 2001. Correlated evolution of morphology and vocal signal structure in Darwin's

finches. Nature 409:185–188. Nature Publishing Group.

R Core Team. 2025. R: A Language and Environment for Statistical Computing. R Foundation

for Statistical Computing, Vienna, Austria.

Rajpurohit, S., L. M. Peterson, A. J. Orr, A. J. Marlon, and A. G. Gibbs. 2016. An Experimental

Evolution Test of the Relationship between Melanism and Desiccation Survival in

Insects. PLOS ONE 11:e0163414. Public Library of Science.

Remolina, S. C., P. L. Chang, J. Leips, S. V. Nuzhdin, and K. A. Hughes. 2012. GENOMIC

BASIS OF AGING AND LIFE-HISTORY EVOLUTION IN DROSOPHILA

MELANOGASTER. Evolution 66:3390–3403.

Rohwer, S. 1975. The Social Significance of Avian Winter Plumage Variability. Evolution

29:593–610. [Society for the Study of Evolution, Wiley].

Roulin, A., and A.-L. Ducrest. 2011. Association between melanism, physiology and behaviour:

A role for the melanocortin system. Eur. J. Pharmacol. 660:226–233.

Ruckman, S. N., E. A. Humphrey, L. Muzzey, I. Prantalou, M. Pleasants, and K. A. Hughes.

2024. Assessing the Association Between Animal Color and Behavior: A Meta-Analysis

of Experimental Studies. Ecol. Evol. 14:e70655.

Sacchi, R., A. J. Coladonato, M. Battaiola, C. Pasquariello, S. Buratti, C. Matellini, M.

Mangiacotti, S. Scali, and M. A. L. Zuffi. 2021. Subjective resource value affects

aggressive behavior independently of resource-holding-potential and color morphs in

male common wall lizard. J. Ethol. 39:179–189.

San-Jose, L. M., and A. Roulin. 2018. Toward Understanding the Repeated Occurrence of

Associations between Melanin-Based Coloration and Multiple Phenotypes. Am. Nat.

192:111–130. The University of Chicago Press.

Scali, S., M. Mangiacotti, R. Sacchi, A. J. Coladonato, M. Falaschi, L. Saviano, M. G. Rampoldi,

M. Crozi, C. Perotti, F. Zucca, E. Gozzo, and M. A. L. Zuffi. 2021. Close encounters of

the three morphs: Does color affect aggression in a polymorphic lizard? Aggress. Behav.

47:430–438.

Schneider, C. A., W. S. Rasband, and K. W. Eliceiri. 2012. NIH Image to ImageJ: 25 years of

image analysis. Nat. Methods 9:671–675. Nature Publishing Group.

Sinervo, B., and C. M. Lively. 1996. The rock–paper–scissors game and the evolution of

alternative male strategies. Nature 380:240–243. Nature Publishing Group.

Slatkin, M. 2008. Linkage disequilibrium — understanding the evolutionary past and mapping

the medical future. Nat. Rev. Genet. 9:477–485. Nature Publishing Group.

Sokolowski, M. B. 2001. Drosophila: Genetics meets behaviour. Nat. Rev. Genet. 2:879–890.

Nature Publishing Group.

Steppan, S. J., P. C. Phillips, and D. Houle. 2002. Comparative quantitative genetics: evolution

of the G matrix. Trends Ecol. Evol. 17:320–327. Elsevier.

Suh, J., and F. R. Jackson. 2007. Drosophila Ebony Activity Is Required in Glia for the

Circadian Regulation of Locomotor Activity. Neuron 55:435–447. Elsevier.

Takahashi, A. 2013. Pigmentation and behavior: potential association through pleiotropic genes

in *Drosophila*. Genes Genet. Syst. 88:165–174.

Taylor, M. J., and R. I. Tuxworth. 2019. Continuous tracking of startled Drosophila as an

alternative to the negative geotaxis climbing assay. J. Neurogenet. 33:190–198. Taylor &

Francis.

Therneau, T. M. 2024. coxme: Mixed Effects Cox Models.

Tibbetts, E. A., and J. Dale. 2004. A socially enforced signal of quality in a paper wasp. Nature

432:218–222. Nature Publishing Group.

Toma, D. P., K. P. White, J. Hirsch, and R. J. Greenspan. 2002. Identification of genes involved

in Drosophila melanogaster geotaxis, a complex behavioral trait. Nat. Genet. 31:349–353.

Nature Publishing Group.

Turner, T. L., P. M. Miller, and V. A. Cochrane. 2013. Combining Genome-Wide Methods to

Investigate the Genetic Complexity of Courtship Song Variation in Drosophila

melanogaster. Mol. Biol. Evol. 30:2113–2120.

Turner, T. L., A. D. Stewart, A. T. Fields, W. R. Rice, and A. M. Tarone. 2011. Population-

Based Resequencing of Experimentally Evolved Populations Reveals the Genetic Basis

of Body Size Variation in Drosophila melanogaster. PLOS Genet. 7:e1001336. Public

Library of Science.

Ueda, A., and Y. Kidokoro. 2002. Aggressive behaviours of female Drosophila melanogaster are

influenced by their social experience and food resources. Physiol. Entomol. 27:21–28.

Vlachos, C., and R. Kofler. 2019. Optimizing the Power to Identify the Genetic Basis of

Complex Traits with Evolve and Resequence Studies. Mol. Biol. Evol. 36:2890–2905.

Wagner, G. P., and J. Zhang. 2011. The pleiotropic structure of the genotype–phenotype map:

the evolvability of complex organisms. Nat. Rev. Genet. 12:204–213. Nature Publishing

Group.

Walsh, B., and M. W. Blows. 2009. Abundant Genetic Variation + Strong Selection =

Multivariate Genetic Constraints: A Geometric View of Adaptation. Annu. Rev. Ecol.

Evol. Syst. 40:41–59.

Weber, K. E. 1996. Large Genetic Change at Small Fatness Cost in Large Populations of

Drosophila melanogaster Selected for Wind Tunnel Flight: Rethinking Fitness Surfaces.

Genetics 144:205–213.

Wittkopp, P. J., and P. Beldade. 2009. Development and evolution of insect pigmentation:

Genetic mechanisms and the potential consequences of pleiotropy. Semin. Cell Dev.

Biol. 20:65–71.

Wittkopp, P. J., S. B. Carroll, and A. Kopp. 2003. Evolution in black and white: genetic control

of pigment patterns in Drosophila. Trends Genet. 19:495–504.

Wittkopp, P. J., G. Smith-Winberry, L. L. Arnold, E. M. Thompson, A. M. Cooley, D. C. Yuan,

Q. Song, and B. F. McAllister. 2011. Local adaptation for body color in *Drosophila*

americana. Heredity 106:592–602.

Yoo, B. H. 1980. Long-term selection for a quantitative character in large replicate populations

of Drosophila melanogaster: 1. Response to selection. Genet. Res. 35:1–17.

Zhong, L., Z. Yang, H. Tang, Y. Xu, X. Liu, and J. Shen. 2022. Differential analysis of negative

geotaxis climbing trajectories in Drosophila under different conditions. Arch. Insect

Biochem. Physiol. 111:e21922.

701