Genomic and phenotypic diversification of *Pseudomonas aeruginosa* during sustained exposure to a ciliate predator

Nayma Romo Bechara[1], Nikhil Bardeskar[1], Heather A. Hopkins[2], Kasie Raymann*[1]

1. Department of Plant and Microbial Biology, North Carolina State University, Raleigh, NC
2. Department of Biology, University of North Carolina Greensboro, Greensboro, NC

*Corresponding author: Kasie Raymann, ktrayman@ncsu.edu

1 Abstract

Opportunistic bacterial pathogens often encounter strong selective pressures outside
their hosts, yet the evolutionary consequences of long-term predator exposure remain
poorly understood. Here, we used experimental evolution to examine how sustained
interaction with a eukaryotic predator shapes genomic adaptation, phenotypic
diversification, and virulence-associated traits in *Pseudomonas aeruginosa*. Replicate
populations of *P. aeruginosa* were evolved for 60 days in the presence or absence of the
ciliate predator *Tetrahymena thermophila*, followed by whole-population metagenomic
sequencing, isolate-level genome sequencing, and quantitative phenotypic assays. We
observed extensive genetic diversification across all populations, with strong signatures of
both positive and purifying selection and pervasive parallel evolution at gene and
nucleotide levels. Predator-exposed populations accumulated mutations enriched in
regulatory, metabolic, and virulence-associated pathways, revealing predictable genomic
targets of selection. However, many parallel mutations were shared between predator-
exposed and predator-free populations, indicating that adaptation to the abiotic
environment represented a dominant selective force. Genotype–phenotype analyses
revealed pleiotropic effects and trade-offs linking motility, growth, and virulence-
associated traits. Despite pronounced genomic adaptation and coordinated phenotypic
shifts, changes in virulence in an *in vivo* host model were modest and context dependent.
Taken together, our results indicate that predator exposure can influence evolutionary
trajectories in *P. aeruginosa* and highlight the value of extending such approaches across
multiple ecological and host contexts.

Significance

Many bacterial pathogens spend much of their evolutionary history outside hosts, where they face intense ecological pressures such as predation. How these pressures shape pathogen evolution and disease potential remains unclear. Using experimental evolution, genomics, and phenotypic analyses, we show that prolonged exposure to a eukaryotic predator drives predictable genetic and phenotypic changes in the opportunistic pathogen *Pseudomonas aeruginosa*. Predator exposure altered regulatory, metabolic, and virulence-associated pathways, yet much adaptation was shared with predator-free populations, highlighting the dominant role of abiotic environments. Although predator-driven evolution reshaped traits linked to motility and growth, its effects on virulence were modest and context dependent. These findings clarify how environmental interactions influence pathogen evolution and underscore the importance of studying pathogens across diverse ecological settings.

Introduction

Pseudomonas aeruginosa is one of the most widely studied opportunistic bacterial pathogens and serves as a model organism for understanding bacterial virulence, host–pathogen interactions, and social evolution in microbial communities. This free-living, Gram-negative species inhabits a remarkably broad range of environments, including soil, freshwater and marine systems, plant surfaces, animals, and humans (Silby et al. 2011; Wu and Li 2015; Crone et al. 2020; Diggle and Whiteley 2020). Its ecological versatility is largely attributed to a large and flexible genome that encodes diverse metabolic pathways, regulatory networks, and virulence factors, enabling rapid adaptation to distinct niches (Stover et al. 2000; Coggan and Wolfgang 2012; Aroca Molina et al. 2024). Clinically, *P. aeruginosa* is a major concern as a leading cause of opportunistic and nosocomial infections and a significant contributor to morbidity and mortality in patients with cystic fibrosis (Kerr and Snelling 2009).

The evolutionary and ecological success of *P. aeruginosa* is shaped by its capacity to respond rapidly to environmental pressures. Experimental evolution studies have shown

that nutrient limitation, growth in biofilms, chronic infection-like conditions, and

interactions with other microbes or phages can drive predictable genetic changes,

including loss-of-function mutations in quorum-sensing regulators such as *lasR*

(Hammond et al. 2015; Mould et al. 2020) and alterations in genes associated with

virulence (Wong et al. 2012; Scribner et al. 2022). Predator-mediated selection, in

particular, influences traits closely linked to pathogenicity, including biofilm formation,

motility, toxin production, and cooperative behaviors (Matz et al. 2008; Friman et al. 2013;

Friman and Buckling 2014). Together, these findings support the hypothesis of coincidental

evolution (Levin 1996), which proposes that virulence in opportunistic pathogens often

arises outside the host as a by-product of traits selected for environmental survival,

including defense against predators such as amoebae, protists, and bacteriophages

(Hosseinidoust et al. 2013; Leong et al. 2022; Hopkins et al. 2024).

Despite growing evidence that environmental selection can shape virulence-related traits,

few studies have examined how long-term exposure to eukaryotic predators influences

both the genomic and phenotypic evolution of *P. aeruginosa*. Coculture experiments with

amoebae, *Acanthamoeba castellanii*, have shown that prolonged predator exposure can

either enhance or attenuate virulence-associated phenotypes depending on ecological

context (Leong et al. 2022), while experimental evolution in the presence of *Caenorhabditis*

elegans demonstrated that host interactions modulate both virulence expression and the

strength of selection (Granato et al. 2018). Beyond these studies, however, the

evolutionary consequences of sustained interaction with eukaryotic predators remain

poorly understood.

Here, we address this gap using experimental evolution to investigate how exposure to a

eukaryotic predator shapes virulence and virulence-associated traits in *P. aeruginosa*.

Populations were evolved for 60 days in the presence or absence of the ciliate

Tetrahymena thermophila, followed by whole-population and isolate-level sequencing to

identify mutations that accumulated during experimental evolution. We detected strong

positive and purifying selection, as well as parallel evolution at both the gene and

nucleotide levels across independent populations. In addition, we quantified fitness and

virulence-related phenotypes in evolved isolates. Our results demonstrate that predator

presence drives predictable genomic changes in key regulatory and virulence-associated

genes and leads to coordinated, context-dependent shifts in traits such as motility, biofilm

formation, and growth. Despite substantial genomic adaptation, changes in virulence in a

host model were modest, highlighting how predator-driven evolution can shape traits

relevant to both environmental survival and pathogenic potential.

Results

Six media-evolved (ME) and six predator-evolved (PE) replicate populations of *P.*

aeruginosa strain NRB were maintained by daily passaging. For the PE lines, 4% of each

bacterial culture was transferred into fresh axenic *T. thermophila* cultures each day to

minimize predator–prey co-evolution. Successful transfer of *P. aeruginosa* was confirmed

by plating on LB agar. The viability and cell density of the *T. thermophila* cultures were

assessed before each passage using light microscopy.

All six PE and six ME populations were subjected to whole-genome shotgun (WGS)

metagenomic sequencing on Day 60. In addition, the ancestral strain and individual

isolates from each of the 12 evolved populations (ME Lines 1–6 and PE Lines 1–6) were

sequenced using WGS sequencing. We then performed phenotypic assays on the evolved

isolates (ME = 6 and PE = 6) to attempt to assign mutations to phenotypes.

Mutation patterns in PE and ME populations

Mutations in the evolved populations were identified by comparison to the ancestral

genome. The ancestral *P. aeruginosa* strain used in this study was obtained from Carolina

Biological Supply; because its precise strain identity was not specified, we sequenced the

isolate to both confirm its identity and generate reference genome. Whole-genome

comparison revealed that the ancestral strain shares all genes with *P. aeruginosa*

FDAARGOS_1041 (GenBank CP065966.1) and is 99.996% identical at the nucleotide level,

corresponding to approximately 274 nucleotide differences. To generate a reliable

reference, the ancestral strain was sequenced using both long-read (Nanopore) and short-

read (Illumina) technologies, and the reads were assembled with hybridSPAdes v3.15.3

(Antipov et al. 2016) to produce a consensus genome. To reduce false-positive variant calls

arising from sequencing or assembly errors, all ancestral short reads (average of 215X

coverage) were mapped back to the consensus genome using breseq v0.39.0 (Deatherage

and Barrick 2014). Any variants detected in the ancestral strain during this mapping were

considered potential sequencing or assembly artifacts and were excluded from the

subsequent analyses of mutations in the evolved populations.

A total of 972 distinct mutations (frequency >5%) were identified across all evolved lines,

excluding same-site parallel mutations. The number of mutations per population ranged

from 10 to 169, with average mutation frequencies between 10% and 79% (Figure 1A;

Dataset S1). The total number of mutations within a population was not correlated with

average mutation frequency (Figure 1A: R^2=0.3209, P=0.0548) or average genome coverage

(Figure 1B, Table S1: R^2=0.0006, P=0.9415). Together, these nonsignificant correlations

suggest that our sequencing depth was sufficient to capture the mutations present in the

populations.

Figure 1. Mutation profiles in evolved populations. A) Relationship between average mutation
frequency and the total number of mutations per evolved population. B) Relationship between
average genome coverage and the total number of mutations per evolved population. For panels A
and B, R^2 values and P-values were calculated using simple linear regression. C) Expected versus
observed proportions of mutation types (nonsynonymous, synonymous, intergenic, and other
mutation types) in each population. D) Functional annotation of coding-region mutations assigned

to KEGG Orthology (KEGG) functions in media-evolved (ME) and predator-evolved (PE) lines; noncoding and unassigned mutations are not shown. See Dataset S1 for a complete list of identified mutations.

In every evolved population, mutations were categorized as nonsynonymous,

synonymous, RNA, or intergenic (Figure 1C; Dataset S1). To estimate the expected

proportions of nonsynonymous, synonymous, intergenic, nonsense, and RNA mutations

under random mutation, we conducted 10,000 independent simulations on the ancestral

genome using the Jukes–Cantor model (Jukes and Cantor 1969). The proportion of

nonsynonymous mutations in the evolved lines ranged from 44.9% to 76.5%, compared

with an expected 64.7% under random mutation (Figure 1C). Five of the six ME lines (all

except ME2) and two PE lines (PE4 and PE5) contained fewer nonsynonymous mutations

than expected, suggesting purifying selection. By contrast, one ME line (ME2) and two PE

lines (PE1 and PE2) showed an excess of nonsynonymous mutations, consistent with

positive selection. Two PE lines (PE3 and PE5) had proportions close to the random

expectation (Figure 1C).

Across the PE lines, 561 total mutations were identified, including those also present in ME

lines (i.e., shared mutations), of which 404 occurred in coding regions and 247 could be

assigned a KEGG Orthology (KO) annotation (Figure 1D). In the ME lines, 457 mutations

were identified, including shared mutations, with 311 in coding regions and 194 assigned

KO annotations (Figure 1D). Overall, the PE lines had a higher proportion of mutations in

genes associated with Signaling and Cellular Processes, Genetic Information Processing,

Metabolism of Cofactors and Vitamins, Nucleotide Metabolism, Human Diseases, and

Organismal Systems than the ME lines.

We additionally detected strong signatures of positive selection in our populations,

reflected by numerous cases of parallel evolution. Parallel evolution events were defined

as mutations that arose at the same site or within the same gene in independently evolved

lineages (ME, PE, or both). Notably, none of these events were shared across all

populations, indicating that the variants were not present in the ancestral genotype. In the

ME populations, we identified 34 site-specific parallel mutations occurring in 7 genes,

along with 12 gene-level parallel mutations distributed across 6 genes (Figure 2A–B). In the

PE populations, we detected 10 site-specific parallel mutations in 4 genes and 35 gene-

level parallel mutations in 17 genes (Figure 2A and 2C). Additionally, ME and PE

populations shared 113 site-specific parallel mutations spanning 10 genes and 287 gene-

level parallel mutations across 40 genes (Figure 2A), 178 of which occurred in the same

gene *blp2* (Dataset S1). Most parallel mutations were present at relatively low frequencies

(<20%); however, several reached high frequencies (>60%) or were fixed, including

mutations in *mgtE*, *phoQ*, and *clpA* in ME lines and in *siaB*, *wbpM*, and *fha1* in PE lines

(Figure 2B–C, Dataset S1). Parallel mutations are extremely unlikely to occur by chance

and constitute strong evidence of positive selection. In addition, the vast majority of the

high frequency (>60%) parallel mutations affected nonsynonymous positions (92%), which

further suggests that these mutations are adaptive.

Figure 2. Parallel evolution events in evolved populations. A) Number of fixed, same-site and same-gene parallel mutations in predator-evolved (PE) and media-evolved (ME) populations, as well as mutations shared between both groups. B–C) Frequency of site-specific and gene-specific parallel mutations in (B) ME and (C) PE populations; genes in which site-specific parallel mutations occurred are indicated in bold font and gene-specific parallel mutations are highlighted in gray. For a list of all mutations identified in the evolved populations see Dataset S1.

161 *Genomic and phenotypic characterization of evolved isolates*

162   A single isolate was collected from each population (*n* = 12). Analysis of each isolate

163   revealed between three and seven mutations per genome, and each isolate carried at least

164   one mutation that was also fixed in its corresponding population (Figure 2A–B and Figure

165   3A). In total, 46 mutations were identified across 24 genes, and several instances of

166   parallel evolution were detected (Figure 3A, Dataset S2).

Figure 3. Genotypic and phenotypic changes in evolved isolates. A) Mutations identified in evolved isolates, highlighting site-specific parallel mutations (bold font) and gene-specific parallel mutations (outlined boxes) unique to predator-evolved (PE) and media-evolved (ME) isolates. NS, nonsynonymous; S, synonymous; I, intergenic; STOP, nonsense; SNV, single-nucleotide variant; Δ, deletion; +, insertion. For a list of all mutations identified in the evolved isolates see Dataset S2. B) Log$_2$ fold change calculated from the mean of all replicates and trials for each phenotype measured in evolved isolates relative to the ancestral strain. Yellow indicates an increase, blue indicates a decrease, and white indicates no change. Asterisks within boxes indicate significant differences when individual isolates were compared to the ancestor, whereas asterisks in the margin indicate significant differences when the mean of all ME lines or all PE lines was compared to the ancestral mean. Bold text denotes phenotypes showing significant changes relative to the ancestor. Statistical significance was determined using a Kruskal–Wallis test with Dunn's multiple-comparison test (P < 0.05). Full data for all replicates, including non-log$_2$ fold-change values, are shown in Figures S1 and S2.

Among the ME isolates, site-specific parallel mutations occurred in *clpA* and *phoQ*, both

present in ME4 and ME6. A site-specific parallel mutation in *mgtE* was also found in ME2,

ME4, and ME6. In addition, a gene-specific mutation in *fleQ*, unique to the ME lines,

occurred in ME1, ME2, and ME5 (Figure 3A, Dataset S2). In the PE isolates, a gene-specific

mutation in *wbpM*, unique to the PE lines, was identified in PE1, PE2, PE5, and PE6. Finally,

two gene-specific mutations, *mgtE* and *phoQ*, were shared between ME and PE isolates

(Figure 3A, Dataset S2).

To assess the impact of predator-driven evolution on *P. aeruginosa*, we performed

phenotypic assays on a single isolate from each evolved population, allowing direct

correlation with mutations identified via genomic analysis. Virulence-associated traits,

including biofilm production, motility, siderophore production, casein hydrolysis, and

hemolytic activity, were measured (Figure 3B, Figure S1 A–F). Fitness was evaluated in two

different conditions: growth after 24 h in media and in coculture with *T. thermophila* (Figure

3B, Figure S1G–H). All assays were performed in at least triplicate.

After correcting for multiple comparisons, only a few isolates differed significantly from the

ancestor for any of the phenotypes tested. Specifically, isolates ME5, PE4, and PE6

exhibited reduced casein hydrolysis, indicating decreased extracellular protease activity

involved in protein degradation (Figure 3B), whereas isolate PE2 showed significantly

increased hemolytic activity compared to the ancestral strain (Figure 3B). When comparing

the mean of all PE or ME isolates to the ancestor, PE isolates showed higher biofilm

production and ME isolates exhibited increased growth in media (Figure 3B; Figure S1G).

Although not statistically significant, additional patterns were observed. Swimming activity

(Figure 3B) decreased in most D60-evolved lines, with the exception of ME3 and ME4.

Swarming activity (Figure 3B) was generally reduced in ME isolates, except ME1, but

increased in most PE isolates relative to the ancestor. For biochemical traits, most evolved

isolates produced slightly more siderophore (Figure 3B), with PE2 as the exception.

Protease activity decreased on average in both ME and PE lines, except for a slight increase

in D60-ME6. Hemolytic activity decreased in all ME lines, whereas PE lines exhibited an

increase (Figure 3B).

To evaluate how predation impacted *P. aeruginosa* virulence, we used a tractable

invertebrate host model, the honey bee (*Apis mellifera*) (Figure 3B, Figure S2). We

performed two types of assays: one in conventional (CV) honey bees, which were healthy

adults with an intact microbiome (Figure S2A), and another in microbiota-depleted (MD)

bees (Figure S2B). The MD bees were newly emerged (~1 day old) adults of the same age.

All PE and ME isolates displayed a decrease in virulence in CV bees, although this was not

significant after multiple comparisons. In MD bees, most ME isolates showed a slight

decrease in virulence compared to the ancestor, with the exception of ME1. Among the PE

isolates, four (PE2, PE4, PE5, and PE6) showed a slight increase, PE1 showed no change,

and PE3 showed a slight decrease. Overall, in CV bees, the average virulence of all PE

isolates did not differ from the ancestor, whereas the average virulence of all ME isolates

was significantly reduced (Figure S2A). In MD bees, no significant differences in virulence

were observed for either group compared to the ancestor (Figure S2B).

To determine whether the attenuated virulence was due to reduced proliferation in the

honey bee gut, we performed growth assays for each isolate in MD bees (Figure 3B, Figure

S2). While the mean growth levels showed trends that loosely resembled the virulence

patterns observed in MD bees (Figure S2A), no statistically significant differences in growth

were detected for either ME or PE isolates. Instead, both groups exhibited substantial

isolate-to-isolate variation. In each group, three of the six replicates showed a slight

increase in growth (ME2, ME3, ME6; PE2, PE4, PE5), whereas the remaining three showed a

slight decrease (ME1, ME4, ME5; PE1, PE3, PE6) relative to the ancestor.

We next analyzed genotype–phenotype correlations and found that a subset of mutations

was strongly associated with changes in multiple functional traits (Figure 4A, Dataset S3).

For example, mutation 28 in *mgtE* was negatively correlated with biofilm production while

mutation 22, also in *mgtE*, was negatively correlated with virulence in CV bees. Mutations

3, 18, 19, and 26 (located in *wbpM*, *morA*, a phage tail protein, and *mgtE*) displayed

identical patterns across phenotypes, showing significant positive correlations with

swarming and significant negative correlations with growth in media and virulence in MD

honey bees.

Figure 4. Correlations between genotypes and phenotypes in evolved isolates. A) Correlations
between individual mutations and phenotypic traits. B) Correlations among identified mutations.
C) Correlations among phenotypic traits. Yellow represents a positive correlation, blue represents

a negative correlation, white means no correlation, and asterisks (*) indicate statistical significance, $P < 0.05$. Statistical significance was assessed using Spearman's rank correlation. See Dataset S3 for complete list of Spearman r and P-values.

Genotype–genotype correlations (Figure 4B, Dataset S3) revealed that several mutations

were positively associated, suggesting preferred evolutionary trajectories. The phenotype–

phenotype correlation matrix (Figure 4C, Dataset S3) showed that certain traits were

significantly positively coupled, including swarming motility and hemolytic activity,

siderophore production and casein hydrolysis, and growth in media and virulence in MD

bees.

In addition to individual fitness assays, we performed competition experiments by

coculturing all isolates for 20 h in media, in the presence of *T. thermophila*, and in honey

bees (Figure 5). Whole-genome metagenomic sequencing was then used to determine

which strains outcompeted the others in each condition. In the presence of the predator,

ME3 and ME4 outperformed all other lines. In media, ME2, ME3, ME4, and ME6

outcompeted the other isolates and the ancestor. In honey bee competitions, ME4

prevailed over all other isolates and the ancestral strain (Figure 5).

Figure 5. Competition assays with the evolved isolates and the ancestral strain in the presence of a predator, in media, or in honey bee host. Each assay was performed in triplicate.

238 Discussion

The emergence of virulence and other adaptations in opportunistic bacterial pathogens,

particularly *P. aeruginosa*, has been widely studied under diverse selective conditions,

including copper exposure (12 days), antibiotic treatment (30 days), oxidative stress (12

days), and in presence of amoeboid and nematode predators (*A. castellanii*, 42 days and

nematode *C. elegans*, 60 days) (Granato et al. 2018; Laborda et al. 2022; Lear et al. 2022;

Leong et al. 2022; Fu et al. 2023). While predator pressure is predicted to indirectly select

for increased virulence, the two studies that have evaluated this in *P. aeruginosa* generally

found reduced virulence (Granato et al. 2018; Leong et al. 2022). However, the impact of a

ciliate predator on *P. aeruginosa* evolution has not been systematically explored. Here, we

investigated the influence of the generalist ciliate predator *T. thermophila* on the virulence,

fitness, and genomic evolution of *P. aeruginosa* over a 60-day experimental evolution

period. By combining WGS metagenomics of evolved populations and isolates with *in vitro*

and *in vivo* phenotypic assays, we provide insight into how predation shapes bacterial

evolution at both the genotypic and phenotypic levels.

Our WGS analyses revealed extensive genetic diversification across the ME and PE

populations. A total of 457 mutations were identified in ME lines and 561 in PE lines, with

numerous cases of parallel evolution at both the site and gene level. The observation of

parallel mutations at high frequency in genes such as *mgtE*, *phoQ*, and *clpA* in ME lines

and in *siaB*, *wbpM*, and *fha1* in PE lines indicates strong selective pressures acting on

specific functional pathways. Interestingly, while several lines showed evidence of

purifying selection, other lines exhibited an excess of nonsynonymous mutations,

consistent with positive selection. These patterns highlight that adaptation can proceed

via both constraint and innovation and that even replicate populations evolving under the

same conditions can follow distinct evolutionary trajectories.

KEGG pathway analysis indicated that PE populations were enriched for mutations in

genes associated with signaling, metabolism, and genetic information processing. This

enrichment suggests that exposure to predation selects for modifications in critical

regulatory and metabolic functions that may enhance survival under predatory pressure.

However, adaptation to the media appears to be the dominant selective force, as reflected

by the large number of parallel mutations shared between ME and PE lines and the

substantial number of mutations unique to ME lines. This may indicate that this strain of *P.*

aeruginosa already possesses effective mechanisms to defend against or avoid predation,

reducing the relative impact of predator-driven selection compared with nutrient-based

adaptation.

Analysis of individual isolates (n = 12) revealed 46 mutations across 24 genes, with

evidence of selective sweeps at identical positions or within the same genes across lines.

Notably, mutations in *mgtE*, *phoQ*, *fleQ*, and *wbpM* were recurrent across multiple lines,

suggesting that these genes are key targets of selection. *FleQ*, a master regulator of

flagellar motility and biofilm formation, was mutated in ME1, ME2, and ME5, coinciding

with reductions in both swimming motility and biofilm production. This is consistent with

previous studies showing that *FleQ* modulates motility and exopolysaccharide

biosynthesis in a context-dependent manner (Arora et al. 1997; Hickman and Harwood

2008). Mutations in *phoQ*, part of the PhoP–PhoQ two-component regulatory system

(Gellatly et al. 2012), were found in several ME and PE lines and likely contributed to

attenuated virulence, particularly in conventional honey bees. Conversely, mutations in

mgtE, a dual-function magnesium transporter linked to type III secretion system regulation

(Anderson et al. 2010; Coffey et al. 2014), were present in all isolates. While *mgtE*

mutations can enhance virulence, in our study most lines exhibited reduced virulence,

except for ME1, PE4, and PE6, suggesting that the phenotypic outcome depends on the

precise mutation location and genetic context. In ME1, a stop codon in *mgtE*, combined

with a mutation in *ladS*, a kinase regulating the acute-to-chronic virulence (Broder et al.

2016), likely contributed to its elevated virulence. Mutations in *wbpM* were observed in

most PE isolates but none of the ME isolates, highlighting a potential role in adaptation to

predation. *WbpM* is involved in O-antigen synthesis (Burrows et al. 2000), and while only

PE6 displayed increased virulence relative to the ancestor, other mutations in *wbpM* may

confer advantages against predation, potentially via enhanced biofilm formation or altered

surface interactions.

Our genotype–phenotype correlation analysis indicates that phenotypic diversification in

this system is driven by a limited number of mutations with broad, pleiotropic effects

rather than by many mutations each affecting a single trait. In particular, distinct mutations

within mgtE were associated with different functional outcomes, with one mutation

negatively correlated with biofilm formation and another specifically linked to reduced virulence in CV bees, highlighting functional heterogeneity within this locus. In contrast, mutations in wbpM, morA, a phage tail protein, and mgtE exhibited highly concordant phenotypic signatures, characterized by increased swarming motility coupled with reduced growth in rich media and decreased virulence in MD honey bees. This repeated association of motility gains with fitness trade-offs suggests that these mutations may shift resource allocation toward dispersal at the expense of host-associated performance. The presence of strong positive genotype–genotype correlations further implies that these mutations do not arise independently but instead form preferred evolutionary combinations, potentially reflecting shared selective pressures or epistatic compatibility. Finally, the observed coupling among phenotypic traits, such as swarming with hemolytic activity and growth in media with virulence, supports the idea that ecological and pathogenic traits are evolutionarily linked, constraining the range of accessible adaptive strategies. Collectively, these patterns suggest that pleiotropic mutations and correlated trait responses interact to bias evolutionary trajectories toward a restricted subset of genetic and phenotypic outcomes.

Using CV and MD honey bees as in vivo models, we observed context-dependent virulence outcomes. While all evolved isolates showed decreased virulence in CV bees, ME isolates were significantly attenuated on average. In MD bees, phenotypic variation was greater, with some PE isolates showing slight increases in virulence. Growth assays in MD bees revealed only nonsignificant trends in gut proliferation, with no consistent or significant differences among isolates, indicating that reduced virulence cannot be attributed to an individual isolate's ability to proliferate in the honey bee gut. Instead, these results suggest that changes in specific virulence factors, rather than overall bacterial fitness, drive host outcomes. It is important to note that *P. aeruginosa* is not a natural pathogen of honey bees, and the virulence effects observed in this host may not fully reflect outcomes in mammalian systems. Indeed, our evolved lines could exhibit differential virulence in mammalian hosts, as some phenotypic changes, such as increased hemolytic activity, would have minimal impact on bees, which lack red blood cells, but could enhance

pathogenicity in vertebrate hosts. However, testing these potential effects in mammals

was beyond the scope of the current study.

Competition assays demonstrated that predator-driven evolution shaped competitive

outcomes in environment-specific ways by decreasing its fitness, as ME isolates

dominated all three competition environments. These findings suggest that adaptation to

predation impair fitness under selective pressure but may incur trade-offs under

alternative conditions, consistent with ecological theory on specialization versus

generalist strategies.

Overall, our results illustrate that long-term exposure to a eukaryotic predator can drive

predictable patterns of parallel evolution in *P. aeruginosa*, with distinct genotypic and

phenotypic consequences. However, the large number of parallel mutations shared

between ME and PE lines suggests that adaptation to the media exerted a stronger

selective force than predator-specific pressures. Comparing these findings to our previous

60-day experimental evolution study of *Serratia marcescens* under *T. thermophila*

predation (Hopkins et al. 2024) highlights both shared and divergent patterns. In *S.*

marcescens, predator-evolved lines maintained or enhanced virulence and grazing

resistance, exhibited increased biofilm production, and accumulated many more

mutations specific to predator-evolved lines, indicating stronger adaptation to predation

than in *P. aeruginosa*. By contrast, *P. aeruginosa* PE populations showed more variable

phenotypic responses, with virulence generally decreasing but biofilm production

increasing. These differences in virulence may also reflect host specificity, as *S.*

marcescens is a natural pathogen of honey bees, whereas *P. aeruginosa* is not known to

naturally infect this host. Together, these results suggest that the relative impact of

predator-specific versus general environmental selection can differ between opportunistic

bacteria, reflecting species-specific differences in ecological strategy, phenotypic

plasticity, and the balance between nutrient adaptation and predation.

Although our 60-day evolution period revealed substantial genetic and phenotypic

changes, longer-term experiments may uncover additional adaptive mutations that require

more generations to become fixed. Future studies could explore how these evolutionary

dynamics manifest in complex microbial communities, across different host species, or

under fluctuating environmental conditions. The functional analyses of key genes, such as

mgtE, *phoQ*, and *wbpM*, will also help elucidate the molecular mechanisms linking

predator-driven evolution to virulence and ecological fitness.

Methods

Experimental evolution

The ancestral *P. aeruginosa* strain we used for our experiments was purchased from

Carolina Biological Supply Company and evolved either in media alone (ME) or in the

presence of the protist predator *T. thermophila* strain SB210 (PE). *T. thermophila* was

obtained from the Tetrahymena Stock Center at Cornell University. Each evolution

treatment included six replicate lines.

Prior to starting the evolution experiment, *T. thermophila* and ancestral *P. aeruginosa* were

cultured individually for 72 h and 48 h, respectively, at 35°C in Neff media. To initiate each

evolved line, 1 OD of *P. aeruginosa* was resuspended in fresh Neff media, and 0.12 mL was

added to either 3 mL of Neff media alone (ME lines) or 3 mL of Neff media containing *T.*

thermophila (PE lines). The starting ratio of bacteria to predator was determined in

preliminary trials to allow both organisms to persist in co-culture for 24 h.

Experimental cultures (3 mL) were incubated at 35°C with shaking. Every 24 h for 180 days,

cultures were vortexed and 4% of each line was transferred into fresh Neff media (ME lines)

or new axenic *T. thermophila* cultures (PE lines). Prior to each transfer, *T. thermophila*

viability and density were assessed using light microscopy, and *P. aeruginosa* growth was

confirmed by plating on LB agar. Throughout the experiment, both organisms were

consistently present, and no noticeable changes in population size were observed based

on daily growth and plating. On day 60, agar plates were inoculated from each culture, and

one colony was randomly selected from each ME (ME1–6) and PE population (PE1–6). Each

colony was grown in 3 mL LB broth for 48 h at 35°C to obtain pure cultures. A 200 µL aliquot

of each culture was frozen at –80°C in 20% glycerol. These 12 isolates were subsequently

sequenced and used for phenotypic assays, including virulence in bees, growth assays,

biochemical tests, biofilm production, and competition experiments.

DNA extraction and sequencing

Cultures of the evolved populations (ME1–6 and PE1–6), evolved isolates (ME1–6 and PE1–

6), and the ancestral strain were diluted to 1 OD, and DNA was extracted using the Zymo

Quick-DNA™ Miniprep Plus Kit (D4069). Libraries for short-read sequencing were prepared

using the Illumina Nextera DNA Flex Library Prep Kit (20018704). The ancestral genome

was sequenced using both long-read (Oxford Nanopore MinION) and short-read (Illumina

iSeq100) technologies. Evolved populations and isolates were sequenced using short-read

Illumina platforms (iSeq100 or MiSeq100) with 2 × 150 bp paired-end reads.

Genome assembly and analysis

The Illumina and Nanopore reads of the ancestral genome were combined to generate a

hybrid assembly using hybridSPAdes v3.15.3 (Antipov et al. 2016). Assembled scaffolds

were annotated with Bakta (Schwengers et al. 2021) on the Department of Energy Systems

Biology Knowledgebase (KBase) platform. Genome sequence comparisons were

performed between the ancestral strain and the three closest related strains, identified

based on BLAST searches against the NCBI database. Average nucleotide identity values

were calculated using FastANI v1.32 (Jain et al. 2018). A pangenome analysis was

conducted using Panaroo v1.5.2 (Tonkin-Hill et al. 2020) with the following parameters: an

initial clustering threshold of 0.98 using CD-HIT, a family threshold of 0.7 for secondary

clustering of orthologous genes into gene families, a length difference percentage cutoff of

0.80, a core gene threshold of 1 (genes present in all genomes), strict clean mode, and

removal of invalid genes. All short reads of the ancestral strain were then mapped back to

the consensus genome using breseq v0.39.0 (Deatherage and Barrick 2014). Any variants

identified during this mapping were considered sequencing or assembly artifacts and were

excluded from the set of candidate mutations in the evolved populations and isolates.

Sequencing reads from the evolved populations and isolates were trimmed using

Trimmomatic v0.40 (Bolger et al. 2014) and subsequently mapped to the assembled

ancestral genome using breseq v0.39.0 (Deatherage and Barrick 2014), with a base-quality

cutoff of PHRED 30 and a minimum frequency cutoff of 0.05 to identify mutations. Once

identified, mutations were compared against the ancestral strain using gdtools

(Deatherage and Barrick 2014).

To assess the expected distribution of mutations under neutrality, we performed 10,000

independent simulations using a custom Python script. Each simulation introduced a

single mutation at random under a Jukes–Cantor model in our *P. aeruginosa* reference

genome. Genome annotations were used to classify mutations as intergenic or coding. The

coding mutations were further classified as synonymous or nonsynonymous by comparing

the translated protein sequence before and after the mutation. The functional

characterization of genes was predicted using KOALA KEGG Orthology and Links

Annotation (Kanehisa 2000; Kanehisa et al. 2016).

Phenotypic assays

Growth in media: Cultures of evolved populations (ME1–6 and PE1–6), evolved isolates

(ME1–6 and PE1–6), and the ancestral strain were diluted to 1 OD. For each isolate, 0.12

mL of culture was added to 3 mL fresh Neff media in triplicate and incubated at 35°C with

shaking at 110 rpm. After 24 h, cultures were diluted and plated on cetrimide agar in

duplicate. Plates were incubated for 48 h at 35°C, and colonies were counted to calculate

CFUs.

Growth in the presence of *T. thermophila*: Axenic 48-h cultures of *T. thermophila* in Neff

media were inoculated with 120 µL of 1 OD cultures of each isolate in triplicate. Cultures

were incubated for 20 h at 35°C and 110 rpm, diluted, and plated on cetrimide agar in

duplicate. Plates were incubated for 48 h at 35°C before counting colonies and calculating

CFUs.

Growth in-host: Pupae were collected 2–3 days before emergence and kept in cages with

sterile sugar syrup (SSS) and sterile pollen. Newly emerged adult bees were hand-fed 10 µL

of either sterile SSS, the ancestor in SSS, or an evolved isolate in SSS (5 bees per cup, 15

bees per treatment). Bees were maintained at 35°C and 95% humidity. After 20 h, guts

were collected, stored in 20% glycerol at –80°C, and dilutions up to 10^{-6} were performed

and plated on cetrimide agar in triplicate. Plates were incubated 48 h at 35°C, colonies

were counted, and CFUs/mL were calculated based on the dilution factor.

Biofilm assay: Biofilm formation was assessed in triplicate. First, 5 µL of 1 OD cultures

were added to 150 µL Neff media in 96-well microplates with an additional sterile

nonskirted plate immersed into the wells to provide a bigger surface attachment; this

setup was incubated at 35°C for 24 h without shaking. Wells and the immersed plate were

washed twice with sterile ddH$_2$O, stained with 150 µL of 0.1% crystal violet for 15 min,

washed again, and air-dried. Biofilms in both microplates (initial and immersed) were

separately solubilized with 150 µL of 30% acetic acid, and absorbance was measured at

490 nm. The measurements were combined to obtain a final reading. Three independent

assays were performed, each with four replicates.

Casein hydrolysis: Milk agar plates were prepared by mixing 2.5 g yeast extract and 7.5 g

agar in 350 mL water and 15 g milk powder in 150 mL water; both solutions were

autoclaved, cooled, mixed, and poured into plates. Sterile discs were inoculated with 4 µL

of 1 OD bacterial culture and incubated at 37°C for 48 h. To assess protease activity, the

plates were photographed and analyzed using ImageJ by calculating the diameter of the

activity and subtracting the 4-mm diameter of the disc.

Siderophore production: CAS-CTAB agar plates were prepared by dissolving 0.065 g

Chromeazurol S (CAS) in 50 mL water and 0.0729 g CTAB in 40 mL water. Those two

solutions were then mixed with 10 mL of 1 mM FeCl$_3$ hexahydrate and 10 mM HCl solution

and added to LB agar medium, followed by pH adjustment to 6.8 and autoclaving. Sterile

discs were placed and inoculated with 4 µL of 1 OD bacterial culture and incubated at

37°C for 48 h. For analyzing siderophore production, plate photographs were uploaded to

ImageJ to measure the diameter of the siderophore production minus the diameter of the

disc (4 mm).

Hemolytic activity: Blood agar plates (LB + 5% sheep blood) were inoculated with 4 µL of 1

OD bacteria and incubated at 37°C for 48 h before assessing hemolysis, as previously

described in the siderophore production protocol.

Swimming and swarming motility: Swim plates contained peptone (5 g), yeast extract (3

g), and 0.3% agar per liter of water; swarm plates used 0.5% agar. Plates were autoclaved,

poured, and allowed to solidify for up to 24 h. The plates were stab-inoculated with 2 µL of

1 OD culture and incubated at 37°C for 48 h before measuring motility by measuring the

diameter of the motion in photographs uploaded to ImageJ.

Virulence assay in honey bees

Conventional bees: Pure 1 OD cultures of each evolved isolate and the ancestor were

centrifuged (10,000 rpm × 2 min) and individually resuspended 1:1 in SSS. Adult worker

bees (*A. mellifera*) were collected from a single hive at the Phytotron, NCSU, Raleigh, NC,

and immobilized at 4°C. Bees were randomly distributed into groups and exposed via the

immersion method (~10 µL per bee) to one of three treatments: 1) sterile SSS only, 2) the

ancestor in SSS, or 3) each evolved isolate in SSS. Bees were maintained in cup cages (20

bees/cup; 100 bees/treatment) under hive-like conditions (35°C, 95% humidity). Mortality

was recorded daily for 5 days. Two independent assays were performed, each with five

replicates, totaling 200 bees per isolate.

Microbiota-depleted bees: Each evolved isolate was first cultured to 1 OD_{600}, followed by

centrifugation to pellet the bacteria and resuspension of the pellet in 1:1 SSS. Newly

emerged honey bees (*A. millifera*) were immobilized at 4°C and randomly distributed in

different 50-mL Falcon tubes. Ten bees were added to a single tube, in which isolate-

containing SSS was added (10 µL per bee) for immersion treatment. Every isolate,

ancestor, and control treatment was performed in triplicate (i.e., 10 bees × 3 = 30 bees per

treatment). The treated bees were then maintained in cup cages under hive-like conditions

(35°C and 95% humidity) for 5 days, and mortality was recorded. Two independent assays

were performed, totaling 60 bees per isolate.

Competition assays

In-host competition: Pupae were collected 2–3 days prior to emergence from a single hive

at the Phytotron (NCSU) and maintained in cages with SSS and sterile pollen. After

emergence, cultures of each evolved isolate and the ancestor were normalized to OD_{600} =

1, centrifuged (10,000 rpm for 2 min), and resuspended in SSS. For inoculation, strains

were pooled such that the ancestor and each evolved isolate (ME1–6 and PE1–6) were

present in equal proportions. A total of 25 bees were hand-fed 10 μL of the bacterial

suspension in SSS and maintained in cup cages (5 bees per cup; 5 cups total) at 35°C and

95% humidity. After 20 h, bee guts were collected under sterile conditions and stored in

20% glycerol at –80°C. Although all bees were sampled, three individual bee gut samples

(each from a different cup cage) were selected for downstream analysis. These samples

were plated on cetrimide agar in triplicate, incubated for 24 h, and ~300 colonies per

sample were pooled in 1× PBS. DNA was extracted from the pooled colonies and analyzed

by WGS metagenomic sequencing (2x150 PE reads on Illumina miSeq100). Reads were

mapped to the ancestral genome using breseq v0.39.0 (Deatherage and Barrick 2014), and

gdtools (part of the breseq pipeline) was used to compare each sample to mutations

present in the ancestor and the individual evolved isolates (ME1–6 and PE1–6). The relative

frequency of each strain was estimated as the average frequency of mutations shared

between the isolate and the sample.

Competition in the presence of *T. thermophila*: Five replicate tubes containing 3 mL of *T.*

thermophila in Neff media were inoculated with 0.12 mL of 1 OD mixture of all evolved

isolates and the ancestor. Cultures were incubated for 20 h at 35°C with shaking at 110

rpm. Three replicate samples were diluted and plated in triplicate on cetrimide agar,

incubated for 48 h at 35°C, and ~300 colonies per sample were collected in 1 mL of 1×

PBS. DNA was extracted from the pooled colonies, sequenced, and analyzed following the

protocol stated above for **in-host competition**.

Competition in media: Five replicate tubes containing 3 mL of Neff media were inoculated

with 0.12 mL of 1 OD mixture of the ancestor and evolved isolates. After 20 h of incubation

at 35°C and 110 rpm, three of the replicate cultures were centrifuged at 10,000 rpm for 2

min. Supernatants were discarded, and pellets were resuspended in 200 μL of 1× PBS.

DNA was extracted from the pooled colonies (~300 colonies per sample), sequenced, and

analyzed following the protocol described above for **in-host competition**.

Statistical analyses

All figures and statistical analyses were performed in GraphPad Prism v10.5.0. Figures

were visually improved using Adobe Illustrator.

Mutations: The relationship between the total number of mutations per evolved

population and (i) average genome coverage and (ii) average number of mutations was

assessed using simple linear regression (see Figure 1A–B).

Phenotypic analyses: For all phenotypic assays, significance between each evolved line

and the ancestral strain was assessed using a Kruskal–Wallis test, followed by Dunn's

multiple-comparison test, incorporating both technical and biological replicates (see

Figure 3B, Figure S1 and Figure S2). In addition, the mean phenotypic values of all ME lines

combined and all PE lines combined were each compared to the ancestral mean using a

Kruskal–Wallis test with Dunn's multiple-comparison test to assess differences between

evolved groups relative to the ancestor (see Figure 3B, Figure S1, and Figure S2).

Genotype and phenotype correlations: **Genotype–phenotype correlations:** All

mutations detected in evolved isolates were scored as binary variables (1 = present, 0 =

absent) for each evolved line. For phenotypic traits, means were calculated across all

biological and technical replicates, and $\log_2$ fold changes relative to the ancestral

phenotype were determined. Binary mutation presence/absence data and phenotypic $\log_2$

fold-change values were then used to identify significant correlations across evolved lines

between (i) mutations and phenotypes (genotype–phenotype), (ii) mutations and other

mutations (genotype–genotype), and (iii) phenotypes and other phenotypes (phenotype–

phenotype) using Spearman's rank correlation coefficient (ρ). All Spearman's ρ values and

corresponding P-values are provided in Dataset S2.

Data availability

Acknowledgements

556 We thank Marcos Garcia for his assistance with the experiments. We are grateful to Dr.

557 Awa Diop for assistance in classifying the ancestral strain used in this study. We also thank

558 Dr. Louis-Marie Bobay, Dr. Parul Bardeskar, and Patrick Gallagher for their careful reading

559 of the manuscript and helpful editorial suggestions. This work was supported by the

560 National Institutes of Health under grant 7R01GM145747 (to K.R.).

References

561 Anderson GG, Yahr TL, Lovewell RR, O'Toole GA. 2010. The Pseudomonas aeruginosa magnesium transporter MgtE inhibits transcription of the type III secretion system. *Infect. Immun.* 78:1239–1249.

562 Antipov D, Korobeynikov A, McLean JS, Pevzner PA. 2016. HYBRID SPADES: an algorithm for hybrid assembly of short and long reads. *Bioinformatics* 32:1009–1015.

Aroca Molina KJ, Gutiérrez SJ, Benítez-Campo N, Correa A. 2024. Genomic differences associated with resistance and virulence in Pseudomonas aeruginosa isolates from clinical and environmental sites. *Microorganisms* 12:1116.

Arora SK, Ritchings BW, Almira EC, Lory S, Ramphal R. 1997. A transcriptional activator, FleQ, regulates mucin adhesion and flagellar gene expression in Pseudomonas aeruginosa in a cascade manner. *J. Bacteriol.* 179:5574–5581.

Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 30:2114–2120.

Broder UN, Jaeger T, Jenal U. 2016. LadS is a calcium-responsive kinase that induces acute-to-chronic virulence switch in Pseudomonas aeruginosa. *Nat. Microbiol.* 2:16184.

Burrows LL, Urbanic RV, Lam JS. 2000. Functional conservation of the polysaccharide biosynthetic protein WbpM and its homologues in Pseudomonas aeruginosa and other medically significant bacteria. *Infect. Immun.* 68:931–936.

Coffey BM, Akhand SS, Anderson GG. 2014. MgtE is a dual-function protein in Pseudomonas aeruginosa. *Microbiology* 160:1200–1213.

Coggan KA, Wolfgang MC. 2012. Global regulatory pathways and cross-talk control Pseudomonas aeruginosa environmental lifestyle and Virulence phenotype. *Curr. Issues Mol. Biol.* 14:47–70.

Crone S, Vives-Flórez M, Kvich L, Saunders AM, Malone M, Nicolaisen MH, Martínez-García E, Rojas-Acosta C, Catalina Gomez-Puerto M, Calum H, et al. 2020. The environmental occurrence of Pseudomonas aeruginosa. *APMIS* 128:220–231.

Deatherage DE, Barrick JE. 2014. Identification of mutations in laboratory-evolved microbes from next-generation sequencing data using breseq. *Methods Mol. Biol. Clifton NJ* 1151:165–188.

Diggle SP, Whiteley M. 2020. Microbe profile: Pseudomonas aeruginosa: opportunistic pathogen and lab rat. *Microbiology* 166:30–33.

Friman V-P, Buckling A. 2014. Phages can constrain protist predation-driven attenuation of
Pseudomonas aeruginosa virulence in multienemy communities. *ISME J.* 8:1820–
1830.

Friman V-P, Diggle SP, Buckling A. 2013. Protist predation can favour cooperation within
bacterial species. *Biol. Lett.* 9:20130548.

Fu T, Gifford DR, Knight CG, Brockhurst MA. 2023. Eco-evolutionary dynamics of
experimental Pseudomonas aeruginosa populations under oxidative stress.
Microbiol. Read. Engl. 169:001396.

Gellatly SL, Needham B, Madera L, Trent MS, Hancock REW. 2012. The Pseudomonas
aeruginosa PhoP-PhoQ two-component regulatory system is induced upon
interaction with epithelial cells and controls cytotoxicity and inflammation. *Infect.*
Immun. 80:3122–3131.

Granato ET, Ziegenhain C, Marvig RL, Kümmerli R. 2018. Low spatial structure and
selection against secreted virulence factors attenuates pathogenicity in
Pseudomonas aeruginosa. *ISME J.* 12:2907–2918.

Hammond JH, Dolben EF, Smith TJ, Bhuju S, Hogan DA. 2015. Links between Anr and
quorum sensing in Pseudomonas aeruginosa biofilms. *J Bacteriol.* 197:2810–2820.

Hickman JW, Harwood CS. 2008. Identification of FleQ from Pseudomonas aeruginosa as a
c-di-GMP-responsive transcription factor. *Mol. Microbiol.* 69:376–389.

Hopkins HA, Lopezguerra C, Lau M-J, Raymann K. 2024. Making a pathogen? Evaluating the
impact of protist predation on the evolution of virulence in Serratia marcescens.
Genome Biol. Evol. 16:evae149.

Hosseinidoust Z, Tufenkji N, van de Ven TGM. 2013. Predation in homogeneous and
heterogeneous phage environments affects virulence determinants of
Pseudomonas aeruginosa. *Appl. Environ. Microbiol.* 79:2862–2871.

Jain C, Rodriguez-R LM, Phillippy AM, Konstantinidis KT, Aluru S. 2018. High throughput ANI
analysis of 90K prokaryotic genomes reveals clear species boundaries. *Nat.*
Commun. 9:5114.

Jukes TH, Cantor CR. 1969. Evolution of protein molecules. In: Munro HN, editor.
Mammalian protein metabolism. New York: Academic Press. p. 21–132.

Kanehisa M. 2000. KEGG: Kyoto encyclopedia of genes and genomes. *Nucleic Acids Res.*
28:27–30.

Kanehisa M, Sato Y, Kawashima M, Furumichi M, Tanabe M. 2016. KEGG as a reference
resource for gene and protein annotation. *Nucleic Acids Res.* 44:D457–D462.

Kerr KG, Snelling AM. 2009. Pseudomonas aeruginosa: a formidable and ever-present
adversary. *J. Hosp. Infect.* 73:338–344.

Laborda P, Martínez JL, Hernando-Amado S. 2022. Evolution of habitat-dependent
antibiotic resistance in Pseudomonas aeruginosa. *Microbiol. Spectr.* 10:e0024722.

Lear L, Hesse E, Buckling A, Vos M. 2022. Copper selects for siderophore-mediated
virulence in Pseudomonas aeruginosa. *BMC Microbiol.* 22:303.

Leong W, Poh WH, Williams J, Lutz C, Hoque MM, Poh YH, Yee BYK, Chua C, Givskov M,
Sanderson-Smith M, et al. 2022. Adaptation to an amoeba host leads to
Pseudomonas aeruginosa isolates with attenuated virulence. *Appl. Environ.*
Microbiol. 88:e02322-21.

Levin BR. 1996. The evolution and maintenance of virulence in microparasites. *Emerg.*
Infect. Dis. 2:93–102.

Matz C, Moreno AM, Alhede M, Manefield M, Hauser AR, Givskov M, Kjelleberg S. 2008.
Pseudomonas aeruginosa uses type III secretion system to kill biofilm-associated
amoebae. *ISME J.* 2:843–852.

Mould DL, Botelho NJ, Hogan DA. 2020. Intraspecies signaling between common variants
of Pseudomonas aeruginosa increases production of quorum-sensing-controlled
virulence factors. mBio 11:e01865–20.

Schwengers O, Jelonek L, Dieckmann MA, Beyvers S, Blom J, Goesmann A. 2021. Bakta:
rapid and standardized annotation of bacterial genomes via alignment-free
sequence identification. *Microb. Genomics* 2021:000685

Scribner MR, Stephens AC, Huong JL, Richardson AR, Cooper VS. 2022. The nutritional
environment is sufficient to select coexisting biofilm and quorum-sensing mutants
of Pseudomonas aeruginosa. *J. Bacteriol.* 204:e00444–21.

Silby MW, Winstanley C, Godfrey SAC, Levy SB, Jackson RW. 2011. Pseudomonas
genomes: diverse and adaptable. *FEMS Microbiol. Rev.* 35:652–680.

Stover CK, Pham XQ, Erwin AL, Mizoguchi SD, Warrener P, Hickey MJ, Brinkman FSL,
Hufnagle WO, Kowalik DJ, Lagrou M, et al. 2000. Complete genome sequence of
Pseudomonas aeruginosa PAO1, an opportunistic pathogen. *Nature* 406:959–964.

Tonkin-Hill G, MacAlasdair N, Ruis C, Weimann A, Horesh G, Lees JA, Gladstone RA, Lo S,
Beaudoin C, Floto RA, et al. 2020. Producing polished prokaryotic pangenomes with
the Panaroo pipeline. *Genome Biol.* 21:180.

Wong A, Rodrigue N, Kassen R. 2012. Genomics of adaptation during experimental
661   evolution of the opportunistic pathogen Pseudomonas aeruginosa. *PLoS Genet.*
662   8:e1002928.

663   Wu M, Li X. 2015. Klebsiella pneumoniae and Pseudomonas aeruginosa. In: Tang Y-W,
664   Sussman M, Liu D, Poxton I, Schwartzman J, editors. Molecular medical
665   microbiology (second edition). Boston: Academic Press. p. 1547–1564.
666

Supplemental Material

Figure S1. Phenotypic characteristics of evolved isolates. Assays performed were: (A) biofilm
production, (B) swimming motility, (C) swarming motility, (D) siderophore production, (E) casein
hydrolysis, (F) hemolytic activity, (G) growth in media after 24 h based on CFU counts, and (H)
growth in co-culture with *T. thermophila* after 24 h based on CFU counts. Statistical significance
was determined using a Kruskal–Wallis test, followed by Dunn's multiple comparisons test. For all
analyses, the mean rank of each group was compared to that of the ancestral strain. Comparisons
were performed for each evolved isolate individually (asterisks on the left) and for the mean of all
ME and PE isolates (asterisks on the right) relative to the mean of the ancestor. *, P < 0.05; **, P <
0.01.

Figure S2. Virulence and growth of evolved isolates in honey bees. Proportion of dead (A)
conventional (CV) honey bees and (B) microbiota-depleted (MD) honey bees after 5 days of

exposure to the ancestral strain or individual PE and ME isolates. (C) Growth of the ancestor, PE, and ME isolates in microbiota-depleted honey bees based on colony-forming unit (CFU) counts; CFU values are shown as $\log_{10}$-transformed counts. In panels A and B, each data point represents a cup cage containing 20 bees for CV trials and 10 bees for MD trials (biological replicates with technical replication). In panel C, each data point represents an individual bee. Statistical significance was tested both between each individual PE and ME isolate and the ancestral strain, and between the mean of PE and ME isolates and the ancestral strain. Statistical significance was determined using a Kruskal–Wallis test followed by Dunn's multiple comparisons test ($*$, $P < 0.05$; $**$, $P < 0.01$).

Sample	Line	Average Coverage	% Aligned
Isolate	PE-1	31.90703	99.45%
Isolate	PE-2	44.70393	99.37%
Isolate	PE-3	32.88026	99.42%
Isolate	PE-4	28.97971	98.98%
Isolate	PE-5	39.65268	98.99%
Isolate	PE-6	24.55709	98.96%
Isolate	ME-1	19.68769	99.06%
Isolate	ME-2	47.9632	99.19%
Isolate	ME-3	93.0591	99.15%
Isolate	ME-4	35.00678	99.02%
Isolate	ME-5	29.0038	98.59%
Isolate	ME-6	55.80903	99.05%
Population	PE-1	38.33171	99.15%
Population	PE-2	282.7614	99.21%
Population	PE-3	49.2864	98.80%
Population	PE-4	66.66734	98.29%
Population	PE-5	244.5668	98.99%
Population	PE-6	91.32875	97.91%
Population	ME-1	81.95025	98.53%
Population	ME-2	39.79088	98.71%
Population	ME-3	89.21901	98.66%
Population	ME-4	38.41933	98.57%
Population	ME-5	30.37836	97.94%
Population	ME-6	71.459	98.28%

Table S1. Average genome coverage of evolved populations and isolates.

Dataset S1. Mutations identified in evolved populations. List of mutations detected across all evolved populations at a frequency greater than 5%.

Dataset S2. Mutations identified in evolved isolates. List of mutations detected across all evolved isolates at a frequency greater than 5%.

Dataset S3. Spearman correlation results for genotype–phenotype, genotype–genotype, and
phenotype–phenotype comparisons. Complete Spearman's rank correlation coefficients (ρ) and
corresponding P-values for all pairwise comparisons across evolved lines. Mutations were
encoded as binary variables (1, present; 0, absent) for each line. Phenotypic values represent
means across all biological and technical replicates and are expressed as $\log_2$ fold changes
relative to the ancestral phenotype.