Multi-omics Profiling Identifies Molecular and Cellular Signatures of Regular
Physical Activity in Human Peripheral Blood

Xuemei Song[1,2]†, Jingzhi Lv[3]†, Siliang Ge[4,5]†, Sihan Xu[6], Yisha Wu[1,7], Yuhui Zheng[1,8],
Wenwen Zhou[1,8], Lifeng Li[9], Yan Zhang[1], Jiaqiang Zhang[10,11], Zhenxia Chen[2], Peng
Gao[12,13]*, Pengbin Yin[4,5]*, Jianhua Yin[1,13,14]*, Chuanyu Liu[1,13,14,15]*

[1]State Key Laboratory of Genome and Multi-omics Technologies, BGI Research,
Shenzhen 518083, China

[2]College of Life Sciences and Technology, Huazhong Agricultural University, Wuhan
430070, China

[3]State Key Laboratory of Molecular Oncology, Key Laboratory of Carcinogenesis and
Translational Research (Ministry of Education/Beijing), Peking University Cancer Hospital
& Institute, Beijing 100021, China.

[4]Department of Orthopedics, Chinese PLA General Hospital, Beijing 100853, China.

[5]National Clinical Research Center for Orthopedics, Sports Medicine & Rehabilitation,
Beijing 100853, China.

[6]Henan Institute of Medical and Pharmaceutical Sciences & BGI College, Zhengzhou
University, Zhengzhou 450052, China

[7]Department of Otolaryngology, Head and Neck Surgery, Second Hospital, Shanxi
Medical University, Taiyuan 030001, China

[8]College of Life Sciences, University of Chinese Academy of Sciences, Beijing 100049,
China

[9]BGI Sports Genomics Institute, Shenzhen 518083, China

[10]Department of Anesthesiology and Perioperative Medicine, People's Hospital of
Zhengzhou University, Henan Provincial People's Hospital, People's Hospital of
Henan Academy of Innovations in Medical Science, Zhengzhou 450003, China

[11]Institute of Electrophysiology, Henan Academy of Innovations in Medical Science,
Zhengzhou 451163, China

[12]BGI, Shenzhen, China.

[13]Shanxi Medical University-BGI Collaborative Center for Future Medicine, Shanxi

Medical University, Taiyuan 030001, China

[14]Shenzhen Proof-of-Concept Center of Digital Cytopathology, BGI Research, Shenzhen

518083, China

[15]Zhangzhou Affiliated Hospital of Fujian Medical University, Zhangzhou 363000, China

*Corresponding author. Email: gaopeng@genomics.cn (P.G.); yinpengbin@gmail.com

(P.Y.); yinjianhua@genomics.cn (J.Y.); liuchuanyu@genomics.cn (C.L.);

†These authors contributed equally to this work.

Abstract

Regular physical activity is well established to protect against metabolic disorders and

bolster immunity; yet, the underlying molecular and cellular mechanisms remain

incompletely understood. We integrated plasma metabolomics and lipidomics with

single-cell transcriptomic and chromatin accessibility profiles to decode the systemic

impact of physical activity on human immunity and metabolism. Our data reveal that

regular physical activity is linked to a coordinated metabolic signature marked by

enhanced fatty acid oxidation and antioxidant defense. In circulating immune cells,

regularly active individuals exhibited synchronous enhancement at both the chromatin

accessibility and transcriptional levels for antigen presentation-related genes in

antigen-presenting cells (APCs), particularly in classical monocytes, naive B cells, and

switched memory B cells. Meanwhile, cytotoxic programs in CD8$^+$ cytotoxic T and mature

NK cells showed epigenetic pre-activation of effector function regulators. Intercellular

communication analysis further revealed that regular exercise enhanced MHC-I/II

signaling between APCs and T cells and suppressed inflammatory signaling networks.

Together, these findings elucidate molecular mechanisms underlying the health benefits of

regular exercise and offer a theoretical basis for enhancing public health and preventing

chronic diseases.

Introduction

Regular physical activity reduces the risk of chronic diseases[1], lowers cancer risk[2, 3], while improving patient outcomes[4], enhances resistance to infection[5], delays aging[6], and mediated by circulating factors[7, 8], ultimately integrating into whole-body benefits. However, existing mechanistic evidence is derived predominantly from human intervention studies or model organisms[9–12]. Such controlled interventions differ substantially from free-living, spontaneously established regular physical activity habits. This discrepancy may lead to an incomplete understanding of the molecular and cellular mechanisms through which regular exercise confers clinical benefits, thereby limiting the external validity of current research findings.

Notably, metabolism and immunity are widely regarded as two central pillars through which exercise exerts its health-promoting effects[13, 14]. Metabolite and lipid profiling studies have identified candidate "exercise mimetics" or exercise-responsive metabolites, such as betaine, which has recently been characterized as an endogenous molecule triggered by sustained exercise that exerts systemic anti-inflammatory and neuroprotective effects[10]; in parallel, flow cytometry and immunophenotyping studies have systematically revealed the remodeling effects of regular physical activity on the human immune system[15]. Regular exercisers exhibit higher diversity of the naive and memory T-cell repertoire, a lower proportion of senescent T cells[16, 17], a more anti-inflammatory phenotype of monocyte subsets[18, 19], and enhanced functional adaptability, reduced senescence markers, and improved metabolic fitness in natural killer (NK) cells[20]. Building on these advances, there is a growing need to move beyond isolated measurements toward a systems-level understanding of how regular physical activity remodels metabolic and immune networks. In particular, we still lack an integrated view of the heterogeneous responses within and across specific immune cell subsets, their dynamic interactions with one another, and how these processes are orchestrated through epigenetic regulation and metabolic reprogramming in response to exercise. Addressing these questions with integrated multi-omics approaches will be essential to construct a more holistic atlas of how regular physical activity reshapes immunometabolic circuits and, ultimately, to translate these insights into more precise and scalable health

interventions.

Here, we conducted a comprehensive multi-omics study leveraging data from the

Chinese Immune Multi-omics Cohort (CIMA)(*21, 22*), a large-scale population-based

cohort. By integrating plasma targeted metabolomics and lipidomics with single-cell

transcriptomic and epigenomic analysis of peripheral blood mononuclear cells (PBMCs),

we systematically mapped the molecular and cellular correlates of regular physical activity

by comparing individuals with regular physical activity with sedentary controls. Our data

reveal that regular physical activity is linked to a coordinated metabolic signature marked

by enhanced fatty-acid oxidation and antioxidant defenses, together with widespread

remodeling of immune cell states mediated by epigenetic reprogramming. Collectively,

these results provide, to our knowledge, multi-dimensional atlas of molecular adaptations

to regularly spontaneous exercise, spanning metabolism, chromatin regulation, and

immunity. Beyond deepening mechanistic insight into how regular physical activity

promotes health, this work highlights candidate biomarkers and regulatory pathways that

may inform prevention and management of chronic diseases.

Results

Study Cohort and Physiological Characterization

Leveraging the CIMA natural cohort, we categorized participants based on self-reported

physical activity (Figure 1A). Our classification criteria were informed by the WHO

Guidelines on Physical Activity and Sedentary Behaviour(*23*). The regular exercise group

(EX, n = 40) was defined as individuals who engaged in physical activity ≥ 3 times per

week, accumulating ≥ 150 minutes of moderate-intensity or ≥ 75 minutes of

vigorous-intensity activity. The sedentary control group (SED, n = 46) was defined as

individuals who reported a weekly exercise duration of 0 minutes and identified "sitting" as

their primary daily posture. The study cohort was characterized by a mean age of 28.2

years (SD ± 3.1) and body mass index of 21.3 kg/m² (SD ± 3.0) (Figure S1A). The two

groups were well-balanced at baseline, with no statistically significant disparities observed

in age, diet, smoking or alcohol status,or sleep duration(Figure S1A and Table S1). The

SED group was predominantly female, consistent with global public health data showing

higher physical inactivity rates among women(*24, 25*). This sex imbalance in our cohort

led to a significantly higher unadjusted mean BMI in EX. Given that sex and Body Mass

Index (BMI) are key covariates for metabolic and immune phenotypes, subsequent

analyses employed multiple linear regression models adjusted for sex and BMI to

independently assess the effects of exercise.

To validate the physiological relevance of the classification based on exercise habits,

we first compared routine blood biochemical markers. Of the 19 indices tested, eight

differed significantly between EX and SED (Figure 1B). These alterations were reflected in

glucose and lipid metabolism indicators: compared to SED, EX showed significantly lower

fasting blood glucose (GLU) and triglyceride (TG) levels, along with significantly higher

high-density lipoprotein cholesterol (HDL-C) levels (Figure S1B). This finding is consistent

with a substantial body of prior research concluding that exercise confers significant

cardiometabolic benefits(*26–28*). Beyond these classic signatures, EX showed

significantly elevated levels of creatinine, uric acid, and bilirubin, all within normal clinical

ranges (Figures 1B and S1C). Notably, creatinine is a product of muscle metabolism, and

its elevation is directly associated with increased muscle mass and heightened metabolic

turnover in exercise-trained individuals(*29*). Meanwhile, uric acid and bilirubin possess

antioxidant properties, and their moderate elevation may reflect an enhancement in

systemic antioxidant defense capacity(*30, 31*).

In summary, classification based on exercise habits effectively captured distinct

physiological states between EX and SED. These phenotypic data indicate that EX

exhibits a multifaceted beneficial profile characterized by improved glucose and lipid

metabolism, enhanced muscle metabolism, and augmented systemic antioxidant

capacity.

Figure 1. Study overview and multi-omics characterization of regular exercisers and sedentary controls.

(A) Overview of the study design. A total of 86 participants from the CIMA cohort were classified into regular exercise (EX, n = 40) and sedentary (SED, n = 46) groups; all data were obtained from this cohort.

(B) Left: pie chart showing the number of significantly changed (n = 8) and unchanged (n = 11) blood biochemical indices. Right: dot plot showing log$_2$ fold change (log$_2$FC) of blood biochemical indices in EX compared with SED, coloured by metabolic category.

(C) Lollipop plot showing the number of significantly up-regulated (red) and down-regulated (blue) plasma metabolites and lipids in EX compared with SED, grouped by molecular class.

(D) UMAP embedding and circular bar plot showing immune cell subtypes and their proportions identified by scRNA-seq (n = 1,317,665 cells), coloured by major lineage.

(E) UMAP embedding and circular bar plot showing immune cell subtypes and their proportions identified by scATAC-seq (n = 644,045 cells), coloured by major lineage.

(F) Dot plot showing the average expression (colour scale) and percentage of expressing cells (dot size) of canonical marker genes across major immune lineages.

(G) Genome browser tracks showing chromatin accessibility at marker gene loci (*MS4A1*, *IL7R*, *CD4*, *CD8B*, *LYZ*, *KLRB1*, and *RORC*) across major immune cell types.

(H) Beeswarmand and box plots showing the distribution of log_2FC differences in cell neighborhoods across major immune cell lineages (EX vs. SED), as determined by Milo differential abundance analysis. Each point represents a cell neighborhood; blue and green indicate decrease and increase in EX, respectively. Box plots show median and interquartile range (IQR); whiskers extend to 1.5×IQR from the hinges.

Multi-omics Profiling of Plasma and Circulating Immune Cells in Exercise versus Sedentary Individuals

To comprehensively elucidate the regulatory effects of regular exercise on systemic metabolism and immune features, we conducted a multi-omics analysis using data from 86 subjects in the CIMA cohort. This analysis integrated targeted plasma metabolomics, lipidomics, single-cell RNA sequencing (scRNA-seq), and single-cell assay for transposase-accessible chromatin sequencing (scATAC-seq) of PBMCs (Figure 1A). First, we performed metabolomic and lipidomic profiling on the plasma samples. Following standardized pre-processing, 321 metabolites and 718 lipid species were utilized for downstream analysis (Table S2). Comparative analysis revealed a distinct class-specific distribution pattern among the differential metabolites between EX and SED (Figure 1C). Specifically, metabolites associated with energy metabolism and substrate turnover (acylcarnitines and organic acids) were broadly upregulated in EX, whereas complex

lipids were downregulated (Figure 1C and Table S2), indicating that regular exercise

significantly remodels human metabolic program.

Subsequently, to investigate the impact of exercise on the immune system, we

conducted scRNA-seq analysis on 1,317,665 high-quality cells from EX and SED (Figure

1D). Simultaneously, we performed an independent re-analysis and stringent filtering of

the matched scATAC-seq raw fragments, ultimately retaining 644,045 high-quality cells

(Figure 1E). Rigorous quality assessment ensured high data quality for both omics

datasets (Figures S2A and S2B), with no significant batch effects detected between EX

and SED (Figure S2C). Regarding cell type identification, we annotated the scRNA-seq

data following the hierarchical annotation system of the CIMA cohort. The first level (L1)

included six major immune cell types: myeloid cells, B cells, CD8$^+$ T cells, CD4$^+$ T cells,

NK cells, and non-canonical T (Unconv_T) cells, which were further divided into 34 distinct

subsets (Figures 1D, 1F and S2D). Subsequently, we annotated the scATAC-seq data

using a cross-modal mapping strategy, achieving high concordance with the

transcriptome at the L1 level and successfully resolving 30 subsets (Figure 1E). Gene

activity scores based on chromatin accessibility aligned closely with transcriptomic cell

type annotations (Figures S2E and S2F), with canonical marker gene promoters (*MS4A1*,

IL7R, *CD4*, *CD8B*, *KLRB1* and *RORC*) exhibiting the expected lineage-specific

accessibility patterns (Figure 1G). The proportions of major cell subsets remained stable

between EX and SED in the scATAC-seq data (Figure S2G). In contrast, scRNA-seq

revealed a significant increase in myeloid cells in EX (Figure S2G), a finding corroborated

by Milo analysis(*32*) (Figure 1H). These findings indicate that regular exercise habits

reshape the innate immune composition primarily at the transcriptomic level.

Systemic Metabolic and Lipidomic Remodeling

To decipher the impact of regular physical activity on systemic metabolic homeostasis, we

performed targeted metabolomics and lipidomics analyses on plasma samples from EX

and SED. A total of 62 differentially abundant molecules were identified (Figure 2A).

Specifically, 86% of the differential metabolites were upregulated in EX, including

O-Adipoylcarnitine and p-Hydroxymandelic acid, while 94% of the differential lipids were

downregulated, such as PG 18:1-18:2 and PC 18:1-20:4 (Figure 2A and Table S2).

Pathway enrichment analysis revealed that the upregulated metabolites were significantly

enriched in core energy metabolism pathways such as the tricarboxylic acid cycle (TCA

cycle), as well as various amino acid conversion pathways involving

phenylalanine/tyrosine metabolism and glycine, serine, and threonine metabolism,

indicating more active energy metabolism and amino acid turnover in the exercising

population (Figure 2B). Furthermore, pathways related to cofactor metabolism and

glutathione metabolism, which is a key antioxidant defense system, were also significantly

enriched.

At the molecular level, we first focused on fatty acid oxidation (FAO), a central

energy-yielding pathway during exercise (Figure 2C). The significant upregulation of

O-adipoylcarnitine, a key acylcarnitine species reflecting mitochondrial fatty acid loading,

indicates an increased flux of fatty acid transport into the mitochondria(33). Concurrently,

elevated levels of 3-hydroxybutyrate, a primary ketone body, indicate the activation of

hepatic fatty acid -oxidation and ketogenesis(34). Furthermore, the accumulation of acetic

acid, a downstream product of fatty acid metabolism, further supports a comprehensive

mobilization of FAO pathways across varying chain lengths(35). The coordinated

elevation of these three core metabolites demonstrates that regular exercise

systematically enhances the body's fatty acid oxidative capacity.

Given that heightened energy metabolism is often accompanied by increased

reactive oxygen species (ROS) production(36), we detailedly examined exercise-induced

modulation of the antioxidant defense system (Figure 2D). In concert with elevated energy

flux, several metabolites involved in redox homeostasis were synchronously upregulated

in EX. Betaine, an important methyl donor, showed significantly higher plasma levels in

EX. Notably, this molecule has recently been characterized as an endogenous exercise

mimetic conferring anti-inflammatory and anti-aging protection(10). Furthermore, the

concurrent elevation of 3-hydroxyanthranilic acid, which possesses radical-scavenging

activity(37), and 2-hydroxybutyrate, an indirect marker of glutathione synthetic flux(38),

collectively suggests that regular exercise reinforces a multi-layered endogenous

antioxidant defense system. This trend was also observed for immunomodulatory

metabolites, as exemplified by the increase in xanthurenic acid (another product of the

tryptophan-kynurenine pathway) and γ-aminobutyric acid (GABA) (Table S2). These

molecules can regulate the function and inflammatory status of immune cells such as T

cells and monocytes/macrophages through the aryl hydrocarbon receptor or GABAergic

signaling pathways(*39, 40*).

Furthermore, we observed that regular exercise also reshapes gut microbiota-related

metabolites (Figure 2E). Levels of microbially derived aromatic amino acid

metabolites—phenyllactic acid and indoleacrylic acid—were significantly elevated in EX.

Indoleacrylic acid has been shown to enhance intestinal barrier function and modulate

innate immunity, thereby contributing to reduced systemic inflammation(*41*). In parallel,

levels of two secondary bile acids, glycodeoxycholic acid and glycochenodeoxycholic acid,

were reduced in EX. Bile acids are signaling molecules that regulate intestinal and hepatic

immunity(*42*), and the altered levels of these secondary bile acids may influence local and

systemic immune activity.

At the lipid level, a broad downregulation was observed in EX. Triacylglycerols (TAG),

the primary energy-storage form, exhibited widespread reduction (Figure 2F), consistent

with the well-established effect of exercise-promoted lipolysis and fatty acid utilization(*43*).

Multiple membrane phospholipid subclasses, including lysophosphatidylethanolamine

(LPE), phosphatidylcholine (PC), phosphatidylglycerol (PG), and phosphatidylinositol (PI),

also showed decreased abundance (Figure S3A), reflecting extensive remodeling of the

lipid metabolic network. Notably, distinct species within the same

phosphatidylethanolamine (PE) subclass displayed opposite abundance changes: while

PE 20:4–22:5 was downregulated, PE 20:4–22:1 was upregulated in EX (Figure S3B).

Overall, these lipid alterations, together with enhanced fatty acid oxidative capacity,

constitute a characteristic metabolic adaptation to regular exercise.

Figure 2. Plasma metabolomic and lipidomic remodeling in regular exercisers.

(A) Volcano plot of differentially abundant plasma metabolites (circles) and complex lipids

(triangles) in EX vs. SED. Pie chart inset shows the proportion of complex lipids and

metabolites among the 62 significantly altered molecules.

(B) Bar plot of enriched pathways of up-regulated metabolites in EX, coloured by

pathway category.

(C–E) Beeswarm and box plots showing normalized plasma levels of key metabolites

related to fatty acid oxidation (C), antioxidant defense (D), and gut microbiota-bile acid

metabolism (E) in SED and EX.

(F) Dot plot showing mean intensity (± SE) of significantly altered triacylglycerol (TAG)

species in SED and EX.

P-values are indicated above each comparison. See also Table S2.

Myeloid Cells Exhibit Enhanced Antigen Presentation Following Regular Exercise

Given the pivotal role of myeloid cells in sensing metabolic signals and regulating

inflammation and antigen presentation(*44*, *45*), elucidating how regular physical activity

remodels these cell populations is essential. Myeloid cells were stratified into five subsets

using canonical markers: classical monocytes (cMono), intermediate monocytes

(intMono), non-classical monocytes (ncMono), classical dendritic cells (cDC), and

plasmacytoid dendritic cells (pDC) (Figures 3A, S4A, and S4F). Conventional cell

proportion analysis showed no significant differences (Figure S5A). Differential

abundance testing with the Milo analysis at the local neighborhood level revealed

significant expansions of intMono, cMono, and ncMono in EX (median log_2FC: +0.37,

+0.31, +0.25; Figure 3B). This suggests that regular physical activity remodels circulating

myeloid cells, with a pronounced effect on the composition of monocyte subsets.

To dissect the functional alterations of myeloid subsets induced by regular exercise,

we performed differentially expressed gene (DEG) and pathway enrichment analyses

(Table S3). Gene Ontology (GO) analysis of upregulated genes in EX revealed

enrichment in antigen processing and presentation, major histocompatibility complex

(MHC) protein complex assembly, and leukocyte chemotaxis, whereas downregulated

DEGs were predominantly linked to oxidative stress and hypoxia responses (Figure 3C).

Pathway scoring analysis further corroborated these shifts, manifesting as enhanced

antigen presentation capabilities alongside a systemic attenuation of inflammatory and

stress-related signatures across multiple subsets in EX (Figure S5B). Consistent with their

enhanced functional state, the core antigen presentation gene *HLA-DQB1* showed

pronounced elevation in cMono, ncMono, and pDC, while *HLA-DQA1* and the innate

immune sensor *FCN1* were markedly upregulated in cDC (Figure 3D). Furthermore, pDC

exhibited increased expression of the key transcriptional regulator *STAT1*, alongside

elevated levels of migration-associated receptors, including *CX3CR1* in cDC and *CXCR3*

in pDC, suggesting a concurrent enhancement of antigen presentation and cellular

migration capacity in myeloid cells.

To investigate the epigenetic mechanisms underpinning these functional changes, we

compared the chromatin accessibility of myeloid cell subsets between EX and SED using

scATAC-seq. Analysis of differentially accessible regions (DARs) revealed that 94.3% of

DARs across all myeloid subsets were specifically enriched in cMono (Figure S5C and

Table S4). GO enrichment analysis of DARs in cMono showed that regions with increased accessibility were enriched for pathways such as antigen presentation and T cell activation, while regions with decreased accessibility were associated with stress response pathways (Figure 3E). Transcription factor (TF) motif enrichment and motif deviation analysis characterized a distinct regulatory signature in EX (Figures 3F and 3G) . Specifically, regions with increased accessibility in EX were primarily enriched for TFs supporting antigen presentation (*IRF1* and *SPI1*)(*46, 47*), viral defense (*STAT2*)(*48*), and immune homeostasis and migration (*KLF2* and *KLF4*)(*49, 50*). Conversely, the accessibility of motifs for regulatory factors mediating stress and inflammatory responses, such as the AP-1 family (*JUN, JUNB, FOS* and *FOSB*) and *BACH1*, was significantly reduced. To determine the epigenetic basis for enhanced antigen presentation, we examined chromatin accessibility at loci linked to this function in cMono. Analysis showed that the promoter regions of multiple key antigen presentation genes (*HLA-DQA1* and *HLA-DQB1*) exhibited significantly enhanced chromatin accessibility in EX (Figure 3H), consistent with their upregulated mRNA expression in this subset (Figure 3D). Additionally, EX exhibited reduced chromatin accessibility at the promoters of *CXCL8* and *CCL7* and a distal regulatory element associated with *CCL2* (Figure S5D). The reduced promoter accessibility of *CXCL8* was consistent with its downregulated mRNA expression (Figure S5E).

Next, we assessed regulon factor activity in cMono at the transcriptional level using the Single-Cell Regulatory Network Inference and Clustering (SCENIC) algorithm(*51*). Results demonstrated that the activities of transcription factors such as *SPI1*, *STAT1*, *IRF1*, *and KLF4* were significantly higher in EX than in SED (Figure 3I), highly consistent with the scATAC-seq data indicating upregulated accessibility of the corresponding motifs (Figures 3F and 3G). To understand how key transcription factors cooperatively regulate cMono function, we identified a core regulatory module driven by *SPI1*, *IRF1*, *KLF4*, *and STAT1* (Figure 3J). This core hub orchestrates diverse immune programs, spanning antigen presentation (*HLA-DQA1* and *HLA-DQB1*) and the regulation of inflammation and chemotaxis (*IL1B*, *IL10*, *CCL20*, and *CCL2*). In summary, regular physical activity remodels the epigenetic landscape and transcriptional profiles of myeloid cells.

Collectively, EX exhibits enhanced antigen presentation and interferon production

alongside altered chemotaxis, with the most pronounced synergistic effects observed in

cMono.

Figure 3. Epigenetic and transcriptional characteristics of myeloid cells in regularly

active individuals.

(A) UMAP visualization of myeloid cell subsets identified by scRNA-seq (left) and

scATAC-seq (right), colored by cell type.

(B) Beeswarmand and box plots showing the distribution of $\log_2$FC differences in cell

neighborhoods across myeloid cell subsets (EX vs. SED). Box plots are created in a

similar fashion as in Figure 1H.

(C) Bubble plot of Gene Ontology (GO) enrichment analysis for differentially expressed

genes (DEGs) in myeloid subsets (EX vs. SED). Dot size represents -$\log_{10}$(adjusted

p-value); color indicates regulation direction (pink: upregulated in EX; blue: downregulated

in EX).

(D) Violin plots overlaid with box plots showing expression levels of selected genes

associated with antigen presentation, innate immunity, and cellular migration across

myeloid subsets in SED and EX. $\log_2$FC and adjusted p-values are indicated. **$p < 0.01$,

***$p < 0.001$.

(E) GO enrichment analysis of differentially accessible regions (DARs) in cMono (EX vs.

SED). Bars extending right (pink) indicate pathways enriched in open chromatin regions in

EX; bars extending left (blue) indicate decreased accessibility in EX. Bar length

represents $-\log_{10}$(p-value).

(F) Transcription factor (TF) motif enrichment analysis of differentially accessible peaks

with increased accessibility in cMono of EX. TFs are ranked by statistical significance;

top-ranked motifs are labeled.

(G) TF motif deviation analysis in cMono. Each point represents a TF motif; x-axis shows

mean chromatin accessibility difference (EX vs. SED); y-axis shows $-\log_{10}$(FDR). Green

and blue points indicate TF motifs with significantly increased and decreased accessibility

in EX, respectively; gray points indicate non-significant motifs.

(H) Genome browser tracks of chromatin accessibility at the *HLA-DQB1* (left) and

HLA-DQA1 (right) loci in cMono from SED and EX. Promoter DARs are highlighted;

$\log_2$FC and FDR values are indicated.

(I) Heatmap showing the scaled mean regulon areas under the curve per cell (AUCell)

scores of selected transcription factors in cMono between SED and EX. Red and green

indicate higher and lower regulon activity, respectively.

(J) Network diagram of the core transcriptional regulatory module in cMono, centered on

SPI1, *IRF1*, *KLF4*, and *STAT1*. Nodes represent target genes involved in antigen

presentation and inflammatory/chemotactic regulation; edge colors indicate regulatory

relationships.

Epigenetic Activation of Effector Functions in T Cells by Regular Exercise

Previous studies have shown that regular exercise can modulate the composition and

functional status of circulating T cells(*14*, *52*), how exercise drives these functional

changes via epigenetic reprogramming at the single-cell level remains to be

systematically elucidated. Using classical marker genes, we identified 17 T cell subsets,

consisting of ten CD8$^+$ and unconv_T cell subsets: CD8$^+$ naive T cells (CD8$^+$ Tn), CD8$^+$

central memory T cells (CD8$^+$ Tcm), CD8$^+$ effector memory T cells (CD8$^+$ Tem), CD8$^+$

cytotoxic T cells (CD8$^+$ CTL), mucosal-associated invariant T cells (MAIT), γδT1, γδT2,

proliferating T, and pre-T-like cells (Figures 4A, S4B and S4G). Additionally, we identified

seven CD4$^+$ T cell subsets (Figures S7A, S4C and S4H). Proportional analysis revealed

that overall compositions of CD8$^+$ and unconv_T cells remained stable between EX and

SED (Figure S6A). However, Milo analysis identified a significant expansion of pre-T-like

cells in EX (Figure 4B). Within the CD4$^+$ T cell compartment, EX exhibited increased Treg

proportions at the epigenetic level (Figure S7B), accompanied by a reduction in CD4$^+$

cytotoxic T lymphocytes (CD4$^+$ CTLs) (Figure S7C).

We identified DEGs and performed functional enrichment analysis (Table S3), broad

transcriptomic alterations were present across these CD8$^+$ and unconv_T cells (Figure

S6B). GO enrichment analysis of upregulated genes in EX revealed significant enrichment

of pathways related to T cell activation, T cell receptor (TCR) signaling, IFN-γ response,

and lymphocyte-mediated immunity (Figure 4C). Within the CD4$^+$ compartment,

transcriptional changes were most pronounced in CD4$^+$ Tn and T Helper cells (Figure

S7D), with upregulated genes enriched in T cell activation, IL-12 production, and type I

interferon signaling (Figures S7E and S7F). Pathway activity scoring further corroborated

these functional shifts, revealing significantly elevated scores in EX across multiple T cell

subsets, including CD8$^+$ CTL, CD8$^+$ Tem, CD8$^+$ Tn, NKT, MAIT, and γδT1 cells, for

processes related to T cell-mediated immunity, cytotoxicity, T cell activation, and type II

interferon production (Figure S6C). Notably, the expression of key cytotoxic genes (*GZMH*,

GZMA and *PRF1*) was significantly increased in CD8$^+$ CTL, CD8$^+$ Tem, and NKT cells

(Figure S6D). Overall, the transcriptomic features collectively indicate a systemic shift in T

cell function, characterized by a broad upregulation of activation programs and a

reinforcement of cytotoxic effector.

To determine how regular exercise drives functional adaptation in T cells through

epigenetic reprogramming, we performed DAR analysis across all subsets (Table S4).

CD8$^+$ CTLs exhibited the highest number of upregulated DARs (Figure 4D), with multiple

DARs shared between two or more subsets. In CD8$^+$ CTLs, upregulated DARs were

enriched for pathways related to cytotoxic T cell differentiation, TCR signaling, and

lymphocyte-mediated immunity. In CD8$^+$ Tn cells, enrichment was observed for pathways

involved in T cell activation and immune response regulation. Pathways associated with

IFN-γ signaling, antigen receptor-mediated signaling, and TCR signaling were commonly

enriched in CD8$^+$ CTLs, MAIT cells, and NKT cells. Additionally, NKT cells showed

enrichment for innate immune response and positive regulation of lymphocyte activation

(Figure 4E).

To resolve the transcriptional regulatory basis of these functional changes, we

performed motif deviation analysis based on scATAC-seq data. In CD8$^+$ CTLs, the

upregulated DARs were significantly enriched for binding motifs of effector function

master regulators, including *EOMES*, *RUNX3*, *RUNX1*, and *TBX21* (Figures 4F and S6E).

TBX21 (T-bet) and *EOMES* are master regulators driving CD8$^+$ T cell differentiation into

effector cells, and their upregulation is directly linked to enhanced cytotoxic function(*53*,

54). Similarly, motif deviation analysis identified increased accessibility for key

transcription factor binding motifs in other subsets. These included motifs for *RORC*,

RUNX3, *STAT1*, and *STAT4* in MAIT cells (Figure S6F), and for *LEF1*, *TCF7*, and *RUNX3*

in CD8$^+$ Tn cells (Figure S6G). Notably, chromatin accessibility was significantly increased

at distal regulatory regions of the effector genes *TBX21* and *GZMB* in CD8$^+$ CTLs (Figure

4G). Similarly, accessibility was elevated at the promoter of *IRF1* in MAIT cells (Figure

S6H) and within an intron of *TCF7* in CD8$^+$ Tn cells (Figure S6I). SCENIC analysis

confirmed elevated regulatory activity of key transcription factors, including *RUNX3* and

TBX21, in CD8$^+$ CTLs of EX (Figure 4H). Given that *RUNX3* showed the most significant

change across both omics layers, we constructed its downstream regulatory network

(Figure 4I), which confirmed that *RUNX3* directly targets key effector genes, including

GZMB, *TBX21*, and *IRF1*.

In summary, single-cell multi-omics data reveal a global remodeling of T cell function

with regular exercise: it not only enhances the cytotoxic execution capacity of effector

cells, with CD8$^+$ CTLs at the core, but also broadly augments the signal responsiveness

and activation preparedness of naive and non-canonical subsets.

Figure 4. Epigenetic and transcriptional signatures of T cells associated with regular exercise.

(A) UMAP visualization of CD8+ and unconventional T cell subsets identified by scRNA-seq (left) and scATAC-seq (right), colored by cell type.

(B) Beeswarmand and box plots showing the distribution of log2FC differences in cell neighborhoods across CD8+ and unconventional T cell subsets (EX vs. SED). Box plots are created in a similar fashion as in Figure 1H.

(C) Bubble plot of GO enrichment analysis for upregulated DEGs in CD8ㄴ and

unconventional T cell subsets (EX vs. SED). Dot size represents gene ratio; color

indicates $-\log_{10}$(FDR).

(D) UpSet plot showing the number and overlap of upregulated DARs across CD8$^+$ and

unconventional T cell subsets in EX. Bar height indicates the number of DARs per subset

or intersection.

(E) Bubble plot of GO enrichment analysis for upregulated DARs across CD8$^+$ and

unconventional T cell subsets in EX. Dot size represents gene ratio; color indicates

$-\log_{10}$(FDR).

(F) Violin plots showing TF motif deviation Z-scores for *RUNX1*, *RUNX3*, *TBX21*, and

EOMES in CD8$^+$ CTL cells from SED and EX. Adjusted *p*-values are indicated. ***$p <$

0.001.

(G) Genome browser tracks of chromatin accessibility at distal regulatory regions of

GZMB (left) and *TBX21* (right) loci in CD8$^+$ CTL cells from SED and EX. Co-accessibility

links and distal DARs are highlighted; $\log_2$FC and FDR values are indicated.

(H) Heatmap showing the scaled mean regulon AUCell scores of selected transcription

factors in CD8$^+$ CTL cells between SED and EX. Red and green indicate higher and lower

regulon activity, respectively.

Network diagram of the *RUNX3* downstream transcriptional regulatory module in CD8$^+$

CTL cells.

Functional Reprogramming of the NK Cell Epigenome and Transcriptome by

Regular Exercise

As key innate effector cells, NK cells are pivotal in antiviral defense, tumor surveillance,

and bridging innate and adaptive immunity(*55*). However, the molecular mechanisms by

which regular exercise enhances NK cell function remain incompletely understood. We

first classified NK cells into six subsets based on classical marker genes: mature NK cells,

transitional NK cells, type 2 innate lymphoid cells (ILC2), CD56^{bright} NK cells, terminal NK

cells, and cycling NK cells (Figures 5A, S4D, and S4I). Notably, the proportion of

CD56^{bright} NK cells was significantly reduced in EX (Figure S8A), consistent with the

results obtained from the Milo algorithm (Figure 5B).

To evaluate functional changes in NK cells, we identified DEGs (Table S3) and

performed pathway scoring analysis (Figure S8B). Elevated scores for NK cell-mediated

cytotoxicity, NK cell activation, and IFN-γ response were observed across multiple NK

subsets in EX, whereas scores for inflammatory response and cellular stress response

were generally reduced (Figure S8B). In EX, subset-specific gene expression changes

were observed in NK cells. *GZMM* was upregulated in terminal NK cells, *IFITM1* in

CD56^{bright} NK cells, and *IFNG* and *CCL3L1* in mature NK cells. In contrast, *FOSB* and

JUN were downregulated in both transitional and mature NK subsets (Figure 5C).

Epigenetically, we identified DARs (Table S4), with mature NK cells exhibiting the

highest number of upregulated DARs (Figure S8C). GO enrichment analysis of these

upregulated DARs revealed that pathways related to "immune defense and killing,"

"immune regulation," and "immune cell mobilization" were significantly enriched in mature

NK cells, whereas inflammation and stress-related pathways were downregulated (Figure

5D). To deeply investigate the transcriptional regulatory basis, we performed motif

deviation analysis based on scATAC-seq data (Figure 5E). Results showed that in EX,

motif accessibility was significantly enhanced for TFs closely related to NK cell immune

function, including key regulators driving cytotoxic differentiation such as *RUNX3*, *EOMES*,

and *TBX21* (T-bet), as well as *ETS1*, which is involved in innate lymphoid cell

development and activation(*56*). Conversely, motif accessibility for inflammation and

stress-related TFs (*RELA*, *RELB*, *NFKB1*, *JUND*, *FOS*, *FOSL2 and BACH1*)(*56*) was

generally reduced. Footprint analysis confirmed the decreased accessibility of *RELA* and

RELB transcription factor binding sites (Figure S8D). Notably, in mature NK cells,

chromatin accessibility was significantly enhanced in EX at the intronic regulatory region

of *STAT1*, a core mediator of type I interferon signaling, and the promoter region of

CXCR4, a known exercise-responsive factor(*57*) (Figure 5F). Furthermore, accessibility

was significantly reduced at the promoter regions of *RELB*, associated with non-canonical

NF-κB pathway activation, and *CCL5*, an important inflammatory chemokine (Figure S8E).

In summary, these results demonstrate that regular exercise reshapes NK cells into a

more robust effector state, characterized by enhanced cytotoxic and mobilization

capacities alongside reduced levels of inflammation and stress.

To investigate whether common regulatory mechanisms exist for the cytotoxic

functions of adaptive and innate immunity under exercise, we compared transcription

factor activities between CD8$^+$ CTL (adaptive immunity) and mature_NK (innate immunity)

cells (Figure 5G). Venn analysis revealed that both cell populations shared a core set of

significantly upregulated transcription factors, including *RUNX3*, *EOMES*, *TBX21*, *FOXO3*,

and *ETS2*. This provides a unified transcriptional regulatory basis for the synergistic

enhancement of cytotoxic functions across different immune branches by regular

exercise.

Regular Exercise Potentiates B Cell Antigen Presentation   via Epigenetic

Reprogramming

Given the positive impact of regular exercise on vaccination responses and antibody

levels(*58*, *59*), we employed single-cell multi-omics technology to conduct an in-depth

analysis of B cells, the core executors of humoral immunity. B cells were annotated into

eight subtypes (Figures S9A, S4E, and S4J): switched memory B cells (Switched_Bm),

naive B cells (Naive_B), pre-switched memory B cells (pre-Switched_Bm), unswitched

memory B cells (Unswitched_Bm), atypical memory B cells (Atypical_Bm), transitional B

cells (Transitional_B), plasma cells, and plasmablasts. Cell proportions did not differ

significantly between EX and SED across these subsets (Figure S9B).

Next, we performed DEG analysis and GO enrichment analysis (Table S3).

Upregulation of pathways related to antigen processing and presentation, humoral

immune response, and chemotaxis was observed in naive_B, switched_Bm, and

transitional_B (Figure S9C). Notably, MHC molecules were significantly upregulated in EX

across multiple B cell subsets: *HLA-DQB1* in naive_B, switched_Bm, and unswitched_Bm;

HLA-DQA1 in transitional_B; and *HLA-F* in unswitched_Bm (Figure S9D). Additionally, the

chemokine *CCL5* and *S100A9* were significantly upregulated in switched_Bm cells, and

the actin regulator *COTL1* was upregulated in transitional_B cells (Figure S9D). To

resolve the epigenetic regulatory basis, we performed DAR analysis (Table S4). Results

showed that changes in chromatin accessibility were highly concentrated in naive_B cells

and switched_Bm (Figure S9E). GO enrichment analysis of the upregulated DARs in

these two subsets revealed that both were enriched in pathways related to antigen

presentation and positive regulation of B cell activation (Figure S9F).

To dissect the transcriptional regulation potentially influenced by these epigenetic

alterations, we analyzed TF activity in naive_B and switched_Bm. In naive_B, we

observed significant upregulation in the activity of key regulators. This included

BCL11A(*60*), a critical determinant of B-cell lineage fate; *RUNX1*(*61*), essential for

lymphocyte development; and the core master regulators of early B-cell commitment,

SPI1 (PU.1)(*62*) and *EBF1*(*63*) (Figure S9G). switched_Bm showed upregulation of

NF-κB family members (*RELA*, *RFLB* and *NFKB1*), *IRF4*, as well as *SPI1* and *EBF1*

(Figure S9H); NF-κB and *IRF4* are master regulators of memory B cell formation, class

switching, and plasma cell differentiation(*64*, *65*). We focused our analysis on the

regulatory regions of genes upregulated in the transcriptome. The results showed that in

EX, chromatin accessibility at the promoter region of the key antigen presentation gene

HLA-DQB1 was significantly enhanced in both naive_B (Figure S9I) and switched_Bm

(Figure S9J). Additionally, in switched_Bm, promoter accessibility of the chemokine gene

CCL5 also showed a synchronous increase (Figure S9K).

Overall, this study demonstrates that although regular exercise does not markedly

alter the proportions of B cell subsets, it may enhance their antigen presentation and

immune activation capabilities through chromatin remodeling and changes in gene

expression. Recent research has highlighted the critical role of B cell antigen presentation

in effective immunization and antiviral defense(*66*). Therefore, our findings suggest that

the robust immune response observed in exercised individuals could be attributed to

exercise-induced chromatin reprogramming and transcriptional modifications in B cells.

Figure 5. Epigenetic and transcriptional profiles of NK cells associated with regular

exercise.

(A) UMAP visualization of NK cell subsets identified by scRNA-seq (left) and scATAC-seq

(right), colored by cell type.

(B) Beeswarmand and box plots showing the distribution of $\log_2$FC differences in cell

neighborhoods across NK cell subsets (EX vs. SED), as determined by Milo differential

abundance analysis. Box plots are created in a similar fashion as in Figure 1H.

(C) Violin plots overlaid with box plots showing expression levels of selected genes across

NK cell subsets in SED and EX. $\log_2$FC adjusted *p*-values and statistical significance are

indicated. *$p < 0.05$, ***$p < 0.001$.

(D) GO enrichment analysis of DARs in Mature_NK cells (EX vs. SED). Bars extending right (pink) indicate pathways enriched in open chromatin regions in EX; bars extending left (blue) indicate decreased accessibility in EX. Bar length represents $-\log_{10}(p\text{-value})$.

(E) TF motif deviation analysis in Mature_NK cells. Each point represents a TF motif; x-axis shows mean chromatin accessibility difference (EX vs. SED); y-axis shows $-\log_{10}(FDR)$. Green and blue points indicate TF motifs with significantly increased and decreased accessibility in EX, respectively; gray points indicate non-significant motifs.

(F) Genome browser tracks of chromatin accessibility at the intronic DAR of *STAT1* (left) and promoter DAR of *CXCR4* (right) in Mature_NK cells from SED and EX. $\log_2FC$ and FDR values are indicated.

(G) Venn diagram showing the overlap of significantly upregulated TFs between CD8[+] CTL and Mature_NK cells in EX. Shared and cell type-specific TFs are listed.

Altered Intercellular Communication

Given the significant enhancement of antigen-presenting cells (APCs) function in EX, particularly in cMono, cDC, and B cells, we employed the CellChat algorithm to analyze whether this change remodeled the immune cell-cell interaction network at the systems level(*67*). Our analysis revealed that myeloid cells mediated the highest number of intercellular interactions among all cell types. Moreover, the most pronounced differential interaction strength between EX and SED was observed for myeloid cell–CD8[+] T cell pairs (Figures 6A and 6B). At the pathway level, we observed a clear pattern: information flows related to inflammation and immune suppression (THBS, RESISTIN and IL6) were dominant in SED, whereas signals driving chemotaxis, antigen presentation, and cell adhesion (CCL, MHC-II and PECAM1) were markedly enhanced in EX (Figure 6C). Human resistin is a key regulator directing inflammatory responses in various immune cells(*68*). Cell communication network showed that myeloid cells were the core signal source for this pathway, and their signal output was significantly lower in EX. (Figure S10A). The interaction intensity of key ligand-receptor axes in this pathway, RETN–CAP1 and RETN–TLR4, was weakened in EX. The downregulation of *RETN* gene expression was also observed at the transcript level in myeloid cells (Figures S10B and S10C).

Exercise significantly enhanced intercellular signaling associated with antigen presentation. Compared to SED, EX exhibited a generalized enhancement of MHC-II signaling from B cells and myeloid cells to CD4$^+$ T cells (Figures 6D and 6E). Subset analysis further pinpointed that interactions involving cMono, naive_B cells, and switched_Bm cells contributed most prominently within this pathway (Figure S10D). Besides, MHC-I signaling from various cell subsets (B cells, CD4$^+$ T cells, myeloid cells, NK cells, and unconv_T cells) to CD8$^+$ T cells showed a global increase.The enhancement of signaling mediated by the HLA-G–CD8A/B pair was the most pronounced (Figure 6F), primarily driven by interactions between cMono and CD8$^+$ Tn cells (Figure S10E). Notably, exercise also potentiated the signaling network mediated by IFN-γ. In EX, IFN-γ signaling from NKT cells to multiple myeloid and B cell subsets was markedly enhanced (Figure 6G). In parallel, cytotoxic T cells (CD4$^+$ and CD8$^+$ CTLs) also strengthened their signaling via the IFNG–receptor axis to these same innate immune cell populations (Figure 6H). IFN-γ, a pivotal cytokine that bridges innate and adaptive immunity by driving CD4$^+$ T cell differentiation toward the Th1 lineage, enhancing the cytotoxicity of CD8$^+$ T cells, and serving as a core effector molecule for NK cell function(*69, 70*).

Notably, exercise also potentiated signaling networks associated with tissue homeostasis and immune microenvironment organization. The COLLAGEN signaling pathway was markedly enhanced in EX, predominantly driven by unconv_T cells(Figures S10F). This was evidenced by the coordinated upregulation of both the key ligand-receptor pair COL6A2–CD44 and its corresponding ligand gene expression (Figures S10F-H). Collagen-mediated signaling is a well-established regulator of cell adhesion, migration, and tissue homeostasis(*71*). In addition, the CCL chemokine signaling pathway was significantly enhanced in EX, manifested primarily as increased signal output from other major immune cell lineages to myeloid cells (Figures S10I and S10J). CCL family chemokines are central orchestrators of immune cell recruitment and spatial positioning within tissues(*72*). In summary, regular exercise remodels the intercellular communication network among immune cells. MHC signaling from APCs to T cells was enhanced, accompanied by increased IFN-γ signaling from NKT and cytotoxic T

644 cells to myeloid and B cell subsets, promoting T cell activation and effector function.

645 Pathways involved in immune recruitment (CCL) and tissue organization (COLLAGEN)

646 were also upregulated, while pro-inflammatory signals such as RESISTIN and IL6 were

647 attenuated. These findings indicate that exercise coordinately enhances antigen

648 presentation and T cell-mediated immunity while reducing chronic inflammation.

Figure 6. Remodeling of immune cell communication networks associated with

regular physical activity.

(A) Bar plot showing the total number of intercellular interactions in SED and EX, with dot

plots indicating the number of interactions mediated by each major immune cell type.

(B) Heatmaps showing the differential number of interactions (left) and differential

interaction strength (right) between major immune cell types (EX vs. SED). Red indicates

increased and blue indicates decreased interactions in EX.

(C) Bar plot of relative information flow for each signaling pathway in SED and EX. Green

and blue bars represent EX and SED, respectively.

(D) Cell-cell communication network plots showing MHC-II signaling among major

immune cell types in SED (top) and EX (bottom). Line thickness represents interaction

strength.

(E) Bubble plot showing communication probabilities of MHC-II ligand–receptor pairs

between B cells and myeloid cells to CD4$^+$ T cells in SED and EX. Dot size represents

communication probability; color indicates statistical significance.

(F) Bubble plot showing communication probabilities of MHC-I ligand–receptor pairs

(HLA-G–CD8A and HLA-G–CD8B) from multiple cell types to CD8$^+$ T cells in SED and EX.

Dot size represents communication probability; color indicates statistical significance.

(G) Cell-cell communication network plots showing IFN-γ signaling among immune cell

subsets in SED (left) and EX (right). Line thickness represents interaction strength.

(H) Bubble plot showing communication probabilities of IFN-γ signaling via

IFNGR1/IFNGR2 from CD4$^+$ CTL and CD8$^+$ CTL to myeloid and B cell subsets in SED

and EX. Dot size represents communication probability; color indicates statistical

significance.

Discussion

Physical inactivity is a risk factor for metabolic disorders, immune dysfunction, and

all-cause mortality(*1*, *73*). In contrast, regular physical activity is a well-established,

effective intervention to mitigate these risks(*74*). However, the underlying cellular and

molecular mechanisms that confer protection in free-living populations remain

incompletely characterized. Our study reveals that regular exercise not only

significantly enhances systemic fatty acid oxidation efficiency and endogenous

antioxidant defense, but also induces widespread epigenetic remodeling at the

single-cell level, selectively optimizing antigen presentation and cytotoxic functions.

Importantly, intercellular communication undergoes a systemic shift. This transition is

characterized by the downregulation of pro-inflammatory mediators, coupled with the

amplification of homeostatic pathways, particularly those related to MHC-mediated

antigen presentation and IFN-γ signaling. Collectively, these findings reveal how regular exercise reshapes the internal milieu across the dimensions of metabolic homeostasis and immune function, providing a novel molecular perspective on its systemic health benefits.

EX exhibit a distinct metabolic signature characterized by elevated FAO intermediates, such as O–Adipoylcarnitine and ketone bodies, alongside a reduction in TAG levels. These findings indicate a significant enhancement in systemic lipid turnover and utilization efficiency. This metabolic phenotype aligns closely with the clinical improvements in lipid profiles (decreased TG and increased HDL-C), collectively forming the physiological basis for exercise-induced cardiometabolic protection(*75*). Furthermore, we observed an increase in endogenous antioxidants, including betaine and 3-hydroxyanthranilic acid, reflecting an augmented buffering capacity against oxidative stress in response to elevated metabolic flux(*36*). Notably, recent evidence has identified betaine as an endogenous TBK1 inhibitor that functions as an "exercise mimetic," exerting anti-inflammatory and pro-longevity effects(*10*). The significantly elevated betaine levels observed in EX not only provide real-world human evidence for this mechanism but also suggest that betaine has the potential to serve as a biomarker of exercise adaptation and a promising therapeutic target for clinical intervention. Several metabolites upregulated in this study have been previously established as potent immunomodulatory signaling molecules or epigenetic regulators. For instance, β-hydroxybutyrate (BHB) functions as an endogenous histone deacetylase (HDAC) inhibitor that suppresses NLRP3 inflammasome activation in macrophages, thereby exerting systemic anti-inflammatory effects(*76*). Furthermore, GABA has been shown to directly regulate lymphocyte proliferation and inflammatory cytokine secretion through functional receptors expressed on the surface of immune cells(*77*). These established links suggest that exercise-induced systemic metabolic remodeling might contribute to the optimization of immune cell functions. Although the direct causal relationship between these metabolites and immune cells was not explicitly tested in this study, these observations provide a robust foundation for hypothesis generation in future

mechanistic research.

Regular exercise not only reshapes systemic metabolic homeostasis but also
orchestrates a functional reinforcement of the professional APCs landscape. Within
the myeloid compartment, monocytes and DCs exhibit elevated expression of
antigen-presentation-related genes. In cMono specifically, we observed enhanced
chromatin accessibility at *HLA-DQB1* and *HLA-DQA1* loci, synchronized with the
increased regulatory activities of master transcription factors governing antigen
presentation and vascular homeostasis, such as *SPI1*, *IRF1*,and *KLF4*. This
enhanced antigen presentation capacity enables more efficient antigen capture and
presentation, facilitating the priming of adaptive immune responses and maintaining
immune cells in a heightened state of alertness. Beyond the myeloid compartment,
such epigenetic remodeling extends to adaptive immune subsets, where naive_B and
switched_Bm cells also demonstrate coordinated increases in *HLA-DQB1* expression
and regulatory element accessibility. Such cross-lineage remodeling aligns with our
cell-cell communication analysis, which reveals a significant amplification of MHC-I
and MHC-II signaling from APCs to T cells in EX. Given that antigen-presentation
efficiency serves as a critical rate-limiting step in adaptive immune responses(*78*, *79*),
the convergent enhancement of antigen presentation capacity across cMono and B
cell subsets, coupled with the amplified APCs-T cell crosstalk, may collectively
underlie the enhanced immune surveillance and reduced susceptibility to infection
documented in physically active populations(*5*, *58*).

In T cell subsets, regular exercise is manifested as a distinct transition in effector
function. For instance, in CD8$^+$ CTLs, increased activities of master transcription
factors driving effector programs, such as *TBX21* and *EOMES*, are observed,
accompanied by the up-regulation of downstream target genes (*GZMB*) and
increased chromatin accessibility at their regulatory elements. These alterations occur
in synchronization with the systemic enhancement of TCR and IFN-γ response
pathways, suggesting that regular exercise may enhance the sensitivity of effector T
cells to activation signals by optimizing the epigenetic landscape. Furthermore, the
increased motif accessibility of factors such as *TCF7* in CD8$^+$ Tn cells suggests that,

while reinforcing the effector reservoir, exercise may also contribute to maintaining or

optimizing the stemness and long-term adaptive potential of the naive T cell pool(*80*,

81). Similar regulatory features, such as the increased openness of *EOMES*/*TBX21*

motifs, are also reflected in mature NK cells.

Cell communication analysis indicates that regular exercise can systematically

remodel the intercellular interaction network, thereby coordinating metabolic

regulation and immune remodeling. This study found that EX simultaneously

enhanced immune surveillance and coordination pathways while suppressing chronic

inflammatory signaling. Specifically, EX upregulated key immune-coordination

pathways such as interferon signaling, antigen presentation (MHC-I/II), and

chemokine activity, which may contribute to improved immune defense and tissue

communication. Concurrently, within this network of cellular interactions, the

pro-inflammatory network centered on resistin signaling and dominated by myeloid

cells was inhibited. Research has established that resistin acts not only as a potent

pro-inflammatory factor, upregulating inflammatory mediators such as tumor necrosis

factor-α (TNF-α) and IL-6(*82*), but also directly targets the vascular wall, inducing

endothelial dysfunction and promoting smooth muscle cell proliferation, thus

accelerating the progression of pathological processes including atherosclerosis and

vascular restenosis(*83*, *84*). Therefore, this study suggests that regular exercise may

provide a protective regulatory foundation for the cardiovascular system at the level of

cellular interactions, potentially through the inhibition of this key pro-inflammatory

network.

In summary, this study employs multi-omics to describe exercise-related changes

in circulating metabolites and lipids, alongside cell-type-resolved transcriptional and

epigenetic landscapes of immune cells. These are characterized by enhanced lipid

oxidation and antioxidant metabolic capacity, alongside functionally optimized

immune states guided by epigenetic reprogramming. We provide multidimensional

map of spontaneous exercise-induced molecular adaptations across metabolic,

epigenetic, and immune dimensions, thereby deepening our understanding of

exercise-mediated health promotion mechanisms and identifying potential biomarkers

and regulatory pathways for chronic disease prevention and management.

Limitations

Several limitations of this study warrant consideration. First, the assessment of physical

activity levels relied on self-reporting; although this is a standard methodology in

large-scale epidemiological research, it remains susceptible to recall and social

desirability biases. Second, the study population consisted primarily of young, healthy

individuals. While this focus facilitates the exclusion of confounding factors and reveals

the core adaptive mechanisms of regular exercise, caution should be exercised when

generalizing these conclusions to elderly, obese, or clinical populations. Third, although

the single-cell multi-omics sample size is substantial relative to comparable studies and all

analyses were statistically adjusted for sex and BMI, the higher proportion of females in

the sedentary group may still introduce residual confounding. Expanding the sample size

in future cohorts will aid in achieving better baseline balance and more granular subgroup

analyses. Nevertheless, our study provides a multi-omics characterization of molecular

adaptations to regular exercise in a natural population cohort, laying a foundation for

future mechanistic and translational studies.

Methods

Data Source

All data for this study were derived from the CIMA project, a publicly available cohort that

enrolled 428 adults with systematic multi-dimensional data collection(*21*). Our analysis

utilized the following data modalities: (1) CIMA sample information metadata and physical

activity questionnaires, (2) clinical blood biochemical markers, (3) targeted plasma

lipidomics and metabolomics profiles, and (4) raw sequencing data matrices from paired

scRNA-seq and scATAC-seq performed on PBMCs.

Assessment of Physical Activity and Grouping

Physical activity was assessed across three dimensions (intensity, frequency, and

duration) using a self-reported questionnaire within the CIMA cohort. Exercise intensity

was determined by the item, "What intensity of exercise do you usually engage in?"

(options: low, moderate, high, or almost none). Frequency (times/week) and session

duration were derived from the items "Exercise habit" and "What is the average daily

exercise duration?", respectively. To obtain a conservative estimate, the lower bounds of

the reported intervals were used for calculation (e.g., "3-4 times/week" was counted as 3;

"1-2 hours" was taken as 60 minutes). Based on the WHO Guidelines on Physical Activity

and Sedentary Behaviour(*23*), participants who engaged in exercise ≥ 3 times per week

and met the recommended exercise volume were classified into the regular exercise

group (EX, n = 40), and those reporting no exercise and a sedentary lifestyle were

classified into the sedentary control group (SED, n = 46).

Differential Analysis of Plasma Metabolites and Lipids

The raw targeted metabolomics and lipidomics data matrices were preprocessed and

normalized using the MetaboAnalyst 6.0 online platform(*85*). Metabolites or lipid species

missing in more than one-third of samples were removed. Remaining missing values were

imputed using the minimum value detected for each molecule. The data were then

subjected to Auto-Scaling (unit variance scaling), i.e., each molecule was mean-centered

and divided by its standard deviation, to eliminate scale differences and give equal

analytical weight to all molecules.To identify molecules associated with exercise habits, a

combined screening strategy was applied. First, linear models were fitted using the limma

package in R, adjusting for sex and BMI, and molecules with $p < 0.05$ and $|\log_2 FC| \geq 0.15$

between exercise groups were selected. Partial least squares discriminant analysis

(PLS-DA) was performed using the MetaboAnalystR package (version 4.0) to calculate

variable importance in projection (VIP) scores(*86*). Ultimately, metabolites and lipids that

satisfied both criteria—VIP scores ≥ 1 and the linear-model thresholds—were defined as

exercise-associated differential molecules (Supplementary Table S2).

Pathway Analysis of Plasma Metabolites

To uncover the potential biological processes underlying the differential metabolites,

Kyoto Encyclopedia of Genes and Genomes (KEGG) pathway enrichment analysis was

conducted for the exercise-associated differential metabolites using the MetaboAnalyst

6.0 platform(*85*). Enrichment significance was evaluated with the hypergeometric test,

and p-values were corrected for multiple testing using the false discovery rate (FDR). To

identify key metabolic alterations, we focused on pathways that passed a significance

threshold of FDR-corrected p-value < 0.05 in our enrichment analysis.

Single-cell RNA-seq data processing

This study used processed single-cell gene expression count matrices. All analyses were

performed using Scanpy (version 1.10.3)(*87*). Quality control retained cells with 500–6000

genes, 1000–25,000 unique molecular identifiers (UMIs), and mitochondrial content < 10%

(Figure S1A). Doublets were identified and removed using Scrublet (expected doublet rate

= 0.06)(*88*). After filtering, data were normalized to 10,000 counts per cell,

log-transformed, and scaled. The top 3,000 highly variable genes were selected for

principal component analysis, and batch effects were corrected using Harmony(*89*).

Unsupervised clustering and subclustering

After batch-effect correction, unsupervised clustering was performed using the Louvain

algorithm(*90*) (resolution = 0.75), and visualized with UMAP. Based on canonical marker

genes, six major cell populations were identified: CD4$^+$ T, CD8$^+$ T, unconventional T, B,

myeloid, and NK cells. Each major population was then extracted and re-clustered

separately with subset-optimized resolutions. Subcluster identities were assigned based

on differential expression and known subtype-specific markers. In total, 34 distinct cell

subpopulations were identified across the dataset.

Differential abundance analysis using scRNA-seq data

To evaluate exercise-associated changes in immune cell composition, differential

abundance analysis was performed using Milo(*32*). A k-nearest neighbor graph (k = 30)

was constructed from the first 50 principal components, and cells were partitioned into

overlapping neighborhoods (makeNhoods, prop = 0.2). Differential abundance testing

between EX and SED was performed using a negative binomial GLM, with TMM

normalization and adjustment for sex and BMI. Significant neighborhoods were defined as

those with FDR < 0.05, and their log_2FC values were used for downstream interpretation.

Differential Expression Analysis of Cell Subpopulations

For the comparison of gene expression differences between the EX and SED groups

across cell subpopulations, we employed the linear model-based limma package for

statistical testing(*91*). For each subpopulation, we constructed a linear model in which

gene expression was the dependent variable, exercise group was the primary predictor,

and sex and BMI were included as covariates. From the fitted model, the log_2FC and raw

p-value were extracted for every gene. To control for false positives, we adopted the

Benjamini-Hochberg (BH) procedure for multiple testing correction of the p-values. In our

analysis, a gene was considered a significant DEG within a given subpopulation if it

showed $|log_2FC| \geq 0.25$ together with an adjusted p-value < 0.05.

Module scoring using scRNA-seq data

To evaluate activity differences in specific biological processes across immune cell

subpopulations, Scoring cells for specific gene modules was based on the Scanpy

sc.tl.score_genes method, which computes the difference between the average

expression of target genes and that of a matched random background per cell. For our

analysis, we sourced gene sets from the Molecular Signatures Database (MSigDB)(*92*).To

assess statistical differences in module scores between the EX and SED groups, we first

computed the mean module score per subpopulation. We then fitted linear models

comparing groups, adjusting for sex and BMI.

Gene regulatory network analysis

Transcriptional regulatory networks in immune cell subsets were inferred using pySCENIC

(v0.12.1)(*93*). GRNBoost was used to identify TF-target gene co-expression modules,

which were then refined against the cisTarget database to retain only genes with relevant

TF-binding motifs(*94*). Regulon activity was quantified by areas under the curve per cell

(AUCell), and differential regulon activity between EX and SED was assessed with limma

(adjusted p < 0.05, |log$_2$FC| ≥ 0.25).

scATAC-seq quality control.

We used ArchR (version 1.0.2) to analyze single-cell ATAC-seq data(*95*). First, Arrow files

were generated for each sample from the raw sequencing fragments following the default

settings. Subsequently, stringent cell-level quality control was applied. Our cell quality

control was based on the parameters below: TSS enrichment ≥ 10, number of unique

fragments between 1×10^3 and 1×10^5, nucleosome signal < 4, and blacklist ratio < 0.05.

Doublets were identified and removed using ArchR's default parameters, resulting in a

high-quality chromatin accessibility matrix for downstream analysis.

scATAC-seq dimensionality reduction and clustering.

To construct a single-cell chromatin accessibility landscape, we first performed

dimensionality reduction on the cell-by-peak matrix using latent semantic indexing (LSI) to

capture dominant patterns of chromatin accessibility variation. To mitigate technical

variability introduced during sample preparation, we applied the Harmony(*89*) algorithm to

integrate the reduced-dimension embeddings across samples.Using the integrated

low-dimensional representation, we partitioned cells into communities via shared nearest

neighbor (SNN)-based graph clustering. An initial clustering step was performed at a

uniform resolution of 0.8 to delineate major cell categories. To further resolve cellular

heterogeneity and enhance the detection of small-scale cell states, we carried out

sub-clustering on selected populations derived from the initial clustering, with resolutions

adapted between 0.3 and 1.5 according to the intrinsic structure of each cell group. As the

final step in our single-cell ATAC-seq analysis, all cells were projected into a

two-dimensional space for visualization using uniform manifold approximation and

projection (UMAP), providing an intuitive representation of cell-state distributions and

community relationships.

scATAC–seq gene activity scores.

Gene activity was quantified with ArchR(v1.0.2), which computes scores by correlating

accessibility across gene-linked regions. Following this, the MAGIC algorithm was applied

to smooth the scores and mitigate noise inherent in sparse scATAC-seq data(*96*).

Integrating scATAC-seq and scRNA-seq Data to Infer Cell Identity

To accurately annotate cell types in the scATAC-seq data, we performed an integrative

analysis against a matched scRNA-seq reference. This was achieved using the

addGeneIntegrationMatrixfunction in ArchR. This process was conducted in two

sequential steps: first, an unconstrained integration was carried out to broadly map each

scATAC-seq cell cluster to the major cell types defined in the scRNA-seq data.

Subsequently, based on this initial mapping, A second, more constrained integration was

then carried out by re-executing addGeneIntegrationMatrix with updated broad group

labels. This refinement enabled finer multimodal alignment and produced final annotations

at high resolution, ensuring consistency with the transcriptomic profiles.To independently

assess the reliability of the annotations, we employed the Gene Activity Score (GAS)

matrix derived from the scATAC-seq data for validation. By comparing the distribution of

annotated cells in the gene-activity space with that of the corresponding scRNA-seq

reference cells in gene-expression space, we further confirmed the accuracy of the

cross-modal annotation.

scATAC–seq pseudobulk replicate generation and peak calling.

To support differential analysis across cell clusters and conditions, we first constructed

pseudobulk samples. This involved running the addGroupCoverages function with tailored

parameters to create non-overlapping replicates for each cell group. These replicates

were then consolidated via addReproduciblePeakSet to generate a consensus peak

matrix. For precise peak identification, we used MACS2(*97*). The resulting unified peak

set provided the foundation for all subsequent differential accessibility testing.

scATAC–seq genomic regions annotation.

For the identified chromatin-accessible regions (peaks), We annotated the called peaks

by applying the annotatePeak function from the ChIPseeker package package with

default parameters(*98*), in order to determine the nearest associated genes and genomic

features for each peak.

scATAC motif enrichment and motif deviation analysis

We carried out transcription factor motif enrichment followed by chromatin accessibility

deviation analysis using the unified pseudobulk peak set. For motif annotation, we utilized

the CIS-BP motif database (via ChromVAR)(*99*), JASPAR2020(*100*), and HOMER(*101*).

Additionally, motif accessibility deviation scores were calculated for each cell using the

chromVAR algorithm implemented in ArchR, enabling systematic assessment of changes

in transcription factor binding activity across different cellular states.

Identification of differential peaks

To detect differentially accessible chromatin regions between groups, we employed two

complementary approaches. First, we used the getMarkerFeatures function with

parameters useMatrix = "PeakMatrix", bias covariates including "TSSEnrichment" and

"log10(nFrags)", and testMethod = "wilcoxon" to identify differential peaks between

groups, applying thresholds of FDR ≤ 0.1 and log_2FC ≥ 0.25. Second, for differential

transcription factor activity analysis, we applied the same function with useMatrix =

"MotifMatrix" and thresholds of FDR ≤ 0.01 and MeanDiff ≥ 0.05.

Gene Ontology Enrichment Analysis of differential peaks

To interpret the potential biological functions of differential chromatin accessible regions,

we performed GO enrichment analysis using the R package rGREAT(*102*). This package

associates genomic regions with nearby genes and evaluates their enrichment

significance across Biological Process, Molecular Function, and Cellular Component

terms using a hypergeometric test. The analysis was conducted with the human genome

as background, applying default parameters, We selected terms passing the significance

threshold of an FDR-adjusted p-value < 0.05 as significantly enriched.

scATAC–seq TF Foot-print analysis

We performed transcription factor footprinting analysis based on Tn5 transposase

cleavage profiles across genomic motif regions. First, the sequence bias of Tn5 was

subtracted from the raw cleavage signal to obtain a corrected footprint profile. Motif

positions were determined using a combination of the CIS-BP database(*103*) (from

chromVAR human_pwms_v1), and the JASPAR2020 database. Subsequently, footprint

signals were normalized using the average insertion density within ± 200–250⌷bp regions

centered on each motif. The normalized results were aggregated by pseudo-replicate, and

their means and standard deviations were calculated and visualized. Finally, to assess

differences in footprint signals between groups, We chose the nonparametric Wilcoxon

rank-sum test for comparisons between two groups, setting the significance level at $p <$

0.05.

scATAC-seq Peak to gene linkage analysis

To identify peak-to-gene links, we performed prediction analysis using the

addPeak2GeneLinks function in ArchR. The analysis was based on the batch-corrected

low-dimensional embedding (specified via the reducedDims parameter) and computed the

correlation between chromatin accessibility and gene expression. A correlation threshold

of corCutOff = 0.4 was applied to screen for high-confidence linkages. The resulting

significant associations were output as a GRanges object for subsequent visualization

and functional interpretation.

1007

Cell-cell interaction analysis

We performed a systematic analysis of potential signaling communication among immune

cell subpopulations using the CellChat R package(*67*). Communication probabilities

between cell types were computed using default parameters. Signaling pathways were

summarized and the global communication network was constructed using the standard

CellChat workflow. To compare communication patterns between EX and SED, differential

interaction strength was assessed with the rankNet function.

1015

Statistical Analysis

Detailed statistical methods for each analysis are provided in the corresponding Method

subsections and figure legends. All statistical analyses were performed in R (v4.1.3) and

Python (v3.9). Unless otherwise specified, statistical significance was defined as FDR <

0.05. In the figures, significance levels are denoted as *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$,

****$p < 0.0001$.

1022

1023 **Acknowledgements**

1024 We thank all of our team members.

1025

1026 **Funding**

1027 Funding for this research was provided by the National Key Research and Development
1028 Program of China (2025YFC3409300), the National Science and Technology Major
1029 Project (2024ZD0530500), and the GuangDong Basic and Applied Basic Research
1030 Foundation (2024B1515230003, 2026A1515012050).

1031

1032 **Author contributions**

1033 J.L., X.S., S.G., J.Y., P.Y., and C.L. conceived the idea; C.L., P.Y., J.Y., Y.Z., L.L., P.G., and

1034 J.Z. supervised the work; X.S., J.L., S.G., S.X., Y.W., Y.Z., and W.Z. analyzed the data and

1035 performed visualization; X.S., J.L., S.G., P.Y., J.Y., Z.C., and C.L. wrote the manuscript.

1036

1037 **Competing interests**

1038 The authors declare no competing interests.

1039

1040 **Data availability**

1041 The scRNA-seq and scATAC-seq data, plasma metabolomics and lipidomics data, and

1042 blood biochemistry data presented in this study were sourced from the CIMA cohort,

1043 hosted on the TrueBlood database within the CNGBdb portal

1044 (https://db.cngb.org/trueblood/cima/).

1045

1046 **Code availability**

1047 The code can be obtained from the corresponding author upon reasonable request.

1048   Information on publicly available software used is provided in the Methods.

1049

Reference

1.   L. Im, S. Ej, L. F, P. P, B. Sn, K. Pt, Effect of physical inactivity on major

non-communicable diseases worldwide: an analysis of burden of disease and life

expectancy. *Lancet (London, England)* **380** (2012).

2.   S. C. Moore, I.-M. Lee, E. Weiderpass, P. T. Campbell, J. N. Sampson, C. M.

Kitahara, S. K. Keadle, H. Arem, A. Berrington de Gonzalez, P. Hartge, H.-O.

Adami, C. K. Blair, K. B. Borch, E. Boyd, D. P. Check, A. Fournier, N. D. Freedman,

M. Gunter, M. Johannson, K.-T. Khaw, M. S. Linet, N. Orsini, Y. Park, E. Riboli, K.

Robien, C. Schairer, H. Sesso, M. Spriggs, R. Van Dusen, A. Wolk, C. E. Matthews,

A. V. Patel, Association of Leisure-Time Physical Activity With Risk of 26 Types of

Cancer in 1.44 Million Adults. *JAMA Intern Med* **176**, 816–825 (2016).

3.   Y. Zhang, D. H. Lee, L. F. M. Rezende, N. Keum, E. L. Giovannucci, Consistent

Adherence to Physical Activity Guidelines and Digestive System Cancer Risk and

Mortality. *JAMA Oncol* **11**, 1488–1496 (2025).

4.   K. S. Courneya, J. L. Vardy, C. J. O'Callaghan, S. Gill, C. M. Friedenreich, R. K. S.

Wong, H. M. Dhillon, V. Coyle, N. S. Chua, D. J. Jonker, P. J. Beale, K. Haider, P. A.

Tang, T. Bonaventura, R. Wong, H. J. Lim, M. E. Burge, S. Hubay, M. Sanatani, K.

L. Campbell, F. Z. Arthuso, J. Turner, R. M. Meyer, M. Brundage, P. O'Brien, D. Tu,

C. M. Booth, CHALLENGE Investigators, Structured Exercise after Adjuvant

Chemotherapy for Colon Cancer. *N Engl J Med* **393**, 13–25 (2025).

5.   S. F. M. Chastin, U. Abaraogu, J. G. Bourgois, P. M. Dall, J. Darnborough, E.

Duncan, J. Dumortier, D. J. Pavón, J. McParland, N. J. Roberts, M. Hamer, Effects

of Regular Physical Activity on the Immune System, Vaccination and Risk of

Community-Acquired Infectious Disease in the General Population: Systematic

Review and Meta-Analysis. *Sports Med* **51**, 1673–1686 (2021).

6.   Y. Qiu, B. Fernández-García, H. I. Lehmann, G. Li, G. Kroemer, C. López-Otín, J.

Xiao, Exercise attenuates the hallmarks of aging: Novel perspectives. *J Sport

Health Sci*, 101108 (2025).

7.   L. S. Chow, R. E. Gerszten, J. M. Taylor, B. K. Pedersen, H. van Praag, S. Trappe,

M. A. Febbraio, Z. S. Galis, Y. Gao, J. M. Haus, I. R. Lanza, C. J. Lavie, C.-H. Lee,

A. Lucia, C. Moro, A. Pandey, J. M. Robbins, K. I. Stanford, A. E. Thackray, S.

Villeda, M. J. Watt, A. Xia, J. R. Zierath, B. H. Goodpaster, M. P. Snyder, Exerkines

in health, resilience and disease. *Nat Rev Endocrinol* **18**, 273–289 (2022).

8.   J. Frampton, K. G. Murphy, G. Frost, E. S. Chambers, Short-chain fatty acids as

potential regulators of skeletal muscle metabolism and function. *Nat Metab* **2**,

840–848 (2020).

9.   C. K, W. S, M. Kj, H. D, A. S, T. Ms, M. Aa, W. E, L.-M. B, Q. Jv, C. S, C. Jw, E. M, B.

B, T. S, D. Mg, K. Da, C. H, A. M, B. A, E. B, A. M, K. Y, A. M, P. D, S.-F. R. Sm, Z.

W, A. Ea, M. Sb, C. H, H. F, S. Mp, Molecular Choreography of Acute Exercise. *Cell*

181 (2020).

10.   L. Geng, J. Ping, R. Wu, H. Yan, H. Zhang, Y. Zhuang, T. Ning, J. Wang, C. Liang, J.

Zhang, Q. Chu, J. Zhang, Y. Wen, Y. Jing, S. Sun, Q. Qiao, Q. Zhao, Q. Ji, S. Ma, Y.

Cai, Y. Zheng, Z. Zou, Z. Diao, M. Li, H. Zhang, J. Hu, L. Fu, W. Kang, R. Bai, H.

Zhao, S. Zhang, Y. Ding, J. Lei, W. Wang, Y. Ji, B. Gou, G. Sun, J. Yin, P. Yan, H. Li,

Z. Wang, S. Ma, Z. Liu, H. Shan, Q. Wang, T. Cao, S. Yang, C. Wang, P. Yang, Y.

Fan, Y. Ye, J. Hu, M. Jiang, Y. Wang, K. Liu, Y. Li, Y. Li, J. Li, W. Ci, Z.-B. Jin, X. Fu,

X. Zhang, G. Zhao, J. C. Izpisua Belmonte, S. Wang, M. Song, W. Zhang, J. Qu,

G.-H. Liu, Systematic profiling reveals betaine as an exercise mimetic for

geroprotection. *Cell* **188**, 5403-5425.e33 (2025).

11.   S. Sun, S. Ma, Y. Cai, S. Wang, J. Ren, Y. Yang, J. Ping, X. Wang, Y. Zhang, H. Yan,

W. Li, C. R. Esteban, Y. Yu, F. Liu, J. C. Izpisua Belmonte, W. Zhang, J. Qu, G.-H.

Liu, A single-cell transcriptomic atlas of exercise-induced anti-inflammatory and

geroprotective effects across the body. *Innovation (Camb)* **4**, 100380 (2023).

12.   MoTrPAC Study Group, Lead Analysts, MoTrPAC Study Group, Temporal

dynamics of the multi-omic response to endurance exercise training. *Nature* **629**,

174–183 (2024).

13.   J. A. B. Smith, K. A. Murach, K. A. Dyar, J. R. Zierath, Exercise metabolism and

adaptation in skeletal muscle. *Nat Rev Mol Cell Biol* **24**, 607–632 (2023).

14.   R. J. Simpson, H. Kunz, N. Agha, R. Graff, Exercise and the Regulation of Immune
Functions. *Prog Mol Biol Transl Sci* **135**, 355–380 (2015).

15.   N. Dc, W. Lm, The compelling link between physical activity and the body's defense
system. *Journal of sport and health science* **8** (2019).

16.   D. Na, P. Rd, L. Nr, H. S, Lord Jm, Major features of immunesenescence, including
reduced thymic output, are ameliorated by high levels of physical activity in
adulthood. *Aging cell* **17** (2018).

17.   G. Spielmann, B. K. McFarlin, D. P. O'Connor, P. J. W. Smith, H. Pircher, R. J.
Simpson, Aerobic fitness is associated with lower proportions of senescent blood
T-cells in man. *Brain Behav Immun* **25**, 1521–1529 (2011).

18.   M. Gleeson, N. C. Bishop, D. J. Stensel, M. R. Lindley, S. S. Mastana, M. A. Nimmo,
The anti-inflammatory effects of exercise: mechanisms and implications for the
prevention and treatment of disease. *Nat Rev Immunol* **11**, 607–615 (2011).

19.   K. L. Timmerman, M. G. Flynn, P. M. Coen, M. M. Markofski, B. D. Pence, Exercise
training-induced lowering of inflammatory (CD14+CD16+) monocytes: a role in the
anti-inflammatory influence of exercise? *J Leukoc Biol* **84**, 1271–1278 (2008).

20.   L. G. Minuzzi, H. Batatinha, C. Weyh, V. S. Balasubramanian Lakshmi, C.
Fiuza-Luces, B. G. Gálvez, A. Lucia, A. M. Teixeira, N. Sommer, J. C. Rosa-Neto, F.
S. Lira, K. Krüger, Natural killer cells from endurance-trained older adults show
improved functional and metabolic responses to adrenergic blockade and mTOR
inhibition. *Sci Rep* **15**, 25380 (2025).

21.   J. Yin, Y. Zheng, Z. Huang, W. Zhou, Y. Yuan, P. Cai, Y. Bai, S. Yang, Y. Gao, S.
Duan, Y. Wang, Z. Xu, W. Zhang, X. Zhang, Y. Wei, Y. Huang, Y. Liu, W. Wang, T.
Yang, Z. Zhang, X. Chen, X. Zhang, J. Lv, F. Li, Y. Zhang, G. Zeng, X. Wang, W. Ma,
G. Hou, S. Hao, C. Liu, Y. Lai, P. Liu, B. Wang, Y. Li, W. Zhang, P. Gao, J. Xie, M. A.
Esteban, Y. Gu, X. Liu, J. Ji, T. Qi, B. Liu, H. Wang, Y. Zhao, X. Yang, X. Wang, R.
Chen, J. Yang, Y. Yin, J. Wang, Y. Cao, X. Xu, L. Liu, X. Jin, C. Liu, Chinese
Immune Multi-Omics Atlas. *Science* **391**, eadt3130 (2026).

22.   J. E. Powell, Charting immune variation through genetics and single-cell genomics.
Gigascience **15**, giaf161 (2026).

23.   F. C. Bull, S. S. Al-Ansari, S. Biddle, K. Borodulin, M. P. Buman, G. Cardon, C.

1139   Carty, J.-P. Chaput, S. Chastin, R. Chou, P. C. Dempsey, L. DiPietro, U. Ekelund, J.

1140   Firth, C. M. Friedenreich, L. Garcia, M. Gichu, R. Jago, P. T. Katzmarzyk, E.

1141   Lambert, M. Leitzmann, K. Milton, F. B. Ortega, C. Ranasinghe, E. Stamatakis, A.

1142   Tiedemann, R. P. Troiano, H. P. van der Ploeg, V. Wari, J. F. Willumsen, World

1143   Health Organization 2020 guidelines on physical activity and sedentary behaviour.

1144   *Br J Sports Med* **54**, 1451–1462 (2020).

1145   24.   R. Guthold, G. A. Stevens, L. M. Riley, F. C. Bull, Worldwide trends in insufficient

1146   physical activity from 2001 to 2016: a pooled analysis of 358 population-based

1147   surveys with 1·9 million participants. *Lancet Glob Health* **6**, e1077–e1086 (2018).

1148   25.   T. Strain, S. Flaxman, R. Guthold, E. Semenova, M. Cowan, L. M. Riley, F. C. Bull,

1149   G. A. Stevens, Country Data Author Group, National, regional, and global trends in

1150   insufficient physical activity among adults from 2000 to 2022: a pooled analysis of

1151   507 population-based surveys with 5·7 million participants. *Lancet Glob Health* **12**,

1152   e1232–e1243 (2024).

1153   26.   S. Mann, C. Beedie, A. Jimenez, Differential effects of aerobic exercise, resistance

1154   training and combined exercise modalities on cholesterol and the lipid profile:

1155   review, synthesis and recommendations. *Sports Med* **44**, 211–221 (2014).

1156   27.   M. Aadahl, L. von Huth Smith, C. Pisinger, U. N. Toft, C. Glümer, K. Borch-Johnsen,

1157   T. Jørgensen, Five-year change in physical activity is associated with changes in

1158   cardiovascular disease risk factors: the Inter99 study. *Prev Med* **48**, 326–331

1159   (2009).

1160   28.   K. S, T. S, S. K, S. M, S. Y, O. F, S. E, S. H, Y. S, K. K, O. Y, Y. N, S. H, Effect of

1161   aerobic exercise training on serum levels of high-density lipoprotein cholesterol: a

1162   meta-analysis. *Archives of internal medicine* **167** (2007).

1163   29.   A. C. Baxmann, M. S. Ahmed, N. C. Marques, V. B. Menon, A. B. Pereira, G. M.

1164   Kirsztajn, I. P. Heilberg, Influence of muscle mass and physical activity on serum

1165   and urinary creatinine and serum cystatin C. *Clin J Am Soc Nephrol* **3**, 348–354

1166   (2008).

30.   A. Bn, C. R, S. E, H. P, Uric acid provides an antioxidant defense in humans

against oxidant- and radical-caused aging and cancer: a hypothesis. *Proceedings*

of the National Academy of Sciences of the United States of America **78** (1981).

31.   R. Stocker, Y. Yamamoto, A. F. McDonagh, A. N. Glazer, B. N. Ames, Bilirubin is an

antioxidant of possible physiological importance. *Science* **235**, 1043–1046 (1987).

32.   E. Dann, N. C. Henderson, S. A. Teichmann, M. D. Morgan, J. C. Marioni,

Differential abundance testing on single-cell data using k-nearest neighbor graphs.

Nat Biotechnol **40**, 245–253 (2022).

33.   M. G. Schooneman, F. M. Vaz, S. M. Houten, M. R. Soeters, Acylcarnitines:

reflecting or inflicting insulin resistance? *Diabetes* **62**, 1–8 (2013).

34.   P. Puchalska, P. A. Crawford, Multi-dimensional Roles of Ketone Bodies in Fuel

Metabolism, Signaling, and Therapeutics. *Cell Metab* **25**, 262–284 (2017).

35.   S. M. Houten, R. J. A. Wanders, A general introduction to the biochemistry of

mitochondrial fatty acid β-oxidation. *J Inherit Metab Dis* **33**, 469–477 (2010).

36.   P. Sk, J. Mj, Exercise-induced oxidative stress: cellular mechanisms and impact on

muscle force production. *Physiological reviews* **88** (2008).

37.   S. Christen, S. R. Thomas, B. Garner, R. Stocker, Inhibition by interferon-gamma of

human mononuclear cell-mediated low density lipoprotein oxidation. Participation

of tryptophan metabolism along the kynurenine pathway. *J Clin Invest* **93**,

2149–2158 (1994).

38.   W. E. Gall, K. Beebe, K. A. Lawton, K.-P. Adam, M. W. Mitchell, P. J. Nakhle, J. A.

Ryals, M. V. Milburn, M. Nannipieri, S. Camastra, A. Natali, E. Ferrannini, RISC

Study Group, alpha-hydroxybutyrate is an early biomarker of insulin resistance and

glucose intolerance in a nondiabetic population. *PLoS One* **5**, e10883 (2010).

39.   B. Stockinger, P. Di Meglio, M. Gialitakis, J. H. Duarte, The aryl hydrocarbon

receptor: multitasking in the immune system. *Annu Rev Immunol* **32**, 403–432

(2014).

40.   R. Bhat, R. Axtell, A. Mitra, M. Miranda, C. Lock, R. W. Tsien, L. Steinman,

Inhibitory role for GABA in autoimmune inflammation. *Proc Natl Acad Sci U S A* **107**,

2580–2585 (2010).

41.   M. Wlodarska, C. Luo, R. Kolde, E. d'Hennezel, J. W. Annand, C. E. Heim, P.

1198    Krastel, E. K. Schmitt, A. S. Omar, E. A. Creasey, A. L. Garner, S. Mohammadi, D. J.

1199    O'Connell, S. Abubucker, T. D. Arthur, E. A. Franzosa, C. Huttenhower, L. O.

1200    Murphy, H. J. Haiser, H. Vlamakis, J. A. Porter, R. J. Xavier, Indoleacrylic Acid

1201    Produced by Commensal Peptostreptococcus Species Suppresses Inflammation.

1202    *Cell Host Microbe* **22**, 25-37.e6 (2017).

1203 42. S. Hang, D. Paik, L. Yao, E. Kim, J. Trinath, J. Lu, S. Ha, B. N. Nelson, S. P. Kelly, L.

1204    Wu, Y. Zheng, R. S. Longman, F. Rastinejad, A. S. Devlin, M. R. Krout, M. A.

1205    Fischbach, D. R. Littman, J. R. Huh, Bile acid metabolites control TH17 and Treg

1206    cell differentiation. *Nature* **576**, 143–148 (2019).

1207 43. L. J. C. van Loon, B. H. Goodpaster, Increased intramuscular lipid storage in the

1208    insulin-resistant and endurance-trained state. *Pflugers Arch* **451**, 606–616 (2006).

1209 44. L. A. J. O'Neill, R. J. Kishton, J. Rathmell, A guide to immunometabolism for

1210    immunologists. *Nat Rev Immunol* **16**, 553–565 (2016).

1211 45. F. Geissmann, M. G. Manz, S. Jung, M. H. Sieweke, M. Merad, K. Ley,

1212    Development of monocytes, macrophages, and dendritic cells. *Science* **327**,

1213    656–661 (2010).

1214 46. K. Storm van's Gravesande, M. D. Layne, Q. Ye, L. Le, R. M. Baron, M. A. Perrella,

1215    L. Santambrogio, E. S. Silverman, R. J. Riese, IFN regulatory factor-1 regulates

1216    IFN-gamma-dependent cathepsin S expression. *J Immunol* **168**, 4488–4494

1217    (2002).

1218 47. N. van der Stoep, E. Quinten, M. Marcondes Rezende, P. J. van den Elsen, E47,

1219    IRF-4, and PU.1 synergize to induce B-cell-specific activation of the class II

1220    transactivator promoter III (CIITA-PIII). *Blood* **104**, 2849–2857 (2004).

1221 48. F. Z. Chowdhury, J. D. Farrar, STAT2: A shape-shifting anti-viral super STAT.

1222    *JAKSTAT* **2**, e23633 (2013).

1223 49. H. Das, A. Kumar, Z. Lin, W. D. Patino, P. M. Hwang, M. W. Feinberg, P. K.

1224    Majumder, M. K. Jain, Kruppel-like factor 2 (KLF2) regulates proinflammatory

1225    activation of monocytes. *Proc Natl Acad Sci U S A* **103**, 6653–6658 (2006).

1226 50. X. Liao, N. Sharma, F. Kapadia, G. Zhou, Y. Lu, H. Hong, K. Paruchuri, G. H.

1227    Mahabeleshwar, E. Dalmas, N. Venteclef, C. A. Flask, J. Kim, B. W. Doreian, K. Q.

Lu, K. H. Kaestner, A. Hamik, K. Clément, M. K. Jain, Krüppel-like factor 4

regulates macrophage polarization. *J Clin Invest* **121**, 2736–2749 (2011).

51. A. S, G.-B. Cb, M. T, H.-T. Va, I. H, H. G, R. F, M. Jc, G. P, A. J,   van den O. J, A.

Zk, W. J, A. S, SCENIC: single-cell regulatory network inference and clustering.

Nature methods **14** (2017).

52. J. Romeo, J. Wärnberg, T. Pozo, A. Marcos, Physical activity, immunity and

infection. *Proc Nutr Soc* **69**, 390–399 (2010).

53. A. M. Intlekofer, N. Takemoto, E. J. Wherry, S. A. Longworth, J. T. Northrup, V. R.

Palanivel, A. C. Mullen, C. R. Gasink, S. M. Kaech, J. D. Miller, L. Gapin, K. Ryan,

A. P. Russ, T. Lindsten, J. S. Orange, A. W. Goldrath, R. Ahmed, S. L. Reiner,

Effector and memory CD8+ T cell fate coupled by T-bet and eomesodermin. *Nat*

Immunol **6**, 1236–1244 (2005).

54. E. L. Pearce, A. C. Mullen, G. A. Martins, C. M. Krawczyk, A. S. Hutchins, V. P.

Zediak, M. Banica, C. B. DiCioccio, D. A. Gross, C.-A. Mao, H. Shen, N. Cereb, S.

Y. Yang, T. Lindsten, J. Rossant, C. A. Hunter, S. L. Reiner, Control of effector

CD8+ T cell function by the transcription factor Eomesodermin. *Science* **302**,

1041–1043 (2003).

55. E. Vivier, E. Tomasello, M. Baratin, T. Walzer, S. Ugolini, Functions of natural killer

cells. *Nat Immunol* **9**, 503–510 (2008).

56. K. Ramirez, K. J. Chandler, C. Spaulding, S. Zandi, M. Sigvardsson, B. J. Graves,

B. L. Kee, Gene deregulation and chronic activation in natural killer cells deficient in

the transcription factor ETS1. *Immunity* **36**, 921–932 (2012).

57. W. Wojakowski, U. Landmesser, R. Bachowski, T. Jadczyk, M. Tendera,

Mobilization of stem and progenitor cells in cardiovascular diseases. *Leukemia* **26**,

23–33 (2012).

58. P. Ar, F. S. Ma, E. Km, The effects of exercise on vaccination responses: a review

of chronic and acute exercise interventions in humans. *Brain, behavior, and*

immunity **39** (2014).

59. J. P. Campbell, J. E. Turner, Debunking the Myth of Exercise-Induced Immune

Suppression: Redefining the Impact of Exercise on Immunological Health Across

the Lifespan. *Front Immunol* **9**, 648 (2018).

60.   P. Liu, J. R. Keller, M. Ortiz, L. Tessarollo, R. A. Rachel, T. Nakamura, N. A. Jenkins,

N. G. Copeland, Bcl11a is essential for normal lymphoid development. *Nat*

Immunol **4**, 525–532 (2003).

61.   W. Seo, T. Ikawa, H. Kawamoto, I. Taniuchi, Runx1-Cbfβ facilitates early B

lymphocyte development by regulating expression of Ebf1. *J Exp Med* **209**,

1255–1262 (2012).

62.   R. P. DeKoter, H. Singh, Regulation of B lymphocyte and macrophage

development by graded expression of PU.1. *Science* **288**, 1439–1441 (2000).

63.   L. H, G. R, Failure of B-cell differentiation in mice lacking the transcription factor

EBF. *Nature* **376** (1995).

64.   S. Gerondakis, U. Siebenlist, Roles of the NF-kappaB pathway in lymphocyte

development and function. *Cold Spring Harb Perspect Biol* **2**, a000182 (2010).

65.   S. R, S. Al, S. Jh, Z. H, S. Lm, S. H, Graded expression of interferon regulatory

factor-4 coordinates isotype switching with plasma cell differentiation. *Immunity* **25**

(2006).

66.   S. Hong, Z. Zhang, H. Liu, M. Tian, X. Zhu, Z. Zhang, W. Wang, X. Zhou, F. Zhang,

Q. Ge, B. Zhu, H. Tang, Z. Hua, B. Hou, B Cells Are the Dominant

Antigen-Presenting Cells that Activate Naive CD4+ T Cells upon Immunization with

a Virus-Derived Nanoparticle Antigen. *Immunity* **49**, 695-708.e4 (2018).

67.   S. Jin, C. F. Guerrero-Juarez, L. Zhang, I. Chang, R. Ramos, C.-H. Kuan, P. Myung,

M. V. Plikus, Q. Nie, Inference and analysis of cell-cell communication using

CellChat. *Nat Commun* **12**, 1088 (2021).

68.   F. M, H. M, G. S, S. L, The role of resistin as a regulator of inflammation:

Implications for various human pathologies. *Clinical immunology (Orlando, Fla.)*

133 (2009).

69.   K. Schroder, P. J. Hertzog, T. Ravasi, D. A. Hume, Interferon-gamma: an overview

of signals, mechanisms and functions. *J Leukoc Biol* **75**, 163–189 (2004).

70.   S. Jr, W. Cb, Regulation of interferon-gamma during innate and adaptive immune

responses. *Advances in immunology* **96** (2007).

71. K. R. Legate, S. A. Wickström, R. Fässler, Genetic and cell biological analysis of
integrin outside-in signaling. *Genes Dev* **23**, 397–418 (2009).

72. J. W. Griffith, C. L. Sokol, A. D. Luster, Chemokines and chemokine receptors:
positioning cells for host defense and immunity. *Annu Rev Immunol* **32**, 659–702
(2014).

73. F. W. Booth, C. K. Roberts, J. P. Thyfault, G. N. Ruegsegger, R. G. Toedebusch,
Role of Inactivity in Chronic Diseases: Evolutionary Insight and Pathophysiological
Mechanisms. *Physiol Rev* **97**, 1351–1402 (2017).

74. H. Arem, S. C. Moore, A. Patel, P. Hartge, A. Berrington de Gonzalez, K.
Visvanathan, P. T. Campbell, M. Freedman, E. Weiderpass, H. O. Adami, M. S.
Linet, I.-M. Lee, C. E. Matthews, Leisure time physical activity and mortality: a
detailed pooled analysis of the dose-response relationship. *JAMA Intern Med* **175**,
959–967 (2015).

75. B. Egan, J. R. Zierath, Exercise metabolism and the molecular regulation of
skeletal muscle adaptation. *Cell Metab* **17**, 162–184 (2013).

76. Y.-H. Youm, K. Y. Nguyen, R. W. Grant, E. L. Goldberg, M. Bodogai, D. Kim, D.
D'Agostino, N. Planavsky, C. Lupfer, T. D. Kanneganti, S. Kang, T. L. Horvath, T. M.
Fahmy, P. A. Crawford, A. Biragyn, E. Alnemri, V. D. Dixit, The ketone metabolite
β-hydroxybutyrate blocks NLRP3 inflammasome-mediated inflammatory disease.
Nat Med **21**, 263–269 (2015).

77. A. K. Bhandage, Z. Jin, S. V. Korol, Q. Shen, Y. Pei, Q. Deng, D. Espes, P.-O.
Carlsson, M. Kamali-Moghaddam, B. Birnir, GABA Regulates Release of
Inflammatory Cytokines From Peripheral Blood Mononuclear Cells and CD4[+] T
Cells and Is Immunosuppressive in Type 1 Diabetes. *EBioMedicine* **30**, 283–294
(2018).

78. J. Neefjes, M. L. M. Jongsma, P. Paul, O. Bakke, Towards a systems understanding
of MHC class I and MHC class II antigen presentation. *Nat Rev Immunol* **11**,
823–836 (2011).

79. R. Pa, F. K, The ins and outs of MHC class II-mediated antigen processing and
presentation. *Nature reviews. Immunology* **15** (2015).

80.   X. Zhou, S. Yu, D.-M. Zhao, J. T. Harty, V. P. Badovinac, H.-H. Xue, Differentiation

and persistence of memory CD8(+) T cells depend on T cell factor 1. *Immunity* **33**,

229–240 (2010).

81.   Y. Wang, J. Hu, Y. Li, M. Xiao, H. Wang, Q. Tian, Z. Li, J. Tang, L. Hu, Y. Tan, X.

Zhou, R. He, Y. Wu, L. Ye, Z. Yin, Q. Huang, L. Xu, The Transcription Factor TCF1

Preserves the Effector Function of Exhausted CD8 T Cells During Chronic Viral

Infection. *Front Immunol* **10**, 169 (2019).

82.   M. Bokarewa, I. Nagaev, L. Dahlberg, U. Smith, A. Tarkowski, Resistin, an

adipokine with potent proinflammatory properties. *J Immunol* **174**, 5789–5795

(2005).

83.   H. S. Jung, K.-H. Park, Y. M. Cho, S. S. Chung, H. J. Cho, S. Y. Cho, S. J. Kim, S. Y.

Kim, H. K. Lee, K. S. Park, Resistin is secreted from macrophages in atheromas

and promotes atherosclerosis. *Cardiovasc Res* **69**, 76–85 (2006).

84.   P. Calabro, I. Samudio, J. T. Willerson, E. T. H. Yeh, Resistin promotes smooth

muscle cell proliferation through activation of extracellular signal-regulated kinase

1/2 and phosphatidylinositol 3-kinase pathways. *Circulation* **110**, 3335–3340

(2004).

85.   Z. Pang, Y. Lu, G. Zhou, F. Hui, L. Xu, C. Viau, A. F. Spigelman, P. E. MacDonald, D.

S. Wishart, S. Li, J. Xia, MetaboAnalyst 6.0: towards a unified platform for

metabolomics data processing, analysis and interpretation. *Nucleic Acids Res* **52**,

W398–W406 (2024).

86.   Z. Pang, J. Chong, S. Li, J. Xia, MetaboAnalystR 3.0: Toward an Optimized

Workflow for Global Metabolomics. *Metabolites* **10**, 186 (2020).

87.   W. Fa, A. P, T. Fj, SCANPY: large-scale single-cell gene expression data analysis.

Genome biology **19** (2018).

88.   S. L. Wolock, R. Lopez, A. M. Klein, Scrublet: Computational Identification of Cell

Doublets in Single-Cell Transcriptomic Data. *Cell Syst* **8**, 281-291.e9 (2019).

89.   I. Korsunsky, N. Millard, J. Fan, K. Slowikowski, F. Zhang, K. Wei, Y. Baglaenko, M.

Brenner, P.-R. Loh, S. Raychaudhuri, Fast, sensitive and accurate integration of

single-cell data with Harmony. *Nat Methods* **16**, 1289–1296 (2019).

90.   V. D. Blondel, J.-L. Guillaume, R. Lambiotte, E. Lefebvre, Fast unfolding of

1349   communities in large networks. *J. Stat. Mech.* **2008**, P10008 (2008).

1350   91.   M. E. Ritchie, B. Phipson, D. Wu, Y. Hu, C. W. Law, W. Shi, G. K. Smyth, limma

1351   powers differential expression analyses for RNA-sequencing and microarray

1352   studies. *Nucleic Acids Res* **43**, e47 (2015).

1353   92.   A. Subramanian, P. Tamayo, V. K. Mootha, S. Mukherjee, B. L. Ebert, M. A. Gillette,

1354   A. Paulovich, S. L. Pomeroy, T. R. Golub, E. S. Lander, J. P. Mesirov, Gene set

1355   enrichment analysis: a knowledge-based approach for interpreting genome-wide

1356   expression profiles. *Proc Natl Acad Sci U S A* **102**, 15545–15550 (2005).

1357   93.   B. Van de Sande, C. Flerin, K. Davie, M. De Waegeneer, G. Hulselmans, S. Aibar,

1358   R. Seurinck, W. Saelens, R. Cannoodt, Q. Rouchon, T. Verbeiren, D. De Maeyer, J.

1359   Reumers, Y. Saeys, S. Aerts, A scalable SCENIC workflow for single-cell gene

1360   regulatory network analysis. *Nat Protoc* **15**, 2247–2276 (2020).

1361   94.   T. Moerman, S. Aibar Santos, C. Bravo González-Blas, J. Simm, Y. Moreau, J.

1362   Aerts, S. Aerts, GRNBoost2 and Arboreto: efficient and scalable inference of gene

1363   regulatory networks. *Bioinformatics* **35**, 2159–2161 (2019).

1364   95.   J. M. Granja, M. R. Corces, S. E. Pierce, S. T. Bagdatli, H. Choudhry, H. Y. Chang,

1365   W. J. Greenleaf, ArchR is a scalable software package for integrative single-cell

1366   chromatin accessibility analysis. *Nat Genet* **53**, 403–411 (2021).

1367   96.   D. van Dijk, R. Sharma, J. Nainys, K. Yim, P. Kathail, A. J. Carr, C. Burdziak, K. R.

1368   Moon, C. L. Chaffer, D. Pattabiraman, B. Bierie, L. Mazutis, G. Wolf, S.

1369   Krishnaswamy, D. Pe'er, Recovering Gene Interactions from Single-Cell Data

1370   Using Data Diffusion. *Cell* **174**, 716-729.e27 (2018).

1371   97.   Y. Zhang, T. Liu, C. A. Meyer, J. Eeckhoute, D. S. Johnson, B. E. Bernstein, C.

1372   Nusbaum, R. M. Myers, M. Brown, W. Li, X. S. Liu, Model-based analysis of

1373   ChIP-Seq (MACS). *Genome Biol* **9**, R137 (2008).

1374   98.   G. Yu, L.-G. Wang, Q.-Y. He, ChIPseeker: an R/Bioconductor package for ChIP

1375   peak annotation, comparison and visualization. *Bioinformatics* **31**, 2382–2383

1376   (2015).

1377   99.   A. N. Schep, B. Wu, J. D. Buenrostro, W. J. Greenleaf, chromVAR: inferring

transcription-factor-associated accessibility from single-cell epigenomic data. *Nat Methods* **14**, 975–978 (2017).

100. O. Fornes, J. A. Castro-Mondragon, A. Khan, R. van der Lee, X. Zhang, P. A. Richmond, B. P. Modi, S. Correard, M. Gheorghe, D. Baranašić, W. Santana-Garcia, G. Tan, J. Chèneby, B. Ballester, F. Parcy, A. Sandelin, B. Lenhard, W. W. Wasserman, A. Mathelier, JASPAR 2020: update of the open-access database of transcription factor binding profiles. *Nucleic Acids Res* **48**, D87–D92 (2020).

101. S. Heinz, C. Benner, N. Spann, E. Bertolino, Y. C. Lin, P. Laslo, J. X. Cheng, C. Murre, H. Singh, C. K. Glass, Simple combinations of lineage-determining transcription factors prime cis-regulatory elements required for macrophage and B cell identities. *Mol Cell* **38**, 576–589 (2010).

102. Y. Tanigawa, E. S. Dyer, G. Bejerano, WhichTF is functionally important in your open chromatin data? *PLoS Comput Biol* **18**, e1010378 (2022).

103. M. T. Weirauch, A. Yang, M. Albu, A. G. Cote, A. Montenegro-Montero, P. Drewe, H. S. Najafabadi, S. A. Lambert, I. Mann, K. Cook, H. Zheng, A. Goity, H. van Bakel, J.-C. Lozano, M. Galli, M. G. Lewsey, E. Huang, T. Mukherjee, X. Chen, J. S. Reece-Hoyes, S. Govindarajan, G. Shaulsky, A. J. M. Walhout, F.-Y. Bouget, G. Ratsch, L. F. Larrondo, J. R. Ecker, T. R. Hughes, Determination and inference of eukaryotic transcription factor sequence specificity. *Cell* **158**, 1431–1443 (2014).