Integrated Multi-omics Analysis Identifies Molecular Signatures of Regular Physical Activity in Human Peripheral Blood

Xuemei Song[1,2]†, Jingzhi Lv[3]†, Siliang Ge[4,5]†, Sihan Xu[6], Yisha Wu[1,7], Yuhui Zheng[1,8], Wenwen Zhou[1,8], Yan Zhang[1], Jiaqiang Zhang[9,10], Pengbin Yin[4,5]*, Jianhua Yin[1,11,12]*, Chuanyu Liu[1,11,12,13]*

[1]State Key Laboratory of Genome and Multi-omics Technologies, BGI Research, Shenzhen 518083, China

[2]College of Life Sciences and Technology, Huazhong Agricultural University, Wuhan 430070, China

[3]State Key Laboratory of Molecular Oncology, Key Laboratory of Carcinogenesis and Translational Research (Ministry of Education/Beijing), Peking University Cancer Hospital & Institute, Beijing 100021, China.

[4]Department of Orthopedics, Chinese PLA General Hospital, Beijing 100853, China.

[5]National Clinical Research Center for Orthopedics, Sports Medicine & Rehabilitation, Beijing 100853, China.

[6]Henan Institute of Medical and Pharmaceutical Sciences & BGI College, Zhengzhou University, Zhengzhou 450052, China

[7]Department of Otolaryngology, Head and Neck Surgery, Second Hospital, Shanxi Medical University, Taiyuan 030001, China

[8]College of Life Sciences, University of Chinese Academy of Sciences, Beijing 100049, China

[9]Department of Anesthesiology and Perioperative Medicine, People's Hospital of Zhengzhou University, Henan Provincial People's Hospital, People's Hospital of Henan Academy of Innovations in Medical Science, Zhengzhou, China

[10]Institute of Electrophysiology, Henan Academy of Innovations in Medical Science, Zhengzhou, China

[11]Shenzhen Proof-of-Concept Center of Digital Cytopathology, BGI Research, Shenzhen, China

[12]Shanxi Medical University-BGI Collaborative Center for Future Medicine, Shanxi

Medical University, Taiyuan 030001, China

[13]Zhangzhou Affiliated Hospital of Fujian Medical University, Zhangzhou, China

*Corresponding author. Email: yinpengbin@gmail.com (Y.P.); yinjianhua@genomics.cn

(Y.J.); liuchuanyu@genomics.cn (C.L.);

†These authors contributed equally to this work.

Abstract

Regular physical activity is well established to protect against metabolic disorders and bolster immunity; yet, the underlying molecular and cellular mechanisms remain incompletely understood. Here, leveraging the Chinese Immune Multi-omics Atlas (CIMA), we integrated plasma metabolomics and lipidomics with single-cell transcriptomic and chromatin accessibility profiles (scRNA-seq and scATAC-seq) to decode the systemic impact of physical activity on human immunity and metabolism. Our data reveal that regular physical activity is linked to a coordinated metabolic signature marked by enhanced fatty acid oxidation and antioxidant defense. In circulating immune cells, it reprograms myeloid chromatin accessibility, activating antigen-presentation pathways while dampening inflammatory programs. Concurrently, effector functional programs in T cells and natural killer cells show extensive enhancement. Moreover, cellular communication analysis highlights a coordinated reinforcement of antigen presentation and chemotaxis signaling. Together, these findings elucidate molecular mechanisms underlying exercise-induced health benefits and offer a theoretical basis for enhancing public health and preventing chronic diseases.

Introduction

Regular physical activity is a powerful, modifiable driver of health, influencing chronic disease risk, cancer outcomes, infection resistance, aging processes, and lifespan[1–6]. Decades of mechanistic research have shown that exercise exerts its effects by inducing broad alterations in molecular signaling across multiple levels of the body, including specific tissues, the whole organism, and circulating body fluids[7–9]. However, most clinical benefits are observed in people who have been naturally and habitually active over many years, whereas much mechanistic evidence arises from short-term, tightly controlled interventions in selective populations or model organisms[10–13]. This mismatch not only obscures our understanding of the true molecular and cellular mechanisms that underlie the clinical benefits of exercise, but also creates gaps in external validity, and limits our ability to directly translate mechanical findings into realistic, population-level public health strategies. A key unmet need is comprehensive, multi-level comparison of habitually active and sedentary individuals, using deep cellular and molecular profiling to better understand how real-world regular exercise shape biology and health.

Notably, metabolism and immunity are widely regarded as two central pillars through which exercise exerts its health-promoting effects[14,15]. Metabolite and lipid profiling studies have identified candidate "exercise mimetics" or exercise-responsive metabolites, such as betaine[12], 12,13-dihydroxy-9Z-octadecenoic acid[16] (12,13-diHOME), β-hydroxybutyrate[17,18], and others, In parallel, flow cytometry–based and related immunophenotyping approaches have characterized how exercise influences immune cell populations, both in terms of their abundance, activation state, and effector functions[19–22]. Building on these important advances, there is now a growing need to move beyond isolated, bulk-level measurements toward a more integrated, systems-level understanding of exercise-induced remodeling of metabolic and immune networks. In particular, we still lack an integrated view of the heterogeneous responses within and across specific immune cell subsets, their dynamic interactions with one another, and how these processes are orchestrated through epigenetic regulation and metabolic reprogramming in response to exercise. Addressing these questions with coordinated multi-omics

approaches will be essential to construct a more holistic atlas of how exercise reshapes immunometabolic circuits and, ultimately, to translate these insights into more precise and scalable health interventions.

Here, we conducted a comprehensive multi-omics study leveraging data from the Chinese Immune Multi-omics Cohort[23] (CIMA), a large-scale population-based cohort. By integrating plasma targeted metabolomics and lipidomics with single-cell multi-omics (scRNA-seq and scATAC-seq) of peripheral blood mononuclear cells (PBMCs)(Figure 1A), we systematically mapped the molecular correlates of habitual physical activity in naturally active individuals compared with sedentary controls. Using both modality-specific and integrative analyses, we characterized exercise-associated alterations in circulating metabolites and lipids, as well as immune cell type–resolved transcriptional and epigenetic landscapes. Our data reveal that long-term regular exercise is linked to a coordinated metabolic signature marked by enhanced fatty-acid oxidation and antioxidant defenses, together with widespread remodeling of immune cell states mediated by epigenetic reprogramming. Collectively, these results provide, to our knowledge, the first multi-dimensional atlas of molecular adaptations to long-term spontaneous exercise, spanning metabolism, chromatin regulation, and immunity. Beyond deepening mechanistic insight into how physical activity promotes health, this work highlights candidate biomarkers and regulatory pathways that may inform prevention and management of chronic diseases, underscoring the translational relevance of our findings.

Results

Physiological Adaptations Induced by EX

This study is based on a secondary analysis of data from the Chinese Immune Multi-omics Cohort (CIMA). We categorized 428 adults according to the World Health Organization (WHO) physical activity guidelines[24]. The regularly exercising group (EX, n=40) was defined as individuals who engaged in physical activity ≥3 times per week, accumulating ≥150 minutes of moderate-intensity or ≥75 minutes of vigorous-intensity activity. The sedentary control group (SED, n=46) was defined as individuals with no

exercise habits whatsoever whose daily life was predominantly sedentary.

The study population had a mean age of 28.2 years (95% CI: 27.2–29.2) and a mean

body mass index of 21.3 kg/m² (SD ± 3.0) (Table S1). Analysis of baseline characteristics

showed no significant differences between the two groups in age, diet, smoking, alcohol

consumption, recent medication history, or sleep duration. However, the sex distribution

differed markedly: SED was heavily skewed toward females, whereas EX had a relatively

balanced sex ratio. This disparity led to a significantly higher unadjusted mean BMI in EX

compared to SED (Figure S1A). Given that sex and BMI are key covariates for metabolic

and immune phenotypes, all subsequent analyses employed multiple linear regression

models adjusted for sex and BMI to independently assess the effects of exercise.

To validate the physiological relevance of the exercise-based classification, we first

compared routine blood biochemical indices. Of the 19 indices tested, eight differed

significantly between EX and SED (Figure 1B). These differences were specifically

reflected in glucose and lipid metabolism indicators: compared to SED, EX showed

significantly lower fasting blood glucose(GLU) and triglyceride(TG) levels, along with

significantly higher high-density lipoprotein cholesterol(HDL-C) levels (Figure 1C). This

finding is consistent with a substantial body of prior research concluding that exercise

confers clear cardiometabolic benefits[25–27].

Beyond these classic signatures, EX showed significantly elevated levels of

creatinine, uric acid, and bilirubin, all within normal clinical ranges, whereas key markers

of hepatocellular injury and function did not differ between groups (Figures 1B, S1B, S1C).

This profile suggests that the observed elevations reflect exercise-induced physiological

adaptations, including increased muscle mass and metabolic turnover[28], as well as

enhanced systemic antioxidant defense[29,30], rather than underlying organ pathology.

In summary, classification based on exercise habits effectively captured distinct

physiological states. EX demonstrated both a healthier metabolic profile and features

indicative of positive physiological adaptation, providing a phenotypic foundation for

subsequent molecular investigations.

Multi-omics Profiling of Plasma and Circulating Immune Cells in EX

To systematically elucidate the regulatory effects of long-term regular exercise on

systemic metabolism and the molecular characteristics of immune cells, we conducted a

multi-omics analysis on 86 subjects (40 in EX and 46 in SED). This analysis integrated

targeted plasma metabolomics, lipidomics, single-cell RNA sequencing (scRNA-seq), and

single-cell assay for transposase-accessible chromatin sequencing (scATAC-seq) of

peripheral blood mononuclear cells (PBMCs) (Figure 1A). First, to dissect the impact of

exercise on systemic metabolism, we analyzed targeted metabolomics and lipidomics

data from all plasma samples. Following standardized pre-processing, we quantified 321

metabolites and 718 lipid molecules (complete list in Table S2). We observed extensive

differences in metabolite and lipid levels between EX and SED (Figure 1C and Table S2),

indicating that regular exercise significantly remodels the human metabolic network.

Subsequently, to investigate the impact of exercise on the immune system, we

proceeded with single-cell multi-omics analysis. Rigorous quality assessment confirmed

high technical reproducibility and the absence of significant batch effects (Figure S2A–C).

After quality control, 1,317,665 cells from scRNA-seq and 644,045 cells from scATAC-seq

were retained for downstream analysis (Figure 1B, 1C). Based on the scRNA-seq data,

we manually annotated cells using key lineage marker genes, identifying six major

immune cell populations: myeloid cells, B cells, CD8$^+$ T cells, CD4$^+$ T cells, NK cells,

and non-canonical T cells. These were further stratified into 34 distinct cell subsets (Figure

1D, 1E, S2E). For the scATAC-seq data, we employed a mapping annotation strategy

aligned with the scRNA-seq data, ultimately resolving 6 major classes and 30 cell subsets

(Figure 1F). To validate the reliability of these annotations, a marker gene heatmap based

on scATAC-seq gene activity scores demonstrated that chromatin accessibility in each

subset was highly specific and consistent with the transcriptomic data (Figure S2D).

Visualization of chromatin openness at key lineage gene promoter regions confirmed

significant accessibility at expected loci: B cells (MS4A1), T cells (IL7R), CD4$^+$ T cells

(CD4), CD8$^+$ T cells (CD8B), NK cells (KLRB1), and non-canonical T cells (RORC)

(Figure 1G, S2F). These chromatin accessibility patterns in known immune marker gene

promoters further verified the accuracy of our annotation. Furthermore, combining our

scRNA-seq data with MiloR[31] analysis revealed that the proportion of myeloid cells in EX

was significantly higher than in SED (Figure 1H-I), suggesting that exercise habits may

influence the cellular composition of the human immune system.

Plasma Metabolic Changes in EX

To decipher the impact of regular physical activity on systemic metabolic homeostasis, we

performed targeted metabolomics and lipidomics analyses on plasma samples from the

EX and SED groups. A total of 62 differentially abundant molecules were identified (Table

S2, Figure 2A). Notably, the majority of differential metabolites (86%) were upregulated in

EX, whereas the majority of differential lipids (94%) were downregulated, suggesting that

habitual exercise induces a global rewiring of the metabolic network.

Pathway enrichment analysis revealed that the upregulated metabolites were

significantly enriched in core energy metabolism pathways such as the tricarboxylic acid

cycle, as well as amino acid conversion pathways including phenylalanine/tyrosine

metabolism, indicating more active energy metabolism and amino acid turnover in the

exercising population (Figure 2B). Concurrently, pathways related to cofactor metabolism

and glutathione metabolism—a key antioxidant defense system—were also significantly

enriched, underscoring the role of redox homeostasis in exercise adaptation.

At the molecular level, we first focused on fatty acid oxidation (FAO), a central

energy-yielding pathway during exercise (Figure 2C). The significant upregulation of

O-adipoylcarnitine, a crucial carrier for mitochondrial long-chain fatty acid import, directly

reflects an increased flux of fatty acid transport[32]. Concurrently, elevated levels of

3-hydroxybutyrate, a hallmark of ketone body metabolism, indicate activation of hepatic

fatty acid β-oxidation and ketogenesis[33]. Furthermore, accumulation of acetic acid, the

end product of short-chain fatty acid oxidation, further supports a comprehensive

mobilization of FAO pathways across varying fatty acid chain lengths[34]. The coordinated

elevation of these three core metabolites demonstrates that regular exercise

systematically enhances the body's fatty acid oxidative capacity through multiple

complementary mechanisms.

Given that heightened energy metabolism is often accompanied by increased

reactive oxygen species (ROS) production[35], we further examined exercise-induced

modulation of the antioxidant defense system (Figure 2D). In concert with elevated energy

flux, several metabolites involved in redox homeostasis were synchronously upregulated

in EX. Betaine, an important methyl donor, showed significantly higher plasma levels in

exercisers, consistent with its recently identified role in mediating exercise-induced

anti-inflammatory and metabolic benefits[12]. The tryptophan derivative 3-hydroxyanthranilic

acid, which possesses free radical-scavenging activity[36], was also elevated, potentially

contributing to exercise-associated antioxidant adaptation. Moreover, increased levels of

2-hydroxybutyrate, an indirect marker of glutathione synthesis, suggest that exercise may

bolster endogenous antioxidant system function[37]. Notably, other metabolites with

immunomodulatory potential followed a similar trend, including xanthurenic acid (another

tryptophan-kynurenine pathway product) and the neuro-immunomodulator γ-aminobutyric

acid (GABA) (Table S2). These molecules can regulate the function and inflammatory

status of immune cells such as T cells and monocytes/macrophages through the aryl

hydrocarbon receptor or GABAergic signaling pathways[38,39], implying that exercise may

co-ordinate antioxidant and immune homeostasis via metabolic reprogramming.

Furthermore, exercise-associated changes extended to the host–microbiota interface

(Figure 2E). Levels of microbially derived aromatic amino acid metabolites—phenyllactic

acid and indoleacrylic acid—were significantly elevated in EX; the latter has been shown

to enhance intestinal barrier function and modulate innate immunity, thereby influencing

systemic inflammation[40]. In parallel, levels of two secondary bile acids, glycodeoxycholic

acid and glycochenodeoxycholic acid, were reduced. Bile acids are important signaling

molecules that regulate intestinal and hepatic immunity[41], and their altered composition

may influence local and systemic immune activity.

At the lipid level, a broad downregulation was observed in EX (Figure 2F-G).

Triacylglycerols, the primary energy-storage form, exhibited widespread reduction,

consistent with the well-established effect of exercise-promoted lipolysis and fatty acid

utilization[42]. Multiple membrane phospholipid subclasses (e.g., LPE, PC, PG, PI) also

showed decreased abundance, reflecting extensive remodeling of the lipid metabolic

network. However, finer-grained comparison revealed molecule-specific regulation; for

instance, distinct species within the same phosphatidylethanolamine subclass displayed

opposite abundance changes (Figure 2H). Overall, these lipid alterations, together with

enhanced fatty acid oxidative capacity, constitute a characteristic metabolic adaptation to

exercise.

Multi-omic Characterization of Myeloid Cells Heterogeneity in EX

Myeloid cells, especially monocytes and macrophages, are recognized as key cell types

that detect metabolic signals and regulate systemic inflammation and immune

homeostasis[43]. In conditions of obesity and other metabolic disorders, the chronic

low-grade inflammation driven by these cells is a key driver of pathological processes,

including insulin resistance[44]. Classic studies have demonstrated that acute or short-term

exercise can rapidly mobilize neutrophils and monocytes[45]. However, Whether and how

regular physical activity remodels the composition, functional states, and epigenetic

landscape of myeloid cell subsets, thereby modulating systemic immune homeostasis,

remains to be systematically elucidated.We first performed fine-grained clustering of

myeloid cells. Based on classical lineage marker genes, cells were manually annotated

into five functional subsets: classical monocytes (cMono), intermediate monocytes

(intMono), non-classical monocytes (ncMono), classical dendritic cells (cDC), and

plasmacytoid dendritic cells (pDC) (Figure 3A, S3A, S3F). Although conventional

abundance analysis revealed no significant inter-group differences at the population level

(Figure S4A), differential abundance testing using the MiloR algorithm at the local

neighborhood level revealed significant increases in intMono, cMono, and ncMono in EX,

with median $\log_2$ fold changes of +0.37, +0.31, and +0.25, respectively (Figure 3B). This

suggests that long-term regular physical activity specifically remodels peripheral myeloid

cells, with a pronounced effect on the composition of monocyte subsets.

To uncover the functional reprogramming underlying these compositional changes,

we performed differential expression gene (DEG) analysis (Table S3) and pathway

enrichment analysis across subsets. Analysis revealed that in cMono, ncMono, cDC, and

pDC from the EX, upregulated pathways were consistently enriched in antigen processing

and presentation, cell adhesion, and chemokine signaling, whereas downregulated

pathways were concentrated in oxidative stress and hypoxia responses (Figure 3C).

Pathway scoring analysis further confirmed enhanced antigen presentation-related

signals and a general attenuation of stress-related pathways across multiple subsets in

EX (Figure S4B, S4C). At the molecular level, the expression of core antigen presentation

genes (e.g., MHC class II molecules) was elevated, alongside increased expression of

migration-associated receptors such as *CX3CR1* in cDC and *CXCR3* in pDC (Figure 3D),

suggesting a concurrent enhancement of myeloid cell immune surveillance and

tissue-homing potential.

To investigate the epigenetic mechanisms underpinning these functional changes, we

performed an in-depth analysis of chromatin accessibility using scATAC-seq. Analysis of

differentially accessible regions (DARs) between the two groups (Table S4) revealed that

94.3% of DARs across all myeloid subsets were specifically enriched in cMono (Figure

S4D); therefore, subsequent investigations focused on this subset. Gene Ontology (GO)

enrichment analysis of DARs in cMono showed that regions with increased accessibility

were enriched for pathways such as antigen presentation and T cell activation, while

regions with decreased accessibility were associated with stress response pathways

(Figure 3E). Transcription factor binding motif enrichment analysis identified significant

enrichment for motifs of KLF2, KLF4, SP1, SPI1 (PU.1), IRF1, and STAT2 (Figure 3F).

Further motif deviation analysis confirmed that the accessibility of binding sites for these

transcription factors was significantly upregulated in EX, whereas accessibility for motifs

of AP-1 family members (e.g., JUN, FOS) and BACH1 was downregulated (Figure 3G).

Notably, IRF1 regulates the expression of CTSS, which participates in MHC class II

antigen presentation[46]. SPI1, a master regulator of myeloid differentiation, can synergize

with IRF4 to drive CIITA expression, thereby activating the antigen presentation pathway[47].

Upregulated STAT2 activity suggests cells may be in an "anti-viral pre-activated" state.

KLF2 and KLF4, important regulators of immune homeostasis highly expressed in

monocytes and macrophages, can inhibit pro-inflammatory activation and promote cellular

residency and migration[48,49]. Conversely, transcription factors with significantly

downregulated binding site accessibility, such as AP-1 and BACH1, are typically

associated with acute/chronic stress, pro-inflammatory responses, and oxidative stress

responses[50,51].

To elucidate how exercise enhances antigen presentation function at the epigenetic

level, we focused on genes linked to differentially accessible peaks. The results showed

that the promoter regions of multiple key antigen presentation genes (e.g., *HLA-DQA1*,

HLA-DQB1) exhibited significantly enhanced chromatin accessibility in EX (Figure 3H),

consistent with their upregulated mRNA expression (Figure 3D). Concurrently, the

chromatin accessibility of the promoter region of the pro-inflammatory gene *CXCL8* was

significantly reduced in EX (Figure 3I), with a corresponding downregulation in its mRNA

expression (Figure 3J). Furthermore, other inflammation-related genes such as *CCL2* and

CCL7 showed a consistent trend of decreased chromatin accessibility (Figure S4E),

further indicating that physical activity may suppress the inflammatory propensity of

myeloid cells through epigenetic regulation.To validate these findings at the single-cell

transcriptomic level, we assessed transcription factor activity in cMono cells using the

SCENIC algorithm. The results demonstrated that the activities of transcription factors

such as SPI1, STAT1, IRF1, and KLF4 were significantly higher in EX than in SED (Figure

3J), highly consistent with the scATAC-seq data indicating upregulated accessibility of the

corresponding motifs (Figure 3F and 3G). To deeply analyze how these key transcription

factors synergistically regulate the functional state of cMono cells, we constructed their

core regulatory network (Figure 3K). Network analysis revealed that SPI1, IRF1, KLF4,

and STAT1 form a synergistic regulatory module, targeting immune-related genes across

multiple functional modules, including antigen processing and presentation (e.g.,

HLA-DQA1, HLA-DQB1, CD74), inflammation and cytokine signaling (e.g., *IL1B, IL10*),

and chemotaxis and cell migration (e.g., *CCL20, CCL2, ITGB1*).In summary, regularly

physical activity remodels the epigenetic landscape and transcriptional regulatory network

of cMono cells, synergistically enhancing antigen presentation function while inhibiting

inflammatory responses, thereby shaping a "high-function, low-inflammation" functional

phenotype in these cells.

Multi-omic Characterization of T Cell Heterogeneity in EX

Although substantial evidence suggests that regular exercise remodels T cell composition

and enhances their function[52], how exercise drives these functional changes via

epigenetic reprogramming at the single-cell level remains to be systematically elucidated.

Using classical marker genes, we identified 17 T cell subsets, including ten CD8$^+$ T cell

subsets and several non-canonical T cell subsets: CD8$^+$ naive T cells (CD8_Tn), central

memory T cells (CD8_Tcm), effector memory T cells (CD8_Tem), cytotoxic T cells

(CD8_CTL), MAIT cells, γδT1 cells, γδT2 cells, proliferating T cells, and pre-T-like cells

(Figure 4A, S3B, and S3G). Additionally, we identified seven CD4$^+$ T cell subsets (Figure

S5A, S3C, and S3H). MiloR differential abundance analysis revealed that the proportion

of CD4_CTL cells was significantly lower in the exercise group (median log$_2$FC = -0.706)

(Figure S5B), while the proportion of pre-T-like cells was significantly increased (median

log$_2$FC = +0.315) (Figure 4B).

To dissect the impact of exercise on the functional state of T cells from an epigenetic

perspective, we performed DAR analysis across all subsets (Table S4). Quantitative

analysis of upregulated DARs indicated that the most significant changes occurred in

CD8_CTL, MAIT, CD8_Tn, NKT, CD8_Tem, and γδT2 cells. Notably, the number of

upregulated DARs in CD8_CTLs was significantly higher than in other subsets (Figure

4C), with multiple DARs shared between two or more subsets. GO enrichment analysis

showed that genes associated with these upregulated DARs were significantly enriched in

pathways related to CD8$^+$ T cell effector functions, such as cell killing and defense

against infection (Figure 4D). Feature gene set scoring of the transcriptomic data further

supported these findings. In the exercise group, scores for gene modules related to

"cytotoxicity," "type II interferon response," and "oxidative phosphorylation" were

significantly upregulated across multiple T cell subsets (Figure 4E and S5C). Conversely,

scores for hypoxia response and inflammatory response gene modules were markedly

downregulated (Figure S5D). This suggests that regular exercise may not only enhance

the effector functions and metabolic adaptations of CD8$^+$ T cells but also attenuate their

inflammatory signaling.

Transcription factors (TFs) serve as core hubs regulating immune cell effector

functions. To resolve the transcriptional regulatory basis of these functional changes, we

performed motif bias analysis based on scATAC-seq data. The results showed that in

CD8_CTLs, which possessed the highest number of upregulated DARs, the accessibility

of binding motifs for TFs closely related to cytotoxic differentiation and effector

function—including EOMES, RUNX3, RUNX1, and TBX21—was significantly upregulated.

Among these, TBX21 (T-bet) and EOMES are master regulators driving CD8[+] T cell

differentiation into effector cells, and their upregulation is directly linked to enhanced

cytotoxic function[53,54]. Similarly, in MAIT cells, motif accessibility for key TFs involved in

activation, cytokine response, and functional regulation (e.g., RUNX3, RORC, STAT1, and

STAT4) was also significantly upregulated (Figure 4F and S5E). STAT1 and STAT4 are

core mediators of the interferon-γ signaling pathway[55], their upregulated activity

corresponds with the enhanced IFN-γ response pathway described above, providing a

transcriptional mechanism for the improved effector function of MAIT cells. Furthermore,

SCENIC analysis based on scRNA-seq data demonstrated that the regulatory activities of

key TFs such as RUNX3 and TBX21 were significantly elevated in CD8_CTL cells in the

exercise group (Figure 4G), consistent with the epigenetic findings. To further unveil the

downstream regulatory networks of these key TFs, we constructed a target gene

regulatory network for RUNX3 (Figure 4H). The analysis indicated that RUNX3 targets

and regulates genes such as GZMB and TBX21, which collectively drive the

enhancement of CD8_CTL effector functions. Further analysis revealed that in CD8_CTL

cells, the chromatin accessibility of distal regulatory regions for the *TBX21* and *GZMB* loci

was significantly enhanced in the exercise group (Figure 4I); in MAIT cells, the promoter

region of the core effector gene *PRF1* exhibited a similar trend of increased accessibility

(Figure 4J). Collectively, both single-cell transcriptomic and ATAC-seq results suggest that

regular exercise enhances T cell effector functions.

Multi-omic Characterization of NK and B Cell Heterogeneity in EX

Extensive research demonstrates that regular aerobic exercise effectively bolsters

immune function and defense against infection[56]. As core effector cells of the innate

immune system, natural killer (NK) cells not only serve as the first line of defense against

viral infections and in tumor immunosurveillance but also play a pivotal role in bridging

innate and adaptive immune responses[57]. However, the specific molecular mechanisms

by which exercise enhances NK cell effector function remain incompletely understood. We

first classified NK cells into six subsets based on classical marker genes: Mature NK cells,

Transitional NKs, type 2 innate lymphoid cells (ILC2), CD56[+]　NK cells, Terminal NK cells,

and Cycling NK cells (Figure 5A, S3D, and S3I). Notably, the proportion of CD56[+]　NK

cells was significantly reduced in the exercise group (Figure S6A), a finding consistent

with MiloR algorithm results (Figure 5B). To evaluate functional changes in NK cells, we

identified differentially expressed genes (DEGs) (Table S3) and performed pathway

scoring analysis (Figure 5C and Figure S6B). The results showed elevated scores for

killing function-related pathways across multiple NK cell subsets in the exercise group,

while scores for inflammation, oxidative stress, and apoptosis signaling pathways were

generally reduced.Epigenetically, we identified differentially accessible regions (DARs)

(Table S4), with Mature NK cells exhibiting the highest number of upregulated DARs

(Figure S6C). Gene Ontology (GO) enrichment analysis of these upregulated DARs

further revealed that pathways related to "immune defense and killing," "immune

regulation," and "immune cell mobilization" were significantly enriched in Mature NK cells,

whereas inflammation and stress-related pathways were downregulated (Figure S6D),

consistent with the transcriptomic findings. To deeply investigate the transcriptional

regulatory basis, we performed motif bias analysis based on scATAC-seq data (Figure

S6E and 5E). Results showed that in the exercise group, motif accessibility was

significantly enhanced for TFs closely related to NK cell effector function, including key

regulators driving cytotoxic differentiation such as RUNX3, EOMES, and TBX21[58] (T-bet),

as well as ETS1, which is involved in innate lymphoid cell development and activation[59].

Conversely, motif accessibility for inflammation and stress-related TFs (RELA, RELB,

NFKB1, JUND, FOS, FOSL2, BACH1)[60] was generally reduced. Footprint analysis

confirmed the decreased accessibility of RELA and RELB transcription factor binding sites

(Figure 5F).

To investigate whether common regulatory mechanisms exist for the cytotoxic

functions of adaptive and innate immunity under exercise, we compared transcription

factor activities between CD8_CTL (adaptive immunity) and Mature_NK (innate immunity)

cells (Figure 5G). Venn diagram analysis revealed that both shared upregulation of a

series of core transcription factors, including RUNX3, EOMES, TBX21, ETS2, and

FOXO3, with FOXO3 involved in maintaining T cell memory and NK cell homeostasis[61].

This provides a unified transcriptional regulatory basis for the synergistic enhancement of

cytotoxic functions across different immune branches by exercise. Concurrently, each

population displayed distinct activation patterns: CD8_CTLs specifically upregulated

RUNX1, STAT6, and BCL11B, which are associated with T cell lineage commitment[62–64];

whereas Mature_NK cells uniquely activated factors such as ETS1, IRF1, STAT1, and

STAT2, which are closely related to interferon signaling responses and rapid NK effector

functions[65–67]. Finally, through track analysis of specific loci, we provided direct evidence

for the epigenetic regulation of key genes. In Mature_NK cells, chromatin accessibility

was significantly enhanced in the exercise group at the intronic regulatory region of *STAT1*,

a core mediator of type I interferon signaling, and the promoter region of *CXCR4*, a known

exercise-responsive factor[68] (Figure 5H). Furthermore, accessibility was significantly

reduced at the promoter regions of *RELB*, associated with non-canonical NF-κB pathway

activation, and *CCL5*, an important inflammatory chemokine[69] (Figure 5J). In summary,

natural killer cells in individuals who exercise regularly exhibit higher effector potential and

lower inflammatory characteristics.

We also investigated changes in B cells in the regular exercise population, given their

crucial role in humoral immunity. B cells were annotated into eight subtypes (Figure S7A,

S3E, and S3J): Switched memory B cells (Switched_Bm), Naive B cells, pre-Switched

memory B cells, Unswitched memory B cells (Unswitched_Bm), Atypical memory B cells

(Atypical_Bm), Plasma cells, and Plasmablast cells. No significant differences in cell

proportions were observed across all subsets (Figure S7B). We identified DEGs (Table S3)

and performed enrichment analysis, which showed upregulation of pathways related to

antigen processing and presentation, humoral immune response, and chemotaxis across

multiple B cell subsets (Figure S7C). Notably, the expression of key genes encoding MHC

Class II molecules was consistently upregulated in multiple B cell subsets, alongside

increased expression of inflammation-regulatory genes and certain chemokine genes

(Figure S7D). To resolve the epigenetic regulatory basis, we performed differential

accessibility region analysis (Table S4). Results showed that changes in chromatin

accessibility were highly concentrated in Naive B cells and Switched Bm cells (Figure

S7E). GO enrichment analysis of upregulated DARs indicated that Naive B cells

upregulated pathways such as T cell costimulation, antigen presentation, and apoptosis

inhibition, while Switched_Bm cells upregulated pathways including lymphocyte activation

and antigen presentation (Figure S7F).

To further determine the molecular mechanisms behind these B cell functional

changes, we identified upregulated TFs in these two subsets. Naive B cells showed

upregulation of BCL11A, RUNX1, SPI1 (PU.1), and EBF1 (Figure S7G), with SPI1 and

EBF1 being master regulators of early B cell development and identity[70,71]. Switched Bm

cells showed upregulation of NF-κB family members, IRF4, as well as SPI1 and EBF1

(Figure S7H); NF-κB and IRF4 are key drivers of memory B cell formation, class switching,

and plasma cell differentiation[72,73]. We focused our analysis on the regulatory regions of

genes upregulated in the transcriptome. The results showed that in the exercise group,

chromatin accessibility at the promoter region of the key antigen presentation gene

HLA-DQB1 was significantly enhanced in both Naive B cells (Figure S7I) and

class-switched memory B cells (Figure S7J). Additionally, in Switched Bm, promoter

accessibility of the chemokine gene *CCL5* also showed a synchronous increase (Figure

S7K). Overall, this study reveals that while regular exercise does not significantly affect B

cell subset proportions, it may enhance their potential for antigen presentation and

immune activation through chromatin reprogramming and transcriptional changes in key

subsets.

Altered Intercellular Communication

Immune cells rely on ligand-receptor interactions for communication to coordinate

immune responses[74]. However, the impact of exercise on this interaction network remains

unclear. To address this, we employed the CellChat algorithm[75]. The results indicated a

significant increase in the number of interactions involving CD8⁺ T cells in the exercise

group, alongside a general strengthening of interaction intensity among B cells, CD8⁺ T

cells, and myeloid cells (Figure 6A–B). Comparison at the pathway level revealed that

information flow for inflammation and immune suppression-related pathways (THBS,

RESISTIN, IL6) was higher in the sedentary group, whereas signals for chemotaxis

migration, antigen presentation, and cell adhesion pathways (CCL, MHC-II, PECAM1)

were significantly enhanced in the exercise group (Figure 6C).

Human resistin is a key regulator directing inflammatory responses in various

immune cells, including macrophages and peripheral blood mononuclear cells[76]. The cell

communication network showed that myeloid cells were the core signal source for this

pathway, with classical monocytes being the primary senders, and their signal output was

lower in the exercise group (Figure 6D and S8B). The interaction intensity of key

ligand-receptor axes in this pathway, *RETN*–CAP1 and *RETN*–TLR4, was weakened in

the exercise group, with a synchronous downregulation of *RETN* gene expression in

myeloid cells (Figure 6E–F). Given that the pro-inflammatory effects of Resistin rely

partially on NF-κB pathway activation[77], to explore changes in the downstream

inflammatory network, we analyzed pathway activity scores in myeloid cell subsets. The

results showed that in classical monocytes, multiple core inflammation-related pathways,

including TNFA_SIGNALING_VIA_NFKB, INFLAMMATORY_RESPONSE, and

IL6_JAK_STAT3_SIGNALING, were significantly suppressed in the exercise group

(Figure S8A).

Regarding adaptive immune regulation, antigen presentation-related signals were

significantly enhanced in the exercise group. The MHC-II signaling pathway between B

cells/myeloid cells and CD4$^+$ T cells was elevated in the exercise group, involving

multiple ligand-receptor pairs (Figure 6G–H). Subset analysis revealed that interactions

between classical monocytes, naive B cells, and memory B cells with naive and helper

CD4$^+$ T cells were particularly enhanced in the exercise group (Figure S8C).

Furthermore, MHC-I signaling pathways between multiple cell subsets and CD8$^+$ T cells

were also activated in the exercise group, especially the enhancement of the

HLA-G–CD8A/B axis (Figure 6I), with interactions between classical monocytes and naive

CD8$^+$ T cells contributing most significantly (Figure S8D). IFN-γ, a member of the type II

interferons, plays a critical role in host defense and immune surveillance . In EX, IFN-γ

signal connections significantly strengthened with NKT cells acting as the source and

multiple myeloid and B cell subsets as targets (Figure 6J). Concurrently, interactions

mediated by the *IFNG*–(IFNGR1+IFNGR2) axis, with cytotoxic T cells (CD4$^+$ CTL and

CD8⁺ CTL) as sources and myeloid and B cell subsets as targets, also exhibited specific

enhancement (Figure 6K).

Notably, exercise also potentiated signaling networks related to tissue repair and

immune microenvironment regulation. On one hand, the COLLAGEN signaling pathway

was dominated by non-canonical T cells (Unconv_T cells) in the exercise group, with

upregulation of both the interaction of the key ligand-receptor axis *COL6A2*–CD44 and

ligand gene expression (Figure S8E–G). Collagen signaling is known to participate in cell

adhesion, migration, and tissue homeostasis maintenance[78]. On the other hand, the

chemokine CCL signaling pathway was also significantly enhanced in the exercise group,

primarily manifested as increased signaling from multiple immune cell subsets to myeloid

cells (Figure S8H-I). CCL chemokines are known to play a core role in immune cell

recruitment and tissue localization[79]. These results suggest that exercise may enhance

adaptive immunity and tissue repair-related signal transmission while specifically inhibiting

myeloid cell-mediated inflammatory pathway activity by influencing ligand-receptor

communication networks among immune cells.

Discussion

Extensive research indicates that exercise exerts its effects by inducing widespread

alterations in molecular signaling across multiple levels of the body, including specific

tissues, the entire organism, and circulating fluids. Regular physical activity serves as an

effective means to improve metabolic health, delay aging, reduce chronic disease risk,

and enhance immune function[1–6]. However, most mechanistic studies originate from

short-term, rigorous exercise interventions or animal models, potentially differing from the

molecular profiles observed in populations that spontaneously maintain long-term

exercise habits in real-world settings[10–13]. This discrepancy limits our understanding of the

true molecular and cellular mechanisms underlying exercise's clinical benefits and

impedes the effective translation of basic research findings into public health strategies.

Therefore, conducting comprehensive, multi-level comparisons between individuals who

engage in regular physical activity and sedentary populations is crucial for understanding

the molecular mechanisms through which exercise reshapes physiology and health in the

real world. Adopting a multi-omics strategy, this study performed a multidimensional

analysis of systemic metabolism and single-cell immunity, thereby delineating a molecular

panorama of the coordinated adaptations that drive the metabolic and immune systems

under long-term spontaneous exercise.

The study first analyzed targeted metabolomics and lipidomics data from all plasma

samples, identifying 62 differentially abundant molecules between the two groups,

including metabolites (72.6%) and complex lipids (27.4%). This revealed a remodeling of

metabolic networks in the plasma of long-term regular exercisers. Notably, the

upregulation of fatty acid oxidation (FAO)-related metabolites (e.g., O-hexanoylcarnitine,

3-hydroxybutyrate) corresponded with a widespread decrease in triglycerides (TG). This

confirmed that regular exercise systematically enhances the body's FAO capacity by

boosting fatty acid transport, ketone body production, and short-chain fatty acid utilization.

This finding is highly consistent with previous studies demonstrating exercise-induced

lipolysis and energy metabolism shifts[34]. Simultaneously, we observed synchronous

upregulation of multiple antioxidant defense-related metabolites, including betaine,

3-hydroxy-o-aminobenzoic acid, and 2-hydroxybutyrate. This suggests that while

improving energy metabolism efficiency, exercise may synergistically activate

endogenous antioxidant networks to counteract potential oxidative stress, further

substantiating the hypothesis that exercise enhances the body's antioxidant capacity[35].

Research over the past decade in the field of immunometabolism has established that

circulating metabolites serve as pivotal signaling molecules and epigenetic modulators

that critically regulate immune cell function and differentiation[80]. This conceptual

framework offers a novel perspective for interpreting how long-term exercise-induced

systemic metabolic reprogramming may influence immune homeostasis. In this study, we

found that several significantly upregulated metabolites after long-term exercise possess

well-defined immunomodulatory properties. For instance, β-hydroxybutyrate (BHB) has

been demonstrated in seminal work to function as an endogenous histone deacetylase

(HDAC) inhibitor, which directly suppresses the assembly and activation of the NLRP3

inflammasome in macrophages, thereby exerting a core anti-inflammatory effect[81].

Concurrently, the significantly altered betaine, serving as a crucial methyl donor, can

modulate methylation metabolism within immune cells such as T cells, thereby influencing

their differentiation and functional responses[82]. Furthermore, γ-aminobutyric acid (GABA)

can directly regulate lymphocyte proliferation and inflammatory cytokine secretion through

functional receptors expressed on the surface of immune cells[83].These findings

collectively suggest that long-term exercise shapes a "systemic metabolic

microenvironment" constituted by specific immunomodulatory metabolites, which may

continuously deliver pro-homeostatic signals to the immune system. Based on this, we

propose a central hypothesis: these characteristic metabolites constitute a critical

metabolic interface linking exercise adaptation to immune optimization and are potential

messengers driving the observed beneficial phenotypes, such as the epigenetic

reprogramming of monocytes. Consequently, the central objective of future research is to

establish the causal relationship between these metabolites and specific immune

phenotypes through functional experiments, thereby completely delineating the

physiological map of the "exercise-metabolism-immunity" axis at the mechanistic level.

We then provides deep insights into exercise-induced remodeling of the immune

system through single-cell multi-omics analysis, revealing that regular exercise profoundly

impacts immune function via epigenetic reprogramming. We found that in classical

monocytes (cMono), the exercise group exhibited increased chromatin accessibility in

regions associated with antigen presentation and migration functions, along with

upregulation of transcription factor motif accessibility. Conversely, motif accessibility for

stress and inflammation-related factors such as AP-1 and BACH1 was downregulated.

This provides an epigenetic explanation for why exercise-trained monocytes exhibit

enhanced antigen-presenting potential and reduced inflammatory propensity. Regarding

adaptive immunity, CD8$^+$ cytotoxic T cells (CD8_CTL) and MAIT cells exhibited

significant epigenetic activation. Their gene module scores for cell killing and interferon

responses were upregulated, and the accessibility of binding motifs for TFs closely

associated with cytotoxic differentiation and effector functions—including EOMES,

RUNX3, RUNX1, and TBX21—was markedly increased. Similarly, NK cells (particularly

mature NK subsets) within the innate immune system exhibited analogous pro-effector

epigenetic signatures. These findings suggest that exercise may synergistically enhance

the killing and defense capabilities of effector cells across distinct immune branches while

suppressing their inflammatory states through a conserved set of epigenetic regulatory

mechanisms.

Cell communication analysis further substantiated the relationship between

metabolic regulation and immune remodeling, establishing a systematic view of exercise

adaptation. We found that exercise enhances signaling in immune cell pathways

represented by MHC-I/II and chemokines (e.g., CCL), including chemotaxis, antigen

presentation, and cell adhesion. Simultaneously, it specifically attenuates the

pro-inflammatory network dominated by myeloid cells and centered on retinoic acid-like

protein (RETN) signaling. The reduction in *RETN* levels depends on the inhibition of its

downstream NF-κB pathway activity. This suggests exercise may influence immune

cell-mediated ligand-receptor communication networks, enhancing adaptive immune and

tissue repair signaling while specifically suppressing myeloid cell-mediated inflammatory

pathway activity. This dynamic metabolic-immune cell interaction network provides crucial

clues and foreshadows future investigations into exercise benefit mechanisms.

In summary, this study employs multi-omics integration to describe exercise-related

changes in circulating metabolites and lipids, alongside cell-type-resolved transcriptional

and epigenetic landscapes of immune cells. These are characterized by enhanced lipid

oxidation and antioxidant metabolic capacity, alongside functionally optimized immune

states guided by epigenetic reprogramming. We provide the first multidimensional map of

long-term spontaneous exercise-induced molecular adaptations across metabolic,

epigenetic, and immune dimensions, thereby deepening our understanding of

exercise-mediated health promotion mechanisms and identifying potential biomarkers and

regulatory pathways for chronic disease prevention and management. This study also has

limitations: First, physical activity assessment relied on self-report. While this is a common

and valid approach in large-scale epidemiological research, it remains susceptible to

recall bias and social desirability bias. Second, our cohort consisted of relatively young

and healthy individuals. This facilitates the identification of core exercise adaptation

signatures in a less confounded context but concurrently raises uncertainty regarding the

direct extrapolation of the discovered molecular profiles to older, obese, or clinical

populations. Third, although the sample size was adequate for in-depth single-cell

multi-omics profiling, a sex imbalance across groups was present. While statistical

adjustments were applied, residual confounding effects may persist. Future studies with

expanded cohorts would enhance statistical power for more granular subgroup analyses

(e.g., by sex or exercise modality) and improve the balancing of baseline characteristics.

Furthermore, the study focused exclusively on peripheral blood, leaving the local

immunomodulatory effects of exercise in key tissues such as muscle and adipose tissue

unaddressed—contexts where tissue-specific immune microenvironments may obey

distinct regulatory principles. Finally, and most critically, the absence of functional

experimental validation for predictive networks, such as the "metabolite-driven immune

remodeling" hypothesis, means the underlying mechanistic relationships await

confirmation in subsequent investigations.

Methods

Data Source

We utilized data from the Chinese Immune Multi-omics Atlas (CIMA) project[23]. This public

cohort initially enrolled 428 adults aged 20–77 and systematically collected

multi-dimensional information, including questionnaire surveys, clinical biochemical

indices, and multi-omics data. The present secondary analysis utilized detailed physical

activity questionnaires, blood biochemical markers, targeted lipidomics data, as well as

the raw data matrices generated by paired single-cell RNA and ATAC sequencing

performed on peripheral blood mononuclear cells (PBMCs).

Assessment of Physical Activity, Grouping, and Final Study Population

Within the CIMA cohort, physical activity was assessed using its standardized

questionnairesacross three dimensions: intensity, frequency, and duration. First, exercise

intensity was evaluated with the question "What intensity of exercise do you usually

engage in?" (options: low intensity [e.g., walking], moderate intensity [e.g., brisk walking,

jogging], high intensity [e.g., running, ball sports], or almost none). Second, exercise

frequency and single-session duration were obtained from the items "Exercise habit"

(options: ≥5 times/week to almost none) and "What is the average daily exercise duration?"

(options: >2 hours to <30 minutes and almost none). To provide a conservative estimate,

the lower bounds of the frequency and duration intervals were used for calculation (e.g.,

"3–4 times/week" was counted as 3, "1–2 hours" as 60 minutes). The total weekly

moderate-to-vigorous physical activity (MVPA) duration was then derived for each

participant. Participants were grouped strictly according to World Health Organization

(WHO) recommendations[24]. The regular exercise group (EX) was defined as meeting both

of the following criteria: (1) engaging in moderate- or high-intensity exercise ≥3 times per

week, and (2) achieving a total weekly duration of ≥150 minutes of moderate-intensity

activity or ≥75 minutes of vigorous-intensity activity. The sedentary control group (SED)

was required to satisfy both conditions: (1) not meeting any of the above exercise criteria,

and (2) self-reporting a daily living pattern dominated by sedentary behavior.

To reduce potential confounding from age and chronic disease status and to establish a

relatively homogeneous analytical sample, the following inclusion and exclusion criteria

were applied: Inclusion criteria: age 20–40 years. Exclusion criteria: (1) self-reported

major chronic diseases (e.g., cardiovascular and cerebrovascular diseases, diabetes,

autoimmune diseases, malignancies) or obesity; (2) missing key physical-activity

questionnaire items or omics data. After applying these screening steps, the above

grouping criteria were implemented. Finally, 86 participants were clearly classified and

included in all subsequent analyses, comprising 40 in EX and 46 in SED.

Differential Analysis of Plasma Metabolites and Lipids

The raw targeted metabolomics and lipidomics data matrices were preprocessed and

normalized using the MetaboAnalyst 6.0 online platform[84]. Metabolites or lipid species

missing in more than one-third of samples were removed. Remaining missing values were

imputed using the minimum value detected for each molecule. The imputed data were

then subjected to Auto-Scaling (unit variance scaling), i.e., each molecule was

mean-centered and divided by its standard deviation, to eliminate scale differences and

give equal analytical weight to all molecules.To identify molecules associated with

exercise habits, a combined screening strategy was applied. First, linear models were

fitted using the limma package in R, adjusting for sex and BMI, and molecules with p < 0.05 and |Log2FC| ≥ 0.15 between exercise groups were selected. In parallel, PLS-DA was performed using the MetaboAnalystR package[85] (version 4.0) to calculate variable importance in projection (VIP) scores. Ultimately, metabolites and lipids that satisfied both criteria—VIP scores ≥ 1 and the linear-model thresholds—were defined as exercise-associated differential molecules (complete list provided in Supplementary Table S2).

Pathway Analysis of Plasma Metabolites

To uncover the potential biological processes underlying the differential metabolites, KEGG pathway enrichment analysis was conducted for the exercise-associated differential metabolites using the MetaboAnalyst 6.0 platform[84]. Enrichment significance was evaluated with the hypergeometric test, and p-values were corrected for multiple testing using the false discovery rate (FDR). To identify key metabolic alterations, we focused on pathways that passed a significance threshold of FDR-corrected p-value < 0.05 in our enrichment analysis.

Single-cell RNA-seq data processing

This study is based on processed single-cell gene expression count matrices, with all data preprocessing and analysis performed using the Scanpy toolkit[86] (version 1.10.3). First, stringent quality control was applied: cells were retained only if they exhibited a total gene count between 500 and 6000, a total UMI count between 1000 and 25000, and a mitochondrial gene expression proportion below 10%(Fig S1A). Subsequently, doublets were identified and removed using the Scrublet algorithm[87]. This method generates simulated doublets as a reference and computes a doublet score for each cell using a k-nearest neighbor classifier. Key parameters were set as follows: expected doublet rate = 0.06 (expected_doublet_rate=0.06), minimum counts per gene = 3 (min_counts=3), minimum cells per gene = 3 (min_cells=3), with log-transformation enabled (log_transform=True). The detection procedure focused on the top 85% most variable genes (min_gene_variability_pctl=85) and used the first 30 principal components

(n_prin_comps=30) to construct the feature space for discrimination. Cells with a doublet

score above the threshold of 0.2 were classified as doublets and excluded from

downstream analysis.Following doublet removal, the top 3000 highly variable genes were

selected from the quality-controlled data using the sc.pp.highly_variable_genes function.

Expression data were then normalized by total count (sc.pp.normalize_total,

target_sum=1e4) and log-transformed (sc.pp.log1p). Only highly variable genes were

scaled (sc.pp.scale), and linear dimensionality reduction was performed via principal

component analysis (sc.tl.pca). Finally, to correct for batch effects across samples, the

Harmony algorithm[88] (sc.external.pp.harmony_integrate) was applied to integrate the

principal components, yielding a batch-corrected low-dimensional embedding for

subsequent clustering and differential expression analysis.

Unsupervised clustering and subclustering

Following batch-effect correction, the integrated low-dimensional embedding was used for

unsupervised cell clustering. Cell communities were identified using the Louvain

algorithm[89] (implemented via sc.tl.louvain in Scanpy) with a resolution parameter set to

0.75. The nonlinear dimensionality reduction method UMAP (sc.tl.umap) was employed to

visualize the clusters in two-dimensional space. Differential expression analysis within

each cluster was performed using the sc.tl.rank_genes_groups function (with default

parameters), and the resulting genes were compared with known cell-type-specific

markers to assign preliminary cell identities. Based on the expression patterns of

canonical marker genes, six major cell populations were initially annotated, including

CD4$^+$ T cells, CD8$^+$ T cells, unconventional T cells, B cells, myeloid cells, and NK

cells.To further resolve the heterogeneity within each major population, subsets

corresponding to each of the above categories (e.g., myeloid, CD4$^+$ T, CD8$^+$ T, NK, and

B cells) were extracted as separate AnnData objects. For each subset, we reapplied

normalization, highly variable gene selection, PCA-based dimensionality reduction, and

sub-clustering using the Louvain algorithm at a resolution tailored to the size and internal

complexity of the subset. Cell-state identities of the sub-clusters were again determined

by differential expression analysis and comparison with established subtype-specific

markers. Ultimately, 34 distinct cell subpopulations were identified across the entire

dataset.

Differential abundance analysis using scRNA-seq data

To systematically evaluate the influence of exercise habits on immune cell composition,

we performed differential abundance analysis using the MiloR package

(https://github.com/MarioniLab/MILOR[31]) on single-cell RNA-sequencing data. First, To

capture the neighborhood relationships among cells, we constructed a k-nearest neighbor

graph (k=30) using the first 50 principal components. Cells were then partitioned into

overlapping neighborhoods using the makeNhoods function (prop = 0.2, k = 30, d = 50,

refined = TRUE). For cell counts in each neighborhood, We used MiloR to perform

differential abundance testing. This method fits a negative-binomial generalized linear

model, and we applied the trimmed mean of M-values (TMM) method for cross-sample

normalization.To isolate the independent effect of exercise status, the model included the

exercise group (EX vs. SED) as the primary variable while adjusting for sex and BMI as

covariates. Hypothesis testing was carried out via the testNhoods function. We classified

neighborhoods as exhibiting significant differential abundance if their false discovery rate

(FDR)-adjusted p-value was less than 0.05. The log2 fold-changes of these significant

neighborhoods were then used for subsequent visualization and biological interpretation.

Differential Expression Analysis of Cell Subpopulations

For the comparison of gene expression differences between the EX and SED groups

across cell subpopulations, we employed the linear model-based limma package[90] for

statistical testing. For each subpopulation, we constructed a linear model in which gene

expression was the dependent variable, exercise group was the primary predictor, and

sex and BMI were included as covariates. From the fitted model, the log2 fold-change

(log2FC) and raw p-value were extracted for every gene. To control for false positives, we

adopted the Benjamini-Hochberg (BH) procedure for multiple testing correction of the

p-values. In our analysis, a gene was considered a significant DEG within a given

subpopulation if it showed |log2FC| ≥ 0.25 together with an adjusted p-value < 0.05.

Module scoring using scRNA-seq data

To evaluate activity differences in specific biological processes across immune cell subpopulations, Scoring cells for specific gene modules was based on the Scanpy sc.tl.score_genes method, which computes the difference between the average expression of target genes and that of a matched random background per cell. For our analysis, we sourced gene sets from the Molecular Signatures Database[91] (MSigDB).To assess statistical differences in module scores between the EX and SED groups, we first computed the mean module score per subpopulation. We then fitted linear models comparing groups, adjusting for sex and BMI.

Gene regulatory network analysis based on SCENIC

We employed the pySCENIC toolkit (v0.12.1)[92] to infer potential transcriptional regulatory networks across distinct immune cell subpopulations. Our analysis began by randomly sampling a subset of 10,000 cells to serve as the training set. Using a comprehensive transcription factor list, we then applied the GRNBoost2 algorithm[93] to infer co-expression relationships between each transcription factor and all potential target genes. This step generated the initial gene regulatory network. Subsequently, the preliminary network was pruned using the cisTarget database. Only target genes whose transcription start sites (TSS) fell within ±10 kb of a corresponding transcription factor binding motif were retained, yielding high-confidence regulons. Next, the AUCell module of pySCENIC was used to calculate the activity level (represented as the area under the curve, AUC) of each regulon in all cells, followed by z-score normalization. To identify regulons with differential activity between the exercise (EX) and sedentary (SED) groups, we compared the AUC z-scores of regulons across cell types using the limma package. Regulons meeting the thresholds of adjusted p-value < 0.05 and |log2FoldChange| ≥ 0.25 were defined as differentially active.

scATAC-seq quality control.

We used ArchR[94] (version 1.0.2) to analyze single-cell ATAC-seq data. First, Arrow files

were generated for each sample from the raw sequencing fragments following the default

settings. Subsequently, stringent cell-level quality control was applied. Our cell quality

control was based on the parameters below: TSS enrichment ≥ 10, number of unique

fragments between $1×10^3$ and $1×10^5$, nucleosome signal < 4, and blacklist ratio < 0.05.

Doublets were identified and removed using ArchR's default parameters, resulting in a

high-quality chromatin accessibility matrix for downstream analysis.

scATAC-seq dimensionality reduction and clustering.

To construct a single-cell chromatin accessibility landscape, we first performed

dimensionality reduction on the cell-by-peak matrix using latent semantic indexing (LSI) to

capture dominant patterns of chromatin accessibility variation. To mitigate technical

variability introduced during sample preparation, we applied the Harmony[88] algorithm to

integrate the reduced-dimension embeddings across samples.Using the integrated

low-dimensional representation, we partitioned cells into communities via shared nearest

neighbor (SNN)-based graph clustering. An initial clustering step was performed at a

uniform resolution of 0.8 to delineate major cell categories. To further resolve cellular

heterogeneity and enhance the detection of small-scale cell states, we carried out

sub-clustering on selected populations derived from the initial clustering, with resolutions

adapted between 0.3 and 1.5 according to the intrinsic structure of each cell group. As the

final step in our single-cell ATAC-seq analysis, all cells were projected into a

two-dimensional space for visualization using uniform manifold approximation and

projection (UMAP), providing an intuitive representation of cell-state distributions and

community relationships.

scATAC–seq gene activity scores.

Gene activity was quantified with ArchR[94] (v1.0.2) , which computes scores by correlating

accessibility across gene-linked regions. Following this, the MAGIC algorithm[95] was

applied to smooth the scores and mitigate noise inherent in sparse scATAC-seq data.

Integrating scATAC-seq and scRNA-seq Data to Infer Cell Identity

To accurately annotate cell types in the scATAC-seq data, we performed an integrative

analysis against a matched scRNA-seq reference. This was achieved using the

addGeneIntegrationMatrixfunction in ArchR. This process was conducted in two

sequential steps: first, an unconstrained integration was carried out to broadly map each

scATAC-seq cell cluster to the major cell types defined in the scRNA-seq data.

Subsequently, based on this initial mapping, A second, more constrained integration was

then carried out by re-executing addGeneIntegrationMatrix with updated broad group

labels. This refinement enabled finer multimodal alignment and produced final annotations

at high resolution, ensuring consistency with the transcriptomic profiles.To independently

assess the reliability of the annotations, we employed the Gene Activity Score (GAS)

matrix derived from the scATAC-seq data for validation. By comparing the distribution of

annotated cells in the gene-activity space with that of the corresponding scRNA-seq

reference cells in gene-expression space, we further confirmed the accuracy of the

cross-modal annotation.

scATAC–seq pseudobulk replicate generation and peak calling.

To support differential analysis across cell clusters and conditions, we first constructed

pseudobulk samples. This involved running the addGroupCoverages function with tailored

parameters to create non-overlapping replicates for each cell group. These replicates

were then consolidated via addReproduciblePeakSet to generate a consensus peak

matrix. For precise peak identification, we used MACS2[96] . The resulting unified peak set

provided the foundation for all subsequent differential accessibility testing.

scATAC–seq genomic regions annotation.

For the identified chromatin-accessible regions (peaks), We annotated the called peaks

by applying the annotatePeak function from the ChIPseeker package [97] package with

default parameters, in order to determine the nearest associated genes and genomic

features for each peak.

scATAC motif enrichment and motif deviation analysis

We carried out transcription factor motif enrichment followed by chromatin accessibility

deviation analysis using the unified pseudobulk peak set. For motif annotation, we utilized

the CIS-BP motif database (via ChromVAR[98]), JASPAR2020[99], and HOMER[100].

Additionally, motif accessibility deviation scores were calculated for each cell using the

chromVAR algorithm implemented in ArchR, enabling systematic assessment of changes

in transcription factor binding activity across different cellular states.

Identification of differential peaks

To detect differentially accessible chromatin regions between groups, we employed two

complementary approaches. First, we used the getMarkerFeatures function with

parameters useMatrix = "PeakMatrix", bias covariates including "TSSEnrichment" and

"log10(nFrags)", and testMethod = "wilcoxon" to identify differential peaks between

groups, applying thresholds of FDR ≤ 0.1 and log2FC ≥ 0.25. Second, for differential

transcription factor activity analysis, we applied the same function with useMatrix =

"MotifMatrix" and thresholds of FDR ≤ 0.01 and MeanDiff ≥ 0.05.

Gene Ontology Enrichment Analysis of differential peaks

To interpret the potential biological functions of differential chromatin accessible regions,

we performed Gene Ontology (GO) enrichment analysis using the R package rGREAT[101].

This package associates genomic regions with nearby genes and evaluates their

enrichment significance across Biological Process, Molecular Function, and Cellular

Component terms using a hypergeometric test. The analysis was conducted with the

human genome as background, applying default parameters, We selected terms passing

the significance threshold of an FDR-adjusted p-value < 0.05 as significantly enriched.

scATAC–seq TF Foot-print analysis

We performed transcription factor footprinting analysis based on Tn5 transposase

cleavage profiles across genomic motif regions. First, the sequence bias of Tn5 was

subtracted from the raw cleavage signal to obtain a corrected footprint profile. Motif

positions were determined using a combination of the CIS-BP database[102] (from

chromVAR human_pwms_v1) and the JASPAR2020 database. Subsequently, footprint

signals were normalized using the average insertion density within ±200–250 bp regions

centered on each motif. The normalized results were aggregated by pseudo-replicate, and

their means and standard deviations were calculated and visualized. Finally, to assess

differences in footprint signals between groups, We chose the nonparametric Wilcoxon

rank-sum test for comparisons between two groups, setting the significance level at p <

0.05.

scATAC-seq Peak to gene linkage analysis

To identify peak-to-gene links, we performed prediction analysis using the

addPeak2GeneLinks function in ArchR. The analysis was based on the batch-corrected

low-dimensional embedding (specified via the reducedDims parameter) and computed the

correlation between chromatin accessibility and gene expression. A correlation threshold

of corCutOff = 0.4 was applied to screen for high-confidence linkages. The resulting

significant associations were output as a GRanges object for subsequent visualization

and functional interpretation.

Cell-cell interaction analysis

We performed a systematic analysis of potential signaling communication among all

immune cell subpopulations using the CellChat R package[75] (v.1.5.0). This approach

leverages the built-in ligand–receptor pair databases of CellChat to identify and

characterize potential signaling interaction networks between different immune cell

subtypes.

Acknowledgements

We express our sincere appreciation to the China National GeneBank (CNGB).

Funding

Funding for this research was provided by the National Key Research and Development

Program of China (2025YFC3409300, 2022YFC3400400) and the National Science and

Technology Major Project (2024ZD0530500).

Author contributions

928 X.S., J.L., S.G., J.Y., P.Y., and C.L. conceived the idea; C.L., P.Y., J.Y., Y.Z., and J.Z.
929 supervised the work; X.S., J.L., S.G., S.X., Y.W., Y.Z., and W.Z. analyzed the data and
930 performed visualization; X.S., J.L., S.G., P.Y., J.Y., and C.L. wrote the manuscript.

Competing interests

933 The authors declare no competing interests.

Data availability

936 The single-cell RNA-seq and single-cell ATAC-seq data, plasma metabolomics and
937 lipidomics data, and blood biochemistry data presented in this study were sourced from
938 the CIMA cohort, hosted on the TrueBlood database within the CNGBdb portal
939 (https://db.cngb.org/trueblood/cima/).

Code availability

942 The code can be obtained from the corresponding author upon reasonable request.
943 Information on publicly available software used is provided in the Methods.

Reference

1. Schmid, D. & Leitzmann, M. F. Association between physical activity and mortality among breast cancer and colorectal cancer survivors: a systematic review and meta-analysis. *Ann Oncol* **25**, 1293–1311 (2014).

2. Moore, S. C. *et al.* Association of Leisure-Time Physical Activity With Risk of 26 Types of Cancer in 1.44 Million Adults. *JAMA Intern Med* **176**, 816–825 (2016).

3. Patel, A. V. *et al.* American College of Sports Medicine Roundtable Report on Physical Activity, Sedentary Behavior, and Cancer Prevention and Control. *Med Sci Sports Exerc* **51**, 2391–2402 (2019).

4. Im, L. *et al.* Effect of physical inactivity on major non-communicable diseases worldwide: an analysis of burden of disease and life expectancy. *Lancet (London,*

England) **380**, (2012).

5.   Meyerhardt, J. A. *et al.* Impact of physical activity on cancer recurrence and
survival in patients with stage III colon cancer: findings from CALGB 89803. *J Clin*
Oncol **24**, 3535–3541 (2006).

6.   Chastin, S. F. M. *et al.* Effects of Regular Physical Activity on the Immune System,
Vaccination and Risk of Community-Acquired Infectious Disease in the General
Population: Systematic Review and Meta-Analysis. *Sports Med* **51**, 1673–1686
(2021).

7.   Hawley, J. A., Hargreaves, M., Joyner, M. J. & Zierath, J. R. Integrative biology of
exercise. *Cell* **159**, 738–749 (2014).

8.   Booth, F. W., Roberts, C. K., Thyfault, J. P., Ruegsegger, G. N. & Toedebusch, R.
G. Role of Inactivity in Chronic Diseases: Evolutionary Insight and Pathophysiological
Mechanisms. *Physiol Rev* **97**, 1351–1402 (2017).

9.   Pedersen, B. K. & Febbraio, M. A. Muscles, exercise and obesity: skeletal muscle
as a secretory organ. *Nat Rev Endocrinol* **8**, 457–465 (2012).

10. Sun, S. *et al.* A single-cell transcriptomic atlas of exercise-induced
anti-inflammatory and geroprotective effects across the body. *Innovation (Camb)* **4**,
100380 (2023).

11. K, C. *et al.* Molecular Choreography of Acute Exercise. *Cell* **181**, (2020).

12. Geng, L. *et al.* Systematic profiling reveals betaine as an exercise mimetic for
geroprotection. *Cell* **188**, 5403-5425.e33 (2025).

13. MoTrPAC Study Group, Lead Analysts, & MoTrPAC Study Group. Temporal
dynamics of the multi-omic response to endurance exercise training. *Nature* **629**,
174–183 (2024).

14. Mathis, D. & Shoelson, S. E. Immunometabolism: an emerging frontier. *Nat Rev*
Immunol **11**, 81 (2011).

15. Pedersen, B. K. & Saltin, B. Exercise as medicine - evidence for prescribing
exercise as therapy in 26 different chronic diseases. *Scand J Med Sci Sports* **25**
Suppl 3, 1–72 (2015).

16. Ki, S. *et al.* 12,13-diHOME: An Exercise-Induced Lipokine that Increases Skeletal
Muscle Fatty Acid Uptake. *Cell metabolism* **27**, (2018).

17. Shimazu, T. *et al.* Suppression of oxidative stress by β-hydroxybutyrate, an
endogenous histone deacetylase inhibitor. *Science* **339**, 211–214 (2013).

18. Jc, N. & E, V. β-Hydroxybutyrate: A Signaling Metabolite. *Annual review of*
nutrition **37**, (2017).

19. Pedersen, B. K. & Ullum, H. NK cell response to physical activity: possible
mechanisms of action. *Med Sci Sports Exerc* **26**, 140–146 (1994).

20. Pedersen, B. K. & Hoffman-Goetz, L. Exercise and the immune system:
regulation, integration, and adaptation. *Physiol Rev* **80**, 1055–1081 (2000).

21. Nieman, D. C. *et al.* The acute immune response to exhaustive resistance
exercise. *Int J Sports Med* **16**, 322–328 (1995).

22. Simpson, R. J., Kunz, H., Agha, N. & Graff, R. Exercise and the Regulation of
Immune Functions. *Prog Mol Biol Transl Sci* **135**, 355–380 (2015).

23. Yin, J., Zheng, Y., Huang, Z. & Zhou, W. Chinese Immune Multi-Omics Atlas.
Science **391**, (2026).

24. Bull, F. C. *et al.* World Health Organization 2020 guidelines on physical activity
and sedentary behaviour. *Br J Sports Med* **54**, 1451–1462 (2020).

25. Mann, S., Beedie, C. & Jimenez, A. Differential effects of aerobic exercise,
resistance training and combined exercise modalities on cholesterol and the lipid
profile: review, synthesis and recommendations. *Sports Med* **44**, 211–221 (2014).

26. Aadahl, M. *et al.* Five-year change in physical activity is associated with changes
in cardiovascular disease risk factors: the Inter99 study. *Prev Med* **48**, 326–331
(2009).

27. S, K. *et al.* Effect of aerobic exercise training on serum levels of high-density
lipoprotein cholesterol: a meta-analysis. *Archives of internal medicine* **167**, (2007).

28. Baxmann, A. C. *et al.* Influence of muscle mass and physical activity on serum
and urinary creatinine and serum cystatin C. *Clin J Am Soc Nephrol* **3**, 348–354
(2008).

29. Waring, W. S. *et al.* Uric acid reduces exercise-induced oxidative stress in healthy
adults. *Clin Sci (Lond)* **105**, 425–430 (2003).

30. Stocker, R., Yamamoto, Y., McDonagh, A. F., Glazer, A. N. & Ames, B. N. Bilirubin
is an antioxidant of possible physiological importance. *Science* **235**, 1043–1046
(1987).

31. Dann, E., Henderson, N. C., Teichmann, S. A., Morgan, M. D. & Marioni, J. C.
Differential abundance testing on single-cell data using k-nearest neighbor graphs.
Nat Biotechnol **40**, 245–253 (2022).

32. Schooneman, M. G., Vaz, F. M., Houten, S. M. & Soeters, M. R. Acylcarnitines: reflecting or inflicting insulin resistance? *Diabetes* **62**, 1–8 (2013).

33. M, E., Ke, C. & B, E. Metabolism of ketone bodies during exercise and training: physiological basis for exogenous supplementation. *The Journal of physiology* **595**, (2017).

34. Tan, J. *et al.* The role of short-chain fatty acids in health and disease. *Adv Immunol* **121**, 91–119 (2014).

35. Sk, P. & Mj, J. Exercise-induced oxidative stress: cellular mechanisms and impact on muscle force production. *Physiological reviews* **88**, (2008).

36. Christen, S., Thomas, S. R., Garner, B. & Stocker, R. Inhibition by interferon-gamma of human mononuclear cell-mediated low density lipoprotein oxidation. Participation of tryptophan metabolism along the kynurenine pathway. *J Clin Invest* **93**, 2149–2158 (1994).

37. Gall, W. E. *et al.* alpha-hydroxybutyrate is an early biomarker of insulin resistance and glucose intolerance in a nondiabetic population. *PLoS One* **5**, e10883 (2010).

38. Stockinger, B., Di Meglio, P., Gialitakis, M. & Duarte, J. H. The aryl hydrocarbon receptor: multitasking in the immune system. *Annu Rev Immunol* **32**, 403–432 (2014).

39. Bhat, R. *et al.* Inhibitory role for GABA in autoimmune inflammation. *Proc Natl Acad Sci U S A* **107**, 2580–2585 (2010).

40. Wlodarska, M. *et al.* Indoleacrylic Acid Produced by Commensal Peptostreptococcus Species Suppresses Inflammation. *Cell Host Microbe* **22**, 25-37.e6 (2017).

41. Hang, S. *et al.* Bile acid metabolites control TH17 and Treg cell differentiation. *Nature* **576**, 143–148 (2019).

42. van Loon, L. J. C. & Goodpaster, B. H. Increased intramuscular lipid storage in the insulin-resistant and endurance-trained state. *Pflugers Arch* **451**, 606–616 (2006).

43. O'Neill, L. A. J., Kishton, R. J. & Rathmell, J. A guide to immunometabolism for immunologists. *Nat Rev Immunol* **16**, 553–565 (2016).

44. Hotamisligil, G. S. Inflammation and metabolic disorders. *Nature* **444**, 860–867 (2006).

45. Simpson, R. J., Kunz, H., Agha, N. & Graff, R. Exercise and the Regulation of Immune Functions. *Prog Mol Biol Transl Sci* **135**, 355–380 (2015).

46. Storm van's Gravesande, K. *et al.* IFN regulatory factor-1 regulates
IFN-gamma-dependent cathepsin S expression. *J Immunol* **168**, 4488–4494 (2002).

47. van der Stoep, N., Quinten, E., Marcondes Rezende, M. & van den Elsen, P. J.
E47, IRF-4, and PU.1 synergize to induce B-cell-specific activation of the class II
transactivator promoter III (CIITA-PIII). *Blood* **104**, 2849–2857 (2004).

48. Das, H. *et al.* Kruppel-like factor 2 (KLF2) regulates proinflammatory activation of
monocytes. *Proc Natl Acad Sci U S A* **103**, 6653–6658 (2006).

49. Liao, X. *et al.* Krüppel-like factor 4 regulates macrophage polarization. *J Clin
Invest* **121**, 2736–2749 (2011).

50. Shaulian, E. & Karin, M. AP-1 as a regulator of cell life and death. *Nat Cell Biol* **4**,
E131-136 (2002).

51. Igarashi, K. & Sun, J. The heme-Bach1 pathway in the regulation of oxidative
stress response and erythroid differentiation. *Antioxid Redox Signal* **8**, 107–118
(2006).

52. Simpson, R. J., Kunz, H., Agha, N. & Graff, R. Exercise and the Regulation of
Immune Functions. *Prog Mol Biol Transl Sci* **135**, 355–380 (2015).

53. Intlekofer, A. M. *et al.* Effector and memory CD8+ T cell fate coupled by T-bet and
eomesodermin. *Nat Immunol* **6**, 1236–1244 (2005).

54. Pearce, E. L. *et al.* Control of effector CD8+ T cell function by the transcription
factor Eomesodermin. *Science* **302**, 1041–1043 (2003).

55. O'Shea, J. J. & Plenge, R. JAK and STAT signaling molecules in
immunoregulation and immune-mediated disease. *Immunity* **36**, 542–550 (2012).

56. Simpson, R. J., Kunz, H., Agha, N. & Graff, R. Exercise and the Regulation of
Immune Functions. *Prog Mol Biol Transl Sci* **135**, 355–380 (2015).

57. Vivier, E., Tomasello, E., Baratin, M., Walzer, T. & Ugolini, S. Functions of natural
killer cells. *Nat Immunol* **9**, 503–510 (2008).

58. Daussy, C. *et al.* T-bet and Eomes instruct the development of two distinct natural
killer cell lineages in the liver and in the bone marrow. *J Exp Med* **211**, 563–577
(2014).

59. Ramirez, K. *et al.* Gene deregulation and chronic activation in natural killer cells
deficient in the transcription factor ETS1. *Immunity* **36**, 921–932 (2012).

60. Hayden, M. S. & Ghosh, S. Regulation of NF-κB by TNF family cytokines. *Semin*

Immunol **26**, 253–266 (2014).

61. Kobayashi, T. & Mattarollo, S. R. Natural killer cell metabolism. *Mol Immunol* **115**,
3–11 (2019).

62. A, C., Dr, L. & I, T. RUNX proteins in transcription factor networks that regulate
T-cell lineage choice. *Nature reviews. Immunology* **9**, (2009).

63. Kaplan, M. H., Schindler, U., Smiley, S. T. & Grusby, M. J. Stat6 is required for
mediating responses to IL-4 and for development of Th2 cells. *Immunity* **4**, 313–319
(1996).

64. Ikawa, T. *et al.* An essential developmental checkpoint for production of the T cell
lineage. *Science* **329**, 93–96 (2010).

65. Ramirez, K. *et al.* Gene deregulation and chronic activation in natural killer cells
deficient in the transcription factor ETS1. *Immunity* **36**, 921–932 (2012).

66. Matsuyama, T. *et al.* Targeted disruption of IRF-1 or IRF-2 results in abnormal
type I IFN gene induction and aberrant lymphocyte development. *Cell* **75**, 83–97
(1993).

67. Platanias, L. C. Mechanisms of type-I- and type-II-interferon-mediated signalling.
Nat Rev Immunol **5**, 375–386 (2005).

68. Wojakowski, W., Landmesser, U., Bachowski, R., Jadczyk, T. & Tendera, M.
Mobilization of stem and progenitor cells in cardiovascular diseases. *Leukemia* **26**,
23–33 (2012).

69. Appay, V. & Rowland-Jones, S. L. RANTES: a versatile and controversial
chemokine. *Trends Immunol* **22**, 83–87 (2001).

70. Scott, E. W., Simon, M. C., Anastasi, J. & Singh, H. Requirement of transcription
factor PU.1 in the development of multiple hematopoietic lineages. *Science* **265**,
1573–1577 (1994).

71. Lin, H. & Grosschedl, R. Failure of B-cell differentiation in mice lacking the
transcription factor EBF. *Nature* **376**, 263–267 (1995).

72. Gerondakis, S. & Siebenlist, U. Roles of the NF-kappaB pathway in lymphocyte
development and function. *Cold Spring Harb Perspect Biol* **2**, a000182 (2010).

73. R, S. *et al.* Graded expression of interferon regulatory factor-4 coordinates
isotype switching with plasma cell differentiation. *Immunity* **25**, (2006).

74. Rn, G., M, M.-S., A, N.-L. & Id, F. Systems biology in immunology: a

computational modeling perspective. *Annual review of immunology* **29**, (2011).

75. Jin, S. *et al.* Inference and analysis of cell-cell communication using CellChat. *Nat*
Commun **12**, 1088 (2021).

76. M, F., M, H., S, G. & L, S. The role of resistin as a regulator of inflammation:
Implications for various human pathologies. *Clinical immunology (Orlando, Fla.)* **133**,
(2009).

77. Silswal, N. *et al.* Human resistin stimulates the pro-inflammatory cytokines
TNF-alpha and IL-12 in macrophages by NF-kappaB-dependent pathway. *Biochem*
Biophys Res Commun **334**, 1092–1101 (2005).

78. Leitinger, B. & Hohenester, E. Mammalian collagen receptors. *Matrix Biol* **26**,
146–155 (2007).

79. Griffith, J. W., Sokol, C. L. & Luster, A. D. Chemokines and chemokine receptors:
positioning cells for host defense and immunity. *Annu Rev Immunol* **32**, 659–702
(2014).

80. O'Neill, L. A. J., Kishton, R. J. & Rathmell, J. A guide to immunometabolism for
immunologists. *Nat Rev Immunol* **16**, 553–565 (2016).

81. Youm, Y.-H. *et al.* The ketone metabolite β-hydroxybutyrate blocks NLRP3
inflammasome-mediated inflammatory disease. *Nat Med* **21**, 263–269 (2015).

82. Yang, C. *et al.* Betaine Ameliorates Experimental Autoimmune Encephalomyelitis
by Inhibiting Dendritic Cell-Derived IL-6 Production and Th17 Differentiation. *J*
Immunol **200**, 1316–1324 (2018).

83. Bhandage, A. K. *et al.* GABA Regulates Release of Inflammatory Cytokines From
Peripheral Blood Mononuclear Cells and CD4+ T Cells and Is Immunosuppressive in
Type 1 Diabetes. *EBioMedicine* **30**, 283–294 (2018).

84. Pang, Z. *et al.* MetaboAnalyst 6.0: towards a unified platform for metabolomics
data processing, analysis and interpretation. *Nucleic Acids Res* **52**, W398–W406
(2024).

85. Pang, Z., Chong, J., Li, S. & Xia, J. MetaboAnalystR 3.0: Toward an Optimized
Workflow for Global Metabolomics. *Metabolites* **10**, 186 (2020).

86. Fa, W., P, A. & Fj, T. SCANPY: large-scale single-cell gene expression data
analysis. *Genome biology* **19**, (2018).

87. Wolock, S. L., Lopez, R. & Klein, A. M. Scrublet: Computational Identification of
Cell Doublets in Single-Cell Transcriptomic Data. *Cell Syst* **8**, 281-291.e9 (2019).

88. Korsunsky, I. *et al.* Fast, sensitive and accurate integration of single-cell data with
Harmony. *Nat Methods* **16**, 1289–1296 (2019).

89. Large-scale network dynamics underlying the first few hundred milliseconds after
stimulus presentation: An investigation of visual recognition memory using iEEG -
PubMed. https://pubmed.ncbi.nlm.nih.gov/37688546/.

90. Ritchie, M. E. *et al.* limma powers differential expression analyses for
RNA-sequencing and microarray studies. *Nucleic Acids Res* **43**, e47 (2015).

91. Subramanian, A. *et al.* Gene set enrichment analysis: a knowledge-based
approach for interpreting genome-wide expression profiles. *Proc Natl Acad Sci U S A*
102, 15545–15550 (2005).

92. S, A. *et al.* SCENIC: single-cell regulatory network inference and clustering.
Nature methods **14**, (2017).

93. Moerman, T. *et al.* GRNBoost2 and Arboreto: efficient and scalable inference of
gene regulatory networks. *Bioinformatics* **35**, 2159–2161 (2019).

94. Granja, J. M. *et al.* ArchR is a scalable software package for integrative single-cell
chromatin accessibility analysis. *Nat Genet* **53**, 403–411 (2021).

95. van Dijk, D. *et al.* Recovering Gene Interactions from Single-Cell Data Using Data
Diffusion. *Cell* **174**, 716-729.e27 (2018).

96. Zhang, Y. *et al.* Model-based analysis of ChIP-Seq (MACS). *Genome Biol* **9**,
R137 (2008).

97. Yu, G., Wang, L.-G. & He, Q.-Y. ChIPseeker: an R/Bioconductor package for
ChIP peak annotation, comparison and visualization. *Bioinformatics* **31**, 2382–2383
(2015).

98. Schep, A. N., Wu, B., Buenrostro, J. D. & Greenleaf, W. J. chromVAR: inferring
transcription-factor-associated accessibility from single-cell epigenomic data. *Nat*
Methods **14**, 975–978 (2017).

99. Fornes, O. *et al.* JASPAR 2020: update of the open-access database of
transcription factor binding profiles. *Nucleic Acids Res* **48**, D87–D92 (2020).

100. Heinz, S. *et al.* Simple combinations of lineage-determining transcription
factors prime cis-regulatory elements required for macrophage and B cell identities.
Mol Cell **38**, 576–589 (2010).

101. Tanigawa, Y., Dyer, E. S. & Bejerano, G. WhichTF is functionally important in
your open chromatin data? *PLoS Comput Biol* **18**, e1010378 (2022).

102.   Weirauch, M. T. *et al.* Determination and inference of eukaryotic transcription
factor sequence specificity. *Cell* **158**, 1431–1443 (2014).

1186

Figure 1. Study design, physiological characteristics, and multi-omics
characterization of habitual exercisers versus sedentary controls.

(A) Overall framework of the present study design, about stratification of 86 participants
from the Chinese Immune Multi-Omics Cohort (CIMA) into a regular active exercise group
(EX, n=40) meeting WHO physical activity guidelines and a sedentary control group (SED,
n=46), followed by integrated multi-omics profiling of plasma and peripheral blood
mononuclear cell fractions and integrative analysis.

(B) Analysis of Differences in Blood Biochemical Indicators Between the regular active
exercise group (EX) and sedentary control group (SED). A pie chart showing changes in
blood biochemical indicators(left). A volcano plot displaying the log2 fold change (Log$_2$FC)
and significance (-log10 p-value) of blood biochemical indicators across groups.

(C) A line chart illustrates the distribution of differentially expressed molecules across
different categories of plasma metabolomics and lipidomics.

(D) Uniform Manifold Approximation and Projection (UMAP) visualization depicting
1,317,665 immune cells derived from single-cell RNA sequencing (scRNA-seq) data
following dimensionality reduction, clustering, and cell annotation, colored by 34 immune
lineages.

(E) A dotplot illustrating the average expression and expression percentage of marker
genes across distinct immune cell subsets, showing both the magnitude and prevalence
of gene expression.

(F) UMAP plot depicting 644,045 immune cells from scATAC-seq datasets, where
annotation labels were transferred from scRNA-seq data and each cell type was assigned
a distinct color code. (match in D).

(G) Visualization of chromatin openness at key lineage gene promoter regions, including
B cells (MS4A1), T cells (IL7R), CD4+ T cells (CD4), CD8+ T cells (CD8B), NK cells
(KLRB1), and non-canonical T cells (RORC).

(H) Box plot showing the percentage of myeloid cells in the SED and EX groups.The
distribution of myeloid cell percentages in the sedentary (SED, blue) and exercise (EX,
green) groups is shown. The P value (P < 0.05) was derived from a linear model adjusted
for age and sex, followed by Benjamini-Hochberg correction.

1219

Figure 2. Plasma metabolic and lipidomic changes induced by habitual physical activity.

(A) A volcano plot illustrating the differential abundance of plasma molecules (metabolites and complex lipids) between the EX and SED groups, the inset pie chart summarizes the composition of 62 differentially abundant molecules.

(B) Pathway enrichment analysis of metabolites up-regulated in the EX group, highlighting significant enrichment in core energy metabolism pathways (e.g., TCA cycle), amino acid metabolism, and antioxidant defense pathways (e.g., glutathione metabolism).

(C) Box plots showing the normalized intensity of key fatty acid oxidation (FAO)-related metabolites between the SED (blue) and EX (green) groups. P-values indicate the significance of group differences.

(D) Regulatory analysis of antioxidants defense system-related metabolites (betaine, 3-hydroxyindoleacetic acid, 2-hydroxybutyric acid) between the SED group (blue) and EX group (green). P-values indicate the significance of intergroup differences.

(E) Changes in molecules associated with gut microbiota-bile acid metabolism between the SED group (blue) and EX group (green). P-values indicate significant differences between groups.

(F) A box plot showing the mean intensity of distinct triglyceride (TAG) species between the SED (blue) and EX (green) groups. P-values indicate the significance of group differences.

(G) Normalized Intensity Distribution of Membrane-Associated Complex Lipids in EX and SED.P-values indicate group difference significance, with violin plots showing intensity distribution.

(H) Bar plots showing the intensity distribution of two PE species between the SED (blue) and EX (green) groups, with P-values denoting significance.

1245

1246 **Figure 3. Multi-omics analysis of Myeloid cells heterogeneity in regular active**

1247 **exercise group (EX).**

1248 (A) UMAP plot displays scRNA-seq-derived Myeloid cells after subclustering and

annotation, colored by five functional subsets(left). UMAP plot shows scATAC-seq-derived Myeloid cells, with annotation labels from corresponding scRNA-seq data and each cell type color-marked(right).

(B) MiloR-based analysis was performed to assess differences in the abundance of Myeloid cell subsets between the EX and SED groups.

(C) Enrichment analysis of Gene Ontology (GO) pathways in differentially expressed genes (DEGs) in each Myeloid cell subset, showing both upregulated and downregulated pathways in the EX group.

(D) The violin plot showing changes in the expression levels of antigen presentation-related genes and cell migration-related genes within specific subsets, based on single-cell RNA data.

(E) GO pathway enrichment analysis of differentially accessible regions (DARs) identified in classical monocytes (cMono) from scATAC-seq data.

(F) Enrichment analysis of transcription factor (TF) binding motifs in DARs of cMono, revealing enriched TF motifs in the up-regulated chromatin-accessible peaks of the EX group.

(G) Motif bias analysis showing the accessibility bias of key transcription factor binding motifs in cMono between the EX and SED groups. Positive values indicate increased accessibility in the EX group.

(H) The chromatin accessibility at differential accessible regions (DARs) of HLA-DQB1 (left) and HLA-DQA1 (right) between the SED and EX groups.

(I) The chromatin accessibility at the differential accessible region (DAR) of CXCL8 between the SED and EX groups.

(J) Violin plot showing expression level of the pro-inflammatory gene CXCL8 in cMono of the EX group based on scRNA-seq data.

(K) A heatmap showing SCENIC regulon AUCell scores (activity) of transcription factors (TFs) in cMono cells between the EX and SED groups.

(L) A regulatory network plot showing the synergistic module of key TFs and their target genes in cMono cells.

1279

1280   **Figure 4. Epigenetic reprogramming enhances T cell functions in EX group.**

(A) UMAP plot displays scRNA-seq-derived T cells after subclustering and annotation, colored by five functional subsets(left). UMAP plot shows scATAC-seq-derived T cells, with annotation labels from corresponding scRNA-seq data and each cell type color-marked(right).

(B) MiloR-based analysis was performed to assess differences in the abundance of T cell subsets between the EX and SED groups.

(C) Bar chart showing the number of upregulated differentially accessible regions (DARs) in each T cell subset from scATAC-seq data. The inset dot plot shows the proportion of DARs shared between two or more subsets.

(D) Enrichment analysis of Gene Ontology (GO) pathways in differentially expressed genes (DEGs) in each T cell subset, showing both upregulated and downregulated pathways in the EX group.

(E) Score distribution of characteristic gene sets in scRNA-seq data from T cell subpopulations differing between SED and EX groups.

(F) Violin plots showing the chromatin accessibility of core transcription factors (TFs) in CD8_CTL and MAIT cells between the SED and EX groups. P-adjusted (Padj) values indicate significance of group differences.

(G) A heatmap showing SCENIC regulon AUCell scores (activity) of transcription factors (TFs) in CD8_CTL cells between the EX and SED groups.

(H) Target gene regulatory network of RUNX3 in CD8_CTL cells.

(I) The chromatin accessibility at the differential accessible region (DAR) of GZMB(left) and TBX21(right) between the SED and EX groups in CD8_CTL cells,All significant peak–gene associations identified for each gene are presented alongside the corresponding track.

(J) The chromatin accessibility at the differential accessible region (DAR) of PRF1 between the SED and EX groups in MAIT cells.

1307

Figure 5. Multi-Omics characterization of NK cell heterogeneity in EX group.

(A) UMAP plot displays scRNA-seq-derived NK cells after subclustering and annotation,

colored by five functional subsets(left). UMAP plot shows scATAC-seq-derived NK cells,

with annotation labels from corresponding scRNA-seq data and each cell type

color-marked(right).

(B) MiloR-based analysis was performed to assess differences in the abundance of NK

cell subsets between the EX and SED groups.

(C) A heatmap showing normalized module scores of functional pathways across NK cell

subsets between the SED and EX groups. Asterisks denote significance of group

differences($*P<0.05, **P<0.01, ***P<0.001$).

(D) Enrichment analysis of Gene Ontology (GO) pathways in differentially expressed

genes (DEGs) in each Mature_nk cell subset, showing both upregulated and

downregulated pathways in the EX group.

(E) Violin plots showing the motif deviation of TFs (EOMES, TBX21) linked to NK cell

effector function between the SED and EX groups. P-adjusted (Padj) values indicate

significance of group differences.

(F) Footprint plots displaying the normalized Tn5 bias (y-axis) around the motif center for

RELA and RELB in the SED and EX groups.

(G) Venn diagram showing the core transcription factors commonly upregulated in both

CD8_CTL (adaptive immunity) and Mature_NK (innate immunity) cells of the EX group, as

well as factors uniquely upregulated in each population.

(H) The chromatin accessibility at the differential accessible region (DAR) of STAT1(left)

and CXCR4(right) between the SED and EX groups in Mature_NK cells.

(I) The chromatin accessibility at the differential accessible region (DAR) of RELB(left)

and CCL5(right) between the SED and EX groups in Mature_NK cells.

1333

Figure 6. The impact of regular active exercise on the communication network

among immune cells.

(A) A chart displaying the number of cell-cell interactions between immune cell subsets in

the SED group and EX group. The top bar chart summarizes total interactions (766 in

SED group, 777 in EX group), while the lower data points represent interaction counts for

each subset.

(B) Heatmap showing the overall interaction strength between cell types in the SED (left)

and EX (right) groups, computed using the CellChat algorithm.

(C) This diagram compares the information flow strength of selected signaling pathways

(e.g., THBS, IL6, RESISTIN) between the SED (sedentary control, blue) and EX (exercise

intervention, green) groups.

(D) A regulatory network plot showing the cellular communication network for the
RESISTIN signaling pathway in SED group(left) and EX group(right).

(E) Analysis of the key ligand-receptor axes in the RESISTIN pathway, comparing the
interaction intensity for the RETN--CAP1 and RETN--TLR4 axes between the EX and
SED groups.

(F) Violin plot comparing the expression level of the ligand gene RETN in myeloid cells
between the two groups. Asterisks denote significance of group differences(***$P<0.001$),
$Log_2FC=-0.721$.

(G) A regulatory network plot showing MHC-II-dependent signaling between immune cell
subsets (nodes: cell types; edges: interaction strength) in the SED (left) and EX (right)
groups.

(H) Cell-cell communication probability of MHC-II-related ligand-receptor pairs. Color
depth reflects communication probability; larger bubbles mean more significant p - values.

(I) Cell-cell communication probability of MHC-I-related ligand-receptor pairs. Color
depth reflects communication probability; larger bubbles mean more significant p - values.

(J) A regulatory network plot showing IFN-γ-dependent signaling (edges: interaction
strength; nodes: cell types) in the SED (left) and EX (right) groups.

(K) A heatmap showing the communication probability (color gradient) and significance
(markers) of the IFNG–(IFNGR1+IFNGR2) ligand-receptor axis: sources are cytotoxic T
cells (CD4[+] CTL/CD8[+] CTL in EX group), and targets are myeloid/B cell
subsets.

A

Participants' Characteristics at Baseline[a]

Characteristic	Physical Activity Pattern, No. (%) of Participants[b]			
	Total (n=86)	SED (n=46)	EX (n=40)	p value
Age, y, weighted mean (95% CI)	28.2 (27.2-29.2)	28.9 (27.5-30.3)	27.4 (26.0-28.8)	0.14
Sex, n (weighted %)				< 0.05
Male	28 (32.6)	9 (19.6)	19 (47.5)	
Female	58 (67.4)	37 (80.4)	21 (52.5)	
BMI, mean (SD)	21.3 (±3.0)	22.0 (±2.8)	20.7 (±3.0)	< 0.05
Lifestyle				
Smoker, n (weighted %)				0.26
Never	82 (95.3)	44 (95.7)	38 (95.0)	
Past	4 (4.7)	2 (4.3)	2 (5.0)	
Current drinker, n (weighted %)				0.15
Yes	16 (21.3)	5 (16.2)	11 (29.2)	
No	70 (78.7)	41 (83.8)	29 (70.8)	
Sleep duration, n (weighted %)				0.56
6-8h	72 (83.8)	40 (87.0)	32 (80.0)	
≥ 8h	14 (16.2)	6 (13.0)	8 (20.0)	
Dietary Habits, n (weighted %)				0.11
Balanced Diet	44 (51.1)	19 (41.3)	25 (62.5)	
Meat-Dominant Diet	12 (14.0)	9 (19.6)	3 (7.5)	
Other	30 (34.9)	18 (39.1)	12 (30.0)	
Medication Usage, n (weighted %)				0.08
No Medication	44(51.1)	28 (60.9)	16 (40.0)	
Used Medication	44(48.9)	18 (39.1)	24 (60.0)	

1366

Supplementary Figure 1. Baseline characteristics and analysis of blood biochemical indices of the study population.

(A) A table summarizing the baseline characteristics of participants (n=86) stratified by physical activity pattern . The table includes metrics such as age, sex, BMI, lifestyle

factors (smoking, drinking, sleep), dietary habits, and medication usage, with weighted

means/percentages and p-values indicating group differences.

(B) Box plots of blood metabolic indices between the EX and SED groups.Levels of

fasting glucose (GLU), triglycerides (TG), and high-density lipoprotein cholesterol (HDL-C)

are shown.

(C) Box plots of blood biochemical indices between the EX and SED groups. Levels of

creatinine (Cr), uric acid (UA), total bilirubin (TBIL), direct bilirubin (DBIL), and indirect

bilirubin (IBIL) are shown.

(D) Box plots showing the concentration of liver-related biochemical indicators (ALB:

albumin; ALT/AST: hepatocellular injury markers; AST/ALT ratio) between the SED (blue)

and EX (green) groups. The values above each plot denote p-values.

1382

Supplementary Figure 2. **Quality control, batch effect evaluation, and annotation**

validation for single-cell multi-omics data.

(A) Violin plots depicting key scRNA-seq quality control metrics (Gene Count, UMI Count,

Mitochondrial Gene Proportion) between the SED (blue) and EX (green) cohorts;

consistent distribution of these metrics verified homogeneous data quality across groups.

(B) Violin plots of scATAC-seq quality control indicators (TSS Enrichment Score, Number

of Fragments, NucleosomeSignal, BlacklistRatio) comparing the SED and EX cohorts.

(C) UMAP plots displaying cell distribution in scRNA-seq (top) and scATAC-seq (bottom)

datasets, with cells colored by group (SED/EX). Overlapping distribution of SED and EX

cells indicates no prominent batch effects.

(D) Heatmap of main marker genes based on gene activity scores from scATAC-seq data,

demonstrating high concordance between chromatin accessibility patterns and the

annotated cellular identities across 30 cell subsets.

(E) UMAP plots showing the expression levels of key marker genes (e.g., CD3D, LYZ) in

scRNA-seq data. The expression patterns align with the annotated cell subsets,

supporting the cell typing results.

(F) UMAP plots showing chromatin accessibility at the promoter regions of key marker

genes (e.g., CD3D, CD8A) in scATAC-seq data. The accessibility patterns match the

corresponding cell subsets, validating the consistency between scATAC-seq and

scRNA-seq annotations.

1403

Supplementary Figure 3. Visualization of marker genes: Dot plot from scRNA-seq

and gene activity score heatmap from scATAC-seq.

(A) Expression dot plot of classical   marker genes (e.g., CD14, FCGR3A, CD1C,

LILRA4) for Myeloid cell subsets.

(B) Expression dot plot of representative subset marker genes (e.g., CCR7, GZMB,

KLRB1, TRDC) for CD8$^+$ T cell and non-conventional T cell subsets.

(C) Expression dot plot of subset marker genes (e.g., FOXP3, IL7R, CXCR5, PRDM1) for

the seven CD4$^+$ T cell subsets.

(D) Expression dot plot of subset marker genes (e.g., NCAM1, KLRB1, FCER1A, IL1RL1)

for NK cell subsets.

(E) Expression dot plot of subset marker genes (e.g., IGHD, CD27, CD38, TNFRSF17)

for B cell subsets.

(F) Gene activity score (GAS) heatmaps for Myeloid cell subsets.

(G) GAS heatmaps for CD8$^+$ T&non-canonical T cells subsets.

(H) GAS heatmaps for CD4$^+$ T cells subsets.

(I) GAS heatmaps for NK cells subsets.

(J) GAS heatmaps for B cells subsets

1421

Supplementary Figure 4. Regulation of myeloid cell subset proportions, pathway activity, and chromatin accessibility by exercise.

(A) Box plots with overlaid scatter points showing the proportion of each myeloid cell subset (cDC, cMono, etc.) in the SED (blue) and EX (green) groups. "ns" indicates no significant difference between groups based on conventional testing.

(B) Bar chart showing the pathway activity scores for "GO:BP_ANTIGEN_PROCES

SING_AND_PRESENTATION" in each myeloid cell subset for the SED and EX gro

ups. The activity of this pathway was significantly upregulated in multiple subsets of

the EX group.

(C) Pathway activity scores for stress-related pathways. The left panel shows scores for

"HALLMARK_HYPOXIA" and the right panel shows scores for

"HALLMARK_REACTIVE_OXYGEN_SPECIES_PATHWAY". The activity of these two

pathways was generally downregulated across myeloid subsets in the EX group.

(D) Pie chart showing the distribution of differentially accessible regions (DARs) across

myeloid cell subsets.

(E) The chromatin accessibility at the differential accessible region (DAR) of

inflammation-related genes between the SED and EX groups.

(F) A network plot illustrating the interaction strength/probability between various immune

cell subsets (nodes: cell types; edges: interaction connections).

(G) A pair of cell-cell interaction network plots showing MHC-I-dependent signaling

between myeloid cells (cMono, ncMono, etc.) and CD8$^+$ T cell subsets (CD8_CTL,

CD8_Tn, etc.) in the SED group (left) and EX group (right).

1444

Supplementary Figure 5. Regulation of CD4⁺ /CD8⁺ T cell subset proportions and

functional pathways by exercise.

(A) UMAP plot displays scRNA-seq-derived CD4⁺ T cells after subclustering and

annotation, colored by five functional subsets(left). UMAP plot shows scATAC-seq-derived

CD4⁺ T cells, with annotation labels from corresponding scRNA-seq data and each cell

type color-marked(right).

(B) MiloR-based analysis was performed to assess differences in the abundance of CD4⁺

T cell subsets between the EX and SED groups.

(C) Bar charts show the pathway activity scores for "GO:BP_POSITIVE_REGULATI

ON_OF_TYPE_II_INTERFERON_PRODUCTION" and "HALLMARK_OXIDATIVE_PHO

SPHORYLATION" in various T cell subsets (e.g., CD8_CTL, MAIT) for the SED an

d EX groups.

(D) .Activity scores for stress-related pathways in T cells. The left panel shows sco

res for "HALLMARK_HYPOXIA" and the right panel shows scores for "HALLMARK_

INFLAMMATORY_RESPONSE".

(E) Motif bias analysis of key transcription factors in CD8 Tn(left), CD8 CDL(middle),

MAIT cells(right).

1462

Supplementary Figure 6. Effects of exercise on NK cell subset proportions,

functional pathways, and epigenetic regulation.

(A) Box plots with overlaid scatter points showing the proportion of each NK cell subset

(CD56_bright_NK, Mature_NK, etc.) in the SED (blue) and EX (green) groups. "ns"

indicates no significant difference. Asterisks denote significance of group

differences(*P<0.05,**P<0.01,***P<0.001).

(B) Multiple bar charts showing pathway activity scores for various NK cell subsets in the

SED and EX groups.

(C) Statistical table of differentially accessible regions (DARs) in NK cell subsets,

showing the number of upregulated (Up) and downregulated (Down) DARs in each

subset.

(D) DARs (e.g., immune defense, killing), and blue bars represent pathways associated

with downreGene Ontology (GO) pathway enrichment analysis of DARs in Mature_NK

cells. Red bars represent enriched pathways associated with upregulategulated DARs

(e.g., stress response, inflammation).

(E) Motif bias analysis of key transcription factors in Mature_NK cells. The plot shows the

distribution of $-\log_{10}$(p-value) versus the mean difference for TF motifs between the SED

and EX groups.

1481

Supplementary Figure 7. Effects of exercise on B cell subset proportions, functional pathways, and epigenetic regulation.

(A) UMAP plot displays scRNA-seq-derived B cells after subclustering and annotation, colored by five functional subsets(left). UMAP plot shows scATAC-seq-derived B cells, with annotation labels from corresponding scRNA-seq data and each cell type

color-marked(right).

(B) MiloR-based analysis was performed to assess differences in the abundance of B cell

subsets between the EX and SED groups.

(C) Gene Ontology (GO) pathway enrichment analysis of DARs in B cells. Red bars

represent enriched pathways associated with upregulated DARs (e.g., immune defense,

killing), and blue bars represent pathways associated with downregulated DARs (e.g.,

stress response, inflammation).

(D) Violin plots showing the chromatin accessibility of antigen presentation-related genes

(e.g., HLA-DQB1, CD86) in various B cell subsets between the SED (blue) and EX (green)

groups; Log2FC values indicate fold changes in accessibility.

(E) Heatmap showing the number of upregulated (Up) and downregulated (Down)

differentially accessible regions (DARs) in each B cell subset from scATAC-seq data.

(F) GO pathway enrichment analysis of upregulated signaling pathways in B cells.

(G) Transcription factor (TF) binding motif enrichment analysis of upregulated DARs in

Naive_B cells of the EX group; the y-axis denotes -log10(p-adj) (motif enrichment

significance), and the x-axis represents rank-sorted enriched TFs.

(H) Left: TF binding motif enrichment analysis of upregulated DARs in Switched_Bm cells

of the EX group. Right: Violin plots showing the motif deviation of key TFs in Switched_Bm

cells between the SED and EX groups; Padj values indicate significance of group

differences.

(I) The chromatin accessibility at the promoter DAR of HLA-DQB1 between the SED and

EX groups in Switched_Bm cells; the track includes peak-gene links for the gene.

(J) The chromatin accessibility at the promoter DAR of HLA-DQA1 between the SED

and EX groups in Switched_Bm cells.

(K) The chromatin accessibility at the promoter DAR of CCL2 between the SED and EX

groups in Switched_Bm cells.

1513

Supplementary Figure 8. Immune cell communication and pathway remodeling in the EX group

(A) Heatmap showing normalized module scores of functional pathways (e.g., TNFA Signaling via NFKB, MHC class II antigen presentation) across myeloid cell subsets (cDC, cMono, etc.) between the SED (blue) and EX (green) groups; asterisks denote significance of group differences (***P<0.001).

(B) Cell-cell interaction network plots of RESISTIN-mediated signaling in the SED (left) and EX (right) groups; nodes represent cell types, and edges represent interaction connections, centered on cMono.

(C) MHC-II-dependent cell communication network plots in the SED (left) and EX (right)

groups; networks show interactions between B cells/Myeloid cells and CD4+ T cell

subsets, centered on cMono.

(D) MHC-I-dependent cell communication network plots in the SED (left) and EX (right)

groups; networks show interactions between various cell types and CD8+ T cell subsets,

centered on cMono.

(E) COLLAGEN-mediated cell communication network plots in the SED (left) and EX

(right) groups; networks show interactions between Myeloid/NK cells and B/T cells,

centered on Unconv_T cells.

(F) Heatmap showing communication probability of the COL6A2-CD44 ligand-receptor

pair (COLAGEN pathway) across cell type interactions in the SED and EX groups; color

intensity represents communication probability.

(G) Violin plot showing the expression level of TCOL6A2 in Unconv_T cells between the

SED and EX groups.

(H) CCL-mediated cell communication network plots in the SED (left) and EX (right)

groups; networks show interactions between Myeloid/B cells and T cells, centered on

Unconv_T cells.

(I) Heatmap showing communication probability of CCL5-CCR1 and CCL3-CCR1

ligand-receptor pairs (CCL pathway) across cell type interactions in the SED and EX

groups; color intensity represents communication probability, and black dots denote

significance (p<0.01).