Geographical gradients in leaky sex expression and reproductive effort in a dioecious plant

are consistent with selection during range expansion

M. T. Nguyen* and J. R. Pannell

Department of Ecology and Evolution

University of Lausanne

1015 Lausanne

Switzerland

* Author for correspondence: maithu.nguyen@unil.ch

Keywords: colonisation, self-fertilisation, reproductive assurance, Baker' Law, life-history trade-

off, r-selection, ruderal life-history strategy, dispersal

Acknowledgements

We thank Kai-Hsiu Chen, Shengman Lyu, Antoine Guissan, Valentin Verdon for their comments

on our analysis; Leonardo Cantoro, Sebastian Cato, Theophanis Constantinidis, Stefano Doglio,

Galina Gavrilenco, Thijs Heijman, Gabriele La Grasta, Ramy Maalouf, Arian Ndoni, Thomas

Schorr, Gertraud Seiser, Kathy Vivian, Jean-Paul Wolff, Walter Wimmer for their help with seed

sampling; Jessica Bainbridge, Kai-Hsiu Chen, Camilo Ferron Martinez, Ivan Giordano, Xia

Jiang, Ehouarn Le Faou, Marjolaine Rousselle, Philipp Schenkel, as well as numerous

undergraduate assistants, for their help in setting up experiments and measuring plant

phenotypes; and the Swiss National Science Foundation (Grant 31003A_163384) and the

University of Lausanne for funding.

Author contributions

Conceived the project and designed the experiment: JRP and MTN; conducted the experiment

and analyzed the data: MTN; revised and wrote the final version of the manuscript: MTN and

JRP. Both authors approved the final version of the manuscript.

Conflict of interest

We have no conflict of interest to declare.

Data availability statement

The scripts and data are provided on https://github.com/maithunguyen/Leakiness-sex-

expression-and-reproductive-effort-Mercurialis-annua-range-expansion.

Abstract

Range expansion involves repeated dispersal and the establishment of new populations through

colonization. Natural selection is therefore expected to favour traits that enhance dispersal and

enable uniparental reproduction when mates are scarce. It is known that range-edge populations

may evolve increased dispersal and, in hermaphroditic species, an enhanced capacity for self-

fertilization. Here, we report results from a large common-garden experiment showing that

males and females of the dioecious annual herb *Mercurialis annua* are more likely to show

'leaky' or inconstant sex expression towards the recently colonized edge of its geographical

distribution in Europe than those from the eastern Mediterranean Basin, from where the species

expanded its range. Range-edge populations also had greater reproductive effort (RE). Leaky sex

expression allows otherwise unisexual individuals of *M. annua* to produce progeny by self-

fertilization, a trait that may have been selected for reproductive assurance during the range

expansion. High RE likely facilitated both colonization and subsequent population

establishment. Our results show that selection during range expansions may modify reproductive

traits towards range margins, extending this notion to the phenomenon of leaky sex expression in

dioecious species. Previous work has shown that leaky sex expression in *M. annua* is strongly

heritable; our results now further suggest that has been enhanced by selection for reproductive

assurance towards range margins and confirm, in a natural setting, that the trait is not just an

expression of developmental instability during floral development or a relic of an incomplete

transition to dioecy.

Introduction

The current distributions of many species, especially at mid to high latitudes, are the legacy of

past range expansions, often from Pleistocene refugia (Hewitt, 2000; Svenning & Skov, 2007).

Such expansions leave characteristic population-genetic signatures, both because populations

encounter new abiotic and biotic environments and because repeated colonization entails serial

bottlenecks (Bridle & Vines, 2007; Sexton *et al.*, 2009). These bottlenecks increase drift and

progressively erode genetic diversity with distance from the ancestral refugium, affecting both

neutral and fitness-related variation (Ibrahim *et al.*, 1996; Hewitt, 2000; Excoffier *et al.*, 2009).

As a result, range expansion can lead to the accumulation of deleterious mutations, changes in

homozygosity and inbreeding depression, and a reduced efficacy of selection towards the range

edge (Pujol *et al.*, 2009; Peischl *et al.*, 2013, 2015; González-Martínez *et al.*, 2017; Willi *et al.*,

2018). At the same time, colonization itself should favour phenotypes that enhance

establishment and spread, so that populations towards a species' range margin may evolve traits

distinct from those in the core range and former refugia. The evolution of such traits is likely to

be an important component of adaptation at range margins.

A key idea is that individuals capable of uniparental reproduction, whether by asexual means

(Delnevo *et al.*, 2024) or self-fertilization (Pannell, 2015), should be more likely to succeed in

colonizing new habitats than those that need to find a mate to reproduce. Baker (1955) linked

observations of an increased capacity for self-fertilization in species on oceanic islands to

colonization following long-distance dispersal to islands. This enrichment on islands of species

with a capacity to self-fertilize as a result of selection for reproductive assurance was later

labelled Baker's Law by Stebbins (1957), highlighting its broad relevance across both plants

(Baker, 1955, 1967) and animals (Longhurst, 1955; Stebbins, 1957). Substantial further evidence

for Baker's Law has accumulated since Baker's initial paper (Grossenbacher *et al.*, 2017), and

the concept has been expanded to explain the evolution of a capacity for uniparental

reproduction in species with a colonizing habit in metapopulations with frequent population

turnover (Pannell, 1997, 2000; Pannell & Barrett, 1998) as well as in the context of species range

expansions (reviewed in Cheptou, 2012; Pannell, 2015; Pannell *et al.*, 2015). In

metapopulations, higher extinction rates and greater patchiness are associated with increased

frequencies of self-fertilizing individuals in species with mixed mating, including *Silene vulgaris*

(Taylor *et al.*, 1999), *Helicteres brevispira* (Franceschinelli & Bawa, 2000), *Eichhornia*

paniculata (Barrett & Husband, 1990; Ness *et al.*, 2010) and *Mercurialis annua* (Obbard *et al.*,

2006b). In the context of range expansion, autonomously apomictic species tend to occupy

broader distributions than their sexually reproducing relatives (Hörandl *et al.*, 2008; Cosendai *et*

al., 2013; Hörandl, 2023), and marginal populations are more likely to possess the ability to

reproduce by self-fertilization (Lloyd, 1965; Moeller & Geber, 2005; Obbard *et al.*, 2006b;

Busch & Delph, 2012; Ruane *et al.*, 2020). For example, range-margin populations of *Clarkia*

xantiana ssp. *parviflora* show significantly lower herkogamy, petal area, and protandry than core

populations, indicating increased selfing ability, a pattern further supported by emasculation

experiments (Ruane *et al.*, 2020). Similarly, in *Leavenworthia* (Cruciferae), floral morphology

and extensive field surveys suggest a breakdown of self-incompatibility in range-marginal

populations (Lloyd, 1965).

Selection on traits that affect colonization ability should also apply to dioecious species, as

originally discussed by Baker (1967). In the context of Baker's Law, the frequently observed

enrichment of dioecious species on islands (Carlquist, 1965, 1967) might appear surprising,

given that dioecy implicates biparental rather than uniparental reproduction. However, Baker

(1967) already drew attention to the fact that the males and females or many dioecious plant

species produce occasional flowers with the opposite sexual function, potentially allowing them

to self-fertilize. This phenomenon, referred to as inconstancy or 'leakiness' in sex expression

(e.g., Baker, 1967; Lloyd & Bawa, 1984; Korpelainen, 1998), is common in dioecious species

(Ehlers & Bataillon, 2007; Cossard & Pannell, 2019). Leakiness in sex expression has tended to

be attributed to developmental noise (Lloyd & Bawa, 1984), a plastic response to environmental

cues such as temperature, drought, herbivory (Charnov & Bull, 1977; Freeman *et al.*, 1980; Case

& Barrett, 2001, 2004; Delph, 2003; Delph & Wolf, 2005; Golenberg & West, 2013; Villamil *et*

al., 2022), or as evidence of still incomplete transitions from combined to separate sexes

(Charlesworth & Charlesworth, 1978; Lloyd, 1980; Ross, 1980, 1982). Ehlers and Bataillon

(2007) and Crossman and Charlesworth (2014) modelled the evolution and maintenance of

leakiness in terms of a response to selection for reproductive assurance, and Cossard et al. (2021)

and Gerchen et al. (Gerchen *et al.*, 2026) have recently demonstrated that variation in leakiness

can have a genetic basis and be responsive to selection. However, empirical evidence that

leakiness can be promoted specifically by selection for reproductive assurance such as in

metapopulations or range expansions is still lacking.

Selection during range expansion might also bring about evolution of dispersal and life-history

traits that promote both colonization and rapid establishment (reviewed in Hargreaves & Eckert,

2014; Chuang & Peterson, 2016; Weiss-Lehman & Shaw, 2022). Increased dispersal may be

selected directly through the process of colonization itself (Travis & Dytham, 1999) and may be

further reinforced by assortative mating among highly dispersive individuals at the expansion

front (Shine *et al.*, 2011). A notable example is provided by the invasive cane toad *Rhinella*

marina in Australia, which has evolved longer legs enabling faster dispersal at the invasion front

(Phillips *et al.*, 2006, 2007). In plants, dispersal-related traits such as enhanced seed pappus or

wing development have been observed in recently colonised compared to older populations

(Cody & Overton, 1996; Cheptou *et al.*, 2008; Darling *et al.*, 2008; Monty & Mahy, 2010), a

phenomenon that has been labelled the 'metapopulation effect' (Olivieri & Gouyon, 1997) but

that also applies to range expansions. For instance, the ragwort *Senecio inaequidens*

exhibits increased pappus size, enhancing wind-dispersal distance toward its range margin

(Monty & Mahy, 2010), and the dune plant *Abronia umbellata* shows greater seed wing index in

more marginal populations, a legacy of selection on colonization ability (Darling *et al.*, 2008).

Successful population establishment then should be enhanced greater allocation to reproduction,

at the expense of traits associated with persistence and competitive ability in crowded

environments, such as vegetative growth, size and competitive performance (Tilman, 1990;

Burton *et al.*, 2010; Urquhart-Cronish *et al.*, 2024). The expected shifts along this trade-off with

range expansion accord, more generally, with classic life-history ideas, including the

colonization-competition trade-off (Tilman, 1990), the r/K continuum (MacArthur & Wilson,

1967; Pianka, 1970; Reznick *et al.*, 2002; Travis *et al.*, 2023), and Grime's C–S–R framework

(Grime, 1974), which predict that populations experiencing repeated colonization should evolve

traits associated with more ruderal, colonizing strategies (MacArthur & Wilson, 1967; Pianka,

1970; Grime, 1974; Stearns, 1983, 1989; Reznick *et al.*, 2002; Travis *et al.*, 2023). Reproductive

effort, the proportion of biomass allocated to flowering and fruiting, is an easily measurable trait

corresponding to this trade-off that should accordingly increase with range expansion.

Here, we test the hypothesis that selection should have favoured increased leakiness in sex

expression and greater reproductive effort in populations towards the margins of the geographic

range of the diploid European annual dioecious plant, *Mercurialis annua* L. (Euphorbiaceae). *M.*

annua is a wind-pollinated annual colonizer of disturbed habitat that expresses leakiness in sex

expression in natural populations for both males and females. The rapid evolution of enhanced

leakiness in experimental populations (Cossard *et al.*, 2021; Gerchen *et al.*, 2026) indicates that

the trait is strongly heritable. Our study involved sampling dioecious *M. annua* from populations

across its geographic range in Europe, growing them in a common garden, and measuring the

size of plants and their allocation to male and female flowers and fruits. We used our measures

to quantify the leakiness in sex expression and the reproductive effort of all individuals sampled.

Leakiness can be measured both by its incidence (the proportion of individuals producing

opposite-sex flowers) and its degree (the number of opposite-sex flowers produced per

individual) (Cossard & Pannell, 2019). We expected reproductive assurance to favour increased

incidence of leakiness in both sexes with distance from the 'expansion origin' (see Methods for

its definition), but an increased degree of leakiness only in males. This is because a mildly leaky

female could self-fertilize many ovules and establish a population, whereas the success of a

colonizing male would depend on it producing many seeds, requiring a substantial degree of

leakiness. An greater degree of leakiness in males than females is also expected as an outcome of

selection towards female-biased sex allocation due to local mate competition (Hamilton, 1967;

Roux *et al.*, 2024).

Methods

<u>Study species and population sampling</u>

Mercurialis annua L. (Euphorbiaceae) is a European, dioecious, wind-pollinated annual plant

with a distribution from the eastern Mediterranean through central and western Europe to the

Atlantic Ocean. The species occupies disturbed ruderal sites and has a life-history typical of

weedy colonizing annuals and a demography characterised by metapopulation dynamics

(Obbard *et al.*, 2006b; Eppley & Pannell, 2007b; Pannell *et al.*, 2008; Pujol *et al.*, 2009; Pannell

et al., 2014; Dorken *et al.*, 2017; González-Martínez *et al.*, 2017). Populations of *M. annua* in

western Europe have lower within-population genetic diversity than those in the east (Obbard *et*

al., 2006b) and show signs of increased genetic load and with fewer genomic signatures of

selective sweeps than those in the east (González-Martínez *et al.*, 2017), consistent with a range

expansion into western Europe from a Pleistocene refugium in the eastern Mediterranean Basin.

M. annua possesses heteromorphic sex chromosomes with an XY sex-determination system

dated back to over 1 MY ago (Russell & Pannell, 2015; Veltsos *et al.*, 2018, 2019; Gerchen *et*

al., 2022) and strong sexual dimorphism (Harris & Pannell, 2008; Hesse & Pannell, 2011;

Sánchez Vilas & Pannell, 2011; Tonnabel *et al.*, 2019). It also shows substantial leakiness in sex

expression in both sexes (Cossard & Pannell, 2019; Cossard *et al.*, 2021). Flowering in *M. annua*

is indeterminate. Females and males of *M. annua* produce pistillate flowers and staminate

inflorescences, respectively, in the axis of each new leaf. Leaky females usually produce one or

more staminate flowers alongside its axillary female flowers. Leaky males produce pistillate

flowers and fruits along with their staminate flowers on extended inflorescence stalks

('peduncles'). Both males and females with leaky sex expression are capable of self-fertilizing

(Eppley & Pannell, 2007a; Li *et al.*, 2019; Cossard *et al.*, 2021) and thus enjoy reproductive

assurance in the absence of mates of the opposite sex, e.g., during colonization or in sparse

stands (Baker, 1967; Pannell *et al.*, 2008, 2015; Cheptou, 2012; Cossard *et al.*, 2021).

For our study, we collected seeds, or had them sent to us, from 26 populations across the

distribution of diploid *M. annua* in autumn 2023 and spring 2024 (Figure 1, Table S1). For each

population, seeds were bulked from at least 20 females, except for the Albanian population,

where only five individuals could be sampled due to its small size. In autumn 2024, we sowed

seeds from all 26 populations in a peat-based substrate in trays set up on a glasshouse at the

University of Lausanne, Switzerland. After one month, when most plants had begun flowering,

we transplanted between 80 and 120 individuals per population into 0.68 L pots filled with

Ricoter 140 soil (2,694 plants in total). Exceptions were Populations 1 and 8, where only 43 and

55 plants germinated, respectively, and all of these were used.

<u>Experimental treatments and design of the common garden</u>

Sex expression in *M. annua* is mediated by the plant hormones auxin (masculinizing) and

cytokinin (feminizing) (Louis & Durand, 1978; Dauphin-Guerin *et al.*, 1980; Hamdi *et al.*, 1987;

Li *et al.*, 2019), and exogenous application of these hormones can bring about altered sex

expression, with cytokinin application causing males to produce female flowers and fruits, and

auxin causing females to produce male flowers. Because we were concerned that we might have

limited power to test our main hypothesis about leakiness if too few plants expressed it, to

amplify levels of leakiness we applied hormone treatments to half of the plants in 24 of the

populations (we excluded the two populations with lower plant numbers, i.e., these were not

treated). Specifically, in addition to automatic watering, half of the males and half the females,

respectively, received 55 mL of cytokinin and 55 ml of auxin twice a week; control plants

received 55 mL of water. Following Louis and Durand (1978), we used 2 µM BAP for the

cytokinin treatment and a mixture of 1×10^{-3} g/L IAA and 1×10^{-2} g/L GA$_\square$ for the auxin

treatment. We were not specifically interested in the effect of the hormone treatment, but rather

in assessing differences in leakiness among populations. As described in the Results, the

hormone treatments ultimately had no effect on flower production in either sex.

Imposition of hormone treatments necessitated separating males and females into different

glasshouses. Our design thus effectively involved two independent experiments, one for each

sex. We were not interested in a direct sex comparison, and we measured a different set of

variables on each sex. In each glasshouse, plants were randomly distributed across seven tables

in a random block design. Randomization ensured a balanced distribution of plants from each

population and treatment across tables.

235
<u>Measurements of phenotypic variation</u>

After three weeks of growth following the onset of the hormone treatments, we harvested plants

table by table in each glasshouse, retaining table number and harvest time as a combined

blocking unit for analysis. For most tables, all plants were harvested within two days of each

other. Because of the indeterminate nature flowering in *M. annua*, our harvest represents a

temporal snapshot of reproductive allocation at the time of sampling. For each plant, we

recorded the number of flowers of the opposite sex (i.e., fruits on males and male flowers on

females), plant height and above-ground biomass. Additionally, for a subset of five plants of

each sex, population and hormone treatment (i.e., 5 x 2 x 25 x 2 for a total of 500 plants), we

counted the number of female fruits on females and male flowers on males.

We measured the incidence of leakiness in a population in terms of the proportion of plants of

each sex that produced at least one flower of the opposite sex. We followed Cossard and Pannell

(2019) to compute the degree of leakiness for females as $d_f=\lambda_f/M$, where λ_f is the given female's

male flower production per biomass unit (i.e., its male RE), and M is the male reproductive

effort of an average male from the same population. Due to phenotyping constraints, M for each

population was calculated as the average of five untreated males selected for detailed

phenotyping. The degree of leakiness for males was computed as $d_m=\lambda_m/F$, where λ_m and F are

defined analogously to λ_f and M, respectively. Dividing λ by M and F served to calibrate the

degree of leakiness in one sex in relation to the typical production of flowers by plants of the

opposite sex, also ensuring that values fall within the range of 0 to 1.

We calculated an index of reproductive effort, defined as the number of male flowers on a male

or the number of female fruits on a female divided by its respective biomass. The reproductive

effort of plants without hormone treatment was used to calculate the population average male or

female reproductive effort used to compute the degree of leakiness.

<u>Statistical analysis</u>

We constructed generalized linear mixed models (GLMMs) and linear models to assess the

dependence of seven plant traits on distance from the assumed point of departure for the range

expansion, which we refer to as the 'expansion origin'; we set this origin as the midpoint

between the two southeastern-most populations sampled (Figure 1, Table S1). The traits were:

plant biomass, plant height, incidence of leakiness, degree of leakiness, and reproductive effort.

We computed each site's distance from the expansion origin (Table S1) using geodesic distance

(Vincenty, 1975).

To analyse trait variation recorded in each of the (male and female) common gardens, we used

GLMMs with a consistent fixed and random effect structure, differing only in response

variables, error distribution, offset terms, and zero-inflation components, when applicable. We

included distance from the expansion origin, hormone treatment (binary), and the interaction

between distance and hormone treatment as fixed effects and population and table number (with

sampling time) as random effects.

We modelled the incidence of leakiness as a binary variable and the degree of leakiness using a

zero-inflated negative binomial GLMM, with counts of leaky flowers (males) or fruits (females)

as the response variable, offset by the log-transformed product of plant biomass and population-

average reproductive effort of the opposite sex. The zero-inflation component included distance

from the expansion origin as a fixed effect, based on results from the binomial model.

Reproductive effort was modelled with a negative binomial GLMM, using counts of fruits or

male flowers, offset by the logarithm of biomass. Plant height and biomass were modelled with

Gaussian GLMMs; biomass was square-root-transformed.

Our model for female biomass as a function of geodesic distance failed to converge. To address

this issue, we removed greenhouse table from the random effects, retaining population as the

only random effect. This approach was taken because variance among tables in female biomass

was effectively zero, while the variance among populations was substantial. We also compared

the AIC of two models, each including only one random effect, either table or population. The

model with population had a lower AIC (10,300) than the model with table number (11,200),

further supporting our choice to remove table number from the final model.

All analyses were performed in R (R Core Team, 2025). All GLMMs and LMs were fitted using

the 'glmmTMB' function from the 'glmmTMB' package in R (Brooks *et al.*, 2017). We checked

the diagnostic fit of all chosen models for significant deviations, overdispersion, and zero-

inflation (when applied) using the 'DHARMa' package (Florian, 2016) in R, with 5,000

simulation points.

Results

Effect of hormone treatment on reproductive and vegetative traits

We found no effect of the hormone treatment on any of the reproductive traits studied, including

leakiness ability, leakiness degree, or reproductive effort in either sex (Table S2). Hormone

treatment did, however, affect plant biomass ($Z = 9.49$, $P < 0.001$ for females; $Z = 3.57$, $P <$

0.001 for males) and height ($Z = 7.57$, $P < 0.001$ for females; $Z = 4.80$, $P < 0.001$ for males), as

well as significant interactions between plant size and distance from the expansion origin (Table

S2). As the effect of hormones on plant size was not the focus of our study, we do not analyze

plant size measurements further, except in terms of its role in the computation of reproductive

effort and its variation with respect to distance from the expansion origin (Table S2).

Greater leakiness with distance from the expansion origin

The incidence of leakiness increased in both sexes with distance from the expansion origin

(females: ~0.012 % per km from the expansion origin, $Z = 4.29$, $P < 0.001$; males: ~0.010 % per

km from the expansion origin, $Z = 2.38$, $P < 0.05$; Figure 2A, Table S2). This trend was further

reflected by the zero-inflation components of models for the degree of leakiness, with the

proportion of excess zeros (i.e., non-leaky individuals) declining significantly with increasing

distance from the expansion origin (females: $Z = -4.76$, $P < 0.001$; males: $Z = -3.29$, $P < 0.01$;

Table S2). The degree of leakiness showed no significant association with the distance from the

expansion origin for either sex (Table S2).

Variation in vegetative traits with distance from the expansion origin

There was no significant association between the distance from the expansion origin and

vegetative traits (biomass or height) in either sex (Table S2).

Greater reproductive effort with distance from the expansion origin

Reproductive effort in both sexes increased with distance from the expansion origin (females:

~8.17 x 10^{-5} flowers/g per km from the expansion origin, $Z = 2.53$, $P < 0.05$; males: ~6.20 x 10^{-5}

flowers/g per km from the expansion origin, $Z = 1.96$, $P = 0.0502$; Figure 2B, C, Table S2).

Discussion

Our study reveals that both the incidence of leaky sex expression and reproductive effort in

dioecious *M. annua* increase with distance from the species' expansion origin in the eastern

Mediterranean Basin and towards its geographical range margin in northern and western Europe.

Both these trends are consistent with expectations for the effects of selection during a past range

expansion. Of course, our results are based on correlations, not a controlled experiment, so that

any causal inference needs to be cautious; in principle, other known or unknown factors could be

responsible for the non-random patterns observed. For instance, the geographic gradient from

southeastern to northwestern Europe coincides with substantial and partially directional changes

in climate, and the observed increases in leakiness and reproductive effort along this gradient are

also strongly associated with variation in climatic variables (see Supporting Information B,

Table S4). Nevertheless, it is difficult to see why a temperate climate should have selected for

increased leakiness. We thus discuss our results here in terms of the hypotheses that motivated

our study, though the reader should bear in mind the correlational nature of our evidence.

Increased leakiness in sex expression with range expansion

We found a significantly greater incidence of leakiness in sex expression in both males and

females of *M. annua* at sites further from the expansion origin, consistent with the hypothesis

that selection for reproductive assurance during repeated colonization has contributed to

phenotypic evolution at the species' range margin. Baker (1955) first suggested that the enriched

capacity for uniparental reproduction on oceanic islands could be a result of selection for

reproductive assurance during colonization, an idea also invoked for the evolution of a selfing

ability in metapopulations (Pannell & Barrett, 1998) and range expansions (Cheptou, 2012;

Pannell, 2015; Pannell *et al.*, 2015). Our results here provide empirical support for this idea in

the context of the selection of a capacity for self-fertilization in dioecious species via leaky sex

expression.

We also expected males to show a greater increase in the degree of leakiness from the expansion

origin than females, because the success of colonization by single males should be enhanced by

the production of larger numbers of female flowers and seeds, whereas a single female should

only produce as many male flowers as necessary to fertilize her ovules. However, our results

revealed no such difference. It is possible that the advantages of a high degree of leakiness in

males during colonization are countered by the disadvantages associated with producing many

seeds in established populations. Because males disperse a great deal of pollen to the wind, any

ovules they produce are likely to be self-fertilized, even in established populations, and self-

fertilized seeds might suffer inbreeding depression (Charlesworth & Charlesworth, 1987; Pannell

& Jordan, 2022).

While Baker (1967) and others (Ehlers & Bataillon, 2007; Crossman & Charlesworth, 2014;

Cossard & Pannell, 2019; Cossard *et al.*, 2021) have recognized that leaky sex expression in

dioecious species should facilitate their colonization potential, our study now provides strong,

albeit correlational, evidence that selection during repeated colonization may have enhanced this

tendency in a wild plant species. An important implication is that leaky sex expression found in

natural populations of dioecious species may be actively maintained by natural selection during

occasional periods where mates are limited, either in low-density populations or during

colonization. In other words, leaky sex expression in dioecious plants may not always be just a

relic of an unfinished transition to fully separate sexes from hermaphroditism or monoecy

(Lloyd, 1980) or an expression of developmental noise (Lloyd & Bawa, 1984). Indeed, dioecy in

the genus *Mercurialis* is ancestral, not derived (Krähenbühl *et al.*, 2002; Obbard *et al.*, 2006a),

and the evolution of leaky sex expression in *M. annua* can almost certainly be regarded as an

evolved adaptation to its colonizing habit that coincided a transition to an annual life history

(Pannell *et al.*, 2008). This adaptation appears to have been enhanced not only by ongoing

colonization in dynamic metapopulations (Obbard *et al.*, 2006b; Eppley & Pannell, 2007b;

Dorken *et al.*, 2017) but also by range expansion.

<u>Increased reproductive effort with range expansion</u>

We found that both male and female reproductive effort (RE) were higher toward the western

range margins of *M. annua*, with female RE showing a stronger increase with distance from the

expansion origin than male RE. To this extent, our results strengthen other correlative evidence

for selection of traits associated with a ruderal life history during colonization in range

expansions (Cwynar & MacDonald, 1987; Phillips *et al.*, 2006; Darling *et al.*, 2008; Monty &

Mahy, 2010; Bufford & Hulme, 2021). Note that we did not find a corresponding decrease in

biomass or height, as expected for a scenario of reduced competition during range expansion.

Given that *M. annua* populations grow quickly after colonization (Dorken *et al.*, 2017) and

plants in established dense populations must be highly competitive, this result is perhaps not

surprising. Trade-offs in competitive ability likely involve multiple traits (Stearns, 1983, 1989)

and may produce patterns that are too weak to detect in height or biomass alone.

<u>Concluding remarks</u>

Our results suggest that selection during range expansion has left a detectable phenotypic signature in *Mercurialis annua*, favouring traits likely to enhance colonization success. Populations further from the inferred expansion origin showed a higher incidence of leaky sex expression in both sexes, consistent with the idea that reproductive assurance can be strengthened during repeated colonization, even in a dioecious species. They also showed greater reproductive effort, in line with broader life-history expectations that expansion-front populations should evolve more ruderal, colonization-oriented strategies. Although our study is correlational and cannot exclude alternative explanations, the patterns we document are nevertheless consistent with long-standing ideas derived from Baker's Law (Baker, 1955, 1967; Stebbins, 1957; Pannell & Barrett, 1998; Cheptou, 2012; Pannell, 2015; Pannell *et al.*, 2015), life-history theory (MacArthur & Wilson, 1967; Pianka, 1970; Grime, 1974; Stearns, 1983, 1989; Reznick *et al.*, 2002; Travis *et al.*, 2023), and models of evolution during range expansion (e.g., reviewed in Hargreaves & Eckert, 2014; Chuang & Peterson, 2016; Weiss-Lehman & Shaw, 2022). More generally, our findings suggest that the evolutionary processes associated with colonization may contribute to shaping phenotypes at species' range margins, where mate limitation and demographic stochasticity may be pronounced. Accordingly, the evolution of traits that conferred reproductive assurance and establishment success on a species during a range expansion may then also influence not only its population and metapopulation dynamics at the range margin but also its capacity to move beyond the range margin, e.g., when a warming climate permits further range expansion.

References

Baker, H.G. 1955. Self-compatibility and establishment after "long-distance" dispersal. *Evolution* **9**: 347–349.

Baker, H.G. 1967. Support for Baker's Law - As a rule. *Evolution* **21**: 853–856.

Barrett, S.C.H. & Husband, B.C. 1990. Variation in outcrossing rates in Eichhornia paniculata: The role of demographic and reproductive factors. *Plant Species Biol.* **5**: 41–55.

Bridle, J.R. & Vines, T.H. 2007. Limits to evolution at range margins: When and why does adaptation fail? *Trends Ecol. Evol.* **22**: 140–147.

Brooks, M.E., Kristensen, K., van Benthem, K.J., Magnusson, A., Berg, C.W., Nielsen, A., *et al.* 2017. glmmTMB balances speed and flexibility among packages for zero-inflated

generalized linear mixed modeling. *R J.* **9**: 378–400. R Foundation for Statistical
Computing.

Bufford, J.L. & Hulme, P.E. 2021. Seed size–number trade-offs are absent in the introduced
range for three congeneric plant invaders. *J. Ecol.* **109**: 3849–3860.

Burton, O.J., Phillips, B.L. & Travis, J.M.J. 2010. Trade-offs and the evolution of life-histories
during range expansion. *Ecol. Lett.* **13**: 1210–1220.

Busch, J.W. & Delph, L.F. 2012. The relative importance of reproductive assurance and
automatic selection as hypotheses for the evolution of self-fertilization. *Ann. Bot.* **109**:
553–562.

Carlquist, S.J. 1965. *Island life: A natural history of the islands of the world*. Garden City,
N.Y.: Published for the American Museum of Natural History [by] the Natural History
Press.

Carlquist, S.J. 1967. The biota of long-distance dispersal. V. Plant dispersal to Pacific Islands.
Bull. Torrey Bot. Club **94**: 129–162.

Case, A.L. & Barrett, S.C.H. 2001. Ecological differentiation of combined and separate sexes of
Wurmbea dioica (Colchicaceae) in sympatry. *Ecology* **82**: 2601–2616.

Case, A.L. & Barrett, S.C.H. 2004. Environmental stress and the evolution of dioecy: *Wurmbea*
dioica (Colchicaceae) in Western Australia. *Evol. Ecol.* **18**: 145–164.

Charlesworth, D. & Charlesworth, B. 1987. Inbreeding depression and its evolutionary
consequences. *Annu. Rev. Ecol. Evol. Syst.* **18**: 237–268.

Charlesworth, D. & Charlesworth, B. 1978. Population genetics of partial male-sterility and the
evolution of monoecy and dioecy. *Heredity* **41**: 137–153.

Charnov, E.L. & Bull, J. 1977. When is sex environmentally determined? *Nature* **266**: 828–830.

Cheptou, P.-O. 2012. Clarifying Baker's Law. *Ann. Bot.* **109**: 633–641.

Cheptou, P.-O., Carrue, O., Rouifed, S. & Cantarel, A. 2008. Rapid evolution of seed dispersal
in an urban environment in the weed *Crepis sancta*. *Proc. Natl. Acad. Sci.* **105**: 3796–
3799.

Chuang, A. & Peterson, C.R. 2016. Expanding population edges: Theories, traits, and trade-offs.
Glob. Change Biol. **22**: 494–512.

Cody, M.L. & Overton, J.McC. 1996. Short-term evolution of reduced dispersal in island plant
populations. *J. Ecol.* **84**: 53–61.

Cosendai, A.-C., Wagner, J., Ladinig, U., Rosche, C. & Hörandl, E. 2013. Geographical
parthenogenesis and population genetic structure in the alpine species *Ranunculus*
kuepferi (Ranunculaceae). *Heredity* **110**: 560–569.

Cossard, G.G., Gerchen, J.F., Li, X., Cuenot, Y. & Pannell, J.R. 2021. The rapid dissolution of
dioecy by experimental evolution. *Curr. Biol.* **31**: 1277-1283.e5.

Cossard, G.G. & Pannell, J.R. 2019. A functional decomposition of sex inconstancy in the
dioecious, colonizing plant *Mercurialis annua*. *Am. J. Bot.* **106**: 722–732.

Crossman, A. & Charlesworth, D. 2014. Breakdown of dioecy: Models where males acquire
cosexual functions. *Evolution* **68**: 426–440.

Cwynar, L.C. & MacDonald, G.M. 1987. Geographical variation of lodgepole pine in relation to
population history. *Am. Nat.* **129**: 463–469.

Darling, E., Samis, K.E. & Eckert, C.G. 2008. Increased seed dispersal potential towards
geographic range limits in a Pacific coast dune plant. *New Phytol.* **178**: 424–435.

Dauphin-Guerin, B., Teller, G. & Durand, B. 1980. Different endogenous cytokinins between
male and female *Mercurialis annua* L. *Planta* **148**: 124–129.

Delnevo, N., Binks, R.M., van Leeuwen, S., Coates, D.J., McArthur, S., Macdonald, B.M., *et al.*
2024. Apomixis goes a long way: Genetic evidence of persistence and long-distance seed
dispersal in an ancient landscape. *J. Biogeogr.* **51**: 694–709.

Delph, L.F. 2003. Sexual dimorphism in gender plasticity and its consequences for breeding
system evolution. *Evol. Dev.* **5**: 34–39.

Delph, L.F. & Wolf, D.E. 2005. Evolutionary consequences of gender plasticity in genetically
dimorphic breeding systems. *New Phytol.* **166**: 119–128.

Dorken, M.E., Freckleton, R.P. & Pannell, J.R. 2017. Small-scale and regional spatial dynamics
of an annual plant with contrasting sexual systems. *J. Ecol.* **105**: 1044–1057.

Ehlers, B.K. & Bataillon, T. 2007. "Inconstant males" and the maintenance of labile sex
expression in subdioecious plants. *New Phytol.* **174**: 194–211.

Eppley, S.M. & Pannell, J.R. 2007a. Density-dependent self-fertilization and male versus
hermaphrodite siring success in an androdioecious plant. *Evolution* **61**: 2349–2359.

Eppley, S.M. & Pannell, J.R. 2007b. Sexual systems and measures of occupancy and abundance
in an annual plant: Testing the metapopulation model. *Am. Nat.* **169**: 20–28.

Excoffier, L., Foll, M. & Petit, R.J. 2009. Genetic consequences of range expansions. *Annu. Rev.*
Ecol. Evol. Syst. **40**: 481–501.

Florian, H. 2016. DHARMa: Residual diagnostics for hierarchical (multi-level / mixed)
regression models. *CRAN Contrib. Packag.*, doi: 10.32614/cran.package.dharma. The R
Foundation.

Franceschinelli, E.V. & Bawa, K.S. 2000. The effect of ecological factors on the mating system
of a South American shrub species (*Helicteres brevispira*). *Heredity* **84**: 116–123.

Freeman, D.C., Harper, K.T. & Charnov, E.L. 1980. Sex change in plants: Old and new
observations and new hypotheses. *Oecologia* **47**: 222–232.

Gerchen, J.F., Veltsos, P. & Pannell, J.R. 2022. Recurrent allopolyploidization, Y-chromosome
introgression and the evolution of sexual systems in the plant genus *Mercurialis. Philos.*
Trans. R. Soc. B **377**: 20210224. The Royal Society.

Gerchen, J.F., Villamil-Buenrostro, N., Li, X., Le Faou, E. & Pannell, J.R. 2026. Sex-allocation
trade-offs and their genetic architecture revealed by experimental evolution. *Proc. Natl.*
Acad. Sci. **123**: e2427240123.

Golenberg, E.M. & West, N.W. 2013. Hormonal interactions and gene regulation can link
monoecy and environmental plasticity to the evolution of dioecy in plants. *Am. J. Bot.*
100: 1022–1037.

González-Martínez, S.C., Ridout, K. & Pannell, J.R. 2017. Range expansion compromises
adaptive evolution in an outcrossing plant. *Curr. Biol.* **27**: 2544-2551.e4.

Grime, J.P. 1974. Vegetation classification by reference to strategies. *Nature* **250**: 26–31.

Grossenbacher, D.L., Brandvain, Y., Auld, J.R., Burd, M., Cheptou, P.-O., Conner, J.K., *et al.*
2017. Self-compatibility is over-represented on islands. *New Phytol.* **215**: 469–478.

Hamdi, S., Teller, G. & Louis, J.P. 1987. Master regulatory genes, auxin levels, and sexual
organogeneses in the dioecious plant *Mercurialis annua*. *Plant Physiol.* **85**: 393–399.

Hamilton, W.D. 1967. Extraordinary sex ratios. *Science* **156**: 477–488.

Hargreaves, A.L. & Eckert, C.G. 2014. Evolution of dispersal and mating systems along
geographic gradients: Implications for shifting ranges. *Funct. Ecol.* **28**: 5–21.

Harris, M.S. & Pannell, J.R. 2008. Roots, shoots and reproduction: Sexual dimorphism in size
and costs of reproductive allocation in an annual herb. *Proc. Biol. Sci.* **275**: 2595–2602.

Hesse, E. & Pannell, J.R. 2011. Sexual dimorphism in a dioecious population of the wind-
pollinated herb *Mercurialis annua*: The interactive effects of resource availability and
competition. *Ann. Bot.* **107**: 1039–1045.

Hewitt, G. 2000. The genetic legacy of the Quaternary ice ages. *Nature* **405**: 907–913.

Hörandl, E. 2023. Geographical parthenogenesis in alpine and arctic plants. *Plants* **12**: 844.

Hörandl, E., Cosendai, A.-C. & Temsch, E.M. 2008. Understanding the geographic distributions
of apomictic plants: A case for a pluralistic approach. *Plant Ecol. Divers.* **1**: 309–320.

Ibrahim, K.M., Nichols, R.A. & Hewitt, G.M. 1996. Spatial patterns of genetic variation
generated by different forms of dispersal during range expansion. *Heredity* **77**: 282–291.

Korpelainen, H. 1998. Labile sex expression in plants. *Biol. Rev.* **73**: 157–180.

Krähenbühl, M., Yuan, Y.-M. & Küpfer, P. 2002. Chromosome and breeding system evolution
of the genus *Mercurialis* (Euphorbiaceae): implications of ITS molecular phylogeny.
Plant Syst. Evol. **234**: 155–169.

Li, X., Veltsos, P., Cossard, G.G., Gerchen, J. & Pannell, J.R. 2019. YY males of the dioecious
plant *Mercurialis annua* are fully viable but produce largely infertile pollen. *New Phytol.*
224: 1394.

Lloyd, D.G. 1965. Evolution of self-compatibility and racial differentiation in *Leavenworthia*
(Cruciferae). *Contrib. Gray Herb. Harv. Univ.* 3–134.

Lloyd, D.G. 1980. The distributions of gender in four angiosperm species illustrating two
evolutionary pathways to dioecy. *Evolution* **34**: 123–134.

Lloyd, D.G. & Bawa, K.S. 1984. Modification of the gender of seed plants in varying conditions.
In: *Evolutionary Biology* (M. K. Hecht, B. Wallace, & G. T. Prance, eds), pp. 255–338.
Springer US, Boston, MA.

Longhurst, A.R. 1955. Evolution in the Notostraca. *Evolution* **9**: 84–86.

Louis, J.P. & Durand, B. 1978. Studies with the dioecious angiosperm *Mercurialis annua* L.
(2n=16): Correlation between genic and cytoplasmic male sterility, sex segregation and
feminizing hormones (cytokinins). *Mol. Gen. Genet. MGG* **165**: 309–322.

MacArthur, R.H. & Wilson, E.O. 1967. *The theory of island biogeography*, REV-Revised.
Princeton University Press.

Moeller, D.A. & Geber, M.A. 2005. Ecological context of the evolution of self-pollination in
Clarkia xantiana: population size, plant communities, and reproductive assurance. *Evol.*
Int. J. Org. Evol. **59**: 786–799.

Monty, A. & Mahy, G. 2010. Evolution of dispersal traits along an invasion route in the wind-
dispersed *Senecio inaequidens* (Asteraceae). *Oikos* **119**: 1563–1570.

Ness, R.W., Wright, S.I. & Barrett, S.C.H. 2010. Mating-system variation, demographic history
and patterns of nucleotide diversity in the tristylous plant *Eichhornia paniculata*.
Genetics **184**: 381–392.

Obbard, D.J., Harris, S.A., Buggs, R.J.A. & Pannell, J.R. 2006a. Hybridization, polyploidy, and the evolution of sexual systems in *Mercurialis* (Euphorbiaceae). *Evolution* **60**: 1801–1815.

Obbard, D.J., Harris, S.A. & Pannell, J.R. 2006b. Sexual systems and population genetic structure in an annual plant: Testing the metapopulation model. *Am. Nat.* **167**: 354–366.

Olivieri, I. & Gouyon, P.-H. 1997. Evolution of migration rate and other traits: The metapopulation effect. In: *Metapopulation biology: Ecology, genetics and evolution* (I. A. Hanski & M. E. Gilpin, eds), pp. 293–323. Academic Press.

Pannell, J.R. 2000. A hypothesis for the evolution of androdioecy: The joint influence of reproductive assurance and local mate competition in a metapopulation. *Evol. Ecol.* **14**: 195–211.

Pannell, J.R. 2015. Evolution of the mating system in colonizing plants. *Mol. Ecol.* **24**: 2018–2037.

Pannell, J.R. 1997. The maintenance of gynodioecy and androdioecy in a metapopulation. *Evolution* **51**: 10–20.

Pannell, J.R., Auld, J.R., Brandvain, Y., Burd, M., Busch, J.W., Cheptou, P.-O., *et al.* 2015. The scope of Baker's Law. *New Phytol.* **208**: 656–667.

Pannell, J.R. & Barrett, S.C.H. 1998. Baker's Law revisited: Reproductive assurance in a metapopulation. *Evolution* **52**: 657–668.

Pannell, J.R., Dorken, M.E., Pujol, B. & Berjano, R. 2008. Gender variation and transitions between sexual systems in *Mercurialis annua* (Euphorbiaceae). *Int. J. Plant Sci.* **169**: 129–139.

Pannell, J.R., Eppley, S.M., Dorken, M.E. & Berjano, R. 2014. Regional variation in sex ratios and sex allocation in androdioecious *Mercurialis annua*. *J. Evol. Biol.* **27**: 1467–1477.

Pannell, J.R. & Jordan, C.Y. 2022. Evolutionary transitions between hermaphroditism and dioecy in animals and plants. *Annu. Rev. Ecol. Evol. Syst.* **53**: 183–201.

Peischl, S., Dupanloup, I., Kirkpatrick, M. & Excoffier, L. 2013. On the accumulation of deleterious mutations during range expansions. *Mol. Ecol.* **22**: 5972–5982.

Peischl, S., Kirkpatrick, M. & Excoffier, L. 2015. Expansion load and the evolutionary dynamics of a species range. *Am. Nat.* **185**: E81-93.

Phillips, B.L., Brown, G.P., Greenlees, M., Webb, J.K. & Shine, R. 2007. Rapid expansion of the cane toad (*Bufo marinus*) invasion front in tropical Australia. *Austral Ecol.* **32**: 169–176.

593 Phillips, B.L., Brown, G.P., Webb, J.K. & Shine, R. 2006. Invasion and the evolution of speed in
594   toads. *Nature* **439**: 803–803.

595 Pianka, E.R. 1970. On r- and K-Selection. *Am. Nat.* **104**: 592–597.

596 Pujol, B., Zhou, S.R., Vilas, J.S. & Pannell, J.R. 2009. Reduced inbreeding depression after
597   species range expansion. *Proc. Natl. Acad. Sci. U. S. A.* **106**: 15379–15383.

598 R Core Team. 2025. R: A language and environment for statistical computing. R Foundation for
599   Statistical Computing, Vienna, Austria. <https://www.R-project.org/>., Vienna, Austria.

600 Reznick, D., Bryant, M.J. & Bashey, F. 2002. r- and K-selection revisited: The role of
601   population regulation in life-history evolution. *Ecology* **83**: 1509–1520.

602 Ross, M.D. 1982. Five evolutionary pathways to subdioecy. *Am. Nat.* **119**: 297–318.

603 Ross, M.D. 1980. The evolution and decay of overdominance during the evolution of
604   gynodioecy, subdioecy, and dioecy. *Am. Nat.* **116**: 607–620.

605 Roux, C., Mullon, C. & Pannell, J.R. 2024. The evolution of sex allocation in metapopulations.
606   bioRxiv.

607 Ruane, L.G., Mangum, S.M., Horner, K.M. & Moeller, D.A. 2020. The opportunity for
608   outcrossing varies across the geographic range of the primarily selfing *Clarkia xantiana*
609   ssp. *parviflora*. *Am. J. Bot.* **107**: 1198–1207.

610 Russell, J.R.W. & Pannell, J.R. 2015. Sex determination in dioecious *Mercurialis annua* and its
611   close diploid and polyploid relatives. *Heredity* **114**: 262–271.

612 Sánchez Vilas, J. & Pannell, J.R. 2011. Sexual dimorphism in resource acquisition and
613   deployment: Both size and timing matter. *Ann. Bot.* **107**: 119–126.

614 Sexton, J.P., McIntyre, P.J., Angert, A.L. & Rice, K.J. 2009. Evolution and ecology of species
615   range limits. *Annu. Rev. Ecol. Evol. Syst.* **40**: 415–436.

616 Shine, R., Brown, G.P. & Phillips, B.L. 2011. An evolutionary process that assembles
617   phenotypes through space rather than through time. *Proc. Natl. Acad. Sci.* **108**: 5708–
618   5711.

619 Stearns, S.C. 1983. The influence of size and phylogeny on patterns of covariation among life-
620   history traits in the mammals. *Oikos* **41**: 173–187.

621 Stearns, S.C. 1989. Trade-offs in life-history evolution. *Funct. Ecol.* **3**: 259–268.

622 Stebbins, G.L. 1957. Self fertilization and population variability in the higher plants. *Am. Nat.*
623   **91**: 337–354.

624 Svenning, J.-C. & Skov, F. 2007. Ice age legacies in the geographical distribution of tree species
625   richness in Europe. *Glob. Ecol. Biogeogr.* **16**: 234–245.

Taylor, D.R., Trimble, S. & McCauley, D.E. 1999. Ecological genetics of gynodieocy in *Silene*
vulgaris: Relative fitness of females and hermaphrodites during the colonization process.
Evolution **53**: 745–751.

Tilman, D. 1990. Constraints and tradeoffs: Toward a predictive theory of competition and
succession. *Oikos* **58**: 3–15.

Tonnabel, J., David, P., Klein, E.K. & Pannell, J.R. 2019. Sex-specific selection on plant
architecture through "budget" and "direct" effects in experimental populations of the
wind-pollinated herb, *Mercurialis annua*. *Evolution* **73**: 897–912.

Travis, J., Bassar, R.D., Coulson, T., Reznick, D. & Walsh, M. 2023. Density-dependent
selection. *Annu. Rev. Ecol. Evol. Syst.* **54**: 85–105.

Travis, J.M.J. & Dytham, C. 1999. Habitat persistence, habitat availability and the evolution of
dispersal. *Proc. R. Soc. Lond. B Biol. Sci.* **266**: 723–728.

Urquhart-Cronish, M., Angert, A.L., Otto, S.P. & MacPherson, A. 2024. Density-dependent
selection during range expansion affects expansion load in life history traits. *Am. Nat.*
203: 382–392.

Veltsos, P., Cossard, G., Beaudoing, E., Beydon, G., Bianchi, D.S., Roux, C., *et al.* 2018. Size
and content of the sex-determining region of the Y chromosome in dioecious *Mercurialis*
annua, a plant with homomorphic sex chromosomes. *Genes* **9**: 277.

Veltsos, P., Ridout, K.E., Toups, M.A., González-Martínez, S.C., Muyle, A., Emery, O., *et al.*
2019. Early sex-chromosome evolution in the diploid dioecious plant *Mercurialis annua*.
Genetics **212**: 815–835.

Villamil, N., Li, X., Seddon, E. & Pannell, J.R. 2022. Simulated herbivory enhances leaky sex
expression in the dioecious herb *Mercurialis annua*. *Ann. Bot.* **129**: 79–86.

Vincenty, T. 1975. Direct and inverse solutions of geodesics on the ellipsoid with application of
nested equations. *Surv. Rev.* **23**: 88–93.

Weiss-Lehman, C. & Shaw, A.K. 2022. Understanding the drivers of dispersal evolution in
range expansions and their ecological consequences. *Evol. Ecol.* **36**: 181–197.

Willi, Y., Fracassetti, M., Zoller, S. & Van Buskirk, J. 2018. Accumulation of mutational load at
the edges of a species range. *Mol. Biol. Evol.* **35**: 781–791.

Figures and Tables

Figure 1. Map of sampling sites in Europe (red points) with their labels (detailed in Table S1).

The point of origin (black point) assumed in our analyses for the expansion origin is midway

between populations 9 and 27.

Figure 2. Measures of leakiness (A) and reproductive effort for females (B) and males (C)

sampled as a function of distance from the expansion origin, based on traits expressed in the

common-garden. Indicated are plant sex and hormone-treatment status (see inset legend). Lines

show GLMM-predicted responses (see Table S2) as functions of distance from the expansion

origin. Statistical significance is indicated in the upper left of each panel.