Hypatia: Comparative Isoform Profiling Across Cell Populations from Long-Read Single-Cell Transcriptomes

Timothy Pan[1, 2, 3], Cheng-Kai Shiau[1, 2], Lina Lu[1, 2], Changhu Wang[4, 5], Minhua Wang[1, 2], Yueying He[1, 2], Arvind Bhimaraj[6], Daniel Brat[7], Jason Huse[8], Jingyi Jessica Li[4, 5], Ruli Gao[1, 2, 3, *]

Affiliations

[1] Department of Biochemistry and Molecular Genetics, Feinberg School of Medicine, Northwestern University, Chicago, IL, USA 60611

[2] Center for Cancer Genomics, Robert H. Lurie Cancer Center, Northwestern University, Chicago, IL, USA 60611

[3] Driskill Graduate Program, Northwestern University, Chicago, IL, USA 60611

[4] Biostatistics Program, Public Health Sciences Division, Fred Hutch Cancer Center, Seattle, WA, USA 98109

[5] Department of Biostatistics, University of Washington, Seattle, WA, USA 98109

[6] Department of Cardiology, Methodist DeBakey Cardiology Associates, Houston Methodist Hospital, Houston, TX, USA-77030

[7] Department of Pathology, Northwestern University – Feinberg School of Medicine, Chicago, IL, USA 60611

[8] Department of Pathology and Translational Molecular Pathology, University of Texas MD Anderson Cancer Center, Houston, TX, USA 77030

* Corresponding author

Email addresses

Timothy Pan, timothy.pan@northwestern.edu

Cheng-Kai Shiau, chengkai.shiau@northwestern.edu

Lina Lu, lina.lu@northwestern.edu

Changhu Wang, wangch156@g.ucla.edu

Minhua Wang, minhua.wang@northwestern.edu

Yueying He, yueying.he@northwestern.edu

Arvind Bhimaraj, ABhimaraj@houstonmethodist.org

Daniel Brat, daniel.brat@northwestern.edu

Jason Huse, jhuse@mdanderson.org

Jingyi Jessica Li, lijy03@fredhutch.org

Ruli Gao, ruli.gao@northwestern.edu

Abstract

High-throughput long-read single-cell RNA-sequencing enables isoform-level study across single cells, yet methods for systematically assessing cell-to-cell variations remain limited. Here, we develop Hypatia, a comprehensive platform for dissecting isoform complexities across cell populations, devising Tsallis entropy and Cramer's V to facilitate robust comparative profiling. Hypatia revealed prominent isoform species variations and usage shifts across cell-types in glioblastoma, renal cell carcinoma, and heart, highlighting clinically relevant applications for studying isoform-derived, cell-specific functions.

Keywords

isoform usage shift, Tsallis isoform diversity, Chi-square correction

Background

Alternative splicing is an essential post-transcriptional mechanism that expands the coding capacity of eukaryotic genomes(1). Despite its critical roles in health and disease, cell-type- and cell-state-specific RNA isoform complexity remains poorly understood due to the limitations of bulk and short-read sequencing technologies. Long-read single-cell RNA-sequencing (LR-scRNAseq)(2-7) achieves high-throughput and single-cell resolution of isoform quantification by coupling single-cell barcoding with long-read sequencing platforms (i.e., ONT and PacBio). However, existing analytical strategies(8-10) lack the desired levels of granularity in calling cell population-specific events relevant to isoform-centric regulatory mechanisms in cells. To address this knowledge gap, we develop Hypatia, a computational platform for the systematic quantification and comparison of population-specific isoform profiles from LR-scRNAseq data.

Results and Discussion

The workflow of Hypatia inputs single cell isoform expression data enabled by LR-scRNAseq to perform three layers of isoform complexity quantification and subsequently compare them across cell populations, including 1) isoform diversity analysis, which classifies genes according to isoform species heterogeneity and identifies differential isoform diversity events (DIVs) between cell populations, 2) isoform usage shift analysis, which identifies differential isoform usage shifts (DIUs) between cell populations, and 3) isoform expression analysis, which detects differentially expressed isoforms (DEIs) across cells (**Fig. 1a**). Each analyses offers unique insights into cell-specific isoforms that are essential for understanding cellular function and disease mechanisms, integrating statistical methods for isoform species diversity, relative usage shifts, and expression abundance in a single platform. In addition, Hypatia leverages object-oriented R programming in the *SingleCellExperiment* class(11), which structures count data, cell-level and feature-level metadata in a synchronized manner, supporting seamless multi-scale data analysis.

We established statistical measurements via rigorous tests in simulation, where isoform

counts are modeled in single cells for two populations by applying a multinomial probabilistic

approach on simulated gene expression, with a subset of genes engineered to bear ground-truth

isoform usage variations (**Fig. 1b**, **Methods**). To support fair comparisons across cells and genes,

we began by assessing isoform species diversity, which represents both the number of isoform

species and their evenness within individual genes. A robust diversity quantification is expected

to mitigate bias from gene-to-gene variation in isoform species number and from the heavy tail of

lowly expressed isoforms. Previous bulk studies have applied entropy-based methods, yet no

unified metric capable of effectively capturing differences across genes and cell-types has been

established. To this end, we evaluated a range of entropic measures (**Methods**) for their reliability

and sensitivity in classifying isoform heterogeneity (e.g., 'monoform' versus 'polyform') and the

relative prevalence of the most dominant transcript (MDT) per gene. Our analysis revealed that

Tallis entropy (q=3) demonstrated the highest association with MDT proportions in both 'monoform'

and 'polyform' genes, outperforming the widely used Shannon entropy(2, 12) (**Fig. 1d**). Of note,

Figure 1. Hypatia workflow and simulation-guided assessment. a, Illustration of the main functional modules and workflow of Hypatia R package. **b,** Diagram outlining the steps in single-cell isoform count simulation. **d,** Relationship between isoform diversity and P$_{MDT}$ of all genes. Pearson correlation coefficients are shown for all data (black), only polyform (orange), and only monoform (cyan) genes. **d,** Precision-recall curve of DIU identification across multiple gene coverage groups with AUCPR shown, colored by selection criteria.

Tsallis entropy facilitates cross-gene comparisons, being independent of isoform species number

compared to other diversity methods (e.g., Shannon, Renyi), minimizing the impact of lowly

expressed isoforms. Together, these data establish Tsallis entropy as a unified metric for

quantifying isoform species diversity for comparative analyses.

Isoform usage shifts provide another critical aspect for examining differential isoform

profiles across cells, while isoform diversity overlooks the relative ranks among isoforms and can

remain unchanged in reciprocal switching events. For this purpose, we evaluated the frequently

adapted Chi-square test, which is highly dependent on sample-size and whose feasibility in

analyzing sparse single-cell data has not been established. Our assessment revealed that Chi-

square P-value alone provided moderate performance (AUCPR 0.460-0.651) in calling true DIU

events across coverage groups (**Fig. 1d**). Coupling with an empirical criterion, namely *Delta* (i.e.,

the greatest difference in isoform proportions per gene between groups) partially rescued the

performance, which however remained low with poor data coverage. Intriguingly, we found that

coupling with effect size (e.g., Cramer's V)(13) (**Methods**) considerably mitigated the spurious

results from sample-size variation and yielded the strongest performance (AUCPR 0.609-0.915)

across all coverage groups (**Fig. 1d**). To summarize, while the combined use of Chi-square

statistics and *Delta* provided a solid baseline for DIU selection, integration with Cramer's V

emerged as a superior approach and consistently outperformed all other selection methods.

To test applications in profiling cell-type-specific isoforms, we applied Hypatia on three LR-

scRNAseq datasets, including an in-house glioblastoma (GBM) dataset and previously published

datasets derived from renal cell carcinoma (RCC)(2) and normal heart(12) tissues (**Table S1,**

Methods). Data showed marked expression of unique isoforms associated with cell-type-specific

genes in all samples (**Fig. 2a-c**). Deeper analysis with Hypatia revealed a bimodal distribution of

Tsallis entropy in all cell-types, corresponding to 'monoform' and 'polyform' genes (**Fig. 2d**).

Notably, we observed a distinct increase in isoform diversity of RCC tumor cells and heart

cardiomyocytes, suggesting that these cells undergo more active splicing programs.

Furthermore, Hypatia identified extensive isoform usage shifts across cell-types that may

reflect underlying regulatory mechanisms of cell-type-specific functions. In GBM, multiple DIU

events were identified in the tumor-suppressor lncRNA gene *MEG3* (**Fig. 2e**). The relative

Figure 2. Hypatia characterizes isoform-level variations across cell-types in LR-scRNAseq. a-c, (left) UMAP determined by gene expression analysis and (right) heatmap of DEIs of each cell-type. **d,** Distribution of cell-type-specific isoform diversity of expressed genes in each sample, color coded as in panels **a-c**. **e-g,** (left) Heatmap showing cell-type-specific isoform proportions of DIU events in each sample. (right) Transcript structures of isoforms corresponding to heatmap rows. **h,** Treemaps showing intersections between three analyses, shaded by overlapping genes in each sample.

abundance of dominant isoform MEG3-209 (ENST00000451743) varied broadly across multiple

cell-types, whereas MEG3-220 (ENST00000524035) and novel isoform MEG3-nic20562 had

distinct expression in neoplastic tumor cells and oligodendrocytes, respectively. Similarly, the

inflammation-related gene, *TMSB4X,* exhibited a marked decrease in protein-coding isoform

TMSB4X-204 (ENST00000451311) specific to microglial cells, accompanied by an increase in

novel isoform TMSB4X-nic1330, which harbors a retained intron, indicating cell-type-specific

splicing regulation. In RCC, significant DIU events were detected in glycolysis gene *ENO1* and

metallothionein-encoding gene *MT2A* (**Fig. 2f**). Tumor cells showed a distinct polyform

heterogeneity of *ENO1*, while a striking dichotomy in the dominant isoform of *MT2A* emerged

across cell-types. In cardiomyocytes of the heart, we observed unique splicing patterns in *MYL3*

and *COX7A1*, which are key regulators in sarcomere contraction and energy production,

respectively (**Fig. 2g**). These data signified that splicing regulation involved key cell-type-specific

functional genes across various tissue types.

While DIU analysis captures changes in relative isoform usage with consideration of the

ranked order of dominant isoforms, DIVs reflect differences in evenness independent of ranking.

Indeed, this was well-reflected in comparative findings from DIU and DIV analysis. In GBM, DIU

and DIV analyses identified isoform complexity changes in 54 and 133 genes, respectively, with

9 overlapped genes (**Fig. 2h**). In RCC, 60 among 297 DIU and 587 DIV events were jointly

detected. In heart, 45 genes were jointly detected among 123 DIU and 96 DIV events. In contrast,

DEI analysis identified fewer genes linked to cell-type-specific isoform profiles (N= 23, 33, and 22

genes in GBM, RCC, and heart, respectively), suggesting that DEI analysis is largely confounded

by overall gene expression levels. Taken together, our results show that DIU and DIV analyses

offer a complementary approach for discovering isoform complexity changes across diverse cell

populations.

Collectively, our results highlight the utility of Hypatia for isoform-resolved analysis of

single-cell transcriptomes. We show that Tsallis entropy robustly quantifies isoform species

heterogeneity and Cramer's V effectively curates Chi-square statistics from long-read single-cell

isoform data. Critically, we show that DIU and DIV tests are complementary in dissecting

differential isoform patterns across cell populations. Results from clinically relevant datasets show

the prevalence of isoform-level variations across cell-types and exemplify the important

applications of Hypatia in revealing comprehensive isoform spectrums of human tissues.

Conclusions

In this study, we present Hypatia, a novel analytical framework for in-depth comparisons of cell-

to-cell variations in isoform complexity from long-read single-cell transcriptomic data. Hypatia

mitigates challenges of data noise and sparsity through structured and systematic approaches,

which greatly extends the profiling capacity of single-cell transcriptomics beyond overall gene

expression levels and provides comprehensive details at isoform-level.

Methods

Simulation of single-cell isoform expression data

Isoform data were simulated for 30,000 genes and 1,000 cells of two populations, each with 500

cells. Generation of single-cell gene expression counts for each group was adapted from

Splatter(14) and utilized the gamma distribution for mean gene expression (k = 0.2, θ = 2) and

the negative binomial distribution with a dispersion parameter of 0.10. The number of isoforms

per gene were drawn from a Poisson distribution (λ = 2) with a minimum of 2 and maximum of 10

possible isoforms. Isoform-level *Deltas* for each gene and corresponding isoform proportions for

both groups were sampled from beta distributions. Genes with at least one *Delta* of $\geq$ 0.10 across

groups were considered ground-truth DIUs. Isoform counts were then generated from the

multinomial distribution using group-specific isoform proportions and gene counts per cell. In

performance evaluation, gene coverage groups were categorized according to total counts per

gene: ≤ 100 counts ('very low'), 100 to 200 ('low'), 200 to 500 ('moderate'), and ≥ 500 ('high').

Sample acquisition and LR-scRNA sequencing

Fresh frozen tissue from a grade 4 glioblastoma of the frontal lobe was collected from the NBST

tissue bank at Northwestern University and subjected to snNanoRNAseq(2). The single-nuclei

suspension was prepared using the Chromium Nuclei Isolation Kit (10X Genomics) following the

manufacturer's protocol. Nuclei were then loaded onto a 10X Genomics Chromium Controller (iX)

with Chip J with an expected recovery of 3,000 nuclei. Full-length barcoded cDNAs were amplified

and enriched for long fragments. The final library was prepared using the SQK-LSK114 ligation

sequencing kit (Oxford Nanopore Technologies) and sequenced on an in-house PromethION with

an R10.4M flow cell (Oxford Nanopore Technologies).

LR-scRNAseq data preprocessing

All processing of raw data was performed using scNanoGPS(2) (version 2.0). Isoform

quantification was performed with IsoQuant(15) (version 3.4.2) using the 'unique only' transcript

quantification setting, and reference genome GRCh38 (GENCODE v44/Ensembl110 annotations),

excluding reads mapping to *MALAT1* and *NEAT1* due to computational restraints. SQANTI3(16)

(version 5.2.2) was used for transcript annotation. Transcripts categorized as full-splice match

were retained. Novel transcripts required support of 5' CAGE peaks in refTSS(17) (version 4.1)

and 3' polyA sites in PolyASite(18) (version 3.0). All transcripts were required to be detected in at

least 3 cells in the sample and free of predicted RT-switching artifacts.

Single-cell gene expression analysis

Single-cell gene expression analysis was performed using Seurat(19) (version 5.0.1). Cells with

expression of more than 200 genes and less than 5% of mitochondrial genes were retained.

Clustering utilized the top 15 principal components and cell-types were annotated using

established gene markers(20).

Differential isoform usage analysis

To calculate isoform usage shifts as implemented in Hypatia's 'RunDIU' function, a contingency

table was constructed for each gene using aggregated isoform counts of each cell group in the

comparison. The Chi-square test was applied, and *P*-values were adjusted for multiple testing

using the Benjamini-Hochberg method. Each cell-type was compared against all other cell-types

for all comparative analyses. Comparisons resulting in an adjusted *P*-value of ≤ 0.05 and

Cramer's V ≥ 0.20 were considered significant. Only genes expressed in at least 5% of cells in

each group and supported by 15 total UMI counts were considered.

Calculation of effect size

215 The effect size of Chi-square test was calculated as Cramer's V (**Eq. 1**),

$$V = \sqrt{\frac{\chi^2}{n(k-1)}},$$

(1)

217 where χ^2 denotes the Chi-square statistic, n denotes the total isoform count of the gene, and k is

218 the number of columns in the contingency table (equal to 2 for two-group comparisons).

Isoform diversity analysis

220 Isoform diversity was calculated using Tsallis (**Eq. 2**, $q = 3$), Shannon (**Eq. 3**), Gini-Simpson (**Eq.**

221 **4**), Renyi (**Eq. 5**, $\alpha = 2$), and Inverse-Simpson (**Eq. 6**) entropies as implemented in the 'RunDIV'

222 function of Hypatia. The entropies were calculated as the following,

$$S = \frac{1}{q-1}\left(1 - \sum_{i}^{n} p_i^q\right),$$

(2)

$$H = -\sum_{i=1}^{2} p_i \, log(p_i),$$ (3)

$$D = 1 - \sum_{i}^{n} p_i^2,$$ (4)

$$H_\alpha = \frac{1}{1-\alpha} \log\left(\sum_{i}^{2} p_i^\alpha\right),$$ (5)

$$D^{-1} = \frac{1}{D},$$ (6)

where n denotes the total number of expressed isoforms of the gene, p is the isoform proportion,

and q and α represent the order of the Tsallis and Renyi entropies, respectively. The Shannon

and Renyi entropies utilized only the top 2 isoform abundances to normalize comparisons across

genes. Cutoffs for monoform and polyform classifications were guided by a P_{MDT} = 0.80, which

corresponded to entropies at 0.243 (Tsallis), 0.500 (Shannon), 0.348 (Gini-Simpson), 0.435

(Renyi), and 1.533 (inverse-Simpson). DIVs were defined as genes whose diversity classifications

diverged across cell groups. Only genes with expression in at least 5% of cells of each group of

the comparison were considered.

Isoform expression analysis

Raw isoform count data were first normalized using 'NormalizeCounts' in Hypatia. Counts were

normalized by the total counts per cell, multiplied by a scale factor of 10,000, and subsequently

Log1p-transformed. The average isoform expression across cell-types were calculated using

'RunDEI', which applies the Wilcoxon Rank Sum Test with Bonferroni corrected *P*-values.

Significant DEIs were those that had |Log2 fold change| $\geq$ 0.60, adjusted *P*-value $\leq$ 0.05, and

detected in at least 5% of cells in comparisons.

Intersection of differential isoform events

Genes presenting differential isoform events were enumerated per cell-type comparison, where

each cell-type was compared against all others. Genes associated with more than one DEI were

only counted once per comparison. Genes that appeared in more than one cell-type comparison

were assigned a separate count for each comparison.

Declarations

Ethics approval and consent to participate

The IRB protocol for studying the multi-omics profiles of human tissues were approved by

Northwestern University (STU00219936). Consent was waved because this study only involves

the secondary analysis of the banked tissues, which has no direct interaction with subjects.

Consent for publication

Not applicable.

Availability of data and materials

Both raw and processed data generated in this study have been deposited in the Gene Expression

Omnibus (GEO) database under accession GSE310974. Previously published sequencing data

were downloaded from SRA under runs SRR21492157 (RCC) and SRR32154432 (heart).

Simulation data have been deposited in Zenodo (doi: 10.5281/zenodo.17781226). All source code

and documentation of Hypatia are open source and are released to the public and freely available

on Github at https://github.com/gaolabtools/Hypatia.

Competing interests

The authors declare that they have no competing interests.

Funding

This work was supported by the National Institute of General Medical Sciences (NIH

R35GM142539) and National Cancer Institute SPORE for Translational Approaches to Brain

Cancer (P50CA221747).

Authors' contributions

TP led the development of Hypatia, performed data analysis, and wrote the manuscript. CKS

processed raw sequencing data and reviewed results. MW performed the sequencing experiment

and edited the manuscript. LL and YH reviewed results. QW and JJL participated in data analysis,

reviewed results, and edited the manuscript. JH, DB and AB assisted with sample collection and

edited the manuscript. RG supervised the study, conceived concepts, designed the computational

tool, and wrote the manuscript.

Acknowledgements

We thank all our funding supports and patients who consented for research usages of their tissue

samples.

References

1.   Jiang W, Chen L. Alternative splicing: Human disease and quantitative analysis from high-

throughput sequencing. Comput Struct Biotechnol J. 2021;19:183-95.

2.   Shiau CK, Lu L, Kieser R, Fukumura K, Pan T, Lin HY, et al. High throughput single cell

long-read sequencing analyses of same-cell genotypes and phenotypes in human tumors. Nat

Commun. 2023;14(1):4124.

3.   Al'Khafaji AM, Smith JT, Garimella KV, Babadi M, Popic V, Sade-Feldman M, et al. High-

throughput RNA isoform sequencing using programmed cDNA concatenation. Nat Biotechnol.

2024;42(4):582-6.

4.   Fan X, Tang D, Liao Y, Li P, Zhang Y, Wang M, et al. Single-cell RNA-seq analysis of

mouse preimplantation embryos by third-generation sequencing. PLoS Biol.

2020;18(12):e3001017.

5.   Lebrigand K, Magnone V, Barbry P, Waldmann R. High throughput error corrected

Nanopore single cell transcriptome sequencing. Nat Commun. 2020;11(1):4025.

6.   Philpott M, Watson J, Thakurta A, Brown T, Jr., Brown T, Sr., Oppermann U, et al.

Nanopore sequencing of single-cell transcriptomes with scCOLOR-seq. Nat Biotechnol.

2021;39(12):1517-20.

7.   Wang Q, Boenigk S, Boehm V, Gehring NH, Altmueller J, Dieterich C. Single cell

transcriptome sequencing on the Nanopore platform with ScNapBar. RNA. 2021.

8.   Guo W, Calixto CPG, Brown JWS, Zhang R. TSIS: an R package to infer alternative

splicing isoform switches for time-series data. Bioinformatics. 2017;33(20):3308-10.

9.   de la Fuente L, Arzalluz-Luque A, Tardaguila M, Del Risco H, Marti C, Tarazona S, et al.

tappAS: a comprehensive computational framework for the analysis of the functional impact of

differential splicing. Genome Biol. 2020;21(1):119.

10.   Wu S, Schmitz U. ScIsoX: a multidimensional framework for measuring isoform-level

transcriptomic complexity in single cells. Genome Biol. 2025;26(1):289.

11.   Amezquita RA, Lun ATL, Becht E, Carey VJ, Carpp LN, Geistlinger L, et al. Orchestrating

single-cell analysis with Bioconductor. Nature Methods. 2020;17(2):137-45.

12.   Pan T, Lu L, Youker K, Shiau CK, Wang M, Lin HY, et al. Single-Cell Splicing Isoform Atlas

of the Adult Human Heart and Heart Failure. Circulation. 2025;152(21):1501-14.

13.   Cramér, Harald. 1946. Mathematical Methods of Statistics. Princeton: Princeton University

Press, page 282 (Chapter 21. The two-dimensional case). ISBN 0-691-08004-6 (table of content

Archived 2016-08-16 at the Wayback Machine).

14.   Zappia L, Phipson B, Oshlack A. Splatter: simulation of single-cell RNA sequencing data.

Genome Biology. 2017;18(1):174.

15.   Prjibelski AD, Mikheenko A, Joglekar A, Smetanin A, Jarroux J, Lapidus AL, et al. Accurate

isoform discovery with IsoQuant using long reads. Nature Biotechnology. 2023;41(7):915-8.

16.   Pardo-Palacios FJ, Arzalluz-Luque A, Kondratova L, Salguero P, Mestre-Tomás J, Amorín

R, et al. SQANTI3: curation of long-read transcriptomes for accurate identification of known and

novel isoforms. Nat Methods. 2024;21(5):793-7.

17.   Abugessaisa I, Noguchi S, Hasegawa A, Kondo A, Kawaji H, Carninci P, et al. refTSS: A

Reference Data Set for Human and Mouse Transcription Start Sites. Journal of Molecular Biology.

2019;431(13):2407-22.

18.   Moon Y, Herrmann Christina J, Mironov A, Zavolan M. PolyASite v3.0: a multi-species

atlas of polyadenylation sites inferred from single-cell RNA-sequencing data. Nucleic Acids

Research. 2024;53(D1):D197-D204.

19.   Butler A, Hoffman P, Smibert P, Papalexi E, Satija R. Integrating single-cell transcriptomic

data   across   different   conditions,   technologies,   and   species.   Nature   Biotechnology.

2018;36(5):411-20.

20.   Couturier CP, Ayyadhury S, Le PU, Nadaf J, Monlong J, Riva G, et al. Single-cell RNA-

seq reveals that glioblastoma recapitulates a normal neurodevelopmental hierarchy. Nature

Communications. 2020;11(1):3406.

335