2-AG as a potential biomarker for predicting response to short-term fluoxetine treatment in a mouse model of depression

Andrea Rodríguez-López [1, 2], El Chérif Ibrahim [3], Victor Gorgievski [4], Miquel Martin [5], Patricia Robledo [6], Rafael Maldonado [1, 6, 7], *, Eleni Tzavara [4, 8], *, Beltrán Álvarez-Pérez [1], *

[1] Laboratory of Neuropharmacology-NeuroPhar, Department of Medicine and Life Sciences, Universitat Pompeu Fabra, Barcelona Biomedical Research Park, 08003 Barcelona, Spain

[2] Present address: Laboratory of Neurobiology of Behavior, Department of Medicine and Life Sciences, Universitat Pompeu Fabra, Barcelona Biomedical Research Park, 08003 Barcelona, Spain

[3] Aix-Marseille Univ, CNRS, INT, Inst Neurosci Timone, 13005 Marseille, France

[4] Université Paris Cité, Inserm, CNRS, HealthFex, F-75006 Paris, France

[5] Department of Morphological Sciences, Unit of Human Anatomy and Embryology, Faculty of Medicine, Universitat Autònoma de Barcelona, Bellaterra, Spain.

[6] Hospital del Mar Medical Research Institute (IMIM), Barcelona, Spain

[7] Université de Strasbourg (UNISTRA), INSERM UMR-S 1329, Strasbourg Translational Neuroscience and Psychiatry, Centre de Recherche en Biomédecine de Strasbourg, France

[8] Hôpital Sainte Marguerite AP-HM, Pôle de Psychiatrie, 13274 Marseille, France

*equal contribution

Correspondence to:

Beltrán Álvarez-Pérez (beltran.alvarez@upf.edu); Rafael Maldonado (rafael.maldonado@upf.edu); Eleni Tzavara (eleni.tzavara@inserm.fr)

Abstract

Major depressive disorder (MDD) is a complex psychiatric condition with significant individual

and social impact and marked variability in treatment response. Identifying biomarkers that

predict antidepressant efficacy remains a major challenge, and metabolomic approaches offer a

powerful tool to uncover biochemical changes associated with depression and treatment

outcome. The prefrontal cortex and the hippocampus are key brain regions involved in the

pathophysiology of MDD and antidepressant response. Fluoxetine, one of the most commonly

prescribed antidepressants, influences not only serotonergic signaling but also the

endocannabinoid system. Accordingly, evaluating endocannabinoid levels in these brain regions

might provide critical insight into their contribution to mood disorders and the effect of

fluoxetine. To identify biomarkers associated with early antidepressant response, male mice

were exposed for 5 weeks to the unpredictable chronic mild stress (UCMS) protocol, with

fluoxetine treatment administered during the final week. Liquid chromatography–tandem mass

spectrometry analyses of hippocampal and prefrontal cortex samples revealed that the

depression-like phenotype induced by UCMS was associated with increased hippocampal

anandamide (AEA) levels. Behavioral improvement following short-term fluoxetine treatment

was accompanied by enhanced hippocampal 2-arachidonoylglycerol (2-AG), which was

associated with reduced GABA and glutamate levels in the prefrontal cortex. These findings

suggest that increased 2-AG levels may serve as a predictor of early response to fluoxetine.

Introduction

Depressive disorders include a cluster of diseases characterized by the presence of sad, empty, and/or irritable mood, associated with alterations that significantly affect the individual's ability to function [1]. Endocannabinoids are central in modulating various neurotransmitter systems [2], including glutamatergic [3], GABAergic, and monoaminergic [4,5] signaling, all of which are implicated in the pathogenesis of major depressive disorder (MDD) [6]. Altered levels of the arachidonic acid derivatives, anandamide (AEA) and 2-arachidonoylglycerol (2-AG), have been associated with stress vulnerability and impaired emotional regulation [2,7]. According to its duration and effects on daily activity, MDD is the most disruptive mental health problem [8]. Almost half of the patients do not respond to antidepressant medication [9,10], which is associated with increased healthcare utilization and cost [11]. Treatments avoiding relapse and achieving full remission remain a major challenge. Selective serotonin reuptake inhibitors (SSRIs), including fluoxetine, are considered a first-line option [12]. Fluoxetine has high acceptability [13,14] and beneficial off-target effects on noradrenaline and dopamine [15], inflammation [16] and the endocannabinoid system [17,18]. However, it has a high variability of response [13], which is a setback in clinical practices, where every additional treatment step decreases the probability of remission and response success [19]. Unlike other SSRIs, fluoxetine also increases noradrenaline and dopamine extracellular levels in the prefrontal cortex [15] through the blockade of 5-HT2C receptors on cortical interneurons [20]. Nonetheless, it is crucial to differentiate acute and chronic effects. Acute fluoxetine increases extracellular serotonin by blocking the transporter [15,21,22]. However, repeated treatment causes an excess of extracellular serotonin that triggers 5-HT1A autoreceptor activity, which subsequently reduces serotonin synthesis and turnover [22,23] and lower serotonin intracellular levels [21]. Yet, chronic fluoxetine treatment (> 2 weeks) leads to 5-HT1A autoreceptor desensitization [23] and restoration of serotonin intracellular levels [21]. Advanced metabolomic techniques, such as liquid chromatography-tandem mass spectrometry (LC-MS/MS), would provide new insight into the role of serotonin and other neurotransmitters in mood disorders and rapid molecular changes in response to fluoxetine. Therefore, the identification of biomarkers of early response to fluoxetine could potentially facilitate and support clinical practice in psychiatry. The unpredictable chronic mild stress (UCMS) model is a reliable mouse model that closely mimics the behavioral deterioration and molecular alterations observed in MDD patients [24]. Our objective was to identify metabolomic alterations associated with short-term fluoxetine response, in two key brain areas playing a key role in this disorder, the hippocampus and the prefrontal cortex [25–28].

Material and methods

Animals and ethics

Male C57BL/6J (8 weeks old) (n = 60) were purchased (Charles River, France). Except for control (n = 8) (CONTROL-VEH), which were grouped in 4, mice were individually housed in Plexiglas cages and maintained in an environment with controlled temperature ($21 \pm 1 \,°C$) and humidity ($55 \pm 10\%$). Food and water were available *ad libitum*. All behavioral experiments were performed during the light phase of a 12-hour cycle (lights on at 7.30 am and off at 7.30 pm). Animal procedures and care were conducted under the European Communities Directive 2010/63/EU and the Spanish Royal Decree 53/201RD, following "Animals in Research: Reporting Experiments" (ARRIVE) guidelines, and approved by the local (Comitè Ètic d'Experimentació Animal-Parc de Recerca Biomèdica de Barcelona, CEEA-PRBB) and autonomous (Generalitat de Catalunya, Departament de Territori i Sostenibilitat) ethical committee. All behavioral and biochemical experiments were performed by a researcher blind to experimental conditions.

Drug administration

Fluoxetine hydrochloride (0927, Tocris) dissolved in saline (0.9 % NaCl) or vehicle (0.9 % NaCl) was injected intraperitoneally (i.p.) at 10 mg/kg daily from 2 pm to 4 pm for 7 consecutive days.

Unpredictable chronic mild stress protocol

After a week of acclimation, mice were subjected to 5 weeks of UCMS as previously described [29]. Stressors were applied twice daily during a 3-h period and overnight in a randomized order. Daytime stressors included: wet bedding, 45°-tilted cages, pair or crowded housing, water deprivation, and confinement in a reduced space with a grid surface, restricted food access, and cage interchange. Overnight stressors included: flashing lights, inversion of the light-dark cycle, 45°-tilted cages, and wet bedding. Before the start of the protocol and once a week afterwards, the animals' weight and apathy (self-grooming and coat state) were evaluated. After 4 weeks, animals under the UCMS protocol were randomly assigned to fluoxetine (UCMS-FLUOX, n = 40) or vehicle (UCMS-VEH, n = 10) groups (Figure 1-(a)). At the end of the experimental protocol and 24 h after the last administration, animals were euthanized by cervical dislocation. After decapitation, brains were removed, placed over ice, and the prefrontal cortex and hippocampus were dissected, weighed, frozen in liquid nitrogen, and stored at -80ºC.

Apathy

Fur coat and grooming time were measured as indicators of apathy [24]. The score of the animal's coat state was calculated as the sum of scores, one for each of 7 different body parts (head,

111 neck, dorsal, ventral, tail, forepaws, hind paws). Each part was given a score on a 3-point scale

112 (1 = poor, 2 = fair, 3 = very good) (26). In the splash test, nesting material was temporarily

113 removed, and animals were splashed with a 20 % glycerol solution on the dorsal surface. Self-

114 directed grooming behavior was recorded for 5 min, and latency, total number of events, and

115 time spent self-grooming were manually measured.

Behavioral despair

Behavioral despair was measured with the forced swim test (FST) [30] after 4 weeks of UCMS

and the tail suspension test (TST) at the end of the experimental protocol. In the FST, mice were

placed in water-filled cylinders (30 cm high and 15 cm diameter) for 6 min. Immobility time,

considered when mice floated in an upright position and made only small movements to keep

their head above, was manually measured during the last 4 min [31,32]. In the TST, mice were

individually suspended by the tail using adhesive tape placed 1 cm from the tip of the tail and

recorded during a 6-min session. Immobility time, considered as the complete absence of

movement, was registered from the last 4 min [31]. Animals falling or climbing over their tails

were discarded.

Quantification of brain endocannabinoids by LC-MS/MS

AEA and 2-AG were quantified as previously described [33] with modifications. Briefly, one

hippocampal hemisphere (21.2 ± 6.43 mg) was homogenized in 700 µL of 50 mM Tris–HCl

(pH 7.4):methanol (1:1) containing deuterated internal standards (AEA-d4, DHEA-d4, 2-AG-

d5, and 2-OG-d5; Cayman Chemical and Toronto Research Chemicals). Chloroform (5 mL)

was added, samples were shaken for 20 min and centrifuged (1700 g, 5 min, room temperature).

The organic phase was evaporated under nitrogen, reconstituted in 100 µL water:acetonitrile

(10:90, v/v) with 0.1% formic acid, and transferred to microvials for LC-MS/MS analysis.

Analyses were performed using a Waters Acquity UPLC system coupled to a Xevo TQ-S micro

mass spectrometer. Chromatographic separation was achieved on a BEH C18 column (2.1 × 100

mm, 1.8 µm) at 40ºC, with a flow rate of 0.4 mL/min and a mobile phase consisting of 0.01%

formic acid in water and acetonitrile. Detection was carried out in multiple reaction monitoring

(MRM) mode, and quantification was performed by isotope dilution.

Quantification of brain metabolites by LC-MS/MS

The remaining hippocampal tissue (21.06 ± 7.16 mg) and prefrontal cortex samples (26.35 ±

8.99 mg) were homogenized in acetonitrile:water (2:1, 0.1% formic acid; 25 mg tissue/600 µL).

After centrifugation (14,500 rpm, 5 min, 4ºC), supernatants were collected and processed using

two analytical procedures.

For GABA and glutamate determination (method A), aliquots (10 µL) were mixed with labelled
internal standards (GABA-d6 and Glu-d5) and acetonitrile, vortexed, centrifuged, and
transferred to microvials. For monoamines, amino acids, and metabolites (method B), larger
aliquots (300 µL) were combined with a labelled internal standard mixture, evaporated under
nitrogen, and derivatized with dansyl chloride in alkaline conditions prior to LC-MS/MS
analysis.

Analyses were conducted on an Acquity I-Class UPLC coupled to a Xevo TQ-S micro mass
spectrometer equipped with an electrospray ionization source operating in positive mode.
Source parameters were set as follows: capillary voltage 3 kV, source temperature 150ºC,
desolvation temperature 600ºC, desolvation gas flow 1200 L/h, and cone gas flow 50 L/h.
Nitrogen was used as nebulizing and drying gas, and argon as collision gas. Data acquisition
and processing were performed using MassLynx v4.1.

For method A, separation was achieved on a BEH Amide column using an acetonitrile/water
gradient containing ammonium formate and formic acid. For method B, a BEH C18 column and
a methanol/water gradient were employed. MRM transitions for each analyte are reported in
Table 1. Calibration curves were generated at 12 concentration levels using external standards
and processed with TargetLynx XS software.

Table 1. MRM transitions used for method B.

Analyte	Precursor ion m/z	CV (V)	Product ion m/z	CE (eV)
5-HT	643	15	170, 408	35, 30
DA	853	15	170, 619	45, 35
NA	869	15	171, 617	45, 35
Gly	309	15	157, 170	30, 15
5-HIAA	425	15	171, 381	30, 20
DOPAC	635	15	170, 401	30, 30
HVA	416	15	156, 170	30, 20
3-MeO-Tyr	634	15	170, 400	35, 35
Phe	399	15	120, 157	20, 30
Tyr	648	15	369, 414	30, 30
Trp	438	15	159, 171	30, 30
5-HT-d4	647	15	412	30
DA-d4	857	15	623	35
NA-d6	875	15	623	35
Gly-13C2,15N	312	15	157	30

5-HIAA-d4	429	15	171	30
DOPAC-d3	638	15	170	30
HVA-13C,d3	420	15	156	30
3-MeO-Tyr-d4	638	15	170	35
Phe-d5	404	15	157	30
Tyr-d4	652	15	373	30
Trp-d5	443	15	159	30

Statistical analysis

Behavioral and metabolomic results are presented as mean ± standard error of the mean (s.e.m).
Based on previous evidence that distinguishes fluoxetine treatment responders from non-
responders [14,34], a global index score was calculated using Z-normalized values of total
grooming time, fur coat score, and body weight. The UCMS-FLUOX group was classified into
responders and non-responders according to the lowest global index value observed in
CONTROL-VEH mice. Serotonin and dopamine turnover were estimated using their respective
metabolite ratios. Data normality was assessed using the Shapiro–Wilk test. Statistical analyses
were performed using two-way ANOVA or mixed-effects models for repeated measures, with
significance set at $p < 0.05$. When appropriate, Bonferroni or Fisher's LSD post hoc tests were
applied. Outliers, defined as values exceeding ±2 standard deviations from the mean, were
excluded. Data analysis and visualization were conducted using IBM SPSS v23 and GraphPad
Prism v10. Pearson's correlation analyses were performed in R (v4.4.1) and visualized using the
ggcorrplot package (v0.1.4.1).

Results

Effects of short-term exposure to fluoxetine on stress-induced depressive-like behavior

Physical condition and behavior were evaluated to assess the effects of UCMS and short-term fluoxetine treatment. For body weight, time, and group differences after mixed-effects analysis were observed (time x group, $F_{[4, 108]} = 4.287$, $p = 0.0029$). Animals exposed to UCMS showed significantly lower body weights than control mice (CONTROL-VEH. *vs.* UCMS-VEH. $p = 0.0123$) after 4 weeks. Fluoxetine did not significantly reverse these effects (CONTROL-VEH. *vs.* UCMS-VEH. $p = 0.0181$; CONTROL-VEH. *vs.* UCMS-FLUOX. $p = 0.0383$) (Figure 1-(b)). There were fur coat time and group differences (time: $F_{[3.333, 181.6]} = 15.17$, $p < 0.0001$; group: $F_{[2, 55]} = 9.347$, $p = 0.0003$; time x group: $F_{[8, 218]} = 6.133$, $p < 0.0001$). Fur was significantly degraded in stressed animals after 4 weeks (CONTROL-VEH. *vs.* UCMS-VEH. $p = 0.0086$; CONTROL-VEH. *vs.* UCMS-FLUOX. $p < 0.0001$), and fluoxetine had no influence (Figure 1-(c)). Mixed-effects analysis showed significant group differences in the total grooming time ($F_{[2, 55]} = 32.61$, $p < 0.0001$). Bonferroni's multiple comparison test exposed no basal differences, but statistically significant differences before treatment between stressed and control animals (CONTROL-VEH. *vs.* UCMS-VEH. $p < 0.0001$, CONTROL-VEH. *vs.* UCMS-FLUOX. $p = 0.0001$). Fluoxetine partially increased total grooming time (UCMS-VEH. *vs.* UCMS-FLUOX. $p = 0.0033$), although not comparable to control animals (CONTROL-VEH. *vs.* UCMS-FLUOX. $p = 0.0252$) (Figure 1-(d)). Immobility time was increased in UCMS-exposed animals (one-way ANOVA $F_{[2, 54]} = 9.42$, $p = 0.0003$; CONTROL-VEH.: *vs.* UCMS-VEH. $p = 0.0005$; and *vs.* UCMS-FLUOX. $p = 0.0007$) (Figure 1-(e)). Short-term fluoxetine treatment significantly decreased immobility time in the TST (one-way ANOVA $F_{[2, 41]} = 7.153$, $p = 0.0022$, UCMS-VEH. *vs.* UCMS-FLUOX. $p = 0.0052$) (Figure 1-(f)). One-way ANOVA in global index (Figure 1-(g)) revealed significant group differences ($F_{[2, 54]} = 29.14$, $p < 0.0001$; CONTROL-VEH. *vs.* UCMS-VEH. $p < 0.0001$, CONTROL-VEH. *vs.* UCMS-FLUOX. $p < 0.0001$; UCMS-VEH. *vs.* UCMS-FLUOX. $p = 0.0027$). Therefore, UCMS induced significant physical and emotional deterioration, which was only partially reversed by short-term fluoxetine treatment. The global index helped distinguish responders from non-responders to treatment.

UCMS exposure and short-term fluoxetine modulate endocannabinoid levels in the hippocampus

Stress and fluoxetine treatment significantly influenced endocannabinoids in the hippocampus (one-way ANOVA for 2-AG: $F_{[3, 51]} = 5.645$, $p = 0.002$; and AEA: $F_{[3,51]} = 6.582$, $p = 0.0008$) (Figure 2-(a), and (b)). Behavioral improvement by short-term fluoxetine treatment in

211   responders was associated with increased 2-AG levels (p = 0.0016 to CONTROL-VEH.; p =
212   0.0005 to UCMS-VEH.; p = 0.0123 to non-responders) (Figure 2-(a)). In UCMS-VEH. animals,
213   AEA levels were significantly increased by stress alone (p = 0.0198 to CONTROL-VEH.),
214   whereas fluoxetine decreased them in both responders and non-responders (responders: p <
215   0.0001; and non-responders p = 0.0007) (Figure 2-(b)).

UCMS and short-term fluoxetine exposure modulate neurotransmitter levels in the hippocampus and the prefrontal cortex

Fluoxetine treatment significantly decreased the serotonin pool (one-way ANOVA F[3, 51] =
4.456, p = 0.0074) (Figure 3-(a)) and turnover (one-way ANOVA F[3, 50] = 27.29, p < 0.0001)
(Figure 3-(b)). Dopamine also decreased in responders compared to CONTROL-VEH. animals
(one-way ANOVA F[3, 51] = 3.725, p = 0.0169, p = 0.0221) (Figure 3-(c)), but the turnover
was not affected (Figure 3-(d)). There were no differences in noradrenaline (Figure 3-(e)),
glutamate (Figure 3-(f)), GABA (Figure 3-(g)), nor glycine (Figure 3-(h)) in the hippocampus.
Stress does not influence neurotransmitters in the hippocampus.

Stress did not significantly alter serotonin nor the turnover in the prefrontal cortex, while
fluoxetine significantly decreased them (levels: one-way ANOVA F[3, 50] = 10.47, responders
and non-responders to CONTROL-VEH. p < 0.0001, and to UCMS-VEH. p < 0.0001; turnover:
one-way ANOVA F[3, 50] = 42.83, p < 0.0001, to CONTROL-VEH. responders p = 0.0233,
and non-responders p = 0.0172, to UCMS-VEH. responders p = 0.0233, and non-responders p
= 0.0172) (Figure 4-(a) and (b)). Oppositely, dopamine and its turnover were not altered by
stress or by short-term fluoxetine (Figure 4-(c) and (d)). As seen in the hippocampus,
noradrenaline levels are unmodified by chronic stress or fluoxetine (Figure 4-(e)). In the
prefrontal cortex, glutamate was decreased by fluoxetine in responders (one-way ANOVA F[3,
50] = 4.231, p = 0.0097, *vs.* CONTROL-VEH.: p = 0.0094, *vs.* UCMS-VEH.: p = 0.0019, *vs.*
non-responders p = 0.0476 by uncorrected Fisher's LSD) (Figure 4-(f)). Likewise, GABA was
decreased by fluoxetine in responders (one-way ANOVA F[3, 50] = 4.489, p = 0.0072, vs.
CONTROL-VEH p = 0.0233; vs. UCMS-VEH p = 0.0009), with a marked tendency compared
to non-responders (p = 0.0594) (Figure 4-(g)). On the other hand, glycine levels were
significantly decreased by fluoxetine to the same extent in responders and non-responders (one-
way ANOVA F[3, 50] = 4.633, p = 0.0062; UCMS-VEH. vs. responders p = 0.0198; vs. non-
responders p = 0.006) (Figure 4-(h)).

Correlations between neurotransmitter levels in the hippocampus and prefrontal cortex

Neurotransmitter interactions are altered in stress and after antidepressant treatment. We
analyzed how UCMS and fluoxetine affect them in the hippocampus and prefrontal cortex.

Pearson correlations revealed that under control conditions, the concentrations of most
neurotransmitters in the hippocampus were significantly positively correlated among them
(Figure 5-(a)) and were significantly reduced in stressed mice (Figure 5-(b)). Thus,
endocannabinoid concentrations no longer correlated with monoamines or excitatory and
inhibitory neurotransmitters; the only remaining correlations were between serotonin and
monoamines, GABA, glutamate, and glycine. Fewer associations between neurotransmitter
levels were observed in the prefrontal cortex of control mice (Figure 5-(a)), while the number of
correlations was higher in stressed mice, mostly between inhibitory and excitatory
neurotransmitters and with serotonin (Figure 5-(b)).

Short-term fluoxetine induced a distinct pattern of associations in the hippocampus of
responders and non-responders. In responders, strong positive correlations were observed,
mainly in the hippocampus. Notably, 2-AG concentrations correlated negatively with AEA in
the hippocampus and with serotonin and noradrenaline in the prefrontal cortex (Figure 5-(c)). In
contrast, in non-responders, dopamine positively correlated with all the other neurotransmitters
in the hippocampus (except serotonin) but not in the prefrontal cortex (Figure 5-(d)). In
addition, the strong negative correlation between 2-AG and AEA found in the hippocampus of
responders is diminished, and the negative correlations between 2-AG and serotonin and
noradrenaline in the prefrontal cortex are lost. In contrast to what was observed in responders,
AEA concentrations positively correlate with most of the hippocampal prefrontal cortex
neurotransmitters (Figure 5-(d)). In the prefrontal cortex of responders, a few significant
associations between neurotransmitters were observed (Figure 5-(c)), opposite to non-
responders, where most neurotransmitters were correlated with each other in the prefrontal
cortex and with neurotransmitters in the hippocampus (Figure 5-(d)).

Stress disrupted hippocampal neurotransmitter associations while increasing correlations in the
prefrontal cortex. Fluoxetine partially restored hippocampal correlations in responders, whereas
non-responders showed distinct patterns, including altered endocannabinoid interactions,
suggesting differential neurochemical adaptations to treatment.

Discussion

Randomized application of mild stressors for 4 weeks resulted in increased despair, physical
deterioration, and reduced body weight. These behavioral and somatic changes are comparable
to the depressive-like phenotypic alterations previously described using similar UCMS
protocols [29,34–36] and demonstrate the reliability of the model. Next, fluoxetine was administered
daily for 7 consecutive days while continuing with the UCMS protocol. The selected dose of 10
mg/kg has previously been demonstrated to be effective in increasing serotonin, dopamine, and
noradrenaline extracellular levels after acute administration [15] and to have an antidepressant
effect after acute [21] and chronic administrations [37]. In the present work, after short-term
fluoxetine treatment, animals had a less pronounced depressive-like phenotype compared to
vehicle-treated stressed mice. This effect was revealed by a healthier coat state and an increased
grooming time, indicative of decreased signs of apathy, and an increased mobility time in the
TST. These behavioral effects observed after 1 week of treatment at a dose of 10 mg/kg were
comparable to those observed with higher doses and longer administration protocols [29,34,38],
suggesting that a shorter treatment protocol is sufficient to elicit antidepressant-like effects in
this animal model of depression. Previous preclinical studies demonstrated that chronic
administration of fluoxetine distinguished two populations: those showing improved behavioral
outcomes (responders) and those that did not experience a significant improvement (non-
responders) [34]. We, as previously reported [34], used total grooming time in the splash test, total
fur coat score, and body weight as variables to distinguish these two populations. The resulting
proportion of responders to non-responders in the present study (9 out of 39) is lower than the
one found in previous literature (9 out of 15) [34]. This discrepancy might be explained by the
differential classification methodologies, the duration of the treatment, and/or the dose of
fluoxetine. While our study employed short-term treatment, previous research has often used
longer treatment periods, typically ranging from 3 to 6 weeks, to assess antidepressant-like
effects [29,34,38]. Working with shorter treatment durations allows us to expedite the treatment
definition process, as behavioral effects may take longer to manifest [13] the underlying molecular
changes occur more quickly. This enables us to identify the effects more rapidly.

The endocannabinoid system is implicated in the pathogenesis of depression and the mechanism
of action of antidepressants [39], and in turn, cannabinoids can elicit antidepressant effects [40]. The
hippocampus is one of the brain regions with the highest CB1R density [41,42] associated with
emotional and stress responses [43]. Acute stress results in increased hippocampal 2-AG [7] and
decreased AEA concentrations, reflecting a divergent response to stressors [2]. Therefore, we
analyzed changes promoted by chronic stress. UCMS did not affect 2-AG hippocampal content,
but increased AEA compared to undisturbed mice. Previous studies have reported contradictory
results. Indeed, chronic stress promoted hippocampal metabolic pathways associated with AEA

degradation [44], although another study reported no alterations in hippocampal AEA following UCMS [45]. The main differences between these previous studies and ours are the duration of the protocol (3 *vs.* 6 weeks) and the time gap between UCMS discontinuation and sacrifice (1 week *vs.* no time gap). Moreover, since samples were collected 24 h after the TST, it could be possible that the observed AEA increase in our study could be the result of an acute response to the TST stressor. These conflicting data suggest that, unlike the known effects of acute stress on AEA in the hippocampus [46,47], chronic stress effects on AEA must be further clarified. Monoaminergic dysfunction has been proposed as a crucial pathophysiological mechanism for depression [48]. In our study, no significant alterations were observed as a result of chronic stress-induced depression, as previously reported [35,36]. Unaltered serotonin levels might be the result of acute stress due to TST exposure on serotonin release [49], but also of the decreased expression of tyrosine hydroxylase and tryptophan hydroxylase enzymes [36].

In the hippocampus of control animals, endocannabinoid concentrations were positively correlated with the levels of most neurotransmitters. In contrast, exposure to UCMS disrupted these relationships, leading to an imbalance between inhibitory and excitatory neurotransmitters. These findings suggest that the modulatory role of the endocannabinoid system is impaired in chronic stress, consistent with previous reports [2], and might contribute to the dysregulation of other neurotransmitter systems. In addition, the endocannabinoid system is involved in synaptic plasticity processes [50], and chronic stress is known to alter synaptic strength and connectivity within the hippocampus [43]. Supporting this view, previous literature linked preservation of proteins involved in vesicular functioning to stress-coping strategies [51]. Taken together, dysfunction of the endocannabinoid system would impact hippocampal neurotransmission and synaptic plasticity, ultimately contributing to stress-related alterations.

Short-term fluoxetine significantly increased hippocampal 2-AG, mainly in mice classified as responders, suggesting that high 2-AG levels are associated with better short-term response. Excitatory neurotransmission could be involved in these changes since rapid fluoxetine-induced serotonin increase evokes endocannabinoid release through 5-HT2R activation [52], and endocannabinoids can act retrogradely to suppress glutamate release [53]. In addition, chronic fluoxetine treatment results in an increase in CB1R binding capacity [54], strengthening endocannabinoid signaling. These findings suggest that fluoxetine effects may be partly mediated by endocannabinoid modulation, particularly through 2-AG, which influences excitatory neurotransmission. As expected, monoamine concentrations in the hippocampus and the prefrontal cortex were affected by fluoxetine. Serotonin decreased, in line with previous results [15,35], an effect likely mediated by the early inhibition of the 5-HT1A autoreceptors on serotonin synthesis [22,23]. Fluoxetine also led to a comparable decrease in serotonin turnover, indicative of reduced MAO activity [55,56], supporting the notion that fluoxetine not only inhibits

serotonin reuptake but also modulates the enzymatic pathways. There is consensus that GABA

and glutamate are decreased in the cerebrospinal fluid [57,58] and brain [59] of patients and

preclinical models [44,60] of MDD. The prefrontal cortex is highly sensitive to stress [61], which

receives enhanced input from subcortical areas whose neuronal activity is impaired during

chronic stress [62,63], resulting in GABAergic and glutamatergic imbalance and impairment of

affective behavior [64]. The present results show that GABA and glutamate are decreased in the

prefrontal cortex of responders, suggesting their association with a better behavioral response.

This effect could be mediated by the alteration of vesicle transport dynamics [65]. Nonetheless,

fluoxetine can also stimulate GABA and glutamate neurotransmission by serotonin activation of

5-HT2AR on GABAergic interneurons and glutamatergic neurons [66], suggesting that fluoxetine

could have multiple effects.

Therefore, the depression-like phenotype induced by UCMS was associated with increased

hippocampal AEA. Behavioral improvement with short-term fluoxetine was accompanied by

increased hippocampal 2-AG, associated with decreased GABA and glutamate in the prefrontal

cortex.

Data Availability

All data are available in the main text or supplementary materials. Correspondence and requests for materials should be addressed to Beltrán Álvarez-Pérez, Eleni Tzavara or Rafael Maldonado.

Bibliography

American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders.* Washington DC, 2022.

Morena M, Patel S, Bains JS, Hill MN. Neurobiological Interactions Between Stress and the Endocannabinoid System. *Neuropsychopharmacol* 2016; **41**: 80–102.

Tzavara ET, Degroot A, Wade MR, Davis RJ, Nomikos GG. CB1 receptor knockout mice are hyporesponsive to the behavior-stimulating actions of d-amphetamine: Role of mGlu5 receptors. *European Neuropsychopharmacology* 2009; **19**: 196–204.

Tzavara ET, Li DL, Moutsimilli L, Bisogno T, Di Marzo V, Phebus LA *et al.* Endocannabinoids Activate Transient Receptor Potential Vanilloid 1 Receptors to Reduce Hyperdopaminergia-Related Hyperactivity: Therapeutic Implications. *Biological Psychiatry* 2006; **59**: 508–515.

El Khoury M-A, Gorgievski V, Moutsimilli L, Giros B, Tzavara ET. Interactions between the cannabinoid and dopaminergic systems: Evidence from animal studies. *Progress in Neuro-Psychopharmacology and Biological Psychiatry* 2012; **38**: 36–50.

Filatova EV, Shadrina MI, Slominsky PA. Major Depression: One Brain, One Disease, One Set of Intertwined Processes. *Cells* 2021; **10**: 1283.

Busquets-Garcia A, Gomis-González M, Srivastava RK, Cutando L, Ortega-Alvaro A, Ruehle S *et al.* Peripheral and central CB1 cannabinoid receptors control stress-induced impairment of memory consolidation. *Proc Natl Acad Sci USA* 2016; **113**: 9904–9909.

Culpepper L, Martin A, Nabulsi N, Parikh M. The Humanistic and Economic Burden Associated with Major Depressive Disorder: A Retrospective Cross-Sectional Analysis. *Adv Ther* 2024; **41**: 1860–1884.

Santomauro DF, Vos T, Whiteford HA, Chisholm D, Saxena S, Ferrari AJ. Service coverage for major depressive disorder: estimated rates of minimally adequate treatment for 204 countries and territories in 2021. *The Lancet Psychiatry* 2024; **11**: 1012–1021.

Thomas L, Kessler D, Campbell J, Morrison J, Peters TJ, Williams C *et al.* Prevalence of treatment-resistant depression in primary care: cross-sectional data. *Br J Gen Pract* 2013; **63**: e852–e858.

Sharman Moser S, Chodick G, Gelerstein S, Barit Ben David N, Shalev V, Stein-Reisner O. Epidemiology of treatment resistant depression among major depressive disorder patients in Israel. *BMC Psychiatry* 2022; **22**: 541.

American Psychological Association. 2019. Clinical practice guideline for the treatment of depression across three age cohorts. https://www.apa.org/depression-guideline.

Cipriani A, Furukawa TA, Salanti G, Chaimani A, Atkinson LZ, Ogawa Y *et al.* Comparative efficacy and acceptability of 21 antidepressant drugs for the acute treatment of adults with major depressive disorder: a systematic review and network meta-analysis. *The Lancet* 2018; **391**: 1357–1366.

Magni LR, Purgato M, Gastaldon C, Papola D, Furukawa TA, Cipriani A *et al.* Fluoxetine versus other types of pharmacotherapy for depression. *Cochrane Database of Systematic Reviews* 2013; **2013**. doi:10.1002/14651858.CD004185.pub3.

15 Koch S. R-fluoxetine Increases Extracellular DA, NE, As Well As 5-HT in Rat Prefrontal
Cortex and Hypothalamus An in vivo Microdialysis and Receptor Binding Study.
Neuropsychopharmacology 2002; **27**: 949–959.

16 Chen H, Zhao X, Ma X, Ma H, Zhou C, Zhang Y *et al.* Effects of esketamine and
fluoxetine on depression-like behaviors in chronic variable stress: a role of plasma
inflammatory factors. *Front Psychiatry* 2024; **15**: 1388946.

17 Hill MN, Ho W-SV, Hillard CJ, Gorzalka BB. Differential effects of the antidepressants
tranylcypromine and fluoxetine on limbic cannabinoid receptor binding and
endocannabinoid contents. *J Neural Transm* 2008; **115**: 1673–1679.

18 Umathe SN, Manna SSS, Jain NS. Involvement of endocannabinoids in antidepressant and
anti-compulsive effect of fluoxetine in mice. *Behavioural Brain Research* 2011; **223**: 125–
134.

19 John Rush A, Trivedi MH, Wisniewski SR, Nierenberg AA, Stewart JW, Warden D *et al.*
Acute and Longer-Term Outcomes in Depressed Outpatients Requiring One or Several
Treatment Steps: A STAR*D Report. 2006.

20 Gobert A, Rivet J-M, Lejeune F, Newman-Tancredi A, Adhumeau-Auclair A, Nicolas J-P
et al. Serotonin2C receptors tonically suppress the activity of mesocortical dopaminergic
and adrenergic, but not serotonergic, pathways: A combined dialysis and
electrophysiological analysis in the rat. *Synapse* 2000; **36**: 205–221.

21 Bianchi M, Moser C, Lazzarini C, Vecchiato E, Crespi F. Forced swimming test and
fluoxetine treatment: in vivo evidence that peripheral 5-HT in rat platelet-rich plasma
mirrors cerebral extracellular 5-HT levels, whilst 5-HT in isolated platelets mirrors
neuronal 5-HT changes. *Exp Brain Res* 2002; **143**: 191–197.

22 Hervás I, Queiroz CMT, Adell A, Artigas F. Role of uptake inhibition and autoreceptor
activation in the control of 5□HT release in the frontal cortex and dorsal hippocampus of
the rat. *British J Pharmacology* 2000; **130**: 160–166.

23 Kaufman J, DeLorenzo C, Choudhury S, Parsey RV. The 5-HT1A receptor in Major
Depressive Disorder. *European Neuropsychopharmacology* 2016; **26**: 397–410.

24 Planchez B, Surget A, Belzung C. Animal models of major depression: drawbacks and
challenges. *Journal of Neural Transmission* 2019; **126**: 1383–1408.

25 MacQueen GM, Yucel K, Taylor VH, Macdonald K, Joffe R. Posterior Hippocampal
Volumes Are Associated with Remission Rates in Patients with Major Depressive Disorder.
Biological Psychiatry 2008; **64**: 880–883.

26 Otte C, Gold SM, Penninx BW, Pariante CM, Etkin A, Fava M *et al.* Major depressive
disorder. *Nature Reviews Disease Primers* 2016; **2**: 16065.

27 Koolschijn PCMP, Van Haren NEM, Lensvelt□Mulders GJLM, Hulshoff Pol HE, Kahn
RS. Brain volume abnormalities in major depressive disorder: A meta□analysis of
magnetic resonance imaging studies. *Human Brain Mapping* 2009; **30**: 3719–3735.

28 Savitz J, Drevets WC. Bipolar and major depressive disorder: Neuroimaging the
developmental-degenerative divide. *Neuroscience & Biobehavioral Reviews* 2009; **33**: 699–
771.

29   Farley S, Apazoglou K, Witkin JM, Giros B, Tzavara ET. Antidepressant-like effects of an
AMPA receptor potentiator under a chronic mild stress paradigm. *International Journal of*
Neuropsychopharmacology 2010; **13**: 1207–1218.

30   Porsolt R. D, Bertin A, Jalfre M. Behavioral despair in mice: a primary screening test for
antidepressants. *Archives Internationales de Pharmacodynamie et de Thérapie* 1977; **229**:
327–336.

31   Can A, Dao DT, Arad M, Terrillion CE, Piantadosi SC, Gould TD. The Mouse Forced
Swim Test. *JoVE* 2011; : 3638.

32   Bura SA, Cabañero D, Maldonado R. Operant self☐administration of pregabalin in a mouse
model of neuropathic pain. *European Journal of Pain* 2018; **22**: 763–773.

33   Navarro-Romero A, Galera-López L, Ortiz-Romero P, Llorente-Ovejero A, De Los Reyes-
Ramírez L, Bengoetxea De Tena I *et al.* Cannabinoid signaling modulation through JZL184
restores key phenotypes of a mouse model for Williams–Beuren syndrome. *eLife* 2022; **11**:
e72560.

34   Apazoglou K, Farley S, Gorgievski V, Belzeaux R, Lopez JP, Grenier J *et al.*
Antidepressive effects of targeting ELK-1 signal transduction. *Nature Medicine* 2018; **24**:
591–597.

35   Estévez-Cabrera MM, Sánchez-Muñoz F, Pérez-Sánchez G, Pavón L, Hernández-
Díazcouder A, Córtes Altamirano JL *et al.* Therapeutic treatment with fluoxetine using the
chronic unpredictable stress model induces changes in neurotransmitters and circulating
miRNAs in extracellular vesicles. *Heliyon* 2023; **9**: e13442.

36   Lu Q, Mouri A, Yang Y, Kunisawa K, Teshigawara T, Hirakawa M *et al.* Chronic
unpredictable mild stress-induced behavioral changes are coupled with dopaminergic
hyperfunction and serotonergic hypofunction in mouse models of depression. *Behavioural*
Brain Research 2019; **372**: 112053.

37   Fang Y, Ding X, Zhang Y, Cai L, Ge Y, Ma K *et al.* Fluoxetine inhibited the activation of
A1 reactive astrocyte in a mouse model of major depressive disorder through astrocytic 5-
HT2BR/β-arrestin2 pathway. *J Neuroinflammation* 2022; **19**: 23.

38   Zhang C-L, Li Y-J, Lu S, Zhang T, Xiao R, Luo H-R *et al.* Fluoxetine ameliorates
depressive symptoms by regulating lncRNA expression in the mouse hippocampus.
Zoological Research 2021; **42**: 28–42.

39   Micale V, Di Marzo V, Sulcova A, Wotjak CT, Drago F. Endocannabinoid system and
mood disorders: Priming a target for new therapies. *Pharmacology & Therapeutics* 2013;
138: 18–37.

40   Bambico FR, Katz N, Debonnel G, Gobbi G. Cannabinoids Elicit Antidepressant-Like
Behavior and Activate Serotonergic Neurons through the Medial Prefrontal Cortex. *J*
Neurosci 2007; **27**: 11700–11711.

41   Flores Á, Maldonado R, Berrendero F. Cannabinoid-hypocretin cross-talk in the central
nervous system: what we know so far. *Front Neurosci* 2013; **7**.
doi:10.3389/fnins.2013.00256.

42   Katona I, Freund TF. Endocannabinoid signaling as a synaptic circuit breaker in
neurological disease. *Nat Med* 2008; **14**: 923–930.

43 Scarante FF, Vila-Verde C, Detoni VL, Ferreira-Junior NC, Guimarães FS, Campos AC.
Cannabinoid Modulation of the Stressed Hippocampus. *Front Mol Neurosci* 2017; **10**: 411.

44 Zhang Y, Yuan S, Pu J, Yang L, Zhou X, Liu L *et al.* Integrated Metabolomics and
Proteomics Analysis of Hippocampus in a Rat Model of Depression. *Neuroscience* 2018;
371: 207–220.

45 Fischer C, Thomas D, Gurke R, Tegeder I. Brain region specific regulation of anandamide
(down) and sphingosine-1-phosphate (up) in association with anxiety (AEA) and resilience
(S1P) in a mouse model of chronic unpredictable mild stress. *Pflugers Arch - Eur J Physiol*
2024; **476**: 1863–1880.

46 Hill MN, Karatsoreos IN, Hillard CJ, McEwen BS. Rapid elevations in limbic
endocannabinoid content by glucocorticoid hormones in vivo. *Psychoneuroendocrinology*
2010; **35**: 1333–1338.

47 Hill MN, McLaughlin RJ, Bingham B, Shrestha L, Lee TTY, Gray JM *et al.* Endogenous
cannabinoid signaling is essential for stress adaptation. *Proc Natl Acad Sci USA* 2010; **107**:
9406–9411.

48 Cui L, Li S, Wang S, Wu X, Liu Y, Yu W *et al.* Major depressive disorder: hypothesis,
mechanism, prevention and treatment. *Sig Transduct Target Ther* 2024; **9**: 30.

49 Fujino K, Yoshitake T, Inoue O, Ibii N, Kehr J, Ishida J *et al.* Increased serotonin release in
mice frontal cortex and hippocampus induced by acute physiological stressors.
Neuroscience Letters 2002; **320**: 91–95.

50 Martínez-Torres S, Bergadà-Martínez A, Ortega JE, Galera-López L, Hervera A, De Los
Reyes-Ramírez L *et al.* Peripheral CB1 receptor blockade acts as a memory enhancer
through a noradrenergic mechanism. *Neuropsychopharmacol* 2023; **48**: 341–350.

51 Henningsen K, Palmfeldt J, Christiansen S, Baiges I, Bak S, Jensen ON *et al.* Candidate
Hippocampal Biomarkers of Susceptibility and Resilience to Stress in a Rat Model of
Depression. *Molecular & Cellular Proteomics* 2012; **11**: M111.016428-1-M111.016428-
12.

52 Best AR, Regehr WG. Serotonin Evokes Endocannabinoid Release and Retrogradely
Suppresses Excitatory Synapses. *Journal of Neuroscience* 2008; **28**: 6508–6515.

53 Castillo PE, Younts TJ, Chávez AE, Hashimotodani Y. Endocannabinoid Signaling and
Synaptic Function. *Neuron* 2012; **76**: 70–81.

54 Hill MN, Carrier EJ, McLaughlin RJ, Morrish AC, Meier SE, Hillard CJ *et al.* Regional
alterations in the endocannabinoid system in an animal model of depression: effects of
concurrent antidepressant treatment. *Journal of Neurochemistry* 2008; **106**: 2322–2336.

55 Mukherjee J, Yang Z-Y. Evaluation of monoamine oxidase B inhibition by fluoxetine
(Prozac): An in vitro and in vivo study. *European Journal of Pharmacology* 1997; **337**:
111–114.

56 Mukherjee J, Yang Z-Y. Monoamine oxidase a inhibition by fluoxetine: An in vitro and in
vivo study. *Synapse* 1999; **31**: 285–289.

57 Gold BI, Bowers Jr MB, Roth RH, Sweeney DW. GABA levels in CSF of patients with
psychiatric disorders. *American Journal of Psychiatry* 1980; **137**: 362–364.

58 Garakani A, Martinez JM, Yehuda R, Gorman JM. Cerebrospinal fluid levels of glutamate
and corticotropin releasing hormone in major depression before and after treatment. *Journal*
of Affective Disorders 2013; **146**: 262–265.

59 Hasler G, Meyers N, Shen J, Drevets WC. Reduced Prefrontal Glutamate/Glutamine and -
Aminobutyric Acid Levels in Major Depression Determined Using Proton Magnetic
Resonance Spectroscopy. *ARCH GEN PSYCHIATRY* 2007; **64**.

60 Liu K, Zhao L, Xu W, Lin Q, Zhou Y, Huang X *et al.* Metabolic Changes Associated with a
Rat Model of Diabetic Depression Detected by Ex Vivo[1] H Nuclear Magnetic Resonance
Spectroscopy in the Prefrontal Cortex, Hippocampus, and Hypothalamus. *Neural Plasticity*
2018; **2018**: 1–12.

61 Popoli M, Yan Z, McEwen BS, Sanacora G. The stressed synapse: the impact of stress and
glucocorticoids on glutamate transmission. *Nat Rev Neurosci* 2012; **13**: 22–37.

62 Xu X, Wu K, Ma X, Wang W, Wang H, Huang M *et al.* mGluR5-Mediated eCB Signaling
in the Nucleus Accumbens Controls Vulnerability to Depressive-Like Behaviors and Pain
After Chronic Social Defeat Stress. *Mol Neurobiol* 2021; **58**: 4944–4958.

63 Qin Z, Zhou X, Pandey NR, Vecchiarelli HA, Stewart CA, Zhang X *et al.* Chronic Stress
Induces Anxiety via an Amygdalar Intracellular Cascade that Impairs Endocannabinoid
Signaling. *Neuron* 2015; **85**: 1319–1331.

64 Shepard R, Coutellier L. Changes in the Prefrontal Glutamatergic and Parvalbumin Systems
of Mice Exposed to Unpredictable Chronic Stress. *Mol Neurobiol* 2018; **55**: 2591–2602.

65 Lazarevic V, Mantas I, Flais I, Svenningsson P. Fluoxetine Suppresses Glutamate- and
GABA-Mediated Neurotransmission by Altering SNARE Complex. *IJMS* 2019; **20**: 4247.

66 Pehrson AL, Sanchez C. Serotonergic modulation of glutamate neurotransmission as a
strategy for treating depression and cognitive dysfunction. *CNS Spectr* 2014; **19**: 121–133.

Acknowledgments

We thank Raquel Martín for her extensive technical assistance throughout the project, and Francisco Porrón for his expert support in the animal experiments. We also thank Rafael de la Torre, Oscar Pozo and Élida Alechaga for their expert assistance with the metabolomic studies. We are also grateful to Araceli Bergadà-Martínez, Tamara Pezzotta, Maria Ruiz Suárez, Alba Piñeiro Justo, and the Neuropharmacology laboratory for their valuable discussions and technical help.

Author Contributions

AR-L contributed to the experimental design, performed the behavioral experiments, conducted sample collection and processing, analyzed the data, and drafted the manuscript. ECI contributed to the study design and supervised data analysis. VG participated in the project development and experimental design. MM, PR, ET, RM, and BA-P contributed to project

funding and manuscript revision. PR and BA-P conceived the study and contributed to
experimental design, data analysis, and manuscript writing and revision.

Funding

This research was framed in the ANTaRES research project, funded by the Instituto de Salud
Carlos III (ISCIII), and Agence Nationale de la Recherche (ANR) on behalf of ERA-NET
Neuron (AC19/00098) to MM, PR, ET and BA-P. This research was supported by the Spanish
Ministry of Science and Innovation through grant PID2020-120029GB-I00
(MICIN/AEI/10.13039/501100011033), and by the Spanish Ministry of Health, Social Services
and Equity (RD16/0017/0020) to RM. Additional support was provided by the Research and
University Department of the Catalan Government (Generalitat de Catalunya) through grant
2017-SGR-669, as well as the ICREA-Acadèmia 2021 and 2025 programs (Institució Catalana
de Recerca i Estudis Avançats, Generalitat de Catalunya, Spain) to RM. AR-L was supported by
a predoctoral fellowship from the Generalitat de Catalunya (2021 FI_B 00116).

Competing Interests

The authors declare that they have no competing interests.

FIGURES

584 **Figure 1. Effects of short-term fluoxetine treatment on depressive-like behavior. (a)** Schematics of

the UCMS protocol. **(b)** Body weight measures, **(c)** fur coat score, and **(d)** grooming time in the splash

test, during UCMS and after 1 week of treatment with fluoxetine. Depressive-like behavior **(e)** before

starting fluoxetine treatment in the FST and **(f)** after 7 days of treatment in the TST. **(g)** Computation of a

global index from the z-score value of total time grooming, body weight, and fur coat score at the end of

the experimental protocol. Data are expressed as mean ± s.e.m. The vertical dashed line in panel (c)

indicates the start of fluoxetine or vehicle treatment. Individual data points are depicted as circles, and n =

indicates the number of male mice per group. The dashed vertical line in (c) indicates the starting point of

fluoxetine treatment. * $p < 0.05$, ** $p < 0.01$, **** $p < 0.001$; #### $p < 0.001$ (time); @@ $p < 0.01$ (time

x group) by one-way ANOVA or mixed-effect analysis followed by Bonferroni *post-hoc*.

Figure 2. Effects of UCMS and short-term treatment with fluoxetine on hippocampal levels of endocannabinoids. Endocannabinoids **(a)** 2-AG and **(b)** AEA quantified by LC-MS/MS. UCMS-subjected mice treated with fluoxetine are separated into responders (Respon.) and non-responders (Nonrespon.). Data are expressed as mean ± s.e.m. Individual data points are depicted as circles and n = indicates the number of mice per group. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$; by one-way ANOVA followed by uncorrected LSD *post-hoc*.

603
604

Figure 3. Effects of UCMS and short-term fluoxetine treatment on hippocampal levels of
neurotransmitters and related metabolites. (**a**) Serotonin (5-HT) and (**b**) serotonin turnover (5-
HIAA/5-HT), (**c**) dopamine (DA) and (**d**) dopamine turnover (DOPAC/DA), (**e**) noradrenaline, (**f**)
glutamate (Glu), (**g**) GABA and (**h**) glycine quantification by LC-MS/MS. Stressed mice treated with
fluoxetine were separated into responders (Respon.) and non-responders (Nonrespon.). Data are
expressed as mean ± s.e.m. Individual data points are depicted as circles and n = indicates the number of
male mice per group. * p < 0.05, ** p < 0.01, *** p < 0.001, **** p < 0.0001; by one-way ANOVA
followed by Bonferroni *post-hoc*.

616
Figure 4. Effects of UCMS and short-term fluoxetine treatment on prefrontal cortex levels of
neurotransmitters and related metabolites. (a) Serotonin (5-HT) and **(b)** serotonin turnover (5-
HIAA/5-HT), **(c)** dopamine (DA) and **(d)** dopamine turnover (DOPAC/DA), **(e)** noradrenaline, **(f)**
glutamate (Glu), **(g)** GABA and **(h)** glycine quantification by LC-MS/MS. Stressed mice treated with
fluoxetine are separated into responders (Respon.) and non-responders (Nonrespon.). Data are expressed
as mean ± s.e.m. Individual data points are depicted as circles, and n = indicates the number of mice per

group. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$; by one-way ANOVA followed by

Uncorrected LSD (panels f and g) and Bonferroni (panels a-e, and h) *post-hoc*.

Figure 5. Correlation matrices of neurotransmitters in hippocampus and prefrontal cortex. Each panel illustrates the significant (p<0.05) correlations found in the hippocampus and the prefrontal cortex between all the neurotransmitter concentrations studied. **(a)** control mice, **(b)** stressed mice treated with vehicle, **(c)** responders to fluoxetine, and **(d)** non-responders to fluoxetine. The numbers inside the squares represent the Pearson correlation coefficient (r). The colors inside the squares indicate the strength and direction of the association, with red and blue for positive and negative correlations, respectively.