Population structure of *Phytophthora infestans* collected from potatoes in Ecuador, Colombia, Peru, Bolivia and Uruguay

Myriam Izarra[1], Mario Coca-Morante[2], Willmer, Pérez[1], Laura Sánchez[3], Soledad Gamboa[1], Diana Valle[4], Viviana Cuarán[5], Beatriz Guerra-Sierra[3*], Jan Kreuze[1*]

[1]Centro Internacional de la Papa (CIP), Avenida La Molina 1558, 15024, Peru.

[2]Universidad Mayor de San Simón (UMSS), Facultad de Ciencias Agrícolas y Pecuarias. Cochabamba, Bolivia.

[3]Universidad de Santander (UDES), Facultad de Ciencias Naturales y Agropecuarias, Grupo Biotecnología Agroambiente y Salud-Microbiota, Bucaramanga, Colombia.

[4]Instituto Nacional de Investigación Agropecuaria (INIA), Sistema Vegetal intensivo, Las Brujas, Canelones Uruguay.

[5]Corporación Colombiana de investigación Agropecuaria (AGROSAVIA), Centro de investigación la Suiza, Km 32, Rionegro, Santander, Colombia

* Correspondence author: Beatriz Guerra, bguerra@udes.edu.co
* Correspondence author: Jan Kreuze, j.kreuze@cgiar.org

Abstract

Late blight, a destructive disease affecting potatoes, is caused by the oomycete *Phytophthora infestans* and remains a major threat to potato production worldwide. Understanding the population structure of this pathogen is essential for effective disease management. We examined the genetic structure of South American *P. infestans* populations using 182 isolates: 97 from Bolivia and southern Peru, 14 from Colombia, 57 from Ecuador (1993-2022), and 14 from Uruguay. The isolates were characterized by clonal lineage, mitochondrial haplotype, and mating type using multilocus genotypes based on microsatellite (SSR) markers. In Bolivia, only the lineage/haplotype 2A1/Ia was identified, whereas Peru exhibited both 2A1/Ia and EC1/IIa, all of which were mating type A1. Puno was the sole department where both lineages were present. In historical Ecuadorian populations, we found US1/Ib and EC1/IIa (*P. infestans*), as well as EC2/Ic and EC3/Ia (*Phytophthora andina*), while recent populations showed only US1. In Colombia, EC1/IIa and a distinct clonal lineage (CO4/Ia) were identified within the present dataset. In Uruguay, 2A1/Ia was predominant.

These results provide updated insights into the genetic diversity and geographic distribution of *P. infestans* across South America and highlight the importance of continued regional surveillance for improved late blight management.

Keywords: Phytophthora infestans, clonal lineage, South America, late blight

Introduction

Phytophthora infestans, the causal agent of potato late blight, has a complex population structure and evolutionary history in South America. The global migration history of *P. infestans* has been the subject of extensive research and debate. There are various hypotheses regarding the origin of *P. infestans*, including Mexican, Andean, and hybrid origin hypotheses. Goss et al. (2014) suggested that Mexico was the center of origin from which the pathogen migrated to South America, then to the United States, and finally to Europe (Andrivon, 1996). A study by Gómez-Alpizar et al. (2007) found that Andean populations harbored greater nucleotide and haplotype diversity than Mexican populations, and that the oldest mutations in *P. infestans* originated in South America, supporting the Andean origin theory. More recently, Coomber et al. (2025) reaffirmed that the Andes is the center of origin, and Patarroyo et al. (2024) suggested a Peruvian origin with a probability of 0.631. The movement of genotypes contributes to shaping population structure and disease dynamics (Guha Roy et al., 2021).

49
The study of the population structure of *P. infestans* is complex and dynamic and has been widely studied across
different regions. It is influenced by factors, such as natural migration or migration associated with the movement of
infected potato seed tubers, sexual reproduction, and geographical isolation. Additionally, *P. infestans* can rapidly
evolve and adapt, posing a challenge for disease management (Cooke & Lees, 2004; Dey *et al.*, 2018;
Ludwiczewska *et al.*, 2025). Microsatellite markers, also known as simple sequence repeats (SSRs), are highly
polymorphic co-dominant nuclear markers, providing valuable insights into the genetic variation and population
dynamics of this devastating pathogen (Lees *et al.*, 2006).
57
P. infestans is potentially able to reproduce sexually when compatible A1 and A2 mating types coexist in the same
population. However, sexual reproduction is not always observed even when both mating types are present, as
compatibility between strains may vary (Runno-Paurson *et al.*, 2012). Until the 1980s, only mating type A1 was
known to occur in Europe, which restricted the pathogen to asexual reproduction. The appearance of A2, first
detected in Switzerland and likely introduced through potato imports from Mexico (Rhouma *et al.*, 2024), changed
the disease dynamics (Drenth et al., 1993; Shattock, 2002). The presence of both A1 and A2 mating types allows
sexual reproduction, potentially leading to increased genetic diversity and the formation of oospores that, as
alternative inoculum sources, can persist in the soil and act as long-term infection sources (Drenth *et al.*, 1993;
Shattock, 2002; Yuen & Andersson, 2013). This supplementary source of inoculum, combined with increased
genetic diversity, enhances the pathogen's capacity to adapt and persist in the absence of a host, potentially
complicating traditional disease management strategies (Drenth et al., 1993, 1994).
69
Clonal lineages of *Phytophthora infestans* are designated using standard nomenclature (e.g., US1, EC1, 2A1), while
mitochondrial haplotypes are indicated by Roman numerals (e.g., Ia, IIa). Combined designations (e.g., US1/Ib,
EC1/IIa, 2A1/Ia) refer to the association between clonal lineage and mitochondrial haplotype. In Peru, the
population structure of *P. infestans* exhibits clonal lineages with varying dominance across regions and over time. In
the southern region, isolates collected between 1994 and 1999 were characterized by clonal lineages and
mitochondrial haplotypes. In Cusco, the identified lineages were US1/Ib, EC1/IIa, PE3/Ia, PE5/IIa, and PE6/IIa. In
Puno, US1/Ib, PE3/Ia, and EC1/IIa were found (Perez *et al.*, 2001).
A subsequent study analyzing isolates collected between 1997 and 2013 reported the presence of PE3/Ia and
EC1/IIa in Cusco, US1/Ib and PE3/Ia in Puno, and EC1/IIa in Apurímac (Lindqvist⬜Kreuze *et al.*, 2020). More
recently, isolates from 2017 were analyzed by Lindqvist⬜Kreuze et al. (2020), revealing EC1/IIa in Cusco and both
EC1/IIa and PE7.1/Ia (a variant of PE7) in Puno, all of which belonged to the A1 mating type.
P. infestans populations in Bolivia were studied using herbarium specimens to determine clonal lineages and
mitochondrial haplotypes, which were found to be US1/Ib, and from modern isolates, which identified BR1/IIa
(Saville & Ristaino, 2021). Gabriel et al. (2018) identified the A1 mating type, in contrast to Goodwin, S.B. et al.
(1994), who identified mating type A2 in Bolivia.
The population structure of *P. infestans* in Colombia has been reported to be dominated by the clonal lineage
EC1/IIa, whereas CO1/IIa and CO2/IIa have also been identified on potato (Vargas *et al.*, 2009). In contrast, US1/Ib
and CO3/Ib were identified on tomato hosts (Céspedes et al., 2013; Gilchrist Ramelli et al., 2009; Vargas et al.,
2009; Olave-Achury et al., 2022).
In Uruguay, the presence of BR1/IIa was previously reported (Deahl *et al.*, 2003), but more recent studies have
revealed the presence of 2A1/Ia (F. Lucca et al., 2023).
In Ecuador, the presence of distinct clonal lineages and mitochondrial haplotypes has been studied over the years,
with EC1/IIa being dominant on potato hosts and US1/Ib on tomato hosts (Forbes *et al.*, 1997; Oyarzun Pj *et al.*,
1998). In addition to *P. infestans*, closely related species such as *Phytophthora andina* have also been reported in the
Andean region. *P. andina* is considered a distinct species within the *P. infestans* complex, likely of hybrid origin,
and is associated with infections on wild and cultivated *Solanum* species (Gómez-Alpizar *et al.*, 2008; Oliva *et al.*,
2010). Additionally, lineages such as EC2 and EC3 have previously been identified as *P. andina* on different
Solanum species (Gómez-Alpizar *et al.*, 2008).
98
The importance of studying South American *P. infestans* populations lies in their considerable diversity (Coomber *et*
al., 2025), which has implications for disease management and breeding resistance. Clonal lineage 2A1, first
reported in Europe in the early 1980s (Cooke *et al.*, 2012), was dominant on potatoes before 2006 (Mariette *et al.*,
2016). The 2A1 lineage has also shown significant expansion beyond Europe including the eastern-Africa region
(EAR), comprising Kenya, Uganda, Tanzania, Burundi, and Rwanda (Njoroge *et al.*, 2019). Its presence in South

America has been associated with the introduction of infected potato tubers. This lineage is now regarded as the
dominant one in Argentina, Chile, Brazil, and Uruguay (Acuña et al., 2019; F. Lucca et al., 2023)
106
We propose that the genetic variation and population dynamics of *Phytophthora infestans* in South America are
influenced by both geographic and agroecological factors, leading to distinct regional patterns. In particular, we
anticipate that populations in ecologically linked or neighboring regions (such as the central Andes—Colombia,
Ecuador, Peru and Bolivia) will exhibit higher genetic similarity, while those geographically distant from this region
(such as southern Uruguay) will show more pronounced genetic differences possibly influenced by the importation
of seed potatoes from Northern Hemisphere countries.
113

Materials and methods

Sampling
A total of 182 *Phytophthora infestans* isolates collected between 1993 and 2025 from five South American countries
(Ecuador, Colombia, Peru, Bolivia and Uruguay) were included in this study. Of these, 127 isolates were from field
collections conducted between 2017 and 2025 in Bolivia (n=67), Peru (n=30), Colombia (n=14), Uruguay (n=14),
and Ecuador (n=2, collected in 2022). Additionally, 55 historical Ecuadorian isolates collected between 1993 and
2008 were obtained from the CIP *Phytophthora* collection and had been preserved in liquid nitrogen.
During 2017-2019, leaf samples with late blight-like symptoms were collected in Bolivia and southern Peru.
Bolivian samples originated from the department of Cochabamba, whereas Peruvian samples were collected in the
departments of Cusco, Apurimac, Ayacucho, and Puno. In Colombia, samples were collected in 2022 from
Santander and the North of Santander. Two Ecuadorian samples were collected in 2022 from Pichincha. In Uruguay,
samples were collected between 2023-2025 from Canelones and San Jose. Each leaf sample yielded a single isolate
for further molecular characterization.
Field samples (Bolivia, Peru, Colombia, Uruguay, and Ecuador-2022) were collected on FTA cards (Whatman FTA
Classic Card, catalog number WB120055; GE Healthcare UK Ltd.) according to the manufacturer's guidelines
(Cooke, 2020). Each sample was placed on FTA paper with the sporulating side down and individually crushed in
the field (Gamboa *et al.*, 2019). The FTA cards were then stored in zip-lock bags and shipped to the pathology
laboratory at the International Potato Center (CIP) in Lima, Peru, for processing.
Complete metadata for all isolates, including collection year and date, geographic origin (country, department,
province, and district), altitude, host, and genotypic characterization, are provided in Supplementary Table S1.
Genotypic Analysis
135
All 182 *Phytophthora infestans* isolates were subjected to molecular characterization. Mitochondrial haplotypes
(mtDNA) and multilocus genotypes (MLGs) were determined for all isolates using PCR-based assays and a 12-plex
SSR panel as described below.
Field-collected samples were obtained on FTA cards as described above. The 55 historical Ecuadorian isolates
(1993-2008), preserved in liquid nitrogen at the CIP collection, were reactivated in culture and transferred onto FTA
cards prior to molecular analysis.
DNA extracted from FTA cards (Qiagen®) was performed following the manufacturer's instructions with minor
modifications: two 3 mm discs were excised (instead of one), rinsed twice in modified Tris (hydroxymethyl)
aminomethane-ethylenediaminetetraacetic acid TE^{-1} buffer (10 mM Tris and 0.1 mM EDTA), and incubated in 50 µl
TE buffer (10mM Tris and 1mM EDTA), at 95 °C for 5 min, and stored at -20 °C until use.
146
147 Mating Type
148
149 Mating type information was available for 178 of 182 isolates included in this study. For field-collected isolates,
150 mating type was determined using W16 CAPS assay (Judelson *et al.*, 1995). For the historical Ecuadorian isolates
151 (1993-2008), mating type information was obtained from previous in vitro pairing assays recorded in the CIP
152 database.
153
W16 CAPS assay
The mating type (A1 and A2) was determined using PCR primers W16-1 (5'-AAC ACG CAC AGG CAT ATA
AAT GTA-3') and W16-2 (5'-GCG TAA TGT AGC GTA ACA GCT CTC-3'), as described by Judelson et al.
(1995) for all isolates except the Ecuadorian isolates. PCR reactions (20 µL) contained 4 µL of 5x buffer (Gotaq,

Promega), 6 μL of MgCl$_2$ (25 mM), 0.2 μL of dNTPs (100 μM), 0.4 μM of each primer, 1 U of Taq DNA
polymerase (Gotaq®, Promega), and 10 ng of template DNA. The thermal cycling protocol included an initial
denaturation at 94 °C for 5 min, followed by 30 cycles of denaturation at 94 °C for 1 min, primer annealing at 53 °C
for 1 min, and elongation at 72 °C for 1 min, with a final elongation step at 72 °C for 10 minutes. Afterwards, the
PCR products were digested for 3h at 37 °C with Hae III (New England Biolabs, #R0108) in a 20 μL reaction
containing 10 μL of PCR product, 2 μL of CutSmart Buffer 10X, 0.3 μL enzyme, and nuclease-free water (Brylińska
et al., 2018). Digested fragments were separated on a 2% agarose gel in TBE buffer at 40 V for 4 h and visualized
under UV light using GelRed (Biotium®) as nucleic acid stain.
166
167 *Pairing Assay*
168 For the historical Ecuadorian isolates, mating type had been previously determined by pairing with A1 (*P. infestans*)
169 and A2 (*P. andina*) tester strains using dual-cultures on 10% clarified V8 agar. Plates were incubated at 15-18 °C in
170 the dark and examined microscopically for oospore formation at the hyphal interfaces within the contact zone.
171 Isolates forming oospores with A1 and A2 testers were classified as A2 and A1, respectively.
172
Mitochondrial haplotype
Mitochondrial haplotypes were determined for all 182 isolates using a PCR-restriction fragment length
polymorphism (PCR-RFLP) assay following Griffith & Shaw (1998). A specific mitochondrial region was amplified
using primer P2, and PCR products were digested with MspI.
Restriction fragments were separated on a 2% agarose gel in TBE buffer at 40 V for 4h and visualized under UV
light using GelRed (Biotium®) as nucleic acid stain. Haplotypes were assigned by comparing with reference isolates
representing the established Ia, Ib, IIa, and Ic mitochondrial haplotypes (Griffith & Shaw, 1998).
180
Molecular Markers
All 182 isolates were genotyped using a 12-plex microsatellite (SSR) panel following a modified protocol of Li et
al. (2013) and Saville et al. (2016). PCR amplification was performed using the Type-It Microsatellite PCR kit
(QIAGEN) according to Saville et al. (2016).
Amplified products were submitted to Arizona State University for each sample, 2 μL of a 1:50 diluted PCR product
was combined with formamide containing LIZ500 size standard ladder and loaded onto 3730 instrument. Alleles
sizes were determined using GENEMARKER v1.9 (SoftGenetics) according to the allele binning parameters
described by Li et al. (2013). Clonal lineage assignment was performed by comparison with reference SSR profiles
representing previously genotyped clonal lineages, including 2A1, EC1, BR1, PE3, PE7, PE7.1, and US8 (Vargas *et*
al., 2009; Martin *et al.*, 2019; Lindqvist□Kreuze *et al.*, 2020; Saville & Ristaino, 2021). Additional Andean
reference lineages included EC2, EC3, and PE8, previously reported within the *P.andina* (Adler *et al.*, 2004; Oliva
et al., 2010; Forbes *et al.*, 2016) which were associated with the Andean species *P.andina*.
193
Data Analysis
All statistical analyses were conducted using R version 4.3.1 (R Core Team, 2019). The following packages were
used: polysat (v.1.7.7), poppr (v.2.9.7), RColorBrewer (v.1.1-3), magrittr (v.2.0.3), ape (v.5.8-1), and ggtree (v.3.12-
0). Locus and population-level statistics, including the number of multilocus genotypes (MLGs), were calculated
using the package poppr (Kamvar *et al.*, 2014).
Clonality was assessed using the index of association (Ia) and the standardized index of association (r□d), with
statistical significance evaluated through 999 permutations using the ia() function in poppr. For diversity analyses
sensitive to repeated multilocus genotypes (MLGs), clone correction was applied by retaining a single representative
of each MLG per population.
Population structure was explored using principal coordinates analysis (PCoA), hierarchical clustering, and
minimum spanning networks (MSN) all based on Bruvo's genetic distance (Bruvo *et al*., 2004). For PCoA, a Bruvo
distance matrix was calculated for triploid SSR genotypes using the meandistance.matrix function in the R package
polysat (Clark & Jasieniuk, 2011), and ordination was performed using classical multidimensional scaling
(cmdscale) in R. Hierarchical relationships were examined using a unweighted pair group method with arithmetic
mean (UPGMA) dendogram generated from a Bruvo distance matrix calculated with bruvo.dist (specifying locus-
specific repeat lengths) and clustered using upgma in poppr. Genotypic relationships were further visualized using a
minimum spanning network (MSN) constructed with bruvo.msn in □poppr (Kamvar *et al.*, 2014). Ordination and
network analyses were performed using the full dataset to reflect genotype frequencies. Reference genotypes
representing known clonal lineages were included across analyses to support lineage assignment.
213

Diversity statistics were computed for each population, including the number of individuals (N), number of
multilocus genotypes (MLG), expected number of multilocus genotypes (eMLG), standard error (SE), Shannon-
Wiener Index (H), Stoddart and Taylor's index (G), Simpson's index (λ), evenness (E.5), and expected
heterozygosity (Hexp), index of association (Ia), and standardized index of association (r⬚D). Additional locus-
level statistics included allelic richness, Simpson's diversity index (1–D), expected heterozygosity (Hexp), and
evenness (Shannon, 1948, 2001; Simpson, 1949; Hedrick, 1999).
220
221 Analysis of molecular variance (AMOVA) was performed using Bruvo's genetic distance to partition genetic
222 variation among clonal lineages and among geographic locations nested within clonal lineages (~ lineage/location).
223 AMOVA was conducted on a clone-corrected dataset, retaining one representative per multilocus genotype (MLG)
224 per population prior to distance matrix computation. The analysis was implemented using the poppr.amova function
225 with the ade4 method. Statistical significance was assessed using 999 Monte Carlo permutations.
226

Results

228 *Samples collected*
229
A total of 182 *Phytophthora infestans* isolates were analyzed in this study. In Bolivia, 67 samples were collected
from the department of Cochabamba, across ten provinces. In Peru, samples were obtained from four departments:
Cusco (n=1), Apurímac (n=4), Ayacucho (n=6), and Puno (n=19). In Colombia, 14 samples were collected from
Norte de Santander (n=6) and Santander (n=8). In Ecuador, 57 isolates were obtained from ten departments: Carchi
(n=11), Bolívar (n=2), Chimborazo (n=2), Cotopaxi (n=5), Imbabura (n=3), Morona Santiago (n=1), Napo (n=9),
Pastaza (n=2), Tungurahua (n=8), and Pichincha (n=14). Two isolates from Pichincha were collected in 2022, while
the remaining Ecuadorian isolates correspond to historical samples preserved in the CIP collection. In Uruguay, 14
samples were collected from the departments of Canelones (n=10) and San José (n=4). Detailed isolate metadata are
provided in Tables 1 and S1.
239
Mating type test
The primer pair W16–1 and W16–2 amplified a 557-bp fragment in all isolates analyzed with this assay. Following
digestion with HaeIII, A1 isolates were identified by the presence of the undigested 557-bp fragment, together with
the expected digested fragment (457-bp), consistent with the expected banding pattern for A1 mating type.
All isolates from Bolivia, Peru, Colombia, and Uruguay were classified as A1 based on the W16 assay (Figure S1,
Table S1). For Ecuadorian isolates, mating type was determined using pairing assays. Most isolates were A1, while
a single isolate (Ec-3365), obtained from the wild species *Solanum brevifolium*, was identified as A2 and
corresponded to *Phytophthora andina* (Table S1).
248
249 *Genotypic characterization*
250
Mitochondrial haplotypes
Analysis of the mitochondrial haplotypes showed that isolates from Bolivia (n=67), Uruguay (n=14), Peru (n=8),
Colombia (n=10), and Ecuador (n=2) belonged to haplotype Ia. In contrast, IIa was detected in 23 isolates from
Peru, 3 from Colombia, and 38 from Ecuador. Haplotype Ib was detected in 16 Ecuadorian isolates, including two
collected in 2022. A single isolate (Ec-3365) obtained from the wild species *Solanum brevifolium* corresponded to
haplotype Ic (Figure S2, Table S1).
257
258 Population structure
259 Principal coordinates analysis (PCoA) based on Bruvo genetic distance revealed clustering patterns among isolates
260 from 19 departments across five countries, together with nine reference genotypes. The analysis showed that the
261 reference genotype PE7.1 clustered with the globally distributed clonal lineage 2A1 (Figure 1).
262
263 Hierarchical clustering using a UPGMA dendrogram based on Bruvo's distance identified six major clonal lineages
264 (2A1, US1, EC1, EC2, EC3, and a distinct clonal lineage, here designated as CO4, identified in Colombia based on
265 its clustering pattern relative to reference genotypes included in the dataset) (Figure 2).
266

Figure 1. Principal coordinate analysis (PCoA) based on Bruvo's genetic distance calculated from SSR marker data for *Phytophthora infestans* and *P.andina* isolates. Colors represent geographic origin: Bolivia (Cochabamba, red), Peru (Cusco, blue; Apurimac, green; Ayacucho, purple; Puno, orange), Colombia (Norte de Santander, bright yellow; Santander, brown), Ecuador (Carchi, pink; Pichincha, gray; Imbabura, teal; Tungurahua, coral; Chimborazo, light fuchsia; Cotopaxi, lime green; Napo, soft yellow; MoronaSantiago ,sandy beige; Pastaza, light gray; Bolivar, forest green); and Uruguay (Canelones, burnt orange; San José, indigo). Reference genotypes used for clonal lineage assignment are indicated: US1(fuchsia), EC1(olive green), 2A1 (dark Gray), PE3 (mint green), PE7.1 (sky blue), PE7 (lavender) and BR1 (pale green), US8 (turquoise) for *P. infestans*; and EC2 (dusty blue), EC3(mustard), PE8 (bronze) for *P. Andina*.

Clonal lineage distribution

In Bolivia (n=67) and Uruguay (n=14), all isolates belonged to the 2A1 clonal lineage. In Peru, isolates from Puno were distributed between EC1 (n=11) and 2A1 (n=8). In contrast, isolates from Cusco (n=1), Apurimac (n=4), and Ayacucho (n=6) were all assigned to EC1, indicating that this lineage predominated in southern Peru (73.3%), followed by 2A1 (26.6%).

In Ecuador, US1 was not detected in cultivated potato but occurred in several host species, including *S. lycopersicum* (18.8%), *S. muricatum* (12.5%), *S. andreanum* (6.2%), *S. quitoense* (56.3%) and *S. caripense* (6.2%). In contrast, EC1 was detected in cultivated potatoes (39.5%) and in several wild *Solanum* species, including *S. colombianum* (18.4%), *S. ochrantum* (23.7%), *S. paucijugum* (10.5%), and *S. andreanum* (7.9%).

Within the Ecuadorian population, EC1 represented 66.7% of isolates, followed by US1 (28.1%) and EC3 (3.5%, all of which were associated with *S. betaceum*). EC2 represented 1.8% of isolates and was detected in *S. brevifolium*. The largest number of isolates were collected in Carchi and Pichincha. In Carchi, all isolates belonged to EC1, whereas in Pichincha the population consisted of EC1 (42.9%) and US1 (57.1%). The isolates collected in 2022

292   from Pichincha belonged to the US1 lineage and represented 14.3% of the isolates from that department. In
293   Tungurahua, three clonal lineages were detected: EC1 (62.5%), EC2 (12.5%), and EC3 (25%).
294   In Colombia, two well-differentiated clonal lineages were detected. The newly identified lineage CO4 represented
295   78.6% of isolates and occurred in both studied departments and 21.4% corresponded to EC1, found only in
296   Santander (Figure 2, 3).

Hierarchical clustering of South American Phytophthora isolates

Bruvo distance

298 **Figure 2**. UPGMA dendrogram of *Phytophthora* isolates from Bolivia, Peru, Colombia, Ecuador, and Uruguay
299 based on Bruvo's distance from 12 SSR loci. Clonal lineage assignment was inferred from clustering with included
300 reference genotypes. Major lineages (US1, EC1, 2A1, EC2, EC3, and CO4) are indicated; EC2 and EC3 correspond
301 to *P. andina*, whereas the remaining lineages belong to *P. infestans*.
302
303
304

305
306
307
308 **Figure 3.** Geographical distribution of *Phytophthora* isolates (n =182) collected in Bolivia and Peru (2017–2019),
309 Ecuador   (1993–2022),   Colombia   (2022),   and   Uruguay   (2023–2025).
310 Departments highlighted in gray indicate sampled locations, and pie charts show the relative composition of clonal
311 lineages 2A1, EC1, CO4, and US1 (*P. infestans*), and EC2 and EC3 (*P. andina*). Pie chart size is proportional to the
312 total number of isolates per department, and numerical values indicate the number of isolates per clonal lineage.
313
314
315
316 Figure 4 illustrates the temporal distribution of *P. infestans* clonal lineages in Ecuador between 1993 and 2022.
317 Throughout this period, EC-1 was the dominant lineage and was detected in most sampling years. In contrast, US1
318 occurred sporadically but was still detected in recent collections, including in 2022.
319

Figure 4. Clonal lineage distribution of *Phytophthora infestans* (EC-1, US-1) *and Phytophthora andina* (EC-2, EC-3) in Ecuador by year of collection (1993–2022)

Multilocus genotypes

Analysis of the 182 isolates yielded 126 multilocus genotypes (MLGs); including reference isolates, a total of 141 MLGs were detected (Table 2). The most frequent genotypes were MLG139 (n=22), MLG80 (n=7), and MLG131 (n=6) (Figure 5).

MLG80 corresponded to six isolates (one from Ayacucho and five from Puno) and one EC1 reference isolate, while MLG67 corresponded to one isolate from Puno and the second EC1 reference isolate. MLG131 included four isolates from Puno and the reference genotype previously designated PE7.1.

Puno showed relatively high genotypic diversity (MLG = 12, Hexp = 0.53) and strong signals of clonality

D = 0.68). Similarly, in Ecuador, Tungurahua and Napo exhibited high genotypic richness (MLG = 7 and 9, respectively) together with elevated linkage disequilibrium indices (Ia = 5.22 and 3.68), indicating predominantly clonal population structure. Bolivia, which represented the largest number of isolates (n=67), showed substantial genotypic diversity with 32 MLGs, although the most frequent genotype (MLG139) corresponded to the clonal lineage 2A1 (Table 2, Figure 5).

D=0.37; p=0.001), confirming significant linkage disequilibrium.

348
Figure 5. Minimum spanning network (MSN) of multilocus genotypes (MLGs) of *Phytophthora infestans* and *Phytophthora andina* populations from Bolivia, Peru, Colombia, Ecuador, and Uruguay. Each node represents a distinct MLG, with node size proportional to its frequency. Colors indicate clonal lineage assigned based on reference genotypes (US1, EC1, 2A1, PE3, PE7.1, PE7, BR1, US8 for *P.infestans;* EC2, EC3, and PE8 for *P.andina*). Edges correspond to Bruvo's genetic distances between MLGs.

Marker performance
The multilocus accumulation curve increased steadily with the addition of markers but did not reach complete saturation at 11 loci, indicating that each SSR marker contributed to genotype discrimination and that full marker coverage was necessary to capture the genetic diversity of isolates analyzed (Figure S3).

Among the SSR markers, G11 and D13 showed the highest numbers of alleles (18 and 17, respectively). The average of alleles per locus was 7.92, indicating moderate allelic diversity. Diversity indices (Simpson's 1-D and expected heterozygosity, Hexp) ranged from 0.38 to 0.81, with mean values of 0.60. Evenness ranged from 0.47 to 0.89 (mean=0.72), indicating moderate variation in allele distribution across loci (Table 3).

AMOVA
AMOVA revealed significant genetic structuring among clonal lineages. A substantial proportion of the total molecular variance (57.77 %) was attributed to differences among clonal lineages, indicating strong genetic differentiation at this level. An additional 10.21% of the variance was explained by differences among geographic

370 locations nested within clonal lineage. The remaining 32.02% of the variation occurred within samples. All
371 hierarchical levels were statistically significant according to the Monte Carlo test (P = 0.001) (Table 4).
372
373

Discussion

The genetic composition of *Phytophthora infestans* populations is dynamic and continuously evolving, with
important implications for late blight management. Previous studies have shown that *P. infestans* populations
worldwide display substantial genetic diversity and undergo rapid evolutionary change (Cooke & Lees, 2004; Dey *et
al.*, 2018). Several studies have investigated *P. infestans* populations in the Andean region, where Peru has been
proposed as a possible center of origin of the pathogen (Gómez-Alpizar *et al.*, 2007; Patarroyo *et al.*, 2024;
Coomber *et al.*, 2025).
381
Perez et al. (2001) analyzed *P. infestans* isolates from Cusco and Puno in Peru and identified EC1, PE3 and US1 as
predominant clonal lineages infecting cultivated and wild potatoes (EC1 and US1) and cultivated potatoes (PE3). In
the Peruvian highlands, host specificity does not appear to strongly influence the composition of pathogen
populations infecting tuber-bearing Solanum species (Garry *et al.*, 2005). More recent studies in these departments
identified only EC1 and a new genotype named PE7.1 (Lindqvist☐Kreuze *et al.*, 2020). EC1 has remained the
predominant genotype in Peru since its first report by Perez et al. (2001), whereas US1 and PE3 have not been
detected in more recent population studies. PE3 was previously associated with native cultivated landraces and wild
tuber-bearing species (Perez *et al.*, 2001; Garry *et al.*, 2005). Our analysis indicates that the genotype previously
designated as PE7.1 does not represent an independent lineage but instead corresponds to a variant within the
broader 2A1 clonal lineage, which has been reported in several South American countries (Acuña et al., 2019; F.
Lucca et al., 2023). US1 was identified in Ecuador in both historical and recent collections, suggests that this lineage
may persist at low frequencies in alternative hosts. Previous studies from the Andean region have reported *P.
infestans* clonal lineages infecting non-potato *Solanum* species (Garry *et al.*, 2005; Lindqvist☐Kreuze *et al.*, 2020),
which may contribute to the persistence of pathogen populations outside cultivated potato fields. More broadly, *P.
infestans* can infect a range of *Solanum* species, including wild hosts (Grünwald & Flier, 2005), and its long-
distance dispersal is strongly associated with human-mediated movement of infected plant material, particularly
seed tubers (Fry *et al.*, 2015), allowing lineages such as US-1 to persist or re-emerge despite being largely displaced
from potato populations. The long-term survival of clonal *P. infestans* lineages likely hinges on a fine balance
between their aggressiveness, adaptability to environmental conditions, and capacity to counteract disease
management strategies (Mariette *et al.*, 2016; Fry, 2020). Our research also corroborated previous studies on the
population structure of certain strains from Ecuador, which had identified the clonal lineages EC1 and
US1(*P.infestans*), EC2 and EC3 (*P.andina*) —using RFLP fingerprinting —as being associated with different host
species: EC1 (cultivated and wild potatoes), EC2 (wild potato), EC3 (cultivated tree tomato), and US1 (cultivated
tomato species, pepino melon, naranjilla, wild potato species, wild relative of naranjilla) (Gómez-Alpizar *et al.*,
2008; Oliva *et al.*, 2010). Subsequent taxonomic studies suggested that isolates previously classified within the EC3
lineage may correspond to *Phytophthora betacei rather than P.andina* (Mideros *et al.*, 2018). These findings were
updated using the 12-SSR multiplex method. The identification of EC1 and the newly discovered CO4 in Colombia
highlights a geographical influence: EC1 has been reported in Ecuador and Venezuela (Forbes *et al.*, 1998; Garry *et
al.*, 2005), while the new clonal lineage CO4 was found only in the department of Norte de Santander, where a
humid tropical ecosystem is present (Dueñas *et al.*, 2007). Norte de Santander borders Venezuela, where the Ia
mitochondrial haplotype and A1 mating type have been documented, which corresponds to the profile observed in
our strains from Norte de Santander (Briceño *et al.*, 2009; Cárdenas *et al.*, 2011). We did not identify the US8
lineage, which was previously reported in Colombia and was isolated from Cape gooseberry and also possesses the
Ia haplotype, but corresponded to the A2 mating type (Vargas *et al.*, 2009).
416
The existence of both mating types promotes gene exchange between populations, as demonstrated by the discovery
of distinct genotypes in fields where both mating types are present (Goodwin, S.B. *et al.*, 1995). In 2001, Perez et al.
(2001) found that the predominant mating type in Peru was A1, whereas in Bolivia, the A2 mating type was detected
in the past (Goodwin, S.B. *et al.*, 1994), which belongs to the clonal lineage BR1. The BR1 reference was added to
our SSR analysis, and we determined that it belonged to a different cluster from the isolates collected from Bolivia
in our study. Sexual reproduction may occur when compatible A1 and A2 mating types coexist in the same location;
however, this was not observed in the populations sampled in this study. However, the risk of sexual reproduction is

always present when both mating types co-exist, which is the case in potato production zones near the border
between Peru and Bolivia.
P. infestans may have a broader host range than previously thought, which could have implications for disease
management. In our study, the A1 mating type was detected using the W16 marker, which has been proven to be
accurate in identifying mating types in clonal lineage EC1 (Brylińska *et al.*, 2018). The pairing test on V8 agar was
chosen for the Ecuadorian population because the W16 marker has shown inconsistent results for the US1 genotype
and also for *P. andina*, as reported by Brylińska et al. (2018). In that study, all US-1 isolates of A1 mating type
amplified the fragment typical of A2 using W16, making this marker unreliable for populations where these lineages
occur.

The high proportion of genetic variation detected among clonal lineages (57.77%) indicates strong genetic
differentiation among lineages. This pattern is consistent with strong genetic differentiation among clonal lineages
identified using SSR markers and with the predominantly clonal population structure reported for *P.infestans* in
many regions (Goodwin, S.B. *et al.*, 1995). In contrast, the lower proportion of variance explained by geographic
location suggests that lineage identity is a stronger determinant of genetic structure than regional distribution.
Lineages EC1, EC2, and EC3 are predominantly associated with Andean countries, whereas lineage 2A1 has been
reported mainly in Europe, while the origin of lineage US1 remains more controversial (Forbes *et al.*, 1997; Mariette
et al., 2016; Martin Md *et al.*, 2016). These distinct evolutionary trajectories have likely generated strong genetic
differentiation among lineages, which is reflected in the high variance component observed in the AMOVA. In a
North American study of 77 isolates, Goodwin, S.B. et al. (1995) found that 63% was due to differentiation among
lineages, and older lineages exhibited more pathogenic variation than recently introduced ones. The US1 clonal
lineage has reportedly been replaced in Peru, with its presence limited to wild Solanaceae species in the 2016-2017
population. This change could be attributed to the lineage's low virulence in cultivated potatoes (Lindqvist☐Kreuze
et al., 2020). Also, EC1 was reported to be adapted to higher temperatures than US1, and to exhibit a high level of
resistance to metalaxyl (Garry *et al.*, 2005), in contrast to US1, which showed over 50% sensitivity to the fungicide
(Lindqvist☐Kreuze *et al.*, 2020). The dynamic nature of *P. infestans populations*, with the emergence and
replacement of dominant clones over time, highlights the importance of continuous monitoring and studying the
genetic variation in this pathogen for effective disease management strategies (Knaus *et al.*, 2020). For example, in
America, the US23 lineage, which is currently widespread in the United States, exhibits genetic similarities with the
BR1 lineage found in Bolivia and Brazil, whereas other lineages in the U.S. are more closely related to populations
in Mexico (Saville & Ristaino, 2019). In terms of management strategies, we need to consider the aggressiveness of
clonal lineage EC1 that currently dominates in the southern regions of Peru (Lindqvist☐Kreuze *et al.*, 2020).
Isolates within this lineage exhibit variation in key epidemiological components, particularly incubation period,
infection frequency, and lesion expansion (RAULEC, a measure of disease progression over time)(Chacón *et al.*,
2007), which may influence disease development and complicate management strategies. Furthermore, 2A1 shows
low or moderate aggressiveness relative to the other coexisting lineages (Mariette *et al.*, 2016). Additionally, the
variations in aggressiveness observed in new populations might result from better adaptation to environmental
conditions like temperature or humidity (Lindqvist☐Kreuze *et al.*, 2020).

Phytophthora infestans can be transmitted through infected seed potatoes, which represent an important source of
primary inoculum (Kool & Evenhuis, 2023). The movement of infected seed tubers has been widely recognized as a
major driver of long-distance dispersal of *P.infestans* lineages and late blight outbreaks worldwide (Guha Roy *et al.*,
2021; Ryley & Drenth, 2024; Saffer *et al.*, 2024; Rhouma *et al.*, 2024).In several Andean production systems, the
use of farm-saved seed potatoes and the exchange of planting material among neighboring countries may facilitate
the persistence and regional spread of clonal lineages such as those identified in this study.

MLGs grouped together from the potato-growing areas of Puno EC1(MLG80) and 2A1 (MLG131) were found
approximately 40 km from the Bolivian border. The current *P. infestans* population of Puno (EC1 and 2A1) is
different from that of the old population (US1, PE3 and EC1) (Perez *et al.*, 2001), suggesting that displacement of
the old dominant lineages has occurred. These shifts in lineage composition, revealed by SSR genotyping, reflect
temporal changes in population structure. The distribution of the 2A1 lineage observed in this study may reflect a
broader regional pattern of introduction and spread. This lineage, originally reported in Europe and now dominant in
several South American countries, may have been introduced into the Southern Cone (e.g., Argentina, Brazil, or
Uruguay) through the importation of infected seed tubers, and subsequently expanded northwards into Bolivia. The
presence of 2A1 in Bolivian populations and near the Peruvian border suggests a possible ongoing process of
regional dissemination, potentially leading to its future establishment in Peruvian populations. Such dynamics

highlight the importance of monitoring the spread of invasive clonal lineages across connected agricultural systems.
Since 2011, a similar shift has been noted in Argentina for the oldest lineages, AR1-AR5_A2, and in Uruguay and
Brazil for BR1_A2 (Forbes *et al.*, 1998; Deahl *et al.*, 2003), and all have been replaced by EU-2_A1 (2A1) (Lucca
et al., 2019, 2023). The most abundant MLG (MLG139) was shared between seven provinces of Cochabamba-
Bolivia, with 64.2% of MLGs being non-unique and distributed across multiple provinces, and the presence of
clonal lineage 2A1, which is present in several South American countries, suggesting that it may be older and have
had more time to spread. In the case of Peru, 56.7% of the MLGs were unique. The high number of unique MLGs
identified using SSR markers highlights the substantial genetic variation within populations despite their
predominantly clonal structure (Forbes *et al.*, 1998; Dey *et al.*, 2018). This difference in diversity could be attributed
to the origin of the pathogen and its history of introduction (Saville *et al.*, 2016).
490
491 Our findings demonstrate that *P. infestans* populations in South America exhibit strong lineage-based genetic
492 structure shaped by regional ecological conditions and the movement of plant material. These results highlight the
493 need for transnational genetic surveillance strategies to monitor the emergence and spread of clonal lineages that
494 may affect late blight management.
495

Author contributions

MI: coordination, sample processing, molecular laboratory work, marker scoring, analysis, interpretation of data,
and drafting of the manuscript.
MC: sampling
WP: coordination, interpretation of data and revision of the manuscript
LS: sampling, sample processing
SG: sample processing.
DV: sampling
VC: coordination
BG: coordination, interpretation of data
JK: interpretation of data and revision of the manuscript.
507

Acknowledgements

509 We would like to thank Freddy Ventura and Marcelo Vinueza for their technical support during the sample
510 processing.

Data availability statement

513 Not applicable
514

Ethic declaration

516 Not applicable
517

Funding

519 This study was supported by the CGIAR Plant Health Initiative (PHI), funded by CGIAR Trust Fund Donors
520 (https://www.cgiar.org/funders/) and the "The Feed the Future Innovation Lab for Current and Emerging Threats to
521 Crops" provided by the United States Agency for International Development (USAID) cooperative agreement No:
522 7200AA21LE00005. Field collection in Bolivia was funded by ASDI-DICYT-UMSS, in Colombia, by Agrosavia
523 under the project BPIN 2018000100186 and in Uruguay, by INIA.

Conflict of Interest

The authors declare that they have no affiliations with or involvement in any organization or entity with any financial interest in the subject matter or materials discussed in this manuscript.

References

Acuña I, Bravo R, Gutiérrez M *et al.*, 2019. *Tizón tardío de la papa: Estrategia de manejo integrado con alertas temprana*. Osorno, Chile: Instituto de Investigación Agropecuaria.

Adler NE, Erselius LJ, Chacón MG *et al.*, 2004. Genetic Diversity of Phytophthora infestans sensu lato in Ecuador Provides New Insight Into the Origin of This Important Plant Pathogen. *Phytopathology®* **94**, 154–162.

Andrivon D, 1996. The origin of populations present in Europe in the 1840s: a critical review of historical and scientific evidence. *Plant Pathology* **45**, 1027–1035.

Briceño A, San Román M, Moreno M *et al.*, 2009. Population structure of *Phytophthora infestans* in the venezuelan andes (2004-2007). *Acta Horticulturae* **834**, 129–140.

Bruvo R, Michiels NK, D'Souza TG, Schulenburg H, 2004. A simple method for the calculation of microsatellite genotype distances irrespective of ploidy level. *Molecular Ecology* **13**, 2101–2106.

Brylińska M, Sobkowiak S, Stefańczyk E, Śliwka J, 2018. Evaluation of PCR markers for Phytophthora infestans mating type determination. *European Journal of Plant Pathology* **152**, 33–44.

Cárdenas M, Grajales A, Sierra R *et al.*, 2011. Genetic diversity of Phytophthora infestans in the Northern Andean region. *BMC Genetics* **12**, 23.

Céspedes MC, Cárdenas ME, Vargas AM *et al.*, 2013. Physiological and molecular characterization of Phytophthora infestans isolates from the Central Colombian Andean Region. *Revista Iberoamericana De Micologia* **30**, 81–87.

Chacón MG, Andrade-Piedra JL, Gessler C, Forbes GA, 2007. Aggressiveness of Phytophthora infestans and phenotypic analysis of resistance in wild Petota accessions in Ecuador. *Plant Pathology* **56**, 549–561.

Clark LV, Jasieniuk M, 2011. POLYSAT: an R package for polyploid microsatellite analysis. *Molecular*

Ecology Resources **11**, 562–566.

Cooke D, 2020. *Protocol for sampling* Phytophthora infestans *DNA using FTA cards*.

Cooke DEL, Cano LM, Raffaele S *et al.*, 2012. Genome Analyses of an Aggressive and Invasive Lineage

of the Irish Potato Famine Pathogen. *PLOS Pathogens* **8**, e1002940.

Cooke DEL, Lees AK, 2004. Markers, old and new, for examining Phytophthora infestans diversity.

Plant Pathology **53**, 692–704.

Coomber AL, Saville AC, Carbone I, Martin M, Bieker VC, Ristaino JB, 2025. A pangenome analysis

reveals the center of origin and evolutionary history of Phytophthora infestans and 1c clade

species. *PLOS ONE* **20**, e0314509.

Deahl KL, Pagani MC, Vilaro FL, Perez FM, Moravec B, Cooke LR, 2003. Characteristics of

Phytophthora infestans Isolates from Uruguay. *European Journal of Plant Pathology* **109**, 277–

281.

Dey T, Saville A, Myers K *et al.*, 2018. Large sub-clonal variation in Phytophthora infestans from recent

severe late blight epidemics in India. *Scientific Reports* **8**, 4429.

Drenth A, Tas ICQ, Govers F, 1994. DNA fingerprinting uncovers a new sexually reproducing population

ofPhytophthora infestans in the Netherlands. *European Journal of Plant Pathology* **100**, 97–107.

Drenth, A., Turkensteen, L. J., & Govers, F. (1993). The occurrence of the A2 mating type of

Phytophthora infestans in the Netherlands; significance and consequences. *Netherlands Journal*

of Plant Pathology, *99*(3), 57–67. https://doi.org/10.1007/BF03041396

Dueñas A, Betancur J, Galindo R, 2007. Estructura y composición florística de un bosque húmedo

trópical del Parque Nacional Natural Catatumbo Barí, Colombia. *Colombia forestal* **10**, 26–39.

Forbes GA, Escobar XC, Ayala CC *et al.*, 1997. Population Genetic Structure of Phytophthora infestans

in Ecuador. *Phytopathology* **87**, 375–380.

Forbes GA, Gamboa S, Lindqvist-Kreuze H, Oliva RF, Perez W, 2016. Identification of an A2 population

of Phythophthora andina attacking tree tomato in Peru indicates a risk of sexual reproduction in

this pathosystem. *Plant Pathology* **65**, 1109–1117.

Forbes GA, Goodwin SB, Drenth A, Oyarzun P, Ordoñez ME, Fry WE, 1998. A Global Marker Database

for Phytophthora infestans. *Plant Disease* **82**, 811–818.

Fry WE, 2020. Phytophthora infestans: the itinerant invader; "late blight": the persistent disease.

Phytoparasitica **48**, 87–94.

Fry WE, Birch PRJ, Judelson HS *et al.*, 2015. Five Reasons to Consider Phytophthora infestans a

Reemerging Pathogen. *Phytopathology®* **105**, 966–981.

Gabriel J, Magne J, Veramendi S *et al.*, 2018a. El tizón de la papa (Phytophthora infestans) en Bolivia:

Retos y oportunidades. *Revista Latinoamericana de la Papa* **22**, 80.

Gabriel J, Magne J, Veramendi S *et al.*, 2018b. El tizón de la papa (Phytophthora infestans) en Bolivia:

Retos y oportunidades. *Revista Latinoamericana de la Papa* **22**, 80.

Gamboa S, Pérez W, Andrade-Piedra JL, Forbes G, 2019. *Laboratory manual for Phytophthora infestans*

work at CIP. International Potato Center.

Garry G, Salas A, Forbes GA, Perez W, Cruz MS, Nelson RJ, 2005. Host Specialization not Detected

Among Isolates of the EC-1 Lineage of Phytophthora infestans Attacking Wild and Cultivated

Potatoes in Peru. *European Journal of Plant Pathology* **113**, 71–81.

Gilchrist Ramelli E, Jaramillo Villegas S, Afanador Kafuri L, Arango Isaza RE, 2009. Characterization of

Phytophthora infestans populations in Antioquia Colombia. *Revista Facultad Nacional de*

Agronomía Medellín **62**, 5031–5037.

Gómez-Alpizar L, Carbone I, Ristaino J, 2007. An Andean origin of Phytophthora infestans inferred from

mitochondrial and nuclear gene genealogies. *Proceedings of the National Academy of Sciences of*

the United States of America **104**.

Gómez-Alpizar L, Hu C-H, Oliva R, Forbes G, and JBR, 2008. Phylogenetic relationships of
601   Phytophthora andina, a new species from the highlands of Ecuador that is closely related to the
602   Irish potato famine pathogen Phytophthora infestans. *Mycologia* **100**, 590–602.

603   Goodwin, S.B., Cohen, B.A., Fry, W.E. WE, 1994. Panglobal distribution of a single clonal lineage of the
604   Irish potato famine fungus.

605   Goodwin, S.B., Sujkowski, L.S., Dyer, A.T., Fry, B.A., Fry, W.E., 1995. Direct Detection of Gene Flow
606   and Probable Sexual Reprroduction of Phytophthora infestans in Northern North América.

607   Goss EM, Tabima JF, Cooke DEL *et al.*, 2014. The Irish potato famine pathogen Phytophthora infestans
608   originated in central Mexico rather than the Andes. *Proceedings of the National Academy of*
609   *Sciences of the United States of America* **111**, 8791–8796.

610   Griffith GW, Shaw DS, 1998. Polymorphisms in Phytophthora infestans: Four Mitochondrial Haplotypes
611   Are Detected after PCR Amplification of DNA from Pure Cultures or from Host Lesions. *Appl.*
612   *Environ. Microbiol.* **64**, 4007–4014.

613   Grünwald NJ, Flier WG, 2005. The biology of Phytophthora infestans at its center of origin. *Annual*
614   *Review of Phytopathology* **43**, 171–190.

615   Guha Roy, S., Dey, T., Cooke, D. E. L., & Cooke, L. R. (2021). The dynamics of Phytophthora infestans
616   populations in the major potato-growing regions of Asia – A review. *Plant Pathology*, *70*(5),
617   1015–1031. https://doi.org/10.1111/ppa.13360

618   Hedrick PW, 1999. Perspective: Highly variable loci and their interpretation in evolution and
619   conservation. *Evolution; International Journal of Organic Evolution* **53**, 313–318.

620   Judelson HS, Spielman LJ, Shattock RC, 1995. Genetic mapping and non-Mendelian segregation of
621   mating type loci in the oomycete, *Phytophthora infestans. Genetics* **141**, 503–512.

622   Kamvar ZN, Tabima JF, Grünwald NJ, 2014. Poppr: an R package for genetic analysis of populations
623   with clonal, partially clonal, and/or sexual reproduction. *PeerJ* **2**, e281.

Knaus BJ, Tabima JF, Shakya SK, Judelson HS, Grünwald NJ, 2020. Genome-Wide Increased Copy

Number is Associated with Emergence of Dominant Clones of the Irish Potato Famine Pathogen

Phytophthora infestans. *mBio* **11**, e00326-20.

Kool J, Evenhuis A, 2023. Early detection of Phytophthora infestans in potato plants using hyperspectral

imaging, local comparison and a convolutional neural network. *Smart Agricultural Technology* **6**,

100333.

Lees AK, Wattier R, Shaw DS, Sullivan L, Williams NA, Cooke DEL, 2006. Novel microsatellite

markers for the analysis of Phytophthora infestans populations. *Plant Pathology* **55**, 311–319.

Li Y, Cooke D, Jacobsen E, van der Lee T, 2013. Efficient multiplex simple sequence repeat genotyping

of the oomycete plant pathogen Phytophthora infestans. *Journal of Microbiological Methods* **92**,

316–322.

Lindqvist☐Kreuze H, Gamboa S, Izarra M *et al.*, 2020. Population structure and host range of the potato

late blight pathogen *Phytophthora infestans* in Peru spanning two decades. *Plant Pathology* **69**,

334–346.

Lucca, F., Acuña, I., Tello, C., & Morales, R. (2023). Mapa poblacional de *P. infestans* en países

miembros. (Actividad 4.5. Elaboración de un mapa poblacional de *P. infestans*). *FONTAGRO*,

Producto 22, 39.

Lucca MF, Restrepo S, Danies G, Acuña I, Zanotta S, 2019. Monitoring of Phytophthora spp on potato

crops and Solanaceous host in Latin America. In: *17th EuroBlight 2019 Workshop*. York, UK,.

Ludwiczewska M, Janiszewska M, Yin Z, Śliwka J, 2025. Populations of Phytophthora infestans in

northern and eastern Europe. *European Journal of Plant Pathology* **171**, 81–95.

Mariette N, Mabon R, Corbière R *et al.*, 2016. Phenotypic and genotypic changes in French populations

of Phytophthora infestans: are invasive clones the most aggressive? *Plant Pathology* **65**, 577–586.

Martin Md, Vieira Fg, Ho Sy *et al.*, 2016. Genomic Characterization of a South American Phytophthora

Hybrid Mandates Reassessment of the Geographic Origins of Phytophthora infestans. *Molecular*

biology and evolution **33**.

Martin FN, Zhang Y, Cooke DEL, Coffey MD, Grünwald NJ, Fry WE, 2019. Insights into evolving global populations of Phytophthora infestans via new complementary mtDNA haplotype markers and nuclear SSRs. *PLOS ONE* **14**, e0208606.

Mideros MF, Turissini DA, Guayazán N *et al.*, 2018. Phytophthora betacei, a new species within Phytophthora clade 1c causing late blight on Solanum betaceum in Colombia. *Persoonia* **41**, 39–55.

Njoroge AW, Andersson B, Lees AK *et al.*, 2019. Genotyping of *Phytophthora infestans* in Eastern Africa Reveals a Dominating Invasive European Lineage. *Phytopathology®* **109**, 670–680.

Olave-Achury A, Cardenas D, Restrepo S *et al.*, 2022. Phenotypic and Genotypic Characterization of Phytophthora infestans Isolates Associated with Tomato and Potato Crops in Colombia. *Phytopathology®* **112**, 1783–1794.

Oliva RF, Kroon LPNM, Chacón G, Flier WG, Ristaino JB, Forbes GA, 2010. Phytophthora andina sp. nov., a newly identified heterothallic pathogen of solanaceous hosts in the Andean highlands. *Plant Pathology* **59**, 613–625.

Oyarzun Pj, Pozo A, Ordoñez Me, Doucett K, Forbes Ga, 1998. Host Specificity of Phytophthora infestans on Tomato and Potato in Ecuador. *Phytopathology* **88**.

Patarroyo C, Lucca F, Dupas S, Restrepo S, 2024. Reconstructing the Global Migration History of Phytophthora infestans Toward Colombia. *Phytopathology* **114**, 2151–2161.

Perez WG, Gamboa JS, Falcon YV, Coca M, Raymundo RM, Nelson RJ, 2001. Genetic Structure of Peruvian Populations of *Phytophthora infestans*. *Phytopathology®* **91**, 956–965.

R Core Team, 2019. R: A language and environment for statistical computing.

Rhouma A, Hajji-Hedfi L, Atallaoui K, 2024. Potato late blight: the pathogen, the menace, the sustainable control. *DYSONA - Life Science* **5**, 37–51.

Runno-Paurson E, Hannukkala A, Trdan S, Williams I, Koppel M, Mänd M, 2012. The structure of mating type, virulence, metalaxyl resistance of Phytophthora infestans in a long-term phenotypic

study in distinct location in Eastern Estonia. *Journal of Plant Diseases and Protection* **119**, 45–

52.

Ryley MJ, Drenth A, 2024. A matter of where and when - the appearance of Late Blight of potato in

Australia. *Historical Records of Australian Science* **35**, 213–222.

Saffer A, Tateosian L, Saville AC, Yang Y-P, Ristaino JB, 2024. Reconstructing historic and modern

potato late blight outbreaks using text analytics. *Scientific Reports* **14**, 2523.

Saville AC, Martin MD, Ristaino JB, 2016. Historic Late Blight Outbreaks Caused by a Widespread

Dominant Lineage of *Phytophthora infestans* (Mont.) de Bary. *PLOS ONE* **11**, e0168381.

Saville A, Ristaino JB, 2019. Genetic Structure and Subclonal Variation of Extant and Recent U.S.

Lineages of Phytophthora infestans. *Phytopathology* **109**, 1614–1627.

Saville AC, Ristaino JB, 2021. Global historic pandemics caused by the FAM-1 genotype of

Phytophthora infestans on six continents. *Scientific Reports* **11**, 12335.

Shannon CE, 1948. A mathematical theory of communication. *The Bell System Technical Journal* **27**,

379–423.

Shannon CE, 2001. A mathematical theory of communication. *ACM SIGMOBILE mobile computing and*

communications review **5**, 3–55.

Shattock R, 2002. Phytophthora infestans: populations, pathogenicity and phenylamides. *Pest*

management science **58**.

Simpson EH, 1949. Measurement of Diversity. *Nature* **163**, 688–688.

Vargas AM, Quesada Ocampo LM, Céspedes MC *et al.*, 2009. Characterization of Phytophthora infestans

populations in Colombia: first report of the A2 mating type. *Phytopathology* **99**, 82–88.

Yuen JE, Andersson B, 2013. What is the evidence for sexual reproduction of Phytophthora infestans in

Europe? *Plant Pathology* **62**, 485–491.

Tables

700   **Table 1.** Summary of samples collected and analyzed in this study.
701

Country	Cultivated potatoes	other species	Department	Species names	Altitude (masl)	collecting year
Bolivia	45	22	Cochabamba	*S. brevicaule, x Solanum sp, S.tuberosum, S. microdontum*	2381-3922	2018-2019
Peru	1		Cusco		3531	2018
	4		Apurimac	*S. tuberosum*	3301-4068	2018
	6		Ayacucho		2732-3460	2018
	19		Puno		3440-3965	2017-2018
Ecuador	3	8	Carchi	*S.tuberosum, S.ochrantum*	2960-3351	1993-2005
	13	1	Pichincha	*S.tuberosum, S.lycopersicum, S.muricatum, S.quitoense, S.ochrantum*	1671-3300	1996-2022
	2	1	Imbabura	*S.lycopersicum, S.caripense*	2018-3117	1996-2002
	7	1	Tungurahua	*S.tuberosum, S.betaceum, S.brevifolium*	2031-2960	2001-2003
	1	1	Chimborazo	*S.tuberosum, S.andreanum*	2961-3282	2000-2004
	1	4	Cotopaxi	*S.tuberosum, S.paucijugum*	2558-3895	2001-2005
	2	7	Napo	*S.colombianum, S.quitoense*	1748-3729	1993-2002
		1	M. Santiago	*S.andreanum*	981	1998
	2		Pastaza	*S.quitoense*	1005-3020	2000-2001
		2	Bolívar	*S.andreanum*	2768-2829	2002
Colombia	6		Norte de Santander	*S.tuberosum*	2574-2842	2022
	8		Santander	*S.tuberosum*	2129-2401	2022
Uruguay	10		Canelones	*S.tuberosum*	30-52	2023-2025
	4		San José	*S.tuberosum*	28-37	2024-2025
Grand total	134	48				

702
703
704
705 **Table 2**. Diversity statistics for microsatellite data across 12 loci in *Phytophthora* populations from Bolivia, Peru,
706 Colombia, Ecuador, and Uruguay categorized by location
707

Pop	N	MLG	eMLG	SE	H	G	lambda	E.5	Hexp	Ia	rbarD
B.Coch	67	32	7.2	1.329	2.814	7.7	0.87	0.428	0.515	0.6031	0.0988
P.Cuz	1	1	1	0	0	1	0	NaN	0.75	NA	NA
P.Apu	4	3	3	0	1.04	2.67	0.625	0.912	0.447	1.193	0.3115
P.Aya	6	6	6	0	1.792	6	0.833	1	0.421	-0.1682	-0.0471

	N	MLG	eMLG	SE	H	G	lambda	E.5	Hexp	Ia	rbarD
P.Pu	19	12	7.22	1.075	2.229	7.08	0.859	0.733	0.528	6.1668	0.6812
C.NSt	6	6	6	0	1.792	6	0.833	1	0.524	-0.0676	-0.0233
C.St	8	6	6	0	1.733	5.33	0.812	0.931	0.469	1.9952	0.4029
E.Pich	14	13	9.51	0.5	2.54	12.25	0.918	0.963	0.572	3.1127	0.3117
E.Car	11	11	10	0	2.398	11	0.909	1	0.338	0.3187	0.0489
E.Imb	3	2	2	0	0.637	1.8	0.444	0.899	0.58	5.2308	1
E.Tung	8	7	7	0	1.906	6.4	0.844	0.943	0.612	5.2196	0.4971
E.Chim	2	2	2	0	0.693	2	0.5	1	0.428	NA	NA
E.Coto	5	5	5	0	1.609	5	0.8	1	0.344	-0.7882	-0.1377
E.Nap	9	9	9	0	2.197	9	0.889	1	0.488	3.6799	0.4186
E.Mor	1	1	1	0	0	1	0	NaN	0.833	NA	NA
E.Pas	2	2	2	0	0.693	2	0.5	1	0.617	NA	NA
E.Bol	2	2	2	0	0.693	2	0.5	1	0.383	NA	NA
U.Can	10	7	7	0	1.748	4.55	0.78	0.747	0.48	0.9565	0.2092
U.SnJ	4	3	3	0	1.04	2.67	0.625	0.912	0.488	0.2264	0.1278

N number of individuals, *MLG* number of multilocus genotypes, *eMLG* expected number of MLGs, *SE* Standard error. The next columns are considering the MLG diversity: *H* Shannon- Weiner Index, *H Stoddart and Taylor's Index (G), lambda* Simpson's index, *E.5* Evenness, H_{exp} Nei's unbiased gene diversity. *Ia* Index of Association, *rbarD* standardized index of association.

Table 3. Population statistics for the 12 microsatellite loci in *Phytophthora* populations from Bolivia, Peru, Colombia, Ecuador, and Uruguay

locus	allele	1-D[b]	Hexp[b]	Evenness[b]
D13	17	0.67	0.67	0.47
SSR8	5	0.64	0.64	0.84
SSR4	12	0.81	0.81	0.8
Pi04	4	0.61	0.61	0.89
Pi70	3	0.4	0.4	0.72
SSR6	7	0.54	0.54	0.77
Pi63	3	0.61	0.61	0.88
G11	18	0.7	0.7	0.53
SSR3	7	0.61	0.61	0.67
SSR11	5	0.62	0.62	0.8
SSR2	3	0.38	0.38	0.63
Pi4B	11	0.59	0.59	0.63
mean	7.92	0.6	0.6	0.72

[b] 1–D, Simpson's diversity index. *H*exp, Nei's gene diversity. Evenness indices of evenness.

Table 4. Analysis of molecular variance (AMOVA) of *Phytophthora* isolates from Bolivia, Peru, Colombia, Ecuador, and Uruguay

	Components of covariance	Monte-Carlo test for significance nrepet=999

	Df	Sum Sq	Mean Sq	Sigma	%	Obs	Std.Obs	Alter	Pvalue
Within samples	114	69.56	0.61	0.61	32.02	0.61	-26.29	less	0.001
Between samples Within clonal lineage	23	33.51	1.46	0.19	10.21	0.19	10.00	greater	0.001
Between clonal lineage	11	129.85	11.80	1.10	57.77	1.10	9.73	greater	0.001
Total	148	232.92	1.57	1.91	100.00				

Supplementary Information

728
729   Supplementary materials are provided as separate files.