Running Head: *Coccidioides* hotspots detected along river and stream environments in the San

Joaquin Valley of California

Title: Urban soils along the Kern River and Los Gatos Creek are hotspots for *Coccidioides* in the

San Joaquin Valley of California

Authors: Robert Wagner[1*], Liliam Montoya[1], Molly Radosevich[2], Justin Remais[2], John W.

Taylor[1*]

[1]Department of Plant & Microbial Biology, University of California Berkeley, Berkeley,

California, USA.

[2]Division of Environmental Health Sciences, University of California Berkeley, Berkeley,

California, USA.

Corresponding authors:

*robertleafwagner@gmail.com, *jtaylor@berkeley.edu

Abstract

Coccidioidomycosis (Valley fever) is acquired through inhalation of spores produced by fungi in

the genus *Coccidioides*. *Coccidioides* is commonly detected in, and cultivated from, the lung

tissue of native rodents and the soils within their burrows. Coccidioidomycosis is acquired by

exposure to environmentally produced spores and is not spread between hosts. Thus, determining

the location of rodent burrows with soils harboring *Coccidioides* will be critical for

understanding coccidioidomycosis incidence and modelling how *Coccidioides* distributions will

be affected by global change. *Coccidioides* is readily detected in rodent burrows on undisturbed

land, has not been detected on agricultural land, and is unstudied on urban land. To test the

hypothesis that *Coccidioides* is in urban soil, we sampled rodent burrow soils from the banks of

two water ways, the Kern River and Los Gatos Creek, which transect two cities, Bakersfield and

Coalinga, in the San Joaquin Valley of California. To test the hypothesis that *Coccidioides* would

not be detected at higher elevations, we extended our sampling of rodent burrows along

waterways into the mountains of California. From 1178 soil and settled dust samples, we find

that *Coccidioides* is found in urban riparian environments in Bakersfield and Coalinga, on

riparian land on the floor of the San Joaquin Valley but not at higher elevations and is negatively

correlated with modeled soil moisture. *Coccidioides* shows significant co-occurrence patterns

with animal-associated fungal taxa, but no broader relationships with the greater fungal

community. Our results warrant caution when excavating urban rodent burrows in the region.

Author Summary

Coccidioidomycosis, or Valley fever, is caused by a fungus that primarily infects people when its

spores become airborne and are inhaled. Although the fungus is found in the burrows of rodents,

its presence in cities has been unclear. We investigated whether urban environments in

California's San Joaquin Valley, an area with high Valley fever incidence, harbor the fungus. By

sampling more than a thousand soils and settled dust samples along riparian areas (waterways

and adjacent environments) in Bakersfield and Coalinga, we found that the fungus that causes

Valley fever, *Coccidioides,* occurs in urban riparian zones. The fungus was restricted to low

elevations and was more likely to be detected in drier soils. We also compared positive and

negative soils using DNA-based surveys of fungi. While the broader fungal community showed

no strong relationship to *Coccidioides*, several species tied to animal activity were more common

in positive samples, supporting a link to rodent hosts. These findings show that urban waterways

can serve as habitat for *Coccidioides*. Because these areas lie within major population centers,

identifying and avoiding disturbance of rodent burrows may reduce exposure risk, and future

models of Valley fever should account for riparian corridors and local ecological conditions.

Introduction

Coccidioides immitis and *Coccidioides posadasii* are Ascomycete fungi, within the Onygenales,

that are responsible for the human disease coccidioidomycosis, colloquially known as "Valley

fever" and "Desert Rheumatism" (Posadas 1892, Ophüls 1905, Kollath et al. 2019).

Coccidioidomycosis is an important disease in arid regions of the western United States, causing

nearly 200 deaths and generating $3.9 billon in healthcare costs per year (Huang et al. 2012,

Centers for Disease Control and Prevention 2018, Gorris et al. 2021). In California, Washington

and areas generally west of the Sierra Nevada-Cascade mountain ranges, only *C. immitis* has

been found, while *C. posadasii* has been observed elsewhere in the United States (Koufopanou et

al. 1997, Johnson et al. 2014, Marsden-Haug et al. 2014, Teixeira and Barker 2016, Chow et al.

2021), as well as in Mexico (Baptista-Rosas et al. 2012), Central America (Engelthaler et al.

2016) and South America (Fisher et al. 2001, De Macêdo et al. 2011, Teixeira et al. 2019).

Although *Coccidioides* has been found most abundantly in undeveloped arid lands, it remains

uncertain whether exposure within cities results from local sources or from travel to nearby rural

areas where *Coccidioides* is established. Understanding possible urban exposure is therefore

critical for public health, especially in cities such as Bakersfield (population ~400,000) and

Coalinga (~17,000), which are major population centers in the SJV.

Humans acquire coccidioidomycosis by inhaling arthroconidia from environmental sources

(Nguyen et al. 2013) and human-to-human transmission is limited to organ transplantation

(Nokes and Blair 2020), which makes determining the geographic location and seasonal timing

of arthroconidial development and dispersal important for understanding coccidioidomycosis

incidence. All fungi are heterotrophs (organisms that obtain energy and carbon from organic

matter rather than producing it themselves), and most fungi, Ascomycota included, consume

plants for nutrition. An association between *Coccidioides* spp. and rodents was discovered in the

1940s, suggesting that rodents may serve as a natural reservoir. (Emmons 1942, Emmons et al.

1942). This relationship has been reaffirmed by recent molecular work (Salazar-Hamm 2021,

Salazar-Hamm et al. 2022). An explanation for this association comes from genomic analyses

showing that *Coccidioides* spp. and closely related genera in the Onygenales evolved away from

an ancestral state of using plant carbohydrates for nutrition and to a derived state of using animal

protein (Sharpton et al. 2009, Whiston and Taylor 2016). *Coccidioides* spp. are significantly

more likely to be found in soil collected from within rodent burrows rather than soil collected

apart from rodent burrows (Egeberg and Ely 1956, Elconin et al. 1957, Kollath et al. 2020,

Wagner et al. 2023, Head et al. 2024).

There is evidence that *Coccidioides* spp. have evolved a commensal relationship with native

rodents, and can persist in the lungs in a dormant state, arrested by the host's immune system

(Sun and Huppert 1976, Rappleye and Goldman 2006, Hsu et al. 2022). This persistence

supports the hypothesis that *Coccidioides* functions as an endozoan (occupies an animal host)

before reverting to a saprotrophic (decomposer) phase following host death (Taylor and Barker

2019). Fungal endozoans may live as benign parasites in animal hosts without observable

symptoms and, following host death, use the host carcass to grow as hyphae and reproduce by

arthroconidia.

We previously found that 37% of 238 soils sampled from burrows on undeveloped land in the

southwestern San Joaquin Valley (SJV) were positive for *Coccidioides* spp., whereas none of 472

agricultural surface soils sampled from four farms in the SJV were positive for *Coccidioides* spp.

(Wagner et al. 2023). There was an observed absence of rodent burrows in agricultural fields at

all agricultural sites, likely owing to cultivation, irrigation, and efforts by farmers to eradicate

rodents (Baldwin et al. 2016, Lloyd and Baldwin 2021). Unlike undisturbed and agricultural soil,

urban soil remains nearly unstudied for *Coccidioides* (Lauer et al. 2012). Having observed that

native rodents excavate burrows in the banks of waterways, we hypothesized (H_1) that burrow

soil from urban riparian habitats would harbor *Coccidioides*. Riparian habitats are vegetated

corridors bordering rivers or streams that differ sharply from surrounding arid landscapes. Here,

we report on the presence of *Coccidioides* in soils sampled from rodent burrows along riparian

corridors traversing two cities in the SJV, Bakersfield and Coalinga. Riparian corridors in arid

regions have higher plant biomass than surrounding areas (Scott et al. 2014), which provides

food for native animals (Case and Kauffman 1997, Pasternak et al. 2013) and increases their

abundance and diversity (Patten 1998, Merritt and Bateman 2012, Free et al. 2015). This

diversity includes all mammals (Falck et al. 2003, Hamilton et al. 2015), including rodents native

to the SJV (Grinnell 1923, Whitford and Kay 1999). Previous research has shown that rodent

burrow soils in arid environments host a rich diversity of fungal species (Hawkins 1996, 1999,

Wagner et al. 2022), many of which could potentially interact with *Coccidioides.*

Studies of *Coccidioides* in California soils have traditionally focused on the SJV rather than in

the adjacent mountains, probably because only one coccidioidomycosis outbreak has been

reported at higher elevation (975m), at a Native American midden near Inyokern, California

(Plunkett and Swatek 1957). In this previous study, excavated soils within the midden were

positive for *Coccidioides*, while soils from outside the midden were negative for *Coccidioides*, as

were the lungs of hundreds of native rodents from the area (Plunkett and Swatek 1957). The

Kern River and Los Gatos Creek both originate in nearby mountains, which provided us the

opportunity to extend our sampling of rodent burrow soils along an elevational gradient. In doing

so, we address a knowledge gap about the distribution of *Coccidioides* and challenge a second

hypothesis, (H_2), that *Coccidioides* would not be found in soils sampled from rodent burrows in

foothill regions between the SJV floor and elevations as high as 900m.

Most previous research investigating coccidioidomycosis incidence, in the context of landscape

environmental variables, has focused on associations with dry, arid soils (Egeberg and Ely 1956,

Elconin et al. 1957, Greene et al. 2000, Kollath et al. 2020), and overlooked riparian and

riparian-adjacent habitats. Our choice to sample soils from riparian rodent burrows situated in the

banks of rivers and streams allowed us to investigate soils that remained relatively dry while

being close to water and food resources that support rodent populations. More generally, riparian

corridors within the SJV represent some of the last continuous, undeveloped habitats in the

region. These waterways bisect large urban areas, potentially situating *Coccidioides* populations

in proximity to dense human populations. Finally, only two other studies have compared soil

moisture with direct detections of *Coccidioides* in soils (Lauer et al. 2012, Head et al. 2024),

allowing us to continue to explore this important knowledge gap.

Here, we survey soil from rodent burrows in two riparian corridors in the SJV, one along the

Kern River, which transects Bakersfield, California, and another along Los Gatos Creek, which

transects Coalinga, California. We report the presence of *Coccidioides* in terms of location along

waterways and, at some locations, seasonality. To understand variation in the presence of

Coccidioides and the composition of the fungal community, we consider the location of burrows,

associations between *Coccidioides* and the soil fungal community, and potential influences of

elevation and estimated soil moisture at the time of sampling. Our results help clarify key

knowledge gaps about the distribution of *Coccidioides* that may impact the location of

arthroconidial development and dispersal. We hope that our research will stimulate research

regarding transmission of *Coccidioides* among native rodents and the dispersal of *Coccidioides*

spores in ambient air and dust, as well as the inclusion of these aspects of the *Coccidioides*

environmental lifecycle into future predictive models regarding spatial and temporal

Coccidioides distributions.

Results

1178 samples (soil and dust) were collected and analyzed for the presence of *Coccidioides* using

the CocciEnv qPCR assay. Soil samples were collected from 625 rodent burrows from 35 sites

along the Kern River, 205 rodent burrows from seven sites in the greater Bakersfield, California

area, 248 rodent burrows or adjacent soils from 16 sites along Los Gatos Creek near Coalinga,

California, and 100 urban settled dust samples collected from public spaces in Coalinga,

California. The soil fungal community was characterized for 68 soil samples (28 from the Kern

River and 40 from Los Gatos Creek), by amplicon sequencing of the ITS2 region of fungal

ribosomal DNA.

Coccidioides detection

Coccidioides was detected in 6.3% (52/830) of rodent burrow soil samples from sites along the

Kern River and in and around the urban Bakersfield area and, of these, was detected in 8.3%,

52/625) of rodent burrow soil samples from sites directly along the Kern River and 0% (0/205)

of sites in Bakersfield away from the Kern River (S2 Table). From sites along Los Gatos Creek,

Coccidioides was detected in 23.2% (53/228) of rodent burrow soil samples, 50.0% (10/20) of

surface soil samples and in 2% (2/100) of settled dust samples collected from public surfaces in

Coalinga (S1 Fig, S2 Table). All surface soil samples were collected from the "Highway 33

West" site at Los Gatos Creek in August 2020, from directly outside the entrances (<1m) from

rodent burrows. It was not determined if these surface soils were otherwise associated with the

burrows they were collected near. All but two *Coccidioides* positive soil samples from Kern

River sites were collected centrally in Bakersfield (Fig 1a), at sites with soil moisture between

0.045 – 0.100 cm^3/cm^3 (Fig 1b) and elevations between 98.56 and 192.59 meters (Fig 1c). The

other two positive samples were collected from Kern River Parkway West End, not far from the

intersection of the Kern River and the California Aqueduct, with soil moisture between 0.058 –

0.115 cm^3/cm^3 and elevations between 97.26 and 99.86 meters. Of *Coccidioides* positive soil

samples from Los Gatos Creek sites, 69.8% (44/63) were collected from areas within 200m of

urban or residential land in the town of Coalinga (Fig 2a). The remaining positive samples came

from either undeveloped land approximately 10km east of Coalinga (7 positive samples), where

soil moisture ranged from 0.052 to 0.136 cm³/cm³ and elevations ranged from 161.72 to 208.48

m (Fig 2a, 2b) or from undeveloped land west of Coalinga (14 positive samples) at elevations

below 373 m (Fig 2c, S2 Fig).

Fig 1. *Coccidioides* detection in rodent burrow soil samples along the Kern River near Bakersfield, California using the CocciENV qPCR assay. Elevation profile (meters above sea level) and qualitative categorization of sample locations (a). Estimated surface soil moisture (upper 5cm) at each timepoint across sampling locations (b). Map of sample coordinates (c). Inset shows map extent. Landscape classification derived from the California Farmland Mapping and Monitoring Program 2018. SMAP = Soil Moisture Active Passive (NASA satellite). Vertical banding in soil moisture is due to varying resolution from sub-9km landscape and vegetation differences, and downscaling algorithms, affecting the SMAP L3 version 3 model. Point colors correspond to positive replicate wells. Map data (lines) were acquired from OpenStreetMap (www.openstreetmap.org) under the Open Data Commons Open Database License (OdbL) (https://www.openstreetmap.org/copyright). Map tiles are by Stamen Design (www.stamen.com) under Creative Commons By Attribution (CC BY 4.0) (https://creativecommons.org/licenses/by/4.0/). California shapefile is public domain data by Natural Earth (http://www.naturalearthdata.com).

Fig 2. *Coccidioides* detection in rodent burrow and surface soil samples along Los Gatos Creek near Coalinga, California using the CocciENV qPCR assay. Elevation profile (meters above sea level) and qualitative categorization of sample locations (a). Estimated surface soil moisture (upper 5cm) at each timepoint across sampling locations (b). Map of sample coordinates (c). Inset shows map extent. Landscape classification derived from the California Farmland Mapping and Monitoring Program 2018. SMAP = Soil Moisture Active Passive (NASA satellite). Vertical banding in soil moisture is due to sub-9km landscape differences affecting model. Point colors correspond to positive replicate wells. Map data (lines) were acquired from OpenStreetMap (www.openstreetmap.org) under the Open Data Commons Open Database License (OdbL) (https://www.openstreetmap.org/copyright). Map tiles are by Stamen Design (www.stamen.com) under Creative Commons By Attribution (CC BY 4.0) (https://creativecommons.org/licenses/by/4.0/). California shapefile is public domain data by Natural Earth (www.naturalearthdata.com).

Of the 35 sites investigated along the Kern River (ignoring non-Kern River sites), more than

80% of samples positive for *Coccidioides* (45/52) were collected from two sites: University

Place (25/106 positive) and Panorama Park West (20/96 positive) (Fig 3a, b). Of the 16 sites

investigated along Los Gatos Creek, three sites provided the bulk of *Coccidioides* positive

samples: Highway 33 West (27/80 positive), Highway 33 East (15/40 positive), and Phelps

Avenue-10km (5/20 positive) (Fig 3c, d). Analyses including soil moisture were limited to

samples from Kern River sites. Across all Kern River sites, *Coccidioides* detection showed a

moderately significant association with sampling timepoint (p = 0.046, 0.039, 0.040, OR = 0.02,

0.02, 2.52E+03) and surface soil moisture (p = 0.049, OR = 0.06), and no significant association

with rootzone soil moisture (p = 0.104) or elevation (p = 0.076) (S5 Table). When excluding

samples collected above 210 meters, which excludes foothill sites east and upland of the Kern

River Canyon entrance, *Coccidioides* detection did not show significant associations with

sampling timepoint, surface moisture and rootzone moisture. This result indicates possible

collinearity between these terms, with higher elevations (within the context of lower elevation

sites) being a significant predictor (p = 0.014, OR = 3.70E+03) of *Coccidioides* detection (S6

Table). Taken together, *Coccidioides* presence did not show any strong or unambiguous

seasonality. Surface soil moisture along the Kern River, increased from west to east, from low to

high elevations, and during spring (April 2021) and winter (January 2022), as compared to

summer (2021) and fall (2021) (Fig 1b). Along Los Gatos Creek, soil moisture was higher in

winter than during the fall (Fig 2b). Despite including site as a random effect to account for site-

to-site variability, it's important to note that 87.5% of all Kern River positive detections for

Coccidioides occurred at two sites, University Place and Panorama Park West, and that

comparing only these two sites showed no significant associations between *Coccidioides*

detection and site, sampling timepoint or soil moisture (Fig 4, S7 Table).

293
Fig 3. *Coccidioides* detection hotspots. Panorama Park West and Panorama Park East, Kern
River (a). University Place, Kern River (b). Phelps Avenue 10km, Lost Gatos Creek (c).
Highway 33 West (in blue rectangle, includes 20 surface samples at the same points as burrow
samples), Highway 33 East and Warthan Creek, Los Gatos Creek (d). All points jittered to avoid
overlapping. Satellite imagery is public domain data provided by the United States Department
of Agriculture (USDA) National Agriculture Imagery Program (NAIP) (https://naip-
usdaonline.hub.arcgis.com).

Fig 4. The predicted probability of detecting *Coccidioides* in rodent burrows as a function of timepoint and site (University Place vs Panorama Park West). Mean predicted probabilities are displayed as a function of timepoint (a) and as a function of site (b). Points jittered to show sampling depth. P_{lrtest} = significance of difference with null model lacking a timepoint variable (a) or a site variable (b) derived from a log-ratio test. "rho squared" (ρ^2) = McFadden's-pseudo-R^2 for full logistic regression model. p = significance of full logistic regression model derived from deviance and null deviance. n = 202. Site error bar for October 2021 perfectly overlap.

Coccidioides and the soil fungal community

For Kern River samples, a significant (p = 0.002) association was found between the structure of

the soil fungal community and the site where soils were collected, representing 22% (r^2 = 0.22)

of the variance in Bray-Curtis dissimilarity among all identified fungal species (S8 Table). A

post-hoc, pairwise comparison showed a significant difference (p = 0.04, r^2 = 0.08) between

fungal communities from rodent burrow soil samples taken from Panorama Park West and

University Place (S9 Table), which, together, account for 23 of the 28 Kern River samples

sequenced for ITS2. However, when examining all sites, no significant associations were found

between the structure of the soil fungal community (Bray-Curtis dissimilarity) and the presence

of *Coccidioides* in soils. This result was true for rodent burrow samples collected from along the

Kern River (Fig 5a, S8 Table), as well as soils collected from within or just outside the entrance

of rodent burrows from Los Gatos Creek (Fig 5b, S10 Table). Significant correlations between

variables of interest and Bray-Curtis dissimilarity, and corresponding clustering on ordination

plots (of which we did not find here), are indicative of ecological relationships (Bray and Curtis

1957, Legendre and Legendre 2012). We found no such correlations here, which does not

preclude, but fails to provide evidence supporting an ecological relationship between the fungal

community and *Coccidioides*. It is notable that, within the examined fungal communities, species

of one genus, *Alternaria*, were most abundant among samples from both Kern River (14.6 ±

2.0%) and Los Gatos Creek (15.4 ± 1.7%) (S3 Fig, S4 Fig). *Alternaria* spp, were also seen to

dominate rodent burrow fungal communities in our previous study of sites along California

Highway 33 (Wagner et al. 2022). Other abundant genera were, from Kern River samples,

Cladosporium (11.5 ± 1.6%), *Trichophaeopsis* (6.5 ± 126%), *Neosetophoma* (5.1 ± 1.3%), and

Penicillium (4.6 ± 0.8%) and, from Los Gatos Creek samples, *Coniochaeta* (7.6 ± 1.8%),

Penicillium (7.4 ± 1.6%), *Cladosporium* (5.8 ± 0.9%), and *Chaetomium* (2.8 ± 0.6%) (S11

Table).

Fig 5. PCoA plot showing the Bray-Curtis dissimilarity within the fungal community between
individual rodent burrow soil samples from Kern River sites in Bakersfield, CA (a) and Los
Gatos Creek in Coalinga, CA (b). Kern River samples were collected from within rodent burrows
in April and July 2021. Los Gatos Creek samples were collected just outside (Entrance) and
within (Interior) rodent burrows in August 2020. Colored points = samples positive for
Coccidioides using the CocciENV qPCR assay. Gray points = samples negative for *Coccidioides*
using the CocciENV qPCR assay. Blue circle = sample positive for *Coccidioides* in the ITS2
rDNA dataset. CT = qPCR cycle threshold value. n = 28 (Kern River) and 40 (Los Gatos Creek).

Indicator species

From the 68 fungal communities sequenced from soils, in which a total of 764 unique fungal

species were detected, 83 fungal indicator species (p ≤ 0.05) were identified, 21 with

Coccidioides positive soils and 62 with *Coccidioides* negative soils. Soils from rodent burrow

interiors yielded 11 positive and seven negative associations from Kern River samples (S12

Table) and seven positive and 51 negative associations from Los Gatos Creek samples (S13

Table, 14 Table). Soils collected from outside rodent burrow entrances along Los Gatos Creek

provided three positive and four negative associations (S15 Table). The most significant ($p \leq$

0.01) associations with *Coccidioides* positive rodent burrow soils were found with *Coniochaeta*

discospora and *Montagnea arenaria* from Kern River samples and *Coniochaeta polymorpha*

from Los Gatos Creek samples. No similarly significant associations were found with

Coccidioides negative rodent burrow soils from Kern River samples. Among Los Gatos Creek

samples, negative associations with *Aspergillus* spp., *Chaetomium* spp., *Holtermanniella* spp.,

Ustilago bullata, *Paraphoma fimeti*, *Septoriella neoarundinis* and *Pseudoechria decidua* were

found for soils within rodent burrows, and with *Dothiora viticola* for soils from outside rodent

burrow entrances, at the same ($p \leq 0.01$) significance level. A single species, *Fusarium solani*,

was identified as an indicator species from both Kern River ($p = 0.03$) and Los Gatos Creek ($p =$

0.03) samples, and it was associated with *Coccidioides* negative samples.

Discussion

Our most important finding is that *Coccidioides* can be found in soils along seasonal waterways

in major population centers of the San Joaquin Valley; Bakersfield (population ~400,000) and

Coalinga (~17,000), where Valley fever incidence is high. This result was true for an ephemeral

stream on the west side of the SJV, Los Gatos Creek, in Coalinga, CA and for a seasonally wet,

major river, the Kern River, that runs east to west through Bakersfield, CA. We rarely found

Coccidioides in urban areas away from a waterway, though our sampling in these areas was more

limited. In Coalinga, *Coccidioides* was found in only two of the 100 settled dust samples we

collected, 0.6 - 3.2 km from Los Gatos Creek. At the Sports Village Complex in Bakersfield, CA,

13 km from the Kern River, none of 87 soils samples taken quarterly over one year was positive

for *Coccidioides*. Taken together, these results fail to disprove H$_1$, that *Coccidioides* would be

present in urban riparian areas. The inverse, that *Coccidioides* is absent from non-riparian urban

areas cannot be claimed because sampling intensity and site selection may have influenced

detection. The close juxtaposition of soils positive and negative for *Coccidioides* within sites

where the fungus was commonly detected, e.g., University Place and Panorama Park West (Fig

3b), suggests that *Coccidioides* distribution may not be uniform, consistent with previous reports

of sporadic detection in soils (Maddy 1965, Greene et al. 2000, Baptista-Rosas et al. 2012,

Litvintseva et al. 2015). While *Coccidioides* is associated with rodents and their burrows in the

SJV (Egeberg and Ely 1956, Elconin et al. 1957, Kollath et al. 2020), our study does not address

Coccidioides dispersal between rodent burrows or among rodent species.

We found a higher probability of detecting *Coccidioides* in drier soils, though this result is

almost certainly tied to other variables such as elevation and site choices. Previous research has

suggested that dry soils are associated with coccidioidomycosis incidence, potentially due to

competition between *Coccidioides* and other soil microbiota (Cordeiro et al. 2006, Kollath et al.

2019, Dobos et al. 2021, Head et al. 2024, Porter et al. 2024). Among our Kern River samples,

no *Coccidioides* positive samples were collected from sites with monthly average soil moisture

above 10% v/v. The higher soil moisture levels associated with drainage and concentrated water

flow in the Kern River Canyon may contribute to the absence of *Coccidioides* in soils at higher

elevations in our study. Downstream areas, where the Kern River is ephemeral due to agricultural

water diversion and flows only during high rainfall years, showed greater prevalence, suggesting

that relatively lower soil moisture may facilitate *Coccidioides* presence. However, regions with

extremely low soil moisture, such as between Bakersfield and Taft, presented limited detection,

possibly due to reduced availability of vegetation and water to support rodent hosts. These

observations suggest that soil moisture may influence *Coccidioides* presence, though other

factors are likely involved. Epidemiologic studies have shown that in typically arid regions,

periods of above-average rainfall are often followed by increased coccidioidomycosis incidence

several months to a year later, likely due to enhanced fungal growth during moist conditions and

subsequent spore release as soils dry (Comrie 2005, Park et al. 2005, Tamerius and Comrie 2011,

Head et al. 2022, Kollath et al. 2022). This pattern supports the idea that *Coccidioides* benefits

from intermittent or moderate moisture, but not persistently wet conditions. Our conclusions

regarding *Coccidioides* population dynamics relative to soil moisture are limited by 1) the

coarse, 9 km^2 grid scale resolution of modelled soil moisture data, 2) the fact that most positive

samples came from only two of the 42 Kern River and greater Bakersfield sites and 3)

covariation of soil moisture with elevation. Despite these limitations, our results provide a

landscape-level comparison of *Coccidioides* presence with modeled soil moisture,

contextualizing previous work linking soil moisture and coccidioidomycosis incidence (Gorris et

al. 2018).

A third observation is the absence of *Coccidioides* at elevations between 373 and 888 m, with

most positive detections below 250 m. This result fails to disprove H$_2$, that *Coccidioides* would

not be detected in foothill and montane sites and is likely influenced by soil moisture and other

environmental variables that correlate with elevation. We commonly observed the diurnal

California ground squirrel (*Otospermophilus beecheyi*) upslope at foothill and montane sites,

similar to our observations at low elevation sites on the valley floor. Deer mouse (*Peromyscus*

maniculatus) and San Joaquin pocket mouse (*Perognathus inornatus*), rodent species known to

harbor *Coccidioides* (Emmons 1942, Catalán-Dibene et al. 2014), are reported to inhabit sites

from Bakersfield to our highest sampling sites at Lake Isabella (Best 1993, Ocampo-Chavira et

al. 2020, Laabs et al. 2022). Thus, observed rodent host populations at higher elevations suggest

that host presence or absence alone cannot explain the absence of *Coccidioides* at these sites.

Fungal Community Characteristics

We did not find consistent associations between *Coccidioides* and overall soil fungal community

structure, consistent with both our previous research along California Highway 33, 60 km west

of Bakersfield, CA (Wagner et al. 2022) and more recent work done in the Carrizo Plain

(Radosevich et al. 2024). Our use of nucleic acids for identification does not distinguish between

dormant spores and actively growing mycelia, nor between living and dead material, limiting

inferences about community interactions. An abundance of spores relative to mycelia could bias

our understanding of the soil fungal community (Nilsson et al. 2019). Additional research will be

needed to determine if meaningful interactions between *Coccidioides* and the broader soil fungal

community exist. Low soil moisture may correlate with spore abundances in arid soils (Liu et al.

2009, Deveautour et al. 2020), and dry conditions can limit fungal growth (Ramirez et al. 2004,

Magan 2007). We did not directly measure soil moisture content, though we qualitatively

observed that the soils we sampled were dry.

Indicator species analysis identified several individual fungi associated with *Coccidioides*

positive samples, including *Coniochaeta discospora* and *Coniochaeta polymorpha*, as well as

Preussia spp., *Gymnoascus dugwayensis*, and *Myxotrichum deflexum*. Many of these taxa are

coprophilous or associated with animal-derived substrates, suggesting a potential link between

Coccidioides presence and rodent activity. This finding agrees with recent research showing that

rodent presence and soil *Coccidioides* presence are correlated in and around giant kangaroo rat

(*Dipodomys ingens*) burrows in the nearby Carrizo Plain (Head et al. 2024).

Coniochaeta spp. are soil saprotrophic and animal pathogenic fungi, of which many species are

coprophilous (live on animal dung) (Richardson 1998, Lytvynenko and Hayova 2018). There is

limited evidence that *Coniochaeta* species are keratinophilic (consumes keratin), having been

isolated from desert soil crusts using sheep wool enriched media (Hamm et al. 2020).

Coniochaeta discospora is commonly found on mammal dung (Cain 1957, Hanks 1963,

Mahoney and LaFavre 1981), while *Coniochaeta polymorpha* is a more recently identified

species isolated from human tracheal (Khan et al. 2013) and gastric (Heydari et al. 2021)

samples, and that morphologically resembles *Coniochaeta hoffmannii*, an opportunistic pathogen

of humans, dogs and cattle (Sakaeyama et al. 2007, Hurst 2019). Some *Coniochaeta* species,

including close relatives of *C. discospora*, are also known as opportunistic wood and tree

pathogens in Prunus and other hosts (Damm et al. 2010), possibly connecting the fungal

communities identified here with the riparian habitats they were isolated from. Most *Preussia*

species are coprophilous (Ahmed and Cain 1972), while *Gymnoascus* spp. are within the

Onygenales (which include many taxa known to degrade animal protein, including *Coccidioides*

spp.), produce keratinase and have been isolated from hair and wool (Błyskal 2009).

Myxotrichum spp., also within the Onygenales, are soil saprotrophs that are mostly associated

with breaking down plant cellulose, though also produce enzymes that can degrade animal tissue

(Rice 2005). Our data do not establish a causal relationship, but may indicate an association

between *Coccidioides* positive samples and actively inhabited rodent burrows, concurring with

research done in the Carrizo Plain of California (Head et al. 2024).

<u>*Coccidioides* biogeography:</u>

Landscape models predict *Coccidioides* distribution based on soil, climate, and case data (Gorris

et al. 2019, Weaver et al. 2020, Dobos et al. 2021). However, no current landscape modeling

efforts take into consideration environmental detections or rodent host ecology. Our findings of

sporadic detection in riparian soils, and our previous finding of absence in agricultural soils

(Wagner et al. 2023) refine understanding of habitat suitability and suggest that water availability

and local environmental conditions may indirectly influence *Coccidioides* presence. Namely, by

suggesting that nearby water resources can influence the distribution of *Coccidioides* in rodent

burrow soils through the provision of ecosystem resources, either directly, or by supporting local

rodent populations (Falck et al. 2003, Hamilton et al. 2015). Indeed, early work isolated

Coccidioides from sites in the SJV along both the Fresno and Chowchilla Rivers in Merced and

Madera counties (Lacy and Swatek 1974). Taking the geography of riparian corridors into

account could influence future modeling efforts regarding the predicted landscape-level

distribution of *Coccidioides* in the environment and add important constraints to modeling efforts

that rely primarily on coccidioidomycosis incidence data. The inclusion of species distribution

modelling, focused on likely rodent reservoirs, can further improve such modelling approaches.

<u>Temporal dynamics and dispersal</u>

495 No clear temporal trends were detected in rodent burrow soil samples at University Place and

496 Panorama Park West, which accounted for most positive Kern River samples. Limited sampling

497 intervals, presence-absence data, and uneven spatial distribution constrain inferences about

498 seasonality or growth dynamics. Our finding a significant association between *Coccidioides*

499 presence and sampling timepoint across all Kern River samples may indicate unmeasured

500 categorical differences between sites where *Coccidioides* is heavily established, and sites where

501 *Coccidioides* may be limited or more recently established.

503 *Coccidioides* appears to be distributed non-uniformly across urban riparian landscapes,

504 consistent with prior studies (Maddy 1965, Lacy and Swatek 1974, Greene et al. 2000).

505 Detection at only a subset of sites with similar environmental features (numerous rodent

506 burrows, observably large populations of *O. beecheyi*, irrigated lawns, proximity to the Kern

507 River and close contact with humans) highlights unmeasured variables likely influencing

508 distribution. Patterns may be influenced by rodent populations or localized microhabitat

509 conditions, though these cannot be confirmed here. The distribution of plague, another microbial,

510 zoonotic disease in California associated with *O. beecheyi* (Eskey and Haas 1940, Lang 2004), is

511 better explained by rodent host populations than by climatic niche data (Fell et al. 2022). The

512 possibility that *Coccidioides* has a non-uniform geographic distribution associated with native

513 rodents, not unlike distribution patterns found in plague, suggests improvements for future

514 landscape-level modelling efforts.

516 <u>Additional methodological limitations</u>

Although we obtained 1178 samples, sample size and the limit of detection remain the two

biggest limitations of our approach (Wagner et al. 2022, 2023). Thus, our inability to detect

Coccidioides at certain sites does not necessarily preclude the presence of *Coccidioides* at those

sites. This limitation is ever greater for settled dust owing to the far smaller size of settled dust

samples compared to soil samples. DNA can come from living or dead fungi and may confound

our interpretation of temporal dynamics. We know that there is a lower limit of detection for the

CocciENV qPCR assay (Bowers et al. 2019), so *Coccidioides* could be present in samples that

we scored as absent (Dineen et al. 2010). Additionally, consideration must be given to unknown

environmental influences that may affect our ability to detect *Coccidioides* in such samples.

Studies of total bacterial and fungal community dynamics at continental or regional scales

(Barberán et al. 2015) have successfully used approaches similar to ours (Wagner et al. 2023),

but understanding the relationships between *Coccidioides* and individual microbial species may

benefit from a targeted approach that investigates a more limited number of fungal taxa. Due to

finite sampling resources, nearly all our samples are from within rodent burrows, as this was the

most straight-forward way to determine if *Coccidioides* was in an area. However, this approach

is unable to address questions regarding *Coccidioides* in other environments. Because

Coccidioides occurs patchily in soil, we prioritized broad spatial coverage across ecological

gradients, complemented by more intensive resampling at sites where detections were frequent.

This design provided both regional context and local replication for temporal analyses. While our

effort exceeds the sampling scale of most prior environmental surveys (S1 Table), we recognize

that further sampling would help quantify detection variability and additional balanced sampling

across regions will be needed to confirm observed trends. Although rodent burrows were a focal

sampling microhabitat, we did not formally collect data on rodent abundance, activity, or species

composition. This study was framed from a microbial ecology perspective to characterize

environmental *Coccidioides* distribution patterns. However, the close ecological association

between *Coccidioides* and rodent hosts suggests that integrating host ecology will be essential

for defining mechanistic interactions. We observed a notably high number of *O. beecheyi* at

several Kern River sites, especially at University Place and Panorama Park West. Dozens were

regularly seen during fieldwork, often retreating into the same burrows we sampled. *O. beecheyi*

have been associated with *Coccidioides* presence in Mojave Desert soils (Lauer et al. 2020), and

related rodents have tested positive for *Coccidioides* in lung tissue (Emmons et al. 1942, Salazar-

Hamm 2021). Many sampled burrows showed small mammal footprints consistent with active

habitation (Grinnell 1923, McNew et al. 2023). While our observations suggest a potential link

between *O. beecheyi* behavior and *Coccidioides* distribution, it is important to emphasize that

future testing of *O. beecheyi* tissues (Baptista-Rosas et al. 2012, Salazar-Hamm 2021) is needed

to confirm of refute their potential role in *Coccidioides* ecology. Our investigation of riparian

zones in urban population centers means that our samples largely came from semi-natural areas

situated in urban centers, making land classification challenging. Finally, our use of 9km^2 grid

scale soil moisture data, derived from satellite measurements, limits the conclusions we can draw

regarding comparisons between *Coccidioides*, the soil fungal community and soil moisture

levels. Future studies should pair direct soil moisture measurements with molecular surveys for

Coccidioides and other fungal species.

Conclusion

Our findings, based on an analysis of 1178 samples, revealed that *Coccidioides* is abundant,

though unevenly distributed, in soils along two waterways that cross major urban population

centers in the SJV, a region high in coccidioidomycosis incidence. The distribution pattern of

Coccidioides showcases a clear elevation-dependent dynamic, suggesting that elevation,

environmental variables that correlate with elevation, or both, play a pivotal role in shaping

Coccidioides biogeography. Our samples were primarily obtained from rodent burrows, and

possible constraints on transmission among rodent groups are suggested by the proximity of

Coccidioides positive and negative soils at single sites throughout the year. Our investigation into

the relationship between *Coccidioides* and the broader soil fungal community yielded consistent

results with our previous study along Highway 33, with both providing no evidence for strong,

community-level associations, challenging assumptions about the activity of *Coccidioides*

outside of animal hosts. However, our identification of a handful of individual fungal taxa that

correlated with *Coccidioides* presence, most of which live on animal derived substrates, provides

evidence that *Coccidioides* presence may be associated with rodent activity. Finally, the

identification of riparian corridors as potential *Coccidioides* habitats emphasizes the need for

nuanced geographical considerations in predictive models, taking into consideration both

landscape topography, landscape use, and distributions of rodent hosts.

Our detection of *Coccidioides* in native rodent burrows situated within and adjacent to major

population centers in the SJV underscores a previously underappreciated risk to public health.

The highest incidence of Coccidioidomycosis in California is in Kern County (Nguyen et al.

2019), nearly half the population of which resides in the city of Bakersfield, where we found

abundant *Coccidioides*. Likewise, Coalinga is the largest population center in western Fresno

County, where we also regularly found *Coccidioides*. Waterways, likely home to potential

Coccidioides rodent hosts, bisect or run adjacent to many other population centers in the SJV.

These findings advocate for similar surveys in population centers elsewhere in the SJV and for

targeted mitigation strategies that prioritize the identification and non-disturbance of rodent

burrows and nearby soils. Where disturbance is unavoidable, stringent adherence to personal

protective protocols is essential (Das et al. 2012). Importantly, efforts to exterminate native

rodents may prove counterproductive; the resultant increase in carcasses could enhance fungal

growth and sporulation, thereby exacerbating the incidence of Valley fever in the area. Similarly,

the effectiveness of trapping and removing rodents is uncertain given the extensive reservoir of

susceptible rodents in cities and surrounding peri-urban landscapes. Preventing access to

discarded human food (and access to other human-derived food resources, i.e. irrigated lawns)

may provide a more sustainable and long-term option for limiting potential *Coccidioides* rodent

host populations. Collectively, our results highlight the need for integrated management

strategies that address both the ecological complexity of *Coccidioides* and public health

imperatives.

Materials and Methods

We sampled soil at 59 individual sites in and around the cities of Bakersfield and Coalinga in

California's SJV (Fig 1), with the intent of obtaining latitudinal transects describing *Coccidioides*

presence across both population centers. Because they are accessible, create natural

uninterrupted transects across landscapes and support native rodent populations whose burrows

are associated with *Coccidioides*, sampling occurred primarily along two seasonally ephemeral

waterways, the Kern River in Kern County, and Los Gatos Creek in Fresno County. The Kern

River originates in the Sierra Nevada mountains in Sequoia National Park (United States

Geological Survey 2021a), travels south to a reservoir, Lake Isabella, and then west through

Bakersfield before terminating at the California Aqueduct near the Buena Vista Aquatic

Recreational Area in the southern SJV. Los Gatos Creek begins in the Clear Creek Management

Area in California's Diablo Range (United States Geological Survey 2021b), flows east through

Coalinga and terminates at the California Aqueduct near Huron, approximately 130km north of

the Kern River terminus. Soils in the vicinity of our study sites along both waterways are aridic

and receive little annual rainfall. Wild vegetation along Los Gatos Creek is dominated by tree

species of *Quercus*, *Acer* and *Pinus* at higher elevations and at lower elevations by grass/shrub

species of *Nassella, Sporobolus,* and *Suaeda nigra* (Griffith et al. 2016). Wild vegetation along

the Kern River is classified in three ecoregions: one characterized by *Allenrolfea occidentalis*,

Atriplex spp., *Nassella spp*. and *Prosopis spp*. from Taft to Bakersfield; a second characterized

by *Atriplex spp*., *Nassella* spp., and some *Quercus douglasii* and *Opuntia basilaris* from

Bakersfield to the Sierra foothills; and a third characterized by *Adenostoma fasciculatum*,

Ceanothus spp., *Nassella spp.*, *Pinus sabiniana* and *Quercus spp.* found east from the Sierra

foothills along an increasing elevation gradient to Lake Isabella (Griffith et al. 2016).

Sampling and DNA Extraction: We collected soil from 625 rodent burrows at 35 sites along the

Kern River and from an additional 205 rodent burrows at seven sites away from waterways (non-

Kern River sites) in and around Bakersfield and Taft, California. In Coalinga we collected soils

from 228 rodent burrows and 20 surface soils (directly outside of burrows) at 16 sites along Los

Gatos Creek, and an additional 100 settled dust samples within the city of Coalinga. Our

sampling effort was limited to accessible public land (parks, roadsides and waterways), and

presents one of the most extensive such efforts among known studies (S1 Table). Coordinates

were obtained using cellular phone GPS data cross-referenced with satellite imagery, and

elevation data were estimated using elevatr version 0.4.2 (Hollister et al. 2022). All soils were

collected as deeply as possible from within rodent burrows using steel hemispherical collectors

attached to 30cm threaded rods, except for 20 soil samples from one site (Highway 33 West)

along Los Gatos Creek that were collected from surface soil located adjacent to entrances of

simultaneously sampled rodent burrows. For all soil samples, ≥25ml of soil was collected in

conical 50ml polypropylene centrifuge tubes and mixed by repeatedly inverting the tubes.

Sampling effort across sites is summarized in S2 Table.

We resampled a subset of locations to explore potential changes in *Coccidioides* presence over

time (S3 Table). Previous research indicates that *Coccidioides* persists in rodent burrows or

adjacent soils at the same location (Vargas-Gastelum et al. 2015, Wagner et al. 2023, Head et al.

2024). Along the Kern River, we sampled five sites (Kern River Parkway West End, University

Place, Rio Kern Park, Yokuts Park, and Panorama Park West) at four time points over one year

(April 2021, July 2021, October 2021 and January 2022). Where sites were sampled more than

once, specific rodent burrows were not necessarily resampled. To investigate *Coccidioides* in

soils unassociated with the Kern River (non-Kern River sites), we sampled four areas along

California Highway 119, 10km south of the Kern River, at one time point each (Buena Vista

Recreation Area, Buena Vista Valley, California State University Bakersfield (CSUB) Campus,

and Sports Village). Along Los Gatos Creek, we sampled from the Clear Creek Management

Area, through Coalinga, CA, to just north of Pleasant Valley State Prison near its confluence with

Jacalitos Creek. Only one site was sampled more than once along Los Gatos Creek, Highway 33

West, in August 2020 and January 2022. In January 2022, we sampled east of California

Highway 33 to the confluence of Los Gatos Creek with Warthan Creek and three sites near the

confluence of Los Gatos Creek with Jacalitos Creek. In April 2024, we sampled ten sites along

Los Gatos Creek Road along an elevational gradient from the Clear Creek Management Area in

the Diablo Range, downhill to Coalinga, CA. In addition to soil samples, settled dust was

sampled from 100 urban surfaces protected from precipitation by architectural features within

Coalinga, CA, in August 2020. Settled dust was sampled by swabbing horizontal public surfaces

using sterile, DNA-free cotton swabs (Puritan, Guilford, ME, USA). DNA was extracted from

0.25g of soil from each collection tube, and from swab tips cut from swab sticks. With both soil

and swab tips, the samples were placed in MoBio Powersoil DNA kit buffer C1 (MoBio,

Carlsbad, CA, USA) and disrupted in a FastPrep-24 5G bead beater (MP Biomedicals, Santa

Ana, CA, USA), followed by DNA recovery using the MoBio Powersoil DNA kit as described

(Wagner et al. 2022, 2023). DNA was diluted to 5ng$\cdot$µl^{-1} following measurement with the Qubit

dsDNA HS Assay kit (Life Technologies Inc., Gaithersburg, MD, USA).

Coccidioides detection: All samples were tested for _Coccidioides_ using the CocciEnv qPCR

assay (Bowers et al. 2019) in quadruplicate reactions, using nuclease-free water as a negative

control and _C. posadasii_ strain Silveira DNA as a positive control (provided by the lab of Anita

Sil at the University of California San Francisco). The CocciENV assay builds upon the

previously developed CocciDx assay (Litvintseva et al. 2015, Saubolle et al. 2018) by expanding

the primer set targeting a unique repeating transposon sequence in the _Coccidioides_ genome

(NCBI BioProject PRJNA46299) and having been comprehensively validated using

environmental soil samples (Bowers et al. 2019). Template DNA, 2µl (50ng$\cdot$µl^{-1}), was added to

10µl of TaqMan Environmental Master Mix2.0 (Applied Biosystems, Waltham, MA, USA), 2µl

CocciEnv primer mix as described in Bowers et al. (2019), and 6µl nuclease-free H_2O for a total

of 20µl per reaction. The qPCR assay was performed on 96-well plates on the Stratagene

Mx3000P platform (Agilent Technologies, Santa Clara, CA, USA) with the following cycling

conditions: 1 cycle at 95°C for 10 minutes, 40 cycles at 95°C for 15 seconds each and 1 cycle at

60°C for 1 minute. Reactions with positive detections were identified at cycle threshold (CT) <

40, a minimum relative fluorescence unit value of 1000, and logarithmic amplification as

described (Wagner et al. 2022, 2023). We were confident that an individual sample contained

Coccidioides if 3 or more of its wells scored positive in the qPCR assay (Wagner et al. 2022,

2023). This threshold was chosen to balance sensitivity and specificity to minimize false

positives while reliably detecting low-level presence. We also report the total number of positive

wells for each sample (Fig 1, 2) (S2 Table, S3 Table). Settled dust recovered by swabbing

provided a much smaller amount of sample compared to sampled soil, making it difficult to

equate a failure to detect *Coccidioides* in settled dust with a failure to detect it in soil (Wagner et

al. 2023). For this reason, following the protocol we established in Wagner et al. (2023), for

settled dust samples having ≥1 positive well, multiple DNA extractions were concentrated using

a vacuum oven operating at 20kPa and 65°C for one hour or until complete volume evaporation.

Following resuspension of the concentrated DNA in nuclease-free H_2O, a second analysis using

the CocciEnv assay in quadruplicate reactions was done, and our initial analyses were updated

for any samples that now showed *Coccidioides* detection in ≥ 3 replicate wells. All initial

processing of qPCR data and generation of amplification curves used Mxpro version 4.1 (Agilent

Technologies, Santa Clara, CA, USA).

<u>Fungal Community Amplicon Sequencing</u>: To characterize the fungal soil community with

regard to *Coccidioides* detection, amplicon sequencing of the internal transcribed (ITS2) region

of fungal rDNA was performed on DNA extracted from 68 rodent burrow soil samples, 28 from

four Kern River sites and 40 from one Los Gatos Creek site (20 within and 20 just outside the

entrance of rodent burrows), all of which were also tested for *Coccidioides* using the CocciENV

qPCR assay. An effort was made to perform sequencing on an equal number of *Coccidioides*

positive and negative soils, as well as across an equal number of soils from two timepoints (April

2021 and July 2021) for Kern River samples and from two burrow locations (burrow interior and

burrow entrance) for Los Gatos Creek soils (S4 Table). As previously described (Gao et al. 2018,

Wagner et al. 2022), the ITS2 region was PCR amplified from extracted DNA using the 5.8SFun

(AACTTTYRRCAAYGGATCWCT) and ITS4Fun (AGCCTCCGCTTATTGATATGCTTAART)

primers (Taylor et al. 2016), with the AccuStart II PCR SuperMix kit (Quantabio, Beverly, MA,

USA), on the Gene Amplification PCR System (Bio-Rad Laboratories, Hercules, CA, USA). The

reaction mixture consisted of 2μl of template DNA, 2.5μl of 50μM forward and reverse primer

each, 12.5μl AccuStart II PCR SuperMix, 2.5μl of nuclease-free water, and 3μl BSA, with the

following thermal cycling conditions: an initial step of 96°C for 2 minutes, followed by 35

cycles of 94°C for 30 seconds, 58°C for 40 seconds and 72°C for 2 minutes, with a final step of

72°C for 10 minutes. The PCR product was quantified using the Qubit dsDNA HS Assay kit

(Life Technologies Inc., Gaithersburg, MD, USA) and then sent to the QB3 Vincent J. Coates

Genomics Sequencing Laboratory (University of California, Berkeley, CA, USA), where the

samples were assigned unique dual indices to prevent barcode tag-jumping (Zinger et al. 2019,

Carøe and Bohmann 2020) and were sequenced using paired-end PE300 chemistry on the MiSeq

platform (Illumina, Inc., CA, USA).

<u>Amplicon Sequencing Data Processing:</u> All sequence data were processed using Qiime 2 version

2019.10.0 (Bolyen et al. 2019). Quality control of the sequencing runs was performed manually

followed by denoising using DADA2 (Callahan et al. 2016). Paired-end reads were then joined,

end bases with a quality score lower than 25 were trimmed, and unpaired reads were removed.

Assignment of operational taxonomic units (OTUs) to binned amplicon sequence variants

(ASVs) was done with a naïve Bayes classifier, trained with the UNITE developer (untrimmed)

database Qiime release version 9.0 (29.11.2022) clustered at 99% similarity (Pedregosa et al.

2011, Bokulich et al. 2018, Bolyen et al. 2019, Abarenkov et al. 2022). Sequences not assigned

to specific taxonomies in the UNITE database were excluded in all downstream analyses. All

supporting sequence data and metadata have been deposited in the NCBI Sequence Read Archive

with BioProject numbers PRJNA1099691 and PRJNA1099703. All code needed to processing

sequencing data and to generate taxonomic tables is included as a supplementary file (Code S1).

<u>Soil Moisture Data</u>: To explore how the presence of *Coccidioides* in soils may coincide with

precipitation and soil water availability (Kolivras and Comrie 2003, Comrie 2005, Tamerius and

Comrie 2011), satellite-derived soil moisture data were acquired. Using the NASA AppEEARS

user interface version 3.3.1 (AppEEARS Team 2022) (https://appeears.earthdatacloud.nasa.gov),

the soil moisture active-passive (SMAP) L3 radar/radiometer global EASE-grid soil moisture

dataset version 3 was downloaded, which is gridded at a 9km resolution using the EASE-Grid

2.0 projection, at a daily resolution (Entekhabi et al. 2016). SMAP L3 version 3 applies active–

passive downscaling, model-based enhancements for vegetated or complex terrain and pixel

overlap during satellite passes, all of which can introduce spatial detail at effective resolutions

below 9km, depending on location. Surface (0-5cm) and rootzone (0-100cm) soil moisture data

(cm^3 water / cm^3 soil) were selected using the "point sample tool" in the AppEEARS user

interface at coordinates, both continuously along Lost Gatos Creek and the Kern River (surface

only), as well as corresponding to soil sampling locations (both surface and rootzone), and mean

values for each sampling month were calculated across sites. Due to the limited number of sites,

and geographic range of sites along Los Gatos Creek, comparisons of site-level soil moisture

data were limited to Bakersfield area sites, encompassing locations both directly along the Kern

River and non-Kern River sites.

<u>Statistical Analysis</u>:

All statistical analyses used R version 4.1.0 (R Core Team 2020), vegan version 2.5.7 (Oksanen

et al. 2019) and lme4 1.1.33 (Bates et al. 2015). Two separate logistic regression models were

employed to determine significant differences in *Coccidioides* detection between sites, sampling

timepoints and across remotely sensed soil moisture levels along the Kern River and surrounding

areas. In the first logistic regression model, differences in *Coccidioides* detection were

investigated as a function of remotely sensed soil moisture across all sites along the Kern River

and the surrounding area. Sampling timepoint and elevation were included as fixed effects and,

given the uneven sampling distribution (and dominant nature of Panorama Park West and

University Place), site was included as a random effect to account for site-to-site variability.

Surface and rootzone soil moisture and elevation were normalized by transforming each

predictor into z-scores due to the large range in absolute values for each. In the second logistic

regression model, differences in *Coccidioides* detection were tested between samples collected at

two sites, Panorama Park West and University Place, which were sampled at all four time

periods, and which had a high number of positive *Coccidioides* samples relative to other sites

along the Kern River. Both sampling timepoint and remotely sensed soil moisture were included

as covariates and z-score normalization was not performed as values between these two sites

were relatively similar.

Given differences in the location and timing of sampling, patterns in soil fungal community

structure were analyzed separately for Kern River and Los Gatos Creek sites. Species-level

taxonomic count data was Wisconsin double transformed (Bray and Curtis 1957, Legendre and

Legendre 2012) following the initial removal of species with only a single count across all

samples, and Bray-Curtis dissimilarity was calculated as a measure of community β-diversity. To

determine the correlation between fungal community β-diversity and *Coccidioides* detection, we

employed a permutational (1000 permutations) multivariate analysis of variance

(PERMANOVA) test, as implemented in the "adonis" function (Anderson 2001). Kern River

samples included covariates for sampling site and timepoint. As the entire subset of Los Gatos

Creek samples that underwent amplicon sequencing was collected from the same site, on the

same day, only the specific sampling location (burrow interior versus burrow entrance) was

included as a covariate. At Los Gatos Creek, soil moisture was excluded from PERMANOVA

analysis due to the limited spatial extent of the subset of samples that was sequenced. We

investigated possible co-occurrences between *Coccidioides* and other fungal taxa, extending our

previous investigation of rodent burrow soils along California highway 33 (Wagner et al. 2023).

An indicator species analysis was employed on untransformed species data using indicspecies

version 1.7.12 (Dufrêne and Legendre 1997, De Cáceres et al. 2010) and significant results ($p <$

0.05, 1000 permutations) with $\geq$ 100 reads were cross-referenced using BLASTN

(https://blast.ncbi.nlm.nih.gov) and Mycobank (https://www.mycobank.org) to determine likely

alternative species identities (> 96% identity).

Figures were made with ggplot2 version 3.3.5 (Kahle and Wickham 2013), Complexheatmap

version 2.8.0 (Gu et al. 2016), ggmap version 3.0.0 (Kahle and Wickham 2013) and osmdata

version 0.1.8 (Padgham et al. 2017). Map data (lines) were acquired from OpenStreetMap

(www.openstreetmap.org) under the Open Data Commons Open Database License (OdbL). Map

tiles are by Stamen Design (www.stamen.com) under Creative Commons By Attribution (CC BY

4.0). California shapefile is public domain data by Natural Earth (www.naturalearthdata.com).

Satellite imagery is public domain data provided by the United States Department of Agriculture

(USDA) National Agriculture Imagery Program (NAIP) (https://naip-

usdaonline.hub.arcgis.com). All non-sequencing data used in this study and all code needed to

replicate statistical analyses are included as supplementary files (Data S1, S2; Code S2).

Acknowledgements: This work was supported by National Institutes of Health, National

Institute of Allergy and Infectious Disease grant R01AI148336, University of California Office

of the President grant VFR-19-633952, and Department of Energy grant DE-SC0014081. We

would like to thank Anita Sil's lab for providing positive control *Coccidioides* DNA. The authors

have no conflict of interest.

References

Abarenkov, K., A. Zirk, T. Piirmann, R. Pöhönen, F. Ivanov, R. H. Nilsson, and U. Kõljalg. 2022.

UNITE QIIME release for Fungi. UNITE Community.

Ahmed, S. I., and R. Cain. 1972. Revision of the genera Sporormia and Sporormiella. Canadian

journal of Botany 50:419–477.

Anderson, M. J. 2001. A new method for non-parametric multivariate analysis of variance.

Austral ecology 26:32–46.

AppEEARS Team. 2022. Application for Extracting and Exploring Analysis Ready Samples

(AppEEARS). Ver. 3.3.1. NASA EOSDIS Land Processes Distributed Active Archive Center

(LP DAAC), USGS/Earth Resources Observation and Science (EROS) Center, Sioux Falls,

South Dakota, USA.

Baldwin, R. A., R. Meinerz, and G. W. Witmer. 2016. Are Cholecalciferol plus Anticoagulant

Rodenticides a Viable Option for Field Rodents? Page Proceedings of the Vertebrate Pest

Conference.

Baptista-Rosas, R. C., J. Catalan-Dibene, A. L. Romero-Olivares, A. Hinojosa, T. Cavazos, and M.

Riquelme. 2012. Molecular detection of Coccidioides spp. from environmental samples

in Baja California: linking Valley Fever to soil and climate conditions. fungal ecology

5:177–190.

Barberán, A., J. Ladau, J. W. Leff, K. S. Pollard, H. L. Menninger, R. R. Dunn, and N. Fierer. 2015.

Continental-scale distributions of dust-associated bacteria and fungi. Proceedings of the

National Academy of Sciences 112:5756–5761.

Bates, D., M. Mächler, B. Bolker, and S. Walker. 2015. Fitting Linear Mixed-Effects Models Using

lme4. Journal of Statistical Software 67:1–48.

Best, T. L. 1993. Perognathus inornatus. Mammalian Species:1–5.

Błyskal, B. 2009. Fungi utilizing keratinous substrates. International Biodeterioration &

Biodegradation 63:631–653.

Bokulich, N. A., B. D. Kaehler, J. R. Rideout, M. Dillon, E. Bolyen, R. Knight, G. A. Huttley, and J.

Gregory Caporaso. 2018. Optimizing taxonomic classification of marker-gene amplicon

sequences with QIIME 2's q2-feature-classifier plugin. Microbiome 6:90.

Bolyen, E., J. R. Rideout, M. R. Dillon, N. A. Bokulich, C. C. Abnet, G. A. Al-Ghalith, H. Alexander,

E. J. Alm, M. Arumugam, F. Asnicar, Y. Bai, J. E. Bisanz, K. Bittinger, A. Brejnrod, C. J.

Brislawn, C. T. Brown, B. J. Callahan, A. M. Caraballo-Rodríguez, J. Chase, E. K. Cope, R.

Da Silva, C. Diener, P. C. Dorrestein, G. M. Douglas, D. M. Durall, C. Duvallet, C. F.

Edwardson, M. Ernst, M. Estaki, J. Fouquier, J. M. Gauglitz, S. M. Gibbons, D. L. Gibson, A.

Gonzalez, K. Gorlick, J. Guo, B. Hillmann, S. Holmes, H. Holste, C. Huttenhower, G. A.

Huttley, S. Janssen, A. K. Jarmusch, L. Jiang, B. D. Kaehler, K. B. Kang, C. R. Keefe, P. Keim,

S. T. Kelley, D. Knights, I. Koester, T. Kosciolek, J. Kreps, M. G. I. Langille, J. Lee, R. Ley, Y.-X.

Liu, E. Loftfield, C. Lozupone, M. Maher, C. Marotz, B. D. Martin, D. McDonald, L. J.

McIver, A. V. Melnik, J. L. Metcalf, S. C. Morgan, J. T. Morton, A. T. Naimey, J. A. Navas-

Molina, L. F. Nothias, S. B. Orchanian, T. Pearson, S. L. Peoples, D. Petras, M. L. Preuss, E.

Pruesse, L. B. Rasmussen, A. Rivers, M. S. Robeson, P. Rosenthal, N. Segata, M. Shaffer, A.

Shiffer, R. Sinha, S. J. Song, J. R. Spear, A. D. Swafford, L. R. Thompson, P. J. Torres, P.

Trinh, A. Tripathi, P. J. Turnbaugh, S. Ul-Hasan, J. J. J. van der Hooft, F. Vargas, Y. Vázquez-

Baeza, E. Vogtmann, M. von Hippel, W. Walters, Y. Wan, M. Wang, J. Warren, K. C.

Weber, C. H. D. Williamson, A. D. Willis, Z. Z. Xu, J. R. Zaneveld, Y. Zhang, Q. Zhu, R.

Knight, and J. G. Caporaso. 2019. Reproducible, interactive, scalable and extensible

microbiome data science using QIIME 2. Nature Biotechnology 37:852–857.

Bowers, J., K. Parise, E. Kelley, D. Lemmer, J. Schupp, E. Driebe, D. Engelthaler, P. Keim, and B.

Barker. 2019. Direct detection of *Coccidioides* from Arizona soils using CocciENV, a highly

sensitive and specific real-time PCR assay. Medical mycology 57:246–255.

Bray, J. R., and J. T. Curtis. 1957. An Ordination of the Upland Forest Communities of Southern

Wisconsin. Ecological Monographs 27:325–349.

Cain, R. F. 1957. Studies of coprophilous Ascomycetes: VI. Species from the Hudson Bay area.

Canadian Journal of Botany 35:255–268.

Callahan, B. J., P. J. McMurdie, M. J. Rosen, A. W. Han, A. J. A. Johnson, and S. P. Holmes. 2016.

DADA2: High-resolution sample inference from Illumina amplicon data. Nature Methods

13:581–583.

Carøe, C., and K. Bohmann. 2020. Tagsteady: a metabarcoding library preparation protocol to

avoid false assignment of sequences to samples. Molecular Ecology Resources 20:1620–

1631.

Case, R. L., and J. B. Kauffman. 1997. Wild ungulate influences on the recovery of willows, black

cottonwood and thin-leaf alder following cessation of cattle grazing in northeastern

Oregon. Northwest science. 71:115–126.

Catalán-Dibene, J., S. M. Johnson, R. Eaton, A. L. Romero-Olivares, R. C. Baptista-Rosas, D.

Pappagianis, and M. Riquelme. 2014. Detection of coccidioidal antibodies in serum of a

small rodent community in Baja California, Mexico. Fungal biology 118:330–339.

Centers for Disease Control and Prevention. 2018. Valley fever (Coccidioidomycosis) Statistics.

https://www.cdc.gov/fungal/diseases/coccidioidomycosis/statistics.html.

Chow, N. A., D. Kangiser, L. Gade, O. Z. McCotter, S. Hurst, A. Salamone, R. Wohrle, W. Clifford, S.

Kim, Z. Salah, and others. 2021. Factors influencing distribution of Coccidioides immitis

in soil, Washington State, 2016. Msphere 6:e00598-21.

Comrie, A. C. 2005. Climate factors influencing coccidioidomycosis seasonality and outbreaks.

Environmental health perspectives 113:688–692.

Cordeiro, R., R. Brilhante, M. Rocha, M. Fechine, A. Costa, Z. Camargo, and J. Sidrim. 2006. In

vitro activities of caspofungin, amphotericin B and azoles against *Coccidioides posadasii*

strains from Northeast, Brazil. Mycopathologia 161:21–26.

Damm, U., P. Fourie, and P. W. Crous. 2010. Coniochaeta (Lecythophora), Collophora gen. nov.

and Phaeomoniella species associated with wood necroses of Prunus trees. Persoonia-

Molecular Phylogeny and Evolution of Fungi 24:60–80.

Das, R., J. McNary, K. Fitzsimmons, D. Dobraca, K. Cummings, J. Mohle-Boetani, C. Wheeler, A.

McDowell, Y. Iossifova, R. Bailey, K. Kreiss, and B. Materna. 2012. Occupational

Coccidioidomycosis in California: Outbreak Investigation, Respirator Recommendations,

and Surveillance Findings. Journal of Occupational & Environmental Medicine 54:564–

571.

De Cáceres, M., P. Legendre, and M. Moretti. 2010. Improving indicator species analysis by

899   combining groups of sites. Oikos 119:1674–1684.

900   De Macêdo, R. C., A. S. Rosado, F. F. Da Mota, M. A. Cavalcante, K. D. Eulálio, L. Martins, M. S.

901   Lazéra, B. Wanke, and others. 2011. Molecular identification of Coccidioides spp. in soil

902   samples from Brazil. BMC microbiology 11:1–9.

903   Deveautour, C., J. Chieppa, U. N. Nielsen, M. M. Boer, C. Mitchell, S. Horn, S. A. Power, A.

904   Guillen, A. E. Bennett, and J. R. Powell. 2020. Biogeography of arbuscular mycorrhizal

905   fungal spore traits along an aridity gradient, and responses to experimental rainfall

906   manipulation. Fungal ecology 46:100899.

907   Dineen, S., R. Aranda IV, D. Anders, and J. Robertson. 2010. An evaluation of commercial DNA

908   extraction kits for the isolation of bacterial spore DNA from soil. Journal of applied

909   microbiology 109:1886–1896.

910   Dobos, R. R., K. Benedict, B. R. Jackson, and O. Z. McCotter. 2021. Using soil survey data to

911   model potential Coccidioides soil habitat and inform Valley fever epidemiology. PloS one

912   16:e0247263.

913   Dufrêne, M., and P. Legendre. 1997. Species assemblages and indicator species: the need for a

914   flexible asymmetrical approach. Ecological monographs 67:345–366.

915   Egeberg, R., and A. F. Ely. 1956. Coccidioides immitis in the soil of the southern San Joaquin

916   Valley. American Journal of Medical Sciences 231:151–4.

917   Elconin, A. F., M. Egeberg, and R. Lubarsky. 1957. Growth Patterns of Coccidioides immitis. Page

918   168 Proceedings of Symposium on Coccidioidomycosis: Held at Phoenix, Ariz.-Feb. 11-

919   13, 1957. US Department of Health, Education, and Welfare Public Health Service ….

Emmons, C., L. Ashburn, and others. 1942. The isolation of Haplosporangium parvum n. sp. and

Coccidioides immitis from wild rodents. Their relationship to coccidioidomycosis. Public

Health Rep 57:1715–1727.

Emmons, C. W. 1942. Isolation of *Coccidioides* from soil and rodents. Public Health Reports

(1896-1970) 57:109–111.

Engelthaler, D. M., C. C. Roe, C. M. Hepp, M. Teixeira, E. M. Driebe, J. M. Schupp, L. Gade, V.

Waddell, K. Komatsu, E. Arathoon, and others. 2016. Local population structure and

patterns of Western Hemisphere dispersal for Coccidioides spp., the fungal cause of

Valley Fever. MBio 7:10–1128.

Entekhabi, D., N. Das, E. Njoku, J. Johnson, and J. Shi. 2016. SMAP L3 radar/radiometer global

daily 9 km EASE-grid soil moisture, version 3. NASA National Snow and Ice Data Center

Distributed Active Archive Center.

Eskey, C., and V. Haas. 1940. Plague in the western part of the United States, US Public Health

bulletin 254. US Government Printing Of fi ce, Washington, DC.

Falck, M. J., K. R. Wilson, and D. C. Andersen. 2003. Small mammals within riparian habitats of a

regulated and unregulated aridland river. Western North American Naturalist:35–42.

Fell, H. G., O. G. Osborne, M. D. Jones, S. Atkinson, S. Tarr, S. H. Keddie, and A. C. Algar. 2022.

Biotic factors limit the invasion of the plague pathogen (Yersinia pestis) in novel

geographical settings. Global Ecology and Biogeography 31:672–684.

Fisher, M. C., G. L. Koenig, T. J. White, G. San-Blas, R. Negroni, I. G. Alvarez, B. Wanke, and J. W.

Taylor. 2001. Biogeographic range expansion into South America by *Coccidioides immitis*

mirrors New World patterns of human migration. Proceedings of the National Academy

of Sciences 98:4558–4562.

Free, C., G. Baxter, C. Dickman, A. Lisle, and L.-P. Leung. 2015. Diversity and community

composition of vertebrates in desert river habitats. PLoS One 10:e0144258.

Gao, C., L. Montoya, L. Xu, M. Madera, J. Hollingsworth, E. Purdom, R. B. Hutmacher, J. A.

Dahlberg, D. Coleman-Derr, P. G. Lemaux, and others. 2018. Strong succession in

arbuscular mycorrhizal fungal communities. The ISME journal 13:214–226.

Gorris, M., L. Cat, C. Zender, K. Treseder, and J. Randerson. 2018. Coccidioidomycosis dynamics

in relation to climate in the southwestern United States. GeoHealth 2:6–24.

Gorris, M. E., J. E. Neumann, P. L. Kinney, M. Sheahan, and M. C. Sarofim. 2021. Economic

valuation of coccidioidomycosis (Valley fever) projections in the United States in

response to climate change. Weather, Climate, and Society 13:107–123.

Gorris, M. E., K. K. Treseder, C. S. Zender, and J. T. Randerson. 2019. Expansion of

Coccidioidomycosis Endemic Regions in the United States in Response to Climate

Change. GeoHealth 3:308–327.

Greene, D. R., G. Koenig, M. C. Fisher, and J. W. Taylor. 2000. Soil isolation and molecular

identification of *Coccidioides immitis*. Mycologia 92:406–410.

Griffith, G. E., J. M. Omernik, D. W. Smith, T. D. Cook, E. Tallyn, K. Moseley, and C. B. Johnson.

2016. Ecoregions of California. US Geological Survey Open-File Report 2016:22–26.

Grinnell, J. 1923. The burrowing rodents of California as agents in soil formation. Journal of

Mammalogy 4:137–149.

Gu, Z., R. Eils, and M. Schlesner. 2016. Complex heatmaps reveal patterns and correlations in

963   multidimensional genomic data. Bioinformatics 32:2847–2849.

964   Hamilton, B. T., B. L. Roeder, K. A. Hatch, D. L. Eggett, and D. Tingey. 2015. Why is small mammal

965   diversity higher in riparian areas than in uplands? Journal of Arid Environments 119:41–

966   50.

967   Hamm, P. S., R. C. Mueller, C. R. Kuske, and A. Porras-Alfaro. 2020. Keratinophilic fungi:

968   Specialized fungal communities in a desert ecosystem identified using cultured-based

969   and Illumina sequencing approaches. Microbiological Research 239:126530.

970   Hanks, D. L. 1963. A study of the Coprophilous Ascomycetes of Utah.

971   Hawkins, L. K. 1996. Burrows of kangaroo rats are hotspots for desert soil fungi. Journal of Arid

972   Environments 32:239–249.

973   Hawkins, L. K. 1999. Microfungi associated with the banner-tailed kangaroo rat, Dipodomys

974   spectabilis. Mycologia 91:735–741.

975   Head, J. R., S. K. Camponuri, A. K. Weaver, L. Montoya, E. Lee, M. T. Radosevich, I. Jones, R.

976   Wagner, A. Bhattachan, G. Campbell, and others. 2024. Small mammals and their

977   burrows shape the distribution of Coccidioides in soils: a long-term ecological

978   experiment. bioRxiv:2024–09.

979   Head, J. R., G. Cooksey, A. K. Heaney, T. Y. Alexander, I. Jones, A. Bhattachan, S. K. Campo, R.

980   Wagner, W. Mgbara, S. Phillips, and others. 2022. Influence of meteorological factors and

981   drought on coccidioidomycosis incidence in California, 2000-2020. medRxiv.

Heydari, S., F. Siavoshi, A. Sarrafnejad, and R. Malekzadeh. 2021. Coniochaeta fungus benefits

from its intracellular bacteria to form biofilm and defend against other fungi. Archives of

Microbiology 203:1357–1366.

Hollister, J., T. Shah, A. L. Robitaille, M. W. Beck, and M. Johnson. 2022. elevatr: Access Elevation

Data from Various APIs.

Hsu, A. P., A. Korzeniowska, C. C. Aguilar, J. Gu, E. Karlins, A. J. Oler, G. Chen, G. V. Reynoso, J.

Davis, A. Chaput, and others. 2022. Immunogenetics associated with severe

coccidioidomycosis. JCI insight 7:e159491.

Huang, J. Y., B. Bristow, S. Shafir, and F. Sorvillo. 2012. Coccidioidomycosis-associated Deaths,

United States, 1990–2008. Emerging Infectious Diseases 18:1723–1728.

Hurst, C. J. 2019. Dirt and disease: the ecology of soil fungi and plant fungi that are infectious for

vertebrates. Understanding terrestrial microbial communities:289–405.

Johnson, S. M., E. L. Carlson, F. S. Fisher, and D. Pappagianis. 2014. Demonstration of

Coccidioides immitis and *Coccidioides posadasii* DNA in soil samples collected from

Dinosaur National Monument, Utah. Sabouraudia 52:610–617.

Kahle, D., and H. Wickham. 2013. ggmap: Spatial Visualization with ggplot2. The R Journal

5:144–161.

Khan, Z., J. Gené, S. Ahmad, J. Cano, N. Al-Sweih, L. Joseph, R. Chandy, and J. Guarro. 2013.

Coniochaeta polymorpha, a new species from endotracheal aspirate of a preterm

neonate, and transfer of Lecythophora species to Coniochaeta. Antonie Van

Leeuwenhoek 104:243–252.

Kolivras, K. N., and A. C. Comrie. 2003. Modeling Valley fever (coccidioidomycosis) incidence on

the basis of climate conditions. International journal of biometeorology 47:87–101.

Kollath, D. R., J. R. Mihaljevic, and B. M. Barker. 2022. PM10 and Other Climatic Variables Are

Important Predictors of Seasonal Variability of Coccidioidomycosis in Arizona.

Microbiology Spectrum 10:e01483-21.

Kollath, D. R., K. J. Miller, and B. M. Barker. 2019. The mysterious desert dwellers: *Coccidioides*

immitis and *Coccidioides posadasii*, causative fungal agents of coccidioidomycosis.

Virulence 10:222–233.

Kollath, D. R., M. M. Teixeira, A. Funke, K. J. Miller, and B. M. Barker. 2020. Investigating the role

of animal burrows on the ecology and distribution of *Coccidioides* spp. in Arizona soils.

Mycopathologia 185:145–159.

Koufopanou, V., A. Burt, and J. W. Taylor. 1997. Concordance of gene genealogies reveals

reproductive isolation in the pathogenic fungus Coccidioides immitis. Proceedings of the

National Academy of Sciences 94:5478–5482.

Laabs, D., M. Allaback, and D. Mitchell. 2022. Distribution, morphology, and karyotype of San

Joaquin pocket mice from the western Mojave Desert. California Fish and Wildlife

Journal 108:e10.

Lacy, G. H., and F. E. Swatek. 1974. Soil ecology of *Coccidioides immitis* at Amerindian middens

in California. Applied microbiology 27:379–388.

Lang, J. D. 2004. Rodent-flea-plague relationships at the higher elevations of San Diego County,

California. Journal of vector ecology: journal of the Society for Vector Ecology 29:236–

247.

Lauer, A., J. D. H. Baal, J. C. H. Baal, M. Verma, and J. M. Chen. 2012. Detection of *Coccidioides*

immitis in Kern County, California, by multiplex PCR. Mycologia 104:62–69.

Lauer, A., V. Etyemezian, G. Nikolich, C. Kloock, A. F. Arzate, F. Sadiq Batcha, M. Kaur, E. Garcia, J.

Mander, and A. Kayes Passaglia. 2020. Valley fever: Environmental Risk Factors and

Exposure Pathways Deduced from Field Measurements in California. International

Journal of Environmental Research and Public Health 17:5285.

Legendre, P., and L. Legendre. 2012. Numerical ecology. Elsevier.

Litvintseva, A. P., N. Marsden-Haug, S. Hurst, H. Hill, L. Gade, E. M. Driebe, C. Ralston, C. Roe, B.

M. Barker, M. Goldoft, and others. 2015. Valley fever: finding new places for an old

disease: *Coccidioides immitis* found in Washington State soil associated with recent

human infection. Clinical Infectious Diseases 60:e1–e3.

Liu, Y., L. He, L. An, T. Helgason, and H. Feng. 2009. Arbuscular mycorrhizal dynamics in a

chronosequence of Caragana korshinskii plantations. FEMS Microbiology Ecology 67:81–

92.

Lloyd, M. G., and R. A. Baldwin. 2021. Burrowing Rodents: Developing a Management Plan for

Organic Agriculture in California.

Lytvynenko, Y. I., and V. P. Hayova. 2018. New and noteworthy records of coprophilous species

of Coniochaeta and Sordaria (Sordariomycetes, Ascomycota) from Ukraine. Український

ботанічний журнал:538–551.

Maddy, K. 1965. Observations on *Coccidioides immitis* found growing naturally in soil. Arizona

Med. 22:281–288.

Magan, N. 2007. Fungi in extreme environments. The mycota 4:85–103.

Mahoney, D., and J. LaFavre. 1981. Coniochaeta extramundana, with a synopsis of other

Coniochaeta species. Mycologia 73:931–952.

Marsden-Haug, N., H. Hill, A. P. Litvintseva, D. M. Engelthaler, E. M. Driebe, C. C. Roe, C. Ralston,

S. Hurst, M. Goldoft, L. Gade, and others. 2014. *Coccidioides immitis* identified in soil

outside of its known range—Washington, 2013. MMWR. Morbidity and mortality weekly

report 63:450.

McNew, L. B., D. K. Dahlgren, and J. L. Beck. 2023. Rangeland Wildlife Ecology and Conservation.

Springer Nature.

Merritt, D. M., and H. L. Bateman. 2012. Linking stream flow and groundwater to avian habitat

in a desert riparian system. Ecological Applications 22:1973–1988.

Nguyen, A., G. S. Cooksey, Y. Djamba, A. Nicolici, S. Jain, and D. Vugia. 2019. Epidemiologic

Summary of Valley fever (Coccidioidomycosis) in California, 2019. California Department

of Public Health.

Nguyen, C., B. M. Barker, S. Hoover, D. E. Nix, N. M. Ampel, J. A. Frelinger, M. J. Orbach, and J. N.

Galgiani. 2013. Recent advances in our understanding of the environmental,

epidemiological, immunological, and clinical dimensions of coccidioidomycosis. Clinical

microbiology reviews 26:505–525.

Nilsson, R. H., S. Anslan, M. Bahram, C. Wurzbacher, P. Baldrian, and L. Tedersoo. 2019.

Mycobiome diversity: high-throughput sequencing and identification of fungi. Nature

Reviews Microbiology 17:95–109.

Nokes, B. T., and J. E. Blair. 2020. Coccidioidomycosis in transplantation. Emerging Transplant

Infections: Clinical Challenges and Implications:1–18.

Ocampo-Chavira, P., R. Eaton-Gonzalez, and M. Riquelme. 2020. Of mice and fungi: Coccidioides

spp. distribution models. Journal of Fungi 6:320.

Oksanen, J., F. G. Blanchet, M. Friendly, R. Kindt, P. Legendre, D. McGlinn, P. R. Minchin, R. B.

O'Hara, G. L. Simpson, P. Solymos, M. H. H. Stevens, E. Szoecs, and H. Wagner. 2019.

vegan: Community Ecology Package.

Ophüls, W. 1905. Further observations on a pathogenic mould formerly described as a

protozoon (*Coccidioides immitis*, *Coccidioides pyogenes*). The Journal of experimental

medicine 6:443.

Padgham, M., B. Rudis, R. Lovelace, and M. Salmon. 2017. osmdata. The Journal of Open Source

Software 2.

Park, B. J., K. Sigel, V. Vaz, K. Komatsu, C. McRill, M. Phelan, T. Colman, A. C. Comrie, D. W.

Warnock, J. N. Galgiani, and others. 2005. An epidemic of coccidioidomycosis in Arizona

associated with climatic changes, 1998–2001. The Journal of infectious diseases

191:1981–1987.

Pasternak, G., S. Kienzle, L. Barrett, P. Henzi, L. R. Brown, and A. Fuller. 2013. Population ecology

of vervet monkeys in a high latitude, semi-arid riparian woodland. Koedoe: African

Protected Area Conservation and Science 55:1–9.

Patten, D. T. 1998. Riparian ecosytems of semi-arid North America: Diversity and human

impacts. Wetlands 18:498–512.

Pedregosa, F., G. Varoquaux, A. Gramfort, V. Michel, B. Thirion, O. Grisel, M. Blondel, P.

Prettenhofer, R. Weiss, V. Dubourg, J. Vanderplas, A. Passos, and D. Cournapeau. 2011.

Scikit-learn: Machine Learning in Python. Journal of Machine Learning Research

12:2825–2830.

Plunkett, O., and F. Swatek. 1957. Ecological studies of *Coccidioides immitis*. Pages 158–160

Proceedings of the Symposium on Coccidioidomycosis. AZ. Public Health Service

Washington DC.

Porter, W. T., L. Gade, P. Montfort, J. R. Mihaljevic, J. R. Bowers, A. Willman, B. A. Klimowski, B. J.

LaFleur, R. H. Sunenshine, J. Collins, and others. 2024. Understanding the exposure risk

of aerosolized Coccidioides in a Valley fever endemic metropolis. Scientific Reports

14:1311.

Posadas, A. 1892. Un nuevo caso de micosis fungoidea con psorospermias. An Circ Med Argent

15:585–97.

R Core Team. 2020. R: A Language and Environment for Statistical Computing. R Foundation for

Statistical Computing, Vienna, Austria.

Radosevich, M., J. Head, L. Couper, A. Gomez-Weaver, S. Camponuri, L. Montoya, J. Taylor, and J.

Remais. 2024, September 28. Characterizing the soil microbial community associated

with the fungal pathogen*Coccidioides immitis*. Cold Spring Harbor Laboratory.

Ramirez, M., S. Chulze, and N. Magan. 2004. Impact of osmotic and matric water stress on

germination, growth, mycelial water potentials and endogenous accumulation of sugars

and sugar alcohols in Fusarium graminearum. Mycologia 96:470–478.

Rappleye, C. A., and W. E. Goldman. 2006. Defining virulence genes in the dimorphic fungi.

Annu. Rev. Microbiol. 60:281–303.

Rice, A. V. 2005. The taxonomy and biology of myxotrichaceous fungi from a Sphagnum bog in

Alberta, Canada.

Richardson, M. 1998. Coniochaeta polymegasperma and Delitschia trichodelitschioides, two

new coprophilous ascomycetes. Mycological research 102:1038–1040.

Sakaeyama, S.-I., A. Sano, Y. Murata, K. Kamei, K. Nishimura, and K. Hatai. 2007. Lecythophora

hoffmannii isolated from a case of canine osteomyelitis in Japan. Sabouraudia 45:267–

272.

Salazar-Hamm, P. S. 2021. Breathing Can Be Dangerous: *Coccidioides* and the Diverse

Community of the Lung Mycobiome. PhD Thesis, The University of New Mexico.

Salazar-Hamm, P. S., K. N. Montoya, L. Montoya, K. Cook, S. Liphardt, J. W. Taylor, J. A. Cook, and

D. O. Natvig. 2022. Breathing can be dangerous: Opportunistic fungal pathogens and the

diverse community of the small mammal lung mycobiome. Frontiers in Fungal Biology

3:996574.

Saubolle, M. A., B. R. Wojack, A. M. Wertheimer, A. Z. Fuayagem, S. Young, and B. A. Koeneman.

2018. Multicenter clinical validation of a cartridge-based real-time PCR system for

detection of Coccidioides spp. in lower respiratory specimens. Journal of Clinical

Microbiology 56:e01277-17.

Scott, R. L., T. E. Huxman, G. A. Barron-Gafford, G. Darrel Jenerette, J. M. Young, and E. P.

Hamerlynck. 2014. When vegetation change alters ecosystem water availability. Global

Change Biology 20:2198–2210.

Sharpton, T. J., J. E. Stajich, S. D. Rounsley, M. J. Gardner, J. R. Wortman, V. S. Jordar, R. Maiti, C.

D. Kodira, D. E. Neafsey, Q. Zeng, C.-Y. Hung, C. McMahan, A. Muszewska, M. Grynberg,

1133   M. A. Mandel, E. M. Kellner, B. M. Barker, J. N. Galgiani, M. J. Orbach, T. N. Kirkland, G. T.

1134   Cole, M. R. Henn, B. W. Birren, and J. W. Taylor. 2009. Comparative genomic analyses of

1135   the human fungal pathogens *Coccidioides* and their relatives. Genome Research

1136   19:1722–1731.

1137   Sun, S. H., and M. Huppert. 1976. A cytological study of morphogenesis in Coccidioides immitis.

1138   Sabouraudia: Journal of Medical and Veterinary Mycology 14:185–198.

1139   Tamerius, J. D., and A. C. Comrie. 2011. Coccidioidomycosis incidence in Arizona predicted by

1140   seasonal precipitation. PloS one 6:e21009.

1141   Taylor, D. L., W. A. Walters, N. J. Lennon, J. Bochicchio, A. Krohn, J. G. Caporaso, and T.

1142   Pennanen. 2016. Accurate Estimation of Fungal Diversity and Abundance through

1143   Improved Lineage-Specific Primers Optimized for Illumina Amplicon Sequencing. Applied

1144   and Environmental Microbiology 82:7217–7226.

1145   Taylor, J. W., and B. M. Barker. 2019. The endozoan, small-mammal reservoir hypothesis and the

1146   life cycle of *Coccidioides* species. Medical mycology 57:S16–S20.

1147   Teixeira, M. M., P. Alvarado, C. C. Roe, G. R. Thompson III, J. S. Patané, J. W. Sahl, P. Keim, J. N.

1148   Galgiani, A. P. Litvintseva, D. R. Matute, and others. 2019. Population structure and

1149   genetic diversity among isolates of Coccidioides posadasii in Venezuela and surrounding

1150   regions. MBio 10:10–1128.

1151   Teixeira, M. M., and B. M. Barker. 2016. Use of population genetics to assess the ecology,

1152   evolution, and population structure of *Coccidioides*. Emerging infectious diseases

1153   22:1022.

United States Geological Survey. 2021a. The National Map US Topo, Mount Brewer Quadrangle

California, 7.5 Minute Series.

United States Geological Survey. 2021b. The National Map US Topo, San Benito Mountain

Quadrangle California, 7.5 Minute Series.

Vargas-Gastelum, L., A. L. Romero-Olivares, A. E. Escalante, A. Rocha-Olivares, C. Brizuela, and

M. Riquelme. 2015. Impact of seasonal changes on fungal diversity of a semi-arid

ecosystem revealed by 454 pyrosequencing. FEMS microbiology ecology 91:fiv044.

Wagner, R., L. Montoya, C. Gao, J. R. Head, J. Remais, and J. W. Taylor. 2022. The air mycobiome

is decoupled from the soil mycobiome in the California San Joaquin Valley. Molecular

Ecology 31:4962–4978.

Wagner, R., L. Montoya, J. R. Head, S. Campo, J. Remais, and J. W. Taylor. 2023. Coccidioides

undetected in soils from agricultural land and uncorrelated with time or the greater soil

fungal community on undeveloped land. PLoS Pathogens 19:e1011391.

Weaver, E., K. N. Kolivras, R. Q. Thomas, V. A. Thomas, and K. M. Abbas. 2020. Environmental

factors affecting ecological niche of Coccidioides species and spatial dynamics of Valley

fever in the United States. Spatial and Spatio-temporal Epidemiology 32:100317.

Whiston, E., and J. W. Taylor. 2016. Comparative phylogenomics of pathogenic and

nonpathogenic species. G3: Genes, Genomes, Genetics 6:235–244.

Whitford, W. G., and F. R. Kay. 1999. Biopedturbation by mammals in deserts: a review. Journal

of Arid Environments 41:203–230.

Zinger, L., A. Bonin, I. G. Alsos, M. Bálint, H. Bik, F. Boyer, A. A. Chariton, S. Creer, E. Coissac, B. E.

Deagle, and others. 2019. DNA metabarcoding—Need for robust experimental designs

to draw sound ecological conclusions. Wiley Online Library.

1177

1178

1179

1180

1181

1182

1183

1184

1185

1186

1187

1188

1189

1190

1191

1192

1193

1194

1195

1196

Supporting Information Captions

S1 Fig. Map of urban surfaces sampled in Coalinga, CA in August 2020 and tested using the CocciENV qPCR assay. n = 100. Point colors correspond to positive replicate wells. Map data (lines) were acquired from OpenStreetMap (www.openstreetmap.org) under the Open Data Commons Open Database License (OdbL) (https://www.openstreetmap.org/copyright). Map tiles are by Stamen Design (www.stamen.com) under Creative Commons By Attribution (CC BY 4.0) (https://creativecommons.org/licenses/by/4.0/).

S2 Fig. Extended map of Los Gatos Creek rodent burrow soil sampling including April 2024 sites. All samples were tested using the CocciENV qPCR assay. Elevation profile (meters above sea level) and qualitative categorization of sample locations (a). Map of sample coordinates (b). Inset shows map extent. Landscape classification derived from the California Farmland Mapping and Monitoring Program 2018. Point colors correspond to positive replicate wells. Map data (lines) were acquired from OpenStreetMap (www.openstreetmap.org) under the Open Data Commons Open Database License (OdbL) (https://www.openstreetmap.org/copyright). Map tiles are by Stamen Design (www.stamen.com) under Creative Commons By Attribution (CC BY 4.0) (https://creativecommons.org/licenses/by/4.0/). California shapefile is public domain data by Natural Earth (www.naturalearthdata.com).

S3 Fig. The mean proportional abundance of the 30 most common fungal genera from sites along the Kern River as a function of site and sampling timepoint. (-) = *Coccidioides* negative samples. (+) = *Coccidioides* positive samples. n = 28.

S4 Fig. The mean proportional abundance of the 30 most common fungal genera from sites along Los Gatos Creek as a function of sampling location (rodent burrow entrance vs interior). (-) = *Coccidioides* negative samples. (+) = *Coccidioides* positive samples. n = 40.

S1 Table. Sampling enumeration for the current study (**bold text**) and all other studies known to the authors where environmental *Coccidioides* was detected in soil, air, or both. The location and the method of detection are shown here. Total studies (including the current study) = 42. (+) = positive samples.

S2 Table. *Coccidioides* detection in soils and settled dust using the CocciEnv qPCR assay as a function of site and detection level. Numbers in parentheses indicate the minimum proportion of replicate wells with CT < 40 and logarithmic amplification. Only samples detected in ≥ 3 replicate wells were considered positive detections. _*_ = Bakersfield area / non-Kern River site

S3 Table. *Coccidioides* detection using the CocciEnv qPCR assay from sites with >1 timepoint as a function of site, date and detection level. Numbers in parentheses indicate the minimum proportion of replicate wells with CT < 40 and logarithmic amplification. * = Bakersfield area / non-Kern River site

S4 Table. Enumeration of subset of soil samples selected for ITS2 amplicon sequencing. n = 35 negative samples; 33 positive samples.

S5 Table. Generalized Linear Mixed Model (GLMM) coefficient table showing *Coccidioides* detection as a function of elevation, sampling month and soil moisture data (fixed effects), with site included as a random effect, across all Kern River sites. n = 830.

S6 Table. Modified generalized Linear Mixed Model (GLMM) coefficient table, excluding samples collected above 210 meters showing *Coccidioides* detection as a function of elevation, sampling month and soil moisture data (fixed effects), with site included as a random effect, across all Kern River sites (below 210 meters). n = 760.

S7 Table. Logistic regression coefficient table showing *Coccidioides* detection as a function of sampling site, sampling month and soil moisture data between Kern River sites at Panorama Park West and University Place. n = 202.

S8 Table. PERMANOVA coefficient table showing the association between the soil fungal community and *Coccidioides* detection (Presence), as a function of sampling site and sampling date, from rodent burrows soils collected from along the Kern River. *Coccidioides* detection determined using the CocciEnv qPCR assay. Fungal community structure modeled as species-level Bray-Curtis dissimilarities. Permutations = 1000. n = 28.

S9 Table. Pairwise fungal community differences (PERMANOVA) between Kern River sites using the 'pairwiseadonis2' function. Number of samples collected at each site shown in parentheses. Permutations = 1000. n = 28.

S10 Table. PERMANOVA coefficient table showing the association between the soil fungal community, *Coccidioides* detection (Presence) and sampling location (burrow interior vs entrance) from rodent burrow soils collected in August 2020 from the Los Gatos Creek "Hwy33 West" site. *Coccidioides* detection determined using the CocciEnv qPCR assay. Fungal community structure modeled as species-level Bray-Curtis dissimilarities. Permutations = 1000. n = 40.

S11 Table. Percent abundance of the 30 most abundant fungal genera at Kern River and Los Gatos Creek Sites. Values are means between replicates. Underlined genera appear uniquely in the top 30 genera at each location.

S12 Table: Indicator species for *Coccidioides* positive and *Coccidioides* negative soil samples collected from within Kern River rodent burrows. Significance was calculated in indicspecies version 1.7.12 with 1000 permutations. Sequences were cross-referenced via NCBI Nucleotide Blast (https://blast.ncbi.nlm.nih.gov/) for each prospective indicator species. • = equally likely alternative species (in subsection). IndVal and p-value apply to all alternative species.

S13 Table: Indicator species for *Coccidioides* positive soil samples collected from within Los Gato Creek rodent burrows. Significance was calculated in indicspecies version 1.7.12 with 1000 permutations. Sequences were cross-referenced via NCBI Nucleotide Blast (https://blast.ncbi.nlm.nih.gov/) for each prospective indicator species. • = equally likely alternative species (in subsection). IndVal and p-value apply to all alternative species.

1288   **S14 Table:** Indicator species for *Coccidioides* negative soil samples collected from within Los
1289   Gatos Creek rodent burrows. Significance was calculated in indicspecies version 1.7.12 with
1290   1000 permutations. Sequences were cross-referenced via NCBI Nucleotide Blast
1291   (https://blast.ncbi.nlm.nih.gov/) for each prospective indicator species. • = equally likely
1292   alternative species (in subsection). IndVal and p-value apply to all alternative species.
1293
S15 Table: Indicator species for *Coccidioides* samples (positive and negative) collected from
surface soils outside Los Gatos Creek rodent burrows. Significance was calculated in
indicspecies version 1.7.12 with 1000 permutations. Sequences were cross-referenced via NCBI
Nucleotide Blast (https://blast.ncbi.nlm.nih.gov/) for each prospective indicator species. • =
equally likely alternative species (in subsection). IndVal and p-value apply to all alternative
species.
1300
1301   **S1_data_qpcr_meta:** All qpcr data and all metadata.
1302
1303   **S2_data_soil_moisture:** All soil moisture data.
1304
1305   **S1_code_sequence_processing:** All code used to process raw sequencing data.
1306
1307   **S2_code_analysis:** All code used for statistical analyses.