Moulting in Pancrustacea is characterised by both deeply conserved and recently evolved gene modules

Kenneth Kim[1,2], Giulia Campli [1,2], Sagane Joye [1,2], Ariel D. Chipman [3], Robert M. Waterhouse [2], Marc Robinson-Rechavi [1,2]

[1] Department of Ecology and Evolution, University of Lausanne, 1015 Lausanne, Switzerland.
[2] SIB Swiss Institute of Bioinformatics, 1015 Lausanne, Switzerland.
[3] Department of Ecology, Evolution & Behavior, The Silberman Institute of Life Science, The Hebrew University of Jerusalem, Jerusalem, Israel

Abstract

Arthropods such as insects and crustaceans, which together form the monophyletic group Pancrustacea, possess a rigid chitinous exoskeleton that must be periodically shed through moulting to allow growth and morphological change. Although moulting is a deeply conserved developmental process across Arthropoda, our understanding of its molecular mechanisms is still largely derived from insect model species. Lineage-specific innovations and losses of moulting related genes raise fundamental questions on the extent of its conservation outside non-insect arthropods. Here, we investigate the evolutionary conservation of moulting gene expression across five representative pancrustacean species. Changes in gene expression during moulting are characterized by both deeply conserved and lineage-specific gene modules. Time-series gene expression analyses reveal that these lineage-specific signatures are not uniformly distributed across the moulting process: the middle transitional phase is more lineage-specific, thereby exhibiting an inverse hourglass pattern. This is likely due to life-history specific processes, development of the cuticle and specialized structures of the exoskeleton. Overall, this study provides evidence for both the evolutionary conservation and divergence of this key post-embryonic developmental process and highlights the modular architecture of the moulting programme.

Introduction

The phylum Arthropoda includes a vast array of extant and extinct taxa, such as insects, crustaceans, arachnids, trilobites, or sea scorpions. These organisms share defining features: segmented bodies, paired jointed appendages, and a chitin-based exoskeleton, often reinforced with proteins or minerals (Chapman et al., 2013a; Daley et al., 2018; Giribet & Edgecombe, 2019). The modular organization of their segmented body plan, combined with a protective exoskeleton, has enabled arthropods to thrive in an extraordinary range of ecological niches. However, their rigid exoskeleton also hinders their ability to grow and must be periodically shed in a process called moulting, which not only enables growth but also facilitates morphological changes during development (Belles, 2020; Chapman et al., 2013a; Truman, 2019; Truman & Riddiford, 2019).

The moulting process is governed by hormonal regulation, primarily through ecdysteroids and sesquiterpenoids (Campli et al., 2024; Cheong et al., 2015; Qu et al., 2018; Truman & Riddiford, 2019). Hormones are synthesized in secretory organs in response to signals from neuropeptides. Once released into the haemolymph, these hormones reach target tissues, where they trigger transcription factors responsible for various moulting pathways (Truman, 2019). As hormone levels decline, neurosecretions activate the "ecdysis motor", a behavioural sequence that leads to the final stages of moulting and exoskeleton hardening (Roer et al., 2015; White & Ewer, 2014; Zieger et al., 2021).

Moulting is coordinated by the increase and decrease of ecdysteroid hormone or its active form 20-hydroxyecdysone (20E) (Truman & Riddiford, 2019) and occurs in distinct moult

stages each characterized by different morphological and behavioural changes (Drach & Tchernigovtzeff, 1967; Gorissen & Sandeman, 2022; Kim et al., 2024; Mykles, 2011). During the pre-moult phase enzymes digest the inner layers of the old cuticle, allowing the exoskeleton to detach from the epidermal cells or also known as apolysis (Roer et al., 2015; X. Zhang et al., 2021). Moulting *sensu stricto*, or ecdysis, involves the rupturing of the old exoskeleton due to body movements and increased haemolymph pressure, enabling the animal to shed the exuvia (Daley & Drage, 2016). In the post-moult phase, the new exoskeleton hardens through processes like sclerotization, mineralization, and pigmentation, which collectively transform the soft cuticle into a mature and hardened exoskeleton.

Most of our knowledge of the molecular and biochemical mechanisms underlying moulting has been derived from a small number of well-established insect model species, such as *Drosophila melanogaster*. As a consequence, molecular insights into moulting largely stem from developmental contexts associated with insect metamorphosis, despite moulting itself being a shared and ancient process across Arthropoda and our understanding of moulting beyond Insecta remains limited (Campli et al., 2024). The increasing availability of genomic resources and sequencing data from a broader range of arthropods, driven in part by large-scale projects such as the i5k initiative (i5K Consortium, 2013) and the European Reference Genome Atlas (ERGA) (Mazzoni et al., 2023) has enabled broader testing of the involvement of genes in moulting that were first identified in insect model species (Campli et al., 2024). These genomic analyses have confirmed the presence of an conserved core set of genes involved in moulting, found

across various lineages (Campli et al., 2024, 2025). While a conserved genetic backbone for moulting exists, lineage-specific modifications are evident such as the incomplete ecdysteroid biosynthetic pathway in barnacles (Dermauw et al., 2020), the missing *Shd* (*cyp314a1*) gene in several fungus-farming ant species (Dermauw et al., 2020), and the loss of certain genes in specific lineages. Such losses include *neverland* in Varroa mites (Cabrera et al., 2015; Campli et al., 2025; Techer et al., 2019) and in beetles (Campli et al., 2025; Perry & Robin, 2025), *phantom (phm)* in chelicerates (Schumann et al., 2018), and *juvenile hormone acid methyltransferase* (*JHAMT*) gene in millipedes (So et al., 2022). Various genes also appear to be restricted to certain clades such as *Nobo* which is only found in Diptera and Lepidoptera (Enya et al., 2015; Koiwai et al., 2020), or *spok* and *Cyp6t3* found only in Drosophilidae (Ono et al., 2006; Ou et al., 2011; Sztal et al., 2007). Collectively, these patterns show that even key genes regulating moulting are not all equally conserved. The conservation of the broader set of genes involved in moulting is not well known. This is partly due to the candidate gene approach commonly used in functional studies, which prioritizes known genes from model organisms and thus leads to a systematic bias towards studying conserved genes. As most genes function within biological pathways or co-regulated modules (Brawand et al., 2011; Harrison et al., 2012; Keidar Haran & Keren, 2022; Oleksiak et al., 2002), investigating the expression patterns of genes during moulting is essential to understand the conservation and evolution of moulting processes across Arthropoda.

Phylogenetic analysis indicates that the hexapod clade, which includes insects, represents an ancient divergence within Pancrustacea, suggesting that insects evolved

from crustacean-like ancestors (Bernot et al., 2023; Rota-Stabelli et al., 2013; Schwentner et al., 2017; Wolfe et al., 2016). This evolutionary relationship makes Pancrustacea an ideal group for studying the conservation and divergence of moulting mechanisms. Moulting in insects and crustaceans is orchestrated by neuropeptide-driven endocrine cascades regulated by specialized steroidogenic glands, the prothoracic gland (PG) in insects and the Y-organ (YO) in crustaceans (Cheong et al., 2015; Mykles, 2021; Mykles & Chang, 2020; Truman & Riddiford, 2019). In insects, moulting is triggered by the release of prothoracicotropic hormone (PTTH), which binds to the Torso receptor on the PG and activates the mitogen-activated protein kinase (MAPK) and mammalian target of rapamycin (mTOR) pathways, leading to the production of ecdysteroids (Truman & Riddiford, 2019). In contrast, crustaceans rely on inhibitory control: moult-inhibiting hormone (MIH), secreted from the eyestalk X-organ/sinus gland complex, suppresses YO activity through a cAMP/Ca^{2+}- and NO/cGMP-dependent signalling cascade (Mykles, 2011). Reduction of MIH release, either through environmental cues or experimental eyestalk ablation, activates the YO and triggers moulting. In addition, the calcium-rich marine environment has enabled crustaceans to evolve heavily calcified, rigid exoskeletons that provide strong protection. In terrestrial environments, where calcium is less readily available, insects have adopted a different strategy by developing lighter, more flexible cuticles that rely on biochemical processes such as laccase-mediated cross-linking for cuticle hardening and tanning (Asano et al., 2019; Volovych et al., 2025). These differences in moulting mechanisms and exoskeletal properties offer a comparative framework for exploring the evolutionary conservation of moulting across Pancrustacea.

In this study, we investigated gene expression conservation of moulting across several insect and malacostracan species, the two largest monophyletic classes of pancrustaceans. We took two approaches to this question. First, in vertical comparison, we examined the gene expression of ecdysis *sensu stricto* and pupariation (larval-prepupal moult) across five representative pancrustacean species (Table 1). Although the terminology for these processes varies in the literature, we specifically refer to moulting modes in this study as either metamorphic, such as larval-prepupal moult in insects, or non-metamorphic, i.e. the instar-instar moult. Of note, specific stages are not necessarily directly homologous across pancrustaceans. We examine shared gene expression patterns, and thus biological processes, during moulting in different species and life-stages. While larval–prepupal transitions in insects are often viewed through the lens of metamorphosis, larval-prepupal moults in insects are also hormonally driven moulting events characterized by a surge in 20-hydroxyecdysone (20E), a period of quiescence (non-feeding phase), cuticle synthesis and involve similar processes to post-ecdysial modifications such as cuticle sclerotization and tanning (Roer et al., 2015; White & Ewer, 2014). Despite the well-established role of 20E in triggering both moulting and larval-prepupal moult, downstream pathways and developmental programs that are potentially conserved between these two processes beyond insects remain largely unexplored. Secondly, in horizontal comparison, we examined gene expression conservation across several timepoints of moulting for *Drosophila melanogaster* and *Litopenaeus vannamei*, the only species in our dataset with time-series data (Supplementary Table 1). This allowed us to examine how gene expression dynamics shift across successive timepoints, and whether expression conservation patterns are consistent across the entire moulting

process. We found that gene expression of moulting is characterized by both highly conserved and lineage-specific signatures, and that this pattern is not uniformly distributed. The mid-phase of the moulting is more lineage specific, exhibiting an inverse hourglass pattern. Our findings highlight the evolutionary conservation of moulting and provide insights into the evolution of moulting strategies in Pancrustacea.

Figure 1. Phylogenetic tree showing the species used in this study. Additional information for each sample used is shown in the table on the right. Icon silhouettes under public domain license were retrieved from Phylopic.

Methods

Data pre-processing

Table 1. Summary of datasets used for gene expression analysis

Taxonomic group	Species	Moult stages	Source
Insecta	*Drosophila melanogaster*	Wandering stage	(Graveley et al., 2011)
		Prepupa	
	Zeugodacus cucurbitae	Wandering stage	(Ward et al., 2021; Wei et al., 2020)
		Prepupa	
	Bactrocera dorsalis	Wandering stage	(S. Liu et al., 2020)
		Prepupa	
		Instar pre-moult	
		Instar post-moult	
Malacostraca	*Litopenaeus vannamei*	Pre-moult	(Gao et al., 2015a)
		Post-moult	
	Scylla paramamosain	Pre-moult	(L. Liu et al., 2022)
		Post-moult	

Table 1 summarizes the RNA sequencing (RNA-seq) data used in this study. Additional time-series datasets were retrieved for *D. melanogaster* from (Ozerova et al., 2025) and for *L. vannamei* from Gao et al., 2015b (Supplementary Table 1). The quality of the reads was checked using FastQC v.0.11.2 (Andrews 2010) and adapter trimming and quality control were performed using Trimmomatic v. 0.36 (Bolger et al., 2014). Trimmed reads were pseudo-aligned against the respective transcriptome of each species using kallisto v0.44.0 (Bray et al., 2016). Transcriptome and annotation files were retrieved from National Center for Biotechnology Information (NCBI) databases (Sayers et al., 2024). The quality and completeness of the genome assembly of each species were assessed with Benchmarking Universal Single-Copy Orthologs (BUSCO) using the Arthropoda lineage dataset (odb10) as reference (Simão et al., 2015) through the A3CAT database (Feron & Waterhouse, 2022). Transcript level expression was imported into R v4.2.2 and summarised to the gene level using the R/tximport v1.10.1 (Soneson et al., 2016).

Construction of orthogroup gene expression matrix

To examine gene expression patterns of moulting stages between groups of species, orthologous genes (orthogroups) were identified using Orthologer v3.0.2 with default parameters (Kuznetsov et al., 2023) after filtering the proteomes for the single longest protein-coding isoform of each gene. A representative gene with the highest mean expression value per species was selected for each orthogroup generating a read count matrix of 5543 orthologues identified across the five species.

Principal component analysis

Prior to conducting principal component analysis (PCA), combined raw read counts of all samples were transformed using variance stabilizing transformation (VST function) implemented in DESeq2 (Love et al., 2014). This transformation generated constant variances across all read counts creating a homoscedastic dataset. To account for the effect of species on gene expression we included species as an explanatory variable in the model when constructing the read counts matrix. A PCA was conducted to examine the overall gene expression pattern of the dataset, using the PCAtools R package (Blighe and Lun 2021) on the VST RNA-seq counts for 5543 orthologues identified across the five species. The top 10 PCs were selected, and individual eigenvalues were tested for correlation with the traits of interest: taxonomic group, moulting stage, and species.

Differential gene expression analysis

Differential gene expression analyses between pre-moult and post-moult stages were performed individually per species using DESeq2 (Love et al., 2014), with P-values adjusted using the Benjamini–Hochberg (Benjamini & Hochberg, 1995) procedure to control for false discovery rate (FDR). An adjusted *P*-value of 0.05 was used as the threshold for identifying differentially expressed genes (DEGs). For simplicity, wandering and prepupal stages of the larval-prepupal moult are referred to as simply pre-moult and post-moult, respectively. The resulting differentially expressed genes from each species were grouped based on orthology of DEGs into the following categories: species specific genes (differential expression not shared with any other species), Pancrustacea shared genes (differential expression in at least one insect and one malacostracan), class specific genes restricted to either class Insecta or Malacostraca (differential expression in at least two species per group), and genes which have differential expression exclusive to metamorphic or to non-metamorphic samples. For each DEG, we also assessed whether its corresponding orthologous group, irrespective of differential expression, was restricted to a single lineage (insect-only or malacostracan-only), shared between insects and malacostracans (Pancrustacea-shared), or lacked detectable orthologs in other taxa (orphan genes). Thus, e.g., a species specific DEG can be either an orphan gene, or the only DEG in an orthogroup shared with other species in the study.

Clustering and functional analysis

Clust (Abu-Jamous & Kelly, 2018) was used to identify and extract distinct groups of genes that display correlated expression patterns across all samples. Briefly, Clust

applies k-means clustering to all genes, using different k-values to generate an initial set of seed clusters. These seed clusters are then refined through M–N scatter plots to isolate large, non-overlapping clusters with low dispersion. Genes with flat expression were filtered out of the analysis. Several cluster tightness values ranging from 0.1 to 1 were evaluated. Smaller tightness values produce larger and more diffuse clusters, while larger values produce smaller and tighter clusters. A tightness value of 1 was used for the study. Gene Ontology enrichment analysis (Ashburner et al., 2000) of genes belonging to each cluster was performed using ClusterProfiler (Wu et al., 2021). Putative cuticular protein families were identified using CutProtFam-Pred (Ioannidou et al., 2014), which uses Hidden Markov Models based on CuticleDB (Magkrioti et al., 2004) to detect and classify cuticular protein families based on conserved sequence motifs, including cuticular protein subfamilies Rebers and Riddiford type 1 (RR-1) and type 2 (RR-2), as well as additional families such as tweedle and cuticular protein analogous to peritrophins types 1 and 3 (CPAP1 and CPAP3).

Transcriptome age analysis and Phylostratigraphy

To examine the conservation of gene expression across different timepoints of moulting, phylostratigraphy analysis was used to estimate the relative transcriptome age of the time-series samples (Supplementary Table 1) (Domazet-Lošo & Tautz, 2010). VST-transformed counts were generated using DESeq2 (Love et al., 2014), and mean expression values for each stage were calculated. The resulting gene expression matrix was then clustered using Clust (Abu-Jamous & Kelly, 2018), and the phylostratum composition for each cluster was determined by using GenEra v1.4.0 (Barrera-Redondo et al., 2023). All genes of a given target genome were queried against a protein database

built from eukaryotic reference genomes retrieved from NCBI using DIAMOND *blastp* in more-sensitive mode with an *e*-value threshold of 10^{-5} (Buchfink et al., 2015). *Blastp* hits were considered homologues of the query protein and were used to determine the phylostrata of the gene.

The Transcriptome Age Index (TAI) of RNA-seq samples was calculated as described previously (Domazet-Lošo & Tautz, 2010) using the *myTAI* (v2.0.0) (Drost et al., 2018) package in R (v4.3.2). TAI values were computed as the weighted mean of the phylostratum rank of a given gene based on its expression level in the transcriptome of an RNA-seq sample. VST transformed gene expression was used, which like log-transform removes the impact of extreme values (see J. Liu & Robinson-Rechavi, 2018; Piasecka et al., 2013). A high TAI value indicates a younger transcriptome age, whereas a lower TAI corresponds to an older transcriptome. Statistical significance of the TAI profile was determined through a permutation-based test implemented in *myTAI* (v2.0.0) (Drost et al., 2018).

Results

Overall gene expression patterns cluster by lineage and moult stages

Figure 2. Principal Component Analysis of RNA-seq data. **A**. PC1 and PC2 cluster samples by taxonomic group and moult stage. **B**. PC2 and PC6 show clustering of samples by moult stage. Left and top portion shows prepupa and post-moult stage while the samples at the centre and bottom show wandering and pre-moult stages. **C**. Eigenvalue correlations of the principal components with respect to taxonomic group, species, and moult stage. Colour indicates the strength of the correlation and asterisks indicate level of significance after Benjamini–Hochberg (Benjamini and Hochberg 1995) correction with $*P < 0.05$, $**P < 0.01$, and $***P < 0.001$.

To examine the gene expression pattern of moulting dataset across all samples, we performed a Principal Component Analysis (PCA) on the VST RNA-seq counts for the 5543 single-copy orthogroups identified across all species (Fig. 2). The first 10 principal

components (PCs) explained 95% of the variance in gene expression, with PC1 and PC2 alone accounting for 54%. We tested whether these PCs were significantly associated with either moult stage, subphylum, or species (Fig. 2C). The strongest correlation with moult stage was PC6 and plotting PC2 and PC6 clustered samples by moult stages. Among the genes which contribute the most to PC 6 are genes known to be involved in moulting, such as hormone receptor 3 (*Hr3)*, ecdysone induces protein 78c (*Eip78c)*, ecdysis triggering hormone receptor (*ETHR)*, and various cuticular proteins and chitinases (Supplementary Table 2). There are also uncharacterized genes, such as CG11892 which has predicted ecdysteroid-22 kinase activity, and CG12009, which has a predicted chitin binding perithophin-A domain. All samples were derived from whole-body extracts, yet despite variable involvement of different tissues in moulting, we detected a clear signal of moulting in the dataset. This likely reflects the large changes in gene expression just before and after ecdysis and pupariation.

Shared and lineage-specific differentially expressed genes in non-metamorphic moults

We first consider only the non-metamorphic moults, i.e. the instar moult of *B. dorsalis* and the two malacostracans (Figure 3, top). While the largest fraction of differentially expressed genes (DEGs) between pre- and post-moult are classified as lineage-specific (i.e. species-specific or class-specific), there are also many DEGs which are shared among pancrustaceans.

Figure 3. Classification of differentially expressed genes (DEGs) and their corresponding orthology distribution. Top: non metamorphic moults; bottom: metamorphic moults. "Class specific" are DEGs restricted to Malacostraca or to insects.

These shared DEGs include known members of the late genes and ecdysis neuromotor pathways, such as *knickkopf* (*Knk*), *obstructor-A* (*Obs-A*), retroactive *(Rtv), serpentine* (*serp*), *krotzkopf verkehrt* (*kkv*), and various chitin deacetylases and chitinases (Campli et al., 2024; Moussian et al., 2015; Pesch et al., 2015, 2016). In addition, many of these DEGs are involved in cuticle development, and include various esterases, ecdysteroid kinases, and structural components such as extracellular matrix proteins, gap junction elements, and other support structures (Supplementary Table 3). We also found cytochrome P450 (CYP) genes and other steroid binding proteins and transporters, some of which have been shown to be involved in ecdysone metabolism and sterol transport, such as organic anion transporter polypeptides, Niemann-Pick type C (*NPC*), δ-

aminolevulinic acid synthase (*Alas*), megalin, and membrane steroid binding protein (*MSBP*) (Fujii-Taira et al., 2009; Huang et al., 2005; Nakaoka et al., 2017; Riedel et al., 2011). Many of these genes have orthologues outside Pancrustacea, indicating that their involvement in moulting might be conserved more broadly.

Comparison of metamorphic and non-metamorphic moults

To examine whether these Pancrustacea-shared DEGs are restricted to non-metamorphic moults or are conserved more broadly across moulting modes, we extended the comparison to include metamorphic moults in insects (Figure 3). Interestingly, only a small fraction of DEGs was exclusively shared among non-metamorphic moults, whereas Pancrustacea-shared DEGs were broadly represented among DEGs in metamorphic moults. This indicates that many conserved components of the moulting programme are deployed in both moulting modes.

There were differences between insects and malacostracans in the number of species-specific compared to class-specific DEGs. Malacostracan species exhibited a higher proportion of species-specific DEGs than insects; splitting them between upregulated and downregulated genes shows similar trends (Supplementary Figure S1). This pattern likely reflects the greater divergence among species within each group. *L. vannamei* and *S. paramamosain* belong to distinct decapod groups Dendrobranchiata and Pleocyemata (Reptants), respectively, which diverged approximately 400 million years ago. They exhibit markedly different life histories and ecology with *L. vannamei* fully aquatic and *S. paramamosain* semi-terrestrial. In contrast, *B. dorsalis*, *Z. cucurbitae* (both Tephritidae),

and *D. melanogaster* (Drosophilidae), diverged only around 150 million years ago (estimated using TimeTree: Kumar et al., 2022).

In addition to differences in DEG numbers, class-specific DEGs differed markedly in their orthology distribution between insects and malacostracans (Figure 3). In the two malacostracans, class-specific DEGs were predominantly composed of malacostraca-only orthologous groups. In contrast, class-specific DEGs in insects included a substantial fraction of pancrustacea-shared orthologous groups. Across all species, species-specific DEGs were strongly enriched in lineage-specific and orphan genes, with the latter being particularly pronounced in insects, indicating that transcriptional novelty during ecdysis is primarily driven by lineage-specific genes rather than by conserved orthologous groups.

Gene co-expression analysis shows conserved gene expression pattern of moulting

Figure 4. Clusters of co-expressed genes in moulting. The strength of expression in *z*-scores (*Y*-axis) is illustrated for the different moult stage replicates per species where coloured lines represent the expression mean while grey lines represent individual gene expression values per replicate (*X*-axis). GO enrichment plots showing representative significant terms for each cluster are shown on the right side.

Most genes function within biological pathways or co-expression networks, and concerted expression changes of distinct gene sets can provide valuable insights into various co-regulated gene modules involved in moulting. We performed a co-expression analysis of all 5543 orthologues across all species and identified gene clusters with similar expression patterns (Figure 4). We recovered two major clusters that exhibit consistent expression across all samples: cluster 1 consists of genes that are down-regulated during the post-moult stage (Figure 4: top panel), and cluster 2 consists of genes that are up-regulated during the post-moult stage (Figure 4: bottom panel). A total of 774 co-expressed genes (14% of all orthologues) were identified across the five species (Supplementary Table 4). When the instar moult sample from *B. dorsalis* was included in the analysis, the number of co-expressed genes decreased to 184 genes (3% of all orthologues) (Figure 5).

Figure 5. Clusters of co-expressed genes in moulting (with *B. dorsalis* instar). The strength of expression in z-scores (*Y*-axis) is illustrated for the different moult stage replicates per species where coloured lines represent the expression mean while grey lines represent individual gene expression values per replicate (*X*-axis). GO enrichment plots showing representative significant terms for each cluster are shown on the right side.

Functional analysis of the 774 co-expressed genes shows that genes in cluster 1 are enriched in terms related to core metabolic and cellular processes, such as cellular

respiration and energy metabolism. Meanwhile, cluster 2 is enriched for terms related to tissue morphogenesis and developmental signalling, suggesting that the post-moult stage is characterized by the activation of pathways associated with cell proliferation, differentiation, and tissue remodelling, as expected. In addition to this we also recovered immune-related genes in cluster 2, including those known to mediate melanotic encapsulation of foreign targets, such as *zizimin-related* (*zir*), *ephexin* (*exn*), *Rho guanine nucleotide exchange factor 3* (*RhoGEF3*), and *Rho-related BTB domain-containing protein* (*RhoBTB*), and Toll and IMD signalling pathway genes including *Relish*, *supernumerary limbs* (*slmb*), *activating transcription factor-2* (*ATF-2*), *spatzle 3* (*spz3*), and *caspase* (*casp*).

The co-expressed gene sets which include the *B. dorsalis* instar-stage are not enriched for genes associated with cellular respiration or signalling pathways (Figure 5). Instead, for cluster 1 the enriched Gene Ontology terms were limited to processes such as protein glycosylation and carbohydrate metabolism. For cluster 2, we continued to recover terms related to tissue morphogenesis, sensory organ development, and muscle development. However, terms associated with key signalling pathways such as *Wnt*, *Hippo*, and *EGFR* were no longer recovered. This suggests that the metabolic and developmental requirements of instar moult in *B. dorsalis* might be different from other samples included in this analysis.

Moulting gene expression exhibits varying patterns of conservation

Figure 6. Transcriptome age index and moulting gene expression clusters in *L. vannamei*

Top panel shows the TAI of various moult stages in *L. vannamei* (*P* < 0.0001). Bottom panel shows the gene expression clusters of *L. vannamei* moult samples and their corresponding phylostratum composition.

Despite recovering a subset of genes that are co-expressed across all species, most genes did not show consistent co-expression patterns across species and groups. This suggests that moulting gene expression is for a large part lineage-specific, with gene expression dynamics varying between species. However, moulting as a process involves more than just the two timepoints around ecdysis. Thus, we expanded our comparison to include additional timepoints from pupariation in *D. melanogaster* and moulting in *L. vannamei*. We then examined transcriptome age patterns (TAI) across these timepoints. We found that conservation of gene expression is not evenly distributed across moulting in *L. vannamei* and larval-prepupal moult in *D. melanogaster*. For both species, TAI during the middle stages tends to be younger (i.e., more lineage-specific) showing an inverse hourglass pattern (Figures 6 and 7). However, it should be noted that while transcriptome age index (TAI) analysis is a widely used method to assess the evolutionary age of gene expression profiles across developmental stages it is also known to be sensitive to certain biases (Piasecka et al., 2013). One of the main challenges is that TAI relies on the mean expression values of genes, which can be skewed by a small number of very highly expressed genes such as housekeeping genes. These genes often have stable and abundant expression across conditions and may mask the contribution of biologically meaningful signals from genes with lower expression. As previously shown, removing or correcting for the influence of highly expressed genes can substantially alter the resulting TAI patterns, often revealing an additional layer of conservation pattern that were otherwise hidden (Piasecka et al., 2013), although this is moderated by the use of transformed expression values (J. Liu & Robinson-Rechavi, 2018). Moreover, the differences in TAI that we observe between stages are very small relative to the range of

possible values. Thus, we performed a co-expression analysis to identify clusters of genes with similar expression patterns.

We identified clusters of genes showing distinct trends across the moulting process (Figures 6 and 7). For each of these clusters we recovered varying patterns of conservation based on their Phylostratum composition. In *L. vannamei*, most of the gene clusters were consistent with TAI analysis. Clusters with higher expression in the middle of the moult process tend to be composed of younger genes, while those with lower expression during this phase are enriched for older, more conserved genes. For example, cluster 1, which shows peak expression in the late pre-moult stage, has a higher proportion of genes assigned to phylostrata dating back to the Pancrustacea node. In contrast, the relative contribution of more recent nodes, particularly those specific to Decapoda, increases toward the post-moult stages. On the other hand, cluster 4, which peaks in early post-moult, deviates from the overall pattern by being composed largely of older genes, suggesting a possible reactivation of conserved programmes following ecdysis. Although most gene clusters are dominated by younger phylostrata, especially during the middle of the moult cycle, it is worth noting the changes of phylostrata coming from Pancrustacea and Decapoda. This shift in phylostratum composition from a greater contribution of Pancrustacea-level genes in the late pre-moult stage to increased representation of Decapoda-level genes in the post-moult stage reflects the stage-specific recruitment of conserved and lineage-specific processes across the moulting process.

Younger gene clusters were enriched for genes involved in immune response, nervous system formation, sensory organ development, peroxidase activity, pigment binding, and

cuticle development (Supplementary Figure S2). Cuticle development is enriched in clusters 1 – 3, especially in cluster 1. We confirmed this with cuticleDB (Ioannidou et al., 2014; Magkrioti et al., 2004) (Supplementary Figure S3). Genes that are predominantly expressed in cluster 3 (post-moult) are enriched in carbonate dehydratase and pigment binding, which implies the stage specific expression of genes involved in calcification and pigmentation. Clusters with older phylostratum composition were enriched for genes related to extracellular matrix organization, fatty acid binding, and conserved metabolic processes such as steroid and carbohydrate metabolism, proteolysis, and lipid biosynthesis.

In *D. melanogaster*, we recovered ten gene expression clusters with varying temporal patterns (Figure 7; Supplementary Figure S4). The gene clusters that we recovered are largely not consistent with the TAI analysis, with mixed patterns (Figure 7). Most of the clusters exhibit a notable contribution of Diptera-specific genes, which is consistent with previous observations on the recruitment of younger genes in life stage transitions of flies (Remmel et al., 2025). Most clusters were dominated by older phylostrata and enriched in cellular and metabolic processes (Supplementary Figure S5). Cluster 1 is enriched in terms related to cilium movement, sperm flagellum assembly, cluster 2 is enriched in gamete development such as microtubule polymerization, spermatogenesis and spermatid differentiation while cluster 3 is enriched in RNA biogenesis, body morphogenesis, and cuticle development.

Figure 7. Transcriptome age index and moulting gene expression clusters in *D. melanogaster*

Top panel shows the TAI of various moult stages in *D. melanogaster* (*P* < 0.0001). Bottom panel shows the representative gene expression clusters of *D. melanogaster* samples and their corresponding phylostratum composition.

Discussion

Moulting is a highly conserved developmental process across Arthropoda. While a core genetic toolkit for moulting has been identified (Campli et al., 2024, 2025), largely derived from insect model systems, comparative genomic studies suggest that this "moulting genetic backbone" has undergone lineage-specific elaborations through gene gains and losses (Campli et al., 2025). The extent to which genes involved in moulting beyond this core toolkit are conserved or lineage-specific remains poorly known. In this study, we analysed gene expression datasets from insects and malacostracans to investigate the conservation of moulting gene expression in Pancrustacea.

We identified differentially expressed genes that are conserved between distant Pancrustacea species including genes involved in ecdysteroid metabolism. In addition, we recovered genes involved in cuticle development and in post-ecdysial processes. They notably include genes involved in the structural organization of the cuticle. For example *Knk*, regulated by *Rtv*, organizes chitin into laminae and protects it from chitinase-mediated degradation, along with chitin deacetylases known to direct proper cuticle organization (Chaudhari et al., 2011, 2013; Moussian, 2013). In the extracellular matrix, *Obst-A* binds the nascent chitin fibres produced by *Kkv* and recruits the *Serp/Knk* complex to the apical plasma membrane for chitin organization (Moussian, 2013; Ostrowski et al., 2002; Pesch et al., 2015; Petkau et al., 2012). This complex is essential for aligning the chitin into an ordered structure and for protecting the developing matrix from premature degradation by chitinases. Loss of these genes results in moulting defects, chitin disorganization and lethality (Chaudhari et al., 2011, 2013; Moussian, 2013; Ostrowski et al., 2002; Pesch et al., 2015; Petkau et al., 2012).

In addition to structural organization and chitin synthesis machinery, we also found genes involved in protein trafficking and membrane transport, that are essential for cuticle turnover. The enrichment of genes related to lysosomal activity, protein trafficking, and endocytosis during ecdysis are likely due to chitin synthesis and degradation. As the old cuticle is degraded, endocytosis enables the uptake of cuticle fragments and extracellular debris, which are processed in lysosomes to recover reusable monomers, while the secretory pathways direct newly synthesized cuticular proteins, chitin synthases, and other structural components to the apical surface for assembly of the new cuticle (Moussian et al., 2007). Consistent with this, we identified core secretion pathway genes such as *ghost* (*Gho*), *Haunted* (*Hau/Sec23*), *Sec24*, and other COAT proteins, which are known to be essential for apical plasma membrane remodelling during cuticle differentiation (Moussian et al., 2007; Norum et al., 2010). The multi-ligand endocytic receptor *Megalin* (*Mgl*) in *Drosophila* illustrates another trafficking-dependent mechanism, where *Mgl*-mediated endocytosis of Yellow protein restricts pigment deposition to the distal procuticle, ensuring proper spatial organization of cuticle components (Christensen & Willnow, 1999; Moestrup & Verroust, 2001; Riedel et al., 2011). In insects, similar roles for endocytosis have been reported. In *Locusta migratoria*, clathrin-mediated endocytosis supports resource recycling during cuticle degradation, and its disruption causes moulting defects, consistent with findings in *Helicoverpa armigera* where a G protein-coupled receptor mediates 20-hydroxyecdysone uptake via clathrin-dependent endocytosis (Kang et al., 2021; Shi et al., 2022). Evidence from *Caenorhabditis elegans* shows that disruption of endocytic trafficking in epidermal cells leads to moulting arrest, defective clearance of the old cuticle, failure to internalize

cargoes such as steroid hormone precursors, and disruption of plasma membrane homeostasis during new cuticle secretion (Lažetić & Fay, 2017; Yochem et al., 2015). The occurrence of such trafficking-dependent mechanisms in nematodes suggests that they represent an ancestral moulting machinery that predates Arthropoda.

Genes co-expressed during ecdysis are not limited to those directly involved in cuticle synthesis and organization. Instead, moulting also recruits evolutionarily conserved pathways supporting broader physiological functions, including metabolism, immune response, cellular differentiation, and tissue morphogenesis. Moulting is an energy-intensive process such that the caloric loss associated with shedding the exuviae is substantial (Harvey et al., 1980; Sweeney & Vannote, 1981). Additionally, insects cease feeding behaviours before, during, and after the moult (Ayres & MacLean, 1987; Kang et al., 2019; Rackauskas et al., 2006; Stamp & Bowers, 1994). Similarly, in decapod crustaceans feeding also declines during the pre-moult phase, stops entirely during ecdysis, and resumes only in the late post-moult period (Chan et al., 1988; Dall, 1986; Harpaz et al., 1987). This feeding behaviour reflects the mechanical and physiological challenges posed by exoskeleton shedding, particularly affecting essential feeding structures such as the buccal cavity, oesophagus, and anterior stomach (Lemos & Weissman, 2021). Previous studies indicate that ATP production is regulated by 20E signalling (Chang et al., 2022) and pre-moult is characterized by high energy metabolism which has been supported both at transcriptomic and proteomic level (Head et al., 2019). In various decapod crustaceans, the duration of pre-moult to post-moult can take a substantial amount of time during its moult cycle (Drach & Tchernigovtzeff, 1967; Gorissen

& Sandeman, 2022; Kim et al., 2024; Lemos & Weissman, 2021) and the expression of these genes ensures sufficient energy supply during ecdysis.

We also identified genes involved in the melanisation pathway, which plays a key role in cuticle hardening during the post-ecdysial period (Hiruma & Riddiford, 2009). Moreover, we recovered immune-related genes, including components of the Toll and IMD pathways. The melanisation pathway itself occupies an interesting intersection between cuticle maturation and innate immunity, functioning both in cuticle hardening and in pathogen defence (Beckage, 2011; Marmaras et al., 1996; Nakhleh et al., 2017; Vavricka et al., 2010). The vulnerability of arthropods to infection is particularly heightened after ecdysis, when the protective cuticle is incompletely formed. Whether the co-expression of immune genes alongside cuticle maturation reflects the vulnerability that accompanies moulting remains unclear, but previous studies suggest that immune activation may provide protective benefits during this period (Clark & Strand, 2013; Louradour et al., 2017; Tang, 2009; J. Zhang et al., 2014). The melanisation pathway is triggered by PAMPs, such as bacterial peptidoglycan or fungal β-glucans. This activates a protease cascade that converts proPO to active PO, which oxidizes phenols into quinones. These reactive intermediates polymerize to form melanin, which is deposited at sites of infection to encapsulate pathogens and generate cytotoxic reactive oxygen species (ROS) (Beckage, 2011; Nakhleh et al., 2017; Shan et al., 2023; Vavricka et al., 2010). Melanisation has been extensively studied in insects such as *D. melanogaster* and *Bombyx mori*, where it enhances survival during pathogenic infection (Louradour et al., 2017; J. Zhang et al., 2014), interacts with hemolymph coagulation (Bidla et al., 2005), and contributes to broader immune defence (Clark & Strand, 2013; Tang, 2009). Similar

immune-related responses during moulting have been observed in decapod crustaceans. In *S. paramamosain*, immune-related genes involved in antimicrobial peptide synthesis, Toll signalling, phenol oxidase system, and antioxidant pathways are significantly upregulated in the hepatopancreas during post-moult and inter-moult stages (Xu et al., 2020). Ortholog co-expression of these gene sets suggests that they play a conserved role during ecdysis.

The co-expression patterns that we observe differ significantly depending on life history strategy. When instar moult stages from holometabolous insect are included, we recover markedly fewer co-expressed genes, particularly those involved in signalling and differentiation pathways. The Hippo and Wnt signalling pathways are important regulators of tissue homeostasis and organ size that coordinate cell proliferation, differentiation and apoptosis (Halder & Johnson, 2011; Komiya & Habas, 2008; Wodarz & Nusse, 1998). In holometabolous insects, growth and differentiation are decoupled (Aldaz & Escudero, 2010; Chapman et al., 2013b; Manthey et al., 2024). During larval development, certain populations of cells, the imaginal primordia, remain in an undifferentiated, pluripotent state with latent embryonic potential and resume differentiation during pupal transition (Aldaz & Escudero, 2010; Chapman et al., 2013b). In holometabolous insects, moulting during the instar stages likely supports only growth and cuticle replacement. Extensive studies over the past decades have established that the transition between moulting and metamorphosis is regulated by a complex interplay of endogenous and environmental factors, involving hormonal feedback between ecdysteroids, *juvenile hormone* (*JH*), and stage-specifying transcription factors such as *Chinmo, Krüppel-homolog 1* (*Kr-h1*), *Broad*,

and *E93* (Suzuki et al., 2013; Truman, 2019; Truman et al., 2006; Truman & Riddiford, 2019, 2022). Gradual elaboration of this hormonal crosstalk might have led to the evolution of metamorphosis, by coupling moult cycles to environmental and physiological signals that determine whether an individual proceeds to another juvenile stage or transitions to the adult form. Consistent with this, our results suggest that the moulting programme can be decoupled from broader developmental processes depending on life history strategy.

Despite identifying a small set of genes which are co-expressed across all five species, most of the genes that are differentially expressed during moulting are lineage-specific, either because they do not have orthologs outside of one lineage, or because the differential expression pattern is restricted to one lineage even though the orthologs are more broadly distributed (Figure 3). Time-series analysis shows that these lineage-specific signatures are not evenly distributed across the moulting process. Mid-moulting gene clusters are enriched for relatively younger genes. In *L. vannamei,* these clusters are primarily involved in the development of sensory structures, nervous system formation, pigment binding, and chitin-based extracellular matrix components. In contrast, older gene clusters are enriched for functions such as steroid metabolic processes, chitin catabolism, and carbohydrate metabolism. This pattern suggests that mid-moult lineage-specific signatures are driven in part by the development of the cuticle and specialized structures of the exoskeleton.

The cuticular surface is remarkably diverse, featuring a wide array of pigmentation and specialized structures (Hallberg & Skog, 2010; Mellon, 2012). This includes various setal structures which line the exoskeleton and are periodically developed with each moult

(Drach & Tchernigovtzeff, 1967; Gorissen & Sandeman, 2022; Kim et al., 2024). Many of these are innervated by sensory neurons which may house mechanoreceptive and chemoreceptive neurons, or both (Hallberg & Skog, 2010; Mellon, 2012; Shields et al., 2008). Studies combining physiological analysis with scanning electron microscopy have shown that many cuticular setae on locomotory, feeding, and grooming appendages are bimodal in function, containing both chemo- and mechanoreceptor cells (Hallberg & Skog, 2010; Shields et al., 2008). These structures often serve species-specific roles, including mating behaviour, mechano-sensation, and predator detection. Their diversity and functional specificity across lineages likely explain the recruitment of younger, lineage-specific genes in their regeneration during moulting.

Lineage specific genes have been implicated in species-specific adaptations for behaviour, environmental factors, reproduction, host pathogen interactions, and speciation (Han et al., 2024; Khalturin et al., 2008, 2009). While it is intuitive to conclude that lineage-specific signatures of moulting are due to adaptive divergence, some studies have shown that lineage-specific genes are not always adaptive. Young, lineage-specific genes can be gained, lost, or integrated into gene regulatory networks through stochastic processes, and may persist without signatures of adaptation (Palmieri et al., 2014; Vakirlis et al., 2022; Wacholder et al., 2023). These lineage specific genes thus present interesting candidates for future tests of adaptive evolution in arthropods.

While the overall transcriptome age pattern is similar between *L. vannamei* and *D. melanogaster*, the underlying gene sets driving these patterns differ substantially. In *L. vannamei*, the mid-moult peak in younger gene expression is primarily driven by genes involved in cuticle development, whereas in *Drosophila*, the same pattern is largely

associated with the expression of reproductive genes. Interestingly, in *Drosophila*, gene modules involved with cuticle development are also composed of younger phylostrata, but they are predominantly expressed earlier during pupariation, rather than in mid-moult as in *L. vannamei*. This heterochrony in cuticle gene expression is likely related to the evolution of the puparium in cyclorrhaphous flies, which retain the larval cuticle as a protective case rather than shedding it during ecdysis. Heterochrony in cuticle development and maturation has been linked to adaptation in other arthropods (Elias-Neto et al., 2014; Falcon et al., 2019) which suggests that timing of cuticle formation can evolve in response to different life history and ecological needs.

Although moulting is regulated by conserved hormonal pathways, our findings suggest that it coordinates both deeply conserved and recently evolved gene modules. The precise coordination of these gene modules reflects a regulatory flexibility that enables moulting to accommodate lineage-specific physiological and developmental requirements. It is likely that the regulatory architecture underlying moulting allows for co-expression of genes not directly tied to the moulting process itself but essential for broad physiological support and reconstructing complex structure of the exoskeleton. Since the exoskeleton forms the primary interface between the organism and its environment, environmental pressures may contribute to its morphology and the gene expression programmes involved in its formation. Divergence time and environmental heterogeneity have been shown to influence convergence in both gene expression and morphological traits (Boisseau et al., 2025; Buckley et al., 2009; Chaturvedi et al., 2022). Testing whether ecological niche similarity predicts convergence in ecdysis gene expression would therefore be an interesting task for future work.

By uncovering both conserved and lineage-specific components of moulting gene expression, this study provides evidence for both evolutionary conservation and divergence of this deeply shared developmental process. This conservation of the moulting programme appears to extend to pupariation. These findings suggest that moulting may involve a broader range of developmental processes than its classical definition of periodic shedding of the exoskeleton which may encompass diverse moulting strategies exhibited by Arthropods, ranging from puparium formation to, potentially, apolysis without ecdysis, as observed in paedomorphic insects such as Strepsipterans (Kathirithamby et al., 1984; Manfredini et al., 2007). In a broader perspective, moulting can be viewed as a developmental process that enables growth and morphological transformation through *shedding or reconstruction* of the exoskeleton. Overall, our findings contribute to a broader understanding of the evolutionary dynamics underlying moulting and highlights the modular architecture of the moulting programme, which may be central to understanding how the vast diversity of arthropod species has adapted their post-embryonic development to different ecological and life-history contexts.

References

Abu-Jamous, B., & Kelly, S. (2018). Clust: Automatic extraction of optimal co-expressed gene

clusters from gene expression data. *Genome Biology*, *19*(1), 172.

https://doi.org/10.1186/s13059-018-1536-8

Aldaz, S., & Escudero, L. M. (2010). Imaginal discs. *Current Biology*, *20*(10), R429–R431.

https://doi.org/10.1016/j.cub.2010.03.010

Asano, T., Seto, Y., Hashimoto, K., & Kurushima, H. (2019). Mini-review an insect-specific system

for terrestrialization: Laccase-mediated cuticle formation. *Insect Biochemistry and*

Molecular Biology, *108*, 61–70. https://doi.org/10.1016/j.ibmb.2019.03.007

Ashburner, M., Ball, C. A., Blake, J. A., Botstein, D., Butler, H., Cherry, J. M., Davis, A. P., Dolinski,

K., Dwight, S. S., Eppig, J. T., Harris, M. A., Hill, D. P., Issel-Tarver, L., Kasarskis, A., Lewis,

S., Matese, J. C., Richardson, J. E., Ringwald, M., Rubin, G. M., & Sherlock, G. (2000).

Gene Ontology: Tool for the unification of biology. *Nature Genetics*, *25*(1), 25–29.

https://doi.org/10.1038/75556

Ayres, M. P., & MacLean, S. F. (1987). Molt as a Component of Insect Development: Galerucella

sagittariae (Chrysomelidae) and Epirrita autumnata (Geometridae). *Oikos*, *48*(3), 273.

https://doi.org/10.2307/3565514

Barrera-Redondo, J., Lotharukpong, J. S., Drost, H.-G., & Coelho, S. M. (2023). Uncovering gene-

family founder events during major evolutionary transitions in animals, plants and fungi

using GenEra. *Genome Biology*, *24*(1), 54. https://doi.org/10.1186/s13059-023-02895-z

Beckage, N. E. (2011). *Insect Immunology*. Academic Press.

Belles, X. (2020). Molting: The basis for growing and for changing the form. In *Insect Metamorphosis* (pp. 199–215). Elsevier. https://doi.org/10.1016/B978-0-12-813020-9.00009-0

Benjamini, Y., & Hochberg, Y. (1995). Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society Series B: Statistical Methodology*, *57*(1), 289–300. https://doi.org/10.1111/j.2517-6161.1995.tb02031.x

Bernot, J. P., Owen, C. L., Wolfe, J. M., Meland, K., Olesen, J., & Crandall, K. A. (2023). Major Revisions in Pancrustacean Phylogeny and Evidence of Sensitivity to Taxon Sampling. *Molecular Biology and Evolution*, *40*(8), msad175. https://doi.org/10.1093/molbev/msad175

Bidla, G., Lindgren, M., Theopold, U., & Dushay, M. (2005). Hemolymph coagulation and phenoloxidase in larvae. *Developmental & Comparative Immunology*, *29*(8), 669–679. https://doi.org/10.1016/j.dci.2004.11.007

Boisseau, R. P., Bradler, S., & Emlen, D. J. (2025). Divergence time and environmental similarity predict the strength of morphological convergence in stick and leaf insects. *Proceedings of the National Academy of Sciences*, *122*(1), e2319485121. https://doi.org/10.1073/pnas.2319485121

Bolger, A. M., Lohse, M., & Usadel, B. (2014). Trimmomatic: A flexible trimmer for Illumina sequence data. *Bioinformatics*, *30*(15), 2114–2120. https://doi.org/10.1093/bioinformatics/btu170

Brawand, D., Soumillon, M., Necsulea, A., Julien, P., Csárdi, G., Harrigan, P., Weier, M., Liechti, A., Aximu-Petri, A., Kircher, M., Albert, F. W., Zeller, U., Khaitovich, P., Grützner, F., Bergmann, S., Nielsen, R., Pääbo, S., & Kaessmann, H. (2011). The evolution of gene expression levels in mammalian organs. *Nature*, *478*(7369), 343–348. https://doi.org/10.1038/nature10532

Bray, N. L., Pimentel, H., Melsted, P., & Pachter, L. (2016). Near-optimal probabilistic RNA-seq quantification. *Nature Biotechnology*, *34*(5), 525–527. https://doi.org/10.1038/nbt.3519

Buchfink, B., Xie, C., & Huson, D. H. (2015). Fast and sensitive protein alignment using DIAMOND. *Nature Methods*, *12*(1), 59–60. https://doi.org/10.1038/nmeth.3176

Buckley, T. R., Attanayake, D., & Bradler, S. (2009). Extreme convergence in stick insect evolution: Phylogenetic placement of the Lord Howe Island tree lobster. *Proceedings of the Royal Society B: Biological Sciences*, *276*(1659), 1055–1062. https://doi.org/10.1098/rspb.2008.1552

Cabrera, A. R., Shirk, P. D., Evans, J. D., Hung, K., Sims, J., Alborn, H., & Teal, P. E. A. (2015). Three H alloween genes from the V arroa mite, *V arroa destructor* (A nderson & T rueman) and their expression during reproduction. *Insect Molecular Biology*, *24*(3), 277–292. https://doi.org/10.1111/imb.12155

Campli, G., Joye, S., Volovych, O., Novikov, A., Chipman, A. D., Robinson-Rechavi, M., & Waterhouse, R. M. (2025). *Evolutionary dynamics of the arthropod moulting machinery*. Evolutionary Biology. https://doi.org/10.64898/2025.12.24.696370

Campli, G., Volovych, O., Kim, K., Veldsman, W. P., Drage, H. B., Sheizaf, I., Lynch, S., Chipman, A. D., Daley, A. C., Robinson-Rechavi, M., & Waterhouse, R. M. (2024). The moulting

arthropod: A complete genetic toolkit review. *Biological Reviews*, brv.13123.

https://doi.org/10.1111/brv.13123

Chan, S.-M., Rankin, S. M., & Keeley, L. L. (1988). Characterization of the Molt Stages in *Penaeus vannamei*: Setogenesis and Hemolymph Levels of Total Protein, Ecdysteroids, and Glucose. *The Biological Bulletin*, *175*(2), 185–192. https://doi.org/10.2307/1541558

Chang, Y., Zhang, B., Du, M., Geng, Z., Wei, J., Guan, R., An, S., & Zhao, W. (2022). The vital hormone 20-hydroxyecdysone controls ATP production by upregulating binding of trehalase 1 with ATP synthase subunit α in Helicoverpa armigera. *Journal of Biological Chemistry*, *298*(2), 101565. https://doi.org/10.1016/j.jbc.2022.101565

Chapman, R. F., Simpson, S. J., & Douglas, A. E. (2013a). *The Insects: Structure and Function*. Cambridge University Press. https://books.google.ch/books?id=NXJEi8fo7CkC

Chapman, R. F., Simpson, S. J., & Douglas, A. E. (2013b). *The insects: Structure and function* (Fifth edition). Cambridge University Press.

Chaturvedi, S., Gompert, Z., Feder, J. L., Osborne, O. G., Muschick, M., Riesch, R., Soria-Carrasco, V., & Nosil, P. (2022). Climatic similarity and genomic background shape the extent of parallel adaptation in Timema stick insects. *Nature Ecology & Evolution*, *6*(12), 1952–1964. https://doi.org/10.1038/s41559-022-01909-6

Chaudhari, S. S., Arakane, Y., Specht, C. A., Moussian, B., Boyle, D. L., Park, Y., Kramer, K. J., Beeman, R. W., & Muthukrishnan, S. (2011). Knickkopf protein protects and organizes chitin in the newly synthesized insect exoskeleton. *Proceedings of the National Academy of Sciences*, *108*(41), 17028–17033. https://doi.org/10.1073/pnas.1112288108

Chaudhari, S. S., Arakane, Y., Specht, C. A., Moussian, B., Kramer, K. J., Muthukrishnan, S., & Beeman, R. W. (2013). Retroactive Maintains Cuticle Integrity by Promoting the Trafficking of Knickkopf into the Procuticle of Tribolium castaneum. *PLoS Genetics*, *9*(1), e1003268. https://doi.org/10.1371/journal.pgen.1003268

Cheong, S. P. S., Huang, J., Bendena, W. G., Tobe, S. S., & Hui, J. H. L. (2015). Evolution of Ecdysis and Metamorphosis in Arthropods: The Rise of Regulation of Juvenile Hormone. *Integrative and Comparative Biology*, *55*(5), 878–890. https://doi.org/10.1093/icb/icv066

Christensen, E. I., & Willnow, T. E. (1999). Essential Role of Megalin in Renal Proximal Tubule for Vitamin Homeostasis. *Journal of the American Society of Nephrology*, *10*(10), 2224– 2236. https://doi.org/10.1681/ASN.V10102224

Clark, K. D., & Strand, M. R. (2013). Hemolymph Melanization in the Silkmoth Bombyx mori Involves Formation of a High Molecular Mass Complex That Metabolizes Tyrosine. *Journal of Biological Chemistry*, *288*(20), 14476–14487. https://doi.org/10.1074/jbc.M113.459222

Daley, A. C., Antcliffe, J. B., Drage, H. B., & Pates, S. (2018). Early fossil record of Euarthropoda and the Cambrian Explosion. *Proceedings of the National Academy of Sciences*, *115*(21), 5323–5331. https://doi.org/10.1073/pnas.1719962115

Daley, A. C., & Drage, H. B. (2016). The fossil record of ecdysis, and trends in the moulting behaviour of trilobites. *Arthropod Structure & Development*, *45*(2), 71–96. https://doi.org/10.1016/j.asd.2015.09.004

Dall, W. (1986). Estimation of routine metabolic rate in a penaeid prawn, Penaeus esculentus

    Haswell. *Journal of Experimental Marine Biology and Ecology*, *96*(1), 57–74.

    https://doi.org/10.1016/0022-0981(86)90013-4

Dermauw, W., Van Leeuwen, T., & Feyereisen, R. (2020). Diversity and evolution of the P450

    family in arthropods. *Insect Biochemistry and Molecular Biology*, *127*, 103490.

    https://doi.org/10.1016/j.ibmb.2020.103490

Domazet-Lošo, T., & Tautz, D. (2010). A phylogenetically based transcriptome age index mirrors

    ontogenetic divergence patterns. *Nature*, *468*(7325), 815–818.

    https://doi.org/10.1038/nature09632

Drach, P., & Tchernigovtzeff, C. (1967). Sur la méthode de détermination des stades d'intermue

    et son application générale aux crustacés. *Vie et Milieu*, 595–610.

Drost, H.-G., Gabel, A., Liu, J., Quint, M., & Grosse, I. (2018). myTAI: Evolutionary transcriptomics

    with R. *Bioinformatics*, *34*(9), 1589–1590.

    https://doi.org/10.1093/bioinformatics/btx835

Elias-Neto, M., Nascimento, A. L. O., Bonetti, A. M., Nascimento, F. S., Mateus, S., Garófalo, C. A.,

    & Bitondi, M. M. G. (2014). Heterochrony of cuticular differentiation in eusocial

    corbiculate bees. *Apidologie*, *45*(4), 397–408. https://doi.org/10.1007/s13592-013-0254-

Enya, S., Daimon, T., Igarashi, F., Kataoka, H., Uchibori, M., Sezutsu, H., Shinoda, T., & Niwa, R.

    (2015). The silkworm glutathione S-transferase gene noppera-bo is required for

    ecdysteroid biosynthesis and larval development. *Insect Biochemistry and Molecular*

    Biology, *61*, 1–7. https://doi.org/10.1016/j.ibmb.2015.04.001

Falcon, T., Pinheiro, D. G., Ferreira-Caliman, M. J., Turatti, I. C. C., Abreu, F. C. P. D., Galaschi-Teixeira, J. S., Martins, J. R., Elias-Neto, M., Soares, M. P. M., Laure, M. B., Figueiredo, V. L. C., Lopes, N. P., Simões, Z. L. P., Garófalo, C. A., & Bitondi, M. M. G. (2019). Exploring integument transcriptomes, cuticle ultrastructure, and cuticular hydrocarbons profiles in eusocial and solitary bee species displaying heterochronic adult cuticle maturation. *PLOS ONE*, *14*(3), e0213796. https://doi.org/10.1371/journal.pone.0213796

Feron, R., & Waterhouse, R. M. (2022). Assessing species coverage and assembly quality of rapidly accumulating sequenced genomes. *GigaScience*, *11*, giac006. https://doi.org/10.1093/gigascience/giac006

Fujii-Taira, I., Yamaguchi, S., Iijima, R., Natori, S., & Homma, K. J. (2009). Suppression of the ecdysteroid-triggered growth arrest by a novel *Drosophila* membrane steroid binding protein. *FEBS Letters*, *583*(4), 655–660. https://doi.org/10.1016/j.febslet.2008.12.056

Gao, Y., Zhang, X., Wei, J., Sun, X., Yuan, J., Li, F., & Xiang, J. (2015a). Whole Transcriptome Analysis Provides Insights into Molecular Mechanisms for Molting in Litopenaeus vannamei. *PLOS ONE*, *10*(12), e0144350. https://doi.org/10.1371/journal.pone.0144350

Gao, Y., Zhang, X., Wei, J., Sun, X., Yuan, J., Li, F., & Xiang, J. (2015b). Whole Transcriptome Analysis Provides Insights into Molecular Mechanisms for Molting in Litopenaeus vannamei. *PLOS ONE*, *10*(12), e0144350. https://doi.org/10.1371/journal.pone.0144350

Giribet, G., & Edgecombe, G. D. (2019). The Phylogeny and Evolutionary History of Arthropods. *Current Biology*, *29*(12), R592–R602. https://doi.org/10.1016/j.cub.2019.04.057

Gorissen, S., & Sandeman, D. (2022). Moult cycle staging in decapod crustaceans (Pleocyemata)

and the Australian crayfish, Cherax destructor Clark, 1936 (Decapoda, Parastacidae).

Crustaceana, *95*(2), 165–217. https://doi.org/10.1163/15685403-bja10180

Graveley, B. R., Brooks, A. N., Carlson, J. W., Duff, M. O., Landolin, J. M., Yang, L., Artieri, C. G.,

Van Baren, M. J., Boley, N., Booth, B. W., Brown, J. B., Cherbas, L., Davis, C. A., Dobin, A.,

Li, R., Lin, W., Malone, J. H., Mattiuzzo, N. R., Miller, D., … Celniker, S. E. (2011). The

developmental transcriptome of Drosophila melanogaster. *Nature*, *471*(7339), 473–479.

https://doi.org/10.1038/nature09715

Halder, G., & Johnson, R. L. (2011). Hippo signaling: Growth control and beyond. *Development*,

138(1), 9–22. https://doi.org/10.1242/dev.045500

Hallberg, E., & Skog, M. (2010). Chemosensory Sensilla in Crustaceans. In T. Breithaupt & M.

Thiel (Eds.), *Chemical Communication in Crustaceans* (pp. 103–121). Springer New York.

https://doi.org/10.1007/978-0-387-77101-4_6

Han, Z., Wang, Z., Rittschof, D., Huang, Z., Chen, L., Hao, H., Yao, S., Su, P., Huang, M., Zhang, Y.-

Y., Ke, C., & Feng, D. (2024). New genes helped acorn barnacles adapt to a sessile

lifestyle. *Nature Genetics*, *56*(5), 970–981. https://doi.org/10.1038/s41588-024-01733-7

Harpaz, S., Kahan, D., Galun, R., & Moore, I. (1987). Responses of freshwater

prawn,Macrobrachium rosenbergii, to chemical attractants. *Journal of Chemical Ecology*,

13(9), 1957–1965. https://doi.org/10.1007/BF01014678

Harrison, P. W., Wright, A. E., & Mank, J. E. (2012). The evolution of gene expression and the

transcriptome–phenotype relationship. *Seminars in Cell & Developmental Biology*, *23*(2),

222–229. https://doi.org/10.1016/j.semcdb.2011.12.004

Harvey, R. S., Vannote, R. L., & Sweeney, B. W. (1980). Life History, Developmental Processes, and Energetics of the Burrowing Mayfly Dolania Americana. In J. F. Flannagan & K. E. Marshall (Eds.), *Advances in Ephemeroptera Biology* (pp. 211–230). Springer US. https://doi.org/10.1007/978-1-4613-3066-0_18

Head, T. B., Mykles, D. L., & Tomanek, L. (2019). Proteomic analysis of the crustacean molting gland (Y-organ) over the course of the molt cycle. *Comparative Biochemistry and Physiology Part D: Genomics and Proteomics*, *29*, 193–210. https://doi.org/10.1016/j.cbd.2018.11.011

Hiruma, K., & Riddiford, L. M. (2009). The molecular mechanisms of cuticular melanization: The ecdysone cascade leading to dopa decarboxylase expression in Manduca sexta. *Insect Biochemistry and Molecular Biology*, *39*(4), 245–253. https://doi.org/10.1016/j.ibmb.2009.01.008

Huang, X., Suyama, K., Buchanan, J., Zhu, A. J., & Scott, M. P. (2005). A *Drosophila* model of the Niemann-Pick type C lysosome storage disease: *Dnpc1a* is required for molting and sterol homeostasis. *Development*, *132*(22), 5115–5124. https://doi.org/10.1242/dev.02079

i5K Consortium. (2013). The i5K Initiative: Advancing Arthropod Genomics for Knowledge, Human Health, Agriculture, and the Environment. *Journal of Heredity*, *104*(5), 595–600. https://doi.org/10.1093/jhered/est050

Ioannidou, Z. S., Theodoropoulou, M. C., Papandreou, N. C., Willis, J. H., & Hamodrakas, S. J. (2014). CutProtFam-Pred: Detection and classification of putative structural cuticular proteins from sequence alone, based on profile Hidden Markov Models. *Insect*

Biochemistry and Molecular Biology, *52*, 51–59.

https://doi.org/10.1016/j.ibmb.2014.06.004

Kang, X.-L., Li, Y.-X., Li, Y.-L., Wang, J.-X., & Zhao, X.-F. (2021). The homotetramerization of a GPCR

transmits the 20-hydroxyecdysone signal and increases its entry into cells for insect

metamorphosis. *Development*, *148*(5), dev196667. https://doi.org/10.1242/dev.196667

Kang, X.-L., Zhang, J.-Y., Wang, D., Zhao, Y.-M., Han, X.-L., Wang, J.-X., & Zhao, X.-F. (2019). The

steroid hormone 20-hydroxyecdysone binds to dopamine receptor to repress

lepidopteran insect feeding and promote pupation. *PLOS Genetics*, *15*(8), e1008331.

https://doi.org/10.1371/journal.pgen.1008331

Kathirithamby, J., Smith, D. S., Lomas, M. B., & Luke, B. M. (1984). Apolysis without ecdysis in

larval development of a strepsipteran, Elenchus tenuicornis (Kirby). *Zoological Journal of

the Linnean Society*, *82*(3), 335–343. https://doi.org/10.1111/j.1096-

3642.1984.tb00647.x

Keidar Haran, T., & Keren, L. (2022). From genes to modules, from cells to ecosystems.

Molecular Systems Biology, *18*(5), e10726. https://doi.org/10.15252/msb.202110726

Khalturin, K., Anton-Erxleben, F., Sassmann, S., Wittlieb, J., Hemmrich, G., & Bosch, T. C. G.

(2008). A Novel Gene Family Controls Species-Specific Morphological Traits in Hydra.

PLoS Biology, *6*(11), e278. https://doi.org/10.1371/journal.pbio.0060278

Khalturin, K., Hemmrich, G., Fraune, S., Augustin, R., & Bosch, T. C. G. (2009). More than just

orphans: Are taxonomically-restricted genes important in evolution? *Trends in Genetics*,

25(9), 404–413. https://doi.org/10.1016/j.tig.2009.07.006

Kim, K., Antcliffe, J., Daley, A. C., & Robinson-Rechavi, M. (2024). Moult cycle and setal development of the Atlantic ditch shrimp *Palaemon varians* Leach, 1814 (Decapoda: Caridea: Palaemonidae). *Journal of Crustacean Biology*, *44*(4), ruae075. https://doi.org/10.1093/jcbiol/ruae075

Koiwai, K., Inaba, K., Morohashi, K., Enya, S., Arai, R., Kojima, H., Okabe, T., Fujikawa, Y., Inoue, H., Yoshino, R., Hirokawa, T., Kato, K., Fukuzawa, K., Shimada-Niwa, Y., Nakamura, A., Yumoto, F., Senda, T., & Niwa, R. (2020). An integrated approach to unravel a crucial structural property required for the function of the insect steroidogenic Halloween protein Noppera-bo. *Journal of Biological Chemistry*, *295*(20), 7154–7167. https://doi.org/10.1074/jbc.RA119.011463

Komiya, Y., & Habas, R. (2008). Wnt signal transduction pathways. *Organogenesis*, *4*(2), 68–75. https://doi.org/10.4161/org.4.2.5851

Kumar, S., Suleski, M., Craig, J. M., Kasprowicz, A. E., Sanderford, M., Li, M., Stecher, G., & Hedges, S. B. (2022). TimeTree 5: An Expanded Resource for Species Divergence Times. *Molecular Biology and Evolution*, *39*(8), msac174. https://doi.org/10.1093/molbev/msac174

Kuznetsov, D., Tegenfeldt, F., Manni, M., Seppey, M., Berkeley, M., Kriventseva, E. V., & Zdobnov, E. M. (2023). OrthoDB v11: Annotation of orthologs in the widest sampling of organismal diversity. *Nucleic Acids Research*, *51*(D1), D445–D451. https://doi.org/10.1093/nar/gkac998

Lažetić, V., & Fay, D. S. (2017). Molting in *C. elegans*. *Worm*, *6*(1), e1330246. https://doi.org/10.1080/21624054.2017.1330246

Lemos, D., & Weissman, D. (2021). Moulting in the grow-out of farmed shrimp: A review.

    Reviews in Aquaculture, *13*(1), 5–17. https://doi.org/10.1111/raq.12461

Liu, J., & Robinson-Rechavi, M. (2018). Developmental Constraints on Genome Evolution in Four

    Bilaterian Model Species. *Genome Biology and Evolution*, *10*(9), 2266–2277.

    https://doi.org/10.1093/gbe/evy177

Liu, L., Liu, X., Fu, Y., Fang, W., & Wang, C. (2022). Whole-body transcriptome analysis provides

    insights into the cascade of sequential expression events involved in growth, immunity,

    and metabolism during the molting cycle in Scylla paramamosain. *Scientific Reports*,

    12(1), 11395. https://doi.org/10.1038/s41598-022-14783-w

Liu, S., Xia, Y., Zhang, Q., Li, W., Li, R., Liu, Y., Chen, E., Dou, W., Stelinski, L. L., & Wang, J. (2020).

    Potential targets for controlling Bᴀᴄᴛʀᴏᴄᴇʀᴀ ᴅᴏʀsᴀʟɪs using cuticle- and hormone-related

    genes revealed by a developmental transcriptome analysis. *Pest Management Science*,

    76(6), 2127–2143. https://doi.org/10.1002/ps.5751

Louradour, I., Sharma, A., Morin-Poulard, I., Letourneau, M., Vincent, A., Crozatier, M., & Vanzo,

    N. (2017). Reactive oxygen species-dependent Toll/NF-κB activation in the Drosophila

    hematopoietic niche confers resistance to wasp parasitism. *eLife*, *6*, e25496.

    https://doi.org/10.7554/eLife.25496

Love, M. I., Huber, W., & Anders, S. (2014). Moderated estimation of fold change and dispersion

    for RNA-seq data with DESeq2. *Genome Biology*, *15*(12), 550.

    https://doi.org/10.1186/s13059-014-0550-8

Magkrioti, C. K., Spyropoulos, I. C., Iconomidou, V. A., Willis, J. H., & Hamodrakas, S. J. (2004). cuticleDB: A relational database of Arthropod cuticular proteins. *BMC Bioinformatics*, *5*(1), 138. https://doi.org/10.1186/1471-2105-5-138

Manfredini, F., Giusti, F., Beani, L., & Dallai, R. (2007). Developmental strategy of the endoparasite *Xenos vesparum* (strepsiptera, Insecta): Host invasion and elusion of its defense reactions. *Journal of Morphology*, *268*(7), 588–601. https://doi.org/10.1002/jmor.10540

Manthey, C., Metcalf, C. J. E., Monaghan, M. T., Steiner, U. K., & Rolff, J. (2024). Rapid growth and the evolution of complete metamorphosis in insects. *Proceedings of the National Academy of Sciences*, *121*(38), e2402980121. https://doi.org/10.1073/pnas.2402980121

Marmaras, V. J., Charalambidis, N. D., & Zervas, C. G. (1996). Immune response in insects: The role of phenoloxidase in defense reactions in relation to melanization and sclerotization. *Archives of Insect Biochemistry and Physiology*, *31*(2), 119–133. https://doi.org/10.1002/(SICI)1520-6327(1996)31:2%253C119::AID-ARCH1%253E3.0.CO;2-V

Mazzoni, C. J., Ciofi, C., & Waterhouse, R. M. (2023). Biodiversity: An atlas of European reference genomes. *Nature*, *619*(7969), 252–252. https://doi.org/10.1038/d41586-023-02229-w

Mellon, D. (2012). Smelling, feeling, tasting and touching: Behavioral and neural integration of antennular chemosensory and mechanosensory inputs in the crayfish. *Journal of Experimental Biology*, *215*(13), 2163–2172. https://doi.org/10.1242/jeb.069492

Moestrup, S. K., & Verroust, P. J. (2001). M EGALIN—AND C UBILIN -M EDIATED E NDOCYTOSIS

OF P ROTEIN -B OUND V ITAMINS , L IPIDS , AND H ORMONES IN P OLARIZED E PITHELIA.

Annual Review of Nutrition, *21*(1), 407–428.

https://doi.org/10.1146/annurev.nutr.21.1.407

Moussian, B. (2013). The apical plasma membrane of chitin-synthesizing epithelia. *Insect*

Science, *20*(2), 139–146. https://doi.org/10.1111/j.1744-7917.2012.01549.x

Moussian, B., Letizia, A., Martínez-Corrales, G., Rotstein, B., Casali, A., & Llimargas, M. (2015).

Deciphering the Genetic Programme Triggering Timely and Spatially-Regulated Chitin

Deposition. *PLOS Genetics*, *11*(1), e1004939.

https://doi.org/10.1371/journal.pgen.1004939

Moussian, B., Veerkamp, J., Müller, U., & Schwarz, H. (2007). Assembly of the Drosophila larval

exoskeleton requires controlled secretion and shaping of the apical plasma membrane.

Matrix Biology, *26*(5), 337–347. https://doi.org/10.1016/j.matbio.2007.02.001

Mykles, D. L. (2011). Ecdysteroid metabolism in crustaceans. *The Journal of Steroid Biochemistry*

and Molecular Biology, *127*(3–5), 196–203.

https://doi.org/10.1016/j.jsbmb.2010.09.001

Mykles, D. L. (2021). Signaling Pathways That Regulate the Crustacean Molting Gland. *Frontiers*

in Endocrinology, *12*, 674711. https://doi.org/10.3389/fendo.2021.674711

Mykles, D. L., & Chang, E. S. (2020). Hormonal control of the crustacean molting gland: Insights

from transcriptomics and proteomics. *General and Comparative Endocrinology*, *294*,

113493. https://doi.org/10.1016/j.ygcen.2020.113493

Nakaoka, T., Iga, M., Yamada, T., Koujima, I., Takeshima, M., Zhou, X., Suzuki, Y., Ogihara, M. H.,

    & Kataoka, H. (2017). Deep sequencing of the prothoracic gland transcriptome reveals

    new players in insect ecdysteroidogenesis. *PLOS ONE*, *12*(3), e0172951.

    https://doi.org/10.1371/journal.pone.0172951

Nakhleh, J., El Moussawi, L., & Osta, M. A. (2017). The Melanization Response in Insect

    Immunity. In *Advances in Insect Physiology* (Vol. 52, pp. 83–109). Elsevier.

    https://doi.org/10.1016/bs.aiip.2016.11.002

Norum, M., Tång, E., Chavoshi, T., Schwarz, H., Linke, D., Uv, A., & Moussian, B. (2010).

    Trafficking through COPII Stabilises Cell Polarity and Drives Secretion during Drosophila

    Epidermal Differentiation. *PLoS ONE*, *5*(5), e10802.

    https://doi.org/10.1371/journal.pone.0010802

Oleksiak, M. F., Churchill, G. A., & Crawford, D. L. (2002). Variation in gene expression within and

    among natural populations. *Nature Genetics*, *32*(2), 261–266.

    https://doi.org/10.1038/ng983

Ono, H., Rewitz, K. F., Shinoda, T., Itoyama, K., Petryk, A., Rybczynski, R., Jarcho, M., Warren, J. T.,

    Marqués, G., Shimell, M. J., Gilbert, L. I., & O'Connor, M. B. (2006). Spook and Spookier

    code for stage-specific components of the ecdysone biosynthetic pathway in Diptera.

    Developmental Biology, *298*(2), 555–570. https://doi.org/10.1016/j.ydbio.2006.07.023

Ostrowski, S., Dierick, H. A., & Bejsovec, A. (2002). Genetic Control of Cuticle Formation During

    Embryonic Development of *Drosophila melanogaster*. *Genetics*, *161*(1), 171–182.

    https://doi.org/10.1093/genetics/161.1.171

Ou, Q., Magico, A., & King-Jones, K. (2011). Nuclear Receptor DHR4 Controls the Timing of

    Steroid Hormone Pulses During Drosophila Development. *PLoS Biology*, *9*(9), e1001160.

    https://doi.org/10.1371/journal.pbio.1001160

Ozerova, A. M., Kulikova, D. A., Evgen'ev, M. B., & Gelfand, M. S. (2025). Temporal Dynamics of

    Gene Expression During Metamorphosis in Two Distant *Drosophila S* pecies. *Genome*

    Biology and Evolution, *17*(6), evaf100. https://doi.org/10.1093/gbe/evaf100

Palmieri, N., Kosiol, C., & Schlötterer, C. (2014). The life cycle of Drosophila orphan genes. *eLife*,

    3, e01311. https://doi.org/10.7554/eLife.01311

Perry, C. L., & Robin, C. (2025). Before the Black Box. *bioRxiv*.

    https://doi.org/10.1101/2025.04.14.648839

Pesch, Y.-Y., Riedel, D., & Behr, M. (2015). Obstructor A Organizes Matrix Assembly at the Apical

    Cell Surface to Promote Enzymatic Cuticle Maturation in Drosophila. *Journal of Biological*

    Chemistry, *290*(16), 10071–10082. https://doi.org/10.1074/jbc.M114.614933

Pesch, Y.-Y., Riedel, D., Patil, K. R., Loch, G., & Behr, M. (2016). Chitinases and Imaginal disc

    growth factors organize the extracellular matrix formation at barrier tissues in insects.

    Scientific Reports, *6*(1), 18340. https://doi.org/10.1038/srep18340

Petkau, G., Wingen, C., Jussen, L. C. A., Radtke, T., & Behr, M. (2012). Obstructor-A Is Required

    for Epithelial Extracellular Matrix Dynamics, Exoskeleton Function, and Tubulogenesis.

    Journal of Biological Chemistry, *287*(25), 21396–21405.

    https://doi.org/10.1074/jbc.M112.359984

Piasecka, B., Lichocki, P., Moretti, S., Bergmann, S., & Robinson-Rechavi, M. (2013). The

    Hourglass and the Early Conservation Models—Co-Existing Patterns of Developmental

Constraints in Vertebrates. *PLoS Genetics*, *9*(4), e1003476.

https://doi.org/10.1371/journal.pgen.1003476

Qu, Z., Bendena, W. G., Tobe, S. S., & Hui, J. H. L. (2018). Juvenile hormone and sesquiterpenoids

in arthropods: Biosynthesis, signaling, and role of MicroRNA. *The Journal of Steroid*

Biochemistry and Molecular Biology, *184*, 69–76.

https://doi.org/10.1016/j.jsbmb.2018.01.013

Rackauskas, C., Koranda, J., Allen, S., Burries, R., Demski, K., Gore, L., Jung, T., Kane, K., Subaitis,

C., Urban, B., & Whitman, D. W. (2006). Molting inhibits feeding in a grasshopper.

Journal of Orthoptera Research, *15*(2), 187–190. https://doi.org/10.1665/1082-

6467(2006)15%255B187:MIFIAG%255D2.0.CO;2

Remmel, A., Käther, K. K., Stadler, P. F., & Lemke, S. (2025). Recently Evolved, Stage-Specific

Genes Are Enriched at Life-Stage Transitions in Flies. *Journal of Experimental Zoology*

Part B: Molecular and Developmental Evolution, jez.b.23317.

https://doi.org/10.1002/jez.b.23317

Riedel, F., Vorkel, D., & Eaton, S. (2011). Megalin-dependent Yellow endocytosis restricts

melanization in the *Drosophila* cuticle. *Development*, *138*(1), 149–158.

https://doi.org/10.1242/dev.056309

Roer, R., Abehsera, S., & Sagi, A. (2015). Exoskeletons across the Pancrustacea: Comparative

Morphology, Physiology, Biochemistry and Genetics. *Integrative and Comparative*

Biology, *55*(5), 771–791. https://doi.org/10.1093/icb/icv080

Rota-Stabelli, O., Daley, A. C., & Pisani, D. (2013). Molecular Timetrees Reveal a Cambrian

Colonization of Land and a New Scenario for Ecdysozoan Evolution. *Current Biology*,

23(5), 392–398. https://doi.org/10.1016/j.cub.2013.01.026

Sayers, E. W., Beck, J., Bolton, E. E., Brister, J. R., Chan, J., Comeau, D. C., Connor, R., DiCuccio,

M., Farrell, C. M., Feldgarden, M., Fine, A. M., Funk, K., Hatcher, E., Hoeppner, M., Kane,

M., Kannan, S., Katz, K. S., Kelly, C., Klimke, W., … Sherry, S. T. (2024). Database resources

of the National Center for Biotechnology Information. *Nucleic Acids Research*, *52*(D1),

D33–D43. https://doi.org/10.1093/nar/gkad1044

Schumann, I., Kenny, N., Hui, J., Hering, L., & Mayer, G. (2018). Halloween genes in

panarthropods and the evolution of the early moulting pathway in Ecdysozoa. *Royal

Society Open Science*, *5*(9), 180888. https://doi.org/10.1098/rsos.180888

Schwentner, M., Combosch, D. J., Pakes Nelson, J., & Giribet, G. (2017). A Phylogenomic Solution

to the Origin of Insects by Resolving Crustacean-Hexapod Relationships. *Current Biology*,

27(12), 1818-1824.e5. https://doi.org/10.1016/j.cub.2017.05.040

Shan, T., Wang, Y., Bhattarai, K., & Jiang, H. (2023). An evolutionarily conserved serine protease

network mediates melanization and Toll activation in *Drosophila*. *Science Advances*,

9(51), eadk2756. https://doi.org/10.1126/sciadv.adk2756

Shi, X., Li, S., Yang, L., Liu, X., Merzendorfer, H., Zhu, K. Y., & Zhang, J. (2022). Clathrin heavy

chain is essential for the development and reproduction of *Locusta migratoria*. *Insect

Science*, *29*(6), 1601–1611. https://doi.org/10.1111/1744-7917.13030

Shields, V. D. C., Gall, L. F., & Rust, M. K. (2008). Ultrastructure of Insect Sensilla. In J. L. Capinera

(Ed.), *Encyclopedia of Entomology* (pp. 4009–4023). Springer Netherlands.

https://doi.org/10.1007/978-1-4020-6359-6_2295

Simão, F. A., Waterhouse, R. M., Ioannidis, P., Kriventseva, E. V., & Zdobnov, E. M. (2015).

BUSCO: Assessing genome assembly and annotation completeness with single-copy

orthologs. *Bioinformatics*, *31*(19), 3210–3212.

https://doi.org/10.1093/bioinformatics/btv351

So, W. L., Nong, W., Xie, Y., Baril, T., Ma, H., Qu, Z., Haimovitz, J., Swale, T., Gaitan-Espitia, J. D.,

Lau, K. F., Tobe, S. S., Bendena, W. G., Kai, Z., Hayward, A., & Hui, J. H. L. (2022).

Myriapod genomes reveal ancestral horizontal gene transfer and hormonal gene loss in

millipedes. *Nature Communications*, *13*(1), 3010. https://doi.org/10.1038/s41467-022-

30690-0

Soneson, C., Love, M. I., & Robinson, M. D. (2016). Differential analyses for RNA-seq: Transcript-

level estimates improve gene-level inferences. *F1000Research*, *4*, 1521.

https://doi.org/10.12688/f1000research.7563.2

Stamp, N. E., & Bowers, M. D. (1994). Effect of temperature and leaf age on growth versus

moulting time of a generalist caterpillar fed plantain (*Plantago lanceolata*). *Ecological

Entomology*, *19*(2), 199–206. https://doi.org/10.1111/j.1365-2311.1994.tb00410.x

Suzuki, Y., Koyama, T., Hiruma, K., Riddiford, L. M., & Truman, J. W. (2013). A molt timer is

involved in the metamorphic molt in *Manduca sexta* larvae. *Proceedings of the National

Academy of Sciences*, *110*(31), 12518–12525. https://doi.org/10.1073/pnas.1311405110

Sweeney, B. W., & Vannote, R. L. (1981). Ephemerella Mayflies of White Clay Creek: Bioenergetic

and Ecological Relationships Among Six Coexisting Species. *Ecology*, *62*(5), 1353–1369.

https://doi.org/10.2307/1937299

Sztal, T., Chung, H., Gramzow, L., Daborn, P. J., Batterham, P., & Robin, C. (2007). Two

independent duplications forming the Cyp307a genes in Drosophila. *Insect Biochemistry*

and Molecular Biology, *37*(10), 1044–1053. https://doi.org/10.1016/j.ibmb.2007.05.017

Tang, H. (2009). Regulation and function of the melanization reaction in Drosophila. *Fly*, *3*(1),

105–111. https://doi.org/10.4161/fly.3.1.7747

Techer, M. A., Rane, R. V., Grau, M. L., Roberts, J. M. K., Sullivan, S. T., Liachko, I., Childers, A. K.,

Evans, J. D., & Mikheyev, A. S. (2019). Divergent evolutionary trajectories following

speciation in two ectoparasitic honey bee mites. *Communications Biology*, *2*(1), 357.

https://doi.org/10.1038/s42003-019-0606-0

Truman, J. W. (2019). The Evolution of Insect Metamorphosis. *Current Biology*, *29*(23), R1252–

R1268. https://doi.org/10.1016/j.cub.2019.10.009

Truman, J. W., Hiruma, K., Allee, J. P., MacWhinnie, S. G. B., Champlin, D. T., & Riddiford, L. M.

(2006). Juvenile Hormone Is Required to Couple Imaginal Disc Formation with Nutrition

in Insects. *Science*, *312*(5778), 1385–1388. https://doi.org/10.1126/science.1123652

Truman, J. W., & Riddiford, L. M. (2019). The evolution of insect metamorphosis: A

developmental and endocrine view. *Philosophical Transactions of the Royal Society B:*

Biological Sciences, *374*(1783), 20190070. https://doi.org/10.1098/rstb.2019.0070

Truman, J. W., & Riddiford, L. M. (2022). *Chinmo* is the larval member of the molecular trinity that directs *Drosophila* metamorphosis. *Proceedings of the National Academy of Sciences*, *119*(15), e2201071119. https://doi.org/10.1073/pnas.2201071119

Vakirlis, N., Vance, Z., Duggan, K. M., & McLysaght, A. (2022). De novo birth of functional microproteins in the human lineage. *Cell Reports*, *41*(12), 111808. https://doi.org/10.1016/j.celrep.2022.111808

Vavricka, C. J., Christensen, B. M., & Li, J. (2010). Melanization in living organisms: A perspective of species evolution. *Protein & Cell*, *1*(9), 830–841. https://doi.org/10.1007/s13238-010-0109-8

Volovych, O., Khananashvili, L., & Chipman, A. D. (2025). Exploring the interplay between tanning, desiccation resistance, and calcium metabolism in cockroaches. *Insect Biochemistry and Molecular Biology*, *185*, 104428. https://doi.org/10.1016/j.ibmb.2025.104428

Wacholder, A., Parikh, S. B., Coelho, N. C., Acar, O., Houghton, C., Chou, L., & Carvunis, A.-R. (2023). A vast evolutionarily transient translatome contributes to phenotype and fitness. *Cell Systems*, *14*(5), 363-381.e8. https://doi.org/10.1016/j.cels.2023.04.002

Ward, C. M., Aumann, R. A., Whitehead, M. A., Nikolouli, K., Leveque, G., Gouvi, G., Fung, E., Reiling, S. J., Djambazian, H., Hughes, M. A., Whiteford, I., Caceres-Barrios, C., Nguyen, T. N. M., Choo, A., Crisp, P., Sim, S. B., Geib, S. M., Marec, F., Häcker, I., … Schetelig, M. F. (2021). White pupae phenotype of tephritids is caused by parallel mutations of a MFS transporter. *Nature Communications*, *12*(1), 491. https://doi.org/10.1038/s41467-020-20680-5

Wei, D., Xu, H.-Q., Chen, D., Zhang, S.-Y., Li, W.-J., Smagghe, G., & Wang, J.-J. (2020). Genome-wide gene expression profiling of the melon fly, Zeugodacus cucurbitae, during thirteen life stages. *Scientific Data*, *7*(1), 45. https://doi.org/10.1038/s41597-020-0387-9

White, B. H., & Ewer, J. (2014). Neural and Hormonal Control of Postecdysial Behaviors in Insects. *Annual Review of Entomology*, *59*(1), 363–381. https://doi.org/10.1146/annurev-ento-011613-162028

Wodarz, A., & Nusse, R. (1998). MECHANISMS OF WNT SIGNALING IN DEVELOPMENT. *Annual Review of Cell and Developmental Biology*, *14*(1), 59–88. https://doi.org/10.1146/annurev.cellbio.14.1.59

Wolfe, J. M., Daley, A. C., Legg, D. A., & Edgecombe, G. D. (2016). Fossil calibrations for the arthropod Tree of Life. *Earth-Science Reviews*, *160*, 43–110. https://doi.org/10.1016/j.earscirev.2016.06.008

Wu, T., Hu, E., Xu, S., Chen, M., Guo, P., Dai, Z., Feng, T., Zhou, L., Tang, W., Zhan, L., Fu, X., Liu, S., Bo, X., & Yu, G. (2021). clusterProfiler 4.0: A universal enrichment tool for interpreting omics data. *The Innovation*, *2*(3), 100141. https://doi.org/10.1016/j.xinn.2021.100141

Xu, Z., Liu, A., Li, S., Wang, G., & Ye, H. (2020). Hepatopancreas immune response during molt cycle in the mud crab, Scylla paramamosain. *Scientific Reports*, *10*(1), 13102. https://doi.org/10.1038/s41598-020-70139-2

Yochem, J., Lažetić, V., Bell, L., Chen, L., & Fay, D. (2015). C. elegans NIMA-related kinases NEKL-2 and NEKL-3 are required for the completion of molting. *Developmental Biology*, *398*(2), 255–266. https://doi.org/10.1016/j.ydbio.2014.12.008

Zhang, J., Lu, A., Kong, L., Zhang, Q., & Ling, E. (2014). Functional Analysis of Insect Molting Fluid

    Proteins on the Protection and Regulation of Ecdysis. *Journal of Biological Chemistry*,

    289(52), 35891–35906. https://doi.org/10.1074/jbc.M114.599597

Zhang, X., Yuan, J., Li, F., & Xiang, J. (2021). Chitin Synthesis and Degradation in Crustaceans: A

    Genomic View and Application. *Marine Drugs*, *19*(3), 153.

    https://doi.org/10.3390/md19030153

Zieger, E., Robert, N. S. M., Calcino, A., & Wanninger, A. (2021). Ancestral Role of Ecdysis-Related

    Neuropeptides in Animal Life Cycle Transitions. *Current Biology*, *31*(1), 207-213.e4.

    https://doi.org/10.1016/j.cub.2020.10.004

Data availability

Acknowledgments

The authors would like to thank all members of the SNSF Sinergia Arthropod Moulting Project: Michele Leone, Valentine Rech de Laval, Olga Volovych, Idan Sheizaf, Asya Novikova, Sinead Lynch, Jonathan Antcliffe, Harriet Drage and Allison Daley, the Robinson-Rechavi Group and Christabel Bucao (EMBL), for their invaluable support, and insightful discussions throughout this study. This work was supported by funding from a Swiss National Science Foundation (SNSF) Sinergia award [grant number 198691]. Silhouette icons used in this study were all retrieved from Phylopic under Universal Public Domain License (CC0 1.0) and Public Domain Mark (PDM).