Aerobic Exercise Preserves Capillary Oxygen Homeostasis and Neurovascular Function in a Mouse Model of Alzheimer's Disease

Alfredo Cardenas-Rivera *[+]; Eda Erdogmus *; Austin Birmingham *[&]; Evan Yee *; Jaime Anton *; Zhengyi Lu *; Yuntao Li *; Matthew Morais *; Nazila Loghmani *; Chang Liu *; Mohammad A. Yaseen *

*Department of Bioengineering, Northeastern University, Boston, MA, USA
[&]Weill Cornell Medical College, New York, NY, USA
[+]Centro de Investigación en Bioingeniería, Universidad de Ingeniería y Tecnología (UTEC), Lima, Peru

Abstract

Age-related cerebrovascular dysfunction is increasingly recognized as a critical contributor to cognitive decline and Alzheimer's disease (AD) progression [1,2], yet the extent to which lifestyle interventions preserve microvascular oxygen delivery remains poorly defined. Physical activity (PA) improves vascular health and cognition in humans [3–7], but its impact on capillary-level oxygenation and functional hyperemia during aging and amyloid pathology is unknown. Here, we longitudinally quantified microvascular oxygen tension and stimulus-evoked oxygen dynamics in awake APP/PS1 mice and wild-type littermates using two-photon phosphorescence lifetime microscopy. Chronic aerobic PA initiated in early adulthood preserved basal arteriolar, capillary, and venular oxygenation, prevented age-dependent increases in microvascular heterogeneity, and mitigated excessive oxygen extraction in preclinical AD mice. While amyloid pathology impaired stimulus-evoked oxygen responses across vascular compartments, PA selectively enhanced capillary dilation and accelerated hyperemic kinetics without altering vascular density or architecture. Notably, sedentary AD mice developed widespread capillary hypoxia and bimodal oxygen distributions, hallmarks of malignant microvascular dysfunction, which were largely absent in physically active animals. These findings demonstrate that routine aerobic PA preserves microvascular oxygen homeostasis and functional responsiveness during aging and amyloid accumulation, supporting a capillary-centric mechanism through which exercise confers neurovascular resilience in preclinical AD.

1. INTRODUCTION

Alzheimer's disease (AD) is traditionally defined by the accumulation of amyloid-β (Aβ) plaques and tau pathology [8–12]; however, mounting evidence indicates that cerebrovascular dysfunction precedes and

accelerates neurodegeneration [1,13–16]. Alterations in cerebral blood flow, oxygen extraction, and neurovascular coupling are detectable in cognitively normal individuals [2,17–21] at risk for AD and are strongly associated with early cognitive impairment [16,22–24]. Functional neuroimaging studies consistently report aberrant blood-oxygen-level-dependent (BOLD) responses during aging and preclinical AD [25–27], yet the microvascular mechanisms underlying these changes remain incompletely understood.

The cerebral microvasculature, particularly the capillary network, plays a central role in oxygen delivery, metabolic exchange, and clearance of toxic metabolites, including Aβ [28,29]. Emerging work suggests that capillary dysfunction—manifested as impaired perfusion, increased flow heterogeneity, and stalled red blood cells—can generate localized hypoxia and metabolic stress even in the absence of overt vascular rarefaction [30,31]. Such microvascular disturbances are not uniformly distributed but instead create microscopic territories of impaired oxygen delivery, a phenomenon increasingly implicated in both normal aging and AD pathogenesis [32]. Despite this, most studies of cerebrovascular aging rely on bulk measures of blood flow or oxygenation, obscuring capillary-scale dysfunction that may critically shape neuronal vulnerability.

Aerobic physical activity (PA) is one of the most robust non-pharmacological interventions known to preserve cognitive function across the lifespan. In humans, PA improves executive function, memory, and processing speed, while reducing the risk of dementia [33–36]. At the vascular level, PA enhances endothelial function, arterial compliance, and systemic oxygen delivery [35,37–42]. In the brain, exercise has been shown to improve cerebral blood flow, reduce neuroinflammation, and attenuate amyloid burden in animal models [43–49]. However, whether PA directly preserves microvascular oxygen delivery and homogenizes capillary oxygenation during aging and AD progression remains unresolved.

Critically, neurovascular coupling—the process by which neuronal activity drives local increases in blood flow and oxygen delivery—relies on coordinated signaling among endothelial cells, vascular smooth muscle cells, and pericytes [18,50–52]. These cellular elements are particularly vulnerable to Aβ, which disrupts vasomotor control, capillary dilation, and oxygen extraction efficiency [16,30,53–59]. At the same time, PA is known to modulate the phenotype and function of these vascular cells, suggesting that exercise may exert its neuroprotective effects, at least in part, through restoration of capillary-level vascular dynamics rather than gross structural remodeling.

Here, we test the hypothesis that chronic aerobic PA preserves microvascular oxygen homeostasis and functional hyperemia during aging and amyloid pathology. Using longitudinal two-photon

phosphorescence lifetime microscopy in awake mice, we directly measured intravascular oxygen tension, oxygen extraction fraction, and stimulus-evoked oxygen responses across arterioles, capillaries, and venules in APP/PS1 mice and wild-type controls. By combining high-resolution oxygen measurements with functional stimulation and vascular morphology analysis, we delineate how PA modifies capillary oxygen delivery, heterogeneity, and responsiveness independently of vascular density. Our findings identify capillary oxygen dysfunction as a key target of PA and provide mechanistic insight into how exercise may counteract early neurovascular impairments in preclinical AD.

3. RESULTS

Physical exercise reportedly enhances systemic vascular function and memory in healthy adult humans [60]. To determine whether physical activity mitigates cerebrovascular alterations during aging and preclinical AD, we longitudinally measured microvascular oxygenation in the somatosensory cortices of awake, head-restrained mice. Male APP/PS1dE9 transgenic (AD) mice and wild-type littermates (WT) were categorized into 4 cohorts. From age 6-12 months, 2 cohorts (AD-PA and WT-PA) underwent structured treadmill running exercise (30 minutes/day, 5 days/week; Fig. 1A), while the remaining cohorts remained sedentary (AD-noPA and WT-noPA). Using two-photon phosphorescence lifetime microscopy and the phosphorescent oxygen sensitizer Oxyphor2P, intravascular oxygen tension (pO_2) was quantified up to 375 μm below the cortical surface in response to somatosensory whisker stimulation after 1 month (phase 1) and 6 months (phase 2). The influences of Aβ pathology, PA, and age were evaluated using linear mixed-effect modelling.

3.a Basal oxygenation

Figure 2 displays baseline intravascular pO_2 from cortical arterioles, capillaries, and venules of each experimental cohort. After one month (Phase 1), both AD and WT mice undergoing PA demonstrated significantly higher levels of arteriolar oxygen compared to their sedentary counterparts (Fig. 1A). Significant differences in arteriolar pO_2 were sustained between PA and sedentary cohorts after 6 months (phase 2); however, arterial pO_2 also declined with age in both AD mouse cohorts. In WT animals, arteriolar pO_2 at phase 2 was significantly higher for the WT-PA animals relative to WT-noPA.

At the capillary level, where the majority of metabolite exchange and Aβ clearance reportedly

transpires [61], PA yielded pronounced differences in AD mice at Phase 2. Both WT-PA and AD-PA

animals had significantly higher baseline capillary oxygenation at phase 2 relative to phase 1. Conversely,

AD-noPA mice demonstrated a pronounced decline in capillary oxygenation over 6 months, and capillary

pO2 did not change significantly in WT-noPA mice. Similar trends were observed in the venous

compartment. In AD-noPA mice, baseline venular oxygenation decreased significantly. In all other

cohorts, venous pO2 increased significantly between phases 1 and 2.

Microvascular heterogeneity has recently emerged as an important indicator of pathological disturbances,

indicating microscopic cortical territories with disrupted metabolic supply [62,63]. The distribution

profiles and coefficients of variations (C.V.) of baseline intravascular pO2 in Figs 1.d-g highlight how Aβ

pathology, age, and PA alter microvascular heterogeneity through their impacts on dispersity of capillary

and venous oxygenation. In all groups during Phase 1, the distributions of capillary oxygen tension

exhibited a unimodal Gaussian profile centered around ~50 mmHg with similar C.V. at 7 months. By 12

months, capillary pO2 within sedentary AD mice differed substantially from the other cohorts, displaying

a shifted, bimodal distribution. The lower, broader distribution is consistent with widespread,

heterogeneous capillary hypoxia. Venular oxygen distributions paralleled these trends, remaining

unimodal (~45 mmHg) at 7 months and becoming bimodal in sedentary AD mice at 12 months, with

distinct modes at ~23 mmHg and ~45 mmHg. In AD-PA mice, PA mitigated increases in malignant

capillary and venous heterogeneity, maintaining values comparable to those observed in age-matched WT

controls (Fig. 2E,G).

We calculated oxygen extraction fraction ($OEF = \frac{SO_{2\_artery} - SO_{2\_vein}}{SO_{2\_artery}} \times 100\%$) to quantify the amount of

oxygen delivered at each cortical depth. Microvascular pO2 measurements were converted to oxygen

saturation using the Hill equation [64]. Figs 2.H-I depict average and depth-resolved OEF in each cohort

at each phase. At phase 1, all groups demonstrated OEFs ranging around 40% that decreased slightly with

cortical depth, in agreement with prior studies of 3-6 month old awake mice [65–67]. Between phases 1

and 2, AD-noPA animals experienced significant increases in OEF across all layers, while all other cohorts

showed slight reductions. At phase 2, OEF in animals with PA appeared lower than their sedentary

counterparts.

Figure 1 Basal microvascular pO₂ in AD and wildtype mice *(A)* Arteriolar cortical oxygenation remains stable between 7 months (phase 1) and 12 months (phase 2) in wild-type (WT) mice, but is significantly reduced in 12-month-old AD mice. Physical activity (PA) induces an early and sustained increase in arteriolar oxygenation. *(B)* Capillary oxygenation increases with aging in WT mice and is markedly reduced in AD-noPA mice at 12 months; this reduction is prevented by PA. *(C)* Venular oxygenation remains stable across ages and groups, except for a significant decline in AD-noPA mice at 12 months. *(D-G)* Basal pO₂ distributions at the capillary and venular levels are similar for all groups at 7 months. AD-noPA mice at 12 months display a left-shifted, bimodal distribution, while other groups maintain a narrow, unimodal profile. *(H-I)* The cortical oxygen extraction fraction (OEF) increases for AD-noPA mice between phases 1 and 2. Routine PA yielded lower cortical OEFs in WT and AD mice at phase 2.

3.b Functional Hyperemia

We applied intravital OISI and 2P-PLIM to assess stimulus-evoked hyperemic responses in microvascular pO₂ and the impacts of physical activity PA, Aβ, and age. We measured transient changes in cortical blood volume and intravascular pO2 in response to 3-second pneumatic whisker deflection. OISI using 532 nm reflectance was first applied to identify the activated cortical region. Stimulation trials were then conducted to measure pO2 in individual microvessels from the cortical surface down to 350 μm in 50 μm increments. Linear mixed effect modeling revealed no significant depth-related differences. As a result, data from cortical layers were combined.

Figs. 3a-c displays OISI-based measurements of cerebral blood volume changes after 0 months (baseline), 1 month (phase 1), and 6-months of aerobic PA. Reductions in 532 nm OISI reflectance intensity correspond to increases in HbT, allowing simple quantification of cerebral blood volume [68]. Linear mixed effects analysis revealed no significant differences between experimental groups or phases for hyperemic blood volume changes. Figs 3d-e highlight group-averaged capillary pO₂ transients at phases 1 and 2, obtained by 2P-PLIM. In all measurements, capillary pO2 reached maximal levels shortly after stimulus cessation, slower than peak changes in blood volume. In WT animals, peak amplitude of capillary pO2 remained unaffected by age or PA. Conversely, both AD-PA and AD-noPA mice demonstrated markedly lower peak responses at 12 months of age compared to 7 months. Figs 3 g-h display maximum stimulus-induced pO2 in arterioles, capillaries, and venules, resolved into baseline pO2 and stimulus-induced hyperemic increases (colored bars and white bars, respectively). Statistical bars in figs 3. g-h indicate significant differences in maximum pO2 during stimulation. In sedentary AD mice, maximal pO2 were significantly lower in older animals (phase 2) within all vascular compartments. While in both sedentary and active WT animals, peak pO2 either increased (capillaries and venules) or remained unchanged.

Stimulus-evoked transients: Blood Volume (HbT)

A 532 nm Reflectance

B Baseline | Phase 1 | Phase 2

- WT−noPA
- WT−PA
- AD−noPA
- AD−PA

C Peak Blood Volume Change

- Baseline
- Phase 1
- Phase 2

Stimulus-evoked transients: pO$_2$

D 2P-PLIM

E Phase 1: Capillary pO$_2$ | Phase 2: Capillary pO$_2$

- WT−noPA
- WT−PA
- AD−noPA
- AD−PA

Basal and Peak pO$_2$ (mmHg)

F Arteriole | **G** Capillary | **H** Venule

- Basal
- Peak
- Phase 1
- Phase 2

To comprehensively assess variations in total blood oxygen, vascular pO2 was converted to oxygen saturation (SO2) using the Hill equation [64,66]. Fig 3 displays group-averaged SO2 transients for each vascular compartment and the calculated metrics for stimulus-evoked hyperemia ($\Delta SO2 = SO2_{peak}$ - $SO2_{baseline}$, AUC = area under curve). In arterioles, baseline SO2 levels at phase 2 were slightly higher for WT-PA and AD-PA animals relative to the noPA groups. In all groups except AD-PA, functional stimulation evoked smaller increases in vascular oxygenation at phase 2, as indicated by $\Delta SO2$ and AUC plots (figs 3.a and b).

Our observations indicate more pronounced alterations related to PA, Aβ, and age within capillary vessels. In WT animals, stimulus-evoked capillary SO2 changes were lower at phase 2 relative to phase 1, and WT-PA animals showed slightly higher SO2 changes compared to WT-noPA animals. While AD-PA animals maintained higher baseline SO2 levels, stimulus-evoked changes in SO2 were smaller in AD-PA animals and reduced further between phases 1 and 2. Interestingly, hyperemic responses in AD-PA animals displayed faster kinetics compared to the other groups, evidenced by lower rise times and fall times. Both WT-PA and AD-PA animals showed slightly faster responses than their sedentary counterparts. These differences were more pronounced at phase 2 compared to phase 1. While further investigation is required, the results could indicate PA- and age-related variations in pericyte-mediated capillary constriction and dilation.

In venules, observations of baseline and stimulus-evoked SO2 responses appeared to follow those of the capillary vessels (figs 3.e-f). AD-noPA animals showed lower baseline SO2 compared to AD-PA animals, and these differences were more pronounced at phase 2. Stimulus-evoked hyperemic SO2 changes were lower at phase 2 relative to phase 1. Interestingly in both PA groups, we observed age-related variations in fall time of the SO2 response. Both PA animals showed faster fall times at phase 2 compared to phase 1.

Figure 4 Microvascular SO₂ transients during whisker stimulation in wild-type and AD mice under aging and physical activity: *(A-B)* Arteriolar oxygen transients show higher basal SO₂, but smaller stimulus-evoked increases, in both AD-PA and WT-PA groups. For all groups, functional stimulation evoked smaller arteriolar SO₂ changes at phase 2 than phase 1. *(C-D)* In capillary vessels, Aβ and aging yielded reductions in basal SO₂ in AD-noPA mice. Routine PA protected AD-PA mice from reductions in basal SO₂. In WT animals, hyperemic increases in SO2 were higher in PA animals compared to sedentary noPA animals, while in AD animals, PA animals responded to stimulation with smaller, yet faster changes in SO2. **(E-F)** In venules, differences in basal SO2 were higher in WT-PA and AD-PA animals at phase 2. Functional stimulation evoked smaller venous SO₂ changes at phase 2 than phase 1.

At Phase 2, after 6-months of PA, we applied rapid, high-speed line-scanning to monitor capillary diameter during whisker stimulation trials (Fig. 4A). Within the activated cortical region, we observed two distinct populations of capillaries. One set demonstrated the conventional stimulus-evoked vasodilation in response to pneumatic air puffs ("responsive" capillaries; Fig. 4C), while the other set of capillaries exhibited no dilatory response ("non-responsive" capillaries; Fig. 4B). Fig 4.d-e display resting, pre-stimulus diameters for both capillary populations, along with their relative proportions for each cohort at phase 2. For all four experimental cohorts, no appreciable differences were detected between responsive and non-responsive capillary populations for resting state diameters and their relative proportions. In all cases, average capillaries diameters remained tightly distributed around 4.9 μm. Fig 4.f displays peak stimulus-induced dilation of responsive capillaries in both WT and AD mice. In both cohorts, mice that underwent six months of PA demonstrated significantly larger peak dilation relative to their sedentary counterparts. Intriguingly, responsive capillaries in AD mice demonstrated larger dilations compared to WT mice, despite having similar baseline capillary diameters. The maximal dilation occurred in response to the first air-puff across all experimental groups (Fig. 4C), while the maximal oxygenation peak occurred following the final air-puff (Fig. 4A). The area under the curve (AUC) of capillary diameter change indicates a greater overall vascular response in PA mice, compared to sedentary animals (Fig. 4E). AD mice exhibited larger AUC values than WT counterparts, with the most pronounced response observed in the AD-PA group. Fig 4.H quantifies the heterogeneity in peak dilation among responsive capillaries. AD mice showed higher, yet not significantly different, heterogeneity in stimulus-evoked diameter changes.

Figure 5 Capillary dilation responses during functional activation in wild-type and AD mice at Phase 2 (age 12 months). *(A)* Line-scan imaging protocol used to measure rapid vascular dynamics during whisker stimulation. *(B–C)* Two distinct subtypes of capillary responses were identified: responsive vessels exhibiting rapid, stimulus-locked dilations to each pneumatic air-puff and (**C**) non-responsive vessels displaying no discernible change in diameter. *(D, E)* Baseline diameters for responsive and non-responsive capillaries at phase 2. *(F-G)* Quantitative analysis of stimulus-evoked capillary dilation, area under the curve (AUC), and heterogeneity revealed larger vasodilatory responses in trained AD-PA and WT-PA mice after 6 months of regular physical activity *(H)* computed coefficients of variation from responsive capillaries indicate no differences in heterogeneity of capillary dilation.

3.c Microvascular morphology

At phase 2, we quantified features of microvascular angiography to assess the influences of Aβ pathology and 6 months routine PA. High-resolution z-stacks of cortical microvessels (~450 μm thick at 1 μm intervals) were acquired using Dextran-labeled fluorescein. We performed segmentation and analysis of the vascular image stacks using Matlab-based VIDA software package [69]. Supplementary figure # displays example stack images for each cohort and computed morphological metrics for the vascular stacks. Although linear mixed effect modeling identified differences at specific cortical depths between some experimental groups, we observed no discernible trends in vascular density, vessel length, cross sectional area of microvessels, or total vascular surface between sedentary and trained groups, irrespective of genotype. These results indicate that neither Aβ accumulation nor chronic exercise appreciably alter microvascular architecture in the barrel cortex at age 12 months.

DISCUSSION

Numerous studies have demonstrated that routine engagement in aerobic physical activity (PA) helps maintain and improve cognition and memory in aging humans and animals. In particular PA, offers benefits for processing speed, attention, and executive control [70,71]. However, a complete understanding of PA's effects on cellular and vascular physiology in the brain remains unclear. Our findings provide compelling evidence that routine aerobic PA alleviates Aβ's detrimental effects on cortical oxygen supply, oxygen extraction fraction, vascular heterogeneity, and vascular responses to functional stimulation. These results offer novel insights into PA's neurovascular implications highlighted in previous studies in healthy humans and AD patients [60,72–74]. Specifically, our findings indicate that both AD pathology and PA do not affect the vascular network uniformly. The results offer greater insight regarding variations between healthy aging and AD progression observed via fMRI BOLD neuroimaging.

We observed age-associated reductions in basal oxygen tension across the cortical vascular network in sedentary APP/PS1 mice (AD-noPA), extending from arteriolar inflow to venular outflow. Though the mechanism remains unclear, the persistent reduction in arteriolar pO_2 is consistent with earlier reports that attribute vascular abnormalities to cerebral amyloid angiopathy (CAA) along cerebral arteries. In the APP/PS1 mouse model, CAA begins to manifest around six months of age and has been shown to promote multiple impairments to arterial structure and function, including degeneration of vascular smooth muscle

cells (vSMCs), compromised integrity of the periarterial basement membrane, impaired vessel wall compliance, and vasomotor function [75]. Our basal pO2 observations in AD-noPA mice demonstrated more substantial alterations in the capillary and venous compartments at age 12 months (phase 2). In addition to significant age-related reductions in basal pO_2, the observed increases in heterogeneity of capillary and venous pO_2 reflect variable, uneven perfusion and functional shunting within the capillary beds. Such alterations purportedly result in microscopic territories of hypoxic tissue as blood flow and oxygen availability drops [62,63,76]. We observed no depth-dependent differences; however, our OEF observations agree well with prior 2P-PLIM observations in awake female C57BL/6, 3–5 months mice [65,66]. In our AD-noPA mice, OEF increased substantially between ages 7 months (phase 1) and 12 months (phase 2). Conversely, based on bolus tracking measurements and Jespersen & Østergaard's 3-compartment model, Gutiérrez-Jimenez et al computed lower OEF values in anesthetized APP/PS1 mice and wildtype littermates at age 18-months [62,77]. The OEF disparities are likely attributable to different measurement methods, anesthesia-induced alterations to cerebral energy metabolism and blood flow, and age-related variations in cerebral blood flow. When taken together, however, the findings from our study and Gutiérrez-Jimenez highlight how capillary function and perfusion each vary independently with age and during AD progression. The resultant age-dependent alterations in cerebral blood flow, metabolism, and oxygen extraction likely account for differences in fMRI BOLD response for pre-symptomatic, subclinical, and demented AD patients [76].

Routine aerobic PA yielded significant increases in basal arteriolar pO2 in both WT and AD mice at phases 1 and 2. In capillaries and venules, PA prevented or mitigated reductions in basal pO2 in AD mice at phase 2. Our analyses indicate no PA-induced changes to microvascular density, morphology, or surface area. For stimulus-induced hyperemic responses, the impacts of both Aβ and PA were most notably pronounced within the capillary bed, which accounts for the majority of vascular resistance and metabolite exchange [61]. In WT mice, PA yielded larger changes in stimulus- evoked capillary dilation and oxygenation changes; while in AD mice, PA yielded increased capillary dilation and faster, but smaller changes in capillary oxygenation. Our observed differences in arterioles and capillaries support prior reports highlighting vulnerability of vascular smooth muscle cells, endothelial cells, and pericytes to Aβ, as well as the potential influences of PA on each of these cell types [59,60,76,76,78,79]. Importantly, while Aβ and PA alter their cellular phenotypes, our data reflects the resultant impact on vascular

physiology. In a pilot study, the impact of PA on capillaries on capillary stalls and other stuff appeared less evident through.

The differential response of capillary oxygenation in WT and AD mice to PA constitutes a notable finding. In WT mice, capillary SO_2 exhibited a transient decline following one month of PA, which fully recovered by six months. This initial dip likely reflects a progressive physiological adaptation, where early increases in oxygen demand could outpace microvascular supply before to completely compensation with prolonged moderate exercise, a process associated with improvements in cerebral blood flow [80]. However, abrupt or excessive exercise may exceed the adaptive capacity of the cortical vasculature, potentially impairing the beneficial effects of moderate exercise [81]. This suggests that the neurovascular benefits of PA are contingent on both duration and intensity, emphasizing the importance of controlled exercise regimens in mitigating cerebrovascular deficits. Further studies are warranted to delineate the threshold at which exercise transitions from being beneficial to detrimental, particularly in the context of AD-related vascular dysfunction

Our findings strongly support emerging hypotheses that emphasize the role of capillary dysfunction for AD progression [76,82–84]. Although it provoked no changes to capillary density or morphology in either WT or AD mice, routine engagement in aerobic PA shows great promise for strongly mitigating disturbances to basal capillary oxygenation and homogenize perfusion. Taken together, our basal pO2 measurements and functional hyperemia measurements offer insight to the "hyperactivation" phenomenon observed in fMRI BOLD studies of AD patients [85]. Although PA did not substantially affect absolute changes in SO2, the relative change is augmented by the reduced baseline in AD-noPA. When compared to other cohorts, our data indicates that reduced basal oxygen in the capillary and venous compartment account for a larger stimulus-evoked relative change.

While Aerobic PA has proven useful for mitigating aging related alterations to vascular function, neuroinflammation, and numerous pathological hallmarks of preclinical AD [73,74,86–89], its protective mechanisms and capacity to prevent functional decline associated healthy aging and AD progression are still highly nuanced, multifaceted, and highly variable. Confounding parameters such as age, sex, intensity levels, and duration drastically affect PA's effectiveness for neuroprotection. Our study highlights the benefits of maintaining routine aerobic PA throughout early adulthood and middle age to mitigate disruptions to cortical arteriolar and capillary oxygenation. Our findings motivate future investigations

exploring PA's effects on endothelial cells, vascular smooth muscle cells, and pericytes within cortical
and subcortical brain regions.

Materials and Methods

Mice and Experimental Design

All male mice used in this study were on a C57BL/6 background. The APP/PS1 transgenic mouse model
(   )85Dbo/Mmjax, MMRRC Strain #034829-JAX, Jackson Laboratory) was utilized. Genotypes
were confirmed via standard PCR reactions performed by TransnetXY Company, employing proprietary
platforms and pipelines to ensure high-quality read depth and reproducibility. Mice were housed in a
temperature- and humidity-controlled animal facility with a 12-hour light/dark cycle and provided ad
libitum access to food and water. Experimental groups consisted of 4–6 mice each, analyzed at 6 months
of age. Wild-type (WT) and Alzheimer's disease (AD) mice were randomly assigned to sedentary (noPA)
or physically active conditions (PA). All mice were bred and housed at the Northeastern University (NEU)
vivarium, and all procedures were conducted in compliance with protocols approved by the NEU
Institutional Animal Care and Use Committee (NE-IACUC).

Treadmill Training

The experimental design was performed as previously described [90]. At the age of 6 months, male
APP/PS1 transgenic mice and wild-type littermate controls were randomly assigned to one of four groups
(n = 4-6 each group), previously assigned. A motor-driven treadmill (Ugo Basile, cat#47303) was used to
train the animals at fixed speeds. To ensure recovery after cranial window surgery, treadmill training
commenced 1-month post-surgery. Exercise mice underwent a 2-day acclimatization period, running for
10 minutes each day (Day 1 at 5 m/min; Day 2 at 8 m/min). Following acclimatization, exercise groups
engaged in treadmill running at progressively increasing speeds (5–11 m/min) for 30 minutes per session,
5 days per week, for 5 months (from 6 to 12 months of age). Each session began at 5 m/min, with the
speed incrementally increased to 11 m/min within the first minute. The maximum speed (11 m/min) was
maintained for the remaining 29 minutes of the session. Mice in the sedentary groups were placed on the
stationary treadmill for the same duration as the exercise groups but were not required to run. Animals
that repeatedly failed to complete exercise sessions were excluded from the experiment. This long-
duration training protocol has been reported to maintain exercise intensity at 45%–55% of VO_2 max [91].

Mapping the center of cortical activation

Cortical activation mapping was performed using optical intrinsic signal imaging (OISI) integrated into a
multimodal two-photon microscope (Ultima 2Pplus, Bruker) equipped with a high-resolution
monochromatic camera (acA1300-200μm, Part No. 106752, Basler aceU). A continuous broadband LED
light source (KL1600 LED, Part No. 1500-600, Schott), filtered at the isosbestic wavelength of oxy- and
deoxyhemoglobin (bandpass filter, 568/4 nm, cat# 65160, Edmund Optics), provided illumination.
Hemodynamic responses were recorded at 5 Hz [92] to capture stimulus-evoked cortical activity.
Activation ratio maps were generated by normalizing the post-stimulus signal (0–1.5 s after stimulus
onset) to the baseline period (1–4 s prior to stimulus onset). The spatial localization of cortical activation
(300×300 μm) was determined using a custom algorithm that mapped the entire cranial window. After
applying the 3D median and 3D Gaussian filters to the image series, the activation center was determined
by averaging the vectorized output of the region of interest (ROI) across successive imaging frames.

Chronical Cranial Window Implantation

All animal handling and surgical procedures were approved by the Northeastern University Institutional
Animal Care and Use Committee (NE-IACUC). Male C57BL/6 mice (16 weeks old, 19–24 g) were used
in this study. The surgical procedure was performed as previously described [93]. Briefly, mice were pre-
treated with dexamethasone (4.8 mg/kg) and cefazolin (0.5 g/kg) 4–6 hours before surgery to prevent brain
edema and infection, respectively. Mice were anesthetized with isoflurane (2% for induction and 1%–2%
during surgery). A circular incision was made on the scalp over the frontal and sagittal bones, and the
exposed periosteum was carefully removed with a No. 15 blade to ensure proper fixation. A custom
holding bar was attached to the skull using cyanoacrylate adhesive (Loctite-401, Henkel, Cat#135429) to
immobilize the animal during imaging. A craniotomy (~3 mm in diameter) was performed on the left
hemisphere over the primary somatosensory cortex (coordinates: A–P: 2.0 mm; M-L: 3.0 mm) while
leaving the dura intact. The craniotomy was sealed with a glass plug and secured with dental cement. Post-
surgical care included administering Sulfamethoxazole/Trimethoprim (0.5 mg/0.1 mg/mL) and
Carboprofen (1 mg/mL) in the drinking water for 5 days. Additionally, cefazolin (0.5 g/kg) was given via
intraperitoneal injections twice daily for 5 days, and Buprenorphine (0.05 mg/kg) was administered
subcutaneously once daily for the first 3 days to manage pain. Mice were provided with softened food
pellets immersed in water in a petri dish placed on the cage floor to ensure adequate nutrition and hydration
during recovery.

Training for awake imaging

Two weeks after surgery, to ensure full recovery, mice were trained to tolerate head fixation in a custom-

designed cradle for imaging sessions lasting up to 2 hours. Training sessions were conducted three times

per week over a 2-week period, with the duration of each session gradually increased. During training and

imaging sessions, mice were rewarded with sweetened milk approximately every 15 minutes. To promote

comfort and natural behavior, mice were placed on a suspended fabric bed, allowing them to engage in

natural grooming behaviors

Functional activation protocol

Functional hyperemia in the somatosensory primary cortex (Barrel cortex) was induced by gentle

stimulation of the whisker pad, following a previously established protocol [94]. Briefly, a Picospritzer

microinjection device (PDES-01DXH-E-LA-4, npi electronic GmbH). Air puffs (10–20 psi) were

delivered to the contralateral whisker pad relative to the cranial window implantation. To ensure effective

activation of the whisker Barrel cortex while minimizing the eye-blink reflex, air puffs were applied in an

inferoforward direction. A 3D-printed custom nozzle directed the air puffs toward the E1 whisker. As

other whiskers were not trimmed, the stimulus affected multiple whiskers simultaneously. Each trial

consisted of a 5-second baseline period followed by a 1.5-second stimulation period. During stimulation,

air puffs were delivered at a frequency of 3.3 Hz, with each puff lasting 300 ms and separated by a 300-

ms interval. The inter-stimulation interval (ISI, onset-to-onset) was set to 35 seconds.

Oxygen Measurements

Systemic oxygen delivery and exchange were quantified using OxyPhor-2P, a phosphorescent oxygen-

sensitive dye injected intravascularly (10 μL of 20 μM solution). Two-photon excitation fluorescence

microscopy was conducted with a scanning microscope (Ultima 2Pplus, Bruker), integrated with a tunable

Ti:sapphire laser (80 MHz, <120 ps pulse width, Insight X3+, Spectra-Physic) tuned to 950 nm for

OxyPhor-2 excitation. The imaging system employed a 25× water-immersion objective (380–1050 nm,

1.1 NA, 2.0 mm WD, N25X-APO-LWD, Nikon), ensuring precise optical focus. System control,

including motorized alignment with the cortical surface, image acquisition, and adjustments to laser power

and PMT voltage, was managed using embedded software (Prairie View, Bruker), fully integrated with

the scan head, laser power and PMT voltage control.

Measurements of partial pressure of oxygen (pO2) were derived from the phosphorescent lifetime of

OxyPhor-2, utilizing the Stern-Volmer equation as previously described [95]. Each phosphorescence

excitation/collection cycle comprised a 10 μs pulse train (800 femtosecond pulses) followed by a 290 μs

interval, culminating in a total cycle duration of 300 μs. Data acquisition was conducted across 10

locations per iteration, with 100 cycles per location. This process was repeated 72 times to accumulate
~20 seconds of measurement per iteration, for a total of 18 iterations (~6 minutes of data collection per
trial). Emission light was routed through an infrared blocker (short-pass filter, 900 nm) and a dichroic
mirror (bins splitter, 565 nm) into a four-channel detection system. The phosphorescence emission was
isolated with a 795±75 nm bandpass filter and detected via a highly sensitive photomultiplier tube (PMT;
H10770PA-50/001, Hamamatsu). The PMT signal was digitized using an analog-to-digital module (SPC-
150N TCSPC, Becker & Hilks). The PMT signal was digitized using an analog-to-digital converter.
Phosphorescence lifetimes were determined by fitting decay curves to single-exponential models using
non-linear least squares regression. Standard errors for lifetime measurements were estimated through a
bootstrap resampling methodology implemented in a custom executable file [95]. The calculated lifetimes
and associated standard errors were converted to pO2 values using the Stern-Volmer relationship. Oxygen
saturation (SO2) was calculated from pO2 measurements using the Hill equation, with a Hill coefficient
(h) of 2.59 and a P50 value of 40.2 mmHg, as previously described [94].

Capillary blood flow imaging and vascular dynamic with two-photon laser-scanning fluorescence
microscopy
Capillary flow dynamics were quantified using a custom-built two-photon laser-scanning microscope as
previously detailed (refer to Oxygen Measurement section). Vascular labeling was achieved through
retroorbital injection of 10 μL FITC–Dextran 70 (50 mg/mL in saline; Sigma-Aldrich, cat# 46945). This
fluorescent marker delineated the plasma as bright regions in the images, with red blood cells (RBCs)
appearing as dark shadows. Imaging was performed within a selected region of interest (600 μm × 600
μm) defining by OISI protocol, targeting approximately 10 capillaries (range: 7–13 capillaries; diameter
<10 μm) across a depth range of 0–400 μm, with measurements obtained at 50 μm intervals. Velocity
calculations involved tracing a straight line parallel to the longitudinal axis of each capillary, while internal
diameter measurements were obtained by drawing perpendicular straight line across the lumen at
approximately 50 points of length each one. Data acquisition was conducted continuously during a
functional activation protocol. Velocity computation was performed using a MATLAB-based dedicated
function: HydroVel [96], and lumen diameter analysis was facilitated by a custom image processing
routine (Supplementary Figure 1). Acquired data were aggregated into 1/100-second time bins for
downstream analyses.

Microvascular Morphology Measurements

Microvascular angiograms were recorded within the same regions of interest (ROIs) used for functional
measurements (e.g., intravascular pO2, blood flow, and arteriolar diameter transients). Imaging was
performed on a separate day in awake mice at rest. Three-dimensional fluorescence intensity stacks were
acquired down to a depth of 400 μm from the cortical surface.

Data Analysis

Oxygen saturation levels were quantified in arterioles, capillaries, and venules, alongside measurements
of blood flow velocity, vascular dynamics, and vessel density, across defined cortical layers. To assess
vascular heterogeneity under both resting and stimulus-evoked conditions, the coefficient of variation
(CV) of microvascular oxygen tension was calculated within 50-μm cortical depth bins for each individual
animal. Statistical comparisons across experimental groups were performed using linear mixed-effects
models implemented with the lme4 package (version 1.1.37) in R (version 4.4.2, Pile of Leaves). The
models included group and vascular segment as fixed effects and animal identity as a random intercept,
accounting for repeated measures within subjects. Post hoc pairwise comparisons were conducted using
estimated marginal means with Tukey adjustment for multiple comparisons, implemented via the
emmeans package (version 1.11.0). All results are reported as mean ± standard error of the mean (SEM),
and statistical significance was defined as p: *< 0.05, **<0.01 and ***<0.001.

ACKNOWLEDGMENTS

This work was performed with generous support from the Northeastern College of Engineering and the
National Institutes of Health: NIH R01AA27097, NIH R56AG058849, and R21AG085655. We thank Dr.
Sava Sakadzic and Allen Alfadhel from the MGH Martinos Center for Biomedical Imaging for
consultations with data processing and interpretation. We thank the Institute for Chemical Imaging of
Living Systems (RRID:SCR_022681) at Northeastern University for consultation and imaging support,
and for the assistance of confocal imaging and wide-field fluorescence imaging.

REFERENCES

1. S. Love and J. S. Miners, "Cerebrovascular disease in ageing and Alzheimer's disease," Acta Neuropathol **131**, 645–658 (2016).
2. B. Zimmerman, B. Rypma, G. Gratton, and M. Fabiani, "Age-related changes in cerebrovascular health and their effects on neural function and cognition: A comprehensive review," Psychophysiology **58**, e13796 (2021).
3. C. H. Hillman, K. I. Erickson, and A. F. Kramer, "Be smart, exercise your heart: exercise effects on brain and cognition," Nat Rev Neurosci **9**, 58–65 (2008).
4. J. N. Barnes and A. T. Corkery, "Exercise Improves Vascular Function, but does this Translate to the Brain?," Brain Plasticity **4**, 65–79 (2018).
5. B. Koščak Tivadar, "Physical activity improves cognition: possible explanations," Biogerontology **18**, 477–483 (2017).
6. M. K. Jedrziewski, V. M.-Y. Lee, and J. Q. Trojanowski, "Physical activity and cognitive health," Alzheimer's & Dementia **3**, 98–108 (2007).
7. S. B. Chapman, S. Aslan, J. S. Spence, M. W. Keebler, L. F. DeFina, N. Didehbani, A. M. Perez, H. Lu, and M. D'Esposito, "Distinct Brain and Behavioral Benefits from Cognitive vs. Physical Training: A Randomized Trial in Aging Adults," Front. Hum. Neurosci. **10**, (2016).
8. U. Sehar, P. Rawat, A. P. Reddy, J. Kopel, and P. H. Reddy, "Amyloid Beta in Aging and Alzheimer's Disease," IJMS **23**, 12924 (2022).
9. G. Gallardo and D. M. Holtzman, "Amyloid-β and Tau at the Crossroads of Alzheimer's Disease," in *Tau Biology*, A. Takashima, B. Wolozin, and L. Buee, eds., Advances in Experimental Medicine and Biology (Springer Singapore, 2019), Vol. 1184, pp. 187–203.
10. J. W. Ashford, "The Dichotomy of Alzheimer's Disease Pathology: Amyloid-β and Tau," JAD **68**, 77–83 (2019).
11. S. Tiwari, V. Atluri, A. Kaushik, A. Yndart, and M. Nair, "Alzheimer's disease: pathogenesis, diagnostics, and therapeutics," IJN **Volume 14**, 5541–5554 (2019).
12. H. Zhang, W. Wei, M. Zhao, L. Ma, X. Jiang, H. Pei, Y. Cao, and H. Li, "Interaction between Aβ and Tau in the Pathogenesis of Alzheimer's Disease," Int J Biol Sci **17**, 2181–2192 (2021).
13. J. Klohs, "An Integrated View on Vascular Dysfunction in Alzheimer's Disease," Neurodegener Dis **19**, 109–127 (2019).
14. B. V. Zlokovic, "Neurovascular pathways to neurodegeneration in Alzheimer's disease and other disorders," Nat Rev Neurosci **12**, 723–738 (2011).
15. P. M. Pimentel-Coelho and S. Rivest, "The early contribution of cerebrovascular factors to the pathogenesis of Alzheimer's disease," Eur J of Neuroscience **35**, 1917–1937 (2012).
16. E. Solis, K. N. Hascup, and E. R. Hascup, "Alzheimer's Disease: The Link Between Amyloid-β and Neurovascular Dysfunction," JAD **76**, 1179–1198 (2020).
17. C. Xing, J. Feng, J. Yao, X.-M. Xu, Y. Wu, X. Yin, R. Salvi, Y.-C. Chen, and X. Fang, "Neurovascular coupling dysfunction associated with cognitive impairment in presbycusis," Brain Communications **6**, fcae215 (2024).
18. A. A. Phillips, F. H. Chan, M. M. Z. Zheng, A. V. Krassioukov, and P. N. Ainslie, "Neurovascular coupling in humans: Physiology, methodological advances and clinical implications," J Cereb Blood Flow Metab **36**, 647–664 (2016).
19. H. Li, C. Chai, Y. Xie, H. Wang, X. Bai, Y. Li, Q. Zhong, M. Xu, X. Wang, and S. Xia, "Disturbed neurovascular coupling in patients with white matter hyperintensities: potential biomarker for cognitive impairment," Neuroradiology **66**, 1967–1978 (2024).

20. Y. Hu, Y. Qiu, Y. Chen, Y. Wang, Y. Dai, Q. Xu, and Y. Zhou, "Neurovascular coupling alterations related to cognitive impairment in cerebral small vessel disease: A multiscale brain network perspective," Progress in Neuro-Psychopharmacology and Biological Psychiatry **138**, 111311 (2025).

21. T.-Y. You, Q. Dong, and M. Cui, "Emerging Links between Cerebral Blood Flow Regulation and Cognitive Decline: A Role for Brain Microvascular Pericytes," Aging Dis **14**, 1276–1291 (2023).

22. J. Li, Q. Zeng, X. Luo, K. Li, X. Liu, L. Hong, X. Zhang, S. Zhong, T. Qiu, Z. Liu, Y. Chen, P. Huang, M. Zhang, and for the Alzheimer's Disease Neuroimaging Initiative, "Decoupling of Regional Cerebral Blood Flow and Brain Function Along the Alzheimer's Disease Continuum," Journal of Alzheimer's Disease **95**, 287–298 (2023).

23. A. Montagne, D. A. Nation, J. Pa, M. D. Sweeney, A. W. Toga, and B. V. Zlokovic, "Brain imaging of neurovascular dysfunction in Alzheimer's disease," Acta Neuropathol **131**, 687–707 (2016).

24. H. Chen, Y. Guo, Z. Zhang, W. Huang, K. Zhang, H. Qin, F. Wu, T. Liu, and F. Chen, "Neurovascular coupling dysfunction associated with cognition and glial fibrillary acidic protein along the Alzheimer's disease continuum," Journal of Alzheimer's Disease **108**, 170–179 (2025).

25. M. D'Esposito, L. Y. Deouell, and A. Gazzaley, "Alterations in the BOLD fMRI signal with ageing and disease: a challenge for neuroimaging," Nat Rev Neurosci **4**, 863–872 (2003).

26. V. Scarapicchia, E. L. Mazerolle, J. D. Fisk, L. J. Ritchie, and J. R. Gawryluk, "Resting State BOLD Variability in Alzheimer's Disease: A Marker of Cognitive Decline or Cerebrovascular Status?," Front. Aging Neurosci. **10**, 39 (2018).

27. S. A. R. B. Rombouts, R. Goekoop, C. J. Stam, F. Barkhof, and P. Scheltens, "Delayed rather than decreased BOLD response as a marker for early Alzheimer's disease," NeuroImage **26**, 1078–1085 (2005).

28. Y. Cheng, D.-Y. Tian, and Y.-J. Wang, "Peripheral clearance of brain-derived Aβ in Alzheimer's disease: pathophysiology and therapeutic perspectives," Transl Neurodegener **9**, 16 (2020).

29. Y.-L. Zhang, J. Wang, Z.-N. Zhang, Q. Su, and J.-H. Guo, "The relationship between amyloid-beta and brain capillary endothelial cells in Alzheimer's disease," Neural Regen Res **17**, 2355 (2022).

30. L. Østergaard, R. Aamand, E. Gutiérrez-Jiménez, Y.-C. L. Ho, J. U. Blicher, S. M. Madsen, K. Nagenthiraja, R. B. Dalby, K. R. Drasbek, A. Møller, H. Brændgaard, K. Mouridsen, S. N. Jespersen, M. S. Jensen, and M. J. West, "The capillary dysfunction hypothesis of Alzheimer's disease," Neurobiology of Aging **34**, 1018–1031 (2013).

31. S. Vorobev, S. Yanishevskiy, S. Efimtsev, A. Sokolov, and V. Dyachuk, "Development of capillary dysfunction in Alzheimer's disease," Front. Aging Neurosci. **16**, 1458455 (2024).

32. A. G. Bryant, M. K. Manhard, D. H. Salat, B. R. Rosen, B. T. Hyman, K. A. Johnson, S. Huang, R. E. Bennett, and Y.-F. Yen, "Heterogeneity of Tau Deposition and Microvascular Involvement in MCI and AD," Curr Alzheimer Res **18**, 711–720 (2021).

33. K. S. Frederiksen, A. Verdelho, S. Madureira, H. Bäzner, J. T. O'Brien, F. Fazekas, P. Scheltens, R. Schmidt, A. Wallin, L. Wahlund, T. Erkinjunttii, A. Poggesi, L. Pantoni, D. Inzitari, G. Waldemar, and on behalf of the LADIS Study, "Physical activity in the elderly is associated with improved executive function and processing speed: the LADIS Study," Int J Geriat Psychiatry **30**, 744–750 (2015).

34. M. I. Cardona, A. Afi, N. Lakicevic, and J. R. Thyrian, "Physical Activity Interventions and Their Effects on Cognitive Function in People with Dementia: A Systematic Review and Meta-Analysis," IJERPH **18**, 8753 (2021).

35. L. M. J. Sanders, T. Hortobágyi, E. G. A. Karssemeijer, E. A. Van Der Zee, E. J. A. Scherder, and M. J. G. Van Heuvelen, "Effects of low- and high-intensity physical exercise on physical and

cognitive function in older persons with dementia: a randomized controlled trial," Alz Res Therapy **12**, 28 (2020).

36. C. M. Hoffmann, M. E. Petrov, and R. E. Lee, "Aerobic physical activity to improve memory and executive function in sedentary adults without cognitive impairment: A systematic review and meta-analysis," Preventive Medicine Reports **23**, 101496 (2021).

37. D. J. Green, A. Maiorana, G. O'Driscoll, and R. Taylor, "Effect of exercise training on endothelium-derived nitric oxide function in humans," The Journal of Physiology **561**, 1–25 (2004).

38. S. Di Francescomarino, A. Sciartilli, V. Di Valerio, A. Di Baldassarre, and S. Gallina, "The Effect of Physical Exercise on Endothelial Function:," Sports Medicine **39**, 797–812 (2009).

39. R. Hambrecht, V. Adams, S. Erbs, A. Linke, N. Kränkel, Y. Shu, Y. Baither, S. Gielen, H. Thiele, J. F. Gummert, F. W. Mohr, and G. Schuler, "Regular Physical Activity Improves Endothelial Function in Patients With Coronary Artery Disease by Increasing Phosphorylation of Endothelial Nitric Oxide Synthase," Circulation **107**, 3152–3158 (2003).

40. B. Hornig, V. Maier, and H. Drexler, "Physical Training Improves Endothelial Function in Patients With Chronic Heart Failure," Circulation **93**, 210–214 (1996).

41. M. J. Joyner, "Effect of Exercise on Arterial Compliance," Circulation **102**, 1214–1215 (2000).

42. A. W. Ashor, J. Lara, M. Siervo, C. Celis-Morales, C. Oggioni, D. G. Jakovljevic, and J. C. Mathers, "Exercise Modalities and Endothelial Function: A Systematic Review and Dose–Response Meta-Analysis of Randomized Controlled Trials," Sports Med **45**, 279–296 (2015).

43. J. P. Fisher, D. Hartwich, T. Seifert, N. D. Olesen, C. L. McNulty, H. B. Nielsen, J. J. Van Lieshout, and N. H. Secher, "Cerebral perfusion, oxygenation and metabolism during exercise in young and elderly individuals," The Journal of Physiology **591**, 1859–1870 (2013).

44. S. Ogoh and P. N. Ainslie, "Cerebral blood flow during exercise: mechanisms of regulation," Journal of Applied Physiology **107**, 1370–1380 (2009).

45. K. J. Smith and P. N. Ainslie, "Regulation of cerebral blood flow and metabolism during exercise," Experimental Physiology **102**, 1356–1371 (2017).

46. J. P. D. Kleinloog, K. M. R. Nijssen, R. P. Mensink, and P. J. Joris, "Effects of Physical Exercise Training on Cerebral Blood Flow Measurements: A Systematic Review of Human Intervention Studies," International Journal of Sport Nutrition and Exercise Metabolism **33**, 47–59 (2023).

47. M. A. Małkiewicz, A. Szarmach, A. Sabisz, W. J. Cubała, E. Szurowska, and P. J. Winklewski, "Blood-brain barrier permeability and physical exercise," J Neuroinflammation **16**, 15 (2019).

48. D. Bao, J. Zhou, Y. Hao, X. Yang, W. Jiao, Y. Hu, and X. Wang, "The Effects of Fatiguing Aerobic Exercise on the Cerebral Blood Flow and Oxygen Extraction in the Brain: A Piloting Neuroimaging Study," Front. Neurol. **10**, 654 (2019).

49. I. D. Braz and J. P. Fisher, "The impact of age on cerebral perfusion, oxygenation and metabolism during exercise in humans," The Journal of Physiology **594**, 4471–4483 (2016).

50. I. M. Devonshire, N. G. Papadakis, M. Port, J. Berwick, A. J. Kennerley, J. E. W. Mayhew, and P. G. Overton, "Neurovascular coupling is brain region-dependent," NeuroImage **59**, 1997–2006 (2012).

51. C. Iadecola, "The Neurovascular Unit Coming of Age: A Journey through Neurovascular Coupling in Health and Disease," Neuron **96**, 17–42 (2017).

52. P. J. Drew, "Neurovascular coupling: motive unknown," Trends in Neurosciences **45**, 809–819 (2022).

53. M. G. Kozberg, L. P. Munting, L. H. Maresco, C. A. Auger, M. L. Van Den Berg, B. Denis De Senneville, L. Hirschler, J. M. Warnking, E. L. Barbier, C. T. Farrar, S. M. Greenberg, B. J.

609 Bacskai, and S. J. Van Veluw, "Loss of spontaneous vasomotion precedes impaired cerebrovascular
610 reactivity and microbleeds in a mouse model of cerebral amyloid angiopathy," (2024).

54. W. M. Zhu, A. Neuhaus, D. J. Beard, B. A. Sutherland, and G. C. DeLuca, "Neurovascular coupling mechanisms in health and neurovascular uncoupling in Alzheimer's disease," Brain **145**, 2276–2292 (2022).

55. K. Kotliar, C. Hauser, M. Ortner, C. Muggenthaler, J. Diehl-Schmid, S. Angermann, A. Hapfelmeier, C. Schmaderer, and T. Grimmer, "Altered neurovascular coupling as measured by optical imaging: a biomarker for Alzheimer's disease," Sci Rep **7**, 12906 (2017).

56. R. Liu, T. Zhang, H. Yang, X. Lan, J. Ying, and G. Du, "The Flavonoid Apigenin Protects Brain Neurovascular Coupling against Amyloid-β25-35-Induced Toxicity in Mice," JAD **24**, 85–100 (2011).

57. L. Y. Di Marco, E. Farkas, C. Martin, A. Venneri, and A. F. Frangi, "Is Vasomotion in Cerebral Arteries Impaired in Alzheimer's Disease?," JAD **46**, 35–53 (2015).

58. K. Niwa, V. A. Porter, K. Kazama, D. Cornfield, G. A. Carlson, and C. Iadecola, "Aβ-peptides enhance vasoconstriction in cerebral circulation," American Journal of Physiology-Heart and Circulatory Physiology **281**, H2417–H2424 (2001).

59. K. Kisler, A. R. Nelson, A. Montagne, and B. V. Zlokovic, "Cerebral blood flow regulation and neurovascular dysfunction in Alzheimer disease," Nature Reviews Neuroscience **18**, 419–434 (2017).

60. D. J. Green and K. J. Smith, "Effects of Exercise on Vascular Function, Structure, and Health in Humans," Cold Spring Harb Perspect Med **8**, a029819 (2018).

61. C. Watanabe, T. Imaizumi, H. Kawai, K. Suda, Y. Honma, M. Ichihashi, M. Ema, and K. Mizutani, "Aging of the Vascular System and Neural Diseases," Front. Aging Neurosci. **12**, (2020).

62. S. N. Jespersen and L. Østergaard, "The Roles of Cerebral Blood Flow, Capillary Transit Time Heterogeneity, and Oxygen Tension in Brain Oxygenation and Metabolism," J Cereb Blood Flow Metab **32**, 264–277 (2012).

63. T. K. Roy and T. W. Secomb, "Functional implications of microvascular heterogeneity for oxygen uptake and utilization," Physiological Reports **10**, e15303 (2022).

64. K. Uchida, M. P. Reilly, and T. Asakura, "Molecular Stability and Function of Mouse Hemoglobins," jzoo **15**, 703–706 (1998).

65. B. Li, T. V. Esipova, I. Sencan, K. Kılıç, B. Fu, M. Desjardins, M. Moeini, S. Kura, M. A. Yaseen, F. Lesage, L. Østergaard, A. Devor, D. A. Boas, S. A. Vinogradov, and S. Sakadžić, "More homogeneous capillary flow and oxygenation in deeper cortical layers correlate with increased oxygen extraction," Elife **8**, e42299 (2019).

66. İ. Şencan, T. Esipova, K. Kılıç, B. Li, M. Desjardins, M. A. Yaseen, H. Wang, J. E. Porter, S. Kura, B. Fu, T. W. Secomb, D. A. Boas, S. A. Vinogradov, A. Devor, and S. Sakadžić, "Optical measurement of microvascular oxygenation and blood flow responses in awake mouse cortex during functional activation," J Cereb Blood Flow Metab **42**, 510–525 (2022).

67. C. Liu, A. Cárdenas-Rivera, S. Teitelbaum, A. Birmingham, M. Alfadhel, and M. A. Yaseen, "Neuroinflammation increases oxygen extraction in a mouse model of Alzheimer's disease," Alzheimer's Research & Therapy **16**, 78 (2024).

68. A. K. Dunn, A. Devor, A. M. Dale, and D. A. Boas, "Spatial extent of oxygen metabolism and hemodynamic changes during functional activation of the rat somatosensory cortex," Neuroimage **27**, 279–290 (2005).

69. T.-H. Tsai, S.-H. Jee, C. Y. Dong, and S.-J. Lin, "Multiphoton microscopy in dermatalogical imaging," Journal of Dermatological Science **56**, 1–8 (2009).

70. P. J. Smith, J. A. Blumenthal, B. M. Hoffman, H. Cooper, T. A. Strauman, K. Welsh-Bohmer, J. N.
Browndyke, and A. Sherwood, "Aerobic Exercise and Neurocognitive Performance: a Meta-
Analytic Review of Randomized Controlled Trials," Psychosom Med **72**, 239–252 (2010).
71. V. A. Holthoff, K. Marschner, M. Scharf, J. Steding, S. Meyer, R. Koch, and M. Donix, "Effects of
physical activity training in patients with Alzheimer's dementia: results of a pilot RCT study," PLoS
One **10**, e0121478 (2015).
72. D. J. Green, M. T. E. Hopman, J. Padilla, M. H. Laughlin, and D. H. J. Thijssen, "Vascular
Adaptation to Exercise in Humans: Role of Hemodynamic Stimuli," Physiol. Rev. **97**, 495–528
(2017).
73. E. Y. Allison and B. K. Al-Khazraji, "Cerebrovascular adaptations to habitual resistance exercise
with aging," American Journal of Physiology-Heart and Circulatory Physiology (2024).
74. L. Alves, D. Hashiguchi, C. M. Loss, H. van Praag, and B. M. Longo, "Vascular dysfunction in
Alzheimer's disease: Exploring the potential of aerobic and resistance exercises as therapeutic
strategies," J Alzheimers Dis **104**, 963–979 (2025).
75. S. H. Kim, J. H. Ahn, H. Yang, P. Lee, G. Y. Koh, and Y. Jeong, "Cerebral amyloid angiopathy
aggravates perivascular clearance impairment in an Alzheimer's disease mouse model," acta
neuropathol commun **8**, 181 (2020).
76. L. Østergaard, R. Aamand, E. Gutiérrez-Jiménez, Y.-C. L. Ho, J. U. Blicher, S. M. Madsen, K.
Nagenthiraja, R. B. Dalby, K. R. Drasbek, A. Møller, H. Brændgaard, K. Mouridsen, S. N.
Jespersen, M. S. Jensen, and M. J. West, "The capillary dysfunction hypothesis of Alzheimer's
disease," Neurobiology of Aging **34**, 1018–1031 (2013).
77. E. Gutiérrez-Jiménez, H. Angleys, P. M. Rasmussen, M. J. West, L. Catalini, N. K. Iversen, M. S.
Jensen, S. Frische, and L. Østergaard, "Disturbances in the control of capillary flow in an aged
APPswe/PS1ΔE9 model of Alzheimer's disease," Neurobiology of Aging **62**, 82–94 (2018).
78. J. N. Barnes, A. G. Pearson, A. T. Corkery, N. A. Eisenmann, and K. B. Miller, "Exercise, Arterial
Stiffness, and Cerebral Vascular Function: Potential Impact on Brain Health," Journal of the
International Neuropsychological Society **27**, 761–775 (2021).
79. R. B. Nielsen, P. Parbo, R. Ismail, R. Dalby, A. Tietze, H. Brændgaard, H. Gottrup, D. J. Brooks, L.
Østergaard, and S. F. Eskildsen, "Impaired perfusion and capillary dysfunction in prodromal
Alzheimer's disease," Alzheimer's & Dementia: Diagnosis, Assessment & Disease Monitoring **12**,
e12032 (2020).
80. J. Liu, L. Min, R. Liu, X. Zhang, M. Wu, Q. Di, and X. Ma, "The effect of exercise on cerebral
blood flow and executive function among young adults: a double-blinded randomized controlled
trial," Sci Rep **13**, 8269 (2023).
81. G. Sangüesa, M. Batlle, E. Muñoz-Moreno, G. Soria, A. Alcarraz, C. Rubies, L. Sitjà-Roqueta, E.
Solana, E. Martínez-Heras, A. Meza-Ramos, S. Amaro, S. Llufriu, L. Mont, and E. Guasch, "Intense
long-term training impairs brain health compared with moderate exercise: Experimental evidence
and mechanisms," Annals of the New York Academy of Sciences **1518**, 282–298 (2022).
82. J. C. Cruz Hernández, O. Bracko, C. J. Kersbergen, V. Muse, M. Haft-Javaherian, M. Berg, L. Park,
L. K. Vinarcsik, I. Ivasyk, D. A. Rivera, Y. Kang, M. Cortes-Canteli, M. Peyrounette, V. Doyeux,
A. Smith, J. Zhou, G. Otte, J. D. Beverly, E. Davenport, Y. Davit, C. P. Lin, S. Strickland, C.
Iadecola, S. Lorthois, N. Nishimura, and C. B. Schaffer, "Neutrophil adhesion in brain capillaries
reduces cortical blood flow and impairs memory function in Alzheimer's disease mouse models,"
Nat Neurosci **22**, 413–420 (2019).

83.   O. Bracko, J. C. Cruz Hernández, L. Park, N. Nishimura, and C. B. Schaffer, "Causes and
consequences of baseline cerebral blood flow reductions in Alzheimer's disease," Journal of
Cerebral Blood Flow & Metabolism **41**, 1501–1516 (2021).
84.   L. S. Madsen, R. B. Nielsen, P. Parbo, R. Ismail, I. K. Mikkelsen, H. Gottrup, L. Østergaard, D. J.
Brooks, and S. F. Eskildsen, "Capillary function progressively deteriorates in prodromal
Alzheimer's disease: A longitudinal MRI perfusion study," Aging Brain **2**, 100035 (2022).
85.   N. Corriveau-Lecavalier, J. N. Adams, L. Fischer, E. N. Molloy, and A. Maass, "Cerebral
hyperactivation across the Alzheimer's disease pathological cascade," Brain Commun **6**, fcae376
(2024).
86.   E. S. Bliss, R. H. Wong, P. R. Howe, and D. E. Mills, "Benefits of exercise training on
cerebrovascular and cognitive function in ageing," J Cereb Blood Flow Metab **41**, 447–470 (2021).
87.   I. Belaya, M. Ivanova, A. Sorvari, M. Ilicic, S. Loppi, H. Koivisto, A. Varricchio, H. Tikkanen, F.
R. Walker, M. Atalay, T. Malm, A. Grubman, H. Tanila, and K. M. Kanninen, "Astrocyte
remodeling in the beneficial effects of long-term voluntary exercise in Alzheimer's disease," J
Neuroinflammation **17**, 271 (2020).
88.   S.-S. Baek, "Role of exercise on the brain," Journal of Exercise Rehabilitation **12**, 380–385 (2016).
89.   C. W. Cotman, N. C. Berchtold, and L.-A. Christie, "Exercise builds brain health: key roles of
growth factor cascades and inflammation," Trends in Neurosciences **30**, 464–472 (2007).
90.   (n.d.).
91.   E. J. Baker and T. T. Gleeson, "The effects of intensity on the energetics of brief locomotor
activity," Journal of Experimental Biology **202**, 3081–3087 (1999).
92.   S. Kura, H. Xie, B. Fu, C. Ayata, D. A. Boas, and S. Sakadžić, "Intrinsic optical signal imaging of
the blood volume changes is sufficient for mapping the resting state functional connectivity in the
rodent cortex," J. Neural Eng. **15**, 035003 (2018).
93.   K. Kılıç, M. Desjardins, J. Tang, M. Thunemann, S. Sunil, Ş. E. Erdener, D. D. Postnov, D. A. Boas,
and A. Devor, "Chronic Cranial Windows for Long Term Multimodal Neurovascular Imaging in
Mice," Front. Physiol. **11**, 612678 (2021).
94.   İ. Şencan, T. Esipova, K. Kılıç, B. Li, M. Desjardins, M. A. Yaseen, H. Wang, J. E. Porter, S. Kura,
B. Fu, T. W. Secomb, D. A. Boas, S. A. Vinogradov, A. Devor, and S. Sakadžić, "Optical
measurement of microvascular oxygenation and blood flow responses in awake mouse cortex during
functional activation," J Cereb Blood Flow Metab **42**, 510–525 (2022).
95.   T. V. Esipova, M. J. P. Barrett, E. Erlebach, A. E. Masunov, B. Weber, and S. A. Vinogradov,
"Oxyphor 2P: A High-Performance Probe for Deep-Tissue Longitudinal Oxygen Imaging," Cell
Metabolism **29**, 736-744.e7 (2019).
96.   P. Y. Chhatbar and P. Kara, "Improved blood velocity measurements with a hybrid image filtering
and iterative Radon transform algorithm," Front. Neurosci. **7**, (2013).
735