Genetic and Glycosylation Variability of the GP5 Glycoprotein in *Betaarterivirus*

americense **NADC34 Variant Isolates from Peru**

Molecular Variability of the GP5 Glycoprotein in *Betaarterivirus americense*

Rony Cotaquispe Nalvarte [a*]   **Orcid:** 0000-0002-8273-4803   r.cotaquispe.n@gmail.com

Edgard De la Cruz Vásquez [b]   **Orcid:** 0000-0002-9229-985X   eddelacruz62@gmail.com

Miriam Legua Barrios [c]   **Orcid:** 0000-0002-4993-0909   miriam.legua@upsjb.edu.pe

Erick Llona García [c]   **Orcid:** 0000-0002-7522-0193   ERICK.LLONA@UPSJB.EDU.PE

Jennifer Toskano Hurtado [c]   jtoskano@corpmontana.com

[a] Research Professor, Universidad Privada Científica del Sur, Lima, Perú.

[b] Professor, Universidad Privada Autónoma de Ica, Ica, Perú.

[c] Professor, Universidad Privada San Juan Bautista, Ica, Perú.

[d] Animal Nutrition and Health Business Unit, Innovation and Development Area, Corporation

Montana S.A., Lima, Peru.

15
16 * **Correspondencia:** r.cotaquispe.n@gmail.com (correo electrónico)

18 ABSTRACT

Betaarterivirus americense causes a clinical condition with a high epidemiological risk in Peru and

worldwide. Therefore, the aim of this study was to identify the genetic diversity of circulating viruses

in different commercial swine farms in Lima. A total of 24 viral sequences from our previously

published work were analyzed, and phylogenetic inference was performed based on amino acid

alignments. The results revealed that all isolates clustered within sublineage 1.5 of the representative

NADC34 variant. Sequence analysis of ORF5 (GP5) identified a high number of amino acid

substitutions in key regions, including aa34 (24/24), aa39 (6/24), aa44 (4/24), aa47 (24/24), aa57

(24/24), aa58 (24/24), and aa59 (24/24), located within epitopes A, B, and C.

N-glycosylation analysis revealed 10 potential N-glycosylation sites: N30, N32, N33, N34, N35, N44,

N51, N57, N128, and N151. Among these, the major conserved sites (N34, N44, and N51) were fully

preserved in strains 40 and 41; in 20 out of 24 strains, only sites N44 and N51 were conserved, whereas

strains 36, 37, 38, and 39 showed complete loss of these glycosylation sites. The signal peptide

cleavage site was identified between amino acids 1 and 31.

Genetic diversity analysis revealed rapid viral evolution and a high mutation rate, with 530 variable

sites and 860 mutations. Tajima's D statistical analysis indicated population subdivision, yielding a D

value of −0.78746. The high number of substitutions in key domains of the GP5 glycoprotein and the variability of potential N-glycosylation sites among the 24 NADC34 variant isolates of *Betaarterivirus americense* suggest that these strains exhibit high genetic variability and recombination potential.

Keywords: *Betaarterivirus americense*, glycoprotein 5, N-glycosylation, NADC34

INTRODUCTION

Porcine reproductive and respiratory syndrome (PRRS) is a viral disease of major sanitary importance that poses a serious threat to the global swine industry [13]. The etiological agent, porcine reproductive and respiratory syndrome virus (PRRSV), has been molecularly and structurally characterized, and two main species have been identified: *Betaarterivirus americense* and *Betaarterivirus europensis* [3,5,6].

The PRRSV genome is a positive-sense single-stranded RNA that contains a 5′ cap structure and a 3′ polyadenylated tail. It includes a 5′ untranslated region (UTR), at least ten open reading frames (ORFs)—ORF1a, ORF1b, ORF2a, ORF2b, ORF3, ORF4, ORF5, ORF5a, ORF6, and ORF7—and a 3′ UTR [27,31]. ORF1a and ORF1b comprise approximately three-quarters of the viral genome and encode the replicase polyproteins pp1a and pp1a/b, which undergo extensive proteolytic processing by virus-encoded proteases to produce at least 16 nonstructural proteins (nsps) involved in viral replication and transcription [10,11]. The remaining ORFs located in the 3′ terminal region encode eight structural proteins, including four membrane-associated glycoproteins (GP2a, GP3, GP4, and GP5), three nonglycosylated membrane proteins (E, ORF5a, and M), and one nucleocapsid protein (N) [8,32].

Phylogenetic analysis of the ORF5 gene in *Betaarterivirus americense* has identified nine monophyletic lineages (L1–L9), as well as three Korea-specific lineages (KOR A, B, and C) and several sublineages [6,20]. This pronounced genetic diversity, attributed to the high heterogeneity and variability of glycoprotein 5 (GP5), supports the hypothesis that PRRSV variants may inherently acquire distinct virulence patterns and clinical presentations.

The ORF5 gene encodes the major viral envelope glycoprotein (GP5), a transmembrane glycosylated protein exposed on the virion surface that plays a critical role in virus attachment to host cell receptors and represents a key target of the humoral immune response [1,7,25]. GP5 consists of a signal peptide, two hypervariable regions (HVR1 and HVR2), three transmembrane domains (TM1, TM2, and TM3), a decoy epitope, the primary neutralizing epitope (PNE), an additional B-cell epitope, and two T-cell epitopes [35]. This protein is essential for viral infectivity and contains important immunological domains associated with viral neutralization, including antigenic determinants and N-glycosylation sites [18].

GP5 glycosylation is an important immune evasion mechanism used by several enveloped viruses to

escape, block, or minimize the activity of neutralizing antibodies [2]. In particular, changes in N-

glycosylation sites associated with the primary neutralizing epitope significantly affect viral

immunogenicity, rendering virions resistant to antibody neutralization in vivo [9]. In addition,

recombination and mutation are the two main evolutionary mechanisms contributing to the emergence

and evolution of new genetic variants of *Betaarterivirus americense*, playing a key role in the

increasing number of circulating genotypes [14,15].

In Peru, following the molecular identification of lineage 1, sublineage 1.5, corresponding to the

IA/2014/NADC34 variant in 2019 and 2025 [6,23], significant knowledge gaps remain regarding the

prevalence of viral species and the genetic diversity of lineages associated with reduced

immunogenicity, linked to the high genetic variability of *Betaarterivirus americense*. Therefore, the

aim of this study was to identify the variability in the molecular architecture of glycoprotein 5 (GP5)

in Peruvian isolates, contributing to a broader understanding of genetic variation, molecular

epidemiology, immune evasion mechanisms, and the limitations in developing effective vaccines

against PRRSV.

MATERIALS AND METHODS

Biological material

86   A total of 24 nucleotide sequences from strains isolated in the study by Cotaquispe [6] were analyzed

87   in this investigation.

RNA extraction, quantification, and in-house conventional RT-PCR of the ORF5 gene

89   The procedures for obtaining viral cDNA of *Betaarterivirus americense* were carried out as described

90   in the study by Cotaquispe [6].

Electrophoresis, purification, and sequencing

Amplified products were visualized using a CS Cleaver Scientific multiSUB™ Choice horizontal

electrophoresis system with 1% molecular-grade agarose gels (SCL AG500, Cleaver Scientific), using

Fluorescent Dye-DNA/Safe-Green™ (ABM®), a 100 bp Opti-DNA Marker (50 bp–1.5 kb, ABM®),

and visualized with a UVP UV Solo Touch photodocumentation system (Analytik Jena). Gel

fragments containing amplicons of the expected size were excised and purified using the innuPREP

Nucleic Acid Purification Kit (Analytik Jena). Purified products were sequenced in both directions by

Macrogen (South Korea) using the BigDye v3.1 Terminator Sequencing Kit (Life Technologies) and

an ABI 3137 XL Genetic Analyzer (Life Technologies).

Phylogenetic analysis of glycoprotein 5 (GP5)

Sequenced chromatograms of glycoprotein 5 from this study were analyzed using Chromas Lite®
v2.6.6 and combined with 58 reference sequences from GenBank. Nine of these corresponded to
Peruvian isolates (MH791386.1; MH791385.1; MH791383.1; MH791388.1; MH791379.1;
MH791378.1; MH791376.1; MH791391.1; MH791390.1), and the remaining sequences corresponded
to pathogenic strains isolated worldwide (MF326985.1; DQ009647.1; DQ306879.1; DQ306878.1;
DQ306877.1; GQ857656.1; FJ349261.1; AY588319.1; DQ489311.1; GQ461593.1; EF484033.1;
AY032626.1; EF635006.1; KF611905.1; KF287140.1; JQ715697.1; EU939312.1; GQ374442.1;
JQ663556.1; HQ315837.1; FJ889130.1; JX512910.2; JQ663545.1; JN411262.1; AY743931.1;
JF802085.1; GU047345.1; EU076704.1; AY366525.1; KC862567.1; AF325691.1; KC862577.1;
DQ473474.1; AB288356.1; JN660150.1; EU807840.1; EU262603.1; AF184212.1; DQ176019.1;
AF331831.1; AF176348.2; U87392.3; MF279253.1; EF536003.1; AF066183.4; DQ988080.1;
AY424271.1; AF494042.1; EF112445.1).

The phylogenetic tree was inferred based on amino acid substitution analysis using the maximum
likelihood method with BioNJ and the Jones–Taylor–Thornton (JTT) model. The tree was drawn to
scale, with branch lengths representing the number of substitutions per site. A total of 201 amino acids
were included in the final dataset. Bootstrap values were calculated using 1,000 alignment replicates.
Evolutionary analyses were performed using DNAMAN 10 and MEGA 6 [28].

Analysis of immunological domains of glycoprotein 5

Amino acid analysis was performed using MEGA 6.0 to determine the viral structure, including the
signal peptide, decoy epitopes I and II, hypervariable regions I and II, the primary neutralizing epitope
(PNE), transmembrane regions 1, 2, and 3, T-cell recognition epitopes 1 and 2, and the B-cell epitope.
This analysis allowed identification of genetic variants, as mutations in viral protein structures may
affect infectivity, pathogenicity, and viral persistence [17]. GP5 glycoprotein length was determined,
amino acid mutations in key immunological domains were identified, and putative N-glycosylation
sites were predicted using the NetNGlyc 1.0 server [12]. In addition, signal peptides were predicted
using SignalP 4.0 [22].

Genetic diversity analysis: Tajima's D test and DnaSP v6

Statistical analysis was based on the neutral theory of molecular evolution, which assumes that the
evolution of slightly deleterious or beneficial, nearly neutral mutations depends primarily on the
mutation rate, selection coefficients, and population size. Tajima's D statistic [29] compares two
estimates of genetic variation (θ) within a group of sequences. D is based on the difference between
these estimates and has a mean of zero with a deviation of ± 1 ($-1 < D < +1$). The D value was compared
with Table 2 [30] according to the number of samples and the closest confidence limit to the obtained
D value. Negative deviations indicate non-neutrality, whereas positive D values may indicate positive

selection or a reduction in population size. The DnaSP v6 program was used to calculate Tajima's D

values and to identify the genetic diversity of our isolates [24].

RESULTS

Serum samples were analyzed using an in-house kit with primers targeting the ORF5 gene, which

encodes glycoprotein 5 (GP5) of the virus. The primers were validated and standardized using vaccine

strains from both species as positive controls and a commercial IDEXX positive control mixture

(Figure 1). Amplification of the gene yielded a product of 723 bp (Figure 2). Sequencing of all partial

amplicons contained the genes encoding glycoprotein 5 (GP5), with lengths of 603 nucleotides and

201 functional amino acids.

Phylogenetic analysis of glycoprotein 5 (GP5)

The phylogenetic relationship of the study isolates, based on amino acid sequence alignment using

representative GenBank strains from lineages 1, 5, and 8, is shown in Figure 3. In addition, nine

isolates reported as Peruvian belonging to the sublineage 1.5 (1-7-4) variant of NADC34 were

included. A 100% support value was observed in the generated tree for the presence of the species

Betaarterivirus europense and *Betaarterivirus americense*, as well as multiple branches corresponding

to groups formed by representative variants of species 2 from Lineage 1: sublineage 1.1 USA (99-

3584, 99-3298, and 98-3403), sublineage 1.5 NADC34, sublineage 1.6 (NADC31), sublineage 1.8

(HENAN), sublineage 1.9 (MN184A); from Lineage 5: sublineage 5.1 (RespPRRS_MLV and

VR2332); and from Lineage 8: sublineage 8.9 (Ingelvac_ATP) and sublineage 8.7 (JXA1).

Representative variants of species 1 from Lineage 1-sublineage 1.A South Korea (SNU1 and KNU-

07), sublineage 1.C (EuroPRRSV, CRESA-VP21, and Porcilis GP5) and from Lineage 3 (BELGA-

Lena) were also included. Genealogy showed that the study isolates (turquoise branch) clustered with

Peruvian isolates reported in 2015, 2016, and 2017 and demonstrated a close relationship with the

highly pathogenic Lineage 1, sublineage 1.5 NADC34 variant first described in North America (Figure

3). Likewise, high divergence of strains 36, 37, 38, 39, and 42 was observed within sublineage 1.5.

Amino acid analysis of GP5

The obtained sequences were analyzed to predict potential N-glycosylation sites within glycoprotein

5 (GP5), as these are involved in viral immune evasion by minimizing the neutralizing capacity of

epitopes. Glycosylation sites N33, N44, and N51 are located within the main neutralizing epitopes.

The results demonstrated the presence of 10 potential N-glycosylation sites (N30, N32, N33, N34,

N35, N44, N51, N57, N128, and N151). Of these, 20/24 isolates presented sites N44 and N51 within

the primary neutralizing epitope (PNE). In addition, all isolates exhibited three or more N-

glycosylation sites in their sequences. Notably, new sites not reported in reference strains (N57, N128,

and N151) were identified, and a higher number of sites was observed in strains 40 and 41 (Table 1).

Furthermore, signal peptide cleavage site prediction analysis was performed to determine the site of

enzymatic cleavage; the peptidase cleavage occurs within residues 1–31 of the protein sequence (Table

1).

Genetic diversity of glycoprotein 5 (GP5)

The structure of the translated amino acid sequences (201 aa) consisted of a signal peptide (aa 1–31),

decoy epitope I (A) (aa 27–36) and decoy epitope II (aa 180–197), specialized in immune response

neutralization, as well as the primary neutralizing epitope (B) (aa 37–45), neutralizing epitope (C) (aa

52–61), transmembrane region 1 (aa 65–82), transmembrane region 2 (aa 96–102), and transmembrane

region 3 (aa 110–127), T-cell recognition epitope 1 (aa 117–131), T-cell recognition epitope 2 (aa

150–163), B-cell epitope (aa 180–200), and the stop codon (aa 201) (Figure 4).

Multiple amino acid substitutions were identified in the antigenic regions of the synthesized

glycoprotein. Specifically, several neutral substitutions were observed in the 27 aa corresponding to

the signal peptide, as well as nonsynonymous substitutions with important functional implications in

the early antibody response mediated by decoy epitope I (A), particularly in hypervariable region 1.

These substitutions included 15/24 at S32N/G/R/E, 24/24 at D34N/S/T, and 5/24 at S35N/I, which

may induce rapid and potent non-neutralizing antibodies. Decoy epitope II also showed important

amino acid substitutions (24/24 at P185A, 24/24 at L189V, 2/24 at T190P, and 24/24 at R191K), which

play an important role in cross-neutralization responses to antibodies generated against species 1 and

2.

In addition, regions responsible for the production of neutralizing antibodies were analyzed.

Neutralizing epitope B (PNE) is characterized by a highly conserved region comprising antigenic

determinant sites (aa 39, 41, and 44). Our results showed substitutions in 7/24 isolates at aa L39F, 6/24

at aa Q40L/R, 3/24 at aa L41Y/V, 4/24 at aa N44I, and 24/24 at aa L47I/V. Neutralizing epitope C,

together with its hypervariable region 2, includes antigenic determinants at positions aa 57, 58, 59, and

61, where substitutions enable escape from neutralizing antibodies. Substitutions were observed in

24/24 isolates at A57N, 24/24 at N58K/E/V/R, and 24/24 at N59H/S/R/N. Amino acid changes in

epitopes A, B, and C of GP5 in field strains may reduce the protective effect of commercial PRRS

vaccines.

Statistical analysis: Tajima's D test and genetic diversity using DnaSP v6

Genetic diversity analysis using the DnaSP program revealed 860 mutations, 530 polymorphic sites,

and 61 conserved sites across the sequences. In addition, Tajima's D test yielded a value of −0.78746,

which was not statistically significant (P > 0.10). This negative value indicates that the hypothesis of

neutral evolution should not be rejected and suggests the presence of recombination or a considerable

incidence of population subdivision with deleterious mutations. The applied genetic diversity analyses

demonstrated viral variability and plasticity, and the results, compared with the Lineage 5 VR2332

strain, are presented in Table 2.

DISCUSSION

The results showed the identification of 24 isolates belonging to the species *Betaarterivirus*

americense. Phylogenetic inference was performed using 58 sequences from major epidemic outbreaks

worldwide retrieved from databases, revealing the presence of lineages 1, 5, and 8. In addition, the

relationship observed between the 24 isolates and the Peruvian reference strains showed genetic

divergence within Lineage 1, specifically sublineage 1.5 of the NADC34 variant, according to

glycoprotein 5 studies and the methods described by Nguyen [19] and Shin [26]. Our data are

consistent with those reported by Ramírez [23], who showed that Peruvian isolates, based on RFLP

typing, are related to the 1-7-4 variant of Lineage 1, specifically sublineage 1.5 of NADC34.

Studies of the structural protein GP5 showed that antigenic domains play an important role in the

immune response due to synonymous and nonsynonymous substitutions occurring in key

immunological domains involved in the production of non-neutralizing antibodies, such as decoy

epitopes I (A) and II, and neutralizing epitopes such as B and C, which contribute to immune evasion

and neutralization [17,21]. In addition, the results demonstrated different degrees of substitution in

other domains, including the signal peptide; transmembrane regions 1, 2, and 3; T-cell recognition

epitopes (1 and 2); and the B-cell epitope, when compared with Lineage 5 of the VR2332 variant.

Our results showed 15/24, 24/24, and 5/24 substitutions at aa S32N/G/R/E, D34N/S/T, and S35N/I,

respectively, in the hypervariable region 1 of the decoy epitope (A). The high variability of this region,

according to Nguyen [19], induces rapid and strong non-neutralizing antibody responses. That is, the

decoy epitopes aa27–35 and aa180–197 fulfill an immune evasion function: when pigs are infected

with the virus, these two decoy epitopes first stimulate immune system cells to rapidly produce large

amounts of non-neutralizing antibodies against these epitopes, thereby delaying the production of

effective antibodies [35]. Furthermore, our findings of 7/24, 6/24, 3/24, 4/24, and 24/24 substitutions

in amino acid regions L39F, Q40L/R, L41Y/V, N44I, and L47I/V of the primary neutralizing epitope

(B) suggest high pathogenicity of our isolates, since substitutions in this region play an important role

in immune neutralization [14]. These changes help PRRS field strains escape vaccine-induced

neutralizing immunity, and mutation at aa39 has been used to classify viral variants according to Zhou

[35]. In addition, studies by Faaberg [9] identified aa38, aa42, aa43, and aa44 as neutralizing activity

sites, and aa39, aa40, and aa41 as antibody-binding sites. According to Chen [4], mutations in these

regions can reduce the sensitivity of PRRSV-2 to neutralizing antibodies in vivo, thereby affecting

immune clearance of viral particles and potentially contributing to vaccination failure and increased
morbidity in swine herds. Finally, our results showed 24/24 substitutions at A57N, 24/24 substitutions
at N58K/E/V/R, and 24/24 substitutions at N59H/S/R/N in the neutralizing epitope C and its
hypervariable region 2, where such substitutions enable escape from neutralizing antibodies [19].
Overall, these data demonstrate a high incidence of mutations located in key immunological domains
within epitopes A, B, and C of GP5, which are responsible for the production of non-neutralizing and
neutralizing antibodies in the field strains isolated in this study. Previous studies by Nguyen [19]
showed that substitutions in these domains reduce the protective effect of commercial PRRS vaccines.
This represents important information regarding the high pathogenicity of Peruvian isolates and the
potential for recombination with vaccine strains if applied, which could lead to the emergence of new,
more pathogenic variants in Peru.

Another major role of GP5 is the glycosylation of membrane proteins. This process involves the
addition or removal of glycan residues at potential N-glycosylation sites in PRRSV species. The N34
site is more prone to mutation, whereas N44 and N51 are highly conserved, and mutations at N44 may
lead to increased virulence [4]. Previous studies, such as those by Ansari [2], have shown that removal
of N-glycosylation sites increases the susceptibility of PRRSV species to neutralization by specific
immune antibodies. Therefore, an increased number of potential N-glycosylation sites may explain the
continuous mutation of viral species. These statements are consistent with the findings of the present
study, as the analysis revealed 10 potential N-glycosylation sites (N30, N32, N33, N34, N35, N44,
N51, N57, N128, and N151) in the extravirion region of the glycoprotein in all 24 isolated strains.

The large number of potential N-glycosylation sites in the study strains was consistent with the results
reported by Li [16], but contrasts with the findings of Xie [33] and Chen [4], who reported lower
numbers of N-glycosylation sites. However, in this study, strains 36, 37, 38, 39, 43, and 44 exhibited
three N-glycosylation sites, whereas strains 40 and 41 showed six N-glycosylation sites with greater
observable variability. According to Ansari [2], the three conserved potential glycosylation sites (N34,
N44, and N51) contribute to viral variability and failure of control by commercial vaccines, and they
indicate that glycan addition at site N44 is particularly critical because it enhances viral escape from
antibodies. Furthermore, these authors argue that PRRSV species containing N-hypoglycosylated
forms of GP5 are sensitive to antibody neutralization, and that removal of glycosylation sites in
laboratory-generated mutant viruses induces significantly higher levels of neutralizing antibodies by
the immune system.

The data obtained in this study suggest that hypoglycosylated wild-type viruses are less common than
highly glycosylated strains, which may explain the expansion of viral variants. The effect of N-
glycosylation suggests that glycans mask one or more neutralizing epitopes, resulting in the failure to
induce neutralizing antibodies when glycans are present [2]. In addition, only strains 40 and 41 retained

all three glycosylation sites (N34, N44, and N51), whereas strains 36, 37, 38, and 39 lost them. Moreover, conserved potential N-glycosylation sites N44 and N51 were present in 20 of the 24 analyzed isolates, and 18 of the 24 had more than three N-glycosylation sites. According to Zhang [34], a higher number of N-glycosylation sites may contribute to reduced immunogenicity of nearby neutralizing epitopes, suggesting that these isolates may escape immune recognition and neutralization by vaccine-induced antibodies. This observation is highly relevant for the design of effective vaccines against emerging variants, suggesting that newly designed vaccines targeting these regions should incorporate modifications in the glycosylation pattern of the GP5 gene to enhance the production of neutralizing antibodies [2]. Nevertheless, further studies are required to determine whether mutation of glycosylation sites affects exposure of viral neutralizing epitopes, as well as immune evasion and viral pathogenicity [4].

Genetic diversity analysis of the 24 strains using the DnaSP program showed rapid evolution and a high mutation rate in Peruvian isolates, with 530 variable sites and 860 mutations. In addition, Tajima's D test revealed a considerable incidence of population subdivision, yielding a D value of -0.78746 [24]. Overall, our results reflect the reality of rapid evolution and spread of PRRSV variants in Peru and help explain the ongoing problem of vaccine inefficacy due to immune evasion mechanisms associated with increased N-glycosylation sites in Peruvian variants.

CONCLUSIONS

- Through phylogenetic analysis, we identified the relationship between the 24 study isolates and the Peruvian reference strains with Lineage 1, sublineage 1.5 of the NADC34 variant, based on glycoprotein 5 alignment.
- We identified a high number of significant substitutions in immunological domains involved in the production of non-neutralizing and neutralizing antibodies: aa34 (24/24), aa39 (6/24), aa44 (4/24), aa47 (24/24), aa57 (24/24), aa58 (24/24), and aa59 (24/24), located in epitopes A, B, and C of GP5.
- We identified 10 potential N-glycosylation sites (N30, N32, N33, N34, N35, N44, N51, N57, N128, and N151) in the 201–amino acid sequence of glycoprotein 5.
- We identified substitutions, losses, and conservation of N-glycosylation sites (N34, N44, and N51), with partial conservation in most strains or complete conservation in strains 40 and 41, and complete loss in strains 36, 37, 38, and 39.
- We identified that the 24 strains exhibit rapid evolution and a high mutation rate in Peruvian isolates, with 530 variable sites and 860 mutations.

305   • Using statistical analysis with Tajima's D test, we determined a considerable incidence of
306   population subdivision, yielding a D value of −0.78746.
307

308 **SUPPLEMENTARY MATERIAL**

309   • Supplementary Material 1. Amino acid sequence alignment of glycoprotein 5 using MEGA 6.
310

311 **CONFLICT OF INTEREST**

312   • The authors declare that they have no conflicts of interest in submitting this article for
313   publication.
314

315 **FUNDING**

319 **ACKNOWLEDGMENTS**

320   • The authors thank Montana S.A. for funding and support in conducting this research, and
321   Bach. Segundo Del Águila Soto for logistical support and valuable suggestions.
322

REFERENCES

[1]. Allende, R., Kutish, G. F., Laegreid, W., Lu, Z., Lewis, T. L., Rock, D. L., Friesen, J., Galeota, J.
A., Doster, A. R., & Osorio, F. A. (2000). Mutations in the genome of porcine reproductive and
respiratory syndrome virus responsible for the attenuation phenotype. *Archives of Virology,*
145(6), 1149–1161. https://doi.org/ 10.1007/s007050070115
[2]. Ansari IH, Kwon B, Osorio FA, Pattnaik AK. 2006. Influence of N-linked glycosylation of porcine
reproductive and respiratory syndrome virus GP5 on virus infectivity, antigenicity, and ability to
induce neutralizing antibodies. J Virol 80(8):3994–4004. https://doi.org/10.1128/JVI.80.8.3994-
4004.2006
[3]. Brinton MA, Gulyaeva AA, Balasuriya U, Dunowska M, Faaberg KS, Goldberg T, Leung F,
Nauwynck HJ, Snijder EJ, Stadejek T, Gorbalenya AE. 2021. ICTV Virus Taxonomy
Profile: *Arteriviridae* 2021. J Gen virol 102(8): 001632. https://doi.org/ 10.1099/jgv.0.001632
[4]. Chen XW, Li L, Yin M, Wang Q, Luo WT, Ma Y, Pu ZH., Zhou JL. 2017. Cloning and molecular
characterization of the ORF5 gene from a PRRSV-SN strain from Southwest China. Microb
pathog 112:295–302. https://doi.org/10.1016/j.micpath.2017.09.011

[5]. Cotaquispe, N. R., De la Cruz, V. E., Quispe, C. E., Sitaras, N., & Toskano, H. J. (2022). Molecular characterization of porcine reproductive and respiratory syndrome (PRRS) based on the ORF7 nucleocapsid gene in swine farms in Lima, Peru. *Revista de Investigaciones Veterinarias del Perú, 33*, e21638. http://dx.doi.org/10.15381/rivep.v33i6.21638.

[6]. Cotaquispe, N. R., Legua, B. M., Escajadillo, L. P., et al. (2025). Molecular characterization of the ORF5 (GP5) gene of the causative agent of porcine reproductive and respiratory syndrome (PRRS) detected in swine farms in Lima, Peru. *Revista Argentina de Microbiología*. https://doi.org/10.1016/j.ram.2025.07.005

[7]. Dea S, Gagnon CA, Mardassi H, Pirzadeh B, Rogan D. 2000. Current knowledge on the structural proteins of porcine reproductive and respiratory syndrome (PRRS) virus: comparison of the North American and European isolates. Arch Virol 145(4): 659–88.

[8]. Dokland T. 2010.The structural biology of PRRSV. Virus Re 154(1–2):86-97. https://doi.org/10.1016/j.virusres.2010.07.029

[9]. Faaberg KS, Hocker JD, Erdman MM, Harris DL, Nelson EA, Torremorell M, Plagemann PG. 2006. Neutralizing antibody responses of pigs infected with natural GP5 N-glycan mutants of porcine reproductive and respiratory syndrome virus. Viral immunol 19(2):294–304. https://doi.org/10.1089/vim.2006.19.294

[10]. Fang Y, Snijder EJ. 2010. The PRRSV replicase: exploring the multifunctionality of an intriguing set of nonstructural proteins. Virus Res 154:61–76 4.

[11]. Fang Y, Trefers EE, Li Y, Tas A, Sun Z, van der Meer Y, de Ru AH, van Veelen PA, Atkins JF, Snijder EJ, Firth AE. 2012. Efcient −2 frameshifting by mammalian ribosomes to synthesize an additional arterivirus protein. Proc Natl Acad Sci USA 109:E2920–E2928

[12]. Gupta R, Brunak S. 2002. Prediction of glycosylation across the human proteome and the correlation to protein function. Pac Symp Biocomput 310–322. PMID: 11928486.

[13]. Holtkamp DJ, Kliebenstein JB, Neumann EJ, Zimmerman JJ, Rotto HF, Yoder TK, Wang C, Yeske PE, Mowrer CL, Haley CA .2013. Assessment of the economic impact of porcine reproductive and respiratory syndrome virus on United States pork producers. J Swine Health Prod 21:72–84.

[14]. Li B, Fang L, Xu, Z, Liu S, Gao J, Jiang Y, Chen H, Xiao S. 2009a. Recombination in Vaccine and Circulating Strains of Porcine Reproductive and Respiratory Syndrome Viruses. Emerg Infect Dis 15(12):2032-2035. https://doi.org/10.3201/eid1512.090390

[15]. Li B, Xiao S, Wang Y, Xu S, Jiang Y, Chen H, Fang L. 2009b. Immunogenicity of the highly pathogenic porcine reproductive and respiratory syndrome virus GP5 protein encoded by a synthetic ORF5 gene. Vaccine 27(13): 1957–1963. https://doi.org/10.1016/j.vaccine.2009.01. 098

[16]. Li B, Fang L, Guo X, Gao J, Song T, Bi J, He K, Chen H, Xiao S. 2011. Epidemiology and Evolutionary Characteristics of the Porcine Reproductive and Respiratory Syndrome Virus in

China between 2006 and 2010. J Clin Microbiol 49(9):3175–3183. https://doi.org/10.1128/JCM.00234-11

[17]. Martínez-Bautista NR, Sciutto-Conde E, Cervantes-Torres J, Segura-Velázquez R, Mercado García MC, Ramírez-Mendoza H, Trujillo Ortega ME, Delgadillo Álvarez J, Castillo-Juárez H, Sanchez-Betancourt JI. 2018. Phylogenetic analysis of ORF5 and ORF7 of porcine reproductive and respiratory syndrome (PRRS) virus and the frequency of wild-type PRRS virus in México. Transbound Emerg Dis 65: 993– 1008. https://doi.org/10.1111/tbed.12831

[18]. Murtaugh MP, Stadejek T, Abrahante JE, Lam TT, Leung FC. 2010. The ever-expanding diversity of porcine reproductive and respiratory syndrome virus. Virus Res 154(1-2):18–30. https://doi.org/10.1016/j.virusres.2010.08.015

[19]. Nguyen NH, Tran H, Nguyen TQ, Nguyen P, Le T, Lai DC, Nguyen MN. 2022. Phylogenetic analysis of porcine reproductive and respiratory syndrome virus in Vietnam, 2021. *Virus genes 58*(4):361–366. https://doi.org/10.1007/s11262-022-01912-w

[20]. Park J, Choi S, Jeon JH, Lee KW, Lee C. 2020. Novel lineage 1 recombinants of porcine reproductive and respiratory syndrome virus isolated from vaccinated herds: genome sequences and cytokine production profiles. Arch virol 165(10):2259–2277. https://doi.org/10.1007/s00705-020-04743-y

[21]. Plagemann PG. 2004. The primary GP5 neutralization epitope of North American isolates of porcine reproductive and respiratory syndrome virus. Vet immunol immunopathol 102(3):263–275. https://doi.org/10.1016/j.vetimm.2004.09.011

[22]. Petersen TN, Brunak S, von Heijne G, Nielsen H. 2011. SignalP 4.0: discriminating signal peptides from transmembrane regions. Nat methods 8(10):785–786. https://doi.org/10.1038/nmeth.1701

[23]. Ramírez M, Bauermann FV, Navarro D, Rojas M, Manchego A, Nelson EA, Diel DG, Rivera H. 2019. Detection of porcine reproductive and respiratory syndrome virus (PRRSV) 1 7 4 type strains in Peru. Transbound Emerg Dis 66: 1107-1113. https://doi.org/10.1111/tbed.13134

[24]. Rozas J, Ferrer-Mata A, Sánchez-DelBarrio JC, Guirao-Rico S, Librado P, Ramos-Onsins SE, Sánchez-Gracia A. 2017. DnaSP 6: DNA Sequence Polymorphism Analysis of Large Data Sets. Mol Biol Evol 34(12):3299–3302. https://doi.org/10.1093/molbev/msx248

[25]. Shi M, Lam TT, Hon CC, Murtaugh MP, Davies PR, Hui RK, Li J, Wong LT, Yip CW, Jiang JW, Leung FC. 2010. Phylogeny-based evolutionary, demographical, and geographical dissection of North American type 2 porcine reproductive and respiratory syndrome viruses. J Virol 84: 8700–8711. https://doi.org/10.1128/JVI.02551-09

[26]. Shin G, Park J, Lee K, Ko M, Ku B, Park C, Jeoung H. 2021. Genetic Diversity of Porcine Reproductive and Respiratory Syndrome Virus and Evaluation of Three One-Step Real-Time RT-PCR Assays in Korea. Research Square 1-20. doi:10.21203/rs.3.rs-1149203/v1. PPR:PPR437459.

[27].   Snijder EJ, Kikkert M, Fang Y. 2013. Arterivirus molecular biology and pathogenesis. J Gen Virol 94: 2141-2163. doi: 10.1099/vir.0.056341-0

[28].   Tamura K, Stecher G, Peterson D, Filipski A, Kumar S. 2013. MEGA6: Molecular Evolutionary Genetics Analysis version 6.0. Mol biol Evol 30(12): 2725–2729.

[29].   Tajima F. 1989. Statistical method for testing the neutral mutation hypothesis by DNA polymorphism. Genetics 123(3):585–595. PMC1203831

[30].   Tajima F. 1993. Statistical analysis of DNA polymorphism. Idengaku zasshi 68(6):567–595. https://doi.org/10.1266/jjg.68.567

[31].   Wootton S, Yoo D. Rogan D. 2000. Full-length sequence of a Canadian porcine reproductive and respiratory syndrome virus (PRRSV) isolate. Arch virol 145(11): 2297–2323.

[32].   Wu WH, Fang Y, Rowland RR, Lawson SR, Christopher-Hennings J, Yoon KJ, Nelson EA. 2005. The 2b protein as a minor structural component of PRRSV. Virus Res 114:177–181

[33].   Xie J, Zhu W, Chen Y, Wei C, Zhou P, Zhang M, Huang Z, Sun L, Su S, Zhang G. 2013. Molecular epidemiology of PRRSV in South China from 2007 to 2011 based on the genetic analysis of ORF5. Microb pathog 63:30–36. https://doi.org/10.1016/j.micpath.2013.05.013

[34].   Zhang L, Feng Y, Martin DP, Chen J, Ma S, Xia P, Zhang G. 2017. Genetic diversity and phylogenetic analysis of the ORF5 gene of PRRSV from central China. Res Vet Sci 115:226–234. https://doi.org/10.1016/j.rvsc.2017.05.013

[35].   Zhou L, Kang R, Ji G, Tian Y, Ge M, Xie B, Yang X, Wang H. 2018. Molecular characterization and recombination analysis of porcine reproductive and respiratory syndrome virus emerged in southwestern China during 2012-2016. Virus genes 54(1):98–110. https://doi.org/10.1007/s11262-017-1519-y

Figure 1. In-house conventional RT-PCR of the ORF5 gene designed in the laboratory. Lane 1 (ORF5-SKA): positive cDNA from the Nebraska Prime Pac® PRRS-2 strain; Lane 4 (ORF5-SKA): positive cDNA from the VP-046 SUIPRABAC® PRRS-1 strain; Lane 7 (ORF5-SKA): positive control cDNA mixture of PRRS-1 and PRRS-2 from the IDEXX commercial kit; Lanes 2, 5, and 8 (avian bronchitis): negative cDNA controls from IBV; Lanes 3, 6, and 9 (PCR blank): nuclease-free water; Lane M (ladder): 100 bp Plus Opti-DNA Marker ABM®. Image adapted from Cotaquispe et al., 2025.

Figure 2. In-house conventional RT-PCR of the ORF5 gene designed in the laboratory. Lane 1 (ORF5-SKA): cDNA from a PRRS field-positive sample; Lane 2 (avian bronchitis): negative control cDNA from IBV; Lane M (ladder): 100 bp Plus Opti-DNA Marker ABM® (50 bp–1.5 kb).

488 Table 1. Results of the genotypic variability of 24 isolates of Porcine Reproductive and Respiratory
489   Syndrome virus (PRRSV) from blood samples of pigs from the Lima region, Peru.

cDNA	GP5 (aa)	Genotyping	N-glycosylation							Signal peptide
			#	1	2	3	4	5	6	
25	201	Lineage 1: Sublineage 1.5	5	30	33	44	51	57		1-31
26	201	Lineage 1: Sublineage 1.5	5	30	33	44	51	57		1-31
27	201	Lineage 1: Sublineage 1.5	5	30	33	44	51	57		1-31
28	201	Lineage 1: Sublineage 1.5	5	30	33	44	51	57		1-31
29	201	Lineage 1: Sublineage 1.5	4	30	33	44	51			1-31
30	201	Lineage 1: Sublineage 1.5	5	32	35	44	51	57		1-31
31	201	Lineage 1: Sublineage 1.5	5	32	35	44	51	57		1-31
32	201	Lineage 1: Sublineage 1.5	5	32	35	44	51	57		1-31
33	201	Lineage 1: Sublineage 1.5	5	32	35	44	51	57		1-31
34	201	Lineage 1: Sublineage 1.5	5	30	33	44	51	57		1-31
35	201	Lineage 1: Sublineage 1.5	5	30	33	44	51	57		1-31
36	201	Lineage 1: Sublineage 1.5	3	33	50	128				1-31
37	201	Lineage 1: Sublineage 1.5	3	33	50	128				1-31
38	201	Lineage 1: Sublineage 1.5	3	33	50	128				1-31
39	201	Lineage 1: Sublineage 1.5	3	33	50	128				1-31
40	201	Lineage 1: Sublineage 1.5	6	30	33	34	44	51	57	1-31
41	201	Lineage 1: Sublineage 1.5	6	30	33	34	44	51	57	1-31
42	201	Lineage 1: Sublineage 1.5	5	33	44	51	57	151		1-31
43	201	Lineage 1: Sublineage 1.5	3	44	51	57				1-31
44	201	Lineage 1: Sublineage 1.5	3	44	51	57				1-31
45	201	Lineage 1: Sublineage 1.5	4	32	33	44	51			1-31
46	201	Lineage 1: Sublineage 1.5	4	32	33	44	51			1-31
47	201	Lineage 1: Sublineage 1.5	4	32	33	44	51			1-31
48	201	Lineage 1: Sublineage 1.5	5	32	33	44	51	57		1-31

Figure 3.Phylogenetic tree of *Betaarterivirus europense* and *Betaarterivirus americense* species based on the glycoprotein 5 (GP5) sequence. (a) 1. Purple branching: *Betaarterivirus suid 1*; 2. Turquoise branching (Lineage 1): sublineage 1.1 (98-3584), sublineage 1.5 (NADC34, study isolates and Peruvian reference strains), sublineage 1.6 (NADC31), sublineage 1.8 (HENAN), sublineage 1.9 (MN184A); 3. Light green branching (Lineage 5): sublineage 5.1 (Res-PRRS MLV and VR2332); 4. Dark blue branching (Lineage 8): sublineage 8.7 (JXA1). (b) Amino acid substitution analysis was performed using the maximum likelihood method applying BioNJ with the Jones–Taylor–Thornton (JTT) model. Bootstrapping with 1,000 alignment replicates was used, and evolutionary analyses were conducted in MEGA6 and DNAMAN 10.

Table 2. Results of the genetic diversity of 24 isolates of Porcine Reproductive and Respiratory Syndrome virus (PRRSV) analyzed using the DnaSP and MEGA 6 programs from blood samples of pigs from the Lima region, Peru.

Strains		ORF 5 (GP5)	C.S.	Variable sites	P. I. S	U. V. S	Mutations
24	nt	603	61	530	48	82	860
	aa	201					

nt: nucleotide; **aa**: amino acid; **C.S**.: conserved sites; **P.I.S**.: parsimony-informative sites; **U.V.S**: unique variable sites.

Figure 4. Alignment of amino acid sequences from antigenic regions in PRRSV GP5; the boxed and colored regions represent antibody-neutralizing antigens. The 24 field isolates and representative variants of sublineages 1.1 (99-3584), 1.5 (NADC34), 1.6 (NADC31), 1.8 (HENAN), 1.9 (MN184A), and sublineage 5.1 (VR2332) were used as reference patterns in the alignment. Dots indicate the position of homologous amino acids, and letters indicate amino acid substitutions. Yellow, orange, and blue boxes indicate the positions of non-neutralizing and neutralizing epitope amino acids.