1 **Bile Acids Regulate Accumbal Cholinergic Circuitry and Dopamine Release through TGR5 Activation**
2

3 AUTHORS:

4 Isabella A. Roque[1], Shimran S. Sharma[1,3], Philipp Mews[7], Summer L. Thompson[4,5,6], Jordan T. Yorgason[1,2,3*]
5

1. Neuroscience Center, Brigham Young University, Provo, Utah, 84602, USA.**

2. Department of Cell Biology and Physiology, Brigham Young University, Provo, Utah, 84602, USA.

3. Department of Translational Neuroscience, Wake Forest University School of Medicine, Advocate Health,
Winston-Salem, North Carolina, 27101, USA.

4. Department of Psychiatry, University of Maryland School of Medicine, Baltimore, Maryland, 21201, USA.

5. Department of Neurobiology, University of Maryland School of Medicine, Baltimore, Maryland, 21201, USA.

6. Kahlert Institute for Addiction Medicine, University of Maryland School of Medicine, Maryland, 21201, USA.

7. Department of Pharmacology, Physiology, and Biophysics, Boston University Chobanian & Avedisian School of
Medicine, Massachusetts, 02118, USA.

*Corresponding author: Jordan Yorgason

**Work performed in this location

18
19 Keywords: striatum, TGR5, dopamine, liver, gut-brain axis, patch-clamp, fatty foods, ethanol, voltammetry,
20 electrophysiology
21
22
23
24
25 Number of Figures: 5
26
27
28 Abstract Word Count: 246
29
30
31 Total Word Count: 5645

Abstract

Background: High calorie foods and alcohol produce strong reinforcing effects. Brain mechanisms for these reinforcers have focused largely on indirect mechanisms on mesolimbic circuits. Bile acid secretion is a major factor during caloric intake, however, the role of biliary acids on mesolimbic circuitry remains unexplored. Brain mesolimbic circuits are modulated in part by altering cholinergic interneuron (CIN) activity and tonic dopamine (DA) transmission within the nucleus accumbens (NAc). Ethanol and fatty foods also both stimulate hepatic and possibly local brain bile acid (BA) synthesis, which raises the possibility that BAs may act as a common upstream regulator of these substances shared mesolimbic effects.

Methods: This study investigated whether BAs can directly alter mesolimbic activity. We collected electrophysiological data from acute mouse brain slices to assess BA effects on accumbal CIN firing and excitatory and inhibitory postsynaptic activity. Bile effects on DA release and clearance in the NAc were measured through voltammetry.

Results: We found that low concentrations of a 1:1 mixture of BAs cholic acid and deoxycholic acid (1-10 μM) increased CIN firing rate whereas high BA concentrations (1-10 mM) decreased CIN firing. We further demonstrated that BA-induced excitatory effects on CIN firing are independently mediated by at least two mechanisms: Takeda G-protein-coupled receptor 5 (TGR5) activation and suppression of inhibitory CIN currents. Additionally, our results indicate that BAs modulate inhibitory input in a complex manner, reducing frequency at low concentrations, but increasing at high concentrations, and increasing amplitude at low concentrations current amplitude, and the distribution of postsynaptic current amplitude sizes across concentrations. Finally, our voltammetry data show that while low BA concentrations enhance NAc DA release without affecting uptake, whereas high concentrations of BA robustly inhibit accumbal DA release.

Conclusion: Our findings provide evidence that BAs exert direct modulatory effects on neural activity in the striatum.

1. Introduction

Fatty foods and ethanol have strong reinforcing effects, implicating changes in mesolimbic activity. High-fat diet (HFD) has been shown to increase cholinergic interneuron (CIN)-evoked accumbal dopamine (DA) release in rat models [1]. In parallel, ethanol has been observed to both directly increase and decrease DA release and CIN firing in the nucleus accumbens (NAc), effects that are largely dose-dependent and appear to involve multiple mechanisms [2]. Together, these findings indicate that both fatty foods and ethanol disrupt CIN-DA interactions in the NAc. Given the established connection between CIN activity, DA release in the striatum, and reward-related behavior [3], identifying a common upstream regulator for fatty food- and ethanol-induced CIN-DA effects may provide insight into the mechanisms underlying their common reinforcing properties.

Notably, both fatty foods and ethanol promote hepatic, and possibly local brain, bile acid (BA) synthesis, making BAs a potential upstream regulator of these substances' shared striatal effects [4-7]. Consumption of 1–2 standard alcohol doses increases hepatic BA production by ~20% and elevates systemic BA precursors 5–15-fold [5, 8]. Alcohol intake also raises cholesterol levels in the brain, providing more precursors for BA synthesis in the Central Nervous System (CNS) [4]. Similarly, HFD raises systemic BA levels by up to 50% in rats and increases levels of brain BA precursor 27-hydroxycholesterol (27-OHC) and BA-synthesizing enzyme cholesterol 27α-hydroxylase (CYP27A1) [6, 7]. While brain-derived BAs can act locally, hepatic BAs must enter systemic circulation and cross the blood-brain barrier (BBB) to influence activity in the CNS [9]. Approximately 10% of BAs exiting the intestines reach systemic circulation [10]. Bile in systemic circulation can subsequently reach the BBB and cross into the CNS either via BA transporters or passive diffusion [11, 12]. There is strong evidence indicating that hepatic BAs cross the BBB and enter the CNS. Brain and serum BA concentrations are significantly correlated, and the presence of BAs such as DCA—a secondary BA that can only be synthesized outside the CNS—has been detected in brain tissue [13, 14]. The liver produces primary BAs, like cholic acid (CA), and intestinal microbiota converts these into secondary BAs, like deoxycholic acid (DCA). In contrast, the brain only expresses enzymes for primary BA synthesis [9, 13]. Once they reach the CNS, both CA and DCA can activate BA receptors present in the brain, such as Takeda G-protein-coupled receptor 5 (TGR5), Farnesoid X receptor (FXR), and Sphingosine-1-phosphate receptor 2 (S1PR2) [9,15-16]. Importantly, there is evidence that TGR5 expression in the NAc contributes to cocaine's rewarding effects, highlighting a putative role for Bas in regulating mesolimbic activity [16].

This study used a 1:1 mixture of CA and DCA to examine general BA effects on mesolimbic circuitry, irrespective of their potential brain- or hepatic-derived source. We assessed BA effects on CIN firing and DA release and evaluated the role of the BA receptor TGR5 in mediating these effects. Our findings demonstrate that BAs exert significant mesolimbic effects and have strong potential to contribute to fatty foods' and ethanol's shared appetitive effects.

2. Materials and Methods

2.1. Animals

Male and female mice on a C57BL/6J background (Jackson Laboratory) were kept in a 12:12-h light/dark cycle and given access to water and standard rodent chow *ad libitum.* Animal care and experimental procedures were approved by Brigham Young University Institutional Animal Care and Use Committee and were performed in accordance with the National Institutes of Health Guide for the Care and Use of Laboratory Animals.

2.2. Brain Slice Preparation and Electrophysiology Recordings

Fully isoflurane (Patterson Veterinary, Devens, MA) anesthetized mice (PND>30) were decapitated and brains were quickly extracted and sliced into 220 µm thick striatal coronal brain slices on a vibratome (Leica VT1000S, Vashaw Scientific, Norcross, GA) surrounded by pre-oxygenated ACSF (95% O_2/5% CO_2; pH≈7.4, 35°C) and 300 µM ketamine to preserve cell health. The artificial cerebrospinal fluid (ACSF) contained (in mM): KCl (2.5), NaCl (126), glucose (11), $NaHCO_3$ (25), NaH_2PO_4 (1.2), $CaCl_2$ (2.4), $MgCl_2$ (1.2). After a short incubation period (~30 minutes), slices were transferred to the recording chamber and perfused with ACSF (34°C) at a rate of ~1.8 ml/min. Cell-attached recordings were made with glass micropipettes (2.5-6 MΩ) pulled on a P97 Pipette puller (Sutter Instruments) and subsequently filled with ACSF (cell attached recordings) or artificial intracellular solution (whole cell recordings). Intracellular solution for whole cell recordings contained (in mM): KCl (145), NaCl (8), MgCl2 (0.2), EGTA (2), HEPES (10), and Mg-ATP (2). Whole-cell recordings were performed to isolate spontaneous inhibitory postsynaptic currents (sIPSCs; in kynurenic acid) and excitatory postsynaptic currents (sEPSCs; in picrotoxin). CINs were identified based on their large size, tonic firing, and sensitivity to muscarine. Drugs were applied as specified and for at least 10 minutes to allow for stabilization. Data

were filtered at the amplifier with a 4-pole high-pass Bessel filter at 2–10 kHz (Axon Multiclamp 700A; Molecular
Devices, Sunnyvale CA, USA) and digitized via a NIDAQ board (PCIe-6321; National Instruments) sampled
continuously at 10kHz using Axograph Software 1.7.6. (Axograph Scientific, Sydney Australia). All cell-attached
experiments consisted of continuous 1 h recordings, with a 5 min baseline and sequential 10 min exposures to each
CA/DCA concentration (1 µM–10 mM). Whole-cell recordings consisted of shorter 15- and 25-min recordings to
prevent confounds from re-sealing of the cell membrane [17]. The following concentrations of drugs were bath
applied for electrophysiology and slice voltammetry experiments where specified: CA/DCA (1 µM - 10 mM;
Millipore Sigma), 5-chloro-2-(ethylsulfonyl)-4-pyrimidinecarboxylic acid 3-methylphenyl ester (SBI-115; 10 µM;
Cayman Chemical), kynurenic acid (KA, 2 mM; Millipore Sigma), picrotoxin (100 µM; Millipore Sigma).
131
2.3. Carbon Fiber Electrode Preparation and Voltammetry Recordings
Slices were prepared as described above, transferred to the recording chamber, and perfused with aCSF
(34°C) at a rate of ~1.8 ml/min. Fast scan cyclic voltammetry recordings were performed and analyzed using Demon
Voltammetry and Analysis software (Yorgason, Zeppenfeld, & Williams, 2017)(RRID:SCR_014468). The carbon
fiber electrodes used for voltammetry experiments were manufactured in-house. Carbon fiber electrodes (7 µm,
Thornel T-650, Cytec, Woodland Park, NJ) were aspirated into borosilicate glass capillaries (TW150, World
Precision Instruments, Sarasota, FL) and pulled on a pipette puller and carbon fibers were cut to a length of 100-150
µm from the tip of the glass tube. Stimulating electrodes were prepared by pulling a glass micropipette and filled
with ACSF. DA release was evoked in the NAc with a 1 pulse (30-100 µA; 0.5 ms) electrical stimulus (1/min) and
detected via fast-scan cyclic voltammetry (100 ms cycle, 400 V/s). Concentration response curve experiments had
an initial baseline (BL) stabilization period (~18-20 min), followed by 10 min exposures to CA/DCA in increasing
concentrations (1 µM–10 mM). For single concentration experiments, a similar BL stabilization period was used,
followed by a ~20 min exposure to 1-10 µM CA/DCA.
145
2.4. Statistical Analysis
Raw patch-clamp (Axograph) and voltammetry (Demon) data were exported to Excel for CIN
firing/current rate, CIN firing variance and DA release averaging. Averages were normalized against respective BLs
and transferred to GraphPad Prism 5 (GraphPad, San Diego, CA) or NCSS 8 (NCSS LLC, Kaysville UT) for
statistical analysis. Since amplitude is an event-level property, sIPSCS/sEPSC current amplitude values were
transferred directly from Excel to GraphPad without averaging. Voltammetry mediation and OLS regression were
run in Python. Mediation analysis
Was used to test how bile affected DA release overall, specifically if the effect was (1) direct (without CIN
involvement), or (2) indirect (with CIN involvement). To test for direct/indirect effects, the regression coefficients
represent the average effect of bile on DA release and CIN firing rate per unit change in bile concentration. c' (direct
effect) represents the effect of bile on DA when you control for CIN firing, and a*b (indirect effect) represents the
portion of bile's effect on DA that occurs due to CIN involvement.
158
This is a summary of all the coefficients:
a = bile → CIN
b = CIN → DA (controlling for bile)
c' = bile → DA (controlling for CIN)
a × b = indirect effect (through CIN)
c = c' + a × b = total effect
165
The predicted lines on the graph are predicted values (based on the original values, calculated through the mediation
analysis). All remaining statistical analyses were performed in GraphPad. One- and two-way ANOVA and two-
tailed unpaired t-tests were used, with Dunnett's test for planned comparisons, or Tukey's HSD or Bonferroni post-
hoc analysis where stated. Outliers were identified using GraphPad's outlier calculator. Significance levels are
indicated on graphs with asterisks *,**,*** and correspond to significance levels $p < 0.05$, 0.01 and 0.001,
respectively. Figures were constructed with Prism 5 (GraphPad, San Diego, CA), Python and Adobe Illustrator
software.
173
3. Results
3.1. Concentration-Dependent Bile Effects on CIN Firing Rate and Variance
Bath application of varying BA concentrations had a dose-dependent effect on CIN firing rate and variance.
Traces displaying CIN firing patterns under each 1:1 CA/DCA mixture concentration are shown (**Fig. 1a**).

Cholinergic interneuron firing rate increased by 178 ±24% with exposure to 1 µM CA/DCA and by 171±25% with
exposure to 10 µM CA/DCA compared to BL (**Fig. 1b**; one-way repeated-measures ANOVA, $F_{6,47}=16.93$,
p<0.0001; Dunnett posthoc p<0.05 for BL vs. 1 µM and 10 µM CA/DCA). Conversely, CIN firing rate decreased by
90±8% with exposure to 10 mM CA/DCA (**Fig. 1b**; Dunnett post hoc p<0.05 for BL vs. 10 mM CA/DCA). Finally,
100 µM and 1 mM CA/DCA did not significantly influence CIN firing rate compared to BL (**Fig 1b**; Dunnett post
hoc p>0.05 for BL vs. 100 µM and 1 mM CA/DCA). Cholinergic interneuron firing variance increased by 275±75%
with exposure to 10 mM CA/DCA, but was not significantly influenced by other CA/DCA concentrations (**Fig. 1c**;
one-way ANOVA, $F_{6,47}=4.869$, p=0.0013; Dunnett post hoc p<0.05 for BL vs. 10 mM bile and p>0.05 for BL vs. 1,
10, 100 µM and 1 mM bile).
187
3.2. Bile Effects on CIN Firing Rate with Exposure to TGR5 Antagonist SBI-115
Treating brain slices with a TGR5 BA receptor antagonist (SBI-115; 10 µM) altered concentration-
dependent bile effects on CIN firing rate and variance (**Fig. 2**). Cholinergic firing traces under each experimental
CA/DCA concentration in SBI-115-treated brain slices are shown (**Fig. 2a**). Low CA/DCA concentrations (1 µM
and 10 µM) did not have a significant effect on CIN firing rate in SBI-115-treated brain slices (**Fig. 2b**; one-way
repeated-measures ANOVA, $F_{6,42}=5.418$, p=0.0008; Dunnett post hoc p>0.05 for BL vs. 1 and 10 µM CA/DCA).
Intermediate and high CA/DCA concentrations (100 µM, 1 mM and 10 mM) all significantly decreased CIN firing
rate by 76±13%, 77±13% and 74±25%, respectively, in SBI-115-treated brain slices compared to BL (**Fig. 2b**;
Dunnett post hoc p<0.05 for BL vs. 100 µM, 1 mM and 10 mM CA/DCA). None of the experimental CA/DCA
concentrations significantly altered CIN firing variance in SBI-115-treated brain slices compared to BL (**Fig. 2c**;
one-way ANOVA, $F_{6,47}=0.3353$, p=0.8887; Dunnett post hoc p>0.05 for BL vs. 1, 10, 100 µM and 1, 10 mM
CA/DCA).
Comparison of BA effects in the presence and absence of SBI-115 showed that bath application of SBI-115
decreased CIN firing rate responses to 1 µM and 10 µM CA/DCA concentrations by 60±12%, but did not
significantly alter CIN firing rate responses to 100 µM, 1 mM and 10 mM CA/DCA concentrations (**Fig. 2d**; two-
way repeated-measures ANOVA (bile vs. bile + SBI-115); concentration: $F_{6,48}=14.59$, p<0.0001; interaction:
$F_{6,48}=5.887$, p=0.0006; Dunnett's post hoc, p<0.05 for 1 µM CA/DCA, p<0.001 for 10 µM CA/DCA and p>0.05 for
100 µM, 1 mM and 10 mM CA/DCA). Although SBI-115 did not alter CIN firing variance at individual CA/DCA
concentrations, when its effects across all BA conditions were pooled, SBI-115 significantly increased overall
variance compared to variance during BA conditions sans SBI-115 (**Fig. 2e**; two-tailed unpaired t-test, $t_{78}=2.631$,
p=0.0103). To test for potential independent SBI-115 effects, CIN firing rate and variance in standard ACSF BL
conditions was compared to CIN firing rate and variance in SBI-115-treated ACSF BL conditions and no significant
differences were found for either firing rate or variance (**Fig. 2f;** firing rate, two-tailed unpaired t-test, $t_{14}=1.009$,
p=0.3301; **Fig. 2g;** firing variance, two-tailed unpaired t-test, $t_{12}=0.5325$, p=0.6041).
212
3.3. Bile Effects on CIN Inhibitory Current Rate and Amplitude
To further investigate how BAs influence CIN firing properties, we conducted whole-cell experiments
assessing BA impact on CIN sIPSC rate and amplitude (**Fig. 3**). BAs had a concentration-dependent effect on CIN
sIPSC rate and amplitude. Representative traces displaying CIN sIPSCs across varying CA/DCA concentrations are
shown for BL, 10 µM, 100 µM, and 1 mM CA/DCA conditions (**Fig. 3a**). sIPSC rate was not significantly affected
by 100 µM CA/DCA, a concentration that also had not previously affected firing rate or variance (**Fig. 1b-c**), yet
was significantly decreased with exposure to 10 µM CA/DCA compared to the 1 mM KA BL (**Fig. 3b**; one-way
ANOVA, $F_{2,17}= 5.083$, p= 0.0186, Dunnett post hoc p<0.05 for BL vs. 10 µM CA/DCA and p>0.05 for BL vs. 100
µM CA/DCA), a dose that had increased firing rate (**Fig. 1b**). sIPSC rate was also not significantly affected by 1
mM CA/DCA (**Fig. 3c**; two-tailed unpaired t-test, $t_7=1.612$, p=0.151), a dose that also had not affected firing rate or
variance (**Fig. 1b-c**).
Additionally, BAs exert a dose-dependent effect on sIPSC amplitude. Exposure to 10 µM CA/DCA
increased, whereas 100 µM and 1 mM CA/DCA decreased sIPSC amplitude (**Fig. 3d**; one-way ANOVA, $F_{3, 8233}=$
281.7, p<0.0001, Dunnett post hoc p<0.0001 for BL vs. 10 µM CA/DCA, p<0.0001 for BL vs. 100 µM CA/DCA,
and p<0.0002 for BL vs. 1 mM CA/DCA). As expected, 10 µM CA/DCA increased the frequency of high-amplitude
inhibitory currents (one-way ANOVA, $F_{4,1521}= 3.88$, p=0.0039, Dunnett post hoc p<0.05 for BL vs. 10 µM
CA/DCA). Interestingly, despite decreasing the average sIPSC amplitude, 1 mM CA/DCA increased the frequency
of high-amplitude sIPSCs (Dunnett post hoc p<0.05 for BL vs. 1 mM CA/DCA). In contrast, the 100 µM CA/DCA
concentration did not significantly alter the frequency distribution of sIPSC amplitudes (Dunnett post hoc p>0.05 for
BL vs. 100 µM CA/DCA).
233

3.4. Bile Acid Effects on Inhibitory and Excitatory Inputs on CINs
Next, the effects of BAs on sIPSCs in the presence of the TGR5 antagonist SBI-115 were examined (**Fig.**
4a,b). The low but effective concentration of 10 μM BA concentration continued to suppress sIPSCs (**Fig. 4b**; two-
tailed unpaired t-test, t_5=2.224, p=0.0384), suggesting that TGR5 is not involved in BA regulation of inhibitory
input frequency. Next, the effects of 10 μM BAs on excitatory inputs was examined to see if a BA induced reduction
in sEPSCs might explain BA effects on CIN firing (**Fig. 4c,d**). The 10 μM CA/DCA BA concentration does not
significantly influence sEPCSs rate on CINs and was not explored further (**Fig. 4d**; two-tailed unpaired t-test,
t_4=1.031, p=0.1804). Since sEPCS seem largely unaffected by BAs at this concentration, and since SBI-115 does not
appear to block BA effects on sIPSCs, this suggests that BAs may be having direct effects on CINs, likely through
modulating intrinsic activity, possibly hyperpolarization activated cation (Ih) channels.
244
3.5. Direct and CIN-Mediated Bile Effects on DA Release in the NAc
Considering the well characterized regulation of DA terminals by cholinergic inputs, next voltammetry
was used to study the effects on BAs on DA transmission (**Fig. 5**). Current traces showing DA release in the
presence of CA/DCA (1 μM-10 mM) are shown (**Fig. 5a**). Similar to the effect of BAs on CIN firing, there is a
slight increase in DA release at low BA concentrations (1-10 μM), but robust decreases in DA release at higher
concentrations (>1mM). Furthermore, applying a single concentration (10 μM) to reduce variability introduced
during longer recordings, resulted in consistent and significant ~20% increases in DA release (**Fig. 5b**; one-way
ANOVA, $F_{13,52}$=3.096, p=0.0019, time course in **Fig. 5f**). Average DA release was significantly altered by CA/DCA
(**Fig. 5c**; one-way repeated-measures ANOVA, $F_{6,35}$=10.75, p<0.0001; Dunnett post hoc p>0.05 for BL vs. 1, 10,
100 μM and 1 mM CA/DCA). However, average DA release was decreased by 96±3% with exposure to 10 mM
CA/DCA (**Fig. 5c**; Dunnett post hoc p<0.05 for BL vs. 10 mM) and DA release was not restored during a wash
condition (Dunnett post hoc p<0.05 for BL vs. 10 mM).
Although low bile concentrations (1–10 μM CA/DCA) initially do not significantly affect average DA
release (OLS regression, total effect: c = 0.024, p = 0.863), prolonged exposure to these concentrations
produces transient increases in DA release (**Fig. 5e-f**).To investigate whether 10 mM bile effects on average DA
release were direct or CIN-mediated, we performed mediation analysis and tested the significance of the indirect
CIN-mediated effect using the Sobel test. Mediation analysis indicated that 10 mM CA/DCA-induced inhibition of
DA release is largely mediated by CIN activity suppression, which accounts for 66.9% of bile's effects at this
concentration (indirect effect a*b = -0.1726, Sobel Z = -2.749, p = 0.006; CIN suppression a = -0.4568, p < 0.001).
Direct bile effects on average DA release were nonsignificant (c'= -0.0852, p=0.272). The solid line represents
observed DA release across varying CIN firing levels, while the dashed lines show predicted DA release after
accounting for indirect (CIN-mediated) and total bile effects. (Fig. 5d; total effect c = -0.2578, p < 0.001).
Specifically, exposure to 1 μM CA/DCA increased DA release by 16.4 ± 5.6 % relative to BL at 8 min (**Fig.**
5e; one-sample two-tailed t-test, t = 2.90, p = 0.0337). To determine whether these effects were direct or
CIN-mediated, we performed mediation analysis comparing time-resolved changes in CIN activity and DA release
during exposure to 1 μM CA/DCA. A time-lagged Pearson correlation was used to account for the expected causal
delay of CIN activity on DA release. The analysis revealed that CIN firing at preceding time points strongly
predicted DA release at 8 min, producing a significant mediated effect (a*b = 0.036, Sobel Z = 2.79,
p=0.0053). In contrast, the direct effect of bile on DA release (c' = –0.0140, p = 0.3020) and the total effect
(c = 0.0216, p = 0.1913) were both nonsignificant. Exposure to 10 μM CA/DCA also produced a transient
increase in DA release, specifically at 14 min after drug application (**Fig. 5f**; one-sample two-tailed t-test, t =
2.90, p = 0.0335). Mediation analysis revealed no significant CIN-mediated contribution (a*b = –0.009, Sobel Z = –
1.41, p = 0.158), an almost significant direct bile effect (c' = 0.0249, p = 0.056), and a significant total bile effect (c
= 0.0163, p = 0.0032).
DA release showed no significant transient alterations with exposure to 100 μM CA/DCA (one-sample
two-tailed t-tests, t = 0.88–1.33, p = 0.242–0.421). Likewise, time-lagged mediation analysis revealed no
evidence for CIN-mediated or direct bile effects on DA at this bile concentration (indirect effect a*b = –0.019,
Sobel Z=-1.009, p = 0.313; direct effect c' = 0.021, p = 0.463; total effect c = 0.003, p = 0.716). DA
release also did not exhibit any transient alterations with exposure to 1 mM CA/DCA (one-sample two-tailed t-
tests, t_5 = –0.56–1.00, p = 0.365–0.914).
285
3.6. Bile Effects on NAc DA Release and Clearance Rate
Obeticholic acid (OCA), a synthetic BA, and lithocholic acid (LCA), a secondary BA made only in the gut,
have been shown to bind non-competitively to dopamine transporters (DATs; Romanazzi 2021). For this purpose,
the clearance of DA (tau), as well as the upward and downward velocity of DA signals were analyzed in the

presence of CA/DCA (1-100 µM; **Fig. 6**). Raw example traces are shown for these experiments, normalized to peak
height for ease of comparison where no easily observable changes were seen either with the clearance of DA or the
decay measure tau (**Fig. 6a**). The measure of upward velocity trended towards significant increases (**Fig. 6b**, one-
way ANOVA; $F_{3,15}$=3.094, p=0.0588), which is consistent with observed increases in DA release results in **Fig. 5**. In
contrast, no significant effect was seen between CA/DCA and tau (**Fig. 6c**, one-way ANOVA; $F_{3,15}$=0.3122,
p=0.8163), and downward velocity (**Fig. 6d**, one-way ANOVA; $F_{3,15}$=1.246, p=0.3281). Thus, CA/DCA does not
affect the clearance of extracellular DA, suggesting some specificity in BA effects on this circuit.

4. Discussion

The present work demonstrates that CA/DCA have potent previously unexplored effects on mesolimbic circuitry.
Cholinergic interneurons and DA terminals are sensitive to low BA concentration, robustly increasing CIN firing
frequency that may explain the increases in DA release. High BA concentrations inhibit both CIN firing and DA
release, likely through membrane dissociating effects. The TGR5 appears to be involved in some of the CIN
excitation effects, but not the high concentration BA effects. Reductions in sIPSC frequency at low BA
concentrations are not prevented by TGR5 antagonism, suggesting that another receptor may also contribute to CIN
excitation through GABA inputs. Excitatory inputs onto CINs were explored minimally but do not appear to be
involved in BA effects at the 10 µM concentration. Importantly, DA terminal activity appears to match that of CINs,
suggesting that CIN effects are robust enough to influence downstream circuits. No direct effects of BAs on DA
clearance were observed suggesting that this interaction may be limited to specific secondary BAs.

Concentration-Dependent Bile Effects on CIN Firing Rate and Variance

Estimates of total brain BA concentrations depend on both peripheral BA influx and potential local BA
synthesis within the brain. Under normal physiological conditions, BAs produced in the liver contribute
approximately 0.2–0.4 µM to total brain BA levels, while brain-resident BA-synthesizing enzymes may contribute
an additional 0.05–0.1 µM, resulting in a net estimated brain BA concentration of ~0.25–0.5 µM [17, 18].
Following moderate ethanol intake (1–2 standard drinks), peripheral BA contributions may rise to ~0.5–2 µM and
local BA synthesis—potentially via increased expression of enzymes like CYP46A1—may increase to ~0.2–
0.5 µM, yielding an estimated total brain concentration of ~0.7–2.5 µM [5, 12]. Similarly, HFD-induced increases
in hepatic BA secretion are estimated to elevate peripheral BA contributions to brain concentrations to
approximately 0.3–0.75 µM, with local synthesis potentially adding another 0.1–0.3 µM. This results in total
brain BA levels in the range of ~0.4–1 µM [6, 7]. Presently, we use a 1:1 CA/DCA mixture to assess general BA
effects on striatal activity, regardless of their possible brain-derived or hepatic origin. Our results show that low
concentrations (1–10 µM) of the 1:1 CA/DCA mixture increase CIN firing rate, whereas high concentrations (10
mM) suppress CIN activity. Our BL brain BA estimates (~0.2–0.4 µM) and post-ethanol/HFD brain BA estimates
(~0.4–2.5 µM) indicate that ethanol and fatty food consumption can elevate brain BA levels into a concentration
range (~1 µM) that, in our *in vitro* experiments, is sufficient to increase CIN activity. Heightened CIN activity in the
NAc could, in turn, promote DA release, providing a potential mechanism by which BAs can influence mesolimbic
signaling and reinforce hedonic feeding and ethanol consumption [2]. Our results also demonstrate that exposure to
10 mM CA/DCA significantly decreases CIN firing rate. This inhibitory effect at moderate to high BA
concentrations may reflect genuine suppression of CIN activity or could result from a compromised cell membrane
potentially leading to cell death, especially at the highest concentration [19]. Additionally, we show that BAs exert a
concentration-dependent effect on CIN firing variance. Specifically, exposure to 10 mM CA/DCA significantly
increases CIN firing variance, which may reflect irregular firing patterns—such as sparse and burst firing—that
sometimes precedes neuronal death in patch clamp experiments.

4.2. Bile Effects on CIN Firing Rate with Exposure to TGR5 Antagonist SBI-115

The TGR5 is activated by several BAs, including CA and DCA [20], and is present in neurons and
astrocytes within the CNS [15]. Furthermore, histological data available on the Allen Mouse Brain Atlas confirm
TGR5 expression in the NAc of adult mouse brain slices [21], and behavioral studies in TGR5 knockout mouse
models demonstrate that this receptor is necessary for cocaine reward [16]. In the present study, we investigated
whether TGR5 activation contributes to BA effects on accumbal CIN firing rate. Bath application of TGR5
antagonist SBI-115 abolished the excitatory effects of low (1–10 µM) CA/DCA concentrations on CIN firing but
had no effect on the inhibitory response observed at high (10 mM) CA/DCA concentrations.
Our findings demonstrate that low, physiologically relevant BA concentrations enhance CIN activity
through TGR5 activation, whereas the inhibitory effects observed at high, supraphysiological concentrations are
mediated by TGR5-independent mechanisms. Supporting this, SBI-115 alone did not alter baseline CIN firing,

indicating that it has no independent effect but instead selectively blocks TGR5-mediated excitation at low BA
concentrations. In contrast, the TGR5-independent nature of high BA concentration effects supports the
interpretation that supraphysiological BA levels act through nonspecific mechanisms, such as compromising
membrane integrity, which leads to neuronal dysfunction or cell death and manifests as apparent inhibition of CIN
activity. The lack of firing rate recovery during washout—regardless of SBI-115 presence—further reinforces the
possibility of membrane disruption and irreversible cell damage.
Our results show that although bath application of SBI-115 did not affect BL CIN firing variance, it
selectively increased variance in response to all experimental BA concentrations (see Fig. 2c,e), suggesting that
TGR5 signaling acts as a key stabilizer of CIN firing during elevated BA exposure. Cholinergic interneurons
normally exhibit irregular firing patterns, and previous studies have shown that activation of specific channels, such
as the small-conductance calcium-activated potassium channel (SK) and the hyperpolarization-activated cyclic
nucleotide-gated channel (HCN, Ih), can reduce variability in CIN activity [22, 23]. Activation of the TGR5 receptor
may trigger many of the same downstream mechanisms that stabilize CIN firing following SK and HCN channel
activation. This stabilizing role for TGR5 would explain why bath application of its antagonist, SBI-115, reveals
stochastic fluctuations in CIN firing that likely arise from both intrinsic ion channel activity and extrinsic synaptic
inputs.
362
4.3. Bile Effects on CIN Inhibitory Current Rate and Amplitude
To further investigate the mechanisms underlying the excitatory effects of low BA concentrations (1 and
10 μM CA/DCA) on CIN firing, we examined the impact of 10 μM CA/DCA exposure on sEPSCs. Intracellular
dialysis, which occurs during whole-cell patch-clamp recordings when the pipette solution mixes with the neuron's
cytoplasm, dilutes endogenous signaling molecules [24], which may on its own disrupt some signaling. Therefore,
we selected a high bath concentration of BA (10 μM) during whole-cell recordings to ensure sufficient BA effects
on CINs and account for possible changes in sensitivity. Although 10 μM CA/DCA exceeds the ~2.5 μM peak
brain concentration observed following ethanol or fatty food consumption, it reliably produced measurable effects
while remaining within a physiologically informative range. Our results showed that 10 μM CA/DCA significantly
suppressed sIPSCs, suggesting that low BA concentrations directly modulate GABAergic signaling. Interestingly,
10 μM BAs also increased sIPSC amplitude despite reducing their frequency. This effect may be due to the partial
inhibition of presynaptic GABA release, which would decrease the number of overlapping inhibitory events, reduce
shunting from concurrent inputs, and thereby allow each remaining inhibitory event to generate larger postsynaptic
currents.
Our experiments further demonstrated that 100 μM and 1 mM CA/DCA did not significantly affect
sIPSC frequency, consistent with cell-attached recordings showing that these concentrations do not alter overall CIN
firing. Notably, the effect of 1 mM CA/DCA on inhibitory currents was highly variable and maintaining the whole-
cell seal was considerably more difficult compared to other BA conditions, causing several experiments to end early.
This is suggestive of well-known cell-membrane–disrupting effects of BAs and may also explain the inhibitory
effects of 10 mM CA/DCA in cell-attached firing experiments. Interestingly, although 100 μM and 1 mM
CA/DCA did not change event frequency, both concentrations reduced the average sIPSC amplitude. The presence
of amplitude alterations in the absence of frequency changes indicates that presynaptic release was likely unaffected,
whereas postsynaptic mechanisms—such as receptor desensitization or subtle membrane alterations—may have
contributed to these inhibitory effects.
Analysis of sIPSCs showed that, beyond exerting independent effects on event frequency and amplitude,
CA/DCA concentrations also differentially modulate average current amplitude and the distribution of event
amplitudes. At 10 μM, both the average amplitude and the occurrence of larger events increased, indicating
selective enhancement of specific inputs via presynaptic facilitation and/or postsynaptic sensitization. At 100 μM,
the average amplitude decreased without altering the amplitude size distribution, consistent with uniform
postsynaptic dampening of all current amplitudes. At 1 mM, although the average amplitude decreased, the
distribution shifted toward a higher occurrence of larger events, reflecting complex, non-uniform postsynaptic
modulation—possibly arising from heterogeneous receptor clustering or differential effects on membrane
microdomains that selectively enhance larger currents while reducing smaller ones. Together, these results highlight
the nuanced role of BAs in inhibitory synaptic transmission.

4.4. Bile Effects on CIN Excitatory Current Rate and Inhibitory Current Rate with Exposure to TGR5 Antagonist
SBI-115
In the cell-attached experiments, the TGR5 antagonist SBI-115 abolished the excitatory effects of low BA
concentrations on CIN firing rate. However, when examining inhibitory currents, CA/DCA continued to suppress

inhibitory input even in the presence of SBI-115, indicating a TGR5-independent mechanism. This suggests that
reduced inhibition contributes to, but does not fully account for, the excitatory effects of low BA concentrations—
which likely involve additional signaling pathways that modulate CIN activity. To investigate whether one of these
additional pathways involves modulation of excitatory transmission, we examined the impact of CA/DCA on
excitatory CIN currents. Excitatory events were isolated by addition of 10 μM picrotoxin, a GABA$_A$ and glycine
receptor antagonist. CA/DCA did not significantly alter sEPSC frequency, indicating that BAs preferentially
modulate GABAergic rather than glutamatergic inputs onto CINs, though these experiments are limited in scope
since only one concentration was examined. In summary, while modulation of GABAergic transmission represents
one mechanism by which BAs alter CIN activity, it does not fully explain the excitatory effects of low BA
concentrations. Instead, BA effects on CIN activity likely result from the convergence of TGR5-dependent signaling
on intrinsic channel activity, and additional TGR5-independent mechanisms. These TGR5-independent mechanisms
may involve the activation of other non- g-protein coupled BA receptors, including FXRs, present in multiple brain
regions [25], and S1PR2, which exhibits RNA expression in the NAc [26].

414
4.5. Direct and CIN-Mediated Bile Effects on DA Release in the NAc
Previous studies have shown that accumbal CIN firing promotes spontaneous DA release in the NAc [2].
Presently, we examined whether BA exposure alters DA release in the NAc, potentially via BA effects on CIN firing
rate observed in our previous cell-attached experiments. Low BA concentrations enhanced DA release, whereas
10 mM CA/DCA markedly decreased DA levels. Mediation analysis suggested that this inhibitory effect is
primarily driven by suppression of CIN activity. However, DA release did not recover during the wash condition,
indicating potential neuronal dysfunction from generalized membrane disruption at 10 mM CA/DCA. Such
dysfunction could inflate the apparent indirect effect, making it appear that 10 mM CA/DCA inhibits DA release
via CIN suppression, when it likely suppresses both CIN activity and DA release directly through membrane
disruption.
While low bile concentrations do not significantly affect average DA release across a larger time frame,
both 1 and 10 μM CA/DCA produce transient DA increases at 8 and 14 min after drug application, respectively.
Mediation analysis indicates that the transient increase at 1 μM CA/DCA is primarily driven indirectly through
bile-induced increases in CIN activity. Despite this significant CIN-mediated effect, the total bile effect remained
largely nonsignificant. Interestingly, the direct effect of bile on DA activity at 1 μM, though nonsignificant, is
moderately inhibitory. We thought this effect may be the result of bile-induced changes in DA reuptake [27],
however, our more direct DA voltammetry experiments measuring reuptake did not observe any changes, reducing
this as a possibility. Overall, these results suggest that 1 μM CA/DCA may exert a subtle direct inhibition of DA
release, which then counteracts bile-induced CIN DA stimulation and ultimately restricts DA increases to transient
periods rather than raising average DA release over time.
Although increased DA release at 1 μM CA/DCA appears to be largely CIN-mediated, transient
elevations during exposure to 10 μM CA/DCA showed no significant CIN-mediated effects. The direct bile effect
on DA activity at 10 μM CA/DCA was excitatory and nearly significant for the concentration response curve,
while the total bile effect was excitatory and significant. Performing the experiment again with 10 μM CA/DCA as
the only concentration resulted in less variability and a more readily statistically detectable effect. Future work
examining a role of CINs in this effect will need to directly assess effects of BAs on DA terminals while controlling
for the presence/absence of CINs. The direct bile effects may involve the activation of BA receptors such as TGR5
and FXR [11]. Although the total bile effect is significant, it remains subtle—likely reflecting opposing inhibitory
and excitatory mechanisms—which may explain the transient nature of DA release increases observed with 10 μM
CA/DCA.
Taken together, our findings indicate that bile modulates DA release in the NAc through distinct,
concentration-dependent mechanisms. At extreme supraphysiological concentrations (1–10 mM), bile suppresses
DA release primarily via inhibition of CIN activity, although generalized membrane disruption likely contributes at
this range. At physiological levels (1 μM), bile enhances DA release through CIN-mediated excitation, whereas at
mildly supraphysiological concentrations (10 μM), bile promotes DA release predominantly through direct
excitatory actions. These results suggest that BA concentrations induced in the brain by ethanol and fatty food
consumption (~1 μM) are sufficient to transiently increase DA release in the NAc, supporting the idea that BAs may
contribute to the reinforcing properties of these substances *in vivo*.

453
454 *5. Conclusion*
455 　　　Fatty foods and ethanol significantly increase hepatic BA secretion, which may contribute to their
456 reinforcing effects in the striatum. The present study demonstrates that BAs modulate striatal activity *in vitro* and

highlights the potential for both hepatic and brain-derived BAs—produced via the classic and alternative synthesis pathways—to influence mesolimbic function *in vivo*. Our results reveal a concentration-dependent effect of BAs on accumbal CIN firing rate and indicate that the excitatory effects of low BA concentrations are mediated by at least two independent mechanisms: TGR5 activation and suppression of inhibitory CIN currents. We further provide evidence that BAs may selectively modulate event frequency, current amplitude, and the distribution of current amplitude sizes. Additionally, voltammetry recordings indicate that although low BA concentrations do not alter average DA release over time, exposure to 1 and 10 μM CA/DCA elicits transient increases in DA levels. These DA effects are likely mediated by both CIN-dependent mechanisms and direct BA actions on dopaminergic terminals in the NAc. Collectively, these results offer compelling evidence for BA modulation of CIN activity and striatal DA signaling.

6. Statements and Declarations

Isabella Roque, Shimran Sharma, Phlipp Mews, Summer L. Thompson, and Jordan Yorgason declare that they have no conflict of interest. All institutional and national guidelines for the care and use of laboratory animals were followed. This work was supported by the US National Institutes of Health (NIH) grant AA030577 to JTY. The authors have no relevant financial or non-financial interests to disclose. Material preparation, data collection, data analysis and figure creation were performed by Isabella Roque and Shimran Sharma. All authors contributed to the experimental design. The first draft of the manuscript was written by Isabella Roque and all authors commented on previous versions of the manuscript and figures. All authors read and approved the final manuscript of figures.

References

1. Stouffer MA, Woods CA, Patel JC, Lee CR, Witkovsky P, Bao L, Machold RP, Jones KT, De Vaca SC,
Reith MEA, Carr KD, Rice ME. Insulin enhances striatal dopamine release by activating cholinergic
interneurons and thereby signals reward. *Nat Commun.* 2015;6:8543. doi:10.1038/ncomms9543

2. Yorgason JT, Rose JH, McIntosh JM, Ferris MJ, Jones SR. Greater ethanol inhibition of presynaptic
dopamine release in C57BL/6J than DBA/2J mice: Role of nicotinic acetylcholine receptors. *Neuroscience.*
2016;307:113–124. doi:10.1016/j.neuroscience.2014.10.052

3. Mohebi A, Collins VL, Berke JD. Accumbens cholinergic interneurons dynamically promote dopamine
release and enable motivation. *eLife.* 2023;12:e85011. doi:10.7554/eLife.85011

4. Alsebaaly J, Dugast E, Favot L, Rabbaa Khabbaz L, Solinas M, Thiriet N. Persistent neuroadaptations in
the expression of genes involved in cholesterol homeostasis induced by chronic, voluntary alcohol intake in
rats. *Front Mol Neurosci.* 2018;11:457. doi:10.3389/fnmol.2018.00457

5. Axelson M, Mörk B, Sjövall J. Ethanol has an acute effect on bile acid biosynthesis in man. *FEBS Lett.*
1991;290(1-2):33–36. doi:10.1016/0014-5793(91)80382-D

6. Wang T, Zhang X, Wang Y, Liu W, Wang L, Hao L, Ju M, Xiao R. High cholesterol and 27-
hydroxycholesterol contribute to phosphorylation of tau protein by impairing autophagy causing learning
and memory impairment in C57BL/6J mice. *J Nutr Biochem.* 2022;106:109016.
doi:10.1016/j.jnutbio.2022.109016

7. Yoshitsugu R, Kikuchi K, Iwaya H, Fujii N, Hori S, Lee DG, Ishizuka S. Alteration of bile acid metabolism
by a high-fat diet is associated with plasma transaminase activities and glucose intolerance in rats. *J Nutr*
Sci Vitaminol (Tokyo). 2019;65(1):45–53. doi:10.3177/jnsv.65.45

8. Nilsson L, Sjövall J, Strom S, Bodin K, Nowak G, Einarsson C, Ellis E. Ethanol stimulates bile acid
formation in primary human hepatocytes. *Biochem Biophys Res Commun.* 2007;364(4):743–747.
doi:10.1016/j.bbrc.2007.10.039

9. Monteiro-Cardoso VF, Corlianò M, Singaraja RR. Bile acids: A communication channel in the gut-brain
axis. *J Mol Neurosci.* 2020;70(11):1643–1656. doi:10.1007/s12017-020-08625-z

10. Mertens KL, Kalsbeek A, Soeters MR, Eggink HM. Bile acid signaling pathways from the enterohepatic
circulation to the central nervous system. *Front Neurosci.* 2017;11:617. doi:10.3389/fnins.2017.00617

11. Xing C, Huang X, Wang D, Yu D, Hou S, Cui H, Song L. Roles of bile acids signaling in neuromodulation

under physiological and pathological conditions. *Cell Biosci.* 2023;13(1):101. doi:10.1186/s13578-023-

01053-z

12. Gong X, Zhang Q, Ruan Y, Hu M, Liu Z, Gong L. Chronic alcohol consumption increased bile acid levels

in enterohepatic circulation and reduced efficacy of irinotecan. *Alcohol Alcohol.* 2020;55(3):264–272.

doi:10.1093/alcalc/agaa005

13. Ridlon JM, Kang D, Hylemon PB. Bile salt biotransformations by human intestinal bacteria. *J Lipid Res.*

2006;47(2):241–259. doi:10.1194/jlr.r500013-jlr200

14. Higashi T, Watanabe S, Tomaru K, Yamazaki W, Yoshizawa K, Ogawa S, Nagao H, Minato K, Maekawa

M, Mano N. Unconjugated bile acids in rat brain: Analytical method based on LC/ESI-MS/MS with

chemical derivatization and estimation of their origin by comparison to serum levels. *Steroids.*

2017;125:107–115. doi:10.1016/j.steroids.2017.07.001

15. Keitel V, Görg B, Bidmon HJ, Zemtsova I, Spomer L, Zilles K, Häussinger D. The bile acid receptor TGR5

(Gpbar-1) acts as a neurosteroid receptor in brain. *Glia.* 2010;58(15):1794–1805. doi:10.1002/glia.21049

16. Reddy IA, Smith NK, Erreger K, Ghose D, Saunders C, Foster DJ, Turner B, Poe A, Albaugh VL,

McGuinness O, Hackett TA, Grueter BA, Abumrad NN, Flynn CR, Galli A. Bile diversion, a bariatric

surgery, and bile acid signaling reduce central cocaine reward. *PLoS Biol.* 2018;16(7):e2006682.

doi:10.1371/journal.pbio.2006682

17. Suchyna TM, Markin VS, Sachs F. Biophysics and structure of the patch and the gigaseal. *Biophys J.*

2009;97(3):738–747. doi:10.1016/j.bpj.2009.05.018

18. Ghaffarzadegan T, Zanzer YC, Östman E, Hållenius F, Essén S, Sandahl M, Nyman M. Postprandial

responses of serum bile acids in healthy humans after ingestion of turmeric before medium/high-fat

breakfasts. *Mol Nutr Food Res.* 2019;63(21):e1900672. doi:10.1002/mnfr.201900672

19. Greenwood J, Adu J, Davey AJ, Abbott NJ, Bradbury MWB. The effect of bile salts on the permeability

and ultrastructure of the perfused, energy-depleted, rat blood-brain barrier. *J Cereb Blood Flow Metab.*

1991;11(5):641–649. doi:10.1038/jcbfm.1991.116

20. Pols TWH, Noriega LG, Nomura M, Auwerx J, Schoonjans K. The bile acid membrane receptor TGR5: A

valuable metabolic target. *Biochem Soc Trans.* 2011;39(4):986–990. doi:10.1042/bst0390986

21. Allen Institute for Brain Science. G protein-coupled bile acid receptor 1 (Gpbar1) expression in the mouse

brain. *Allen Mouse Brain Atlas.* Accessed October 28, 2025. https://mouse.brain-map.org/gene/show/86541

22. Bennett BD, Callaway JC, Wilson CJ, College CN. Intrinsic membrane properties underlying spontaneous

tonic firing in neostriatal cholinergic interneurons. *J Neurosci.* 2000;20(22):8493–8503.

doi:10.1523/JNEUROSCI.20-22-08493.2000

23. Wilson CJ. The mechanism of intrinsic amplification of hyperpolarizations and spontaneous bursting in

striatal cholinergic interneurons. *Neuron.* 2005;45(2):199–208. doi:10.1016/j.neuron.2004.12.053

24. Aiba I, West AK, Sheline CT, Shuttleworth CW. Intracellular dialysis disrupts Zn^{2+} dynamics and enables

selective detection of Zn^{2+} influx in brain slice preparations. *J Neurochem.* 2013;125(6):979–989.

doi:10.1111/jnc.12246

25. Huang C, Wang J, Hu W, Wang C, Lu X, Tong L, Wu F, Zhang W. Identification of functional farnesoid X

receptors in brain neurons. *FEBS Lett.* 2016;590(18):3233–3242. doi:10.1002/1873-3468.12373

26. Human Protein Atlas. S1PR2 expression in the nucleus accumbens. *Human Protein Atlas.* Accessed

October 3, 2025. https://www.proteinatlas.org/ENSG00000267534-S1PR2/brain/nucleus%2Baccumbens

27. Romanazzi T, Zanella D, Bhatt M, Di Iacovo A, Galli A, Bossi E. Bile acid interactions with

neurotransmitter transporters. *Front Cell Neurosci.* 2023;17:1161930. doi:10.3389/fncel.2023.1161930

Fig. 1 BAs have a concentration-dependent effect on CIN firing rate and variance. (a) Sample CIN firing traces during BL and 1 μM-10 mM CA/DCA conditions. **(b)** 1 μM and 10 μM CA/DCA significantly increased, 10 mM CA/DCA significantly decreased, and 100 μM and 1 mM CA/DCA did not significantly alter CIN firing rate compared to BL. **(c)** Only 10 mM CA/DCA significantly altered CIN firing variance, increasing variance significantly above BL. Statistical significance represented as * ($p < 0.05$), ** ($p < 0.01$).

Fig. 2 Treating brain slices with TGR5 antagonist SBI-115 alters bile effects on CIN firing rate and variance. (a) Sample CIN firing traces during BL and 1 μM-10 mM CA/DCA conditions in SBI-115-treated brain slices. **(b)** In SBI-115-treated brain slices, 100μM, 1mM and 10mM CA/DCA significantly decreased, and 1 and 10μM CA/DCA did not significantly alter CIN firing rate compared to BL. **(c)** None of the experimental CA/DCA concentrations significantly altered CIN firing variance in SBI-115-treated brain slices. **(d)** Bath application of SBI-115 significantly altered 1μM and 10μM CA/DCA effects on CIN firing rate, but not 100μM, 1mM and 10mM CA/DCA effects on CIN firing rate. **(e)** Addition of SBI-115 to varying CA/DCA concentration conditions significantly increased overall CIN firing variance. **(f)** CIN firing rate in standard ACSF BL conditions was not significantly different from CIN firing rate in SBI-115-treated BL ACSF conditions. **(g)** CIN firing variance was also not significantly different between standard ACSF BL conditions and SBI-115-treated BL conditions. Statistical significance represented as * ($p < 0.05$), ** ($p < 0.01$), *** ($p < 0.001$).

Fig. 3 BAs have a concentration-dependent effect on CIN inhibitory current rate and amplitude. (a) Sample CIN firing traces with 1s expanded views of whole-cell currents during a 1mM KA BL condition, and with the addition of 10 μM-1 mM CA/DCA to BL conditions. **(b)** 10 μM CA/DCA significantly decreased and 100 μM CA/DCA did not significantly influence CIN inhibitory current rate. **(c)** 1 mM CA/DCA did not have a significant or consistent effect on CIN inhibitory current rate. **(d)** 10 μM CA/DCA significantly increases, and 100 μM and 1mM CA/DCA significantly decrease average CIN inhibitory current amplitude. **(e)** 10 μM and 1mM CA/DCA significantly increase the frequency distribution of high-amplitude inhibitory CIN currents. Statistical significance represented as * ($p < 0.05$), ** ($p < 0.01$), *** ($p < 0.001$), **** ($p < 0.0001$).

Fig. 4 Bath application of 10 μM CA/DCA continues to suppress CIN inhibitory current rate during exposure to SBI-115 and does not significantly influence excitatory CIN current rate. (a) Sample CIN firing traces with enlarged 1s views of whole-cell currents at a 1mM KA BL condition and during exposure to 10 μM CA/DCA and 10 μM SBI-115 in addition to BL conditions. **(b)** During exposure to SBI-115, 10 μM CA/DCA continues to suppress CIN inhibitory current rate. **(c)** Representative CIN firing traces with 1s expanded views of whole-cell currents recorded at a 100 μM picrotoxin BL condition and during exposure to 10 μM CA/DCA in addition to BL conditions. **(d)** 10 μM CA/DCA does not have a significant effect on CIN excitatory current rate. Statistical significance represented as * ($p < 0.05$).

Fig. 5 Bile modulates DA release both directly and indirectly by enhancing CIN activity at physiological concentrations (1 μM CA/DCA), suppressing CIN firing at extreme supraphysiological levels (1-10mM CA/DCA), and directly influencing DA release at minimally supraphysiological concentrations (10 μM CA/DCA). (a) Sample current traces of DA release during BL, and 1 μM-10 mM CA/DCA conditions. **(b)** Representative current trace at BL and after 14 min of exposure to 10 μM CA/DCA. **(c)** 10 mM CA/DCA significantly decreased DA release, but no other experimental CA/DCA concentration conditions significantly altered average DA release. DA release was not restored during the wash condition. **(d)** Bile's effects on average DA release are largely indirect and mediated by changes in CIN activity. Dashed lines indicate predicted DA release after removing indirect (I) and both direct and indirect (D/I) bile effects. **(e)** DA release exhibits a transient increase at 8 min following bath application of 1 μM CA/DCA. **(f)** Following 10 μM CA/DCA application, DA release transiently rises at 14 min. Dashed line extending from the solid CIN firing trace represents the linear regression projection of firing beyond 10 min. Statistical significance represented as * ($p < 0.05$), ** ($p < 0.01$), *** ($p < 0.001$).

Fig. 6 CA/DCA does not affect DA clearance (tau). (a) Overlay of sample traces for DA signals at varying concentrations of CA/DCA (1-100μM) with alignment at peak heights. **(b)** Dopamine clearance (tau) from the synaptic cleft across different concentrations of bile. No significant reuptake at any concentration. **(c)** Upward velocity, a measure of DA release that helps separate it from the clearance value, showed no significance. **(d)** Downward velocity, showing no significance for the clearance of DA from the cleft via DA transporters (DATs).

A

BL   CA/DCA (1 µM)   CA/DCA (10 µM)

CA/DCA (100 µM)   CA/DCA (1 mM)   CA/DCA (10 mM)

60 sec

B

Firing Rate (Norm)

**   *

BL

[CA/DCA] (Log M)

C

CIN Firing Variance

BL

[CA/DCA] (Log M)

A

BL KA (1mM)

55 pA

KA + 10 µM CA/DCA

150 pA

KA + 100µM CA/DCA

80 pA

KA + 1mM CA/DCA

90 pA

60s   1s

B

*

Current Rate (Norm)

BL KA

-5   -4
[CA/DCA] (Log M)

C

Current Rate (Norm)

BL KA

-3
[CA/DCA] (Log M)

D

****   ****   ***

Amplitude (Norm)

BL KA

-5   -4   -3
[CA/DCA] (Log M)