Expression of *Calca* gene-derived peptides in the murine taste system

Salin Raj Palayyan[1*], Abdul Hamid Siddiqui[1*], Sunil Kumar Sukumaran[1,2]

[1]Department of Nutrition and Health Sciences

University of Nebraska-Lincoln

Lincon, NE, 68583

*Equal contribution

[2]Corresponding author: ssukumaran2@unl.edu

Abstract

The Calcitonin Related Polypeptide Alpha (*Calca*) gene is a source of four biologically active

peptides with varied physiological roles. Alternative splicing of the *Calca* messenger RNA

generates either prepro calcitonin gene related peptide (CGRP) or preprocalcitonin encoding

transcripts. Proteolytic processing of preprocalcitonin generates procalcitonin, calcitonin and

katacalcin. Calcitonin is a ligand for the G-protein coupled receptor calcitonin receptor (CALCR)

while CGRP is a ligand for the CGRP receptor (CGRP1R) formed by the calcitonin receptor like

receptor (CALCRL)receptor activity modifying protein 1 (RAMP1) complex. Interestingly,

procalcitonin too, is a ligand for the CGRP1R where it can antagonize CGRP. CGRP expression

in taste neurons has been documented and is posited to regulate taste signaling. Single cell and

bulk RNASeq of taste papillae revealed that the preprocalcitonin but not the CGRP transcript is

expressed in *Tas1r3*- expressing type II taste cells, while *Calcrl* (but not *Calcr*) and *Ramp1* are

expressed in stem/progenitor and type I cells in the circumvallate papillae. The CGRP1R is also

expressed by fibroblasts in the lingual mesenchyme. We confirmed this expression pattern

using quantitative polymerase chain reaction (qPCR), RNAScope and immunohistochemistry.

qPCR of geniculate and nodose-petrosal ganglia revealed that both express *Cgrp* and CGRP1R

subunit mRNAs*,* but not procalcitonin and *Calcr*. This interesting expression patterns suggests

that procalcitonin and CGRP might reciprocally regulate the CGRP1R in taste cells and lingual

fibroblasts and thereby influence taste signaling, taste cell regeneration and the taste

microbiome.

Keywords

Neuropeptide, Procalcitonin, CGRP, Gustation

Introduction

The *Calca* gene is a source of up to four peptide signaling molecules. Tissue specific alternative

splicing of *Calca* transcript generates mRNAs coding for either calcitonin gene related peptide

(preproCGRP, proteolytically processed to alpha CGRP) or preprocalcitonin (prePCT).[1,2]

PrePCT is proteolytically processed to generate procalcitonin (PCT), which is further processed

to generate calcitonin and katacalcin.[3] All four belong to a larger group of peptides that include

beta CGRP, amylin, adrenomedullin and intermedin, which are ligands for receptors formed by

the GPCRs calcitonin receptor (CALCR) or calcitonin receptor like receptor (CALCRL).[4] CALCR

and CALCRL combine with one of three receptor activity modifying proteins (RAMP1, RAMP2 or

RAMP3) to form receptor complexes specific to a subset of these peptides. For example,

calcitonin is a ligand for CALCR (without an associated RAMP subunit), while CGRP is a ligand

for both CALCR+ RAMP1 and CALCRL+RAMP1 (this being its primary receptor, henceforth

designated CGRP1R).[4,5] CGRP is an exceptionally well-studied neuropeptide; it is expressed in

virtually all peripheral sensory neurons, many central neurons, and some non-neuronal cells.[6-8]

It serves a wide variety of functions including nociception, neurogenic inflammation, immunity,

vasodilation, wound healing, regulation of the microbiome etc.[6-8] It is a potent neuro-immune

modulator and was recently shown to modulate antigen sampling by M cells in the Peyer's

patch.[9] It displays antimicrobial activity against several gram-negative and gram-positive

bacteria and Candida albicans, which adds a further dimension to its immunomodulatory and

wound healing roles.[10,11] From a clinical standpoint, its prominent role in triggering migraine has

generated significant scientific interest.[6] This has led to the development of small molecule- and

monoclonal antibody- based CGRP1R antagonists to treat migraines.[12,13] Interestingly, clinical

studies have suggested that one-quarter of migraine patients have altered taste perception.[14]

Indeed, CGRP was shown to shape taste transduction by inducing 5-HT secretion by type III

taste cells in a phospholipase C-dependent manner.[15]

Calcitonin and katacalcin are key regulators of calcium and phosphate levels in blood and

bone.[3,16,17] PCT is a much less studied neuropeptide. It is not widely secreted in the steady

state; it may regulate bone density by suppressing osteoclast (macrophage) migration and

maturation.[18] Interestingly, sepsis-associated cytokine storm is preceded by ubiquitous

upregulation of PCT expression, leading to its adoption as an early sepsis marker.[19-22] PCT is a

partial agonist of CGRP1R and exerts its mediator role in sepsis through this receptor. It also

partially antagonizes CGRP at this receptor.[23-25] However, its biological role(s) in both the steady

state and in sepsis remains enigmatic. Single cell RNASeq (scRNASeq) and bulk RNASeq of

the circumvallate papillae (CVP) done in our lab showed that the prePCT encoding *Calca*-

transcript is highly expressed in type II taste cells, while subunits of the CGRP1R are expressed

in taste stem cells, type I taste cells and lingual mesenchymal fibroblasts. The CGRP transcript

and *Calcr* were not expressed in taste tissues. This was confirmed using qPCR, RNAScope and

immunohistochemistry. On the other hand, the nodose- petrosal and geniculate ganglia that

innervate taste buds were shown to express the CGRP- transcript and CGRP1R subunits, but

not the prePCT transcript and *Calcr*. This intriguing expression pattern suggests that taste cell

derived PCT and taste nerve derived CGRP may reciprocally modulate CGRP1R in the taste

papillae.

Methods

Animals.

8-10 weeks old C57BL/6J mice (The Jackson Laboratory, Bar Harbor, ME) were used for this study. Animals were housed in a specific pathogen free vivarium with a 12-h light/dark cycle and open access to food and water. All animal experiments were performed in accordance with the National Institutes of Health guidelines for the care and use of animals in research and reviewed and approved by the Institutional Animal Care and Use Committee at University of Nebraska-Lincoln (protocols: 2610 and 2366).

Bulk RNASeq of taste papillae.

Pooled taste buds from CVP and fungiform papillae (FFP) of C57BL/6 mice (n=3 each) were excised from CVP sections using laser micro dissection, and full length bulk RNASeq libraries were prepared using the Ovation RNASeq system V2 (Tecan biosystems, Morgan Hill, CA) per manufacturer instructions. Indexed illumina sequencing libraries were prepared and sequenced in a Hiseq 2000 sequencer (Illumina Inc, San Diego, CA). Raw sequences were aligned to the mouse reference genome (version GRCm38.p3) using the STAR program with default settings and Gencode M24.gtf as the splice junction annotation file, and the reads mapping to genes were counted using the featureCounts package.[26,27] Sashimi plots to show alignment of reads were generated using the integrative genome viewer.[28]

scRNASeq of taste papillae.

Chromium™ Single Cell 3′ Solution (10x Genomics Inc, Pleasanton, CA, Cat. no. PN-1000268) was used for scRNASeq analysis.[29] Single cell preparation from CVP was done as previously described.[30,31] A protease cocktail was injected under the lingual epithelium of excised tongue (n=16 mice) and incubated at 37°C for ten minutes. The epithelia were peeled, the CVP were excised and minced to form single cells. Single cell capture, library preparation, sequencing, and primary analyses of sequencing data were done using 10X genomics protocols, and secondary analysis was done using the Seurat package in R.[32]

RNAScope Hiplex assay.

RNAscope assay was done using the Hiplex fluorescent assay kit for mice (Advanced Cell Diagnostics, Hayward, CA, Cat. no. 324443) with indicated probes (Table S1) as previously described using the manufacturer's instructions.[33] Positive and negative control probes were run in parallel to test probes to ensure proper hybridization and imaging conditions were attained in

our experiments. Confocal images were captured using a Nikon A1R-Ti2 confocal laser

scanning microscope using NIS-Elements A1R software image acquisition and analysis

software, using 40/60X objectives. Images were taken using a sequential channel series setting

to minimize cross-channel signal, and the channels used were: GFP 488, TxRed 550, Cy5

650 nm. Z-series stack with 10 images per stack was captured at a step size of 1 μm.

Acquisition parameters [i.e., gain, offset, photomultiplier tube (PMT) setting] were held constant

for experiments. Colocalization counts were made using QuPath software.[34] Cell boundaries

were detected automatically based on DAPI staining. The fidelity of each cell boundary was

confirmed by manual inspection. The number of fluorescent spots in each channel (that

corresponds to individual mRNA molecules) per cell were extracted and used for colocalization

counting. Data from more than two non-consecutive sections from two mice were pooled.

Immunohistochemistry

Standard immunohistochemical techniques were used as previously described.[33,35] Briefly,

frozen sections were rehydrated with PBS. Nonspecific binding was blocked with SuperBlock

Blocking Buffer (Thermo scientific, Waltham, MA, Cat. no. 37580) at room temperature for 1 h.

Sections were incubated with primary antibodies overnight at 4 °C in a humidified chamber.

After three 15-min washes with PBST, slides were incubated for 1 h at room temperature with

one of the following fluorescent secondary antibodies in blocking buffer. All double-

immunofluorescent labeling was done with combinations of the secondary antibodies along with

DAPI (1:1,000; Invitrogen™, Thermo scientific, Waltham, MA, Cat. no. D1306) to label cell nuclei

for cell counting. The primary and secondary antibodies and their concentrations used in this

study are listed in Table S2.

Double labeling with two antibodies made from the rabbit (LRMP+PCT and T1R3+PCT), was

done as described before.[36] After blocking, the sections were incubated in succession: first

primary anti-rabbit antibody overnight at 4 °C, first fluorescence Fab fragment secondary for 1 h

at room temperature, unlabeled anti-rabbit antisera [AffiniPure Fab fragment donkey anti-rabbit

IgG (H+L)] for 3 h at room temperature to ensure that all binding sites in the first primary

antibody are occupied, second primary anti-rabbit antibody overnight at 4 °C, and then second

fluorescence Fab fragment secondary for 1 h at room temperature. Controls for the double

immunohistochemistry experiments with two rabbit primary antibodies were done without

donkey Fab fragment incubation following the first secondary antibody and omitting the second

primary antibody (Figure S6 I & II, control A) or with donkey Fab fragment incubation and

omitting the second primary antibody (Figure S6 I & II, Control B) to show adequate blocking of

rabbit IgG by donkey Fab fragment. Confocal imaging was done as described above for
RNAScope. Colocalization for each taste cell marker was done using images from at least 2
sections showing entire CVP and FOP (n=1 mouse). Only those taste cells for which the entire
cell body and nucleus could be visualized were counted.
175
Western Blot
CVP tissue from three animals was isolated and homogenized in RIPA lysis and extraction
buffer (Thermo scientific, Waltham, MA, Cat. no. 89900). Protein quantification was done using
BCA method (Thermo scientific, Waltham, MA, Cat. no. 23227). Lysate were then mixed with 4
X laemmli buffer (Bio-Rad Laboratories Inc, Hercules, CA, Cat. no.1610747), and 30 μg of
protein run on SDS-PAGE, followed by transfer onto nitrocellulose membrane. Blots were
blocked in Blocker (Thermo scientific, Waltham, MA, Cat no. 37520) and incubated with PCT
primary antibody (Table S2) at 4 °C overnight. Incubation with secondary antibody was done for
1 h and imaged on Odyssey F Imager (LI-COR Inc. Lincoln, NE).
185
PCR and qPCR.
Total RNA was isolated from freshly dissected geniculate and nodose-petrosal ganglia, taste
papillae or NT epithelium using the Quick-RNA Microprep kit (Zymo Research Corp, Irvine, CA,
Cat. no. R1050) with on-column DNA digestion. cDNAs were prepared from total RNAs using
SuperScript IV VILO Master Mix kit (Thermo Fisher, Waltham, MA, Cat. no. 11756050). End
point PCR and qPCR were done as previously described. Exon-exon junction spanning primers
were designed whenever possible to avoid amplification from any contaminating genomic DNA.
A minimum of three biological replicates were used for all cDNA samples. The ratio of the log10
of the average δ-cycle threshold (Ct) value (difference between Ct values of *Bact* and each
gene of interest was plotted. Primers used are shown in Table S3.
196
197   **Results**
198
Bulk RNASeq and PCR results to show the expression of Calca transcripts and receptors
in taste papillae and ganglia.
Analysis of bulk RNASeq data from CVP taste buds isolated by laser microdissection showed
that the *Calca* transcript is strongly expressed in CVP. Alignment of the RNASeq reads to the
Calca genomic locus showed that no reads mapped to the preproCGRP specific exon 5, while
large number of reads aligned to prePCT specific exon 4 (Figure S1). This observation was

confirmed using end point PCR using primers for the *Cgrp* and prePCT transcripts (Figure S2A),

which showed amplification for the prePCT transcript in CVP and foliate papillae (FOP), but not

the FFP. *Cgrp* transcript is expressed at very low levels or were undetectable in all three taste

papillae. We could readily detect transcripts for *Calcrl* and *Ramp1* in all three taste papillae and

for *Ramp2* in CVP, but not for *Calcr* and *Ramp3*. The transcript for *Calcb* that encodes beta

CGRP is expressed in all three taste papillae. None of the transcripts were expressed in non-

taste lingual epithelium (NT, Figure S2A). Next, we checked the expression of these transcripts

in the geniculate and nodose- petrosal ganglia that innervate the FFP and CVP and FOP

respectively. The prePCT transcript and *Calcr* are not expressed in either of them, while the

Cgrp transcript, *Calcrl, Calcb*, *Ramp1*, and *Ramp2* were detected in both (Figure S2B).

Next, we quantified the expression of these transcripts in all four tissues using qPCR.

Consistent with the end point PCR results, robust expression of prPCT is detected in both

papillae and weak expression of *Cgrp* is detected in the CVP but not FOP (Figure 1A).

Conversely, *Cgrp*, but not prPCT transcript is strongly expressed in both taste ganglia (Figure

1B). Robust expression of *Calcrl, Calcb and Ramp1* were detected in both the taste papillae and

ganglia, while *Calcr and Ramp3* are not expressed in any sample. Weak *Ramp2* expression is

detected in the two ganglia, while it is not expressed in the taste papillae (Figure 1).

scRNASeq of CVP identifies Calca and CGRP1R subunit expressing taste and

mesenchymal cells. To identify taste cells that express *Calca* and CGRP1R subunits, we

turned to scRNAseq data from CVP. We found that *Calca* is expressed only in mature and

immature sweet taste cells. *Calcrl* is expressed in subtypes of type I cells and taste stem cells.

Ramp1 expression was low overall, while the regulatory subunit of the CGRP1R, *Crcp* is

expressed widely across all cell types (Figure S3).

Western blot and histological analyses confirm the expression of Calca and Calcrl in

taste cells. Using western blot analysis using an antibody specific to PCT, we detected its

expression in CVP (Figure S4). Next, we turned to histological analysis to confirm the cell type

specific patterns of *Calca* and *Calcrl*. RNAscope Hiplex analysis of CVP confirmed the

scRNASeq results: *Calca* is strongly coexpressed with the sweet taste receptor subunit *Tas1r3*,

with 148/164 (90%) of *Tas1r3* expressing cells coexpressing *Calca,* and 148/149 (99%) *Calca*

expressing cells coexpressing *Tas1r3* (Figure 2A- A″)*. Calca* expressing cells also coexpressed

the pan type II taste cell marker *Trpm5,* with 131/437 (30%) of *Trpm5* expressing cell

coexpressing *Calca*, while 131/149 (88%) of *Calca* expressing cells coexpress *Trpm5* (Figure 2

B- B″). Most type II cells that do not express *Calca* appear to be bitter taste receptor cells, as

less than one tenth of *Gnat3* (primarily bitter taste cell marker in CVP) expressing cells

expressed *Calca* (Figure 2C- C″). *Calca* is not expressed in type III taste cells marked by *Ddc*.

(Figure 2D- D″). *Calcrl* staining is observed in basal (presumably stem/progenitor) cells in the

CVP taste buds (Figure 2E, E′). Significant staining for *Calcrl* is also observed in mesenchymal

cells close to basal cells of taste buds that stained strongly for the fibroblast marker *Sparc*

(Figure 2E, E′). Comparable results were obtained using double labeled immunohistochemistry.

Using a PCT specific antibody, we saw strong co expression of PCT with the sweet taste

receptor subunit T1R3 (100% in both directions) in CVP (Figure 3 A-E) and FOP (Figure 3 F-J).

In addition, PCT is strongly coexpressed with the pan-type II marker LRMP, with all PCT

expressing cells expressing LRMP and 33/45 (73%) of LRMP expressing cells coexpressing

PCT in the CVP (Figure 3 K-O). In case of LRMP, colocalization is observed primarily with cells

that stained strongly with LRMP antibody. The weaker LRMP positive cells primarily expressed

GNAT3, which, as alluded to before, is primarily expressed in bitter taste receptor cells in the

CVP (Figure S5).

Discussion

The *Calca* derived peptides CGRP, calcitonin and procalcitonin play key roles in health and

disease.[3,6-8,19,21,37,38] Using a slew of techniques probing the expression of the corresponding

mRNAs and proteins, we show that the prePCT encoding transcript is strongly expressed in

CVP and FOP, almost exclusively in the *Tas1r3* expressing type II (primarily sweet and umami

receptor expressing) taste cells. Curiously, *Calca* expression (both prePCT and *Cgrp* mRNAs) is

not observed in FFP (Figure S2). The related *Calcb* gene is expressed in all three taste papillae.

All three taste papillae express the CGRP1R subunits *Calcrl* and *Ramp1* (Figure S2, Figure 1).

Interestingly, the taste ganglia that innervate the papillae expresses the *Cgrp* but not the

prePCT transcript and expresses both the CGRP1R subunits (Figure S2, Figure 1). A small

amount of *Cgrp* transcript is detected in the CVP using endpoint PCR and qPCR, although it is

likely derived from the nerve endings that innervate the taste papillae rather than taste cells

themselves (Figure S2, Figure 1). This is supported by the absence of *Cgrp* transcript in the

bulk RNASeq data from CVP, derived from laser microdissected taste buds devoid of

contamination from nerve bundles in the CVP core that would be present in the CVP samples

used for PCR experiments (Figure S1). CGRP expression in trigeminal neurons is well studied,

and it is very likely that trigeminal neurons that innervate the taste papillae also express CGRP,

although this has not been experimentally demonstrated to our knowledge.[7] Thus, it is possible

that the bulk of alpha CGRP in the taste papillae is nerve derived. This is also supported by

immunolocalization studies of CGRP in rodent, pig and human taste papillae, all of which show

CGRP expression restricted to nerve bundles and their termini in and around taste buds.[15,39-41]

PrePCT is processed to generate PCT and then to calcitonin and katacalcin. We were able to

detect the robust expression of prePCT transcript using end point and qPCR and could detect

PCT protein expression using western blot (Figures S2, S4 and 1). Although the RNAScope

probe for *Calca* does not distinguish between the two alternatively spliced mRNAs, we obtained

strong staining in TAS1R3- and LRMP- expressing taste cells with the PCT antibody (Figure 2,

Figure 3). However, these experiments do not allow us to determine if calcitonin and katacalcin

are produced by taste cells. Since both peptides are part of PCT, they cannot be readily

distinguished using antibody staining. Notably, their biological roles are distinct from that of PCT;

both are well known hormones produced by the parathyroid gland that bind to the calcitonin

receptor to regulate bone and serum calcium and phosphorus levels.[16,17] PCT on the other

hand, is an early marker of sepsis, which is only expressed at very low levels in healthy

individuals.[19,24] Notably, we did not detect the calcitonin receptor mRNA *Calcr,* in the taste

papillae or the geniculate and nodose-petrosal ganglia (Figure S2, Figure 1). Thus, the available

evidence indicates that calcitonin and katacalcin are either not produced by taste cells and

nerves or will not be biologically effective in case they are produced. PCT on the other hand,

can stimulate CGRP1R, and it also antagonizes CGRP at this receptor.[23-25] Considering

abundant CGRP1R receptor expression in taste papillae and ganglia, PCT is the only peptide

generated from taste cell expressed prePCT transcript capable of exerting its biological role.

What are the likely roles of PCT and CGRP in the taste system? The amount of peptides

produced by the taste cells will not be sufficient to elevate their circulating levels, and their

effects will be mediated by paracrine signaling within the taste papillae. We know a lot about the

biological roles of CGRP. One study that looked at the effects of CGRP in isolated taste buds

using calcium imaging and bioassay showed that CGRP may regulate taste signaling by

regulating 5-HT signaling by type III cells.[15] We (and the original study) did not detect CGRP1R

expression in type III cells. We detected *Crcp* expression in many taste cell types using

scRNASeq, which agrees with their findings (Figure S2). Using scRNASeq, we detected

CGRP1R expression in type I cells (Figure S2). Type I cells can regulate taste signaling, which

is an alternative explanation for CGRP's ability to regulate taste signaling.[42] We found that

Calcrl is expressed in taste stem cells using scRNASeq and RNAScope (Figure S3, Figure 2).

However, *Ramp1 was not detected by scRNASeq* which is likely a false negative due to the

limitations of scRNASeq, as we could detect it using qPCR. CGRP regulates stem cell

maintenance, and CGRP1R expression in basal cells in the taste buds and adjoining fibroblasts

raises the possibility that taste nerve derived CGRP regulates taste stem cells directly, and

indirectly by regulating adjoining fibroblasts (Figure 2).[43,44] CGRP can also regulate immunity at

the taste cells through its effects on immune cells, epithelial cells and fibroblasts, and can shape

the oral microbiome directly through its microbicidal effects.[8,11,37,45-47] It is also capable of

regulating microfold cells that mediate microbial transcytosis in the Peyer's patch.[9] We have

shown that type II taste cells might mediate immune surveillance similar to microfold cells, and it

is possible that CGRP plays a similar role in taste papillae.[33] Sepsis-associated cytokine storm

is preceded by ubiquitous PCT expression, leading to its adoption as an early sepsis marker.

However, very little is known about its role in normal physiology. In the skeletal system, it may

regulate bone density by suppressing osteoclast (macrophage) migration and maturation.[18] It's

expression in CVP and FOP but not FFP might provide some clues to its function in taste cells.

Unlike the FFP, the CVP and FOP have deep trenches around which the taste buds are

arranged. The trenches are relatively less exposed to salivary flux and may be more hospitable

to the oral microbiota, including those responsible for halitosis, and plausibly, pathogenic as

well. This raises the possibility that PCT might regulate the taste papillae microbiome in much

the same manner as CGRP. In addition, it has the potential to regulate taste signaling and taste

cell regeneration by regulating type I and stem/progenitor cell expressed CGRP1R. As stated

before, PCT is a partial agonist of the CGRP1R, and it can partially antagonize the effects of

CGRP at this receptor. Thus, it is plausible that PCT and CGRP reciprocally shape the biological

effects CGRP1R signaling in the taste papillae. Thus, the taste papillae might be a suitable

model system to determine the biological roles of PCT and its cross talk with CGRP.

Acknowledgements

We would like to thank Brian Lewandoski for helping with taste cell isolation for scRNASeq, and

Ichiro Matsumoto for the T1R3 antibody. The Microscopy work was carried out at the

Microscopy Research Core Facility of the Center for Biotechnology at UNL, which is partially

funded by the Nebraska Center for Integrated Biomolecular Communication COBRE grant (P20

GM113126 and NIGMS) and the Nebraska Research Initiative.

Funding

This work was supported by National Science Foundation CAREER award # 2443659,

NIH/NIDCR New Investigator RO3 award #1R03DE032417, a Project leader award from

Nebraska Center for the Prevention of Obesity Diseases (NIH-NIGMS grant no.

P20GM104320), and PA State tobacco grant STA019A01SUKUM to SKS.

Conflict of Interest

344   The authors declare no conflict of interest.

Author Contributions

SRP and AHS: performing experiments, acquiring the data, data analysis, interpretation of the

results, creation of the figures, and revision of the manuscript. SKS: conceptualization and

design of the study, supervision of the data acquisition, interpretation of results, writing of the

manuscript.

Data availability.

The underlying bulk and scRNASeq data are being uploaded to NCBI's short read archive and

will be available very soon.

References

1   Lou, H. & Gagel, R. F. Alternative RNA processing--its role in regulating expression of
calcitonin/calcitonin gene-related peptide. *J Endocrinol* **156**, 401-405,
doi:10.1677/joe.0.1560401 (1998).
2   Lou, H. & Gagel, R. F. Mechanism of tissue-specific alternative RNA processing of the
calcitonin CGRP gene. *Front Horm Res* **25**, 18-33, doi:10.1159/000061000 (1999).
3   Naot, D., Musson, D. S. & Cornish, J. The Activity of Peptides of the Calcitonin Family in
Bone. *Physiol Rev* **99**, 781-805, doi:10.1152/physrev.00066.2017 (2019).
4   Hay, D. L., Garelja, M. L., Poyner, D. R. & Walker, C. S. Update on the pharmacology of
calcitonin/CGRP family of peptides: IUPHAR Review 25. *Br J Pharmacol* **175**, 3-17,
doi:10.1111/bph.14075 (2018).
5   Pioszak, A. A. & Hay, D. L. RAMPs as allosteric modulators of the calcitonin and
calcitonin-like class B G protein-coupled receptors. *Adv Pharmacol* **88**, 115-141,
doi:10.1016/bs.apha.2020.01.001 (2020).
6   Russo, A. F. & Hay, D. L. CGRP physiology, pharmacology, and therapeutic targets:
migraine and beyond. *Physiol Rev* **103**, 1565-1644, doi:10.1152/physrev.00059.2021
(2023).
7   Russell, F. A., King, R., Smillie, S. J., Kodji, X. & Brain, S. D. Calcitonin gene-related
peptide: physiology and pathophysiology. *Physiol Rev* **94**, 1099-1142,
doi:10.1152/physrev.00034.2013 (2014).
8   Lu, Y. Z., Nayer, B., Singh, S. K., Alshoubaki, Y. K., Yuan, E., Park, A. J., Maruyama, K.,
Akira, S. & Martino, M. M. CGRP sensory neurons promote tissue healing via
neutrophils and macrophages. *Nature* **628**, 604-611, doi:10.1038/s41586-024-07237-y
(2024).

9   Lai, N. Y., Musser, M. A., Pinho-Ribeiro, F. A., Baral, P., Jacobson, A., Ma, P., Potts, D.
E., Chen, Z., Paik, D., Soualhi, S., Yan, Y., Misra, A., Goldstein, K., Lagomarsino, V. N.,
Nordstrom, A., Sivanathan, K. N., Wallrapp, A., Kuchroo, V. K., Nowarski, R., Starnbach,
M. N., Shi, H., Surana, N. K., An, D., Wu, C., Huh, J. R., Rao, M. & Chiu, I. M. Gut-
Innervating Nociceptor Neurons Regulate Peyer's Patch Microfold Cells and SFB Levels
to Mediate Salmonella Host Defense. *Cell* **180**, 33-49 e22,
doi:10.1016/j.cell.2019.11.014 (2020).

10   El Karim, I. A., Linden, G. J., Orr, D. F. & Lundy, F. T. Antimicrobial activity of
neuropeptides against a range of micro-organisms from skin, oral, respiratory and
gastrointestinal tract sites. *J Neuroimmunol* **200**, 11-16,
doi:10.1016/j.jneuroim.2008.05.014 (2008).

11   N'Diaye, A. R., Leclerc, C., Kentache, T., Hardouin, J., Poc, C. D., Konto-Ghiorghi, Y.,
Chevalier, S., Lesouhaitier, O. & Feuilloley, M. G. Skin-bacteria communication:
Involvement of the neurohormone Calcitonin Gene Related Peptide (CGRP) in the
regulation of Staphylococcus epidermidis virulence. *Sci Rep* **6**, 35379,
doi:10.1038/srep35379 (2016).

12   Caronna, E., Alpuente, A., Torres-Ferrus, M. & Pozo-Rosich, P. CGRP monoclonal
antibodies and CGRP receptor antagonists (Gepants) in migraine prevention. *Handb*
Clin Neurol **199**, 107-124, doi:10.1016/B978-0-12-823357-3.00024-0 (2024).

13   Edvinsson, L., Haanes, K. A., Warfvinge, K. & Krause, D. N. CGRP as the target of new
migraine therapies - successful translation from bench to clinic. *Nat Rev Neurol* **14**, 338-
350, doi:10.1038/s41582-018-0003-1 (2018).

14   Kelman, L. Osmophobia and taste abnormality in migraineurs: a tertiary care study.
Headache **44**, 1019-1023, doi:10.1111/j.1526-4610.2004.04197.x (2004).

15   Huang, A. Y. & Wu, S. Y. Calcitonin Gene-Related Peptide Reduces Taste-Evoked ATP
Secretion from Mouse Taste Buds. *J Neurosci* **35**, 12714-12724,
doi:10.1523/JNEUROSCI.0100-15.2015 (2015).

16   Zdrojewicz, Z. & Januszewski, A. [Katacalcin--structure, secretion and clinical
significance]. *Postepy Hig Med Dosw* **48**, 371-380 (1994).

17   Kaneider, N. C., Egger, P., Wiedermann, F. J., Ritter, M., Woll, E. & Wiedermann, C. J.
Involvement of cyclic adenosine monophosphate-dependent protein kinase A and
pertussis toxin-sensitive G proteins in the migratory response of human CD14+
mononuclear cells to katacalcin. *J Bone Miner Res* **17**, 1872-1882,
doi:10.1359/jbmr.2002.17.10.1872 (2002).

18   Baranowsky, A., Jahn, D., Jiang, S., Yorgan, T., Ludewig, P., Appelt, J., Albrecht, K. K.,
Otto, E., Knapstein, P., Donat, A., Winneberger, J., Rosenthal, L., Kohli, P., Erdmann, C.,
Fuchs, M., Frosch, K. H., Tsitsilonis, S., Amling, M., Schinke, T. & Keller, J. Procalcitonin
is expressed in osteoblasts and limits bone resorption through inhibition of macrophage
migration during intermittent PTH treatment. *Bone Res* **10**, 9, doi:10.1038/s41413-021-
00172-y (2022).

19   Muller, B. & Becker, K. L. Procalcitonin: how a hormone became a marker and mediator
of sepsis. *Swiss Med Wkly* **131**, 595-602, doi:10.4414/smw.2001.09751 (2001).

20   Nagayama, D., Imamura, H., Endo, K., Saiki, A., Sato, Y., Yamaguchi, T., Watanabe, Y.,
Ohira, M., Shirai, K. & Tatsuno, I. Marker Of Sepsis Severity Is Associated With The
Variation In Cardio-Ankle Vascular Index (CAVI) During Sepsis Treatment. *Vasc Health*
Risk Manag **15**, 509-516, doi:10.2147/VHRM.S228506 (2019).

21   Tang, H., Huang, T., Jing, J., Shen, H. & Cui, W. Effect of procalcitonin-guided treatment
in patients with infections: a systematic review and meta-analysis. *Infection* **37**, 497-507,
doi:10.1007/s15010-009-9034-2 (2009).

22 Leli, C., Ferranti, M., Moretti, A., Al Dhahab, Z. S., Cenci, E. & Mencacci, A. Procalcitonin
levels in gram-positive, gram-negative, and fungal bloodstream infections. *Dis Markers*
2015, 701480, doi:10.1155/2015/701480 (2015).
23 Sexton, P. M., Christopoulos, G., Christopoulos, A., Nylen, E. S., Snider, R. H., Jr. &
Becker, K. L. Procalcitonin has bioactivity at calcitonin receptor family complexes:
potential mediator implications in sepsis. *Crit Care Med* **36**, 1637-1640,
doi:10.1097/CCM.0b013e318170a554 (2008).
24 Baranowsky, A., Appelt, J., Kleber, C., Lange, T., Ludewig, P., Jahn, D., Pandey, P.,
Keller, D., Rose, T., Schetler, D., Braumuller, S., Huber-Lang, M., Tsitsilonis, S., Yorgan,
T., Frosch, K. H., Amling, M., Schinke, T. & Keller, J. Procalcitonin Exerts a Mediator
Role in Septic Shock Through the Calcitonin Gene-Related Peptide Receptor. *Crit Care*
Med **49**, e41-e52, doi:10.1097/CCM.0000000000004731 (2021).
25 Messerer, D. A. C., Datzmann, T., Baranowsky, A., Peschel, L., Hoffmann, A., Groger,
M., Amling, M., Wepler, M., Nussbaum, B. L., Jiang, S., Knapstein, P., Donat, A., Calzia,
E., Radermacher, P. & Keller, J. Systemic calcitonin gene-related peptide receptor
antagonism decreases survival in a porcine model of polymicrobial sepsis: blinded
randomised controlled trial. *Br J Anaesth* **128**, 864-873, doi:10.1016/j.bja.2021.11.042
(2022).
26 Dobin, A., Davis, C. A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P.,
Chaisson, M. & Gingeras, T. R. STAR: ultrafast universal RNA-seq aligner.
Bioinformatics **29**, 15-21, doi:10.1093/bioinformatics/bts635 (2013).
27 Liao, Y., Smyth, G. K. & Shi, W. featureCounts: an efficient general purpose program for
assigning sequence reads to genomic features. *Bioinformatics* **30**, 923-930,
doi:10.1093/bioinformatics/btt656 (2014).
28 Thorvaldsdottir, H., Robinson, J. T. & Mesirov, J. P. Integrative Genomics Viewer (IGV):
high-performance genomics data visualization and exploration. *Brief Bioinform* **14**, 178-
192, doi:10.1093/bib/bbs017 (2013).
29 Zheng, G. X., Terry, J. M., Belgrader, P., Ryvkin, P., Bent, Z. W., Wilson, R., Ziraldo, S.
B., Wheeler, T. D., McDermott, G. P., Zhu, J., Gregory, M. T., Shuga, J., Montesclaros,
L., Underwood, J. G., Masquelier, D. A., Nishimura, S. Y., Schnall-Levin, M., Wyatt, P.
W., Hindson, C. M., Bharadwaj, R., Wong, A., Ness, K. D., Beppu, L. W., Deeg, H. J.,
McFarland, C., Loeb, K. R., Valente, W. J., Ericson, N. G., Stevens, E. A., Radich, J. P.,
Mikkelsen, T. S., Hindson, B. J. & Bielas, J. H. Massively parallel digital transcriptional
profiling of single cells. *Nat Commun* **8**, 14049, doi:10.1038/ncomms14049 (2017).
30 Lewandowski, B. C., Sukumaran, S. K., Margolskee, R. F. & Bachmanov, A. A.
Amiloride-Insensitive Salt Taste Is Mediated by Two Populations of Type III Taste Cells
with Distinct Transduction Mechanisms. *J Neurosci* **36**, 1942-1953,
doi:10.1523/JNEUROSCI.2947-15.2016 (2016).
31 Sukumaran, S. K., Lewandowski, B. C., Qin, Y., Kotha, R., Bachmanov, A. A. &
Margolskee, R. F. Whole transcriptome profiling of taste bud cells. *Sci Rep* **7**, 7595,
doi:10.1038/s41598-017-07746-z (2017).
32 Satija, R., Farrell, J. A., Gennert, D., Schier, A. F. & Regev, A. Spatial reconstruction of
single-cell gene expression data. *Nat Biotechnol* **33**, 495-502, doi:10.1038/nbt.3192
(2015).
33 Qin, Y., Palayyan, S. R., Zheng, X., Tian, S., Margolskee, R. F. & Sukumaran, S. K. Type
II taste cells participate in mucosal immune surveillance. *PLoS Biol* **21**, e3001647,
doi:10.1371/journal.pbio.3001647 (2023).
34 Bankhead, P., Loughrey, M. B., Fernandez, J. A., Dombrowski, Y., McArt, D. G., Dunne,
P. D., McQuaid, S., Gray, R. T., Murray, L. J., Coleman, H. G., James, J. A., Salto-Tellez,
M. & Hamilton, P. W. QuPath: Open source software for digital pathology image analysis.
Sci Rep **7**, 16878, doi:10.1038/s41598-017-17204-5 (2017).

35 Qin, Y., Sukumaran, S. K., Jyotaki, M., Redding, K., Jiang, P. & Margolskee, R. F. Gli3 is
a negative regulator of Tas1r3-expressing taste cells. *PLoS Genet* **14**, e1007058,
doi:10.1371/journal.pgen.1007058 (2018).
36 Sukumaran, S. K., Yee, K. K., Iwata, S., Kotha, R., Quezada-Calvillo, R., Nichols, B. L.,
Mohan, S., Pinto, B. M., Shigemura, N., Ninomiya, Y. & Margolskee, R. F. Taste cell-
expressed alpha-glucosidase enzymes contribute to gustatory responses to
disaccharides. *Proc Natl Acad Sci U S A* **113**, 6035-6040, doi:10.1073/pnas.1520843113
(2016).
37 Xie, W., Fisher, J. T., Lynch, T. J., Luo, M., Evans, T. I., Neff, T. L., Zhou, W., Zhang, Y.,
Ou, Y., Bunnett, N. W., Russo, A. F., Goodheart, M. J., Parekh, K. R., Liu, X. &
Engelhardt, J. F. CGRP induction in cystic fibrosis airways alters the submucosal gland
progenitor cell niche in mice. *J Clin Invest* **125**, 2179, doi:10.1172/JCI82138 (2015).
38 Zhou, Y., Feng, Y., Liang, X., Gui, S., Ren, D., Liu, Y., She, J., Zhang, X., Song, F., Yu,
L., Zhang, Y., Wang, J., Zou, Z., Mei, J., Wen, S., Yang, M., Li, X., Tan, X. & Li, Y.
Elevations in presepsin, PCT, hs-CRP, and IL-6 levels predict mortality among septic
patients in ICU. *J Leukoc Biol*, doi:10.1093/jleuko/qiae121 (2024).
39 Montavon, P. & Lindstrand, K. Immunohistochemical localization of neuron-specific
enolase and calcitonin gene-related peptide in pig taste papillae. *Regul Pept* **36**, 235-
248, doi:10.1016/0167-0115(91)90059-p (1991).
40 Montavon, P. & Lindstrand, K. Immunohistochemical localization of neuron-specific
enolase and calcitonin gene-related peptide in rat taste papillae. *Regul Pept* **36**, 219-
233, doi:10.1016/0167-0115(91)90058-o (1991).
41 Kusakabe, T., Matsuda, H., Gono, Y., Furukawa, M., Hiruma, H., Kawakami, T., Tsukuda,
M. & Takenaka, T. Immunohistochemical localisation of regulatory neuropeptides in
human circumvallate papillae. *J Anat* **192 (Pt 4)**, 557-564, doi:10.1046/j.1469-
7580.1998.19240557.x (1998).
42 Park, G. Y., Lee, G., Yoon, J., Han, J., Choi, P., Kim, M., Lee, S., Park, C., Wu, Z., Li, Y.
& Choi, M. Glia-like taste cells mediate an intercellular mode of peripheral sweet
adaptation. *Cell* **188**, 141-156 e116, doi:10.1016/j.cell.2024.10.041 (2025).
43 Michot, B., Casey, S. M. & Gibbs, J. L. Effects of Calcitonin Gene-related Peptide on
Dental Pulp Stem Cell Viability, Proliferation, and Differentiation. *J Endod* **46**, 950-956,
doi:10.1016/j.joen.2020.03.010 (2020).
44 Zhao, X., Wu, G., Zhang, J., Yu, Z. & Wang, J. Activation of CGRP receptor-mediated
signaling promotes tendon-bone healing. *Sci Adv* **10**, eadg7380,
doi:10.1126/sciadv.adg7380 (2024).
45 Kim, Y. J. & Granstein, R. D. Roles of calcitonin gene-related peptide in the skin, and
other physiological and pathophysiological functions. *Brain Behav Immun Health* **18**,
100361, doi:10.1016/j.bbih.2021.100361 (2021).
46 Li, W., Zhang, Z., Li, X., Cai, J., Li, D., Du, J., Zhang, B., Xiang, D., Li, N. & Li, Y. CGRP
derived from cardiac fibroblasts is an endogenous suppressor of cardiac fibrosis.
Cardiovasc Res **116**, 1335-1348, doi:10.1093/cvr/cvz234 (2020).
47 Wee, N. K. Y., Novak, S., Ghosh, D., Root, S. H., Dickerson, I. M. & Kalajzic, I. Inhibition
of CGRP signaling impairs fracture healing in mice. *J Orthop Res* **41**, 1228-1239,
doi:10.1002/jor.25474 (2023).

Figure 1. qPCR profiling of *Calca and Calcb* transcripts and their receptors in cDNA from taste papillae and sensory ganglia. A) Strong expression of *PrPCT, Calcb*, *Calcrl* and *Ramp1* is observed in both CVP and FOP. Weak expression of *Cgrp* is observed in CVP, while it is undetectable in FOP. *Calcrl and Ramp1* are expressed in both CVP and FOP while *Calcr, Ramp2 and Ramp3 are* not detected in either CVP or FOP. *Tas1r3* is used as a control to demonstrate the quality of taste cDNA. B) qPCR of above transcripts in geniculate and nodose-petrosal ganglia. *Cgrp* and *Calcb* are expressed in both ganglia, with stronger expression observed in the nodose-petrosal ganglion. *Calcrl, Ramp1 and Ramp2* are expressed in both ganglia and similar levels. Expression of *Ramp3*, *PrPCT* and *Calcr* is not observed in either ganglion. The taste ganglion marker genes *Tubb3, Shh, Phox2b*, and *P2rx3* are used to demonstrate the quality of ganglia cDNA. The expression of each gene is plotted as the logarithm of the ratio between its cycle threshold value and that of *Bact*. ND= not detected.

538
539
540
541

Figure 2. RNAScope analysis of *Calca* and *Calcrl* gene expression in CVP. RNAscope Hiplex
fluorescence assay was used to determine the coexpression of *Calca* with the taste cell markers *Tas1r3*
(A-A′), *Trpm5* (B-B′), *Gnat3* (C-C′) and *Ddc* (D-D′). The areas highlighted in red boxes in the top row is
magnified in the bottom rows for each set. Taste buds are highlighted by white dotted lines. A″-D″ are
venn diagrams showing the number of taste cells that co express or singly express the indicated marker
genes and *Calca*. Data are from two non-consecutive sections from two mice. E & E′ shows the
expression of *Calcrl* in fibroblasts adjacent to basal taste cells marked by *Sparc*. Strong coexpression of
Calca is observed with *Tas1r3* and *Trpm5*, less strong coexpression is observed with *Gnat3* and
negligible coexpression is observed with *Ddc*. *Calcrl* expression is observed in basal (presumably
stem/progenitor) cells in the taste buds and adjacent fibroblasts. Filled white arrowheads highlights single
positive cells, filled red arrowheads highlights double positive cells, and open red arrowheads depicts
basal cells in taste buds expressing *Calcrl*. Scale bars = 30 µm.

Figure 3. PCT is expressed in type II taste cells. Double labelled immunofluorescence confocal
microscopy of CVP and FOP sections with antibodies against PCT (green) and type II taste cell markers
T1R3 (red; A–D, F-I) or LRMP (red; K–N). Nuclei are counterstained with DAPI (blue). A, F, K are lower
magnification images and dashed red boxes indicate regions shown at higher magnification in panels in
the right. (D, H, L) Higher-magnification views of the boxed region showing double positive cells with solid
arrows and single positive cells hollow arrows. Colocalization counts are shown in Venn diagrams (E,J,O).
Scale bars = 20 µm.

Supplementary data

Figure S1. Sashimi plot of the *Calca* gene locus (transcribed from bottom strand in right to left direction) from bulk RNASeq data from CVP. The reads aligning to exons 1-6 (Ex1-Ex6, right to left) are shown in red, with the height of the red bars showing the strength of expression, and the number of reads aligning across splice junctions indicated in the loops between exons. No reads align to the preproCGRP specific exon 5, while large number of reads align to prePCT specific exon 4.

Figure S2. End point PCR showing expression of Calca- related mRNAs. **A)** Expression of indicated transcripts in taste papillae and non-taste lingual epithelium (NT). *PrPCT, Calcb*, *Calcrl* and *Ramp1* expression is observed in CVP, FFP and FOP. Weak *Cgrp expression* is observed in CVP and FOP, while it is undetectable in FOP and FFP. *Calcrl and Ramp1* are expressed in all taste papillae while *Calcr, Ramp2 and Ramp3 are* not detected in any tissue. None of the tested mRNAs are detected in NT. *Tas1r3* is used as a control to demonstrate the quality of taste cDNA, and *Bact* is used as cDNA synthesis control. **B)** qPCR of above transcripts in the geniculate and nodose-petrosal ganglia. *Cgrp, Calcb, Calcrl, Ramp1 and Ramp2* are expressed in both ganglia. Expression of *Ramp3, PrPCT* and *Calcr* is not observed in either ganglion. *Bact* is used as cDNA synthesis control, while the taste ganglion marker genes *Tubb3, Shh, Phox2b*, and *P2rx3* are used to demonstrate the quality of ganglia cDNA.

Figure S3. Dot plot from CVP. Type I a-d, type IIIa-c and type IV, sweet and bitter taste receptor cells are
detected. BitterPrecursor and sweetPrecursor are clusters of Immature type II immature cells. TasteStem
is a cluster of stem cells. Strong *Calca* (=prePCT) mRNA expression is seen in sweet taste receptor cells
and immature type II cells, and *Calcrl* expression is observed in type I and taste stem cells, and *Crcp*
expression is found in multiple cell types. *Ramp1* could not be detected in this dataset.

Figure S4. Western blot analysis of protein extract from CVP of male and female mice with anti-PCT
antibody showing expression of PCT.

Figure S5. Double labeling for GNAT3 and LRMP. A-F) Unlike PCT staining shown in Figure 3, GNAT3 staining is restricted to weaker LRMP expressing cells with less intense staining for LRMP. **G)** Venn diagram shows quantification of coexpression data.

Figure S6. Controls for immunostaining with two rabbit primary antibodies. Double indirect immunofluorescence confocal microscopy of CVP shows cross-labeling when unlabelled donkey Fab fragment is omitted (control A, *I & II*), whereas adequate blocking of rabbit IgG is achieved with excess donkey Fab fragment (control B, I and II).

Probe	CAT number	LOT number
RNAscope™ HiPlex Probe- Mm-Ddc-T1	318681-T1	22363A
RNAscope™ HiPlex Probe- Mm-Tas1r3-T2	515431-T2	22174B
RNAscope™ HiPlex Probe- Mm-Sparc-T4	466781-T4	24017A
RNAscope™ HiPlex Probe- Mm-Calcrl-T5	452281-T5	24017A
RNAscope™ HiPlex Probe- Mm-Calca-T6	578771-T6	223634
RNAscope™ HiPlex Probe- Mm-Gnat3-T8	531661-T8	22171A
RNAscope™ HiPlex Probe- Mm-Trpm5-T9	453251-T9	22363A

614 **Table S1:** List of RNAscope HiPlex Probes used in this study.

Antibody	Host Species	Catalog No.	RRID	Source	Dilutions
PCT	Rabbit	LS-C296040	-	LS Bio, Newark, CA	1: 50
GNAT3	Goat	OAEB00418	AB_1088282	Aviva Systems Biology, San Diego, CA	1:500
LRMP	Rabbit	ORB166443	-	Biorbyt, Durham, NC	1:600
T1R3	Rabbit	-	-	Gift from Dr. Ichiro Matsumoto, Monell Chemical Senses center	1:500
Alexa Fluor 488 Conjugated AffiniPure Fab Fragmant Goat anti-Rabbit IgG (H+L)	Goat	111-547-003	-	Jackson Immuno Research Inc. West Grove, PA	1:500

Alexa Fluor 647 Conjugated AffiniPure Fab Fragmant Donkey anti-Rabbit IgG (H+L)	Donkey	711-607-003	-	Jackson Immuno Research Inc. West Grove, PA	1:500
Alexa Fluor 647 Conjugated AffiniPure Donkey anti-Goat IgG (H+L)	Donkey	705-606-147	-	Jackson Immuno Research Inc. West Grove, PA	1:500
Alexa Fluor 488 Conjugated Donkey anti-Goat IgG	Donkey	A11055	-	Jackson Immuno Research Inc. West Grove, PA	1:500
AffiniPure Fab Fragment Donkey anti-Rabbit IgG(H+L)	Donkey	711-007-003	-	Jackson Immuno Research Inc. West Grove, PA	1:500

617 **Table S2:** Primary and secondary antibodies for IHC and western blot used in this study.

Gene	Forward Primer	Reverse Primer
Bact	GGCTGTATTCCCCTCCACG	CCAGTTGGTAACAATGCCATGT
Calcrl	CATCGTGGTGGCTGTGTTT	GTAATACAAGCTTCTGGCAATGG
Calcr	GCTGAGTGCAGAAACCCACT	TTTGCCTCATCTTGGTCACA
Ramp1	AGCCGCTTCAAGGAGAACAT	CGTGCTTGGTGCAGTAAGTG
Ramp2	TGAGGACAGCCTTGTGTCAA	CAGCACAGCAGAAAGGTTCC
Ramp3	AAGTTGGTTTTGGACGGTGA	GCATACCTGGGCACACTCA
Tubb3	GAACCTGGAACCATGGACAG	GTTGTTGCCAGCACCACTCT

Shh	AAGGATGAGGAAAACACGGG	GGTCACTCGCAGCTTCACTC
Phox2b	CACTTTTGGGGCCACGTC	CGTGGTCGGTGAAGAGTTTG
P2rx3	CCAAATATTCCTTCACTCGGC	GCTGCCATTCTCCATCTTGT
PrePCT	GGAGCAGGAGGAAGAGCAGG	GCCAGGTGCTTCAACCCCAA
Cgrp	TATGCAGATGAAAGCCAGGG	GTGGCAGTGTTGCAGGATCT
Calcb	CAGGAAGCTGGAACAGGAGG	AAGGCTTCAGAGCCCACATC
Tas1r3	CAAGTTCTTCAGCTTCTTCC	GGCGGCCACCCAGTTCCAGC

621 **Table S3:** List of primers used for PCR and qPCR.