Minimal Mimics and Maps of Natural Light for Mammals

Philippe Morquette[*] and Michael Tri H. Do[*]

F. M. Kirby Neurobiology Center and Department of Neurology, Boston Children's Hospital and Harvard Medical School. Boston, MA 02115, USA

*Correspondence: philippe.morquette@childrens.harvard.edu,
michael.do@childrens.harvard.edu

ABSTRACT

Light drives processes that include perception and the regulation of circadian rhythms, sleep, metabolism, and development. These processes are initiated by photopigment molecules, each preferentially absorbing particular wavelengths. Light of a given spectrum stimulates an animal's set of photopigments in a specific profile. Natural light and its variations produce stimulation profiles that promote normal physiology. To mimic these profiles using artificial light, we consider the thermally stable, photoconvertible states of relevant photopigments: ground states of rhodopsin and cone photopigments, and three states of melanopsin. This gives a high-dimensional representation of illumination. Nevertheless, we find that two wavelengths suffice to closely mimic the effects of natural light for mammals, including humans and mice. Adjusting the wavelength ratio allows mimicry of natural light's variations, such as those from twilight to noon. Ratio adjustments also compensate for light's filtering by elements like the eye's optics and laboratory cages. Adding a third wavelength makes natural light mimicry nearly perfect. By contrast, common artificial lighting—designed for low-dimensional, human color space—stimulates photopigments in unnatural proportions. We conclude by providing species-specific maps of photopigment stimulation profiles under natural and artificial illumination, which make our observations intuitive while providing insight into the diverse visual ecologies of mammals.

SIGNIFICANCE STATEMENT

Humans sense light for vital processes like sight and physiological regulation. These processes are normal in natural light, where they evolved. However, much of modern life is spent in artificial light, which is unlike natural light in many of its biological effects. This disparity has been linked to disorders that range from cardiovascular disease to cancer. This manuscript introduces simple forms of artificial lighting that replicate the effects of natural light on photoreceptors of humans and other species. It also demonstrates how the biological effects of natural and artificial lights can be captured in simple maps, facilitating the choice and further design of illumination that is beneficial.

KEYWORDS

Natural light, Daylight, Artificial light, Opsin, Photopigment, Photoreceptor, Metamer, Image-forming vision, Non-image-forming vision, Cone Opsin (Opn1), Rhodopsin (Opn2), Melanopsin (Opn4), Neuropsin (Opn5), Bistability, Tristability, Multistability

INTRODUCTION

Humans and other animals rely on light for perception and the regulation of processes that include development, cardiovascular function, circadian rhythms, sleep, metabolism, cognition,

and mood (1, 2). Mammals sense light using photopigments, G protein coupled receptors
(opsins) bound to chromophore. Three types are known to trigger electrical signals: rhodopsin,
cone photopigments, and melanopsin (2, 3). Each absorbs light of particular wavelengths
between ultraviolet and near-infrared. Thus, various spectra of illumination produce specific
profiles of photopigment stimulation. A given spectrum of illumination produces profiles that
differ among species whose photopigments vary in their absorption spectra.
58
Mammalian photopigments evolved under ambient illumination from the sun. The spectrum and
intensity of this natural light support normal biological processes. However, most light in modern
society is artificial, which tends to be unnatural in its spectrum, too bright at night, and too dim
during the day (4, 5). Even natural light, when passed through windows and other materials,
usually becomes unnatural. Abnormal lighting impairs the development of retinal structure and
circadian period, increases mortality from cardiovascular events, dysregulates mood, and
misaligns circadian rhythms to raise the risk of disorders such as cancer (2, 4, 6, 7). Conversely,
light is therapeutic when properly designed and controlled (8, 9). For these reasons, it is
important to develop artificial light that stimulates photoreceptive mechanisms naturally.
68
An artificial mimic of natural light should produce the same photopigment stimulation profile as
ambient solar illumination. Most artificial lights appear natural to humans because their design is
based on cone photopigments, which drive color perception. Rhodopsin is generally thought to
operate only in dim light; however, it may have a broader role (10). Melanopsin is also essential
(2, 11). Indeed, a new lighting standard incorporates melanopsin (12). However, it only concerns
melanopsin's ground state in darkness. In light, melanopsin photoconverts among three states
that are spectrally distinct and fundamentally shape melanopsin's actions; for instance, the
active state may switch into an inactive state (13-16). Thus, a photostimulation profile of six
photopigment states should be considered for dichromatic mammals and seven for trichromatic
mammals.
79
A natural light mimic should be tunable to compensate for elements in its path to the
photopigments, a process called prereceptoral filtering. Common prereceptoral filters include
window materials, lamp coverings, and the eye's optics. Their effects can be severe. For
example, aging of the human lens reduces transmission, especially of short wavelengths,
potentially contributing to the dysregulation of circadian rhythms and sleep (17). Most laboratory
cages attenuate the ultraviolet wavelengths that mice and many other species sense for
perception and physiological regulation (18). Conversely, even if daylight were given directly to
the retina, as in *ex vivo* physiological studies, it would act differently and potentially unnaturally
due to the absence of filtering by the eye. The inability to deliver the same, naturalistic stimulus
ex vivo and *in vivo* impedes efforts to connect biological mechanisms to behavior.
Compensation is challenging because most light sources and prereceptoral filters have complex
spectra.
92
We sought to define how natural light—defined here as ambient solar illumination at nautical
twilight and above—stimulates rhodopsin, cone photopigments, and melanopsin's three states
for various species. We then asked if these photopigment stimulation profiles could be mimicked
with simple forms of artificial light. We found that two wavelengths are effective and three
appear nearly perfect. Adjusting the ratio of wavelengths compensates for prereceptoral filtering
and enables mimicry of natural light and its variations. We constructed minimal natural light
mimics for humans and found that their performances are close to our theoretical predictions.
Finally, we found that photopigment stimulation profiles in natural light are captured in two-
dimensional maps for each species tested, facilitating the study of photoreception in an
ecological context and further improvements in artificial lighting design.

103
104 **RESULTS**
105
We will start by explaining our process for estimating spectra for natural light, ocular
transmission, and photopigment absorption. We will use these spectra to design natural light
mimics using two or three wavelengths delivered in the correct ratio for mice, rats, cats, rabbits,
dogs, and humans. We then compare the accuracy of wavelength mimics in theory and practice
for humans. Throughout, we provide simple maps of photopigment stimulation profiles under
natural and artificial illumination. We will focus our analyses on photopigments that are known to
trigger electrical responses in the mammalian retina, those of rods (rhodopsin), cones (cone
photopigments), and intrinsically photosensitive retinal ganglion cells (ipRGCs; melanopsin).
114
115 **Choosing photopigment absorption spectra**
116
We consider photopigment states that are stable and thus likely to absorb a photon and produce
a biological effect (i.e., we ignore short-lived intermediate states). For rhodopsin and cone
photopigments, which are monostable, this is only the ground state; following activation, the
opsin loses its chromophore and becomes insensitive to light (19). Mammalian melanopsin is
multistable (13-16). Its three known conformations are melanopsin (the R state, non-signaling),
extramelanopsin (E, non-signaling), and metamelanopsin (M, signaling). Photon absorption by
R or E can produce M, while absorption by M can produce R or E. Melanopsin's actions depend
on the photoequilibration of these states.
125
Several absorption spectra have been reported for each photopigment state, with varying
wavelengths of peak sensitivity (λ_{max}). We prioritize *in vitro* measures because they are direct
and choose λ_{max} values that are most consistent across spectrophotometry (purified
photopigment), microspectrophotometry (photopigment in cells), and electrophysiology (the
electrical activity of photopigment in cells)(**Table S1**). We use the mathematical function of
Govardovskii and colleagues to reconstruct each photopigment state's absorption spectrum,
with only the λ_{max} as a free parameter (20). These reconstructed spectra are highly accurate
across log units of variation in absorbance, though complexities exist (15, 21). We also
corrected absorption spectra for self-screening of photopigments in mouse and human
photoreceptors (22, 23).
136
137 **Mimicking natural light for mouse photopigments**
138
We begin with the mouse because of its widespread use as a model system and detailed
knowledge of its photoreceptive mechanisms. We use photopigment λ_{max} values of 502
(rhodopsin, Opn2)(24), 358 (ultraviolet-sensitive cone photopigment, Opn1SW)(25) and 508 nm
(medium-wavelength-sensitive cone photopigment, Opn1MW)(24, 25). For the R, E, and M
states of melanopsin (Opn4-R, Opn4-E, and Opn4-M), these values are 471, 453, and 476 nm,
respectively (13-15). We take the R and E state λ_{max} values from electrophysiological
measurements of mouse ipRGCs, as they concern melanopsin in its native environment (14).
These values are slightly red-shifted from those of purified mouse melanopsin (467 and 446 nm,
respectively)(13). The M state's λ_{max} is only known for purified melanopsin, so we use it as is
(i.e., without imposing a red shift that may be inferred from electrophysiological assessments of
the other states).
150
The light that reaches the retinal photopigments is shaped by the transmission spectrum of the
eye, primarily the cornea and lens. A spectrum has been published for the mouse eye but only
for a limited wavelength range (26). Thus, we dissected the mouse eye, removed a portion of

the posterior pole, and mounted the remainder in a custom cuvette for spectrophotometry
(**Supplementary Methods**). We worked swiftly and used oxygenated media to maintain tissue
health. Compared to the published spectrum, ours has higher transmission of short wavelengths
(**Fig. S1**).
158
Multiplying light's spectrum (photons μm^{-2} s^{-1} nm^{-1}), the ocular transmission spectrum
(normalized to its peak), and a photopigment state's absorption spectrum (normalized to its
peak) yields the light that is available to that state (**Fig. 1A**). Downstream events—for instance,
the light that the photopigment actually absorbs, the probability that the photopigment activates,
and the probability that activated photopigment triggers its G protein cascade—follow from
photon availability (**Supplementary Methods**)(27-29). Integrating the product gives the total
photon flux density that is available for the photopigment state in question (**Fig. 1B**). For brevity,
we refer to this integral as a photopigment's stimulation.
167
We first consider a standard daylight spectrum, ASTM International's G173-tilted (30), which
captures both direct and diffuse illumination at mid-day under a cloudless sky. We examined the
range from 300 nm (photon flux is scarce at shorter wavelengths, especially after attenuation by
the cornea and lens) to 700 nm (at which point the mouse photopigment sensitive to the longest
wavelength only absorbs at ~10^{-5} of its peak) at a resolution of <1 nm. The G173 spectrum, with
consideration of the eye's optics and each photopigment's absorption, stimulates Opn1SW,
Opn4-E, Opn4-R, Opn4-M, Opn2, and Opn1MW photopigments at $2.38×10^8$, $1.37×10^9$,
$1.61×10^9$, $1.68×10^9$, $2.28×10^9$, and $2.19×10^9$ photons μm^{-2} s^{-1}, respectively (**Fig. 1B**; we order
photopigments from shortest to longest λ_{max}). This photopigment stimulation profile reveals a
nearly 10-fold difference in daylight's stimulation across the mouse photopigments.
178
Inspection of the mouse's photopigment absorption spectra (**Fig. 1A**) suggests that only two
wavelengths may suffice to mimic the daylight photopigment stimulation profile. That is because
one wavelength can be chosen to intersect most absorption spectra at various levels, and a
second wavelength can provide fine tuning. To test this idea, we searched for a pair of
wavelengths that produce an artificial profile with the lowest deviation from the G173 profile
(**Supplementary Methods**). We focused on the spectrum of light rather than its intensity by
normalizing the profiles of G173 and potential 2-wavelength mimics (i.e., comparing their unit
vectors). We found that optimal wavelengths were 403.3 and 512.2 nm, given at an intensity
ratio of 1.26:1 (in units of photons; **Fig. 1C-D**). This 2-wavelength G173 mimic has a stimulation
profile that deviates <1.1% from that of G173 for each photopigment; averaged over all
photopigments, its deviation is 0.60% (relative root mean squared error, RRMSE). The overall
intensity of the two wavelengths can be adjusted to match the photon flux density of G173 or
any desired illumination level, from the threshold of perception to maximum daylight.
192
193 **Mimicking natural light's variations for mouse photopigments**
194
To ask if two wavelengths can mimic natural light's variations, we compiled a large number of
published spectra. We chose spectra that have a range of at least 300 to 700 nm (see above).
We also selected for a minimum resolution of 5 nm. A ≤10-nm band is often considered
monochromatic in vision studies. Extending the wavelength range or resolution would greatly
reduce the available spectra. We arrived at a set of 3,567 spectra measured in Pennsylvania,
USA (31) and Granada, Spain (32). Together, they cover solar elevations from nautical twilight
to daylight (**Fig. 2A**). These spectra are in units of photon flux density (photons μm^{-2} s^{-1} nm^{-1})(28). We multiplied each spectrum by those of the mouse eye and its photopigments to yield a
large set of photopigment stimulation profiles, which we refer to as the 3K profiles. Again, to
large set of photopigment stimulation profiles, which we refer to as the 3K profiles. Again, to

focus on the spectrum rather than intensity of light, we converted each profile to its unit vector
(**Fig. 2B**).
206
We asked if the wavelengths we found for mimicking G173 (403.3 and 512.2 nm) could be used
to mimic the 3K profiles. Adjusting the ratio of these wavelengths from 0.35 to 3.57 allowed
mimicry of 90.1% of the 3K profiles with <5% deviation (**Fig. 2C**). We also optimized the two
wavelengths for all 3K profiles, obtaining values of 402.5 and 508.9 nm. Changing their intensity
ratio suffices to mimic 98.0% of the 3K profiles with <5% deviation (**Fig. S5A**). The overall
intensity of these wavelengths can be adjusted to match any level of environmental illumination.
Therefore, two wavelengths in the correct ratio largely suffice to mimic the spectrum of natural
light as it varies across parameters that include geographical location and meteorological
conditions.
216
Daily variations in the spectrum of natural light are salient cues for processes that include
synchronization of the circadian clock with the day-night cycle (27, 33, 34). These spectral
changes are more pronounced from astronomical to civil twilight, when the sun approaches the
horizon (31). Therefore, we assessed the effectiveness of a 2-wavelength mimic that is
specifically designed to capture these variations. We pooled 3K profiles in 1° bins of solar
elevation, from -13 to 76° (just below nautical twilight to the highest elevation in the 3K spectra).
This gave the average trajectory of daily spectral variations. We found two optimal wavelengths,
402.6 and 510.8 nm which, given in ratios between 1.00 to 1.40, have <2% deviation from
profiles of spectra measured at solar elevations above 0° (**Fig. 2D**). Accuracy declined below 0°,
to a maximum of 4.1% deviation. In practice, this decrease in accuracy has little consequence
because it is most prominent when only rods are sensitive enough to drive photoreception (i.e.
photon flux density lower than ~200 photons μm^{-2} s^{-1} between 300 and 700 nm, corresponding
to light intensities below nautical twilight, **Fig. 2E**). In this regime, one only needs to consider
how the wavelength of light stimulates rhodopsin, and the target stimulation level can be
achieved simply by adjusting the intensity. Hence, two wavelengths given in varying proportions
suffice to mimic natural light's variations from twilight to full daylight for mouse photopigments.
233
Given the effectiveness of two wavelengths, we asked if three wavelengths provided a
substantial improvement (**Fig. S8D**). With three wavelengths, mimicry of the G173 profile
appears virtually perfect, with RRMSE values of <1×10^{-6}%. Three wavelengths also increased
accuracy for mimicking any of the 3K natural light profiles (**Fig. S6A**). Thus, we consider three
wavelengths to be ideal for mimicking natural light, and two wavelengths to be both accurate
and practical.
240
241 **A map of mouse photopigment stimulation profiles**
242
We found that the 3K profiles were almost entirely captured in two dimensions, with 99.9% of
their variance explained by the first two principal components (PC). Thus, the projection of the
3K profiles onto these PCs provides a simple and complete map of how natural light's variations
stimulate the mouse photopigments. Indeed, changes in solar elevation produce profiles that
follow an orderly trajectory through the map (**Fig. 3A**). Profiles of other available natural light
spectra fell into the cloud formed by the 3K profiles (35)(not shown). The profiles of G173, and
all aforementioned mimics were also within the cloud (**Fig. 3B**).
250
We then investigated how artificial lights stimulate mouse photopigments. Given the vast variety
of artificial lights, we began by examining 42 standard spectra that represent typical sources
(36). All produced photopigment stimulation profiles that deviated substantially from that of
G173 (>25% RRMSE; **Table S2**). However, many of these standards lack shorter wavelengths;

they were designed for human visual perception, which is negligible for those wavelengths due
to strong prereceptoral filtering and the relatively red-shifted absorption spectra of cone
photopigments (see below). Shorter wavelengths are relevant for mouse photoreception (**Fig.**
1A)(18). Therefore, we measured a selection of xenon, LED, halogen, incandescent,
fluorescent, and monochromatic (10-nm band) sources and calculated their photopigment
stimulation profiles (**Fig. S2A-B**). Xenon was closest to G173. Its maximum deviation, for
Opn1SW, was 36%. All other artificial profiles had deviations of >71% for at least one
photopigment (**Fig. S2B-C**). Finally, we considered monochromatic lights in theoretical, 0.01-nm
bands from 380-700 nm. Each of their profiles deviated from that of G173 by >97% for one or
more photopigments. Substantial deviations are also evident when considering the average of
all photopigment stimulations in each profile (RRMSE; **Fig. S2D**). Thus, all artificial lights
examined differed substantially from a natural light standard.
267
To consider how artificial lights compare to a range of natural lights, we projected their
photopigment stimulation profiles onto the 3K profile map. Most broadband, artificial profiles
were far from or on the periphery of the natural profile cloud; xenon was an exception (**Fig. 3C,**
and **Supplementary Methods**). The profiles of monochromatic light were distant from the
cloud, delineating a broad arc around it (**Fig. 3C**). Therefore, even when considering the many
varieties of ambient solar illumination, most artificial lights are outliers in their actions on the
mouse visual system.
275
We also investigated how artificial lights stimulate mouse photopigments under two conditions
that are common in research environments. The first incorporates additional prereceptoral
filtering by a "clear" cage (37). The cage-filtered G173 and xenon profiles fell outside the 3K
cloud. We also considered a condition of less prereceptoral filtering: When the retina has been
dissected free of the eye, as for *ex vivo* physiological study. The less-filtered xenon and G173
profiles fell outside the 3K cloud or shifted within it, respectively (**Fig. 3C**). These analyses
underscore the importance of prereceptoral filtering even for natural light. Compensating for
prereceptoral filtering of broadband illumination is impractical due to the complexity of the filter
and the light. Compensating for monochromatic light is simple, requiring only an adjustment of
intensity, but monochromatic light remains unnatural in its effect on the mouse photopigments.
On the other hand, the ratios of our 2-wavelength mimics can be adjusted for prereceptoral
filters of any shape, if the wavelengths are transmitted sufficiently.
288
289 **Mimicking and mapping natural light for other dichromatic mammals**
290
We investigated the construction of mimics and maps for other dichromatic mammalian species:
rats, cats, rabbits, and dogs. We used published estimates of ocular transmission spectra and
λ_{max} values of rod and cone photopigments (38-40)(**Table S1**). Melanopsin's individual states
have not been defined for these species. However, we have found that the R and E states have
similar λ_{max} values in mouse and macaque monkey, suggesting broad evolutionary conservation
(14, 16). Furthermore, melanopsin's λ_{max} appears similar across many mammalian species, from
humans to whales (41); as estimated, these λ_{max} values are likely to represent activation from
both R and E states, suggesting that their individual λ_{max} values are conserved (15). For these
reasons, we adopt the λ_{max} values reported for mouse melanopsin's R, M, and E states (13-15).
300
We tested the possibility of accurately mimicking natural light and its variations for these species
using a minimal spectrum. We found that two wavelengths in the correct proportion mimicked
the G173 profile with deviation of <5% across all species (RRMSE; **Fig. S3**). For each species
except dog, the mimic profile was more similar to the G173 profile than the artificial broadband
or monochromatic profiles (**Fig. S4** and **Table S2**). Finally, for all species, adjusting the intensity

ratio of the two wavelengths (optimized for the 3K profiles) sufficed to replicate variations of
natural light in the 3K cloud (**Fig. S5**) and to correct for different prereceptoral filters (not
shown). Thus, a 2-wavelength natural-light mimic performs well for the dichromatic mammals
tested here. As for the mouse, a third wavelength increased accuracy, though two wavelengths
appear sufficient (**Figs. S6*B-E*** and **S8*D***).

Using the 3K natural light spectra (**Fig. 2*A***), relevant ocular transmission spectra and
photopigment λ_{max} values, we constructed maps of natural photopigment stimulation profiles for
these species (31, 32, 38-40)(**Fig. S7**). We found that these maps resembled one another and
that of the mouse. For each species, two PCs captured >99.8% of the variance, 3K profiles
showed orderly variation across the maps with solar elevation (not shown), profiles for most of
our measured, broadband lights bordered or were outside the 3K cloud, and profiles of
monochromatic lights surrounded the 3K-profile cloud at a distance (**Fig. S7**). Thus, these maps
are largely conserved across dichromatic species despite their varied ocular transmission
spectra and photopigment λ_{max} values. A trivial explanation for this conservation is that we used
the same λ_{max} values for melanopsin, which accounts for three of the six photopigment states.
However, conservation remains when omitting melanopsin (not shown; **Supplementary**
Methods).

Mimicking and mapping natural light for humans

Most humans are trichromatic. We asked whether a small set of wavelengths could reproduce
natural light's stimulation of their seven photopigment states (compared to the six of dichromatic
mammals). We used a standard transmission spectrum of the eye (23), and published λ_{max}
values for the rod and cone photopigments (**Table S1**; **Supplementary Methods**)(42, 43). Two
variants of the Opn1LW photopigment are common in the population, and we used their
average λ_{max} (554.55 nm)(43). For melanopsin, we adopted the λ_{max} values for macaque
melanopsin's R and E states, which are slightly shorter than those of the mouse (16). The M
state's λ_{max} is undefined, so we used that of mouse melanopsin (13). The photoswitching of
macaque melanopsin is similar to that of mouse melanopsin (14-16); because this property
depends on the M state's λ_{max}, any difference in this value between macaque and mouse is likely
to be small. We accounted for self-screening of rod and cone photopigments (**Supplementary**
Methods)(22). We omitted macular pigment to focus on the majority of the retina (but see
below).

For a 2-wavelength mimic of G173, we found that the optimum was 470.2 and 557.7 nm in a
ratio of 0.64:1. Its profile deviated from that of G173 by <8% for any given photopigment; across
photopigments, the RRMSE was 5.53% (**Fig. S8*A*** and ***C***). We also considered a common
standard for human daylight perception, CIE D65 (36). It is a mathematical model with a
correlated color temperature of ~6500 K. Its optimum 2-wavelength mimic comprised 469.3 and
556.0 nm in a ratio of 0.71:1. The profile of the D65 mimic deviated from that of D65 itself by
<8% for any given photopigment and 5.20% when averaged across all photopigments (not
shown).

Next, we tested mimicry with three wavelengths. For G173 and D65, the optimal wavelengths
were 461.7, 523.3, and 592.6 nm in a 0.64:0.77:1 ratio, and 460.9, 522.4, and 591.4 nm in a
0.75:0.85:1 ratio, respectively. These 3-wavelength mimics deviated from G173 or D65 by
<0.3% for any photopigment; the average deviations were 0.13 and 0.12%, respectively (**Fig.**
S8*B-C*, not shown for D65). Given this high accuracy, we asked if a fixed set of three
wavelengths could mimic any of the 3K natural light profiles (**Supplementary Methods**). We
found that a combination of 460.2, 520.7 and 591.0 nm, with the relative intensity of any one

wavelength ranging from 0.21-2.27 (normalized to the last wavelength), was optimal. Across all
3K profiles, this 3-wavelength mimic (with varying ratio) had a photopigment stimulation profile
that deviated <1.83% for any photopigment and 0.36 ± 0.16% on average (**Fig. S6F**). Thus, a 3-
wavelength mimic can replicate natural light's meteorological and daily variations.
361
We constructed a map of natural photopigment stimulation profiles for humans, using the set of
3K natural light spectra. It differed in shape from those of dichromatic species. Nevertheless,
two PCs captured 99.9% of the variance (**Fig. 4A**). Moreover, 3K profiles showed orderly
variation across the maps with solar elevation (not shown); profiles for most artificial, broadband
lights bordered or lay outside the cloud; and profiles for monochromatic lights surrounded the
cloud at a distance. Hence, most artificial lights, including many that are designed to appear
natural to humans, stimulate the retinal photopigments unnaturally. By contrast, the profiles of
the 2- and 3-wavelength mimics of G173 and D65 spectra were near or within the cloud of 3K
profiles, with relatively low deviation (RRMSE values; **Figs. 4A** and **S8F**).
371
372   **Comparison of natural and artificial photopigment stimulation profiles**
373
To assess differences between natural and artificial profiles that may lie outside the two
dimensions of the 3K maps, we applied cluster analysis to all dimensions; six for dichromatic
species and seven for humans (**Supplementary Methods**). We detail the outcomes in **Table**
S3. For each species, >99% of the 3K profiles were classified as part of one cluster, along with
the G173 standard. Across all species, profiles of most artificial lights were excluded from this
core: >95.2% for standard spectra and all except xenon for our set of measured spectra.
Furthermore, when G173 was filtered by cage material or lacked filtering by the eye (see
above), its profile was often excluded from the core. This analysis underscores the abnormal
effect of most artificial light on the mammalian photopigments, and the importance of
prereceptoral filtering on natural light.
384
The core 3K profiles clustered with profiles of all 3-wavelength mimics for all species and across
all conditions assessed (i.e., *in vivo*, *ex vivo*, and cage) and the 2-wavelength mimics of mouse
and cat (for all conditions assessed; **Table S3**). The profiles of the remaining mimics deviated
little from those of natural light (e.g., <5% RRMSE for G173). Their exclusion likely reflects the
clustering method's sensitivity to covariation; photopigment stimulations covary across natural
spectra in a way that can be discontinuous with the profile induced by the mimic. This
discontinuity is subtle compared to the qualitatively different profiles under most of the common
sources of artificial light.
393
394   **Construction and evaluation of human natural light mimics**
395
We designed, built, and tested 2- and 3-wavelength natural-light mimics for humans. Actual
illumination spectra are broader than the theoretical, single wavelengths given above.
Therefore, we asked how deviations from theoretical wavelength and ratio optima affected
mimic accuracy (as illustrated for the mouse in **Fig. 1C-D**). We found regions of shallowly
changing, relatively low average deviation (RRMSE <10%) and designed our mimics around
them (**Supplementary Methods**). We selected two or three LEDs, collimated their outputs, sent
the beams through bandpass filters, combined the beams with dichroic mirrors, focused the
combined beam onto one end of a liquid light guide, and connected the opposite end to a
custom Ganzfeld sphere (**Fig. 4B**; **Supplementary Methods**).
405
We used a Ganzfeld because it produces diffuse illumination, like ambient natural light. This
stimulus is also precisely quantifiable, as each point in the retina receives an intensity that is

nearly the product of the Ganzfeld's output, the attenuation by the eye's optics (cornea, iris, and
lens), and the inverse of the retinal surface area. Our Ganzfelds had a 30.5- or 12.7-cm outer
diameter, a thick internal shell of $BaSO_4$ (which reflects relevant wavelengths evenly), and a 13-
or 3.4-cm diameter viewing port (the former filling much of one's field of view and the latter
achieving higher intensities; **Supplementary Methods**). We measured light at the viewing port
of the Ganzfelds with a calibrated spectrometer and power meter, converting units to photons
μm^{-2} s^{-1} nm^{-1}, and used those measurements for evaluation.
415
We focused on the D65 daylight spectrum because it is a common standard for human vision
(11). We found that the constructed mimics performed close to our theoretical predictions. For
the constructed, 2-wavelength D65 mimic, the outputted spectra had peak wavelengths at 467.7
nm (full-width at half-maximum, FWHM, of 4.7 nm) and 554.5 nm (10.5 nm) in a ratio of 0.69:1
(integrals of the photon flux density across wavelength; not shown). Compared to the D65
standard, this constructed mimic had a maximum deviation of <8.5% for any given
photopigment and an average deviation of 5.35% (compared to 5.20% for the theoretical 2-
wavelength D65 mimic, see earlier). For the constructed, 3-wavelength D65 mimic, the peak
wavelengths were 460.0 nm (FWHM of 8.6 nm), 522.6 nm (9.6 nm), and 592.6 nm (9.8 nm)
(**Fig. 4C**). Their ratios were 0.72:0.92:1. This mimic had a maximum deviation of <1.6% for any
given photopigment and an average deviation of 0.86% (compared to 0.12% for the theoretical
3-λ D65 mimic, see earlier). The constructed 2- and 3-wavelength mimics had profiles that fell
near or within the cloud of 3K natural light profiles, respectively (**Fig. 4A**), and the latter
clustered with the core.
430
The smaller Ganzfeld gave a maximum, total output of 6.5×10^7 photons μm^{-2} s^{-1} (integrated
across the measured spectrum). Using the CIE photopic luminosity function (44), this
corresponds to ~11,000 lux and is equivalent to ambient illumination at ~10-20° of solar
elevation (45). The maximum output of the 30.5-cm Ganzfeld was ~1,100 lux (ca. 0° solar
elevation). Both outputs meet safety standards (46). They can be controlled finely over a broad
range—for example, by using currents of different amplitude, pulse-width modulation, and
neutral density filters—and their maximum outputs increased, if desired, using more intense
sources.
439
We designed our natural light mimics for all photopigments that are known to trigger electrical
signals in humans, thus replicating light's actions on both image vision (perception, which is
primarily mediated by rod and cone photopigments) and non-image vision (e.g., circadian
photoregulation, for which melanopsin is vital)(1, 2, 11). We consider nearly as many
photopigment states for the former (four) as the latter (three). How does optimizing for all
photopigments influence visual perception? We addressed this question by tuning our D65
mimics for the central 10° of visual field, akin to the standard observer commonly used to
evaluate color space (23). Macular pigment is substantial in this field so we considered its
prereceptoral filtering (**Supplementary Methods**)(23). We kept the wavelengths constant and
adjusted their ratios. We found that the optimal 2- and 3-wavelength mimics had ratios of 0.72:1
and 0.73:0.87:1, respectively. We tested the outputs of the constructed mimics and found that
the 2-wavelength version deviated by <8.9% for any photopigment and 5.75% across
photopigments (RRMSE), while the 3-wavelength version deviated by <1.29% for any
photopigment and 0.70% across photopigments. Next, we mapped the outputs of these mimics
to color space (**Fig. 4E** and **Supplementary Methods**)(36). The distances between D65 and
the 2- and 3-wavelength mimics were 25.69 and 2.46 (CIEDE2000). The latter is considered
barely perceivable. Collectively, these analyses indicate that three wavelengths mimic daylight
well for both image and non-image vision.
458

DISCUSSION

We designed and built artificial light sources that mimic photopigment stimulation profiles produced by natural light. These sources perform well with only two wavelengths and almost perfectly with three, and are readily adjusted to mimic variations of natural light and to compensate for prereceptoral filtering. We have also examined how thousands of natural light variations are represented by the photopigment complements possessed by humans, mice, and other mammals. We report that the representation for each species is captured in a 2-dimensional map that has orderly and informative substructure. Here, we place our work into context and discuss how it may be extended.

Mimicking natural light's stimulation of mammalian photopigments

By accounting for how an animal's complement of photopigments is stimulated by light, one can manipulate light to produce desired stimulation patterns. This is the basis of silent substitution, in which light's spectrum and intensity are varied to selectively modulate one or more photopigments while keeping the others in a constant state (27, 47). Silent substitution has provided insight into the roles of specific photopigments, even without manipulations like gene knockouts that are unfeasible in humans and other species.

We have used similar principles to develop artificial lights that mimic natural light's effects on the mammalian photopigments. Different spectra that produce the same percept are called metamers. Our mimics are metamers for perception as well as for non-perceptual actions of light. They differ in several respects from prior work in a similar vein (27): we mimicked natural light for species from mice to humans and considered prereceptoral filtering for all; incorporated melanopsin's three stable conformations; included many natural light variations; developed spectra whose wavelength bands are narrow and few in number; identified a simple means of correcting for prereceptoral filtering; omitted potentially harmful, ultraviolet wavelengths; and verified human natural light mimicry using psychophysical color space. The natural light mimics that we describe are accurate, safe, and flexible. We anticipate that they can be crafted for many species.

An advantage of mimicking natural light with only two or three wavelengths is the simplicity of correcting for prereceptoral filtering. A disadvantage is that a minimal mimic's reflected light is, in many contexts, unnatural. Minimal mimics of the kind we introduce here are most effective when delivered directly, such as for light therapy and research; when the intensity of direct illumination outstrips that of reflected light; or when light is altered little after reflection (as in a white room). Achieving natural reflectance spectra is likely to require illumination with broader spectra. One might find a middle ground depending on the application. For example, many LEDs that appear white and produce naturalistic reflectance spectra have one or more broad peaks. Broadening one peak of our human D65 mimic increases its accuracy (from an average deviation across photopigments of 5.75 to 3.1%; not shown).

Incorporating additional photopigments into mimics of natural light

The natural light mimics we present are comprehensive in that they include melanopsin's three states, but they omit photopigments that are not known to trigger electrical signals. The most notable is arguably neuropsin (Opn5), which has two stable states (λ_{max} of 380 and 470 nm), is found in the mammalian retina and brain, and is implicated in processes that include photoregulation of the retina clock and thermogenesis (48). We considered how minimal natural light mimics perform when adding neuropsin's two states, which brings the total number of

510 states to eight for mice and nine for humans. We found that 2- and 3-wavelength mimics of
511 G173 have average deviations of 11.11 and 0.69% for mice, and of 14.3 and 1.9% for humans,
512 respectively. Therefore, minimal mimics appear possible even for large numbers of
513 photopigment states.
514
515 **Representing light in terms of photopigment stimulation**
516
Expressing light in terms of its photopigment effects is a basic practice in vision research. For
example, light intensity is often given as photoactivated rhodopsin molecules (R*) per rod. One
obtains this value by using the rod's effective collecting area (in units of μm^2), which
incorporates light's spectrum (in photons μm^{-2} s^{-1} nm^{-1}) and prereceptoral filtering, as well as
rhodopsin's orientation, extinction coefficient, quantum efficiency of isomerization, and density
(19). The resulting units of R*/rod focuses on the level of biological activation rather than
properties of the light itself. This measure is possible because rods are stereotyped and have
been characterized extensively across species; it is also available for some cones. A measure
that requires fewer parameters is the α-opic irradiance, obtained by integrating the product of
the absorption spectrum of each photopigment (α) of a species (corrected for prereceptoral
filtering) and the power spectral density of the light (watts per unit area and unit time) at the
retina. For intuition, an additional transformation yields the intensity of standard daylight (D65)
that would give the same photopigment-weighted irradiance (α-opic equivalent daylight
illuminance; EDI)(11).
531
Our work extends these traditions. Like R*/rod, we use photons rather than watts to focus on
biological effects. Like α-opic irradiance, we consider several photopigment types across
species. We go further by incorporating melanopsin's three stable and spectrally distinct
conformations, which were defined relatively recently and have stark effects on melanopsin-
expressing cells (13-16). Like α-opic EDI, we examine natural light. However, rather than
examining an idealized daylight spectrum like D65, we analyze thousands of spectra that detail
the actual, ambient illumination from the sun. In this manner, we produced maps of
photopigment stimulation profiles across variations of natural light for mice, rats, cats, dogs,
rabbits, and humans. The profile map is likely to be useful. For example, by situating artificial
lights within it, one sees how naturalistically they influence perception and regulate physiology.
542
We expect that our approach can be brought to non-mammalian species, where interesting
complications are found. For instance, prereceptoral filtering may be complex; some birds and
reptiles have oil droplets of different colors that shape the light received by cones, and other
species situate their photopigments deep in the brain or beneath other tissues (48, 49). Some
species have many photopigment types—the mantis shrimp has a dozen in its compound eye,
and a fruit fly has eight in its compound eye and dorsal ocellus (50, 51). Some photopigments
are associated with molecules that broaden their spectra, and others change their absorption
spectra by switching their chromophore type (52, 53). Examining the photopigment stimulation
maps of these species would provide insight into their visual ecologies.
552
Future maps might add ambient solar illumination spectra from other geographical locations,
meteorological conditions, and seasons; include spectra of the reflected light that an organism
tends to experience; consider the daylight available to aquatic organisms; and broaden this
analysis to include other natural illumination such as the spectra of moonlight, starlight,
bioluminescence, fluorescence, and chemiluminescence (28). These maps would support
investigations of the diversity of photoreceptive mechanisms.
559

560   Photopigment stimulation maps reveal how different photic environments influence a given
561   species. It would be interesting and likely useful to determine how the maps of interacting
562   species overlap, and how those overlaps vary with natural conditions as well as with the type
563   and prevalence of artificial illumination.
564
565   **Closing remarks**
566
Newton found that passing white daylight through a prism yields a spectrum of colors. Helmholtz
discovered that combining complementary colors reproduces the perception of white light (54).
Rod and cone photopigments provided an explanation, but psychophysical experiments
suggested that this explanation may be incomplete (55). Behavioral and physiological
experiments also alluded to an additional photopigment, which is melanopsin (2). We provide
evidence that two wavelengths suffice to mimic the effect of natural light even when considering
its many variations and melanopsin's complexities. One advantage of such minimal mimics is
easy compensation for prereceptoral filtering, which allows it to be precisely matched across
varied conditions—for instance, to study animals and *ex vivo* tissues using the same
illumination, and to overcome age-related changes in the eye's optics and restore the
perception and physiological actions of light from earlier life.

MATERIALS AND METHODS

All experiments were conducted according to the protocols approved by the Institutional Animal Care and Use Committee at Boston Children's Hospital. All other experimental details are provided in Supplementary Information, Materials and Methods.

DATA, MATERIALS, AND SOFTWARE AVAILABILITY.

All study data are included in the article and/or Supplementary Information.

DATA SHARING PLAN

All data are available upon request from Philippe Morquette (philippe.morquette@childrens.harvard.edu).

AUTHOR CONTRIBUTIONS

Conceptualization: 2-wavelength mimics of natural light with compensation for arbitrary prereceptoral filtering, M.T.H.D; maps of photopigment stimulation profiles, P.M.; the remainder: P.M. and M.T.H.D. Experiments and analyses: P.M. Writing: P.M. and M.T.H.D.

DISCLOSURES

Philippe Morquette and Michael Tri H. Do declare that a patent application related to the work described in this manuscript has been filed through Boston Children's Hospital.

ACKNOWLEDGMENTS

Javier Hernández-Andrés kindly provided metadata of the Granada natural light dataset. Andreas Liu gave an example of wavelength optimization. Franklin Caval-Holme, Sophia Wienbar, and Navid Mousavi provided critical discussion. Grigory Loginov and Navid Mousavi assisted with construction of the diaphragm adaptor for spectrophotometry. Samuel Brill-Weil, Franklin Caval-Holme, Grigory Loginov, Nguyen-Minh Viet, Victoria Amstrup Vold, and Sophia Wienbar provided critiques of the manuscript. This research was funded by NIH grants EY023648, EY025555, EY032731, EY030628, and EY034089 (to M.T.H.D.). P.M. is a Warren Alpert Distinguished Scholar in Neuroscience, a Lefler Fellow, and a Tommy Fuss Fellow

REFERENCES

1. R. J. Lucas *et al.*, Measuring and using light in the melanopsin age. *Trends Neurosci* **37**, 1-9 (2014).
2. M. T. H. Do, Melanopsin and the Intrinsically Photosensitive Retinal Ganglion Cells: Biophysics to Behavior. *Neuron* **104**, 205-226 (2019).
3. S. P. D'Souza *et al.*, Retinal patterns and the cellular repertoire of neuropsin (Opn5) retinal ganglion cells. *J Comp Neurol* **530**, 1247-1262 (2022).
4. D. P. Windred *et al.*, Brighter nights and darker days predict higher mortality risk: A prospective analysis of personal light exposure in >88,000 individuals. *Proc Natl Acad Sci U S A* **121**, e2405924121 (2024).
5. J. F. Harmsen *et al.*, Natural daylight during office hours improves glucose control and whole-body substrate metabolism. *Cell Metab* 10.1016/j.cmet.2025.11.006 (2025).
6. L. Lazzerini Ospri, G. Prusky, S. Hattar, Mood, the Circadian System, and Melanopsin Retinal Ganglion Cells. *Annu Rev Neurosci* **40**, 539-556 (2017).
7. S. Sahar, P. Sassone-Corsi, Metabolism and cancer: the circadian clock connection. *Nature Reviews Cancer* **9**, 886-896 (2009).
8. D. L. Wescott, B. P. Hasler, J. E. Foust, K. A. Roecklein, Circadian realignment and depressed mood: A systematic review. *Sleep Med Rev* **79**, 102022 (2025).
9. Z. J. Voysey, R. A. Barker, A. S. Lazar, The Treatment of Sleep Dysfunction in Neurodegenerative Disorders. *Neurotherapeutics* **18**, 202-216 (2021).
10. K. Reinhard, K. Powell, M. Rizzi, Beyond night vision: the expanding role of rod photoreceptors in bright light. *Vision Res* **240**, 108744 (2025).
11. R. J. Lucas *et al.*, Recommendations for measuring and standardizing light for laboratory mammals to improve welfare and reproducibility in animal research. *PLoS Biol* **22**, e3002535 (2024).
12. CIE (2018) International Standard: CIE System for Metrology of Optical Radiation for iPRGC-Influenced Responses to Light.
13. T. Matsuyama, T. Yamashita, Y. Imamoto, Y. Shichida, Photochemical Properties of Mammalian Melanopsin. *Biochemistry* **51**, 5454-5462 (2012).
14. A. J. Emanuel, M. T. H. Do, Melanopsin tristability for sustained and broadband phototransduction. *Neuron* **85**, 1043-1055 (2015).
15. A. J. Emanuel, M. T. H. Do, The multistable melanopsins of mammals. *Frontiers in Ophthalmology* **3** (2023).
16. A. Liu *et al.*, Encoding of environmental illumination by primate melanopsin neurons. *Science* **379**, 376-381 (2023).
17. T. Eto, S. Higuchi, Review on age-related differences in non-visual effects of light: melatonin suppression, circadian phase shift and pupillary light reflex in children to older adults. *J Physiol Anthropol* **42**, 11 (2023).
18. F. van Oosterhout *et al.*, Ultraviolet light provides a major input to non-image-forming light detection in mice. *Curr Biol* **22**, 1397-1402 (2012).
19. D.-G. Luo, V. Kefalov, K.-W. Yau, "1.10 - Phototransduction in retinal rods and cones." in The Senses: A Comprehensive Reference*, A. I. Basbaum, Ed. (Elsevier/Academic Press, 2008), vol. 1-6.
20. V. I. Govardovskii, N. Fyhrquist, T. Reuter, D. G. Kuzmin, K. Donner, In search of the visual pigment template. *Visual Neuroscience* **17**, 509-528 (2000).
21. D. G. Luo, W. W. Yue, P. Ala-Laurila, K. W. Yau, Activation of visual pigments by light and heat. *Science* **332**, 1307-1312 (2011).
22. M. Alpern, A. B. Fulton, B. N. Baker, "Self-screening" of rhodopsin in rod outer segments. *Vision Res* **27**, 1459-1470 (1987).
23. CIE, *Fundamental chromaticity diagram with physiological axes - Part 1* (2006).

24.   K. Sakurai *et al.*, Physiological properties of rod photoreceptor cells in green-sensitive
cone pigment knock-in mice. *J Gen Physiol* **130**, 21-40 (2007).

25.   K. Tsutsui, H. Imai, Y. Shichida, Photoisomerization efficiency in UV-absorbing visual
pigments: protein-directed isomerization of an unprotonated retinal Schiff base.
Biochemistry **46**, 6437-6445 (2007).

26.   J. T. Henriksson, J. P. Bergmanson, J. E. Walsh, Ultraviolet radiation transmittance of the
mouse eye and its individual media components. *Exp Eye Res* **90**, 382-387 (2010).

27.   J. W. Mouland, F. Martial, A. Watson, R. J. Lucas, T. M. Brown, Cones Support
Alignment to an Inconsistent World by Suppressing Mouse Circadian Responses to the
Blue Colors Associated with Twilight. *Curr Biol* **29**, 4260-4267 e4264 (2019).

28.   S. Johnsen, *The Optics of Life: A Biologist's Guide to Light in Nature* (Princeton
University Press, Princeton, New Jersey, 2012).

29.   A. Stockman, A. T. Rider, Formulae for generating standard and individual human cone
spectral sensitivities. *Color Res Appl* **48**, 818-840 (2023).

30.   ASTM, Standard Tables for Reference Solar Spectral Irradiances: Direct Normal and
Hemispherical on 37° Tilted Surface. 10.1520/G0173-23 (2023).

31.   M. Spitschan, G. K. Aguirre, D. H. Brainard, A. M. Sweeney, Variation of outdoor
illumination as a function of solar elevation and light pollution. *Sci Rep* **6**, 26756 (2016).

32.   J. Hernandez-Andres, J. Romero, J. L. Nieves, R. L. Lee, Jr., Color and spectral analysis
of daylight in southern Europe. *J Opt Soc Am A Opt Image Sci Vis* **18**, 1325-1335
(2001).

33.   L. Walmsley *et al.*, Colour as a signal for entraining the mammalian circadian clock.
PLoS Biol **13**, e1002127 (2015).

34.   E. Solessio, G. A. Engbretson, Antagonistic chromatic mechanisms in photoreceptors of
the parietal eye of lizards. *Nature* **364**, 442-445 (1993).

35.   S. Johnsen *et al.*, Crepuscular and nocturnal illumination and its effects on color
perception by the nocturnal hawkmoth Deilephila elpenor. *The Journal of Experimental
Biology* **209**, 789-800 (2006).

36.   CIE (2018) Colorimetry, 4th Edition.

37.   R. T. Dauchy *et al.*, Effect of spectral transmittance through red-tinted rodent cages on
circadian metabolism and physiology in nude rats. *J Am Assoc Lab Anim Sci* **52**, 745-
755 (2013).

38.   J. C. Merriam *et al.*, An action spectrum for UV-B radiation and the rat lens. *Invest
Ophthalmol Vis Sci* **41**, 2642-2647 (2000).

39.   R. H. Douglas, G. Jeffery, The spectral transmission of ocular media suggests ultraviolet
sensitivity is widespread among mammals. *Proc Biol Sci* **281**, 20132995 (2014).

40.   T. G. Gorgels, D. van Norren, Spectral transmittance of the rat lens. *Vision Res* **32**,
1509-1512 (1992).

41.   R. J. McDowell, M. J. Gatt, S. Notash, R. J. Lucas, Spectral Tuning in Mammalian
Melanopsins. *Mol Biol Evol* **42** (2025).

42.   J. K. Bowmaker, H. J. Dartnall, Visual pigments of rods and cones in a human retina. *J
Physiol* **298**, 501-511 (1980).

43.   S. L. Merbs, J. Nathans, Absorption spectra of human cone pigments. *Nature* **356**, 433-
435 (1992).

44.   CIE (2019) CIE spectral luminous efficiency for photopic vision.

45.   T. W. Cronin, S. Johnsen, N. J. Marshall, E. J. Warrant, *Visual Ecology* (Princeton
University Press, 2014).

46.   IES, ANSI/IES RP-27-20: Recommended Practice: Photobiological Safety for Lighting
Systems. (2020).

47.   M. Spitschan *et al.*, The human visual cortex response to melanopsin-directed
715   stimulation is accompanied by a distinct perceptual experience. *Proc Natl Acad Sci U S
716   A* **114**, 12291-12296 (2017).

717   48.   H. Calligaro, O. Dkhissi-Benyahya, S. Panda, Ocular and extraocular roles of neuropsin
718   in vertebrates. *Trends Neurosci* **45**, 200-211 (2022).

719   49.   M. B. Toomey, J. C. Corbo, Evolution, Development and Function of Vertebrate Cone Oil
720   Droplets. *Front Neural Circuits* **11**, 97 (2017).

721   50.   M. P. Christenson, S. N. Mousavi, E. Oriol, S. L. Heath, R. Behnia, Exploiting colour
722   space geometry for visual stimulus design across animals. *Philos Trans R Soc Lond B
723   Biol Sci* **377**, 20210280 (2022).

724   51.   H. H. Thoen, M. J. How, T. H. Chiou, J. Marshall, A different form of color vision in mantis
725   shrimp. *Science* **343**, 411-413 (2014).

726   52.   T. Isayama *et al.*, An accessory chromophore in red vision. *Nature* **443**, 649 (2006).

727   53.   J. C. Corbo, Vitamin A(1)/A(2) chromophore exchange: Its role in spectral tuning and
728   visual plasticity. *Dev Biol* **475**, 145-155 (2021).

729   54.   H. v. Helmholtz, J. P. C. Southall, *Helmholtz's treatise on physiological optics* (The
730   Optical Society of America, Rochester, N.Y., 1924).

731   55.   M. V. Danilova, J. D. Mollon, Bongard and Smirnov on the tetrachromacy of extra-foveal
732   vision. *Vision Res* **195**, 107952 (2022).

733   56.   CIE (2019) CIE 1964 colour-matching functions , 10 degree observer.

734   57.   W. A. Rushton, Pigments and signals in colour vision. *J Physiol* **220**, 1P-P (1972).

735   58.   A. Fein, E. Z. Szuts, *Photoreceptors: Their Role in Vision*, IUPAB Biophysics Series
736   (Cambridge University Press, 1982), vol. 5.

737   59.   M. T. H. Do *et al.*, Photon capture and signalling by melanopsin retinal ganglion cells.
738   *Nature* **457**, 281-287 (2009).

739   60.   M. Alpern, E. N. Pugh, Jr., The density and photosensitivity of human rhodopsin in the
740   living retina. *J Physiol* **237**, 341-370 (1974).

741   61.   F. Naarendorp *et al.*, Dark light, rod saturation, and the absolute and incremental
742   sensitivity of mouse cone vision. *J Neurosci* **30**, 12495-12507 (2010).

743   62.   Y. V. Wang, M. Weick, J. B. Demb, Spectral and temporal sensitivity of cone-mediated
744   responses in mouse retinal ganglion cells. *J Neurosci* **31**, 7670-7681 (2011).

745   63.   A. L. Lyubarsky, B. Falsini, M. E. Pennesi, P. Valentini, E. N. Pugh, Jr., UV- and
746   midwave-sensitive cone-driven retinal responses of the mouse: a possible phenotype for
747   coexpression of cone photopigments. *J Neurosci* **19**, 442-455 (1999).

748   64.   A. J. Emanuel, K. Kapur, M. T. H. Do, Biophysical Variation within the M1 Type of
749   Ganglion Cell Photoreceptor. *Cell Reports* **21**, 14 (2017).

750   65.   E. S. Milner, M. T. H. Do, A Population Representation of Absolute Light Intensity in the
751   Mammalian Retina. *Cell* **171**, 11 (2017).

752   66.   C. Zhu, R. H. Byrd, P. Lu, J. Nocedal, Algorithm 778: L-Bfgs-B: Fortran Subroutines for
753   Large-Scale Bound-Constrained Optimization. *ACM Transactions on Mathematical
754   Software* **23**, 10 (1997).

755   67.   H. Strasburger, I. Rentschler, M. Juttner, Peripheral vision and pattern recognition: a
756   review. *J Vis* **11**, 13 (2011).

757   68.   IEC (2003) Multimedia systems and equipment - Colour measurement and management
758   - Part 2-1: Colour management - Default RGB colour space - sRGB.

759   69.   M. R. Luo, G. Cui, B. Rigg, The Development of the CIE 2000 Colour-Difference
760   Formula: CIEDE2000. *COLOR research and application* **26**, 11 (2000).

761   70.   M. Ester, H.-P. Kriegel, J. Sander, X. Xu, A density-based algorithm for discovering
762   clusters in large spatial databases with noise. *KDD-96 Proceedings*, 6 (1996).

763   71.   F. V. Pedregosa, Gaël; Gramfort, Alexandre; Michel, Vincent; Thirion, Bertrand; Grisel,
764   Olivier; Blondel, Mathieu; Prettenhofer, Peter; Weiss, Ron; Dubourg, Vincent;

VanderPlas, Jake; Passos, Alexandre; Cournapeau, David; Brucher, Matthieu; Perrot,
Matthieu; Duchesnay, Édouard, Scikit-Learn: Machine Learning in Python. *Journal of*
Machine Learning Research **12**, 5 (2011).
72. J. Sander, M. Ester, H.-P. Kriegel, X. Xu, Density-Based Clustering in Spatial Databases:
The Algorithm GDBSCAN and Its Applications. *Data Mining and Knowledge Discovery* **2**,
25 (1998).
73. E. Schubert, J. Sander, M. Ester, H.-P. Kriegel, X. Xu, DBSCAN Revisited, Revisited.
ACM Transactions on Database Systems **42**, 20 (2017).

Figure 1

Figure 1. 2-wavelength mimicry of natural light for a dichromatic mammal (mouse).
(*A*) Multiplying the spectrum of a daylight standard (ASTM G173) by the normalized
transmission spectrum of the mouse eye (see **Fig. S1**) and the normalized absorption spectrum
of each mouse photopigment yields the photon availability for each photopigment (or, in the
case of melanopsin, photopigment state). (*B*) Integrating each photon availability spectrum
gives the G173 photopigment stimulation profile ("profile"). Photopigments are ordered by peak
wavelength sensitivity, from shortest to longest. The three stable states of melanopsin (Opn4)
are given individually. (*C*) Varying two wavelengths (first and second λ) and their ratio (vertical
axis) reproduces the G173 profile with varying accuracy (the deviation is given as a relative root
mean squared error, RRMSE, across photopigments). The G173 profile is mimicked optimally
by wavelengths of 403.3 nm and 512.2 nm, given in a ratio of 1.26:1. The color code for
deviation is shown on the right. (*D*) The profiles of G173 (gray bars) and its 2-wavelength mimic
(black diamonds). Profiles are expressed in terms of their unit vectors to focus on spectrum
rather than intensity.

Figure 2

788
Figure 2. 2-wavelength mimicry of natural light's variations for a dichromatic mammal
(mouse). *(A)* Spectra of the G173 daylight standard (black) and 3,567 (3K) samples of natural
light (gray traces)(31, 32). Highlighted are spectra taken at the lowest (-13°, blue) and highest
(76.2°, red) solar elevations in the 3K set. *(B)* Photopigment stimulation profiles generated from
the blue and red spectra in '*A*' compared with that of G173. *(C)* The two wavelengths for optimal
G173 mimicry (403.3 and 512.2 nm), with their ratio adjusted (0.35-3.57), mimic most 3K
profiles (gray points) with <10% deviation. Points are spread along the x axis for visualization.
Shown are the interquartile range (IQR, box), median (line), mean (diamond), and last point
within 1.5×IQR. The -13° (blue) and 76.2° (red) solar elevations are highlighted. *(D)* Two
wavelengths, optimized for an average trajectory of daily spectral variations (402.6 and 510.8
nm) with ratio adjustment (black), mimic that trajectory with little deviation (RRMSE, gray).
Deviation is negligible at lower elevations where only rhodopsin requires consideration (gray
box); the stimulation of one photopigment can be set precisely just using intensity. *(E)*
Photopigment stimulation as a function of solar elevation, considering both the spectrum and
intensity of illumination. The gray box indicates nautical twilight, below which only rhodopsin
requires consideration.

Figure 3. Mapping photopigment stimulation profiles under natural and artificial lights for a dichromatic mammal (mouse). (*A*) The 3K set of natural spectra produces mouse photopigment stimulation profiles (dots) that are well-described by their first two principal components (>99.9% of the variance explained). Profiles are color-coded by solar elevation, showing their systematic distribution within this 3K map. This distribution is also evident in the trajectory of profiles during an average day (black dots and line). (*B*) Profiles of the G173 daylight standard (white circle) and its 2- and 3-wavelength mimics (black and green arrowheads, respectively), projected onto the 3K map. By changing the intensity ratio of the two G173 mimic wavelengths, different regions of solar elevation can be replicated (blue dots; note that the y axis accounts for only 1.69% of the variance). Using three wavelengths increases the precision of mimicry (pink dots). (*C*) Profiles of artificial light (red circles, closed for measurements and open for CIE standards) projected onto the 3K map. Also shown are G173 profiles with different prereceptoral filtering: adding a cage (yellow circle) and removing the eye's optics (blue circle). Inset: profiles of monochromatic lights on the periphery of the 3K profiles (gray dots inside the dashed box). For each arrow/arrowhead, the apex point indicates the position of the mimic.

Figure 4

822
Figure 4. Mimics and maps of natural light for trichromatic humans. (*A*) The map of 3K
natural light profiles for humans, shown with profiles of the G173 daylight standard (white circle),
2- and 3-wavelength mimics of G173 (black and green arrowheads), D65 standard (yellow
square), optimal 2- and 3-wavelength mimics of D65 (light blue and yellow arrows), constructed
2- and 3-wavelength mimics of D65 (blue and orange arrows), measured and standard artificial
lights (closed and open red circles), and monochromatic lights (inset, with the 3K profiles shown
as gray dots inside the dashed box; wavelengths shorter than 400 nm are scarcely transmitted
by the human eye). For each arrow/arrowhead, the apex point indicates the position of the
mimic. See **Table S3** for additional information. (*B*) A constructed mimic of the D65 daylight
spectrum, shown in its 3-wavelength configuration with the 12.7-cm Ganzfeld. Labeled are three
narrowband light sources (**1**, **3**, and **5**), bandpass filters (**2**, **4**, and **6**), beamsplitters (**7**, **8**), a lens
(**9**), a light guide (**10**), and a Ganzfeld diffusing sphere (**11**; aperture is outlined). (*C*) Measured

output spectra of the 2- and 3-wavelength mimic systems. (*D*) Photographs of the Ganzfeld
aperture showing the individual light sources and their combination, for the 2- and 3-wavelength
mimics. (*E*) Projection of D65 and its constructed 2- and 3-wavelength mimics (2λ D65 mimic
and 3λ D65 mimic in '*C*') onto the CIE chromaticity diagram (a map of human color
perception)(56)**.** (*F*) Comparison of the D65 photopigment stimulation profile with those of
measured artificial lights. Deviation is expressed as the relative root mean square error,
RRMSE, across photopigments. Also see **Fig. S8*F***.

SUPPLEMENTARY INFORMATION

Supplementary Methods

Estimating photopigment stimulation profiles in natural and artificial light

All spectra were expressed in units of photon flux density (photons μm^{-2} s^{-1} nm^{-1}). Each light spectrum was measured for this study, published previously, or is a standard. The light spectrum was multiplied by the normalized transmission spectrum of each prereceptoral filter considered. The product was multiplied with the spectral sensitivity of a photopigment, then integrated to give the total photons available to stimulate the photopigment (57). For brevity, this integral is referred to as the photopigment's "stimulation." The spectral sensitivity of a photopigment was given by its λ_{max} and the mathematical template (nomogram) of Govardovskii and colleagues (20), which includes the main α band and the secondary β band. The γ band was not considered because it is a non-specific absorption peak that is common to most proteins, not specified in existing nomograms, and excites the photopigment modestly; furthermore, photons at these wavelengths are sparse in natural light spectra (30-32, 35, 58).

Repeating this procedure for each relevant photopigment state produced a series of values. For example, for mice, six photopigment states were considered: Opn2, Opn1SW, Opn1MW, and Opn4's R, M, and E conformations. These photopigment stimulations were combined into a six-element vector, which was converted into a unit vector to compare spectra of different intensities. For brevity, this vector is referred to as the "photopigment stimulation profile" or simply "profile" of a natural or artificial light. Each stimulation profile is specific to one species' complement of photopigments under one light spectrum and one condition of prereceptoral filtering.

Photopigment self-screening and co-screening

Photopigment self-screening was accounted for in humans and mice, for which data are available. Each photopigment's absorption spectrum was adjusted using this equation (23):

$$F(\lambda) = 1 - 10^{-OD*a(\lambda)}$$

where a(λ) is the photopigment's Govardovskii nomogram (normalized) and OD is its axial optical density in the phototransducing outer segment (OS). The corrected spectrum was normalized by its peak. Self-screening was omitted for melanopsin because of its low expression density in ipRGCs (59).

For the human, OD values were 0.35 for Opn2 (60), 0.30 for Opn1SW, and 0.38 for Opn1MW and Opn1LW (23).

For the mouse, the OD of Opn2 was 0.456 (61). Accounting for Opn1SW and Opn1MW was more complex, as most cones co-express them in reciprocal gradients across the retina (62). Thus, in single cones, these photopigments screen themselves and one another to different degrees. A simplification was used in which individual cones were considered to express either Opn1SW or Opn1MW alone, each with an OD of 0.195, computed as 0.015 OD μm^{-1} × 13.0 μm (the OS length)(63).

To assess the fairness of this simplification, mutual screening was assessed. Opn1SW and Opn1MW were considered to be intermixed in the OS. The total absorbance of the OS was

893
894
$$A_{tot}(\lambda) = OD_S * a_S(\lambda) + OD_M * a_M(\lambda)$$
895
896 where OD_S and OD_M are the optical densities of Opn1SW and Opn1MW, respectively. aS(λ) and
897 aM(λ) are Govardovskii templates for their absorption spectra.
898
899 From the Beer–Lambert law, the total fraction of incident photons absorbed is
900
901
$$F_{tot}(\lambda) = 1 - 10^{-A_{tot}(\lambda)}$$
902 Each photopigment absorbed photons in proportion to its relative absorbance at each
903 wavelength. The mutually-screened photopigment absorption spectra are
904
905
$$F_S^{(mix)}(\lambda) = \left(\frac{OD_S * a_S(\lambda)}{\left(OD_S * a_S(\lambda) + OD_M * a_M(\lambda) \right)} \right) * \left(1 - 10^{-\left(OD_S*a_S(\lambda) + OD_M*a_M(\lambda) \right)} \right),$$
906
907
$$F_M^{(mix)}(\lambda) = \left(\frac{OD_M * a_M(\lambda)}{\left(OD_S * a_S(\lambda) + OD_M * a_M(\lambda) \right)} \right) * \left(1 - 10^{-\left(OD_S*a_S(\lambda) + OD_M*a_M(\lambda) \right)} \right)$$

908
The most extreme case is for Opn1MW within a cone containing 99% Opn1SW (62). Under the
2-wavelength G173 mimic, stimulation of this mutually-screened Opn1MW deviates from that of
Opn1MW (self-screening only) by -3.5%. For Opn1SW, its smallest proportion is within a cone
containing 65% Opn1MW (62). Under the 2-wavelength G173 mimic, stimulation of this
mutually-screened Opn1SW deviates from that of Opn1SW (self-screening only) by -10.9%. A
case was then considered where Opn1SW and Opn1MW are present in equal proportions.
Under the 2-wavelength G173 mimic, stimulation of these mutually-screened photopigments
deviate from those with self-screening by 8.5 and -1.8%, respectively, resulting in an overall
RRMSE across all photopigments of 3.4%. Mutual screening thus appears to modestly affect
the accuracy of natural light mimicry.
919
920 **Transmission spectrum of the mouse eye**
921
Male and female adult (>P56) C57BL/6J mice were deeply anesthetized with intraperitoneal
Avertin and their pupils dilated with tropicamide (0.5-1%). Their eyes were removed into Ames'
medium (US Biological or Millipore Sigma) that was supplemented with 1.8 g/L of $NaHCO_3$ and
equilibrated with 95% O_2/5% CO_2 to achieve a pH of 7.4. The posterior pole of the eye was cut
away near the ora serrata to open a light path through the vitreous, lens, aqueous, and cornea.
The eye was mounted in a silica cuvette (500-µl volume) filled with HEPES-buffered Ames'
medium at 35 °C (in mM, 140 NaCl, 3.1 KCl, 0.5 KH_2PO_4, 1.2 $CaCl_2$, 1.2 $MgSO_4$, 6 glucose, and
10 HEPES; pH adjusted to 7.4 with NaOH)(59, 64, 65). The transmission spectrum (300–700
nm) was obtained using a Shimadzu 1800 UV-Vis spectrophotometer. The background was
subtracted using the same cuvette with no eye. For each eye, 5-10 spectra were averaged,
ensuring no systematic change in the spectra over time. These average transmission spectra
were consistent across eyes when acquired immediately after dissection (<5 min; transmission
decreased, primarily over shorter wavelengths, over time). The ocular transmission spectrum
showed a longpass quality, declining by 18.1 ± 0.6% from 700 to 400 nm (11 eyes, mean ± SD).
Nevertheless, transmission of shorter wavelengths is greater than previously reported (26).
937
938 **Determining optimal wavelengths and intensities for natural light mimicry**

939
To design mimics of natural light's effects on the photopigments of a given species, a matrix was
constructed with each row corresponding to a photopigment and each column to a wavelength.
Thus, for the mouse 2-wavelength mimic, the matrix was 6×2. Wavelengths were selected
randomly within a defined range (e.g., 380-700 nm for the mouse). To determine the intensities
of each wavelength that would provide the closest match of photopigment stimulations to natural
light, an error function was defined:
946

$$E(I) = ||D - MI||_W^2$$

948
Where E is the error, D is the normalized (unit norm) photopigment stimulation profile under the
daylight spectrum in question (e.g., G173), and the mimic profile is the vector $\widehat{D} = MI$ generated
by a given wavelength combination and its intensities. M is the matrix that captures the
stimulation profiles at intensity 1 for a given set of wavelengths and I is a vector containing the
intensities of each wavelength. The weights (W) are the inverse square of each photopigment's
target stimulation profile value, ensuring that error contributions are relative and no
photopigment dominates the fit. The optimal intensity vector (I_{opt}) that minimizes E is derived
analytically as
957

$$I_{\text{opt}} = (M^T W M)^{-1} M^T W D$$

958
960   Using I_{opt}, the predicted profile $\widehat{D} = MI_{\text{opt}}$ is computed and then normalized (to unit norm) before
961   being compared to D using

$$\text{RRMSE} = 100 \sqrt{\frac{1}{P} \sum_{i=1}^{P} \left(\frac{\widehat{D}_i - D_i}{D_i} \right)^2}$$

963
964   where RRMSE is relative root mean squared error and P is the number of photopigments. Each
965   component was divided by D_i to express the error for each photopigment in relative (fractional)
966   units rather than absolute units. This makes RRMSE a component-wise relative RMSE that
967   matches the weighting $W = \text{diag}(1/D_i^2)$ used in $E(I)$.
968
This process was repeated 1000-10,000 times, discarding any I containing negative values, and
the ten wavelength combinations with the lowest RRMSE were selected for further optimization
using gradient descent (using the L-BFGS-B algorithm)(66), which further refined the
wavelengths (with intensities determined analytically as I_{opt} for each candidate wavelength set).
Each intensity is the integral of the light's spectrum (in units of photons). Mimic accuracy
depends on the ratio of intensities. The overall intensity (keeping the ratio constant) may be set
to the desired level.
976
977   **Determining practical ranges of wavelength and intensity for natural light mimicry**
978
Because real light sources have spectra of discrete widths, it is useful to identify ranges of
wavelengths that support adequate natural light mimicry. For a 2-wavelength mimic, a matrix
was constructed whose dimensions are the first stimulus wavelength (λ1), the second stimulus
wavelength (λ2), the ratio of their intensities (λ ratio = intensity λ1 / intensity λ2), and the
complement of photopigments (i.e., six additional dimensions in mouse). The wavelengths were
from 300–700 nm at 0.2-nm resolution, and the ratio ranged from 0.1 to 10 in increments of

0.01. Each value in the matrix is a photopigment's stimulation by two wavelengths at a given
ratio. Each profile (e.g., the vector containing stimulation of each photopigment, for a given
choice of $\lambda 1$, $\lambda 2$, and λ ratio) was compared to that of the reference spectrum (e.g., D65 for
humans) to generate an RRMSE. This operation was performed for all vectors, resulting in a 3-
dimensional matrix ($\lambda 1$, $\lambda 2$, and λ ratio) of RRMSEs. This matrix was used to generate a color
map, in which parameters producing an RRMSE <10% were highlighted. The 3-wavelength
mimic had two ratios, λ ratio$_1$ (intensity $\lambda 1$ / intensity $\lambda 3$) and λ ratio$_2$ (intensity $\lambda 2$ / intensity $\lambda 3$).

Construction of natural light mimics for humans

A Ganzfeld diffusing sphere was constructed from a hollow acrylic globe of 30.5-cm diameter
(Crown Plastics, 20012-SM-5N). A smaller version was built using 2 aluminum hemispheres of
12.7-cm diameter (Wagner, B4162). The larger globe had a 13-cm opening that served as the
viewing port, covering ~130-140° of visual angle (>80% of a human eye's visual field)(67). The
smaller Ganzfeld had a viewing port of 3.4 cm. For both spheres, a second, smaller port was
added to accommodate a liquid light guide (5-mm core, Thorlabs, LLG05-4H). This port was
positioned such that direct illumination from the light guide was outside the field of view; all
viewed light was internally reflected. For diffuse internal reflection that is even across
wavelengths, the globes were internally coated with layers of $BaSO_4$ until uniform and opaque
(Edmund Optics, 83-890).

2-wavelength D65 mimic. Light from LEDs (Solis-470C and Solis-565D) was conditioned with
bandpass filters (FBH460-10 and FL457.9-10 in series for the short-wavelength beam, and
FBH550-10 for the medium-wavelength beam), and combined using a beamsplitter
(DMSP490R; all parts from Thorlabs).

3-wavelength D65 mimic. Light from LEDs (Solis-470C, Solis-525C and Solis-565D) was
conditioned with bandpass filters (FBH455-10, FBH520-10 and FBH590-10, respectively) and
combined with beamsplitters (DMSP490R and DMSP550R; all parts from Thorlabs).

For both mimics, the combined beam was focused on the end of the liquid light guide using an
aspherical lens. Power was controlled by current level (DC20, Thorlabs). The output of each
Ganzfeld sphere was measured using a radiometer (PM100D and S130VC, Thorlabs) and
spectrometer (Flame-T-UV-VIS, Ocean Optics; coupled to a 400-μm diameter fiber). The
sensors were placed directly in front of the viewing apertures for measurement. The power of
each LED was adjusted until the desired spectrum was obtained.

Human perception of natural light mimics

To mimic natural light for both foveal and peripheral vision, we used the CIE 1964 color
matching functions (CMFs; 10° standard observer covering a large field around the fovea)(56).
Following standard procedure, the power spectral density of the mimic was multiplied by the
corresponding $\bar{x}$, $\bar{y}$, and $\bar{z}$ values from the CMFs, and the products integrated to obtain the raw
X, Y, and Z tristimulus values; the normalization constant was set such that Y=100. From these
normalized values, the chromaticity coordinates (x, y) were computed by dividing each of X, Y,
and Z by their sum (i.e., X+Y+Z)(36). The mimics and D65 coordinates were then projected onto
a CIE 1964 chromaticity diagram. For screen visualization, a regular grid of (x, y) coordinates
within the diagram was linearly transformed into sRGB color space, and gamma correction (68)
was applied to align the rendered colors with human perception.

To quantify the color differences between the D65 illuminant and its 2- and 3-wavelength
mimics, their XYZ tristimulus values were converted into the CIELAB color space, and the
CIEDE2000 formula was used to estimate their color difference (ΔE_{00}) under our Ganzfeld
viewing conditions (36, 69).
1039
1040   **Maps of photopigment stimulation profiles in natural light**
1041
Photopigment stimulation profiles (unit vectors) were computed for 2600 natural light spectra
measured at solar angles of -8-76° in Granada, Spain (only 56 spectra below 0°), and for 967
natural light spectra measured at solar angles of -13-24° (744 spectra below 0°) in rural
Pennsylvania, USA (31, 32). These were the most complete, readily available sets of natural
light that met inclusion criteria (range of at least 300–700 nm, resolution of ≤5 nm, and more
than one measurement at a given solar elevation in a given terrestrial location).
1048
1049   Principal component analysis (PCA) was performed on a matrix containing the 3,567 profiles
1050   (using the scikit-learn package in Python), where each row corresponded to a profile obtained
1051   for a spectrum and each column to one photopigment's stimulation (see above) in the set of
1052   photopigment present in a species. The first two principal components were used for
1053   visualization.
1054
1055   **Cluster analysis**
1056
DBSCAN, a density-based clustering algorithm, was used to assess the closeness of natural
and artificial photopigment stimulation profiles (70, 71). It was applied to all dimensions of these
profiles (e.g. 6 for the dichromatic mouse). DBSCAN is advantageous because it does not
require specifying the number of clusters and can detect clusters of arbitrary shape.
Furthermore, its ability to classify low-density points as noise makes it robust to outliers, which
is ideal when comparing diverse natural and artificial light sources. HDBSCAN was also
considered but found to produce clusters that were not robust to small changes in clustering
parameters or to subsampling of the data set.
1065
The parameters of DBSCAN are MinPts, the minimum number of neighbors that a point must
have to be the core of a cluster, and Eps, the maximum distance over which a point can be
considered a neighbor. These parameters were initialized using heuristics that are based, for
transparency and reproducibility, solely on the properties of the data (72, 73). MinPts was set to
2*dimensions. Eps was first set based on the distance to the k^{th} nearest neighbor (kNN), where
k = (2*dimension-1); for 6-dimensional mouse data, this parameter is 11. For each data point,
the kNN distance was then calculated. Data points were ordered by their kNN distance and
plotted as a k-distance graph. The value at the point of maximum curvature, where the k-
distance begins to increase steeply, was identified as the updated Eps. Points with kNN
distances smaller than the updated Eps were assigned to a cluster, and others considered noise
(72, 73). Most of the 3,567 natural stimulation profiles were within a single cluster, with <1.3%
excluded.
1078
Bootstrapping was used to ask if artificial profiles clustered with the natural profiles. In each
iteration, 80% of the 3K profiles were randomly sampled. An artificial profile was then added to
the resampled data and DBSCAN applied (maintaining the Eps and MinPts values determined
for the full data set). This process was repeated for 1,000 iterations, with a new resampling for
each. Resampling was performed without replacement to mitigate bias from varied local
densities of natural profiles. If an artificial profile clustered with the natural profiles in ≥95% of
iterations, it was classified as being similar.

Species	Photopigment	λ$_{max}$(nm)	References
Mouse	Opn1SW	358	Tsutsui et al., 2007
	Opn4 (E,R,M)	453.0, 471.0, 476.0	Emanuel and Do, 2015; Matsuyama et al., 2012
	Opn2	502	Sakurai et al., 2007
	Opn1MW	508	Sun et al., 1997
Rat	Opn1SW	358	Yokoyama et al., 1998
	Mel (E,R,M)	453.0, 471.0, 476.0	Emanuel and Do, 2015; Matsuyama et al., 2012
	Opn2	498	Bridges, 1959
	Opn1MW	509	Radlwimmer and Yokoyama, 1998
Cat	Opn1SW	450	Guenther & Zrenner, 1993
	Opn4 (E,R,M)	453.0, 471.0, 476.0	Emanuel and Do, 2015; Matsuyama et al., 2012
	Opn2	501	Guenther & Zrenner, 1993
	Opn1MW	553	Sun et al., 1997
Rabbit	Opn1SW	421	McDowell et al., 2024
	Opn4 (E,R,M)	453.0, 471.0, 476.0	Emanuel and Do, 2015; Matsuyama et al., 2012
	Opn2	502	Bridges, 1959
	Opn1MW	509	Radlwimmer and Yokoyama, 1998
Dog	Opn1SW	428	Jacobs, 1993; McDowell et al., 2024; Neitz et al., 1989
	Opn4 (E,R,M)	453.0, 471.0, 476.0	Emanuel and Do, 2015; Matsuyama et al., 2012
	Opn2	506	Parkes et al., 1982
	Opn1MW	554	Jacobs, 1993; McDowell et al., 2024; Neitz et al., 1989
Human	Opn1SW	426.3	Merbs and Nathans, 1992
	Opn4 (E,R,M)	453.51, 467.21, 476.02	Liu et al., 2023; Matsuyama et al., 2012
	Opn2	497.6	Bowmaker and Dartnall, 1980
	Opn1MW	529.7	Merbs and Nathans, 1992
	Opn1LW	552.4 and 556.7	Merbs and Nathans, 1992

1086 **Supplementary Table 1.** Wavelengths of peak absorption (λ$_{max}$) for the photopigments used in
1087 this study. Those of melanopsin states are from the mouse (13, 14), with the exception of the
1088 human R and E states, which are taken from the macaque (16).

	Deviation from G173 profiles (%, RRMSE)					
	Mouse	**Rat**	**Cat**	**Rabbit**	**Dog**	**Human**
2-wavelength G173 mimic	0.60	1.52	0.60	0.8	4.17	5.53
3-wavelength G173 mimic	$<10^{-6}$	$<10^{-6}$	$<10^{-6}$	$<10^{-6}$	$<10^{-6}$	0.13
Measured broadband lamp	**Mouse**	**Rat**	**Cat**	**Rabbit**	**Dog**	**Human**
Xenon	14.54	14.53	2.90	1.57	**3.26**	**3.52**
Halogen	40.87	40.12	38.90	28.80	42.11	40.58
Fluo1	38.74	38.65	14.18	14.92	18.33	13.94
Fluo2	39.55	37.09	46.60	25.14	44.69	49.56
Fluo3	44.93	43.05	47.40	30.43	44.70	48.55
White LED1	39.95	40.00	1.77	8.00	7.06	7.58
White LED2	39.98	40.00	5.60	9.29	10.12	**4.44**
White LED3	42.90	42.56	35.49	26.06	39.16	35.05
White LED4	40.40	40.33	15.57	11.69	18.2	13.50
CIE standard illuminants	**Mouse**	**Rat**	**Cat**	**Rabbit**	**Dog**	**Human**
A (Tungsten-filament)	32.30	31.49	33.36	24.22	36.12	36.12
LED-B1 (Phosphor-type LED)	45.63	45.05	41.79	28.07	44.50	41.23
LED-B2 (Phosphor-type LED)	44.36	43.89	35.92	24.56	38.89	35.38
LED-B3 (Phosphor-type LED)	41.69	41.50	19.03	13.30	21.68	17.75
LED-B4 (Phosphor-type LED)	40.99	40.86	11.68	6.83	13.43	10.12
LED-B5 (Phosphor-type LED)	40.60	40.58	2.48	4.15	4.85	8.45
LED-BH1 (Hybrid-type LED)	45.92	45.39	40.55	25.54	42.01	40.72
LED-RGB1 (RGB-type LED)	46.74	46.69	36.77	36.47	41.90	37.04
LED-V1 (Violet-pumped phosphor-type LED)	38.37	37.89	29.08	13.25	29.63	36.84
LED-V2 (Violet-pumped phosphor-type LED)	36.47	36.18	10.38	5.03	11.41	15.12
HP1 (Stand. high pressure sodium)	49.66	47.79	70.48	36.94	70.48	66.73
HP2 (Color-enhanced high-pressure sodium)	45.68	44.41	48.66	30.57	50.16	48.72
HP3 (High-pressure metal halide)	37.91	36.89	32.69	16.87	33.18	36.73
HP4 (High-pressure metal halide)	30.85	29.47	17.35	10.15	16.88	23.44
HP5 (High-pressure metal halide)	35.97	35.44	13.78	5.52	14.51	16.85
FL1 (Standard fluorescent)	33.55	32.89	1.93	3.75	**1.92**	**3.91**
FL2 (Standard fluorescent)	34.63	33.70	18.27	6.43	18.15	21.51
FL3 (Standard fluorescent)	36.56	35.30	31.22	11.40	30.70	35.08
FL4 (Standard fluorescent)	38.90	37.31	42.16	16.12	41.13	45.41
FL5 (Standard fluorescent)	33.46	32.77	3.09	3.17	**3.07**	**4.09**
FL6 (Standard fluorescent)	35.50	34.46	22.71	8.53	22.53	26.15
FL7 (Broadband fluorescent)	32.42	31.66	1.47	3.92	**1.72**	**4.52**
FL8 (Broadband fluorescent)	35.97	35.51	7.99	6.22	9.84	7.80
FL9 (Broadband fluorescent)	36.65	36.10	15.39	8.10	16.66	17.20
FL10 (Narrowband fluorescent)	37.58	37.08	11.00	4.79	11.54	11.85
FL11 (Narrowband fluorescent)	38.40	37.73	20.00	9.08	20.51	22.29
FL12 (Narrowband fluorescent)	40.84	39.74	36.77	18.67	37.00	39.86
FL3.1 (Stand. halophosphate fluorescent)	39.01	37.39	42.76	17.18	41.43	46.54
FL3.2 (Stand. halophosphate fluorescent)	34.53	33.58	19.21	7.51	18.92	23.20
FL3.3 (Stand. halophosphate fluorescent)	33.32	32.61	3.44	5.05	**3.59**	**4.96**
FL3.4 (DeLuxe type fluorescent)	40.13	39.07	37.68	18.42	37.45	41.50
FL3.5 (DeLuxe type fluorescent)	36.70	36.22	13.61	7.88	15.25	15.17
FL3.6 (DeLuxe type fluorescent)	36.49	36.07	8.50	6.91	10.63	8.41
FL3.7 (Three-band fluorescent)	41.05	39.96	37.51	18.33	37.42	41.00
FL3.8 (Three-band fluorescent)	38.67	37.95	21.70	10.11	22.02	24.44
FL3.9 (Three-band fluorescent)	37.99	37.52	12.24	5.43	12.43	13.89
FL3.10 (Three-band fluorescent)	38.82	38.60	8.86	5.32	10.43	8.63
FL3.11 (Three-band fluorescent)	37.68	37.32	6.25	4.32	6.33	8.02
FL3.12 (Multi-band fluorescent)	40.09	39.49	31.90	18.88	33.57	35.53
FL3.13 (Multi-band fluorescent)	37.88	37.56	14.82	10.67	17.49	16.65
FL3.14 (Multi-band fluorescent)	37.86	37.64	7.57	10.95	12.02	6.90
FL3.15 (D65 simulator)	25.26	23.76	2.52	5.13	**4.86**	**2.73**

1089

Supplementary Table 2. Comparison of photopigment stimulation profiles (%, RRMSE)
between G173, 2- and 3-wavelength G173 mimics, measured artificial light sources, and CIE
artificial light standards (36). A range of 300-700 nm was considered. CIE standards for

fluorescent lamps do not extend below 380 nm, so the range from 300-380 nm was set to 0 for
these calculations. This has little effect for humans, whose eyes scarcely transmit these
wavelengths, but may skew the RRMSE for other species.

	mouse			rat			cat			rabbit			dog			human	
	in	cage	ex	in	cage	ex	in	cage	ex	in	cage	ex	in	cage	ex	in	ex
G173	✓	x	✓	✓	x	x	✓	x	✓	✓	x	x	✓	x	✓	✓	x
2λ G173 mimic[1]	✓	✓	✓	x	x	x	✓	✓	✓	x	x	x	x	x	x	x	x
3λ G173 mimic[1]	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Xenon	✓	x	x	✓	x	x	✓	x	✓	✓	x	x	✓	x	✓	✓	x
Halogen	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
Fluo1	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	✓
Fluo2	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
Fluo3	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
White LED1	x	x	x	x	x	x	✓	x	✓	x	x	x	x	x	x	x	x
White LED2	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
White LED3	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
White LED4	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
mono 405	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
mono 480	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
mono 500	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
CIE-A	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
CIE-LED-B1	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
CIE-LED-B2	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
CIE-LED-B3	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
CIE-LED-B4	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
CIE-LED-B5	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
CIE-LED-BH1	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
CIE-LED-RGB1	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
CIE-LED-V1	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
CIE-LED-V2	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
CIE-HP1	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
CIE-HP2	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
CIE-HP3	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
CIE-HP4	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
CIE-HP5	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
CIE-FL1	x	x	x	x	x	x	x	x	x	x	x	x	✓	x	x	x	x
CIE-FL2	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
CIE-FL3	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
CIE-FL4	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
CIE-FL5	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
CIE-FL6	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
CIE-FL7	x	x	x	x	x	x	✓	x	✓	✓	x	x	✓	x	✓	✓	x
CIE-FL8	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
CIE-FL9	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
CIE-FL10	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
CIE-FL11	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
CIE-FL12	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
CIE-FL3.1	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
CIE-FL3.2	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
CIE-FL3.3	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
CIE-FL3.4	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
CIE-FL3.5	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
CIE-FL3.6	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
CIE-FL3.7	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
CIE-FL3.8	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
CIE-FL3.9	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
CIE-FL3.10	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
CIE-FL3.11	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
CIE-FL3.12	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
CIE-FL3.13	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
CIE-FL3.14	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	✓	✓
CIE-FL3.15	x	x	x	x	x	x	x	x	✓	x	x	✓	x	x	✓	✓	x

1096

Supplementary Table 3. Most photopigment stimulation profiles for artificial light (x) do not
cluster with the profiles for natural light (3K set). Some do (✓) but rarely for all conditions (*in*
vivo, cage, and *ex vivo*). Profiles for the 2-wavelength (2λ) G173 mimics cluster with natural
profiles for all conditions in mouse and cat. The 3λ G173 mimics cluster with natural profiles for
all conditions and species. [1]Note that the mimics have wavelengths and wavelength ratios that

are specific to each species and condition; for a given species, the wavelengths were fixed and
the ratios changed according to condition.

Supplementary Figure 1

1104

Supplementary Figure 1. Transmission spectrum of the mouse eye. (*A*) Schematic of the
optical path, inside a spectrophotometer, for measuring the transmission spectrum of the
anterior chamber. The anterior chamber was mounted in a custom cuvette filled with
oxygenated media, and all transmitted light was captured by the detector. (*B*) Illustration of the
anterior chamber's mounting within the cuvette. (*C*) Raw transmission spectra from 11 adult,
wild-type eyes (from 7 males and 4 females), normalized at 700 nm.

Supplementary Figure 2

1111
Supplementary Figure 2. Photopigment stimulation profiles of artificial lights. (*A*) Example
spectra measured from artificial lights. (*B*) Photopigment stimulation profiles from the artificial
lights in '*A*' (red x symbols) and the G173 standard daylight (gray bars). (*C*) Compared to G173,
measured artificial lights produce higher (+) or lower (-) stimulation of individual photopigments.
(*D*) Comparison of the G173 profile with those of measured artificial lights (deviation is
expressed as the relative root mean square error, RRMSE, across photopigments). The 2- and
3-wavelength G173 mimics outperform all artificial lights, including broadband and
monochromatic ('mono') sources.

Supplementary Figure 3

1120
Supplementary Figure 3. 2-wavelength mimicry of natural light for dichromatic mammals
(rat, cat, rabbit, and dog). Shown are mimics of the G173 photopigment stimulation profile for
rat (*A*), cat (*B*), rabbit (*C*), and dog (*D*). *Left panels:* Effect of varying the two wavelengths on
mimic accuracy (their ratios are fixed to optimal values). *Right panels:* The photopigment
stimulation profiles of G173 (gray bars) compared to those of the optimal 2-wavelength mimics
(black diamonds).

1127
1128 **Supplementary Figure 4.** Comparing photopigment stimulations for the G173 daylight standard
1129 and example artificial lights for rat (*A*), cat (*B*), rabbit (*C*) and dog (*D*). Artificial lights produced
1130 more (+) or less (-) stimulation than G173.

Theoretical 2-wavelength mimic

A Mouse

Min: 0.53%
Median: 2.04%
Mean: 2.32%
Max: 9.70%
Std: 1.13%

Deviation (%, RRMSE)

λ1:402.5 nm, λ2:508.9 nm
λ ratio (First/Second): 0.32-3.27

B Rat

Min: 1.25%
Median: 2.28%
Mean: 2.49%
Max: 9.56%
Std: 0.90%

λ1:403.3 nm, λ2:507.7 nm
λ ratio (First/Second): 0.32-2.94

C Cat

Min: 0.52%
Median: 1.00%
Mean: 1.08%
Max: 4.45%
Std: 0.46%

λ1:467.0 nm, λ2:552.2 nm
λ ratio (First/Second): 0.42-2.38

D Rabbit

Min: 0.69%
Median: 0.99%
Mean: 1.02%
Max: 4.06%
Std: 0.26%

Deviation (%, RRMSE)

λ1:447.8 nm, λ2:521.0 nm
λ ratio (First/Second): 0.39-1.95

E Dog

Min: 3.52%
Median: 4.13%
Mean: 4.10%
Max: 9.62%
Std: 0.40%

λ1:455.3 nm, λ2:544.7 nm
λ ratio (First/Second): 0.35-2.04

F Human

Min: 4.37%
Median: 5.20%
Mean: 5.19%
Max: 9.64%
Std: 0.47%

λ1:468.7 nm, λ2:554.8 nm
λ ratio (First/Second): 0.29-1.64

1131
Supplementary Figure 5. 2-wavelength mimicry of natural light's variations for
dichromatic and trichromatic mammals. Two optimal wavelengths, with their intensity ratio
adjusted (wavelengths and ratio range are given below each plot), mimic most 3K daylight
profiles (gray points) with <10% deviation. Points are spread along the x axis for visualization.
Shown are the interquartile range (IQR, box), median (line), mean (diamond), and last point
within 1.5×IQR. Analyses are shown for mouse (*A*), rat (*B*), cat (*C*), rabbit (*D*), dog (*E*), and
human (*F*).

Supplementary Figure 6

Theoretical 3-wavelength mimic

A Mouse

Min: 0.05%
Median: 0.22%
Mean: 0.21%
Max: 0.78%
Std: 0.07%

λ1:393.0 nm, λ2:470.8 nm, λ3:535.4 nm
λ ratio₁ (First/Third): 0.15-2.80
λ ratio₂ (Second/Third): 0.57-2.69

B Rat

Min: 0.05%
Median: 0.21%
Mean: 0.20%
Max: 0.88%
Std: 0.10%

λ1:395.0 nm, λ2:474.4 nm, λ3:540.4 nm
λ ratio₁ (First/Third):0.14-2.82
λ ratio₂ (Second/Third): 0.58-2.72

C Cat

Min: 0.00%
Median: 0.04%
Mean: 0.05%
Max: 0.41%
Std: 0.04%

λ1:444.3 nm, λ2:507.6 nm, λ3:570.4 nm
λ ratio₁ (First/Third):0.33-3.25
λ ratio₂ (Second/Third): 0.56-1.87

D Rabbit

Min: 0.01%
Median: 0.07%
Mean: 0.07%
Max: 0.31%
Std: 0.04%

λ1:436.1 nm, λ2:492.1 nm, λ3:548.9 nm
λ ratio₁ (First/Third): 0.30-2.56
λ ratio₂ (Second/Third): 0.63-1.80

E Dog

Min: 0.03%
Median: 0.18%
Mean: 0.20%
Max: 1.30%
Std: 0.12%

λ1:439.8 nm, λ2:502.7 nm, λ3:567.5 nm
λ ratio₁ (First/Third): 0.28-3.01
λ ratio₂ (Second/Third): 0.55-1.92

F Human

Min: 0.12%
Median: 0.33%
Mean: 0.36%
Max: 1.83%
Std: 0.16%

λ1:460.2 nm, λ2:520.7 nm, λ3:591.0 nm
λ ratio₁ (First/Third): 0.21-2.27
λ ratio₂ (Second/Third): 0.55-1.61

1139
1140
1141 **Supplementary Figure 6. Three-wavelength mimicry of natural light's variations for**
1142 **dichromatic and trichromatic mammals.** As for **Fig. S5**, but for 3-wavelength mimics.

Supplementary Figure 7

1143
Supplementary Figure 7. Mapping photopigment stimulation profiles under natural and
artificial lights for dichromatic mammals. *Left:* Maps of photopigment stimulation profiles
(from the 3K set of natural spectra), with monochromatic lights projected (380–700 nm). *Right:*
The boxed region on the left, expanded to show profiles of the 3K spectra (gray dots), the G173
standard (white circle), artificial lights that fall nearby (measured, closed red circles; CIE
standards, open circles); and 2- and 3-wavelength mimics (black and green arrowheads). These
analyses are shown for rat (*A* and *B*), cat (*C* and *D*), rabbit (*E* and *F*), and dog (*G* and H).

Supplementary Figure 8

1151
Supplementary Figure 8. Mimicking natural light and assessing artificial light for
trichromatic humans. (*A*) Effect of varying wavelengths on the accuracy of a 2-wavelength
G173 mimic (their ratios are fixed at optimal values). (*B*) Effect of varying ratios on the accuracy
of a 3-wavelength G173 mimic (the three wavelengths are fixed at optimal values). (*C*) The
photopigment stimulation profile of G173 profile (gray bars) compared to those of the optimal 2-
and 3-wavelength mimics (black and cyan diamonds). (*D*) For all tested species, mimic
performance peaks with three wavelengths. (*E*) Artificial lights stimulate human photopigments
more (+) or less (–) than the G173 daylight spectrum. (*F*) Deviation of photopigment stimulation
profiles of artificial lights from that of G173. The 3-wavelength mimic is more accurate than any
tested, artificial spectrum. The 2-wavelength mimic has higher deviation than xenon or one type
of white LED, but is preferable to these broadband sources when compensating for different
degrees of prereceptoral filtering.