Co-option of Lysosomal Machinery for Biomineralization

Maitri Rangarajan-Paul,[1,*] Scott A. Nichols,[2] Martin Tresguerres[1,*]
[1] Scripps Institution of Oceanography, University of California San Diego, 9500 Gilman Drive, La Jolla, CA 92093,
USA
[2] University of Denver, Department of Biological Sciences, 2101 E. Wesley Ave SGM 203, Denver, CO 80210-5210
*Correspondence: mapaul@ucsd.edu (M.R.P.), mtresguerres@ucsd.edu (M.T.)

Summary

Biomineralization first emerged ~600 million years ago, giving rise to novel strategies for
buoyancy, locomotion, and defense.[1] Sponges are the only metazoans that build a silica-based
skeleton,[2] and the cellular and molecular mechanisms underlying their silicification process are
much less understood than in diatoms and land plants, the other two major biosilicifiers. Here,
we show that the cells specialized for sponge biosilicification, termed sclerocytes, highly
express lysosomal-associated genes, including the V-H$^+$-ATPase (VHA) proton pump,
oculocutaneous albinism type 2 (OCA2) anion channels, cathepsin-L proteins, silicateins,
TMEM55A, TMEM192, TMEM199, and multiple other proteins involved with lysosomal
maintenance and degradation. Further, we demonstrate that VHA protein localizes to the
sclerocyte silica deposition vesicle (SDV) and that VHA-dependent SDV acidification is essential
for silica spicule formation. This function mirrors the localization and role of VHA in diatom
biosilicification,[3] while in land plants, acidification occurs extracellularly in the xylem and is
driven by a different proton pump, the plasma membrane P-type H$^+$-ATPase.[4,5] These results
indicate that vesicle acidification is a shared mechanism for silica handling in sponges, diatoms,
and plants. We also found that genes homologous to the plant silicon efflux transporter Lsi2,[6]
which belong to the same gene family as OCA2,[7,8] are upregulated in sclerocytes. Using
Hybridization Chain Reaction Fluorescence In Situ Hybridization (HCR-FISH), we found that
sponge Lsi2 genes are expressed in sclerocytes, corroborating prior phylogenetic and
transcriptomic evidence that Lsi2 homologs are involved in sponge and diatom silicic acid
transport.[7–10] Taking into account previous work showing that VHA, cathepsins, and other
lysosomal-associated genes have been associated with calcification across a remarkable range
of eukaryotes including coccolitophores,[11] foraminifera,[12] sea urchins,[13] bryozoans,[14]
crustaceans,[15] mollusks,[16] and mammals,[17] we posit that the various biomineralization
mechanisms observed throughout the tree of life arose convergently through the co-option of
an ancestral lysosomal gene toolkit.

Results and Discussion
Upregulation of lysosomal genes in sclerocytes
Analysis of single-cell RNA sequencing (RNASeq) data of demosponge *Spongilla*
lacustris[18] revealed that mRNA transcripts of genes involved in lysosomal machinery were

upregulated (log2fold change > 1) in sclerocytes (Figure 1); including most subunits of VHA, an

ATP-driven proton pump essential for lysosomal acidification,[19] OCA2, also known as the pink-

eyed dilution protein or P protein,[20] which localizes to lysosome related organelles called

melanosomes in vertebrate pigment-producing cells,[21,22] lysosomal cysteine protease

cathepsin-L proteins, cathepsin L-like proteins called silicateins that have been found to

involved in spiculogensis,[23] TMEM55A/B and TMEM192, proteins involved in lysosomal

autophagy, phagocytosis, and lysosome repair,[24–26] TMEM199, a VHA assembly factor,[27] SNX8, a

protein that enables lysosome reformation,[28]and CLIP4 and RNF11, proteins that mediate

processes leading to lysosomal degradation.[29,30] VHA is highly conserved across all eukaryotes.[31]

In addition to acidifying lysosomes, endosomes and phagosomes, VHA plays a critical role in

diatoms silica cell wall formation by acidifying the silica deposition vesicle (SDV)[3]. Further, we

found that two of the upregulated sponge genes annotated as OCA2 bore greater structural

resemblance to plant silicon efflux transporter Lsi2 (Supplementary Figure 1A) and are thus

referred to as sponge Lsi2 genes. Due to the links of VHA and Lsi2, to diatom and plant

biosilicification, we specifically explored their roles in sponge biosilicification.

<u>VHA is abundant in sclerocytes and is essential for skeleton formation</u>

In recent decades, VHA has been shown to play a wide range of physiological roles in

marine organsims.[3,32–39] Notably, VHA is essential for biosilicification in diatoms, allowing for

silicic acid accumulation in the SDV.[3] The protein has been linked to biomineralization in

calcifying organisms as well. In foraminifera, VHA maintains the proton flux responsible for

calcification.[12] The gene has been shown to be upregulated in calcifying strains of

coccolithophores.[11] A transcriptomic study on bryozoans identified VHA as a candidate

biomineralization protein as it was upregulated in younger zooids, which are at the distal

growth margin of the colony, compared to older zooids.[14] In sea urchins, VHA has been shown

to play an essential role in skeleton re-calcification.[13] The protein also contributes to the

mineralization-demineralization cycle of terrestrial isopods.[15]

Our analysis of published single-cell RNASeq data from *S. lacustris*[18] found that all

subunits of VHA were upregulated in sclerocytes compared to other cell types, with the

exception of integral domain V0 subunit e which was more highly expressed in

myopeptidocytes (Supplementary Figure 2). However, this subunit does not seem to be

essential for proton pump activity and may instead play a role in endomembrane specific VHA-

assembly or targetting.[40] Some subunit isoforms such as V0 subunit a gene 3 and V0 subunit c

gene 2 were specifically upregulated in sclerocytes compared to other cell types and compared

to other isoforms suggesting that these isoforms may only localize to sclerocyte-specific

organelles like the SDV. Immunostaining with specific antibodies against VHA subunit B (VHA$_B$)

revealed a striking VHA localization within sclerocytes and around spicules in *S. lacustris* as well

as in *Ephydatia muelleri* (Figure 2). In both cases, the VHA protein appeared to be surrounding

the SDV of sclerocytes, as well as within the cytoplasm, potentially on vesicles (Supplementary

Figure 3). Airyscan 'super-resolution' imaging showed VHA surrounding spicules fluorescently

labelled with PDMPO [2-(4-pyridyl)-5-((4-(2-dimethylaminoethylaminocarbamoyl)methoxy)-

phenyl)oxazole], an acidotropic dye that is selectively incorporated and co-deposited with Si

into the newly polymerized silica (Figure 3A).[41]

To test the functional role of VHA in spiculogenesis, sponges were incubated with the

specific pharmacological inhibitor of VHA, concanamycin (concA; 100 nM),[42,43] for 24 hours

along with PDMPO or with vehicle DMSO as a control (Figure 3B). VHA inhibition significantly

decreased the number of spicules per mm^2 of sponge tissue by 41.6% $\pm$ 10.6% in *E. muelleri* and

77.5% $\pm$ 7.1% in *S. lacustris*. The proportion of spicules labeled with PDMPO decreased by

85.8% $\pm$ 4.7% in *E. muelleri* and 98.3%$\pm$ 1.7% in *S. lacustris* indicating that spiculogenesis is

stunted by VHA inhibition (Figure 3C-D). Given that *S. lacustris* and *E. muelleri* are

phylogenetically distinct sponges with divergent life history, these observations suggest that

VHA-dependent silicification is a widespread, conserved mechanism in sponges (Figure 2, 3).

These results show that, as reported in diatoms[3], VHA-dependent SDV acidification is

critical for sponge biosilicification. By acidifying the SDV in both diatoms and sclerocytes, VHA

drives silicic acid accumulation and prevents its rapid polymerization into silica.[44,45] Given prior

evidence that VHA plays a role in calcification across a wide range of phyla, VHA appears to

have been co-opted independently by multiple lineages for biomineralization. Interestingly,

plants also use a proton pump to acidify the xylem sap where silicic acid is stored prior to

hardening. The xylem is acidified by the plant and fungi-specific[46] plasma membrane P-type H$^+$-

ATPase.[4,5] The co-option of proton pumps for maintenance of accumulated silicic acid appears

to be a shared feature for biosilicification over vastly diverged lineages.

Lsi2 and OCA2 are highly expressed in sclerocytes

Lsi2 and OCA2 belong to a family of transmembrane transporters with similarity to

SLC13. [7,8,47–49] Lsi2 is a silicic acid/H$^+$ antiporter that mediates silicic acid efflux into the plant

xylem.[6] OCA2 is a protein that localizes to melanosomes, lysosome related organelles, in

pigment-producing cells called melanocytes[21] which specifically occur in vertebrates.[22] These

proteins play a role in mediating chloride-selective anion conductance, providing charge

compensation and increasing proton-pumping efficiency of VHA.[50,51] A phylogenetic analysis of

the evolutionary history of Si transport reported that Lsi2 has homologs across a broad range of

eukaryotes which resolve into two main groups: one containing vertebrate OCA2-like

sequences, and the other containing the land plant Lsi2s together with Lsi2-like genes from

diverse prokaryotic and eukaryotic organisms.[7] This Lsi2-like clade was dominated by homologs

from silicifying organisms such as land plants, sponges, loricate choanoflagellates, haptophytes,

chrysophytes/synurophytes, radiolarians, and diatoms indicating that Lsi2-like genes play a role

in biosilicification. In alignment with this, Lsi2-like genes have been found to be upregulated in

marine hexactinellid sponge *Vazella pourtalesii* upon exposure to dissolved silicon.[8] In another
marine hexactinellid sponge *Aphrocallistes vastus*, an analysis of differentially expressed genes
between the osculum, the vertical growth zone where new skeletal growth is concentrated,[52]
and the main body, reported that Lsi2 was upregulated with other mineralization-related
genes.[9] There is also some transcriptomic evidence that an Lsi2-like gene may be involved in
silicic acid transport in diatoms.[10]

Our analysis of published single-cell RNASeq data from *S. lacustris*[18] showed that Lsi2
genes were upregulated in sclerocytes (Figure 4A). Alignment of the protein sequences of the
Lsi2 genes in *S. lacustris* and their equivalent homologs in *E. muelleri* against rice *Oryza sativa*
Lsi2 (*Os*Lsi2) showed that transmembrane regions are conserved (Supplementary Figure 1B).
OCA2 genes were also upregulated in sclerocytes but OCA2_1 was expressed in a smaller
percentage of sclerocyte cells than the Lsi2 genes and OCA2_2 was expressed in other cell types
as well indicating it may play a more generic housekeeping role in sponge lysosomes. Sponge
OCA2 genes may play a role similar to their function in vertebrate melanocytes of providing
counter-ion balance and improving efficiency of the VHA acidification mechanism. Notably, we
found that Lsi2 gene expression was much higher in sclerocytes than the previously described
sponge silicic acid uptake protein the $\{Na^+/HCO_3^-[Si(OH)_4]\}$ co-transporter or NBCSA,[53] (Figure
4A) which matches recent claims that NBCSA is not critical for silicic acid transport in all
sponges.[8] Confirming the single-cell RNASeq data, HCR-FISH probes designed against the Lsi2
genes in *S. lacustris* and *E. muelleri* distinctly labeled sclerocytes (Figure 4B). Interestingly, the
two Lsi2 homologs showed heterogeneous expression in sclerocytes: i.e. Lsi2_1 signal was
stronger in some sclerocytes, while Lsi2_2 signal was stronger in others; overall, Lsi2_2 was
more ubiquitously expressed around spicules than Lsi2_1 (Supplementary Figure 4). As we only
visualized mRNA, we could not definitively conclude whether the Lsi2 proteins localize to the
SDV membrane or the plasma membrane. We attempted to design antibodies against these
proteins but were unsuccessful. However, we speculate that they would localize to the SDV
membrane as the Lsi2 silicic acid/H^+ antiporter activity relies on a proton gradient and VHA
would only be able to pump H^+ ions from the cytosol to an intracellular compartment.[54]

These findings strongly suggest that the Lsi2 genes are involved in sponge silicic acid
uptake. The dependance of Lsi2 on an acidification mechanism and the critical role of OCA2 in a
lysosome-related organelle, point to lysosomal origins of a gene family with clear association to
sponge spiculogenesis.

<u>Links between lysosomal detoxification and biomineralization</u>

In their phylogenetic analysis of silicon transport genes, Marron *et al*. propose that the Lsi2
protein evolved in the original eukaryote as a detoxification mechanism to export silicic acid out
of cells in the high Si oceans of the Precambrian to prevent uncontrolled silica deposition and
were later co-opted for biosilicification by a specific set of lineages at the

Precambrian/Cambrian boundary.[7] This hypothesis implies a switch of Lsi2 orientation within
the membrane from exporting silicic acid out of cells to transporting it in. However, it is
possible that Lsi2 detoxified silicic acid by sequestering it within lysosomes rather than by
pumping it out of cells. Such a mechanism would seamlessly lend itself to the accumulation of
silicic acid into the SDV of diatoms and sponges, while also inserting at the plasma membrane
of plant cells in an orientation that mediates silicic acid extrusion into the xylem.
Lysosomes and detoxification are intrinsically linked. For example, calcium and heavy metal
ions including zinc, copper, and iron are sequestered in lysosomes and vacuoles in many
invertebrate phyla into structures termed concretions or granules, rendering them insoluble
and detoxified.[55–62] These organelles are often acidified by VHA.[63,64] The sequestered
concretions found in oysters,[56] bivalves,[58] snails,[59,60] insects,[61] and crustaceans[55,64] could be
considered precursors to skeletons. In fact, it has been proposed that metal detoxification was
a selective pressure that shaped the evolution of biomineralization.[65–67] Calcium ions are
disruptive to cellular functioning as their precipitation intracellularly would be deleterious;
therefore, sequestration within a pre-existing acidified organelle may have provided an
energetically effective solution,[65] while also offering a parsimonious mechanistic explanation at
the evolutionary level. Recently, it was shown that osteoblasts, cells responsible for secreting
calcium phosphate mineral precursors for bone formation, transport amorphous calcium
phosphate via lysosomes.[17] Additionally, pharmacological inhibition of VHA prevented the
formation of electron-dense granules and extracellular calcified nodules,[17] matching the
importance of VHA for silicification in sponges (current study) and in diatoms. [3] Furthermore,
other studies have shown that increased expression of the master regulator of lysosomal
biogenesis, *Tfeb*, enhances osteoblastic differentiation[68] and increases bone mass and
strength.[69] Altogether, the evidence suggests that ancient links between detoxification and
lysosomal machinery set the stage for the repeated evolution of biomineralization.
176
<u>Lysosomal machinery as part of ancestral genetic toolkit for biomineralization</u>
It has been proposed that genes involved in biomineralization were independently co-
opted from an ancestral genetic toolkit,[70] however, the specific genes within this toolkit have
not been clearly identified. Here, we found high expression of lysosomal genes in sclerocytes
including cathepsins, silicateins, TMEM55A/B, TMEM192, and TMEM199. Cathepsins are
lysosomal proteases speculated to play a role in making precise cuts in matrix and
mineralization proteins for the process of biosilicification.[9] They are upregulated during shell
repair in mussels,[16] cathepsin B/X has been identified in sea urchin spine extracts,[71] and our
analysis of single-cell RNASeq dataset of a stony coral revealed that cathepsin L and cathepsin V
are upregulated in skeleton-forming cells, calicoblasts.[72] Silicatein was discovered in the protein
filaments of spicules from demosponge *Tethya aurantia.*[23] Silicateins are highly similar to
members of the cathepsin L and papain family of proteases, but the cysteine at the active site in

the proteases is replaced by serine.[23] They have been proposed to retain protease activity

similar to other cathepsins,[73] catalyze the polymerization of silicic acid, [74] or serve as scaffolding

for spicule formation.[75] TMEM55A/B and TMEM192, involved in lysosomal autophagy,

phagocytosis, and lysosome repair,[24–26] and TMEM199,[27] a VHA assembly factor, have not been

previously described within the context of biosilicification to our knowledge. We found

upregulation of VHA mRNA in sclerocytes, VHA protein localization around spicules, and

showed that inhibition of VHA disrupts skeleton formation. This aligns with similar findings on

biosilicification in diatoms,[3] and calcification in bryozoans,[14] sea urchins,[13] foraminifera,[12]

coccolitophores,[11] and crustaceans.[15] We report that Lsi2 genes were upregulated in sponge

sclerocytes, likely playing a role in silicic acid uptake similar to plants.[6] In the marine

hexactinellid sponge *Aphrocallistes vastus*, VHA, Lsi2, and various cathepsins were upregulated

in the osculum region of indicating a potential link to sponge skeleton formation.[9] Exposure of

hexactinellid sponge *Vazella pourtalesii* to dissolved silicon (DSi), led to the upregulation of

cathepsins and Lsi2.[8] Further, a proteomic analysis found that spicule contents of marine

demosponge *Petrosia ficiformis* were dominated by proteins involved in lysosomal function

including cathepsins and silicatein.[76] While these studies used methods of bulk RNASeq and

proteomics of isolated spicules, a major advance of our study is the usage of single-cell RNASeq

data, *in situ* protein and mRNA localization, and functional assays to conclusively show the

importance of lysosomal genes in sponge skeleton formation. The machinery responsible for

the biogenesis of lysosome-related organelles is present across the eukaryotic tree of life and is

likely to have been in place prior to the last eukaryotic common ancestor.[77] Given that

lysosomal genes have been shown to be involved in biomineralization across a broad sweep of

taxa, the combined evidence suggest that lysosomal machinery was co-opted independently

multiple times for skeleton formation, forming a key set of genes within the ancestral toolkit for

biomineralization.

Conclusion

Skeleton formation was a key innovation that catapulted the trajectory of body plan

evolution during the Cambrian. It provided organisms with new possibilities for protection,

support, locomotion, and defense.[1] Biomineralization is intrinsically linked with global calcium

and silica cycling. Importantly, many marine organisms are ballasted by their silica and calcium

skeletons for burial in the seafloor, allowing for carbon sequestration.[78,79] Here, we describe the

co-option of lysosomal genes for biosilicification in sponges, the earliest diverged branch of

metazoans.[80] We discuss parallels to biomineralization pathways across the tree of life, from

protists to mammals. Although biomineralization evolved independently multiple times, the co-

option of genes from a conserved toolkit is remarkable.[70] Our findings provide novel evidence

for the co-option of ancestral lysosomal machinery for biomineralization.

Methods

Organisms
Ephydatia muelleri and *Spongilla lacustris* gemmules were obtained from the Nichols
lab. They were collected from the upper peninsula of Michigan in 2023. Gemmules were stored
at 4°C and hatched in spring water (Crystal Geyser) at room temperature in cover-slip bottom
dishes as previously described[81].

RNASeq analysis
Datasets acquired from Musser et al[18] and used Seurat in R to plot the expression of
sclerocytes markers. We referenced the *S. lacustris* transcriptome and proteome provided in
Musser et al[18] to identify homologs and design HCR-FISH probes. R code and lists of *S. lacustris*
genes analyzed can be found in our github repository.

Antibodies
Sponges were probed for VHA using a custom rabbit polyclonal antibody developed
using a peptide antigen matching a conserved region of VHA's B subunit (VHA$_B$;
AREEVPGRRGFPGY; GenScript Biotech Corporation, Piscataway, NJ, USA) which is conserved in
species ranging from sponges to humans.[34,35,82–84] We confirmed the epitope is 100% conserved
in *E. muelleri* and *S. lacustris* (Supplementary Figure 5A).

Immunohistochemistry
Immunohistochemical localization of anti-VHA$_B$ antibody in sponges was done following
previously described protocol.[85] Briefly, 4-7 day old gemmule-hatched *E. muelleri* and *S.
Lacustris* sponges were fixed with 4% paraformaldehyde in 95% EtOH for 45 minutes at room
temperature. Sponges were rinsed with 0.1% Tween 20 (v/v) in phosphate buffered saline
(PTw) three times before incubation with blocking buffer (3% Bovine Albumin Serum in PTw) for
45 minutes at room temperature. Samples were incubated overnight (4°C) with either 0.962 µg
ml^{-1} polyclonal anti-VHA$_B$ primary antibody in blocking buffer or 0.962 µg ml^{-1} polyclonal anti-
VHA$_B$ primary antibody with 60x peptide as a "peptide preabsorption control". Sponges were
rinsed with PTw three times and subsequently incubated with 4 µg ml^{-1} secondary antibody
(goat anti–rabbit–Alexa Fluor 488; A-11008; Invitrogen, Carlsbad, CA, USA) and 4 µg ml^{-1} 4',6-
diamidino-2-phenylindole (DAPI) DNA stain (Invitrogen) in blocking buffer for 45 minutes in the
dark at room temperature. Excess secondary antibody and DAPI were washed from samples by
rinsing three times with PTw and they were mounted for imaging using ProLong Glass Antifade
Mountant (P36982; Invitrogen). Confocal microscopy was performed on a Zeiss AxioObserver
Z1 connected to a laser scanner equipped with 405, 488, 561, and 640 nm laser lines (Zeiss LSM
800 with Airyscan, Carl Zeiss AG, Oberkochen, Baden-Württemberg, Germany). Samples were
imaged with the following excitation and emission wavelengths: Alexa Fluor 488 — ex: 488 nm,
em: 488-561 nm; DAPI—ex: 405 nm, em: 400-488 nm.

Western blots
Custom polyclonal antibodies against VHA subunit B (VHA$_B$) were used to investigate
VHA localization in sponge tissues of demosponges *Ephydatia muelleri* and *Spongilla lacustris*.

Approximately 50-80 *E. muelleri* and *S. lacustris* gemmules were hatched in separate
petri dishes. At 9-11 days old, sponges were scraped off the dishes with a razor blade and
dounce homogenized in 100 μL of 1x protease inhibitor cocktail (P2714; Sigma-Aldrich) in spring
water. Samples were sonicated for 3x10 second bursts with 1-minute rests on ice.
Homogenates were spun down for two minutes and the supernatant was retained. The protein
concentrations of samples were obtained using Bradford assay performed with a bovine serum
albumin standard curve (BioRad, Hercules, CA, USA). Samples were incubated in Laemmli
Sample buffer with 5% (v/v) β-mercaptoethanol for 5 min at 100 °C and increasing amounts of
protein were loaded into a sodium dodecyl sulphate–polyacrylamide gel electrophoresis gel in
duplicate (*E. muelleri*: 0.7 μg, 3.5 μg, 7 μg, 13.9 μg; *S. lacustris*: 0.7 μg, 3.5 μg, 7 μg, 13.9 μg).
The gel was run in in Tris-glycine running buffer (25 mM Tris-base, 192 mM glycine, 0.1% (w/v)
sodium dodecyl sulphate, pH 8.3; Sigma-Aldrich) at 47V for 35 minutes followed by 150V for 35-
40 minutes until the proteins had reached the bottom of the gel. Proteins were transferred
from the gel onto a polyvinylidene difluoride (PVDF) membrane using a Mini Tans-Blot Cell (Bio-
Rad) filled with Towbin transfer buffer (25 mM Tris, 192 mM glycine, 20% (v/v) methanol, pH
8.3; Sigma-Aldrich) at 90 mAmps overnight. PVDF membranes were incubated with blocking
buffer (1g non-fat dry milk powder in 10 mls Tris-buffered saline with 0.1% Tween (TBS-T)) for 1
hour at room temperature on an orbital shaker and subsequently incubated overnight at 4 °C
with either 0.2405 μg/mL anti-VHA$_B$ primary antibody or anti-VHA$_B$ primary antibody with 60x
excess peptide ("peptide preabsorption control") in blocking buffer. Membranes were washed
with 3 x 10 min TBS-T and then incubated with secondary antibody (goat anti–rabbit–
horseradish peroxidase diluted 1:5000; Bio-Rad) for 1 hour on an orbital shaker at room
temperature. Membranes were washed again with 3 x 10 min TBS-T and imaged with Clarity
Max Western ECL Substrate (Bio-Rad) using a Chemidoc Imaging system (Bio-Rad). Western
blotting specifically detected the expected ~55 kDa band (Supplementary Figure 5B).
295
HCR-FISH protocol
Probes were designed using publicly available HCR-FISH probe generator.[86–89] Sequences
against which probes were designed are provided in supplementary table 2. *E. muelleri* Lsi2
homologs were identified by blasting Lsi2 sequences against an unpublished transcriptome. 6-
10 day old gemmule-hatched *E. muelleri* and *S. Lacustris* sponges were stained using HCR FISH
probes following an adapted previously described protocol.[90] Briefly, sponges were fixed in cold
4% PFA in 1/4x HF buffer (147.5mM NaCl, 0.1675 mM KCl, 0.19 mM CaCl$_2$, 0.06mM NaHCO in
RNAse-free water) on a gently rocking platform at 4C overnight. Samples were rinsed with ¼ x
HF buffer and then incubated with ¼ x HF buffer containing an RNAse inhibitor (1 U/μl final
concentration) at room temperature for 10 minutes. Sponges were rinsed again with ¼ x HF
buffer and progressively dehydrated into 100% methanol. Subsequently, 100% methanol was
replaced with 100% ethanol, sponges were progressively rehydrated into 0.1% Tween 20 (v/v)
in phosphate buffered saline (PTw) and incubated with of 5 μg/mL Proteinase K in PTw for 90
seconds at room temperature. The proteinase K solution was removed and sponges were
incubated with 2mg/ml glycine in PTw for 5 minutes at room temperature twice. Samples were
rinsed with PTw and re-fixed with 4% PFA in PTw for one hour at room temperature. Following
two rinses with PTw, samples were incubated with 2x Saline Sodium Citrate (SSC) for 10
minutes at room temperature twice. Sponges were then pre-hybridized by incubation with

Hybridization buffer (Molecular Instruments), first at room temperature for 10 minutes and
then at 37°C for 30 minutes. Subsequently, samples were incubated with 0.04 µM probe in
hybridization buffer or just hybridization buffer for the "no probe, hairpin only control"
(Supplemnetary figure 6) at 37°C overnight. Sponges were rinsed with probe wash buffer
(Molecular Instruments) for 5 mins at 37°C three times, then incubated with 2x SSC for 5
minutes at room temperature twice. Samples were pre-amplified with amplification buffer
(Molecular Instruments) for 5 minutes at room temperature twice, then incubated with 60 nM
of snapcooled (incubated at 95°C for 90 seconds, then cooled in the dark for at least 30
minutes) hairpins (Molecular Instruments) in amplification buffer at room temperature in the
dark overnight. Specimens were rinsed with 2x SSC for 5 minutes at room temperature twice.
2x SSC was replaced with 10 µg ml^{-1} 4',6-diamidino-2-phenylindole (DAPI) in PBS for 20 minutes
at room temperature. Sponges were rinsed with PBS and then mounted using ProLong Glass
Antifade Mountant (P36982; Invitrogen). Confocal microscopy was performed on a Zeiss
AxioObserver Z1 connected to a laser scanner equipped with 405, 488, 561, and 640 nm laser
lines (Zeiss LSM 800 with Airyscan, Carl Zeiss AG, Oberkochen, Baden-Württemberg, Germany).
Samples were imaged with the following excitation and emission wavelengths: Alexa Fluor 555
— ex: 561 nm, em: 561- 585 nm; Alexa Fluor 488 — ex: 488 nm, em: 488-561 nm; DAPI—ex:
405 nm, em: 400-488 nm.
332
PDMPO experiments
4-11 day old gemmule-hatched *E. muelleri* and *S. Lacustris* sponges were incubated with
1 uM PDMPO [2-(4-pyridyl)-5-((4-(2-
dimethylaminoethylaminocarbamoyl)methoxy)phenyl)oxazole] or LysoSensor Yellow/Blue
DND-160 (Invitrogen L7545) and either 100nM concanamycinA (concA) or an equivalent volume
of DMSO for 24 hours (*E. muelleri* n=4 and *S. lacustris* n=3 for each condition). PDMPO
fluorescence (ex: 405 nm, em: 490-515 nm) and brightfield images were taken in five 0.408
mm^2 regions around the periphery of each sponge. The number of spicules per region and the
number of spicules per region exhibiting PMDPO fluorescence were counted in ImageJ. The
proportion of spicules with PDMPO fluorescence was calculated by dividing number of spicules
exhibiting PDMPO fluorescence by the total number of spicules per region. The number of
spicules per mm^2 and the proportion of spicules with PDMPO were averaged across regions to
get the final mean values per sponge. The statistical differences for both aforementioned
parameters between sponges treated with concA versus DMSO were calculated using a student
t-test. Stereoscope images of all sponges in the experiment were taken before and after
treatment to confirm viability of the sponges (Supplementary Figure 6).
349
350   Resource Availability
351   Lead Contact
352   Requests for further information and resources should be directed to and will be fulfilled by the
353   lead contact, Maitri Rangarajan-Paul (mapaul@ucsd.edu).
354
Materials availability
Antibodies used for immunostaining of sponges will be shared by the lead contact upon
request.

Data and Code availability
Data and code used in this study are available at https://github.com/maitripaul/Co-option-of-
Lysosomal-Machinery-for-Biomineralization
361

362 Acknowledgments
363 We thank Jake Pawley and Haley Womack for their assistance in providing us with sponge
364 gemmules and advice on sponge-rearing. M.R.P. was funded by NSF-GRFP 2023360321.

365
Author contributions
Conceptualization: M.R.P., M.T.; Data curation: M.R.P., S.N.; Formal analysis: M.R.P., S.N., M.T.;
Funding acquisition: M.R.P., M.T.; Investigation: M.R.P.; Methodology: M.R.P., S.N., M.T.;
Project administration: M.R.P., M.T.; Resources: S.N., M.T.; Software: M.R.P.; Supervision: M.T.;
Validation: M.R.P.; Visualization: M.R.P., S.N., M.T.; Writing – original draft: M.R.P., M.T.;
Writing – editing: M.R.P., S.N., M.T.
372
373 Declaration of interests
374 The authors declare no competing interests.
375

References
1. Gilbert, P.U.P.A., Bergmann, K.D., Boekelheide, N., Tambutté, S., Mass, T., Marin, F., Adkins,
J.F., Erez, J., Gilbert, B., Knutson, V., et al. (2022). Biomineralization: Integrating mechanism
and evolutionary history. Science Advances *8*, eabl9653.
https://doi.org/10.1126/sciadv.abl9653.

2. Müller, W.E.G., Wang, X., Cui, F.-Z., Jochum, K.P., Tremel, W., Bill, J., Schröder, H.C., Natalio,
F., Schloßmacher, U., and Wiens, M. (2009). Sponge spicules as blueprints for the
biofabrication of inorganic–organic composites and biomaterials. Appl Microbiol Biotechnol
83, 397–413. https://doi.org/10.1007/s00253-009-2014-8.

3. Yee, D.P., Hildebrand, M., and Tresguerres, M. (2020). Dynamic subcellular translocation of
V-type H+-ATPase is essential for biomineralization of the diatom silica cell wall. New
Phytologist *225*, 2411–2422. https://doi.org/10.1111/nph.16329.

4. Jahn, T., Baluska, F., Michalke, W., Harper, J.F., and Volkmann, D. (1998). Plasma membrane
H+-ATPase in the root apex: Evidence for strong expression in xylem parenchyma and
asymmetric localization within cortical and epidermal cells. Physiologia Plantarum *104*, 311–
316. https://doi.org/10.1034/j.1399-3054.1998.1040304.x.

5. Sattelmacher, B. (2001). The apoplast and its significance for plant mineral nutrition. New
Phytologist *149*, 167–192. https://doi.org/10.1046/j.1469-8137.2001.00034.x.

6. Ma, J.F., Yamaji, N., Mitani, N., Tamai, K., Konishi, S., Fujiwara, T., Katsuhara, M., and Yano,
M. (2007). An efflux transporter of silicon in rice. Nature *448*, 209–212.
https://doi.org/10.1038/nature05964.

7. Marron, A.O., Ratcliffe, S., Wheeler, G.L., Goldstein, R.E., King, N., Not, F., de Vargas, C., and
Richter, D.J. (2016). The Evolution of Silicon Transport in Eukaryotes. Mol Biol Evol *33*, 3226–
3248. https://doi.org/10.1093/molbev/msw209.

8. Maldonado, M., López-Acosta, M., Beazley, L., Kenchington, E., Koutsouveli, V., and Riesgo,
A. (2020). Cooperation between passive and active silicon transporters clarifies the
ecophysiology and evolution of biosilicification in sponges. Science Advances *6*, eaba9322.
https://doi.org/10.1126/sciadv.aba9322.

9. Francis, W.R., Eitel, M., Vargas, S., Garcia-Escudero, C.A., Conci, N., Deister, F., Mah, J.L.,
Guiglielmoni, N., Krebs, S., Blum, H., et al. (2023). The genome of the reef-building glass
sponge Aphrocallistes vastus provides insights into silica biomineralization. Royal Society
Open Science *10*, 230423. https://doi.org/10.1098/rsos.230423.

10. Shrestha, R.P., Tesson, B., Norden-Krichmar, T., Federowicz, S., Hildebrand, M., and Allen,
A.E. (2012). Whole transcriptome analysis of the silicon response of the diatom Thalassiosira
pseudonana. BMC Genomics *13*, 499. https://doi.org/10.1186/1471-2164-13-499.

11. Mackinder, L., Wheeler, G., Schroeder, D., von Dassow, P., Riebesell, U., and Brownlee, C.
(2011). Expression of biomineralization-related ion transport genes in Emiliania huxleyi.
Environmental Microbiology *13*, 3250–3265. https://doi.org/10.1111/j.1462-
2920.2011.02561.x.

12. Toyofuku, T., Matsuo, M.Y., de Nooijer, L.J., Nagai, Y., Kawada, S., Fujita, K., Reichart, G.-J.,
Nomaki, H., Tsuchiya, M., Sakaguchi, H., et al. (2017). Proton pumping accompanies
calcification in foraminifera. Nat Commun *8*, 14145. https://doi.org/10.1038/ncomms14145.

13. Hu, M.Y., Petersen, I., Chang, W.W., Blurton, C., and Stumpp, M. (2020). Cellular
bicarbonate accumulation and vesicular proton transport promote calcification in the sea
urchin larva. Proceedings of the Royal Society B: Biological Sciences *287*, 20201506.
https://doi.org/10.1098/rspb.2020.1506.

14. Achilleos, K., Smith, A.M., Kenny, N.J., and Brown, C.M. (2024). A new transcriptome
resource for Cellaria immersa (Phylum: Bryozoa) reveals candidate genes and proteins
related to biomineralization. Front. Mar. Sci. *11*.
https://doi.org/10.3389/fmars.2024.1389708.

15. Ziegler, A., Weihrauch, D., Hagedorn, M., Towle, D.W., and Bleher, R. (2004). Expression and
polarity reversal of V-type H+-ATPase during the mineralization–demineralization cycle in
Porcellio scabersternal epithelial cells. J Exp Biol *207*, 1749–1756.
https://doi.org/10.1242/jeb.00953.

16. Yarra, T., Ramesh, K., Blaxter, M., Hüning, A., Melzner, F., and Clark, M.S. (2021).
Transcriptomic analysis of shell repair and biomineralization in the blue mussel, Mytilus
edulis. BMC Genomics *22*, 437. https://doi.org/10.1186/s12864-021-07751-7.

17. Iwayama, T., Okada, T., Ueda, T., Tomita, K., Matsumoto, S., Takedachi, M., Wakisaka, S.,
Noda, T., Ogura, T., Okano, T., et al. (2019). Osteoblastic lysosome plays a central role in
mineralization. Sci Adv *5*, eaax0672. https://doi.org/10.1126/sciadv.aax0672.

18. Musser, J.M., Schippers, K.J., Nickel, M., Mizzon, G., Kohn, A.B., Pape, C., Ronchi, P.,
Papadopoulos, N., Tarashansky, A.J., Hammel, J.U., et al. (2021). Profiling cellular diversity in
sponges informs animal cell type and nervous system evolution. Science *374*, 717–723.
https://doi.org/10.1126/science.abj2949.

19. Mindell, J.A. (2012). Lysosomal Acidification Mechanisms*. Annual Review of Physiology *74*,
69–86. https://doi.org/10.1146/annurev-physiol-012110-142317.

20. H. Brilliant, M. (2001). The Mouse p (pink-eyed dilution) and Human P Genes,
Oculocutaneous Albinism Type 2 (OCA2), and Melanosomal pH. Pigment Cell Research *14*,
86–93. https://doi.org/10.1034/j.1600-0749.2001.140203.x.

21. Sitaram, A., Piccirillo, R., Palmisano, I., Harper, D.C., Dell'Angelica, E.C., Schiaffino, M.V., and
Marks, M.S. (2009). Localization to Mature Melanosomes by Virtue of Cytoplasmic Dileucine
Motifs Is Required for Human OCA2 Function. MBoC *20*, 1464–1477.
https://doi.org/10.1091/mbc.e08-07-0710.

22. Sauka-Spengler, T., and Bronner-Fraser, M. (2008). A gene regulatory network orchestrates
neural crest formation. Nat Rev Mol Cell Biol *9*, 557–568. https://doi.org/10.1038/nrm2428.

23. Shimizu, K., Cha, J., Stucky, G.D., and Morse, D.E. (1998). Silicatein α: Cathepsin L-like
protein in sponge biosilica. Proceedings of the National Academy of Sciences *95*, 6234–6238.
https://doi.org/10.1073/pnas.95.11.6234.

24. Morioka, S., Nigorikawa, K., Okada, E., Tanaka, Y., Kasuu, Y., Yamada, M., Kofuji, S.,
Takasuga, S., Nakanishi, H., Sasaki, T., et al. (2018). TMEM55a localizes to macrophage
phagosomes to downregulate phagocytosis. J Cell Sci *131*, jcs213272.
https://doi.org/10.1242/jcs.213272.

25. Jeong, E., Willett, R., Rissone, A., La Spina, M., and Puertollano, R. (2024). TMEM55B links
autophagy flux, lysosomal repair, and TFE3 activation in response to oxidative stress. Nat
Commun *15*, 93. https://doi.org/10.1038/s41467-023-44316-6.

26. Nguyen, T.L., Schneppenheim, J., Rudnik, S., Lüllmann-Rauch, R., Bernreuther, C., Hermans-
Borgmeyer, I., Glatzel, M., Saftig, P., and Schröder, B. (2017). Functional characterization of
the lysosomal membrane protein TMEM192 in mice. Oncotarget *8*, 43635–43652.
https://doi.org/10.18632/oncotarget.17514.

27. Miles, A.L., Burr, S.P., Grice, G.L., and Nathan, J.A. (2017). The vacuolar-ATPase complex and
assembly factors, TMEM199 and CCDC115, control HIF1α prolyl hydroxylation by regulating
cellular iron levels. eLife *6*, e22693. https://doi.org/10.7554/eLife.22693.

28. Li, X., Xiang, C., Zhu, S., Guo, J., Liu, C., Wang, A., Cao, J., Lu, Y., Neculai, D., Xu, P., et al.
(2024). SNX8 enables lysosome reformation and reverses lysosomal storage disorder. Nat
Commun *15*, 2553. https://doi.org/10.1038/s41467-024-46705-x.

29. Tang, T., Yang, Z., Wang, D., Yang, X., Wang, J., Li, L., Wen, Q., Gao, L., Bian, X., and Yu, S.
(2020). The role of lysosomes in cancer development and progression. Cell Biosci *10*, 131.
https://doi.org/10.1186/s13578-020-00489-x.

30. Mattioni, A., Boldt, K., Auciello, G., Komada, M., Rappoport, J.Z., Ueffing, M., Castagnoli, L.,
Cesareni, G., and Santonico, E. (2020). Ring Finger Protein 11 acts on ligand-activated EGFR
via the direct interaction with the UIM region of ANKRD13 protein family. The FEBS Journal
287, 3526–3550. https://doi.org/10.1111/febs.15226.

31. Kibak, H., Taiz, L., Starke, T., Bernasconi, P., and Gogarten, J.P. (1992). Evolution of structure
and function of V-ATPases. J Bioenerg Biomembr *24*, 415–424.
https://doi.org/10.1007/BF00762534.

32. Tresguerres, M. (2016). Novel and potential physiological roles of vacuolar-type H+-ATPase
in marine organisms. J Exp Biol *219*, 2088–2097. https://doi.org/10.1242/jeb.128389.

33. Tresguerres, M., Parks, S.K., Wood, C.M., and Goss, G.G. (2007). V-H+-ATPase translocation
during blood alkalosis in dogfish gills: interaction with carbonic anhydrase and involvement
in the postfeeding alkaline tide. American Journal of Physiology-Regulatory, Integrative and
Comparative Physiology *292*, R2012–R2019. https://doi.org/10.1152/ajpregu.00814.2006.

34. Tresguerres, M., Katz, S., and Rouse, G.W. (2013). How to get into bones: proton pump and
carbonic anhydrase in Osedax boneworms. Proceedings of the Royal Society B: Biological
Sciences *280*, 20130625. https://doi.org/10.1098/rspb.2013.0625.

35. Barott, K.L., Venn, A.A., Perez, S.O., Tambutté, S., and Tresguerres, M. (2015). Coral host
cells acidify symbiotic algal microenvironment to promote photosynthesis. Proceedings of
the National Academy of Sciences *112*, 607–612. https://doi.org/10.1073/pnas.1413483112.

36. Barott, K.L., Thies, A.B., and Tresguerres, M. (2022). V-type H+-ATPase in the symbiosome
membrane is a conserved mechanism for host control of photosynthesis in anthozoan
photosymbioses. Royal Society Open Science *9*, 211449.
https://doi.org/10.1098/rsos.211449.

37. Thies, A.B., Rangarajan-Paul, M., Wangpraseurt, D., and Tresguerres, M. (2025). Co-option
of immune and digestive cellular machinery to support photosymbiosis in amoebocytes of
the upside-down jellyfish Cassiopea xamachana. Journal of Experimental Biology *228*,
jeb249849. https://doi.org/10.1242/jeb.249849.

38. Hargadon, A.C., Viliunas, J.W., Koehler, S., Thies, A.B., Chen, G.Y., Ladinsky, M.S., Kuwabara,
J., Avila-Magana, V., Ruby, E.G., Tresguerres, M., et al. (2024). An acidic microenvironment

produced by the V-type ATPase of Euprymna scolopes promotes specificity during Vibrio
fischeri recruitment. Commun Biol 7, 1642. https://doi.org/10.1038/s42003-024-07348-2.

39. Allard, C.A.H., Thies, A.B., Mitra, R., Vaelli, P.M., Leto, O.D., Walsh, B.L., Laetz, E.M.J.,
Tresguerres, M., Lee, A.S.Y., and Bellono, N.W. (2025). A host organelle integrates stolen
chloroplasts for animal photosynthesis. Cell 188, 5266-5277.e13.
https://doi.org/10.1016/j.cell.2025.06.003.

40. Seidel, T., Schnitzer, D., Golldack, D., Sauer, M., and Dietz, K.-J. (2008). Organelle-specific
isoenzymes of plant V-ATPase as revealed by in vivo-FRET analysis. BMC Cell Biol 9, 28.
https://doi.org/10.1186/1471-2121-9-28.

41. Shimizu, K., Amo, Y.D., Brzezinski, M.A., Stucky, G.D., and Morse, D.E. (2001). A novel
fluorescent silica tracer for biological silicification studies. Chemistry & Biology 8, 1051–1060.
https://doi.org/10.1016/S1074-5521(01)00072-2.

42. Dröse, S., and Altendorf, K. (1997). Bafilomycins and concanamycins as inhibitors of V-
ATPases and P-ATPases. J Exp Biol 200, 1–8. https://doi.org/10.1242/jeb.200.1.1.

43. Huss, M., Ingenhorst, G., König, S., Gaßel, M., Dröse, S., Zeeck, A., Altendorf, K., and
Wieczorek, H. (2002). Concanamycin A, the Specific Inhibitor of V-ATPases, Binds to the Vo
Subunit c *. Journal of Biological Chemistry 277, 40544–40548.
https://doi.org/10.1074/jbc.M207345200.

44. Vrieling, E.G., Gieskes, W.W.C., and Beelen, T.P.M. (1999). SILICON DEPOSITION IN
DIATOMS: CONTROL BY THE pH INSIDE THE SILICON DEPOSITION VESICLE. Journal of
Phycology 35, 548–559. https://doi.org/10.1046/j.1529-8817.1999.3530548.x.

45. Mejía, L.M., Isensee, K., Méndez-Vicente, A., Pisonero, J., Shimizu, N., González, C.,
Monteleone, B., and Stoll, H. (2013). B content and Si/C ratios from cultured diatoms
(Thalassiosira pseudonana and Thalassiosira weissflogii): Relationship to seawater pH and
diatom carbon acquisition. Geochimica et Cosmochimica Acta 123, 322–337.
https://doi.org/10.1016/j.gca.2013.06.011.

46. Baxter, I.R., Young, J.C., Armstrong, G., Foster, N., Bogenschutz, N., Cordova, T., Peer, W.A.,
Hazen, S.P., Murphy, A.S., and Harper, J.F. (2005). A plasma membrane H+-ATPase is
required for the formation of proanthocyanidins in the seed coat endothelium of Arabidopsis
thaliana. Proceedings of the National Academy of Sciences 102, 2649–2654.
https://doi.org/10.1073/pnas.0406377102.

47. Mesdaghi, S., Murphy, D.L., Simpkin, A.J., and Rigden, D.J. (2023). Structural insights into
pink-eyed dilution protein (Oca2). Biosci Rep 43, BSR20230060.
https://doi.org/10.1042/BSR20230060.

48. Kondo, T., Namiki, T., Coelho, S.G., Valencia, J.C., and Hearing, V.J. (2015). Oculocutaneous
albinism: Developing novel antibodies targeting the proteins associated with OCA2 and

OCA4. Journal of Dermatological Science *77*, 21–27.
https://doi.org/10.1016/j.jdermsci.2014.11.006.

49. Coskun, D., Deshmukh, R., Shivaraj, S.M., Isenring, P., and Bélanger, R.R. (2021). Lsi2: A
black box in plant silicon transport. Plant Soil *466*, 1–20. https://doi.org/10.1007/s11104-
021-05061-1.

50. Bellono, N.W., Escobar, I.E., Lefkovith, A.J., Marks, M.S., and Oancea, E. (2014). An
intracellular anion channel critical for pigmentation. eLife *3*, e04543.
https://doi.org/10.7554/eLife.04543.

51. Stauber, T., and Jentsch, T.J. (2013). Chloride in Vesicular Trafficking and Function. Annual
Review of Physiology *75*, 453–477. https://doi.org/10.1146/annurev-physiol-030212-183702.

52. Kahn, A.S., and Leys, S.P. (2017). Spicule and flagellated chamber formation in a growth
zone of Aphrocallistes vastus (Porifera, Hexactinellida). Invertebrate Biology *136*, 22–30.
https://doi.org/10.1111/ivb.12155.

53. Schröder, H.-C., Perović-Ottstadt, S., Rothenberger, M., Wiens, M., Schwertner, H., Batel, R.,
Korzhev, M., Müller, I.M., and Müller, W.E.G. (2004). Silica transport in the demosponge
Suberites domuncula: fluorescence emission analysis using the PDMPO probe and cloning of
a potential transporter. Biochem J *381*, 665–673. https://doi.org/10.1042/BJ20040463.

54. Forgac, M. (2007). Vacuolar ATPases: rotary proton pumps in physiology and
pathophysiology. Nat Rev Mol Cell Biol *8*, 917–929. https://doi.org/10.1038/nrm2272.

55. Ahearn, G.A., Sterling, K.M., Mandal, P.K., and Roggenbeck, B. (2010). Heavy Metal
Transport and Detoxification by Crustacean Epithelial Lysosomes. In Epithelial Transport
Physiology, G. A. Gerencser, ed. (Humana Press), pp. 49–71. https://doi.org/10.1007/978-1-
60327-229-2_3.

56. Wang, W.-X., Yang, Y., Guo, X., He, M., Guo, F., and Ke, C. (2011). Copper and zinc
contamination in oysters: subcellular distribution and detoxification. Environ Toxicol Chem
30, 1767–1774. https://doi.org/10.1002/etc.571.

57. Engel, D.W. (1983). The intracellular partitioning of trace metals in marine shellfish. Sci
Total Environ *28*, 129–140. https://doi.org/10.1016/s0048-9697(83)80013-8.

58. Subcellular compartmentalization of Cd and Zn in two bivalves. I. Significance of metal-
sensitive fractions (MSF) and biologically detoxified metal (BDM) | U.S. Geological Survey
https://www.usgs.gov/publications/subcellular-compartmentalization-cd-and-zn-two-
bivalves-i-significance-metal-sensitive.

59. Howard, B., Mitchell, P.C., Ritchie, A., Simkiss, K., and Taylor, M. (1981). The composition of
intracellular granules from the metal-accumulating cells of the common garden snail (Helix
aspersa). Biochem J *194*, 507–511. https://doi.org/10.1042/bj1940507.

60. Desouky, M.M.A. (2006). Tissue distribution and subcellular localization of trace metals in
the pond snail Lymnaea stagnalis with special reference to the role of lysosomal granules in
metal sequestration. Aquat Toxicol *77*, 143–152.
https://doi.org/10.1016/j.aquatox.2005.11.009.

61. Ballan-Dufrançais, C. (2002). Localization of metals in cells of pterygote insects. Microsc Res
Tech *56*, 403–420. https://doi.org/10.1002/jemt.10041.

62. Pigino, G., Migliorini, M., Paccagnini, E., and Bernini, F. (2006). Localisation of heavy metals
in the midgut epithelial cells of Xenillus tegeocranus (Hermann, 1804) (Acari: Oribatida).
Ecotoxicol Environ Saf *64*, 257–263. https://doi.org/10.1016/j.ecoenv.2005.12.012.

63. Blaby-Haas, C.E., and Merchant, S.S. (2014). Lysosome-related Organelles as Mediators of
Metal Homeostasis. J Biol Chem *289*, 28129–28136.
https://doi.org/10.1074/jbc.R114.592618.

64. Sterling, K.M., Mandal, P.K., Roggenbeck, B.A., Ahearn, S.E., Gerencser, G.A., and Ahearn,
G.A. (2007). Heavy metal detoxification in crustacean epithelial lysosomes: role of anions in
the compartmentalization process. J Exp Biol *210*, 3484–3493.
https://doi.org/10.1242/jeb.008300.

65. Simkiss, K. (1977). Biomineralization and detoxification. Calc. Tis Res. *24*, 199–200.
https://doi.org/10.1007/BF02223316.

66. Carney, C.K., Harry, S.R., Sewell, S.L., and Wright, D.W. (2007). Detoxification Biominerals. In
Biomineralization I: Crystallization and Self-Organization Process, K. Naka, ed. (Springer), pp.
155–185. https://doi.org/10.1007/128_050.

67. Li, X., Sun, M., Zhang, L., Finlay, R.D., Liu, R., and Lian, B. (2022). Widespread bacterial
responses and their mechanism of bacterial metallogenic detoxification under high
concentrations of heavy metals. Ecotoxicol Environ Saf *246*, 114193.
https://doi.org/10.1016/j.ecoenv.2022.114193.

68. Tsukuba, T., Sakai, E., Nishishita, K., Kadowaki, T., and Okamoto, K. (2017). New functions of
lysosomes in bone cells. Journal of Oral Biosciences *59*, 92–95.
https://doi.org/10.1016/j.job.2017.01.004.

69. James, A., Hendrixson, J.A., Kadhim, I., Marques-Carvalho, A., Laster, J., Crawford, J.,
Thostenson, J., Wanchai, V., Sato, A.Y., Nookaew, I., et al. (2025). Elevation of master
autophagy regulator *Tfeb* in osteoblast lineage cells increases bone mass and strength. JCI
Insight *10*. https://doi.org/10.1172/jci.insight.191688.

70. Murdock, D.J.E. (2020). The 'biomineralization toolkit' and the origin of animal skeletons.
Biological Reviews *95*, 1372–1392. https://doi.org/10.1111/brv.12614.

71. Sciani, J.M., Antoniazzi, M.M., Neves, A. da C., and Pimenta, D.C. (2013). Cathepsin B/X is
secreted by Echinometra lucunter sea urchin spines, a structure rich in granular cells and
toxins. J Venom Anim Toxins Incl Trop Dis *19*, 33. https://doi.org/10.1186/1678-9199-19-33.

72. Levy, S., Elek, A., Grau-Bové, X., Menéndez-Bravo, S., Iglesias, M., Tanay, A., Mass, T., and
Sebé-Pedrós, A. (2021). A stony coral cell atlas illuminates the molecular and cellular basis of
coral symbiosis, calcification, and immunity. Cell *184*, 2973-2987.e18.
https://doi.org/10.1016/j.cell.2021.04.005.

73. Müller, W.E.G., Wang, X., Kropf, K., Ushijima, H., Geurtsen, W., Eckert, C., Tahir, M.N.,
Tremel, W., Boreiko, A., Schloßmacher, U., et al. (2008). Bioorganic/inorganic hybrid
composition of sponge spicules: Matrix of the giant spicules and of the comitalia of the deep
sea hexactinellid *Monorhaphis*. Journal of Structural Biology *161*, 188–203.
https://doi.org/10.1016/j.jsb.2007.10.009.

74. Povarova, N.V., Barinov, N.A., Baranov, M.S., Markina, N.M., Varizhuk, A.M., Pozmogova,
G.E., Klinov, D.V., Kozhemyako, V.B., and Lukyanov, K.A. (2018). Efficient silica synthesis from
tetra(glycerol)orthosilicate with cathepsin- and silicatein-like proteins. Sci Rep *8*, 16759.
https://doi.org/10.1038/s41598-018-34965-9.

75. Görlich, S., Samuel, A.J., Best, R.J., Seidel, R., Vacelet, J., Leonarski, F.K., Tomizaki, T.,
Rellinghaus, B., Pohl, D., and Zlotnikov, I. (2020). Natural hybrid silica/protein superstructure
at atomic resolution. Proceedings of the National Academy of Sciences *117*, 31088–31093.
https://doi.org/10.1073/pnas.2019140117.

76. Pozzolini, M., Mikšík, I., Ghignone, S., Oliveri, C., Tassara, E., and Giovine, M. (2022). The
Lysosome Origin of Biosilica Machinery in the Demospongiae Model Petrosia ficiformis
(Poiret, 1789). Front. Mar. Sci. *9*. https://doi.org/10.3389/fmars.2022.850653.

77. More, K.J., Kaufman, J.G.G., Dacks, J.B., and Manna, P.T. (2024). Evolutionary origins of the
lysosome-related organelle sorting machinery reveal ancient homology in post-endosome
trafficking pathways. Proc Natl Acad Sci U S A *121*, e2403601121.
https://doi.org/10.1073/pnas.2403601121.

78. Armstrong, R.A., Lee, C., Hedges, J.I., Honjo, S., and Wakeham, S.G. (2001). A new,
mechanistic model for organic carbon fluxes in the ocean based on the quantitative
association of POC with ballast minerals. Deep Sea Research Part II: Topical Studies in
Oceanography *49*, 219–236. https://doi.org/10.1016/S0967-0645(01)00101-1.

79. Aldridge, D., Beer, C.J., and Purdie, D.A. (2012). Calcification in the planktonic foraminifera
Globigerina bulloides linked to phosphate concentrations in surface waters of the North
Atlantic Ocean. Biogeosciences *9*, 1725–1739. https://doi.org/10.5194/bg-9-1725-2012.

80. Steenwyk, J.L., and King, N. (2025). Integrative phylogenomics positions sponges at the root
of the animal tree. Science *390*, 751–756. https://doi.org/10.1126/science.adw9456.

81. Nichols, S. (2023). Growing freshwater sponges from gemmules in the laboratory.

82. Armstrong, E.J., Roa, J.N., Stillman, J.H., and Tresguerres, M. (2018). Symbiont
photosynthesis in giant clams is promoted by V-type H+-ATPase from host cells. J Exp Biol
221, jeb177220. https://doi.org/10.1242/jeb.177220.

83. Damsgaard, C., Lauridsen, H., Harter, T.S., Kwan, G.T., Thomsen, J.S., Funder, A.M., Supuran,
C.T., Tresguerres, M., Matthews, P.G., and Brauner, C.J. (2020). A novel acidification
mechanism for greatly enhanced oxygen supply to the fish retina. eLife *9*, e58995.
https://doi.org/10.7554/eLife.58995.

84. Roa, J.N., and Tresguerres, M. (2016). Soluble adenylyl cyclase is an acid-base sensor in
epithelial base-secreting cells. American Journal of Physiology-Cell Physiology *311*, C340–
C349. https://doi.org/10.1152/ajpcell.00089.2016.

85. Nichols, S. (2023). Fixation and staining of gemmule-hatched Ephydatia muelleri for
fluorescence microscopy v1. Preprint, https://doi.org/10.17504/protocols.io.j8nlkwnx6l5r/v1
https://doi.org/10.17504/protocols.io.j8nlkwnx6l5r/v1.

86. Kuehn, E., Clausen, D.S., Null, R.W., Metzger, B.M., Willis, A.D., and Özpolat, B.D. (2022).
Segment number threshold determines juvenile onset of germline cluster expansion in
Platynereis dumerilii. J Exp Zool B Mol Dev Evol *338*, 225–240.
https://doi.org/10.1002/jez.b.23100.

87. Choi, H.M.T., Schwarzkopf, M., Fornace, M.E., Acharya, A., Artavanis, G., Stegmaier, J.,
Cunha, A., and Pierce, N.A. (2018). Third-generation in situ hybridization chain reaction:
multiplexed, quantitative, sensitive, versatile, robust. Development *145*, dev165753.
https://doi.org/10.1242/dev.165753.

88. Wang, Y., Liu, X., Zeng, Y., Saka, S.K., Xie, W., Goldaracena, I., Kohman, R.E., Yin, P., and
Church, G.M. (2024). Multiplexed in situ protein imaging using DNA-barcoded antibodies
with extended hybridization chain reactions. Nucleic Acids Res *52*, e71.
https://doi.org/10.1093/nar/gkae592.

89. Tsuneoka, Y., and Funato, H. (2020). Modified in situ Hybridization Chain Reaction Using
Short Hairpin DNAs. Front Mol Neurosci *13*, 75. https://doi.org/10.3389/fnmol.2020.00075.

90. Nichols, S. (2023). HCR-fluorescent in situ hybridization (HCR-FISH) of gemmule-hatched
freshwater sponges.

Figure 1. Single-cell RNA-Seq expression of lysosome-related sclerocyte markers. (A) Heatmap showing expression of genes upregulated (log2fold>1) in sclerocytes that are linked with lysosomal function across *S. lacustris* cell types (single-cell RNASeq data from Musser et al., 2021). Note: not all VHA subunits are present here as they did not all make the cutoff log2fold cutoff. (B) Cartoon of VHA showing *S. lacustris* genes encoding each subunit.

Figure 2. VHA localization in *Ephydatia muelleri* and *Spongilla lacustris*. Immunohistochemical localization of VHA$_B$ in *E. muelleri* and *S. lacustris* sponges incubated with anti-VHA$_B$ antibody or anti-VHA$_B$ antibody with excess peptide (peptide pre-absorption control). VHA$_B$ in green, DAPI in blue. VHA$_B$, DAPI, and DIC channels are also shown separately. All images except DIC are orthogonal projections. Scale bars: 20 µm. sp; spicules.

Figure 3. Essential role of VHA in spiculogenesis in *Ephydatia muelleri* and *Spongilla lacustris*.
(A) Airyscan confocal image showing VHA$_B$ localization around *E. muelleri* spicule labelled with PDMPO (B) Representative images of PDMPO fluorescence of spicules in sponges incubated with DMSO or 100 nM concA for 24 hours. (C) Dot plot depicting the number of spicules per mm^2 of sponge tissue incubated with DMSO or 100 nM concA for 24 hours with error bars. (D) Dot plot depicting average proportion of spicules that showed PDMPO fluorescence in sponges incubated with DMSO or 100 nM concA for 24 hours with error bars. *E. muelleri* in red (n=4) and *S. lacustris* in cyan (n=3). Asterisks denote statistical significance (t test, p-value < 0.05). VHA$_B$ in green, PDMPO in magenta. Scale bars in (A): 5 μm and (B): 50 μm.

Figure 4. Gene expression of Lsi2 genes in *Ephydatia muelleri* and *Spongilla lacustris*. (A)
Heatmap showing expression of Lsi2, OCA2, and [Na$^+$/HCO$_3^-$[Si(OH)$_4$]] co-transporter (NBCSA)
genes in *S. lacustris* cell types. (B) HCR-FISH in *E. muelleri* and *S. lacustris* sponges incubated
with probes designed against Lsi2_1 and Lsi2_2 genes. Lsi2_1 in green, Lsi2_2 in orange, DAPI in
blue. Lsi2_1, Lsi2_2, DAPI, and DIC channels are also shown separately in grayscale. All images
are orthogonal projections. Scale bars: 20 μm. sp; spicules

Supplementary figures

Supplementary Figure 1. Phylogenetic and structural analysis of Lsi2/OCA2 homologs. (A) Lsi2 and OCA2 homologs were identified in *Ephydatia muelleri* (abbreviated Em) and *Spongilla lacustris* (abbreviated Sl) genomic and transcriptomic datasets by BLAST searches using choanoflagellate sequences from Marron *et al.* and by extraction of candidate orthogroups from OrthoFinder. Protein sequences were aligned with MAFFT using the DASH structure-guided alignment mode, after which DASH profile sequences were removed. The alignment was trimmed with ClipKIT to retain phylogenetically informative sites, and a maximum-likelihood phylogeny was inferred using IQ-TREE with automated model selection and branch support assessed by 1,000 ultrafast bootstraps and 1,000 SH-aLRT replicates. (B) Sequence alignment of *OsLsi2* gene from rice (*Oryza sativa),* Lsi2_1 and 2 from *E muelleri*, and Lsi2_1 and 2 from *S. lacustris.* Alignment shaded from light blue to dark blue based on percentage identity similarity. 11 predicted transmembrane (TM) domains in red (based on Ma et al., 2007). Sequence alignments were done on Clustal Omega and visualized using Jalview.

Supplementary Figure 2. Single-cell RNA-Seq expression of VHA subunits in *S. lacustris*. (A) Heatmap showing expression of genes encoding VHA subunits in *S. lacustris* cell types (single-cell RNASeq data from Musser et al., 2021). Highest expression of almost all VHA subunits occurs in sclerocytes. (B) Cartoon of VHA showing *S. lacustris* genes encoding each subunit.

Supplementary Figure 3. VHA localization at sclerocyte SDV and within cytoplasm in *S. lacustris*. (A, B) Confocal microscopy showing VHA$_B$ immunolabeling in *E. muelleri* sclerocytes surrounding spicules and within cytoplasm. (C) Airyscan confocal image showing VHA$_B$ localization around *E. muelleri* spicule labelled with PDMPO. Arrows indicate vesicles surrounded by VHA$_B$. VHA$_B$ in green, DAPI in blue, PDMPO in magenta. Scale bars: 20 µm

Supplementary Figure 4. HCR-FISH in *E. muelleri* sponges incubated with probes against Lsi2_1 and Lsi2_2 genes. Representative images showing more ubiquitous Lsi2_2 expression around spicules than Lsi2_1. Lsi2_1 in green, Lsi2_2 in orange, DAPI in blue. Lsi2_1, Lsi2_2, DAPI, and DIC channels are also shown separately in grayscale. Lsi1_1 and Lsi2_2 probes imaged at same laser intensity and gain. All images are orthogonal projections.

A

```
Homo sapiens    135  KGPVVMAEDFLDINGQPINPHSRIYPEEMIQTGISPIDVMNSIARGQKIPIFSAAGLPHNEIAAQICRQAGLVKKS-KAVLDYHDDNFAIVFAAMGVNMETARFFKSDFEQNGTMGNVCL  253
S. pistillata   124  KGPAVMAEDFLDIQGQPINPWSRIYPEEMIQTGISAIDTMNSIARGQKIPIFSAAGLPHNDIAAQICRQAGLVKFK-KGVMDDHSDNFAIVFAAMGVNMETARFFKQDFEENGSMENVCL  242
S. lacustris    133  KGPNILAEDFLDIQGQPINPWSRIYPEEMIQTGISAIDTMNSIARGQKIPIFSAAGLPHNDIAAQICRQGGLVKQTPKGVIDDHEANFAIVFAAMGVNMETARFFKQDFEENGSMENVCL  252
E. muelleri     133  KGPNILAEDFLDIQGQPINPWSRIYPEEMIQTGISAIDTMNSIARGQKIPIFSAAGLPHNDIAAQICRQGGLVKQTPKGVIDDHEANFAIVFAAMGVNMETARFFKQDFEENGSMENVCL  252

Homo sapiens    254  FLNLANDPTIERIITPRLALTTAEFLAYQCEKHVLVILTDMSSYAEALREVSAAREEVPGRRGFPGYMYTDLATIYERAGRVEGRGGSITQIPILTMPNDDITHPIPDLTGFITEGQIYV  373
S. pistillata   243  FLNLANDPTIERIITPRLALTTAEFLAYQCENHVLVILTDMSSYAEALREVSAAREEVPGRRGFPGYMYTDLATIYERAGRVEGRNGSITQIPILTMPNDDITHPIPDLTGYITEGQIYV  362
S. lacustris    253  FLNLANDPTIERIITPRLALTTAEFLAYQCEKHVLVILTDMSSYAEALREVSAAREEVPGRRGFPGYMYTDLATIYERAGRVEGRNGSITQIPILTMPNDDITHPIPDLTGYITEGQIYV  372
E. muelleri     253  FLNLANDPTIERIITPRLALTTAEFLAYQCEKHVLVILTDMSSYAEALREVSAAREEVPGRRGFPGYMYTDLATIYERAGRVEGRNGSITQIPILTMPNDDITHPIPDLTGYITEGQIYV  372

Homo sapiens    374  DRQLHNRQIYPPINVLPSLSRLMKSAIGEGMTRKDHGDVSNQLYACYAIGKDVQAMKAVVGEEALTSEDLLYLEFLQKFEKNFINQGPYENRSVFESLDLGWKLLRIFPKEMLKRIPQAV  493
S. pistillata   363  DRQLHNRQVYPPINVLPSLSRLMKSAIGEGMTRKDHADVSNQLYANYAIGKDVQAMKAVVGEEALTSEDLLYLEFLSKFEKNFITQGTYENRSVFDSLDIGWSLLRIFPKEMLKRIPQSI  482
S. lacustris    373  DRQLHNRQIYPPINVLPSLSRLMKSAIGDGMTRKDHADVSNQLYANYAIGKDVQAMKAVVGEEALTSEDKLYLEFLIKFEKNFITQGVYENRTVFESLEIGWQLLRIFPKEMLKRIPQSV  492
E. muelleri     373  DRQLHNRQIYPPINVLPSLSRLMKSAIGDGMTRRDHADVSNQLYANYAIGKDVQAMKAVVGEEALSSEDKLYLEFLQKFEKNFIAQGVYENRTVFESLEIGWQLLRIFPKEMLKRIPQAV  492

Homo sapiens    494  IDEFYSREG  502
S. pistillata   483  LAEYYPRDG  491
S. lacustris    493  LDEYYARD   500
E. muelleri     493  LDEYYARD   500
```

B

Supplementary Figure 5. VHA expression and conservation in sponges. (A) Alignment of the human VHA$_B$ protein sequence with the coral *Stylophora pistillata, S. lacustris, and E. muelleri* VHA$_B$ protein sequences. (B) Western blot detection of VHA in E. muelleri and S. lacustris using anti-VHA$_B$ antibody, and anti-VHA$_B$ antibody preabsorbed with 60x excess peptide on blots loaded with 0.7 µg, 3.5 µg, 7 µg, and 13.9 µg protein (*E. muelleri*) and 0.53 µg, 5.3 µg, and 10.7 µg protein (*S. lacustris*).

Supplementary Figure 6. Stereoscope images of *E. muelleri* and *S. lacustris* before and after concA treatment to assess viability. 4-day old sponges before and after 24 hr treatment with 1 uM PDMPO and DMSO or 100 nM concA followed by washout and replacement with spring water + 1uM PDMPO in (A) *E. muelleri* and (B*) S. lacustris.* Scale bars: 1 mm