Fibrin Selective Alteplase with Improved Thrombolysis and Inhibition Resistance Engineered by Rational Design

Toul M.[1,2,#], Slonkova V.[1,#], Mican J.[1,2,#], Thalerova S.[2,3], Peskova M.[2,3,4], Kittova P.[2,4,5], Scheer P.[2,6], Hlozkova J.[2,6], Brhelova E.[2,7], Aksu D.A.[2,6], Biskupic J.[2,7], Ondrus J.[2,8], Kasparek P.[9], Batkova T.[2,9,10], Marek M.[1,2], Vitecek J.[2,3], Kubala L.[2,3], Mikulik R.[2,10], Damborsky J.[1,2], Bednar D.[1,2,*], Prokop Z.[1,2,*]

1. Loschmidt Laboratories, Department of Experimental Biology and RECETOX, Faculty of Science, Masaryk University, Kamenice 5/C13, 625 00 Brno, Czech Republic
2. International Clinical Research Center, St. Anne's University Hospital, Pekarska 53, 656 91 Brno, Czech Republic
3. Department of Biophysics of Immune System, Institute of Biophysics of the Czech Academy of Sciences, Kralovopolska 135, 612 65 Brno, Czech Republic
4. Department of Biochemistry, Faculty of Science, Masaryk University, Kamenice 5, 625 00 Brno, Czech Republic
5. Department of Biochemistry, Faculty of Medicine, Masaryk University, Kamenice 5, 625 00 Brno, Czech Republic
6. Department of Pharmacology and Toxicology, Faculty of Pharmacy, Masaryk University, Palackeho trida 1946/1, 612 00 Brno, Czech Republic
7. Department of Chemical Drugs, Faculty of Pharmacy, Masaryk University, Palackeho trida 1946/1, 612 00 Brno, Czech Republic
8. Veterinary Research Institute, Hudcova 296/70, 621 00 Brno, Czech Republic
9. BioVendor LM, Karasek 1767/1, 621 00 Brno, Czech Republic
10. Department of Neurology, Tomas Bata Regional Hospital, Havlickovo nabrezi 600, 762 75 Zlin, Czech Republic

[#] These authors contributed equally to the work
[*] Corresponding authors

Abstract

AIMS: Thrombolytic enzymes represent an accessible and streamlined treatment for acute ischemic stroke, yet current FDA-approved agents, alteplase and tenecteplase, are limited by incomplete recanalization, hemorrhagic risk, and neurotoxicity. We aimed to develop a multidisciplinary workflow for rational design and multi-level testing of thrombolytic proteins to generate a next-generation thrombolytic with improved potency and safety.

METHODS AND RESULTS: Our pipeline integrated *in silico* design, biochemical characterization, *in vitro* thrombolysis, and *in vivo* validation. Computational screening included rational mutagenesis to reduce side effects, reconstructed ancestral sequences, and database-mined homologs. Selected candidates were biochemically assessed for enzymatic activity, fibrin stimulation, selectivity and affinity, inhibition resistance, and clot penetrability. The most promising variants underwent *in vitro* clot thrombolysis and *in vivo* efficacy and safety studies in rats. Brnoteplase emerged as the lead candidate, exhibiting 80-fold enhanced fibrin selectivity, 4-fold higher inhibition resistance, and superior clot penetrability compared to alteplase. *In vitro*, Brnoteplase demonstrated deeper intrathrombus penetration and a progressive dose–response profile, unlike alteplase, which declined at higher concentrations. *In vivo*, Brnoteplase achieved higher thrombolysis rates and recanalization frequency across all doses, with extended biological half-life enabling bolus administration. Safety assessments revealed the lowest incidence of severe hemorrhagic transformation and minimal hemispheric asymmetry.

CONCLUSIONS: We established a robust pipeline for efficient generation and selection of novel thrombolytics and identified Brnoteplase as a promising candidate combining improved fibrin selectivity, inhibition resistance, biological half-life, and clot penetration. These features translate into higher thrombolytic efficacy and a promising safety profile *in vivo*, warranting further preclinical evaluation to confirm its potential as a next-generation thrombolytic for acute ischemic stroke.

Introduction

Cardiovascular diseases, including ischemic stroke, represent the leading cause of morbidity and mortality worldwide, and the burden continues to rise[1]. Ischemic stroke therapy based on mechanical thrombectomy is restricted to specialized stroke centers and will therefore never become a universal solution[2]. In contrast, drug-induced thrombolysis treatment is readily accessible, more affordable, and straightforward to administer, so it remains the first-line, widely applicable therapy[3]. Unfortunately, current thrombolytic drugs still possess major limitations[4] and their further improvement is critical to reverse the alarming statistics and growing socioeconomic burden of stroke[5].

For almost 30 years, alteplase was the only Food and Drug Administration (FDA)-approved thrombolytic drug for acute ischemic stroke. Despite its proven effectiveness, it exhibits several drawbacks, including lower than 50 % recanalization rates, short half-life, and an increased risk of intracranial hemorrhage[6,7]. In March 2025, tenecteplase was approved by the FDA as the second drug for acute ischemic stroke treatment[8]. Tenecteplase is a mutant variant of alteplase with higher fibrin specificity and improved inhibition resistance, resulting in an extended half-life that enables bolus administration[9]. However, tenecteplase does not show clear superiority over alteplase in terms of efficacy or safety[10-13]. Therefore, there is a continuous need to develop a more efficient and safer thrombolytic drug with attenuated side effects.

An ideal thrombolytic should be not only more active but primarily more fibrin-selective and with alleviated binding to receptors, namely low density lipoprotein receptor-related protein 1 (LRP1) and N-methyl-D-aspartate receptor (NMDAR)[14]. These properties would make the thrombolytic safer by decreasing hemorrhagic and neurotoxicity risks[15,16]. Furthermore, reduced receptor binding combined with enhanced plasminogen activator inhibitor-1 (PAI-1) resistance would prolong the agent biological half-life, increasing overall efficacy throughout the therapeutic window and allowing convenient bolus administration, as achieved with tenecteplase[17].

Here, we report the development of an experimental pipeline for the design, comprehensive characterization, and selection of novel thrombolytic drugs (**Figure 1**). The pipeline allows for

initial screening of numerous candidate thrombolytics and their systematic filtering at multiple

levels to select only the most promising variants for demanding and ethically sensitive *in vivo*

animal and preclinical studies. We validated the robustness of this pipeline by generating

improved variants of current thrombolytics based on plasminogen activation. The final selection

resulted in an alteplase variant, denoted Brnoteplase, which exhibited favorable therapeutic

properties, making it a promising candidate for clinical studies and a valuable starting point for

follow-up improvements.

Figure 1: Schematic illustration of the research workflow established in this study to develop a novel
improved thrombolytic. The pipeline involves (i) *in silico* rational design of new thrombolytic protein sequences,
(ii) their biochemical characterization, (iii) testing in *in vitro* thrombolysis and penetration assays, and (iv)
validation in *in vivo* animal rat models. The numbers in the circles correspond to the number of thrombolytic
protein variants advanced to each step (grey rings) of the workflow, enabling initial screening of more than
10,000 variants and their comprehensive filtering down to one selected thrombolytic to be verified in resource-
demanding preclinical *in vivo* models.

Methods

Computer-aided selection of mutations alleviating binding of tPA to LRP1

Computational design of alteplase variants with reduced binding to the LRP1 receptor was performed using (i) structure-based and (ii) sequence-based strategies.

In the structure-based strategy, we performed conservation and correlation analysis of lysine residues on the first four domains of alteplase, which are responsible for LRP1 binding, using the HotSpot Wizard web server[18]. Afterwards, we docked the structure of LRP1 complement-like binding repeat to tissue-type plasminogen activator (tPA) using ClusPro and RosettaDock[19]. We clustered the docked structures and evaluated distances of conserved and exposed lysines of tPA to the lysine-binding site of LRP1. Lastly, we have performed molecular dynamics (MD) of tPA and evaluated which pairs of lysines are in the required distance of 21 to 29 Å for binding LRP1[20]. We designed mutant tPAs lacking specific lysines. These lysines fulfil the following criteria: (i) conserved in homologous sequences, (ii) exposed on the protein surface, (iii) located on the first four domains of tPA responsible for LRP1 binding, (iv) located on the interface with LRP1 complement-binding repeat in cluster centres of docked structures, and (v) being within 21 – 29 Å of another lysine residue in tPA during MD simulations.

In the sequence-based strategy, we compared the tPA sequence to its homolog, desmoteplase, which competitively binds LRP1[21]. In the pairwise sequence alignment, we identified lysines in both tPA and desmoteplase at homologous positions. These lysines are likely candidates to mediate LRP1 binding.

Lysine positions identified by structure- and sequence-based strategies were subjected to *in silico* saturation mutagenesis using the Rosetta ddg_monomer application. Mutated residues that retained thermodynamical stability were selected for experimental characterization.

Combining beneficial tPA mutations from literature:

To identify mutations for enhancing the efficacy and safety of tPA, we performed a comprehensive literature review on the structural biology and protein engineering

of thrombolytics. We then sought an innovative combination of mutations to improve tPA's

efficacy and safety and combined them into a multiple mutant for experimental

characterization.

Ancestral sequence reconstruction of tPA-related proteins

The sequence of tPA, including the signal peptide, was submitted to the FireProt[ASR] web server[22].

The calculation was performed using default settings. Ancestral sequences for experimental

characterization were selected using the following criteria: (i) as few insertions or deletions

compared to tPA as possible, (ii) preserved catalytic residues, and (iii) preserved functionally

important residues known to influence different tPA functions.

Mining novel enzymes from sequence databases

EnzymeMiner was used to search for novel thrombolytic enzymes in protein sequence

databases[23]. Two separate runs were performed using tPA and desmoteplase as query sequences.

Proteins with transmembrane regions or mutated catalytic residues were discarded. Moreover,

proteins had to contain key domains for plasminogen activation and fibrin binding and score a

predicted solubility higher than the query sequence. After filtering for these characteristics,

proteins with interesting properties were selected manually.

Construction of mutant gene sequences

For each designed thrombolytic variant, a synthetic gene (Genscript) was cloned into the

expression vector pcDNA3.1 (Thermo Fisher Scientific). *E. coli* strain TOP10F (Thermo Fisher

Scientific) was transformed with the plasmid construct. Plasmid DNA was isolated for transient

transfection using Plasmid Giga Kit (Qiagen).

Heterologous gene expression and protein isolation

HEK-293E cells were adapted to suspension growth in HyClone™ CDM4HEK293 serum-free medium

(Cytiva) supplemented with 4 mM L-glutamine (AppliChem). Prior to transient transfection, HEK-

293E cells were centrifuged at 250 g for 10 minutes and resuspended at the density of 2×10^7

cells/ml in the Gibco™ FreeStyle™ 293 expression medium (Thermo Fisher Scientific). HEK-293E

cells were transiently transfected by plasmid DNA and linear 25-kDa polyethyleneimine

(Polysciences) at a 1:2 ratio and a final concentration of 25 and 50 ug/mL, respectively. Plasmid

pmaxGFP at 5 % of the total DNA content was added as a positive control of the transfection

process. Four hours after transfection, cells were diluted with HyClone medium containing 2 mM

L-glutamine. The cells were cultured at 37 °C and 130 rpm in an incubator shaker (New Brunswick

Innova® 44/44R). The culture supernatants were harvested 7 days post-transfection. The

supernatant was purified on Ni-NTA column (Protino, Macherey-Nagel) connected to the BioLogic

DuoFlow FPLC system (Bio-Rad) and equilibrated with the purification buffer (10 mM Tris-HCl pH

7.5, 200 mM NaCl). The thrombolytic protein bound to the column was washed and eluted with

10 mM and 250 mM imidazole in buffer 10 mM Tris-HCl buffer pH 7.5 with 200 mM NaCl and 200

mM L-Arginine. Subsequently, the protein was purified on the Superose 6 SEC column (GE

Healthcare) equilibrated with PBS buffer pH 7.4, 200 mM L-Arginine. The protein was

supplemented with 0.01 % of Tween 80 and stored at 4 °C.

Protein thermostability measurement

Protein thermostability was determined by analyzing changes in protein intrinsic tryptophan

fluorescence with increasing temperature using the nanoscale differential scanning fluorimetry

(nanoDSF) instrument Prometheus NT.48 (NanoTemper Technologies). Each measurement was

performed in triplicate in phosphate-buffered saline PBS pH 7.4 (10 mM Na_2HPO_4, 1.8 mM KH_2PO_4,

2.7 mM KCl, 137 mM NaCl) containing 3.5 % L-arginine, and 0.01 % Tween 80. The fluorescence

was monitored at temperatures ranging from 20 °C to 100 °C with a temperature ramp of 1

°C.min^{-1} and the values of the fluorescence ratio at 350/330 nm were plotted against

temperature. The values of onset temperature (T_{onset}) and melting temperature (T_m) were

determined using the manufacturer-provided software after deriving the obtained fluorescence

curves. The final values were expressed as averages of individual replicates and their respective

standard deviations.

Determination of enzymatic activity and fibrin(ogen) stimulation and selectivity

The activities of thrombolytic enzymes were measured using the fluorogenic substrate D-VLK-AMC (AAT Bioquest), which is converted by the generated plasmin molecules to the fluorescent product 7-amino-4-methylcoumarin (AMC), whose fluorescence was monitored. All the experiments were performed at 37 °C in phosphate-buffered saline PBS pH 7.4 (10 mM Na_2HPO_4, 1.8 mM KH_2PO_4, 2.7 mM KCl, 137 mM NaCl) containing 1 mM $CaCl_2$, 0.0035 % L-arginine, and 0.01 % Tween 80 and each measurement was repeated in three replicates. The activities were determined in the absence but also in the presence of fibrinogen or fibrin that served as stimulation agents. In the case of fibrin stimulation, small fibrin clots were formed in a microplate well of a black clear bottom 96-well plate by mixing 8 μL of human fibrinogen (HYPHEN BioMed) with 2 μL of human thrombin (Sigma-Aldrich). The wells were sealed with an adhesive foil to prevent evaporation and kept at room temperature for an hour to ensure complete polymerization and clot formation. In the case of fibrinogen stimulation, the microplate wells were filled with 8 μL of human fibrinogen mixed with 2 μL of the reaction buffer only. In the absence of any stimulant, only 10 μL of the reaction buffer was injected. After that, all the microplate wells were added 50 μL of human plasminogen (Roche Diagnostics), 30 μL of D-VLK-AMC, and 10 μL of the tested thrombolytic to initiate the reaction. The resulting concentrations in microplate wells were 3.0 μM fibrin (where applicable), 0.04 UN/mL thrombin (where applicable), 3.0 μM fibrinogen (where applicable), 4.5 μM plasminogen, 200 μM D-VLK-AMC, and 0.08 nM tested thrombolytic. The microplate containing all the tested conditions was then sealed with a transparent adhesive foil and the enzymatic activity was measured by monitoring the increase in fluorescence with time using the microplate reader FLUOstar OPTIMA (BMG Labtech). The emission was detected every 60 seconds at 460 nm upon excitation at 360 nm with the instrument gain set to 1,100 and the bottom reading position.

Activity and stimulation kinetic data analysis

All the curves of increasing AMC fluorescence with time were first recalculated to AMC concentration changes with time based on the calibration curve that was obtained by measuring fluorescence of various increasing concentrations of the AMC standard (Sigma-Aldrich) ranging

200 from 0 to 1000 µM. The concentration dependence exhibited an exponential trend, so the

201 calibration curve was constructed by fitting the calibration points with the exponential **Equation**

202 **1** to obtain the calibration factors F_a, F_b, and F_c.

$$\text{Fluorescence} = F_a \cdot (1 - e^{-F_b \cdot [\text{AMC}]}) + F_c \tag{1}$$

Due to the coupled nature of the applied assay, the initial phase of each recalculated kinetic

curve was fitted with the quadratic **Equation 2**, where v_0 corresponds to the determined initial

rate of the enzymatic reaction, t_0 to the dead-time between the manual mixing of the assay

components and the start of the measurement, and c to the initial y-axis offset. Final values of

initial rates (activities) were expressed as average ± standard deviation for each measured

sample and condition.

$$[\text{AMC}] = v_0 \cdot (t + t_0)^2 + c \tag{2}$$

Stimulation factors and selectivities were determined according to the previously postulated

definitions[24]. Stimulation factor for the fibrinogen stimulation ($Stim_{FG}$) or the fibrin stimulation

($Stim_{FN}$) was calculated as a ratio of the activity in the presence of the stimulant ($v_{0,FG}$ or $v_{0,FN}$)

over the activity in the absence of the stimulant (v_0), as summarized in **Equation 3 and 4**.

Selectivity (Sel_{FN}) towards fibrin (with respect to fibrinogen) was then calculated as a ratio of

stimulation factors or initial rates in the presence of fibrin over fibrinogen according to **Equation**

5. The resulting value was relativized to alteplase and statistically compared using two-sample

t-test.

$$Stim_{FG} = \frac{v_{0,FG}}{v_0} \tag{3}$$

$$Stim_{FN} = \frac{v_{0,FN}}{v_0} \tag{4}$$

$$Sel_{FN} = \frac{Stim_{FN}}{Stim_{FG}} = \frac{v_{0,FN}}{v_{0,FG}} \tag{5}$$

Plasminogen activator inhibitor 1 (PAI-1) inhibition resistance

The potency of thrombolytic inhibition was analyzed by determining the fraction of active thrombolytics at different PAI-1 concentrations. The experiment was performed at 37 °C in phosphate-buffered saline PBS pH 7.4 (10 mM Na_2HPO_4, 1.8 mM KH_2PO_4, 2.7 mM KCl, 137 mM NaCl) containing 1 mM $CaCl_2$, 0.0035 % L-arginine, and 0.01 % Tween 80, and each measurement was repeated in three replicates. 40 µL of PAI-1 (Sigma-Aldrich) was mixed with 10 µL of a tested thrombolytic and the sealed mixture was incubated at 37 °C for 60 min while shaking at 180 rpm to ensure that the interaction with PAI-1 reached equilibrium. The resulting concentrations in the inhibition mixture were 0–1.5 µM PAI-1 and 0.16 µM thrombolytic. Residual activity was determined by transferring 10 µL of the inhibition mixture into a microplate well of a black clear bottomed 96-well plate and mixing it with 40 µL of the tPA fluorogenic substrate Boc-L-(p-F)FPR-ANSN-H-C_2H_5 (US Biological) to initiate the enzymatic reaction. The resulting fluorogenic substrate concentration was 200 µM. The microplate was sealed with a transparent adhesive foil and the enzymatic activity was detected by monitoring the increase in fluorescence over time using the microplate reader FLUOstar Omega (BMG Labtech). The emission was detected every 60 seconds at 460 nm upon excitation at 360 nm with the instrument gain set to 580 and the bottom reading position.

Inhibition resistance data analysis

The initial phase of each fluorescence kinetic curve was fitted with the linear **Equation 6** where the slope v_i corresponds to the initial rate of the inhibited enzymatic reaction (residual activity) and F_0 to the initial fluorescence offset. Since the fluorogenic substrate was converted directly by the tested thrombolytic enzymes with no reaction coupling and since the fluorescence of the generated product was directly proportional to its concentration, fluorescence curves could be fitted directly with the linear equation to reliably obtain relative values of initial rates without a need for recalculation to concentration changes. Final values of initial rates were expressed as average ± standard deviation for each measured sample and condition.

$$\text{Fluorescence} = v_i \cdot t + F_0 \tag{6}$$

Concentration dependence of determined residual activities v_i with increasing PAI-1

concentration was fit with **Equation 7** for a tightly bound inhibitor where c_I corresponds to the

total concentration of PAI-1, c_E to the total concentration of a thrombolytic enzyme, v_i to the

residual activity at a given concentration c_I of PAI-1, v_0 to the non-inhibited activity in the

absence of PAI-1, and IC_{50} to the concentration of PAI-1 causing 50 % inhibition of a tested

thrombolytic. The values of IC_{50} were determined for each tested thrombolytic, relativized to

the value for alteplase, and statistically compared using a two-sample t-test.

$$c_I = c_E \cdot \left(1 - \frac{v_i}{v_0}\right) + \left(IC_{50} - \frac{c_E}{2}\right) \cdot \left(\frac{v_0}{v_i} - 1\right) \tag{7}$$

Fibrin binding and affinity

Affinities of thrombolytics towards fibrin were analyzed by indirectly determining the level of

fibrin-bound thrombolytics at various fibrin concentrations. The experiment was performed at

37 °C in phosphate-buffed saline PBS pH 7.4 (10 mM Na_2HPO_4, 1.8 mM KH_2PO_4, 2.7 mM KCl, 137

mM NaCl) containing 1 mM $CaCl_2$, 0.0035 % L-arginine, and 0.01 % Tween 80 and each

measurement was repeated in three replicates. 320 μL of plasminogen-free human fibrinogen

(HYPHEN BioMed) was premixed with 40 μL of a tested thrombolytic and the clotting was initiated

by the addition of 40 μL of human thrombin (Sigma-Aldrich). The resulting concentrations in the

polymerization mixture were 0–38 μM fibrinogen, 16 nM (alteplase, Alt01, Alt02, Alt03, AltAnc02,

AltEM03, DesEM01) or 235 nM (tenecteplase, Brnoteplase, DesEM02, desmoteplase) thrombolytic,

and 2 UN/mL thrombin. This mixture was incubated at 37 °C for 60 min while shaking at 180 rpm

to ensure that a complete clot formation was achieved. The clots with bound thrombolytics were

separated by centrifugation at 4 °C and 15,000 rpm for 30 minutes and the amount of unbound

thrombolytic in the supernatant was determined by residual activity measurement. 20 μL of the

supernatant was carefully transferred into a microplate well of a black clear bottom 96-well

plate and mixed with 50 μL of human plasminogen (Roche Diagnostics) and 30 μL of the

fluorogenic substrate D-VLK-AMC (AAT Bioquest) to initiate the enzymatic reaction. The resulting

concentrations in a microplate well were 2.3 μM plasminogen and 200 μM D-VLK-AMC. The

microplate was sealed with a transparent adhesive foil and the enzymatic activity was detected

by monitoring an increase of fluorescence with time using the microplate reader FLUOstar

OPTIMA (BMG Labtech). The emission was detected every 60 seconds at 460 nm upon excitation

at 360 nm with the instrument gain set to 1,100 and the bottom reading position.

Fibrin binding equilibrium data analysis

The values of residual plasminogen activation initial rates (activities) $v_{0,res}$, corresponding to the

relative amount of an unbound thrombolytic agent, were determined by fitting the kinetic curves

as described above in the corresponding 'Activity and stimulation kinetic data analysis' section.

These values were then relativized to the initial rate in the absence of fibrin ($v_{0,abs}$) according to

Equation 8 in order to obtain a fraction of bound plasminogen activator (*Bind*) at a given

concentration of fibrin. Concentration dependence of the determined bound fraction *Bind* with

increasing fibrin concentration was fit with the hyperbolic **Equation 9** to obtain the value of

dissociation constant K_d and the percentage of the maximal fibrin binding $Bind_{lim}$. The values

were determined for each tested thrombolytic, relativized to the value for alteplase, and

statistically compared using a two-sample *t*-test.

$$\text{Bind} = \frac{v_{0,abs} - v_{0,res}}{v_{0,abs}} \cdot 100\ \% \tag{8}$$

$$\text{Bind} = \frac{Bind_{lim} \cdot [\text{fibrin}]}{K_d + [\text{fibrin}]} \tag{9}$$

Preparation of clots and fibrin gel for *in vitro* assays

Semi-synthetic clots were prepared by mixing human fibrinogen (Tisseel Lyo kit, Baxter, dissolved

in water for injection to reach the concentration of 52 mg/mL), human thrombin (Tisseel Lyo

kit, Baxter, dissolved in 40 µM CaCl$_2$ solution to the concentration of 3.3 IU/mL), and red blood

cells (RBCs) isolated from healthy donors' blood in the volume ratio of 1:1:1. The incubation and

clotting were performed in plastic Eppendorf tubes (0.5 mL for static model; 0.2 mL for flow

model) for 30 min at 37 °C and >90 % relative humidity under static conditions. The prepared

clots were immediately introduced into the model. The final concentrations of components in

the prepared clots were 17.3 mg/mL fibrinogen, 1.1 IU/mL thrombin, and $4.2 \cdot 10^9$ RBCs/mL and

the total clot volume was 90 μL for the static model and 60 μL for the flow model.

RBC dominant clots were prepared from 200 μL of healthy donors' whole blood without

anticoagulant by clotting in borax glass tubes (inner diameter 8 mm) for 3.5 h at 37 °C and >90

% relative humidity under static conditions. The prepared clots were immediately introduced

into the model.

Fibrin gel was prepared from the mixture of fibrinogen (Tisseel Lyo kit, Baxter, dissolved in water

for injection to the concentration of 34.6 mg/mL) and thrombin (Tisseel Lyo kit, Baxter, dissolved

in 40 μM $CaCl_2$ solution to the concentration of 2.2 IU/mL) in the volume ratio of 1:1. The

incubation and the clotting was performed inside micro-slide channels for 30 min at 37 °C and

>90 % relative humidity under static conditions. The final concentrations of the components in

the prepared fibrin gel were 17.3 mg/mL fibrinogen and 1.1 IU/mL thrombin and the total volume

per micro-slide channel was 30 μL.

All blood donors had agreed to donate blood samples on the premise of signed informed consent

for the collection of blood. Individuals, who had received acetylsalicylic acid, non-steroidal anti-

inflammatory, or antiplatelet drugs within 7 days before blood collection, were excluded.

In vitro **thrombolytic models**

Thrombolytic experiments were performed in static and flow *in vitro* models that were described

in contemporary literature[25,26] but were optimized to determine the suitably measurable effect

of alteplase in a highly reproducible manner, as reported in previous publications[27-30]. Briefly,

the static model consisted of 1.5 mL plastic Eppendorf tubes filled with fresh frozen human

plasma (177RM, Monobind Inc.) medium to a total volume of 500 μL, in which the clots were

individually incubated. The tubes were incubated at 37 °C in a dry-block incubator for 60 minutes

(the same amount of time as indicated for alteplase treatment of stroke patients).

The flow model consisted of silicone chips made of Sylgard 184 Silicone Elastomer (Dow Corning),

which were prepared according to human middle cerebral artery anatomy, with narrowings

dimensionally based on patient CT scans (n = 4). The bifurcation was included to enable

permanent circulation in the system. Each silicone chip was connected by plastic pipes (internal

diameter 3.1 mm) to the 8-channel pump head peristaltic pump MINIPULS 3 (Gilson Inc.), and

the entire system was maintained at 37 °C for incubation lasting 180 min. This arrangement

maintained the hydrodynamic forces involved in clot removal[31], whereby a pressure gradient of

10 mm Hg was generated across the occlusion.

Both semi-synthetic and red blood cell dominant clots (preparation described in the

corresponding 'Preparation of clots and fibrin gel for in vitro assays' section above) were

separately employed in both thrombolytic models and the thrombolytic efficiencies were

compared. The final concentrations of tested thrombolytic enzymes were selected to be

consistent with the clinically relevant dosage indicated for patients with ischemic stroke (1.3

μg/mL), according to the manufacturer's instructions and supporting pharmacokinetic data[32].

Alternatively, 10-fold lower and 2-, 5-, 10-, and 50-fold higher concentrations were tested as

well.

In vitro **lytic efficacy analysis**

Clot lysis in the static model was determined by measuring clot mass loss[26-28,30,33] and by

spectrophotometric determination of RBC release into the incubation medium at 575 nm[27,28,30].

Clot lysis in the flow model was determined by measuring recanalization time, clot length

changes, and spectrophotometric determination of RBC release[27,29,30]. Recanalization frequency

was determined as a percentage ratio of complete recanalization to the total number of samples

in the respective treatment group. Relative clot reduction was determined as a percentage of

clot area reduction prior to clot release from the narrowed occlusion site. In addition, the

thrombolysis rate was calculated using linear regression of relative clot reduction at 10-minute

intervals. Data were expressed as mean ± standard deviation, if not otherwise indicated. The

generated box plots were presented as mean value (cross), median (line), interquartile range

(box), and minimum and maximum value (whiskers). ANOVA with post-hoc Tukey's HSD test was

used to compare data. P-values ≤ 0.05 were considered to be statistically significant. All analyses

were performed using GraphPad Prism (GraphPad Software Inc.).

In vitro penetration microarray

Penetration experiments were performed in the microarray model outlined in recent literature[34-38] and optimized to provide a suitably measurable effect of alteplase in a highly repeatable manner. The microarray consisted of the microslide µ-Slide VI 0.4 (ibidi GmbH) containing channels that were filled with a prepared fibrin gel (procedure described in the corresponding 'Preparation of clots and fibrin gel for in vitro assays' section above). Fluorescently labelled plasminogen activators or bovine serum albumin (A9418, Sigma-Aldrich), as a relevant fibrin non-interacting control, were applied to individual microslide channels and fluorescence intensities were continuously recorded in 5 min intervals for 180 min, at 0.25 mm distance intervals up to 1.00 mm from the application site. The fluorescence signal was recorded using the microscope ZEISS Axio Observer Z1 (Carl Zeiss AG) equipped with the metal halide fluorescence light source HXP 120 (Carl Zeiss AG) and the high efficiency Filter Set 43 HE (Carl Zeiss AG). The detection settings were as follows: excitation BP 550/25; beam splitter FT 570; emission BP 605/70. Thrombolytic proteins and bovine serum albumin control were fluorescently labelled with rhodamine B isothiocyanate (283924, Sigma-Aldrich) prior to the penetration assay according to the standard labeling protocol provided by the manufacturer. The final concentrations of the labelled proteins were the same as for thrombolytic experiments, i.e., 1.3 µg/mL. Thrombolytic efficiencies of labelled plasminogen activators were verified in the static model.

Penetration rate analysis

Plasminogen activator penetration rate through the fibrin gel was determined by continuous measurements of fluorescence intensities at a distance of up to 1 mm from the front of the plasminogen activator application site[30]. The penetration rate was expressed as the relative fluorescence intensity (RFI) at a given distance and time. RFI was determined using a custom image analysis script in the Python programming language. Penetration rate data were further normalized to relevant controls. Data were expressed as mean ± standard deviation, if not otherwise indicated. The generated box plots were presented as mean value (cross), median (line), interquartile range (box), and minimum and maximum value (whiskers). ANOVA with post-hoc Tukey's HSD test was used to compare data. P-values ≤ 0.05 were considered to be

statistically significant. All analyses were performed using GraphPad Prism (GraphPad Software

Inc.).

Preparation of clots for *in vivo* assays

Details of the characteristics and nature of artificial clots are the subject of a previous project[39], and their preparation is therefore described only briefly. Commercial human fibrin glue Tisseel kit (Baxter) was used to generate artificial clots for *in vivo* experiments. Saline solution (1.5 mL) and $BaSO_4$ suspension (0.5 mL) (Micropaque, Guerbet) were injected into the original "TISSEEL" ampoule. 1 mL of the original calcium chloride solution and 5 mL of bovine serum were injected into the "Thrombin 500 I.U." original ampoule. Finally, "TISSEEL" and "Thrombin 500 I.U." preparations were injected in a 1:1 ratio into a PE60 and PE10 tubes (Braintree Scientific) for the thrombolytic effectivity study and the safety study, respectively, using the original Duploject system (Baxter). Both ends of the cannula containing clot components were hermetically sealed to prevent any evaporation or desiccation, and the mixture was incubated at 37 °C for 120 min, followed by storage at 4–8 °C for additional 24 hours. The resulting artificial clot was cut into 10 mm fragments for the effectivity study and 1 mm fragments for the safety study and the corresponding samples were stored in the saline solution in an Eppendorf tube at −20 °C.

Thrombolytic effectivity study *in vivo*

Adult male outbreed Wistar rats (n=68; body weight 120–180 g) were selected for the *in vivo* lysis experiments and were randomized into groups with different treatment regimens (variable thrombolytic types, doses, and administration routes) as described in **Table 1**.

Table 1: **Overview of animal experimental groups for in vivo effectivity study based on different treatment regimes.** CRI = continuous rate infusion.

Experimental group	Number of animals in the group
Control group (saline solution)	10
Alteplase (0.9 mg/kg, CRI)	16
Tenecteplase (0.25 mg/kg, bolus)	12
Brnoteplase (0.9 mg/kg, CRI)	4
Brnoteplase (0.9 mg/kg, bolus)	12
Brnoteplase (2.5 mg/kg, bolus)	12
Brnoteplase (5.0 mg/kg, bolus)	2

The exclusion criterion for the experiment was death during the anesthesia or the surgical

procedures before the clot delivery. Throughout the thrombolysis experiment, rats were under

general anesthesia induced by inhalation of 2.5 % isoflurane in filtered room air as a carrier gas.

After the intramuscular application of xylazine and ketamine (5 mg/kg and 35 mg/kg of body

weight, respectively), a mixture of urethane (80 mg/mL) and alpha-chloralose (6 mg/mL) at a

dose of 8 mL/kg of body weight was applied into the peritoneal cavity. Subsequently, the left

common carotid artery (CCA) was dissected, visualized, and cannulated in a retrograde direction

with a PE50 (Braintree Scientific) cannula.

Three individual barium-labeled artificial clots were injected via the PE50 catheter cannula into

the aortic arch of each tested rat. Micro-fluoroscopy using the C-arm LabScope system

(Glenbrook Technology, Inc.) was used to verify the localization of the introduced clots in the

branches of the abdominal aorta and the position of the animals was then adjusted to achieve

the optimal visibility of the clots with the maximum area of their size and without the shielding

of bone structures. The initial image was captured at time zero (before the thrombolytic drug

administration). Subsequently, a bolus injection/continuous rate infusion of a tested

thrombolytic was initiated. The continuous rate infusion treatment consisted of an initial bolus

of 10 % of the total dose volume, followed by an infusion of the remaining 90 % volume for 60

minutes. In the bolus treatment groups, the entire volume of the thrombolytic agent was

administered at once. The control group received an equivalent volume of intravenous saline

solution. The therapy was monitored for 60 minutes by the C-arm micro-fluoroscopy and

radiographs were acquired and stored every 5 minutes.

In vivo **thrombolytic effectivity study outcome analysis**

The outcome was measured as a relative area change of each monitored artificial clot, and the

lysis rate was then calculated as a linear fit of relative area change over time. All collected

radiographs were analyzed using the ImageJ 1.52a software (Wayne Rasband, National Institutes

of Health), and the area of identified clots in pixels was manually determined at each time point.

All images of one animal were measured simultaneously in one reading sequence. Two

independent observers performed the image analysis, blinded to one another's results and to

animal group assignment. Anatomical determination of arteries with clots was performed after

all measurements. A standardized protocol was established and followed for non-standard clot

positions or movements: (1) If the area of two clots in different arteries overlapped, the size was

measured as a single clot. (2) If the clot changed position due to animal/intestinal movement or

projection angle, the area of the clot was recalculated to the adjusted change of the nearest

bone shadow. The image was excluded if no bone was present. (3) If the lysis of the clot caused

its movement but the clot was still visible on the image, the shadow area of the clot was included

in the original area of the clot. The image was excluded if the clot moved out of the image.

The change in the artificial clot area (CAC) was calculated from the clot area at a calculated

time (a_t) relative to the clot area at time zero baseline before thrombolytics administration (a_0),

i.e., at the beginning of the experiment, based on **Equation 10**. The lysis rate in $\%.min^{-1}$

represents a relative change in the area of the artificial clot (CAC) divided by the minute in which

the area was measured (t) based on **Equation 11**. Results of the thrombolysis were used to

generate the thrombolytic curve. The area under the thrombolytic curve (AUC) was calculated

and used as a comparative parameter among the tested and control groups. All analyses were

performed using Excel for Microsoft 365 MSO version 2405 (Microsoft Corporation) and GraphPad

Prism version 8.0.0 (GraphPad Software Inc.).

$$CAC = \frac{a_0 - a_t}{a_0} \cdot 100 \ \% \tag{10}$$

$$\text{lysis rate} = \frac{CAC}{t} \tag{11}$$

Safety study

The safety study was based on a thromboembolic Middle Cerebral Artery Occlusion (MCAO)

model, utilizing the same clot type as in the thrombolytic effectivity study. However,

embolization to all major cerebral arteries, namely the anterior, middle, and posterior cerebral

arteries, was included. This approach is referred to as the **C**ircle **o**f **W**illis **B**ranches **O**cclusion -

CoWBOi model. Male outbred Wistar rats, weighing between 480 and 980 grams and aged 7 to 9

months, were housed individually under standard conditions, with a natural light/dark cycle and

unrestricted access to food and water. Animals were randomly assigned to one of three

experimental groups, receiving either alteplase (0.9 mg/kg, continuous rate infusion, n=25,

included n=22), tenecteplase (0.25 mg/kg, bolus, n=25, included n=24), or Brnoteplase (2.5

mg/kg, bolus, n=23, included n=22). Rats, that did not develop cerebral stroke or died within

120 minutes of thrombolytic treatment, were excluded from the study.

General anesthesia was induced via inhalation of 2.5 % isoflurane, followed by intramuscular

administration of xylazine (5 mg/kg), ketamine (35 mg/kg), and intraperitoneal diazepam (2

mg/kg). Embolization was performed in a modified version of the standard MCAO model, with

cannulation of the left external, and internal carotid arteries. Two embolic clots (1.5 mm × 0.3

mm) were introduced into the intracranial internal carotid artery. Thrombolytic treatment was

initiated 4 hours post-embolization. After 24 hours, the animals were euthanized, and their

brains were collected in cold saline without flushing. As described previously, the extracted

brains were fixed in formalin for 24 hours and stained with iodine[40] . In contrast with the original

set-up, the brains were not examined in agarose but in paraffin oil. Micro-computed tomography

(miCT) imaging was performed using a SkyScan1276 CMOS (Bruker) scanner, and the acquired

scans were analyzed using the Amira software (Thermo Scientific). Examiners were blinded to

group assignments to ensure unbiased assessment.

In vivo **safety study outcome analysis**

To evaluate hemorrhagic transformation (HT), a classification system based on ECASS II criteria[11]

was applied. Hemorrhagic transformation was categorized into four subtypes (HI1, HI2, PH1, and

PH2) according to the criteria listed in **Table 2**. Considering that thrombolytic enzymes were

administered 4 hours after clot embolization, by which time all collateral vessels had already

closed due to ischemic damage, successful recanalization in this model was consistently

associated with hemorrhagic transformation.

Table 2: Hemorrhagic transformation (HT) classification based on the ECAS II study. The classification recognizes four HT subtypes (HI1, HI2, PH1, and PH2) that differ in the listed criteria.

Grade	Description
HI1	Petechial hemorrhages at infarct margins
HI2	Petechial hemorrhages throughout the infarct without mass effect
PH1	Hemorrhage affecting ≤30 % of the infarcted area with minor mass effect
PH2	Hemorrhage affecting >30 % of the infarcted area with substantial mass effect

To quantify hemispheric asymmetry (HA), we calculated the affected hemisphere volume (*affected*) and the intact hemisphere volume (*intact*), excluding the olfactory bulbs and cerebellum from the analysis. The percentage of HA was then derived using **Equation 12**. A hemispheric asymmetry of 6 % was identified as the threshold distinguishing normal from pathological brains. Values exceeding this threshold were consistently associated with the presence of cerebral edema.

$$HA = \frac{affected - intact}{intact} \cdot 100\ \% \tag{12}$$

Quantification of residual plasminogen, plasmin, and thrombolytic levels in rat plasma

At the end of the *in vivo* thrombolytic effectivity study, whole blood from the common carotid was collected to a K3-EDTA tube (2.7 mL, S-Monovette, Sarstedt AG). The collected blood was centrifuged for 15 min at 1,500 rpm and the resulting plasma was separated into an Eppendorf tube, aliquoted, and stored at -20 °C. The collected plasma samples were tested for levels of plasmin, plasminogen, and a thrombolytic enzyme using ELISA Rat Plasmin Kit (MyBioSource), ELISA Rat Plasminogen Kit (MyBioSource), and ELISA Human Active tPA Kit (Molecular Innovations), respectively. The determination of the concentration was performed according to the protocol provided by the manufacturer with no modifications. In the case of Human tPA Kit, purified solutions of alteplase and Brnoteplase (the same as administered in rats) with known concentrations were used to generate individual calibration curves to prevent any discrepancies caused by different sources of the administered and the calibration samples. The final plasma levels were calculated from the measured absorbance values based on the constructed calibration curves.

Consent for animals used in experiments

All procedures and animal experiments were performed in full compliance with the ARRIVE and

the European Community Council Directive (2010/63/EU) for Protection of Vertebrate Animals

Used for Experimental and other Scientific Purposes guidelines. The project was approved by the

Masaryk University Brno Institutional Committee for the Protection of Experimental Animals and

by the Committee for the Protection of Experimental Animals of the Ministry of Education Youth

and Sports in the Czech Republic (MEYS CZ) with the protocol number MSMT-35804 / 2019-2.

Results

Computational design selects the most promising thrombolytic candidates

The initial pool of novel thrombolytic variants for experimental testing was designed and selected

in silico using three independent computational strategies: (i) rational design, (ii) ancestral

sequence reconstruction, and (iii) database mining (**Table 3**, **Figure S1**).

Rational design focused on reducing interactions with clearance and signaling receptors,

particularly LRP1 and NMDAR, which are implicated in short half-life and neurotoxicity. This was

achieved by introducing Tyr67Asn and Ser69Ala mutations shown to prevent glycosylation

responsible for LRP1 binding[41] (resulting in variant Alt03), and by combining these mutations with

additional substitutions of lysine residues (variants Alt01 and Alt02). Lysine residues are known

to mediate LRP1 interaction. Alt01 includes additional Lys10Glu, Lys49Ile, and Lys152Gln

mutations which were selected using structure-based computational tools, namely protein-

protein docking, molecular dynamics, surface exposure analysis, and thermodynamic stability

predictions (**Supplementary Note 1**). Alt02 incorporates additional Lys82Ala, Lys159Ser, and

Lys162Ile mutations selected based on their sequence conservation in desmoteplase, which has

been shown to compete with alteplase for the LRP1 binding site[42] (**Supplementary Note 2**).

Finally, the Brnoteplase variant was constructed by combining the two original substitutions

preventing LRP1-binding glycosylation[41] (Tyr67Asn and Ser69Ala) with mutations decreasing PAI-

1 inhibition and increasing fibrin specificity[45] (Lys296Ala, His297Ala, Arg298Ala, Arg299Ala),

disabling NMDAR-mediated neurotoxicity[44] (Trp253Arg), and removing high-mannose
glycosylation responsible for the low alteplase biological half-life[43] (Asn117Gln).

Ancestral sequence reconstruction (ASR) strategy utilized the FireProt[ASR] webserver to
explore evolutionary and functional diversity while preserving thrombolytic activity. The analysis
produced a phylogenetic tree comprising 150 extant alteplase homologs and 149 ancestral
sequences (**Figure S2**). Two ancestral nodes close to alteplase, named AltAnc01 and AltAnc02,
were selected based on sequence identity and conservation of catalytic residues (**Table 3**).
AltAnc01 is ancestral to alteplase and four other extant sequences annotated as plasminogen
activators (PAs), while AltAnc02 is ancestral to alteplase, ten additional annotated PAs, and one
hypothetical protein predicted to be a PA (**Supplementary Note 3**).

Table 3: Summary of *in silico* designed and selected thrombolytics for experimental testing. Variants
highlighted in grey could not be experimentally produced and therefore were not further characterized. Full
sequences of all variants are provided in **Figure S1**. ASR = ancestral sequence reconstruction.

Variant naming	Modification strategy	Parental Protein	Sequence modification
Alt01	Rational design	Alteplase	K10E/K49I/Y67N/S69A/K152Q
Alt02	Rational design	Alteplase	Y67N/S69A/K82A/K159S/K162I
Alt03	Rational design	Alteplase	Y67N/S69A
Brnoteplase	Rational design	Alteplase	Y67N/S69A/N117Q/W253R/K296A/H297A/R298A/R299A
AltAnc01	ASR	Alteplase	76.6 % identity to alteplase; ancestral to 5 extant genes
AltAnc02	ASR	Alteplase	71.0 % identity to alteplase; ancestral to 12 extant genes
AltEM01	Database mining	Alteplase	78.4 % identity to alteplase; gene isoform X1 from bottlenose dolphin
AltEM02	Database mining	Alteplase	82.7 % identity to alteplase; gene isoform X2 from bottlenose dolphin
AltEM03	Database mining	Alteplase	92.0 % identity to alteplase; gene from Bolivian squirrel monkey
DesEM01	Database mining	Desmoteplase	68.0 % identity to alteplase; gene isoform X3 from Angolan colobus monkey
DesEM02	Database mining	Desmoteplase	80.0 % identity to alteplase; gene from white-winged vampire bat

535 **Database mining** using the EnzymeMiner webserver identified naturally occurring homologs
536 of alteplase and desmoteplase with potentially favorable biochemical features. Over 10,000
537 mined sequences were filtered based on predicted solubility, absence of transmembrane regions,

and preservation of essential catalytic domains and residues, resulting in 198 alteplase and 615 desmoteplase homologs meeting these criteria. From these, we manually selected five candidates (**Table 3**): three homologous to alteplase (AltEM01–03) and two to desmoteplase (DesEM01–02). The selected sequences originate from bottlenose dolphin, Bolivian squirrel monkey, Angolan colobus, and vampire bat (**Supplementary Note 4**).

Biochemical characterization identifies an improved alteplase variant Brnoteplase

Gene sequences encoding putative and mutant plasminogen activators selected *in silico* were experimentally constructed, and 8 out of 11 selected thrombolytic proteins were successfully produced recombinantly. Our comprehensive biochemical characterization (**Figure 2**, **Table S1**) revealed that all the proteins exhibited clear thrombolytic activity (**Figure 2a**, **Figure S3**, **Table S1**), validating the robustness of our computational approach for accurately predicting and designing plasminogen activator enzymes. Thermal stabilities of all characterized proteins (**Figure 2d**, **Figure S4**, **Table S1**) were highly above the physiological temperature, allowing their administration to the human body without a risk of rapid thermal inactivation.

Fibrinogen and fibrin stimulation analysis (**Figure 2b**, **Figure S3**, **Table S1**) yielded unimpaired profiles for Alt01–03 compared to alteplase (p=0.37, p=0.35, p=0.49, respectively), whereas other variants showed significantly lower fibrin stimulation (p<0.001) and activity (p<0.001), suggesting inefficient thrombolysis on fibrin clots. Remarkably, the variant Brnoteplase displayed minimal activity with no stimulation and when stimulated by fibrinogen, but it was massively stimulated by fibrin, 5.2-fold more than alteplase (p<0.001). This resulted in an extraordinary fibrin selectivity of Brnoteplase (**Figure 2c**, **Figure S3**, **Table S1**) that was similar to highly selective thrombolytics desmoteplase and tenecteplase.

All four sufficiently fibrin-active variants, Alt01–03 and Brnoteplase, further exhibited significantly increased resistance to PAI-1 inhibition (p<0.001) and reduced fibrin binding (p<0.001) compared to alteplase, indicating improved potential for clot penetration and lysis throughout the entire volume from inside. The highest inhibition resistance was observed for the variant Brnoteplase (**Figure 2e**, **Figures S5 and S6**, **Table S1**), while the most pronounced

reduction in fibrin binding was observed for Alt01, followed by Brnoteplase (**Figure 2f**, **Figures**

S7 and S8, **Table S1**).

In summary, the set of biochemical assays identified an exceptional alteplase variant,

Brnoteplase, with several favorable clinical implications over alteplase: (i) 80-fold enhanced

fibrin selectivity, (ii) 4-fold increased resistance to PAI-1 inhibition, and (iii) 4-fold improved

penetration potential.

Figure 2: Biochemical characterization of thrombolytic variants. (**a**) Comparison of plasminogen activation enzymatic activities (dark solid bars) and their values in the presence of fibrinogen (hatched bars) and fibrin (light solid bars) stimulants. (**b**) Comparison of fibrinogen (solid bars) and fibrin (hatched bars) stimulation factors. The factors are defined as fibrin(ogen)-stimulated activity over non-stimulated activity ratios. (**c**) Comparison of fibrin selectivity factors. (**d**) Comparison of melting temperatures T_m values, defining protein thermal stabilities. At T_m, half of the protein molecules are unfolded. (**e**) Relative comparison of inhibition resistance towards plasminogen activator inhibitor-1 (PAI-1) calculated as a ratio of the IC_{50} value for a tested protein over the IC_{50} value for alteplase (reference benchmark protein). (**f**) Relative comparison of fibrin penetration potential calculated as a ratio of the fibrin dissociation constant (K_d) value for a tested protein over the fibrin K_d value for alteplase (reference benchmark protein). Clinically used variants alteplase and tenecteplase are highlighted in blue and yellow, respectively. The variant Brnoteplase, which was selected for further *in vitro* and *in vivo* testing, is highlighted in green. Brnoteplase exhibited superior fibrin stimulation, fibrin selectivity, PAI-1 inhibition resistance, and penetration potential. Please note the logarithmic scale of the x-axis in (**a**), (**b**), and (**c**). The bars represent mean values and the whiskers correspond to standard errors.

586 *In vitro* **thrombolysis and penetration confirm enhanced Brnoteplase properties**

587 Alt01–03 and Brnoteplase, the four variants identified as highly fibrin-active in biochemical

588 assays, were subjected to thrombolysis testing on *in vitro* prepared human clots and compared

589 to alteplase and tenecteplase (**Figure 3**). Treatment with all the tested thrombolytics in the

590 static model containing semi-synthetic clots provided significantly greater clot mass loss

591 (p<0.001) and red blood cell (RBC) release (p<0.001) compared to the untreated group (**Figure**

592 **3a**, **Figure S9**, **Table S2**). The results of lysis efficiencies were consistent with the biochemical

593 characterization data (**Figure 2a**).

594
595 **Figure 3:** *In vitro* **analysis of thrombolysis efficacy and fibrin penetration.** (**a**) Comparison of thrombolytic
596 efficacy in the static model with semi-synthetic clots expressed as clot mass loss. (**b**) Concentration dependence
597 profile of thrombolytic efficacy with increasing applied concentration of a tested thrombolytic in the static
598 model. (**c**) Comparison of thrombolytic efficacy at multiple concentrations in the flow model expressed as red
599 blood cell release. (**d**) Comparison of relative penetration rates at multiple distances of a fibrin gel. Box plots
600 illustrate mean values (cross), median (line), interquartile range (box), and minimum/maximum values
601 (whiskers). Scatter plots illustrate mean values (circle) and standard errors (whiskers).

The concentration dependence experiment with alteplase and the most promising variant

Brnoteplase (**Figure 3b**, **Table S3**) revealed notable differences between the two tested

proteins. Alteplase showed no significant enhancement of thrombolysis with increasing

concentration, as clot mass loss at 2.6, 6.5, and 13 µg/mL was comparable to the clinically

relevant concentration of 1.3 µg/mL ($p=0.93$, $p>0.99$, $p=0.97$, respectively) and it even

decreased at the highest applied concentration of 65 µg/mL ($p=0.002$). In contrast, Brnoteplase

exhibited a clear trend of increasing thrombolytic efficacy with increasing dosing: clot mass loss

was comparable to the clinically relevant concentration only at 2.6 µg/mL ($p=0.96$) while

significantly greater lysis was observed at 6.5, 13, and 65 µg/mL ($p=0.02$, $p<0.001$, $p<0.001$,

respectively). At higher concentrations of 6.5 and 13 µg/mL, Brnoteplase reached the

comparable level of lysis as for alteplase ($p>0.99$, $p=0.20$, respectively), and it even surpassed

alteplase efficiency at the highest concentration of 65 µg/mL ($p=0.009$).

Similar trends were observed for RBC release analysis with semi-synthetic clots (**Figure S10a**,

Table S3) and RBC-dominant clots (**Figure S10bc**, **Table S4**), confirming the validity of the

measured data (details in **Supplementary Note 5**). In the flow model (**Figure 3c**, **Figure S11**,

Table S5), alteplase provided lower recanalization time ($p=0.005$), greater clot reduction rate

($p<0.001$), and greater RBC release ($p=0.006$) compared to the untreated group only at the

clinically relevant concentration (1.3 µg/mL), while higher concentrations yielded no significant

effect compared to the untreated group. In contrast, Brnoteplase was significantly more

effective than the untreated group across all tested concentrations (details in **Supplementary**

Note 6), supporting the concentration-dependent trend observed in the static model (**Figure**

3b).

No significant changes in Brnoteplase thrombolytic efficacy were detected even after 3

months of storage at 4 °C ($p=0.40$), as opposed to alteplase ($p=0.02$), concluding the high shelf-

life of Brnoteplase and its suitability for long-term storage applications (**Figure S12**, **Table S6**).

Fibrin gel penetration experiments (**Figure 3d, Table S7**) confirmed detectable interaction

with fibrin for both alteplase and Brnoteplase, as both exhibited significant changes in

penetration rates ($p<0.001$ at all distances) compared to the non-interacting control bovine

serum albumin (BSA). Comparison of alteplase and Brnoteplase showed similar penetration rates for both proteins up to the distance of 0.50 mm (p=0.07 at 0.00 mm; p=0.08 at 0.25 mm; p=0.06 at 0.50 mm). However, a significantly higher amount of Brnoteplase penetrated beyond 0.75 mm (p=0.008 at 0.75 mm; p<0.001 at 1.00 mm), confirming the superior clot penetration potential identified during biochemical characterization (**Figure 2f**).

Altogether, *in vitro* analysis confirmed the strong potential of Brnoteplase due to multiple additional advantageous properties over alteplase: (i) thrombolytic efficacy increase with concentration, (ii) preserved stability for over 3 months at 4 °C, and (iii) faster penetration rate and greater distribution within the clot. Combined with its enhanced fibrin selectivity (**Figure 2c**) and PAI-1 inhibition resistance (**Figure 2e**), the Brnoteplase profile was indicating improved safety, recanalization potential, and biological half-life, which would enable bolus administration and increased dosing in the subsequent *in vivo* testing.

Animal testing verifies increased Brnoteplase efficiency and safety *in vivo*

Our selected variant Brnoteplase and approved drugs alteplase and tenecteplase were tested *in vivo* for thrombolysis effectivity and safety in rat models (**Figure 4**). Thrombolytic effectivity was assessed by the lysis rate of administered thrombi and the area under the curve (AUC) of lytic curves. The Brnoteplase group (2.5 mg/kg, bolus) showed a lysis rate of 1.3 ± 0.4 %/min (**Figure 4a**, **Table S8**), which was the highest among groups when compared to 0.9 ± 0.5 %/min lysis rate for the alteplase (0.9 mg/kg, continuous rate infusion) and 1.1 ± 0.5 %/min for the tenecteplase (0.25 mg/kg, bolus) groups. Similarly, the AUC in Brnoteplase-treated rats was 2350 ± 720, significantly surpassing values observed in the alteplase (1520 ± 810) and tenecteplase (2180 ± 1060) groups (**Figure 4a**, **Table S8**). Statistical analysis confirmed that Brnoteplase demonstrated superior thrombolytic efficacy across all tested doses (0.9, 2.5, and 5.0 mg/kg; **Figure S13**, **Table S8**) by providing significantly higher lysis rates and AUC values than both control and alteplase groups (p ≤ 0.01). Interestingly, Brnoteplase (0.9 mg/kg) demonstrated activity at the level of untreated control when applied in the continuous rate infusion mode (**Table S8**).

The safety study assessed the overall incidence of hemorrhagic transformation (HT) and its

severity according to ECASS II criteria[46] to evaluate the safety profile of each thrombolytic agent

(**Figure 4b**, **Figure S14**, **Table S9**). Given the experimental model, in which the blood-brain

barrier was damaged and collaterals collapsed due to prolonged ischemia, bleeding could not

occur without recanalization[47], making overall HT incidence an indicator of successful

recanalization. HT occurred most frequently (86 %) in the Brnoteplase group (2.5 mg/kg),

followed by the 71 % incidence in the tenecteplase (0.25 mg/kg), and 64 % in the alteplase (0.9

mg/kg) groups, being positively correlated with recanalization frequencies (87 %, 68 %, and 56

%, respectively; **Figure S14**, **Table S9**). Among HT subtypes, Brnoteplase treatment led

predominantly to mild petechial hemorrhages (HI1), occurring in 60 % of treated rats, while only

15 % resulted in severe hemorrhages (PH1). These results substantially differed from both

alteplase (43 % HI1 vs. 21 % PH1) and tenecteplase (24 % HI1 vs. 35 % PH1) groups where the

proportion of severe PH1 was more prevalent (**Figure 4b**, **Table S9**).

Figure 4: *In vivo* **analysis of alteplase, tenecteplase, and Brnoteplase performance in animal rat models.** (**a**)
Comparison of the thrombolytic effectivity expressed as lysis rate (% of the clot dissolved per minute) and the
area under the curve (AUC) of the lytic curve. (**b**) Proportion of hemorrhagic transformation severities according
to ECASS II scoring system[46]. For a detailed description, see the Methods section. (**c**) Comparison of hemispheric
asymmetry of rat brains expressed as a relative change to an intact hemisphere. Box plots illustrate mean values
(cross), median (line), interquartile range (box), and minimum/maximum values (whishers). Bar plots illustrate
mean values (bars) and standard errors (whiskers).

Hemispheric asymmetry analysis (**Figure 4c**, **Table S9**) yielded the lowest value of 11 ± 8 %

for Brnoteplase-treated rats (2.5 mg/kg), followed by 13 ± 5 % for tenecteplase-treated (0.25

mg/kg) and 15 ± 11 % for alteplase-treated (0.9 mg/kg) groups. The lowest asymmetry in the

Brnoteplase group is coupled with the highest overall HT incidence, reflecting the highest

recanalization frequency (**Figure S14**), and the predominance of the least dangerous HT grade

(HI1). Together with the lowest observed proportion of severe HT (PH1; **Figure 4b**), this

outcome indicates that Brnoteplase treatment was associated with reduced brain swelling and

infarct expansion compared to the administration of alteplase or tenecteplase, suggesting a

potentially more favorable safety profile.

Rat plasma analysis corroborates superior Brnoteplase properties

Rat plasma collected at the end of the thrombolysis effectivity experiments was analyzed to

determine circulating levels of plasminogen, plasmin, and the administered thrombolytic (**Figure**

5, **Table S10**). Residual thrombolytic concentrations provided information about residence time

and half-life (**Figure 5a**, **Table S10**). The results yielded 37-, 11-, and 30-fold increased levels

of active Brnoteplase over alteplase at doses of 0.9, 2.5, and 5.0 mg/kg, respectively. Notably,

the residual active Brnoteplase concentration after administering the 0.9 mg/kg dose (0.22 ±

0.06 mg/L) was significantly higher even than the alteplase concentration after 5.0 mg/kg (0.09

± 0.01 mg/L). The distinct administration regimes—bolus for Brnoteplase vs. continuous rate

infusion for alteplase—underscore this difference and support that the improved inhibition

resistance of Brnoteplase (**Figure 2e**) and potentially alleviated binding to clearance receptors

(**Table 3**) substantially extended its biological half-life.

Figure 5: Determination of pharmacological marker levels in rat plasma samples collected at the end of the
thrombolysis effectivity experiments. (**a**) Comparison of residual active concentrations of administered
thrombolytics as an indicator of the protein residence time and biological half-life. (**b**) Comparison of residual
plasminogen-to-plasmin ratios as an indicator of thrombolytic selectivity and safety potential. The bars represent
mean values and the whiskers correspond to standard errors.

Plasminogen and plasmin levels were quantified to evaluate the thrombolytic pharmacological profile. Substantial systemic plasminemia is associated with increased hemorrhagic risk, whereas preserving high plasminogen levels and restricting plasmin generation to occluded areas promises more favorable treatment outcomes[17]. Accordingly, the plasminogen/plasmin ratio serves as an indicator of thrombolytic selectivity and safety potential. Across all tested doses (0.9, 2.5, and 5.0 mg/kg), Brnoteplase-treated rats exhibited a consistent trend of higher plasminogen/plasmin ratios than the alteplase group (**Figure 5b**, **Table S10**). Furthermore, Brnoteplase treatment at the dose of 5.0 mg/kg retained a comparable or even higher ratio (2700 ± 130) relative to the alteplase group at the clinically relevant dose of 0.9 mg/kg (1970 ± 510). Together with the significantly increased *in vivo* thrombolytic efficacy of Brnoteplase (**Figure 4a**), these findings corroborate previously observed enhanced selectivity (**Figure 2c**) and favorable safety profile (**Figure 4bc**) of Brnoteplase.

Discussion

In this study, we established a comprehensive protein engineering pipeline integrating *in silico* rational design, biochemical characterization, *in vitro* clot lysis and penetration assays, and *in vivo* animal models. This workflow enables initial high-throughput screening of numerous thrombolytic candidates, followed by systematic filtering to prioritize variants for resource-intensive *in vivo* testing. Through this approach, we identified Brnoteplase, a novel thrombolytic with improved pharmacological properties, representing a promising candidate for future clinical evaluation.

Biochemical characterization revealed that Brnoteplase exhibited (i) markedly enhanced fibrin selectivity (80-fold higher than alteplase), (ii) superior resistance to PAI-1 inhibition (4-fold higher), and (iii) increased clot penetration potential (4-fold higher). These parameters are critical determinants of safer and more effective thrombolysis and provide a mechanistic basis for the favorable outcomes observed in subsequent *in vitro* and *in vivo* experiments.

The pronounced fibrin selectivity of Brnoteplase ensures activation of plasminogen predominantly within occluded regions with fibrin-containing thrombi, while minimizing systemic

plasminemia associated with hemorrhagic risk. As a result, Brnoteplase maintains a higher

plasminogen/plasmin ratio relative to alteplase, as confirmed by ELISA analysis of rat plasma

collected after thrombolysis. Additionally, the *in silico* design of Brnoteplase aimed to reduce

affinity for LRP1 and NMDAR which is expected to mitigate off-target central nervous system

interactions and thereby further enhance the safety profile[15,16]. All these findings were

corroborated *in vivo*, where the Brnoteplase-treated group exhibited a lower proportion of severe

hemorrhagic transformations compared to alteplase and tenecteplase.

Improved inhibition resistance, another major characteristic of Brnoteplase, aligns with

expectations from the rational design phase. This variant incorporates mutations previously

shown to reduce interaction with PAI-1, a primary inhibitor of tissue plasminogen activator in

human blood. Together with mutations designed to minimize binding to clearance receptors such

as LRP1 and mannose receptors, these combined features were intended to modulate protein

pharmacokinetics[6,48,49], and this expectation was confirmed *in vivo*. Rat plasma samples collected

at the end of thrombolysis experiments revealed significantly higher residual levels of active

Brnoteplase compared to alteplase, indicating an extended biological half-life. This enabled

efficient bolus administration, a clinically meaningful advantage over alteplase.

Finally, superior clot penetration potential of Brnoteplase in biochemical assays was derived

from decreased fibrin affinity. This assumption was validated by *in vitro* assays directly

monitoring thrombolytics motion through fibrin matrix gels. Compared to alteplase, Brnoteplase

exhibited a faster penetration rate and greater distribution within the clot. This facilitates lysis

from within the clot rather than relying solely on surface degradation, which is characteristic of

conventional thrombolytics. By promoting more uniform and deeper fibrinolysis, Brnoteplase may

accelerate recanalization, particularly in large or densely structured thrombi.

In addition to these mechanistic advantages, Brnoteplase demonstrated a favorable dose–

response profile *in vitro*. Thrombolytic efficacy progressively increased with concentration until

plateau, whereas alteplase exhibited an early peak with a decline at elevated doses. Combined

with the Brnoteplase selectivity and reduced affinity for receptors, this profile suggests that

higher dosing could enhance thrombolytic potency without compromising safety and hemorrhagic

complications. Consistent with these findings, *in vivo* experiments in rat models confirmed that Brnoteplase achieved significantly higher thrombolysis rates and recanalization frequency across all tested doses compared to alteplase. Importantly, safety assessments demonstrated the lowest incidence of severe hemorrhagic transformation and minimal hemispheric asymmetry in Brnoteplase-treated animals relative to both alteplase and tenecteplase. Nevertheless, given the low overall occurrence of hemorrhagic events in the tested cohorts, larger-scale studies will be required to confirm statistical significance of the safety study outcome.

Considering that Brnoteplase incorporates several mutations originally introduced in tenecteplase, it is not unexpected that they share a similar profile in biochemical testing, namely fibrin selectivity and PAI-1 resistance. However, Brnoteplase additionally exhibited significantly greater fibrin clot penetration and a lower rate of severe hemorrhagic transformation, positioning it as a potentially advantageous alternative within the class of current state-of-the-art thrombolytics.

In summary, we developed a comprehensive and scalable workflow for the rational design, rapid initial screening and prioritization, and the final selection of the most viable and clinically relevant novel thrombolytic proteins. By applying this pipeline, we generated Brnoteplase, an alteplase variant combining enhanced fibrin selectivity, improved inhibition resistance, prolonged biological half-life, superior clot penetration, and a favorable dose–response profile. These attributes collectively translate into higher thrombolytic efficacy and a promising safety profile in preclinical *in vivo* models. While further validation in larger animal cohorts is essential, Brnoteplase represents a strong candidate for follow-up (pre)clinical studies. By enabling bolus administration due to extended half-life, reducing hemorrhagic risk via diminished systemic plasminemia, and accelerating recanalization through deeper intrathrombus penetration, Brnoteplase has the potential to advance acute ischemic stroke therapy beyond the limitations of current thrombolytic agents.

Funding

This work was supported by the Czech Ministry of Education (INBIO - CZ.02.1.01/0.0/0.0/16_026/0008451, ENOCH - CZ.02.1.01/0.0/0.0/16_019/0000868); Grant Agency of Czech Republic (20-15915Y); and National Institute for Neurology Research (LX22NPO5107 MEYS) Financed by European Union – Next Generation EU. MT, ST, and JSM are supported by the scholarship Brno Ph.D. Talent and MM is supported by the Masaryk University (MUNI/H/1561/2018). ST, MP, and JV received additional support from the Ministry of Health of the Czech Republic in cooperation with the Czech Health Research Council under project No. NW24-08-00064. Computational resources were provided by the e-INFRA CZ project (ID:90254), supported by the Ministry of Education, Youth and Sports of the Czech Republic, and by the ELIXIR-CZ project (ID:90255), part of the international ELIXIR infrastructure.

Author Contribution Statement.

J.M and D.B. performed the *in silico* analysis and selection of the initial thrombolytic pool. **Pe.Ka. and T.B.** produced the designed thrombolytic enzymes. **M.T. and V.S.** performed the biochemical characterization of selected proteins and ELISA measurements of rat plasma samples and analyzed the corresponding data. **S.T., M.P., Pa.Ki., and J.V.** performed the *in vitro* thrombolysis and fibrin penetration assays and analyzed the corresponding data. **P.S., J.H., E.B., A.D.A., J.B., and J.O.** performed the *in vivo* animal testing and the corresponding data analysis and statistical evaluation. **M.T., V.S.** drafted the initial version of the manuscript with the help of **J.M., S.T., and J. V..** **J.D., Z.P., D.B., M.M., J.V., L.K., and R.M.** conceived and designed the study, supervised the project, and acquired funding. **All authors** reviewed and interpreted the data, critically revised the manuscript, and approved the final version.

Acknowledgements

We thank the laboratory technicians Marketa Stuchla, Veronika Novakova, Irena Halikova, and Klara Markova for their assistance with the preparation of common reagents and sample storage.

Conflict of Interest

None declared.

References

1. Feigin VL, Brainin M, Norrving B, et al. World Stroke Organization: Global Stroke Fact Sheet 2025. *Int J Stroke*. 2025;20(2):132-144. doi:10.1177/17474930241308142

2. Aguiar de Sousa D, von Martial R, Abilleira S, et al. Access to and delivery of acute ischaemic stroke treatments: A survey of national scientific societies and stroke experts in 44 European countries. *Eur Stroke J*. 2019;4(1):13-28. doi:10.1177/2396987318786023

3. Xiong Y, Wakhloo AK, Fisher M. Advances in Acute Ischemic Stroke Therapy. *Circ Res*. 2022;130(8):1230-1251. doi:10.1161/CIRCRESAHA.121.319948

4. Yaghi S, Willey JZ, Cucchiara B, et al. Treatment and Outcome of Hemorrhagic Transformation After Intravenous Alteplase in Acute Ischemic Stroke: A Scientific Statement for Healthcare Professionals From the American Heart Association/American Stroke Association. *Stroke*. 2017;48(12):e343-e361. doi:10.1161/STR.0000000000000152

5. Mosconi MG, Paciaroni M. Treatments in Ischemic Stroke: Current and Future. *Eur Neurol*. 2022;85(5):349-366. doi:10.1159/000525822

6. Mican J, Toul M, Bednar D, Damborsky J. Structural Biology and Protein Engineering of Thrombolytics. *Comput Struct Biotechnol J*. 2019;17:917-938. doi:10.1016/j.csbj.2019.06.023

7. Potla N, Ganti L. Tenecteplase vs. alteplase for acute ischemic stroke: a systematic review. *Int J Emerg Med*. 2022;15(1):1. doi:10.1186/s12245-021-00399-w

8. Burwell JM, Howay JR, Wasko L, et al. Tenecteplase is here: navigating the shift of a stroke thrombolytic in the United States prior to FDA approval: a mini-review on rationale, barriers, and pathways. *Front Neurol*. 2025;16:1563423. doi:10.3389/fneur.2025.1563423

9. Rodriguez M, Sidebottom C, Wells DA, et al. A Systematic Review of the Efficacy and Safety of Tenecteplase Versus Alteplase in Acute Ischemic Stroke: A Time to Pass the Torch. *Stroke Vasc Interv Neurol*. 2024;4(4):e001110. doi:10.1161/SVIN.123.001110

10. Ma Y, Xiang H, Busse JW, et al. Tenecteplase versus alteplase for acute ischemic stroke: a systematic review and meta-analysis of randomized and non-randomized studies. *J Neurol*. 2024;271(5):2309-2323. doi:10.1007/s00415-024-12243-1

11. Rose D, Cavalier A, Kam W, et al. Complications of Intravenous Tenecteplase Versus Alteplase for the Treatment of Acute Ischemic Stroke: A Systematic Review and Meta-Analysis. *Stroke*. 2023;54(5):1192-1204. doi:10.1161/STROKEAHA.122.042335

12. Warach SJ, Dula AN, Milling TJ. Tenecteplase Thrombolysis for Acute Ischemic Stroke. *Stroke*. 2020;51(11):3440-3451. doi:10.1161/STROKEAHA.120.029749

13. Meng X, Li S, Dai H, et al. Tenecteplase vs Alteplase for Patients With Acute Ischemic Stroke: The ORIGINAL Randomized Clinical Trial. *JAMA*. 2024;332(17):1437-1445. doi:10.1001/jama.2024.14721

14. Van de Werf FJ. The ideal fibrinolytic: can drug design improve clinical results? *Eur Heart J*.
1999;20(20):1452-1458. doi:10.1053/euhj.1999.1659

15. Maier W, Bednorz M, Meister S, et al. LRP1 is critical for the surface distribution and
internalization of the NR2B NMDA receptor subtype. *Mol Neurodegener*. 2013;8(1):25.
doi:10.1186/1750-1326-8-25

16. Baron A, Montagne A, Cassé F, et al. NR2D-containing NMDA receptors mediate tissue
plasminogen activator-promoted neuronal excitotoxicity. *Cell Death Differ*. 2010;17(5):860-
871. doi:10.1038/cdd.2009.172

17. Nikitin D, Choi S, Mican J, et al. Development and Testing of Thrombolytics in Stroke. *J
Stroke*. 2021;23(1):12-36. doi:10.5853/jos.2020.03349

18. Lenka S, Jan S, T M, David B, Jiri D. HotSpot Wizard 3.0: web server for automated design of
mutations and smart libraries based on sequence input information. *Nucleic Acids Res*.
2018;46(W1):W356-W362. doi:10.1093/nar/gky417

19. Alford RF, Leaver-Fay A, Jeliazkov JR, et al. The Rosetta All-Atom Energy Function for
Macromolecular Modeling and Design. *J Chem Theory Comput*. 2017;13(6):3031-3048.
doi:10.1021/acs.jctc.7b00125

20. Prasad JM, Young PA, Strickland DK. High Affinity Binding of the Receptor-associated Protein
D1D2 Domains with the Low Density Lipoprotein Receptor-related Protein (LRP1) Involves
Bivalent Complex Formation: CRITICAL ROLES OF LYSINES 60 AND 191*. *J Biol Chem*.
2016;291(35):18430-18439. doi:10.1074/jbc.M116.744904

21. Freeman R, Niego B, R. Croucher D, Pedersen LO, Medcalf RL. t-PA, but not desmoteplase,
induces plasmin-dependent opening of a blood-brain barrier model under normoxic and
ischaemic conditions. *Brain Res*. 2014;1565:63-73. doi:10.1016/j.brainres.2014.03.027

22. M M, Rt K, A B, et al. FireProtASR: A Web Server for Fully Automated Ancestral Sequence
Reconstruction. *Brief Bioinform*. 2021;22(4):bbaa337. doi:10.1093/bib/bbaa337

23. J H, S B, J S, et al. EnzymeMiner: automated mining of soluble enzymes with diverse
structures, catalytic properties and stabilities. *Nucleic Acids Res*. 2020;48(W1):W104-W109.
doi:10.1093/nar/gkaa372

24. Toschi L, Bringmann P, Petri T, Donner P, Schleuning WD. Fibrin selectivity of the isolated
protease domains of tissue-type and vampire bat salivary gland plasminogen activators. *Eur
J Biochem*. 1998;252(1):108-112. doi:10.1046/j.1432-1327.1998.2520108.x

25. Harpaz D, Chen X, Francis CW, Marder VJ, Meltzer RS. Ultrasound enhancement of
thrombolysis and reperfusion in vitro. *J Am Coll Cardiol*. 1993;21(6):1507-1511.
doi:10.1016/0735-1097(93)90331-T

26. Prasad S, Kashyap RS, Deopujari JY, Purohit HJ, Taori GM, Daginawala HF. Development of
an in vitro model to study clot lysis activity of thrombolytic drugs. *Thromb J*. 2006;4:14.
doi:10.1186/1477-9560-4-14

27. Thalerová S, Pešková M, Kittová P, et al. Effect of Apixaban Pretreatment on Alteplase-
Induced Thrombolysis: An In Vitro Study. *Front Pharmacol*. 2021;12:740930.
doi:10.3389/fphar.2021.740930

28. Víteček J, Wünschová AV, Thalerová S, et al. Factors influencing the efficacy of recombinant
tissue plasminogen activator: Implications for ischemic stroke treatment. *PLOS ONE*.
2024;19(6):e0302269. doi:10.1371/journal.pone.0302269

29. Thalerová S, Wünschová AV, Kittová P, et al. A collateral circulation in ischemic stroke
accelerates recanalization due to lower clot compaction. *PLOS ONE*. 2024;19(11):e0314079.
doi:10.1371/journal.pone.0314079

30. Nikitin D, Mican J, Toul M, et al. Computer-aided engineering of staphylokinase toward
enhanced affinity and selectivity for plasmin. *Comput Struct Biotechnol J*. 2022;20:1366-
1377. doi:10.1016/j.csbj.2022.03.004

31. Diamond SL. Engineering Design of Optimal Strategies for Blood Clot Dissolution. *Annu Rev*
Biomed Eng. 1999;1(Volume 1, 1999):427-461. doi:10.1146/annurev.bioeng.1.1.427

32. Acheampong P, and Ford GA. Pharmacokinetics of alteplase in the treatment of ischaemic
stroke. *Expert Opin Drug Metab Toxicol*. 2012;8(2):271-281.
doi:10.1517/17425255.2012.652615

33. Elnager A, Abdullah WZ, Hassan R, et al. In Vitro Whole Blood Clot Lysis for Fibrinolytic
Activity Study Using D-Dimer and Confocal Microscopy. *Adv Hematol*. 2014;2014:814684.
doi:10.1155/2014/814684

34. Rottenberger Z, Komorowicz E, Szabó L, et al. Lytic and mechanical stability of clots
composed of fibrin and blood vessel wall components. *J Thromb Haemost*. 2013;11(3):529-
538. doi:10.1111/jth.12112

35. Marcos-Contreras OA, Ganguly K, Yamamoto A, et al. Clot penetration and retention by
plasminogen activators promote fibrinolysis. *Biochem Pharmacol*. 2013;85(2):216-222.
doi:10.1016/j.bcp.2012.10.011

36. Longstaff C. Measuring fibrinolysis: from research to routine diagnostic assays. *J Thromb*
Haemost. 2018;16(4):652-662. doi:10.1111/jth.13957

37. Longstaff C, Varjú I, Sótonyi P, et al. Mechanical Stability and Fibrinolytic Resistance of Clots
Containing Fibrin, DNA, and Histones*. *J Biol Chem*. 2013;288(10):6946-6956.
doi:10.1074/jbc.M112.404301

38. Morrow GB, Whyte CS, Mutch NJ. Functional plasminogen activator inhibitor 1 is retained on
the activated platelet membrane following platelet activation. *Haematologica*.
2019;105(12):2824-2833. doi:10.3324/haematol.2019.230367

39. Vítečková Wünschová A, Novobilský A, Hložková J, et al. Thrombus Imaging Using 3D Printed
Middle Cerebral Artery Model and Preclinical Imaging Techniques: Application to Thrombus
Targeting and Thrombolytic Studies. *Pharmaceutics*. 2020;12(12):1207.
doi:10.3390/pharmaceutics12121207

40. Kavkova M, Zikmund T, Kala A, et al. Contrast enhanced X-ray computed tomography imaging
of amyloid plaques in Alzheimer disease rat model on lab based micro CT system. *Sci Rep*.
2021;11(1):5999. doi:10.1038/s41598-021-84579-x

41. Bassel-Duby R, Jiang NY, Bittick T, et al. Tyrosine 67 in the epidermal growth factor-like
domain of tissue-type plasminogen activator is important for clearance by a specific hepatic
receptor. *J Biol Chem*. 1992;267(14):9668-9677.

42. López-Atalaya JP, Roussel BD, Ali C, et al. Recombinant Desmodus rotundus salivary
plasminogen activator crosses the blood-brain barrier through a low-density lipoprotein
receptor-related protein-dependent mechanism without exerting neurotoxic effects. *Stroke*.
2007;38(3):1036-1043. doi:10.1161/01.STR.0000258100.04923.84

43. Narita M, Bu G, Herz J, Schwartz AL. Two receptor systems are involved in the plasma
clearance of tissue-type plasminogen activator (t-PA) in vivo. *J Clin Invest*. 1995;96(2):1164-
1168. doi:10.1172/JCI118105

44. Goulay R, Naveau M, Gaberel T, Vivien D, Parcq J. Optimized tPA: A non-neurotoxic
fibrinolytic agent for the drainage of intracerebral hemorrhages. *J Cereb Blood Flow Metab*.
2018;38(7):1180-1189. doi:10.1177/0271678X17719180

45. Keyt BA, Paoni NF, Refino CJ, et al. A faster-acting and more potent form of tissue
plasminogen activator. *Proc Natl Acad Sci U S A*. 1994;91(9):3670-3674.
doi:10.1073/pnas.91.9.3670

46. Hacke W, Kaste M, Fieschi C, et al. Randomised double-blind placebo-controlled trial of
thrombolytic therapy with intravenous alteplase in acute ischaemic stroke (ECASS II). *The*
Lancet. 1998;352(9136):1245-1251. doi:10.1016/S0140-6736(98)08020-9

47. Ma J, Ma Y, Shuaib A, Winship IR. Improved collateral flow and reduced damage after remote
ischemic perconditioning during distal middle cerebral artery occlusion in aged rats. *Sci Rep*.
2020;10(1):12392. doi:10.1038/s41598-020-69122-8

48. Biessen EAL, van Teijlingen M, Vietsch H, et al. Antagonists of the Mannose Receptor and the
LDL Receptor–Related Protein Dramatically Delay the Clearance of Tissue Plasminogen
Activator. *Circulation*. 1997;95(1):46-52. doi:10.1161/01.CIR.95.1.46

49. Chandler WL, Alessi MC, Aillaud MF, Henderson P, Vague P, Juhan-Vague I. Clearance of
Tissue Plasminogen Activator (TPA) and TPA/Plasminogen Activator Inhibitor Type 1 (PAI-1)
Complex. *Circulation*. 1997;96(3):761-768. doi:10.1161/01.CIR.96.3.761

50. Meiners K, Hamm P, Gutmann M, et al. Site-specific PEGylation of recombinant tissue-type
plasminogen activator. *Eur J Pharm Biopharm*. 2023;192:79-87.
doi:10.1016/j.ejpb.2023.09.017

51. Toul M, Slonkova V, Mican J, et al. Identification, characterization, and engineering of
glycosylation in thrombolytics. *Biotechnol Adv*. 2023;66:108174.
doi:10.1016/j.biotechadv.2023.108174

52. Ma H, Jiang Z, Xu J, Liu J, Guo ZN. Targeted nano-delivery strategies for facilitating
thrombolysis treatment in ischemic stroke. *Drug Deliv*. 28(1):357-371.
doi:10.1080/10717544.2021.1879315

53. Zenych A, Fournier L, Chauvierre C. Nanomedicine progress in thrombolytic therapy.
Biomaterials. 2020;258:120297. doi:10.1016/j.biomaterials.2020.120297

54. Rijken DC, Barrett-Bergshoeff MM, Jie AFH, Criscuoli M, Sakharov DV. Clot penetration and
fibrin binding of amediplase, a chimeric plasminogen activator (K2tu-PA). *Thromb Haemost*.
2017;91:52-60. doi:10.1160/TH03-07-0435

55. Tsikouris JP, White CM. Tenecteplase - A novel fibrin-specific thrombolytic agent for
treatment of acute myocardial infarction. *Formulary*. 2000;35(8):647-653.

56. Tanswell P, Modi N, Combs D, Danays T. Pharmacokinetics and pharmacodynamics of
tenecteplase in fibrinolytic therapy of acute myocardial infarction. *Clin Pharmacokinet*.
2002;41(15):1229-1245. doi:10.2165/00003088-200241150-00001

57. Logallo N, Novotny V, Assmus J, et al. Tenecteplase versus alteplase for management of
acute ischaemic stroke (NOR-TEST): a phase 3, randomised, open-label, blinded endpoint
trial. *Lancet Neurol*. 2017;16(10):781-788. doi:10.1016/S1474-4422(17)30253-3

58. Campbell BCV, Mitchell PJ, Churilov L, et al. Tenecteplase versus Alteplase before
Thrombectomy for Ischemic Stroke. *N Engl J Med*. Published online 2018:10.

59. Huang J, Zheng H, Zhu X, Zhang K, Ping X. Tenecteplase versus alteplase for the treatment
of acute ischemic stroke: a meta-analysis of randomized controlled trials. *Ann Med*.
56(1):2320285. doi:10.1080/07853890.2024.2320285

60. Xu Y, Chen D, Liu P, et al. A triple fusion tissue-type plasminogen activator (TriF-ΔtPA)
enhanced thrombolysis in carotid embolism-induced stroke model. *Int J Pharm*.
2023;637:122878. doi:10.1016/j.ijpharm.2023.122878

61. Chen S, Fang S, Zhou Y, et al. A low bleeding risk thrombolytic agent: citPA5. *Cardiovasc
Res*. 2024;120(10):cvae060. doi:10.1093/cvr/cvae060