# *In vivo* isogenic modelling unveils a *TP53*-mediated relapse phenotype in T-cell acute lymphoblastic leukemia

Stéphanie Gachet[1,2,3,#], Samuel Quentin[1,2,4*], Lucie Hernandez[1,2*], Loïc Maillard[1,2], Marie Passet[1,2,4], Rathana Kim[1,2,3,4], Hugo Bergugnat[1,2], Melha Benlebna[1,2], Maxime Boy[1,2], Véronique Parietti[1,2], Pierre Fenaux[5], André Baruchel[6], Hervé Dombret[1,4,5], Nicolas Boissel[1,2,3,7], François Sigaux[1,2,3,4], Hugues de Thé[1,2,3,4], Emmanuelle Clappier[1,2,3,4] and Jean Soulier[1,2,3,4].

[1]Institut de Recherche Saint-Louis (IRSL), Université Paris Cité, 75010 Paris, France;
[2]INSERM UMR1342 / CNRS EMR8000, Paris, France;
[3]Institut de la Leucémie, Hôpital Saint-Louis, APHP, Université Paris Cité, Paris, France;
[4]Hôpital Saint-Louis, Laboratoire d'Hématologie, APHP, Paris, France;
[5]Hôpital Saint-Louis, Département Hématologie Senior, APHP, Paris, France;
[6]Hôpital Robert Debré, APHP, Université Paris Cité, Paris, France;
[7]Hôpital Saint-Louis, Service d'Hématologie Adolescents et Jeunes Adultes, APHP, Paris, France.
[*]These co-authors contributed equally

**Corresponding authors:**
Dr. Stéphanie Gachet, PhD: stephanie.gachet@inserm.fr and Pr. Jean Soulier, MD, PhD: jean.soulier@aphp.fr

[#]Present address : Laboratoire des cellules Souches Hématopoïétiques et des Leucémies, Equipe Niche, Cancer et Radiations dans l'Hématopoïèse, Unité Mixte de Recherche (UMR) 1274-E008, INSERM, CEA, Université Paris-Saclay.

**Running title:**
A relapse-associated program driven by *TP53* alterations in T-ALL

**Keywords:** T-cell acute lymphoblastic leukemia, somatic *TP53* alterations, relapse, leukemia initiating cells, metabolic adaptation, p53 therapeutic targeting, preclinical models

**Footnotes:**

Supplementary informations for this article (Supplementary Figures, 5 and Supplementary Tables, 4) are available online.

**ABSTRACT**

Although most patients with T-cell acute lymphoblastic leukemia (T-ALL) achieve remission with chemotherapy, many relapse with a poor prognosis, treatment-resistant disease. The mechanisms driving chemoresistance remain largely elusive, in part due to the lack of faithful experimental models. Here, we generated isogenic patient-derived xenograft (PDX) pairs at diagnosis and relapse from T-ALL cases harboring *TP53* alterations at relapse (*TP53*^alt^, *TP53* mutation and/or deletion). These models faithfully captured the genomic and functional features of both the diagnosis and the relapse primary leukemias. Despite inter-T-ALL variability, comparative functional analyses revealed a unique, cell-intrinsic relapse phenotype that includes enhanced leukemia-initiating capacity and which can be experimentally conferred to diagnosis cells by *TP53* silencing. Deregulated metabolic pathways and increased MYC signaling are prominent in the relapse phenotype. Ultra-sensitive backtracking of *TP53*-mutant cells and single-cell expression profiling revealed cellular heterogeneity at diagnosis, with sequential progression from a minor subpopulation toward a biallelic *TP53*^alt^ relapse. Our findings support a model in which T-ALL relapse originates from a pre-existing transcriptional state with deregulated metabolism that is selected under therapeutic pressure, opening monitoring prospects for precision medicine.

# INTRODUCTION

T-cell acute lymphoblastic leukemia (T-ALL) is an aggressive hematologic malignancy that represents 15% of pediatric and 25% of all adult ALL cases[1,2]. Multistep deregulation of key transcription factors and additional genes controlling self-renewal, senescence, apoptosis, differentiation or proliferation in a T-cell precursor leads to clonal transformation and expansion of blast cells in the lymphoid organs and the bone marrow[1,3–7]. Although most T-ALL patients at diagnosis achieve remission with intensive chemotherapy and glucocorticoids, 15% to 40% of the patients subsequently experience relapse into a resistant leukemia with dismal prognosis[8–10]. Therefore, comprehensive understanding of the functional mechanisms underlying relapse and resistance is a major unmet need to develop more efficient therapeutic strategies. While exhaustive genetic and immunophenotypic landscape studies have enabled to identify a large variety of T-ALL subtypes[2,7,11,12], these subtypes only partially correlate with the onset of relapse. It rather appears that resistance initiates upon treatment by the selection of residual T-ALL cells with cellular features driving cell survival despite therapy and re-expansion[13–19]. Selection of *NT5C2* somatic mutations leading to cell resistance to 6-mercaptopurine is a well-established, although rare, example of resistance-driven mechanism in T- and B-ALL patients[20–24], and alternative mechanisms remain elusive.

Here, we investigated *TP53* alterations (TP53[alt], *TP53* mutation and/or deletion) as a factor of resistance acquisition in T-ALL. While rarely detected at diagnosis (2-3%), *TP53*[alt] is frequently found at relapse (10-25%) and is associated to poor clinical outcomes[25–30]. *TP53*[alt] cells are genetically trackable, allowing to monitor their functional relapse-associated phenotype. We unveiled a *relapse phenotype* that included increased leukemia initiating activity and modulation of key signaling and metabolic pathways. Sequential selection from a minor cell population expressing this *relapse phenotype* at diagnosis can prime relapse upon therapy.

# RESULTS

**Establishment of isogenic, patient-derived xenografts (PDX) faithfully capturing *TP53*-altered T-ALL diagnosis and relapse genomic patterns**

We aimed at uncovering functional mechanisms involved in leukemia resistance and relapse. We focused on *TP53*-altered T-ALL since *TP53* alterations are well-recognized, genetically-trackable, drivers of relapse but are still poorly mechanistically

understood in T-ALL[25–29,31]. We selected four cases of T-ALL with *TP53* alterations at

relapse only for which cryopreserved diagnosis, remission (i.e., non-tumoral) and

relapse patient samples were available. To overcome the limitations of *ex vivo*

functional studies on primary samples or cell lines, we established patient-derived

xenografts (PDXs) of paired diagnosis (Dx) and relapse (Rel) leukemia in

immunodeficient NOD.Cg-*Prkdc$_{scid}$ IL2rg$_{tm1Wjl}$*/SzJ (NSG) mice (**Figure 1A**). Genomic

analyses of the resulting isogenic, Dx-Rel PDX pairs showed that these PDXs largely

captured the genomic pattern of the patient cells, both at diagnosis and at relapse,

respectively (**Figure 1B** and **Supplementary Tables 1-2**). *TP53* alterations in patient

T-ALL cells were missense mutations in the *TP53* DNA-binding domain within

mutational hotspot residues R175 and R248 (cases TALL42; and TALL11 and TALL43,

respectively) and a less classical but already reported pathogenic mutation H214 (case

TALL30) (**Figure 1C**). The second allele was lost by deletion in 3 cases (TALL30,

TALL42 and TALL43) or by 17p unidisomy of the mutated allele in 1 case (TALL11)

(**Figure 1B**). The resulting PDXs carried the original *TP53* missense mutations

occurring in the DNA-binding domain in association with the loss of the remaining wild-

type (WT) allele, as in primary patient samples. Of note, in one T-ALL case, the

resulting PDX (PDX43R) captured one of the two *TP53*-altered clones from the relapse

patient sample (**Figure 1B**).

Interestingly, other somatic CNVs and point mutations (i.e., in addition to *TP53*[alt]) were

largely similar in both the diagnosis and relapse samples in the 4 T-ALL pairs,

suggesting that *TP53*-driven T-ALL relapse did not display stronger genomic instability

(**Figure 1B**).

Therefore, these T-ALL PDX models faithfully captured the diagnosis and the relapse

patient samples, respectively, enabling further functional comparison experiments

using this unlimited ressources of isogenic, human leukemic cells.

**Relapse cells display a cell-intrinsic, stable, *relapse phenotype* compared to**

**their isogenic diagnosis cells**

To uncover functional mechanisms involved in leukemia resistance and relapse, we

compared the leukemia propagation ability of the diagnosis and relapse T-ALL pairs.

First, primary leukemic cells from paired Dx-Rel patient samples were transplanted in

parallel at equivalent cell doses in NSG mice, and leukemia onset and mice survival

were monitored. Despite strong inter-patient T-ALL heterogeneity, we found in each

pair that the relapse cells propagated leukemia more efficiently than their diagnosis counterparts (**Supplementary Figure 1A**), which translated into a significantly shorter mice survival (**Figure 2A** and **Supplementary Figure 1B**). This increased leukemia propagating activity at relapse was conserved in the PDXs established from the 4 pairs. We performed multiple iterative transplantations using decreasing cell doses (**Figures 2B-C** and **Supplementary Figures 1C-F**) and found that the propagation phenotype was stable upon serial reinjection, without additional acceleration (**Supplementary Figures 1D,1F**). Limiting dilution *in vivo* assays (injection of $10^6$ to 1 leukemic cell/mouse) showed that the leukemia initiating cell (LIC) frequencies were increased in the Rel PDXs compared to isogenic Dx counterparts (**Figure 2D** and **Supplementary Table 3**). Consistently, *in vitro* long-term culture-initiating cell assays in coculture with MS5-DL1 stromal cells[32] showed a significantly increased clonogenicity in Rel PDXs from all T-ALL pairs, demonstrating that increased LIC activity is a cell-intrinsic feature of relapse cells (**Supplementary Figure 1G** and **Supplementary Table 3**). Real-time and end-point *in vitro* proliferation using Bromodeoxyuridine (BrdU) pulse labelling and apoptosis experiments, in particular cleaved caspase 3 staining, showed increased proliferation and decreased apoptosis in Rel versus Dx leukemic cells for all tested PDX pairs (**Figures 2E-F** and **Supplementary Figures 1H-I**). Moreover, homing of the leukemic cell in the bone marrow, spleen and liver at 24 hrs was increased in PDXs from Rel for the 4 T-ALL pairs (**Supplementary Figures 1J-K**). However, when homing was bypassed by direct injection of leukemic cells into the femoral bone marrow cavity, leukemic cells from Rel PDXs still propagated leukemia more efficiently than Dx PDXs (**Supplementary Figures 1L-M**).

Collectively, our data indicate that despite inter-T-ALL heterogeneity, *TP53*[alt] mediates a unique cell-intrinsic *relapse phenotype* that is associated with increased proliferation, cell survival, and cell homing and post-homing advantage, resulting in higher LIC activity.

**Experimental *TP53* silencing confers the cell-intrinsic *relapse phenotype* to diagnosis leukemia cells**

To model *TP53* deregulation as a driver of the *relapse phenotype*, we inactivated *TP53* using a lentiviral shRNA approach in Dx PDX leukemic cells followed by transplantation in NSG mice (**Figure 3A** and **Supplementary Figure 2A**). *TP53* downregulation

increased the leukemia propagation capacities of Dx PDX cells when transplanted

under parallel and competitive settings (**Figure 3B** and **Supplementary Figures 2B,**

**2D-E**), and resulted in enhanced leukemia spreading and shorter survival of the

recipient mice (**Figure 3C** and **Supplementary Figure 2C**). This phenotype was

retained in secondary transplants, both in parallel and competitive settings (**Figures**

**3D-F** and **Supplementary Figures 2F-H**).

*In vitro* limiting dilution and proliferation experiments on the *TP53*-silenced Dx PDX

cells confirmed the *relapse phenotype*, indicating a *TP53*-dependent increased

clonogenic activity (**Figure 3G**, **Supplementary Table 3** and **Supplementary Figure**

**2I**).

Finally, *in vivo p53*-silenced Dx cells also displayed increased homing and post homing

capacities (**Supplementary Figures 2J-M**).

Altogether, these results demonstrate that *TP53* silencing confers functional features

of the *relapse phenotype* to patient diagnosis cells.

***TP53* alterations deregulate MYC signaling and metabolism pathways in T-ALL**

**cells and directly contributes to the *relapse phenotype***

To dig into the biological pathways deregulated by *TP53*[alt] in T-ALL relapse, we

compared transcriptional profiles of isogenic Dx and Rel T-ALL PDX samples. While

91 genes were upregulated at relapse, 181 genes were downregulated including, as

expected, *TP53* itself (except in PDX11R where *TP53* was point-mutated on both

alleles) (**Figure 4A** and **Supplementary Figures 3A-B**). Geneset enrichment analysis

(GSEA) identified the MYC, E2F, and G2M checkpoint pathways as enriched in Rel

cells along with several cell metabolism pathways, including oxidative phosphorylation

(OXPHOS), glycolysis and cholesterol homeostasis, defining a *relapse expression*

*signature* (**Figure 4B**). Moreover, we interrogated genesets linked to cell proliferation,

invasiveness and stemness, and evidenced their upregulation at relapse

(**Supplementary Figure 3C**). Analysis in secondary transplant PDXs showed

consistent expression patterns, demonstrating functional stability (**Supplementary**

**Figures 3D-G** and data not shown). The *relapse expression signature* was also

retrieved from the corresponding primary T-ALL samples, further outlining that our Dx-

Rel PDX models faithfully captured the features of the patient cells (**Figure 4C**).

Considering the role of TP53 in regulating Myc signaling and metabolism

pathways[33,34], we focused on OXPHOS and glycolysis and performed real-time

measurements of oxygen consumption rate (OCR) and extra cellular acidification rate
(ECAR) in isogenic Dx-Rel PDX pairs. We found increased basal and maximal
mitochondrial respiration, along with increased glycolysis, in the Rel vs Dx T-ALL cells
(**Figures 4D-E** and **Supplementary Figures 3H-K**). Consistently, experimental
silencing of *TP53* in the diagnosis PDX cells conferred a relapse mitochondrial and
glycolytic profile, demonstrating a driver role of *TP53* inactivation into deregulation of
metabolic functions (**Figures 4F-G** and **Supplementary Figures 3L-N**).
To evaluate the relevance of the *relapse signature* beyond *TP53* genomic alterations,
we analyzed gene expression data from eleven additional primary T-ALL sample pairs
lacking *TP53* alterations. Notably, the *relapse expression signature* was enriched in
seven of these cases, suggesting that a conserved relapse-associated phenotype may
be activated through alternative mechanisms independent of *TP53* genetic alterations.
(**Supplementary Figures 3O-P**).
150

**Backtracking pre-relapse *TP53*alt clones at diagnosis**
As T-ALL relapse can emerge from a minor preexisting subclone at
diagnosis[13,17,19,22,35,36], we next aimed to backtrack rare *TP53*-mutated cells that would
constitute a pre-relapse reservoir. We performed ultra-high sensitivity approach,
namely droplet digital PCR (ddPCR), on the isogenic Dx-Rel primary and PDX
samples. We found the expected *TP53* mutation in each case at diagnosis, with mutant
allele frequencies corresponding to approximately 1/1300 to 1/5000 total cells, i.e.,
below the usual medical diagnosis detection thresholds (**Supplementary Table 4**).
This demonstrates the presence in each case of a very minor *TP53*-mutant clone at
diagnosis that was further selected in the post-treatment relapse sample. Interestingly,
a remission sample was available in one case (TALL11), in which we detected the
expected *TP53-p.R248Q* mutation with a frequency of 1 out of 2500 total cells,
demonstrating persistence of the cells reservoir after treatment. Strikingly, when the
remission sample was injected into NSG mice, a full-blown leukemia developed from
these rare cells. All the resulting leukemic cells carried the *TP53-p.R248Q* missense
mutation, definitively showing the LIC ability of the minor cell population from
diagnosis. However, this leukemia, raised from the remission sample, retained a
wildtype allele (*TP53-p.R248Q* allele frequency, 45.7% in sorted leukemia cells) by
contrast with the relapse cells that demonstrated copy-neutral loss of the wildtype allele
(*TP53-p.R248Q* allele frequency, 91.76%), highlighting the advantage conferred by the

monoallelic point mutation in the leukemia expanded from the patient remission sample
while the relapse cells displayed an additional loss of the wildtype *TP53* allele with
duplication of the *TP53-p.R248Q* allele through acquisition of an uniparental disomy.
Those data demonstrates that relapse emerges from a very minor (that is, undetected
by standard medical screen) that expands upon therapy with sequential progression
towards bi-allelic *TP53^{alt}* relapse.
177

**Single-cell profiling can uncover a pre-relapse minor cell population at diagnosis**
**that expresses features of the *relapse phenotype***
In an attempt to profile functionally the pre-relapse cell reservoir from diagnosis, we
performed single-cell RNAseq analysis on the four T-ALL pairs. The 16 resulting
samples (8 primary and 8 from PDXs) did not globally step out by their diagnosis or
relapse nature (i.e., all diagnosis *versus* all relapse samples), nor by their origin (i.e.,
all primary *versus* all PDX samples), or by cell cycle repartition, but rather by individual
T-ALL case (**Figure 5A** and **Supplementary Figures 4A-B**), underlying prominent
inter T-ALL biological heterogeneity. GSEA confirmed oxidative phosphorylation
signature as the most differentially expressed pathway in all relapse samples along
with cell proliferation, cell signaling and cell metabolism pathways (**Supplementary**
**Figures 4C-F** and **Figures 5B-C**).
Integrated comparison of the diagnosis and relapse samples uncovered a minor cell
population in two diagnosis samples that clustered nearby their paired relapse
counterparts, suggesting a "pre-relapse" cells reservoir (**Supplementary Figures 4C,**
**4E**).
*In case TALL42*, the small cluster of cells represented approximately 10% of the total
leukemic cells (9.6% and 15% of the primary and the PDX diagnosis samples,
respectively) (**Figures 5D-E** and **Supplementary Figures 4G-P**). This cluster was
enriched in most of the *relapse expression signature* (**Figure 5F** and **Supplementary**
**Figure 4Q**) including OXPHOS, MTORC1 signalling, MYC targets V1, E2F targets,
cholesterol homeostasis and DNA repair, these pathways being even more
upregulated in the matched relapse cells. In contrast, the G2M checkpoint, MYC target
V2 and unfolded protein response (UPR) pathways were downregulated in the minor
subclone as compared to the remaining diagnosis cells and to the relapse cells,
suggesting an intermediate less proliferative state (**Figure 5F** and **Supplementary**
**Figure 4Q**). We interrogated additional geneset collections and found a significant

enrichment of stemness, invasiveness and OXPHOS signatures combined with downregulation of cell proliferation pathways in the subclone, in line with the properties attributed to a quiescent LIC-activity enriched, pre-relapse leukemia reservoir (**Figure 5G** and **Supplementary Figure 4R**). Specifically, interrogation of several leukemia stem cell (LSC) signatures[37–40] showed that the pre-relapse subclone from primary and PDX diagnosis samples had increased LSC signature scores compared to diagnosis cells, as in relapse cells (**Figure 5H**).

*In the second case, TALL30*, the deregulation of the *relapse signature* was also detected in a minor subclone at diagnosis. Although this case eventually diverged from the previous T-ALL42 case by the late acquisition of a *NT5C2* mutation, known to drive resistance to maintenance therapy, the initiation of the relapse through a pre-existent population at diagnosis displaying features of the *relapse signature* suggests a common mechanism (**Supplementary Figure 5**).

Intriguingly, *TP53* mutation was found in only 0.03% (1 out of 3000) total cells of the two diagnosis samples TALL30 and TALL42, contrasting with approximately 5 and 10% of the cells expressing the relapse signature, respectively). Copy-number variations (CNVs) screen using inference analysis showed that the chromosome 17p was not deleted in the pre-relapse subclone in the two cases (**Figure 5I Supplementary Figure 4S** and **Supplementary Figure 5H, 5O)**. This suggests that only a small fraction of the minor cell population, i.e. the cells harboring the *TP53* mutation, is further selected towards relapse by chemotherapy, with an additional acquisition of del17p.

## DISCUSSION

Treatment failure and subsequent patient death are largely related to the emergence, within the first 2-years, of a relapse much less responsive to therapies than the initial diagnosis leukemia. Comprehensive longitudinal genomic studies on patient cohorts showed that a few cancer genes can be mutated at T-ALL relapse, *TP53* alterations being the most prominent[25–29]. However, no clear genomic driver of relapse can be detected in most cases and the underlying mechanisms, maybe non-genetic, remain elusive.

Here we leveraged *TP53*[alt] T-ALL relapse cases, i.e. leukemia with a clear, genetically-trackable driver of resistance and relapse to develop faithfull *in vivo* models based on

isogenic diagnosis-relapse paired xenografts. Contrasting with many other T-ALL
cases where a minor pre-relapse subclone at diagnosis can be selected by the
xenotransplantation process[13], our current PDXs faithfully captured the features of both
the diagnosis and the relapse T-ALLs, respectively. This substantial difference may be
due to the *TP53* lesions nature confering an *in vivo* advantage upon treatment only, as
seen in other premalignant conditions[18,41]. Importantly, our isogenic PDX models
enable to overcome fundamental limitations of *ex vivo* functional studies on primary
samples or cell lines, i.e. rarity and practical workability for primary samples; highly
rearranged and drifted cells for cell lines; and, more generally, the absence of a
complex microenvironnement crosstalk. In addition, isogenic PDXs also allow to avoid
experimental bias related to inter-patient heterogeneity, see for instance the markedly
distinct proliferation range between our 4 T-ALL cases, faithfully captured by the PDX
isogenic pairs (**Supplementary Figures 1 A-E**).

Interestingly, we found that the *TP53*-driven T-ALL relapse cells did not display strong
genomic instability features, suggesting a direct oncogenic role of *TP53* alterations in
those leukemias as "gatekeeper" rather than its commonly recognized "caretaker" role.
This was confirmed by *TP53* silencing experiments evidencing a cell-intrinsic gain of
malignancy in relapse cells largely driven by the *TP53* lesion itself. The *relapse*
*phenotype* we identied includes a cell-intrinsic functional gain of leukemia initiating
activity and proliferation, along with an expression profile that involved cell metabolism,
signaling, MYC pathways, and stemness, consistently with a role of *TP53* in the
regulation of normal and pathologic stem cell function in hematopoiesis[42]. We
especially evidenced that increased p53-dependent OXPHOS was crucial to the
*relapse phenotype,* as seen in other acute leukemia subtypes such as acute myeloid
leukemia[43]. Collectively, the metabolic *relapse signature* found at relapse, and in pre-
relapse minor reservoir at diagnosis, suggests that *TP53* alterations primed T-ALL
relapse in a subpopulation resistant to chemotherapy and enriched in LIC activity,
along with metabolic and MYC pathway reprogramming.

Like in the vast majority of cancers, *TP53*-altered relapses in T-ALL are associated
with very poor clinical outcomes and limited therapeutic options, representing a major
clinical challenge. While T-ALL stem cell identification by usual immunophenotypic
techniques is still debated[44–46,40], our data of a subpopulation resistant to
chemotherapy and enriched in LIC activity through enhanced oxidative
phosphorylation and MYC pathway activation with concurrent sequential acquisition of

bi-allelic *TP53* alterations raises therapeutic propects. Detecting very rare pre-existing

*TP53*-mutant cells at diagnosis yields potential strategies of early detection, monitoring

and adapted treatment of these cells by early administration of targeted therapies in

combination with classical chemotherapy and timely and risk-adapted hematopoietic

stem cell transplantation.

Interestingly, many T-ALL relapses did not display any additional "relapse-associated

genes"[17], suggesting that leukemia resistance and relapse may also be driven by non-

genetic mechanisms. In line with this observation, several studies have raised a

spectrum of key epigenetic, transcriptionnal remodeling, cell plasticity and

microenvironmental/niche mechanisms that may contribute to relapse[17,47,48,35,37,40].

Interestingly, the enrichment of the p53-associated *relapse expression signature* in

several T-ALL cases lacking *TP53* alterations at relapse suggests that other, possibly

epigenetic, mechanisms may activate the *relapse phenotype* we evidenced in this

work. This raises the possibility that these biological patterns of resistance may be of

wide relevance, including for strategies of early detection, monitoring and targeting of

pre-relapse cell populations.

## METHODS

**Patient cohort, collection of biological material and annotations.** Primary samples from 42 T-ALL patients who have relapsed were obtained from blood or bone marrow (BM) aspirates/biopsies at initial diagnosis, remission (considered as germline reference) and first relapse. The cohort included children and young adults (age, 1 to 42 years old). Samples were stored in the Tumor Bank of Saint-Louis Hospital, Paris (BiRTH, Biobank for Research in Translational Hematology) accordingly to the French law, including informed consent. Annotations of T-ALL mutations were obtained using the NGS sequencing and analysis pipeline from Saint-Louis Hospital, using a VAF threshold of 2%[49]. Seven of the 42 patients (16.7%) had *TP53* alterations at relapse, including 4 for which live cells from both diagnosis and relapse could be recovered and injected to mice. IRB approval from the Inserm was given to this study under the number #12–078.

**Animal studies.** Mice experiments described in this study comply with the current ethical regulations. Authorizations for animal experimentation were obtained from the *Ministère de l'enseignement supérieur et de la recherche* with the No. APAFIS#19958-2019032615074625v2 and APAFIS#47448-2024020909144467v3. Mice were housed under a dark-light cycle at ambient temperature ($\approx$18-23°C) with a humidity of 40-60% and handled in a pathogen-free animal facility in accordance with the guidelines of the Animal Care and Use Committee (IRSL, Université Paris Cité). Animals were excluded from experiments if any sign of stress were observed without clinical signs of full-blown leukemia development (absence of human leukemic cells in blood, bone marrow and spleen).

**Patient derived xenografts (PDXs) of isogenic diagnosis-relapse T-ALL samples.** Isogenic diagnosis (Dx) and relapse (Rel) T-ALL patient samples were thawed, qualified and injected intravenously (unless otherwise stated) at equivalent cell doses into sublethally irradiated immunodeficient NOD.Cg-Prkdc$_{scid}$ IL2rg$_{tm1Wjl}$/SzJ (NSG) mice from the Jackson Laboratory that are bred locally, as described previously[49]. Leukemia development was monitored monthly by flow cytometry (FACS) analysis of blood or BM aspirates using hCD45 and hCD7-based staining protocol detailed in the 'Flow cytometry' section to confirm disease engraftment.

Human T-ALL cells from both patients and derived PDXs (BM of fully leukemic mice

with >95% blasts) were used for genomics, molecular and functional experiments. To

evaluate the stability of the phenotype from PDX models, leukemic cells from PDXs at

first passage were serially transplanted into NSG recipient mice by intravenous

injection (unless otherwise stated) at equivalent cell doses. In some circumstances to

evaluate post homing involvement, leukemic cells from isogenic Dx-Rel PDXs were

injected into sublethally irradiated mice by intrafemoral route. Recipient mice were

monitored daily by visual and clinical observations and monthly for biological

parameters, i.e. signs of stress (weakness, weight loss, ruffled coat, lethargy or ataxia)

and the presence of human leukemic cells in blood and/or BM. To monitor survival,

mice were sacrificed at or before reaching ethical endpoints. Pathology and flow

cytometry analyses confirmed full-blown T-ALL disease with marrow, spleen and liver

infiltration by leukemic cells.

**Flow cytometry.** Leukemic cells obtained from peripheral blood after Ficoll separation

and from BM aspirates were washed and resuspended in PBS (*Gibco*) supplemented

with 2% FBS (*Gibco*) before staining. Human leukemic cells infiltration in NSG recipient

mice was measured in legs, spleen and liver that were crushed or mashed. Leukemic

cell suspensions were washed and resuspended in PBS supplemented with 2% FBS

for staining. Human leukemic cells were then stained for 30 min at 4°C with anti-human

CD45 (APC, VioBlue) and CD7 (PE, PE-Vio770) antibodies (*Miltenyi Biotec*) (1:100

dilution). Cells were then washed and analysed on a FACS Canto II or LSR Fortessa

analyzer (*BD Biosciences*). Flow cytometry analyses were performed using Flowjo

10.10 (*TreeStar*).

**Whole Exome Sequencing Analysis of primary and PDX T-ALL samples.** Primary

samples from matched tumoral (initial diagnosis and first relapse) and non-tumoral

(remission) obtained from T-ALL patients and the corresponding isogenic Dx-Rel PDXs

were subjected to deep whole-exome sequencing (WES). DNA libraries were prepared

using SureSelect XT Target Enrichment according to the manufacturer's instructions

(*Agilent Technologies*, Santa Clara, CA, USA). Exon capture was performed with

Agilent SureSelect Human All Exon V5 kit. The samples were sequenced on an

Illumina NextSeq 500 platform (*Illumina*, San Diego, CA, USA) in a 2 × 100 bp paired-

end sequencing run. The average sequencing depth was 110X. Raw data were first

submitted to quality check using Trim_Galore (v0.6.6) to perform quality and adapter trimming for paired-end reads. The paired end clean reads were aligned to the human reference genome (hg19) using BWA (v0.7.17-r1188). Duplicates reads were flagged using Picard-tools (v2.3.0) and local realignment around InDels and base quality score recalibration was performed using Genome Analysis ToolKit (GATK v4.2.5.0).

Somatic mutations were called by VarScan somatic (v2.4.2) and annotated by ANNOVAR (version 2022-04-13) before filtering. Somatic copy number variants (CNVs) were detected by facets (v0.16.0-1) and by an in-house R script. The final retained abnormalities were validated by visual inspection considering a number of markers ≥2 and log2ratio with respect to the individual background noise.

Exon capture was performed with Agilent SureSelect Human All Exon V6 UTR kit for the isogenic Dx-Rel PDX pairs. The same procedure as above was used to analyse PDX samples (n= 8) using the corresponding primary non-tumour sample as reference for somatic calling variant step. The overlap between V5 and V6_UTR capture was performed for the upstream analysis.

***In vivo* limiting dilution experiments.** Leukemic cells from paired Dx-Rel PDXs (primografts) were intravenously injected under limiting dilution conditions from $10^6$ to 1 cell per sublethally irradiated mouse (up to 6 mice per condition). Leukemia development in recipient NSG mice was monitored monthly by blood or BM aspirates. All mice were sacrificed at or before reaching ethical endpoints. Blast infiltration was confirmed as detailed above. Pathology analysis confirmed full-blown T-ALL diagnosis, with BM, spleen and liver infiltration by lymphoblasts expressing human CD45 and CD7 markers. Leukemia initiating cell (LIC) frequencies of isogenic Dx-Rel T-ALL PDXs were calculated from the number of engrafted mice for each cell dose in each condition using the WEHI bioinformatic ELDA (Extreme Limiting Dilution Analysis) software[50]. *P* values were calculated with overall tests for differences in LIC frequencies between any of the group tested.

***In vitro* limiting dilution experiment.** Leukemic cells from isogenic Dx-Rel T-ALL PDX samples (primografts) and from paired control mCherry⁺ and *TP53* KD GFP⁺ PDX samples at diagnosis were sorted by flow cytometry based on CD45⁺CD7⁺ (+/- GFP⁺ or mCherry⁺) and seeded on irradiated (30 Gy) confluent stromal cells under limiting dilution conditions from 78.125 to 1 cell per well (24 wells per condition). Leukemic

cells were then cocultured on stromal feeder cells for 2 to 4 weeks under hypoxic
condition (3% $O_2$) as previously described[49]. Cell medium was partially changed twice
a week. Positive wells for leukemic cells outgrowth were visually scored at regular time
intervals. Leukemic cells detection was confirmed by flow cytometry analysis of
recovered wells after 2 to 4 weeks of culture upon hypoxia as detailed above to confirm
the presence of leukemic cells. LIC frequencies were calculated from the number of
positive wells for leukemic cells outgrowth for each cell dose in each condition using
the WEHI bioinformatic ELDA software[50]. *P* values were calculated with overall tests
for differences in LIC frequencies between any of the group tested.

**Real-time measurement of leukemic cells proliferation and apoptosis with the**
**IncuCyte® Live-Cell Analysis System.** Real-time measurement of *in vitro* leukemic
cell proliferation and apoptosis was performed using the IncuCyte® Live-Cell Analysis
System (*Sartorius*). Human $CD45^+CD7^+$ leukemic cells from isogenic Dx-Rel T-ALL
PDX samples (primografts) were sorted by flow cytometry and 100.000 cells were
seeded on 96-well flat bottom plate (4 wells per condition) previously coated with 0.4$\mu$g
of purified human Delta4-Fc fusion protein (*PX'Therapeutics*) according to
manufacturer's recommendation. Leukemic cells were cultured as previously
described[49] for up to 96 hours under normoxia in the IncuCyte® system (5% $CO_2$ and
37 °C). Real-time *in vitro* leukemic cell growth was monitored using live-cell time-lapse
imaging by capturing phase contrast images every 2 hours and analysed using the
integrated confluence algorithm provided. Cell proliferation was represented as the
proportion of phase objects confluence over time, without labels. Leukemic cell
apoptosis was detected using the IncuCyte® Caspase-3/7 green reagent for apoptosis
(*Sartorius*). Briefly, leukemic cells were cultured as previously described in the
presence of 5$\mu$M Caspase-3/7 green reagent. Real-time live-cell imaging of apoptosis
was monitored by capturing phase contrast and green fluorescence images every 2
hours and analysed using the integrated algorithm provided. Leukemic cell apoptosis
was represented as the proprotion of green fluorescently labelled nuclei after activated
caspase-3/7 cleavage of inert Incucyte® Caspase-3/7 dye.

**Apoptosis and cell proliferation end-point assays.** Leukemic cells from isogenic
Dx-Rel PDX T-ALL samples were co-cultured on stromal cells[49] for 48 hours under
normoxia in a humidified 5% $CO_2$ incubator at 37°C. To perform apoptosis assay,

leukemic cells from co-cultures were then collected, washed and stained as detailed above. Leukemic cells were then fixed and stained with PE or Biotin anti-Cleaved Caspase 3 antibody (*BD Biosciences*) as recommended by the manufacturer's recommendations, washed and then analyzed by flow cytometry on FACS Canto II or a LSR Fortessa analyzers (*BD Biosciences*). Cell cycle assays were performed using the APC BrdU Flow kit (*BD Pharmigen*) as recommended by the manufacturer's recommendations. Briefly, leukemic cells co-cultured on stromal cells for 48 hours were pulse-labeled for one hour at 37°C with 1mM Bromodeoxyuridine (BrdU). Leukemic cells from the recovered co-cultures were stained with cell surface antibodies as above. Leukemic cells were then fixed, permeabilized, incubated with DNase and stained with APC-conjugated anti-BrdU antibody (1:100 dilution) for 20 min at room temperature and incubated with Vybrant violet (1:1000 dilution, *Invitrogen*) for 30 min at 37°C before FACS analyses.

***In vivo* homing assay.** Leukemic cells from isogenic Dx-Rel PDXs or from matched *TP53*KD and controls Dx PDXs were injected intravenously ($2\text{-}6\times10^6$ cells/mouse) into sublethally irradiated immunodeficient NSG recipient mice (n = 2 to 4 mice per group). Homing capacities of human leukemic cells were evaluated 24 hours after intravenous injection by FACS analysis of BM, spleen and liver.

***TP53* gene silencing in human leukemic cells and *in vivo* transplantations.** The Hu ShRNA anti-*TP53* was previously described[51]. The human *TP53* cDNA was cloned under the H1 promoter in the backbone pRRL.SIN.cPPT.PGK.GFP lentiviral vector (*Addgene* plasmid #22038). The pRRL_H1shCTL_PGK_mCherry lentivector was constructed in house from the pRRL_H1shLacZ_PGK_eGFP mentioned above. Lentiviral particles were produced as described[52] with jetPRIME reagents (*Polyplus transfection*) by co-transfection of viral production plasmids, according to the manufacturer's protocol. Lentiviral supernatants titration was performed using the qPCR Lentivirus Titer Kit (*Applied Biological Materials*) according to the manufacturer's recommendation.

Leukemic cells from primary PDX samples at diagnosis were cultured on 24-wells plates previously coated with 2.5µg of purified human Delta4-Fc fusion protein (*PX'Therapeutics*) and transduced for 48-72 hours with the lentiviral supernatants including controls (at MOI 20-50) in the presence of 4ug/ml Protamine sulfate (*Sigma*).

Transduced leukemic cells were then washed , GFP- and mCherry-FACS sorted and

immediately intravenously injected into sub-lethally irradiated NSG mice, at equivalent

dose in parallel settings. For competitive experiments, leukemic cells transduced with

a shRNA CTL mCherry or shRNA *TP53*-GFP lentivirus were FACS-sorted, and mixed

in equivalent number before intravenous injection into sub-lethally irradiated NSG

mice. The starting GFP$^+$/mCherry$^+$ ratio was re-checked by flow cytometry before IV-

injection. Gene silencing efficiency was checked by immunoblot (human p53 D0-1,

*Santa Cruz*; Actin,*Santa Cruz*).

**Large-scale gene expression analysis of primary and PDX T-ALL samples by**

**bulk RNA sequencing analysis.** Total RNAs were extracted from frozen pellets of

leukemic blasts at initial diagnosis and first relapse (n= 8 samples) and from BM of fully

leukemic NSG mice for the corresponding derived PDXs (n= 24 PDX samples, first

injection) using Qiagen RNA extraction kit (*Qiagen,* miRNeasy Tissue cell advance)

according to the manufacturer's recommendations. cDNA libraries were prepared

using the TruSeq Stranded mRNA kit (*Illumina*, San Diego, CA, USA). Alignment to

the hg19 (GRCh37) reference genome was performed using STAR aligner (Version

2.7.8a) with quantMode GeneCounts parameter to build raw count matrix for gene-

level summarisation based on the in-built hg19 RefSeq-based annotation.

Normalisation by the method of trimmed mean of M-values (TMM) was performed

using the calcNormFactors function in edgeR. A linear model fitted in limma package

was performed to identify differential gene expression between relapse and diagnosis

samples in basal conditions. Differentially expressed genes were identified based on

the adjusted *p*-value < 0.05 and filtered with a fold change absolute value > 0.5. The

relationship between samples was visualized by hierarchical clustering and principal

component analysis (PCA). Over Representation Analysis (ORA) and Gene set

enrichment analyses (GSEA) were then performed to identify significant - hallmark,

positional (C1), curated (C2), and ontology (C5) - genes sets using the fgsea and

msigdbr packages. Additional analyses on PDX samples from secondary transplants

were performed using the ClariomTM D Assay plateform (*Thermo Fisher Scientific*)

using the manufacturer's recommendations.

**Mitochondrial Respiration and Glycolytic Function Measurements.** Oxygen

consumption rate (OCR) and extracellular acidification rate (ECAR) were respectively

measured with the Seahorse XFp Cell Mito Stress and XFp Glycolysis Stress Test kits on the Seahorse XFp Analyser (*Agilent, Seahorse Bioscience*) in accordance with the manufacturer's recommendations. Leukemic cells recovered from 24h co-culture on stromal cells were seeded at $4 \times 10^5$ cells per well (optimal cell seeding density was determined in preliminary experiments) in an XFp-well miniplate previously coated with Cell-Tak (*Corning, Fisher Scientific*) in 180μl of XF DMEM Medium (pH 7.4), supplemented with 2 mM L-glutamine (*Gibco*) (for OCR and ECAR), 1 mM fresh sodium pyruvate (*Gibco*) (for OCR only) and 10 mM D-glucose (*Sigma*) (for OCR only) on the day of the assay. For OCR measurements, sequential compound injections of Oligomycin (1μM), FCCP (1μM), Rotenone (0.5 μM) and Antimycin A (0.5μM) enabled measurements of basal respiration, ATP production, proton leak, maximal respiration, spare respiratory capacity, and non-mitochondrial respiration. For ECAR measurements, sequential compound injections of Glucose (10mM), Oligomycin (1μM) and 2-Deoxy-glucose (2-DG, 50mM) enabled measurements of glycolysis, glycolytic capacity, glycolytic reserve and non-glycolytic acidification. Key parameters of mitochondrial respiration and glycolytic functions were recorded for 90 minutes after cells were placed into the analyzer. OCR (pmol/min) and ECAR (mpH/min) were measured simultaneously in all wells, three times at each step, by optical fluorescent $O_2$ sensors, and a minimum of three replicates were analyzed per condition in any given experiment. Unless otherwise stated, all compounds and material were obtained from *Agilent Seahorse Bioscience*.

**Cell lines and human T-ALL cells culture.** HEK-293T cell line was purchased from ATCC (CRL-3216). HEK-293T cells were grown in DMEM medium (*Gibco*) supplemented with 2mM L-glutamine (*Gibco*), 1% penicillin/streptomycin (*Gibco*), and 10% heat-inactivated fetal bovine serum (FBS) (*Gibco*). Mouse MS5-Delta-like1[13,53] and human hTERT-immortalized primary bone marrow MSCs[54] derived stromal cells were maintained in αMEM medium (*Gibco*) supplemented with 2mM L-glutamine (*Gibco*), 1% penicillin/streptomycin (*Gibco*), 1μM Hydrocortisone (*Sigma*, for MSCs only) and 10% heat-inactivated FBS (*Gibco*). All cell lines were grown in a humidified 5% $CO_2$ incubator at 37 °C and tested monthly negative for Mycoplasma using MycoAlert PLUS Mycoplasma Detection Kit (*Lonza,*) or MycoStrip Mycoplasma Detection Kit (*Invivogen*).

Primary and PDX human T-ALL leukemic cells were cocultured on stromal cells (MS5-

DL1 or hMSC cells) in αMEM medium (*Gibco*) supplemented with 2mM L-glutamine

(*Gibco*), 1% penicillin/streptomycin (*Gibco*), 10% heat-inactivated FBS (*Stem Cell*

*Technologies*), 10% heat-inactivated human AB serum (*Institut de Biotechnologies*

*Jacques Boy*), 20nM Human Recombinant Insuline (*SAFC Sigma*), 50ng/ml Human

Recombinant SCF (*Miltenyi Biotec*), 10ng/ml Human Recombinant IL7 (*Miltenyi*

*Biotec*), 20ng/ml Human Recombinant Flt3 ligand (*Miltenyi Biotec*) as already

described[13,49].

**High-throughput short read single-cell RNA sequencing analysis of paired Dx-**

**Rel primary and PDX T-ALL samples.** Single-cell RNA-sequencing (scRNAseq)

studies were performed on viable leukemic cells prepared from cryopreserved BM

samples from T-ALL patients and the corresponding derived PDXs. Libraries from

10.000 leukemic cells were prepared using the Chromium Next GEM Single Cell 3'

v3.1 reagent kits (*10x Genomics*) and subjected to short read sequencing on the

Illumina NovaSeq 6000 platform (*Illumina*, San Diego, CA, USA) in a 28-8-91 paired-

end sequencing run (50.000 reads per cell) allowing the captured expression of ~2000

genes per cell. FASTQ files were aligned to the human reference sequence (hg38)

using CellRanger count pipeline (*10× Genomics*, v3.1.0). The output gene-barcode

matrices were imported into the Seurat R package (v4.3.0) for scRNAseq data analysis

and visualization. Cells with nFeature_RNA <500 or >20% mitochondrial genes

content were filtered out. After removing of low-quality cells, data were normalized

using the Normalized-Data function with "LogNormalize" and a scale factor of 10,000.

Highly variable features were identified using the FindVariableFeatures function with

selection method "vst". Data were scaled using the ScaleData function and S.Score

and G2M.Score obtained from CellCycleScoring function were regressed out. Cells

were clustered using FindNeighbors (dims =1:50) with the FindClusters function

(resolution =0.6) based on the shared nearest neighbour. Datasets were visualized

through non-linear dimensional reduction using the Uniform Manifold Approximation

and Projection (UMAP). To identify a list of differentially expressed genes, gene

expression comparisons between clusters and/or samples were performed using the

fast Wilcoxon rank-sum test and area under the receiver operating characteristic curve

(auROC) analysis implemented in the R package Presto (v1.0.0). Gene signature

scores were computed using the AddModuleScore_UCell function from the UCell R

package (v2.6.2), which applies a rank-based scoring method for robust and scalable analysis. To detect large-scale copy number variations in tumor cells, we utilized the inferCNV package (v1.20.0)[55,56]. Gene expression profiles of all relapse cells, or of a defined subset representing the pre-relapse subclone, were contrasted against diagnostic cells to infer chromosomal variations in tumor cell clusters by analyzing expression intensities across genomic locations and thereby explore potential chromosomal alterations.

**TP53-mutant cells backtracking in isogenic primary and PDX T-ALL samples by droplet-digital PCR.**

Droplet-digital PCR (ddPCR) reactions were performed following Bio-Rad recommendation's with 2X ddPCR SuperMix for probes No dUTP and commercially designed ddPCR Mutation Assays for each *TP53* mutation (*Bio-Rad*). Mutation assays are a mix of two fluorescent probes in competition for the annealing on the same target sequence: one specific probe for the wild-type sequence (FAM probe) and a HEX probe, specific of the mutated sequence. Two wet-lab validated ddPCR MUT FAMHEX Assay were used for detection of patient's specific mutations (*TP53 p.R175H* (Assay ID dHsaMDV2010105) and *TP53 p.R248Q* (Assay ID dHsaMDV2010127)) and a non-wet-lab validated assay IS ddPCR MUT FAMHEX Assay was used for *TP53 p.H214R* mutation detection (Assay ID dHsaMDS2510824). Experiments were performed on genomic DNA from isogenic primary and PDX T-ALL samples. Several wells were loaded with DNA from each sample to test enough material. Droplets were generated using QX200 Droplet Generator (*Bio-Rad*). PCR reactions were performed on an Eppendorf Nexus Gradient thermocycler following cycling conditions after validation of optimum annealing/extension temperature: 95°C for 10 min, 40 cycles of two steps PCR 94°C for 30 sec, 55°C for 1 min with a 2°C/sec ramping, an enzyme denaturation step at 98°C for 10 min and a final hold at 4°C. FAM and HEX fluorescence's of each droplet were then read on the QX200 Droplet reader. Visualization, analysis and VAFs calculation were performed with QXOne Manager 1.2 Standard edition software (*Bio-Rad*). Thresholds for each assay were determined aiming maximum specificity, despite sensitivity and quantification accuracy by only incorporated double-positive droplets of a well when at least a single positive-droplet was spotted to avoid potential false double-negative droplets (none as "rain droplets") in the analysis. Non-template controls containing all reagents and the corresponding amount of carrier DNA to be

equivalent to the tested samples were included in the analysis. VAFs were calculated

by considering all the wells analysed.

**Statistical analyses.** Statistical analyses were performed with GraphPad Prism

(version 10.3; *GraphPad Software Inc.*) or R packages using Mann-Whitney, Anova,

Wilcoxon and Kruskal-Wallis tests, as indicated in figure legends. Unless otherwise

stated, the data are expressed as mean±s.d. of *n*=3 or more determinations.

For xenotransplantation analysis, Kaplan–Meier survival curves were performed and

the statistical significance was determined with the Log-rank (Mantel-Cox) test. Overall

survival was defined as the time from cell injection to euthanasia or death at predefined

endpoints. Animals were censored if the cause of death/sacrifice was not leukemia

related. Number of mice and median survival are indicated into brackets.

For limiting dilution assays, statistical analysis was performed using the WEHI

bioinformatic ELDA (Extreme Limiting Dilution Analysis) software[50]). Differences were

considered statistically significant at *p*<0.05 (*), *p*<0.01 (**), *p*<0.001 (***) or *p*<0.0001

(****).

**Authors' Contributions**

Conception and design of the study: SG, EC and JS

Development of methodology: SG, SQ, LH and JS

Acquisition of data (provided materials and animals, acquired and managed patients, provided facilities): SG, SQ, LH, LM, MP, RK, HB, MB, VP, PF, AB, HD, NB, EC and JS

Analysis and interpretation of data (biostatistics, computational analysis, statistical analysis): SG, SQ, LH, LM, MB, FS, EC and JS

Writing of the manuscript - original draft preparation, review and editing: SG, HdT and JS

Grants acquisition and study supervision: SG and JS

All authors critically reviewed the manuscript.

**Disclosure of potential conflicts of interest**

The authors declare no potential conflicts of interest.

**Acknowledgments**

The authors thank Niclas Setterblad, Christelle Doliger, Antonio Alberdi, Sophie Duchez and Claire Maillard from the Technological core facility of the Saint-Louis Research Institut, Inserm UMR1342 / CNRS EMR8000; the ICGex NGS platform of the Curie Institut for high-throughput sequencing of *10X* cDNA librairies.

**Fundings**

This work was supported by grants to Jean Soulier from the ERC St Grant Consolidator #311660, the team FRM 2022 program from the Fondation pour la Recherche Médicale #00107939, the Ligue Contre le Cancer IDF, the Fondation Leucémie Espoir, and by grants to Stéphanie Gachet from the Fondation de France, the Fondation ARC pour la Recherche sur le Cancer (PJA 20191209702), and the Paris Saint-Louis Leukemia Institute program (a grant managed by the French National Research Agency under the France 2030 program, ANR-23-IAHU-0005, Institut de la Leucémie Paris Saint-Louis). The ICGex platform was supported by the grants from the French National Research Agency, the Canceropole Ile-de-France and by tge SiRIC-Curie program.

645 **Data-sharing statement**

646 The data that support the findings of this study are available from the corresponding

647 authors, [SG and JS], upon reasonable request.

## REFERENCES

1. Van Vlierberghe, P. & Ferrando, A. The molecular basis of T cell acute lymphoblastic leukemia. *J Clin Invest* **122**, 3398–406 (2012).
2. Buckley, M., Yeung, D. T., White, D. L. & Eadie, L. N. T-cell acute lymphoblastic leukaemia: subtype prevalence, clinical outcome, and emerging targeted treatments. *Leukemia* **39**, 1294–1310 (2025).
3. Weng, A. P. *et al.* Activating mutations of NOTCH1 in human T cell acute lymphoblastic leukemia. *Science* **306**, 269–71 (2004).
4. Ferrando, A. A. *et al.* Gene expression signatures define novel oncogenic pathways in T cell acute lymphoblastic leukemia. *Cancer Cell* **1**, 75–87 (2002).
5. Soulier, J. *et al.* HOXA genes are included in genetic and biologic networks defining human acute T-cell leukemia (T-ALL). *Blood* **106**, 274–86 (2005).
6. Girardi, T., Vicente, C., Cools, J. & De Keersmaecker, K. The genetics and molecular biology of T-ALL. *Blood* **129**, 1113–1123 (2017).
7. Pölönen, P. *et al.* The genomic basis of childhood T-lineage acute lymphoblastic leukaemia. *Nature* **632**, 1082–1091 (2024).
8. Desjonqueres, A. *et al.* Acute lymphoblastic leukemia relapsing after first-line pediatric-inspired therapy: a retrospective GRAALL study. *Blood Cancer J* **6**, e504 (2016).
9. Heikamp, E. B. & Pui, C. H. Next-Generation Evaluation and Treatment of Pediatric Acute Lymphoblastic Leukemia. *J Pediatr* **203**, 14-24 e2 (2018).
10. Pagliaro, L. *et al.* Acute lymphoblastic leukaemia. *Nat. Rev. Dis. Primer* **10**, 41 (2024).
11. Brady, S. W. *et al.* The genomic landscape of pediatric acute lymphoblastic leukemia. *Nat. Genet.* **54**, 1376–1389 (2022).
12. Pölönen, P., Mullighan, C. G. & Teachey, D. T. Classification and risk stratification in T-lineage acute lymphoblastic leukemia. *Blood* **145**, 1464–1474 (2025).
13. Clappier, E. *et al.* Clonal selection in xenografted human T cell acute lymphoblastic leukemia recapitulates gain of malignancy at relapse. *J Exp Med* **208**, 653–61 (2011).
14. Mullighan, C. G. *et al.* Genomic analysis of the clonal origins of relapsed acute lymphoblastic leukemia. *Science* **322**, 1377–80 (2008).
15. Tosello, V. *et al.* WT1 mutations in T-ALL. *Blood* **114**, 1038–1045 (2009).
16. Anderson, K. *et al.* Genetic variegation of clonal architecture and propagating cells in leukaemia. *Nature* **469**, 356–61 (2011).
17. Kunz, J. B. *et al.* Pediatric T-cell lymphoblastic leukemia evolves into relapse by clonal selection, acquisition of mutations and promoter hypomethylation. *Haematologica* **100**, 1442–50 (2015).
18. Wong, T. N. *et al.* Role of TP53 mutations in the origin and evolution of therapy-related acute myeloid leukaemia. *Nature* **518**, 552–555 (2015).
19. Oshima, K. *et al.* Mutational landscape, clonal evolution patterns, and role of RAS mutations in relapsed acute lymphoblastic leukemia. *Proc Natl Acad Sci U A* **113**, 11306–11311 (2016).
20. Meyer, J. A. *et al.* Relapse-specific mutations in NT5C2 in childhood acute lymphoblastic leukemia. *Nat. Genet.* **45**, 290–294 (2013).
21. Tzoneva, G. *et al.* Activating mutations in the NT5C2 nucleotidase gene drive chemotherapy resistance in relapsed ALL. *Nat Med* **19**, 368–71 (2013).
22. Tzoneva, G. *et al.* Clonal evolution mechanisms in NT5C2 mutant-relapsed acute lymphoblastic leukaemia. *Nature* **553**, 511–514 (2018).
23. Dieck, C. L. *et al.* Structure and Mechanisms of NT5C2 Mutations Driving Thiopurine Resistance in Relapsed Lymphoblastic Leukemia. *Cancer Cell* **34**, 136-147.e6 (2018).
24. Barz, M. J. *et al.* Subclonal NT5C2 mutations are associated with poor outcomes after relapse of pediatric acute lymphoblastic leukemia. *Blood* **135**, 921–933 (2020).
25. Diccianni, M. B. *et al.* Clinical significance of p53 mutations in relapsed T-cell acute lymphoblastic leukemia. *Blood* **84**, 3105–12 (1994).

26. Hof, J. *et al.* Mutations and deletions of the TP53 gene predict nonresponse to treatment
and poor outcome in first relapse of childhood acute lymphoblastic leukemia. *J Clin*
*Oncol* **29**, 3185–93 (2011).
27. Hof, J. *et al.* NOTCH1 mutation, TP53 alteration and myeloid antigen expression predict
outcome heterogeneity in children with first relapse of T-cell acute lymphoblastic
leukemia. *Haematologica* **102**, e249–e252 (2017).
28. Richter-Pechańska, P. *et al.* Identification of a genetically defined ultra-high-risk group in
relapsed pediatric T-lymphoblastic leukemia. *Blood Cancer J.* **7**, e523 (2017).
29. Simonin, M. *et al.* Prognostic value and oncogenic landscape of TP53 alterations in adult
and pediatric T-ALL. *Blood* **141**, 1353–1358 (2023).
30. Kempter, T. *et al.* Subclonal TP53 and KRAS variants combined with poor treatment
response identify ultrahigh-risk pediatric patients with T-ALL. *Blood Adv.* **9**, 1267–1279
(2025).
31. Li, B. *et al.* Therapy-induced mutations drive the genomic landscape of relapsed acute
lymphoblastic leukemia. *Blood* https://doi.org/10.1182/blood.2019002220 (2020)
doi:10.1182/blood.2019002220.
32. Armstrong, F. *et al.* NOTCH is a key regulator of human T-cell acute leukemia initiating
cell activity. *Blood* **113**, 1730–1740 (2009).
33. Liu, Y., Su, Z., Tavana, O. & Gu, W. Understanding the complexity of p53 in a new era of
tumor suppression. *Cancer Cell* **42**, 946–967 (2024).
34. Dong, Y., Tu, R., Liu, H. & Qing, G. Regulation of cancer cell metabolism: oncogenic
MYC in the driver's seat. *Signal Transduct. Target. Ther.* **5**, 124 (2020).
35. Richter-Pechańska, P. *et al.* Pediatric T-ALL type-1 and type-2 relapses develop along
distinct pathways of clonal evolution. *Leukemia* **36**, 1759–1768 (2022).
36. Albertí-Servera, L. *et al.* Single-cell DNA amplicon sequencing reveals clonal
heterogeneity and evolution in T-cell acute lymphoblastic leukemia. *Blood* **137**, 801–811
(2021).
37. Calvo, J. *et al.* High CD44 expression and enhanced E-selectin binding identified as
biomarkers of chemoresistant leukemic cells in human T-ALL. *Leukemia* **39**, 323–336
(2025).
38. Xu, J. *et al.* A multiomic atlas identifies a treatment-resistant, bone marrow progenitor-
like cell population in T cell acute lymphoblastic leukemia. *Nat. Cancer* **6**, 102–122
(2025).
39. Ebinger, S. *et al.* Characterization of Rare, Dormant, and Therapy-Resistant Cells in
Acute Lymphoblastic Leukemia. *Cancer Cell* **30**, 849–862 (2016).
40. Costea, J. *et al.* Role of stem-like cells in chemotherapy resistance and relapse in
pediatric T-cell acute lymphoblastic leukemia. *Nat. Commun.* **16**, 5413 (2025).
41. Hsu, J. I. *et al.* PPM1D Mutations Drive Clonal Hematopoiesis in Response to Cytotoxic
Chemotherapy. *Cell Stem Cell* **23**, 700-713.e6 (2018).
42. Chen, S. *et al.* Mutant p53 drives clonal hematopoiesis through modulating epigenetic
pathway. *Nat. Commun.* **10**, 5649 (2019).
43. Rodriguez-Zabala, M. *et al.* Combined GLUT1 and OXPHOS inhibition eliminates acute
myeloid leukemia cells by restraining their metabolic plasticity. *Blood Adv.* **7**, 5382–5395
(2023).
44. Chiu, P. P. L., Jiang, H. & Dick, J. E. Leukemia-initiating cells in human T-lymphoblastic
leukemia exhibit glucocorticoid resistance. *Blood* **116**, 5268–5279 (2010).
45. Belmonte, M., Hoofd, C., Weng, A. P. & Giambra, V. Targeting leukemia stem cells:
which pathways drive self-renewal activity in T-cell acute lymphoblastic leukemia? *Curr*
*Oncol* **23**, 34–41 (2016).
46. Gerby, B. *et al.* High-throughput screening in niche-based assay identifies compounds to
target preleukemic stem cells. *J. Clin. Invest.* **126**, 4569–4584 (2016).
47. Yadav, B. D. *et al.* Heterogeneity in mechanisms of emergent resistance in pediatric T-
cell acute lymphoblastic leukemia. *Oncotarget* **7**, 58728–42 (2016).
48. Ariës, I. M. *et al.* PRC2 loss induces chemoresistance by repressing apoptosis in T cell
acute lymphoblastic leukemia. *J. Exp. Med.* **215**, 3094–3114 (2018).

49. Gachet, S. *et al.* Deletion 6q Drives T-cell Leukemia Progression by Ribosome
Modulation. *Cancer Discov* **8**, 1614–1631 (2018).
50. Hu, Y. & Smyth, G. K. ELDA: extreme limiting dilution analysis for comparing depleted
and enriched populations in stem cell and other assays. *J. Immunol. Methods* **347**, 70–78
(2009).
51. Ceccaldi, R. *et al.* Bone marrow failure in Fanconi anemia is triggered by an exacerbated
p53/p21 DNA damage response that impairs hematopoietic stem and progenitor cells.
*Cell Stem Cell* **11**, 36–49 (2012).
52. Sebert, M. *et al.* Clonal hematopoiesis driven by chromosome 1q/MDM4 trisomy defines
a canonical route toward leukemia in Fanconi anemia. *Cell Stem Cell* **30**, 153-170.e9
(2023).
53. Gerby, B. *et al.* Optimized gene transfer into human primary leukemic T cell with NOD-
SCID/leukemia-initiating cell activity. *Leukemia* **24**, 646–9 (2010).
54. Mihara, K. *et al.* Development and functional characterization of human bone marrow
mesenchymal cells immortalized by enforced expression of telomerase. *Br. J. Haematol.*
**120**, 846–849 (2003).
55. Patel, A. P. *et al.* Single-cell RNA-seq highlights intratumoral heterogeneity in primary
glioblastoma. *Science* **344**, 1396–1401 (2014).
56. Zhu, Q. *et al.* Single cell multi-omics reveal intra-cell-line heterogeneity across human
cancer cell lines. *Nat. Commun.* **14**, 8170 (2023).
776
777

**FIGURE LEGENDS**

**Figure 1. *In vivo* isogenic modelling of diagnosis and relapse *TP53*-altered T-ALL. A.** Experimental design (Created in BioRender. Gachet, S. (2026) https://BioRender.com/zczr8er). Isogenic Dx-Rel PDX pairs were established after injection into immunodeficient NSG mice of paired primary diagnosis and *TP53*-altered (*TP53*$^{alt}$) relapse samples from the same T-ALL patients. **B.** Comparative Whole Exome Sequencing (WES) analysis of the isogenic Dx-Rel PDX samples established from 4 T-ALL patients who relapsed with *17p/TP53* alterations. In each case, PDX samples (PDXD and PDXR) are compared side-by-side to the corresponding primary samples (TALLD and TALLR, respectively). **C.** Lollipop plot of TP53 mutations observed in four primary relapse samples (TALL11R, TALL30R, TALL42R and TALL43R).

**Figure 2. *TP53*$^{alt}$ T-ALL relapse cells display a cell-intrinsic gain of malignancy phenotype. A.** Kaplan-Meier survival curves of PDX mice engrafted with primary isogenic Dx-Rel leukemic cells from 4 T-ALL patients (TALL11, TALL30, TALL42 and TALL43). N=4-5 recipient mice injected at equivalent cell doses per sample. **B-C.** Parallel serial transplantations of paired Dx-Rel T-ALL PDX samples. Kaplan-Meier survival curves of mice injected with 3 cell doses ($10^6$, $10^5$ and $10^4$ cells) of isogenic Dx-Rel PDX pairs (n = 4 TALL cases) raised from primary (**B**) and secondary (**C**) xenografts. **D.** LIC frequencies of isogenic Dx-Rel PDX sample pairs calculated from *in vivo* limiting dilutions. **E-F.** Real-time *in vitro* cell proliferation and apoptosis of leukemic cells from isogenic Dx-Rel PDX pairs, measured with the Incucyte® Live-Cell Analysis System. Proliferation was quantified by measuring the proportion of phase objects confluence over time, without labels **(E)**, and apoptosis by the quantification of green fluorescently labelled nuclei after activated caspase-3/7 cleavage of inert Incucyte® Caspase-3/7 dyes **(F)**.

Statistical data in this figure are represented as mean ± s.d. Significance is indicated by *P* values; A-C: Log-rank test (Mantel-Cox); D: overall tests for differences using ELDA software; E-F: two-way Anova test. Number of recipient mice and median survival are indicated into brackets on Kaplan-Meier survival curves. ns, not significant; *, *p* < 0.05; **, *p* < 0.01; ***, *p* < 0.001; ****, *p* < 0.0001.

**Figure 3.** *TP53* **silencing confers a** *relapse phenotype* **to diagnosis leukemic cells. A.** Experimental design of *TP53* silencing in diagnosis cells from PDX samples using a lentiviral shRNA approach (Created in BioRender. Gachet, S. (2026) https://BioRender.com/usb6t9c). **B-C.** Primary transplantation of paired control mCherry⁺ (shCTL mCherry) and *TP53* KD GFP⁺ (sh *TP53* GFP) leukemic cells from PDX30D injected in parallel settings into recipient mice. **B.** Engraftment kinetics in the bone marrow (BM) of recipient mice (n=6) injected with paired control and *TP53* KD leukemic cells from PDX30D. **C.** Kaplan-Meier survival curves of mice injected with paired control and *TP53* KD leukemic cells from PDX30D. **D-F.** Secondary transplant of paired control mCherry⁺ (shCTL mCherry) and *TP53* KD GFP⁺ (sh *TP53* GFP) leukemic cells from PDX30D injected either in parallel (panels D-E) or competitive (panel F) settings into recipient mice. **D.** Engraftment kinetics in the BM of recipient mice (n=5) injected in parallel settings with cells from control and *TP53* KD conditions. **E.** Kaplan-Meier survival curves of mice injected in parallel settings with sorted control or *TP53* KD leukemic cells from PDX30D. **F.** Engraftment kinetics in the BM of recipient mice (n=4) injected with sorted cells from both control and *TP53* KD in competitive settings (ratio 1:1). Data are represented as ratios of *TP53* KD GFP⁺ over control mCherry⁺ leukemic cells within BM of the same recipient mouse and compared to the input. **G.** LIC frequencies of paired control-*TP53* KD samples calculated from *in vitro* limiting dilutions of sorted GFP⁺ or mCherry⁺ CD45⁺CD7⁺ leukemic cell from diagnosis PDX samples (PDX30D and PDX42D). LIC frequencies are estimated based on the proportions of positive wells for GFP⁺/mCherry⁺ CD45⁺CD7⁺ leukemic cells evaluated after 2 and 4 weeks of culture.

Statistical data in this figure are represented as mean ± s.d. Significance is indicated by *P* values; B and D: Mann-Whitney test; C and E: Log-rank test (Mantel-Cox); F: Kruskal-Wallis test; G: overall tests for differences using ELDA software. Number of recipient mice and median survival are indicated into brackets on Kaplan-Meier survival curves. ns, not significant; *, $p < 0.05$; **, $p < 0.01$; ****, $p < 0.0001$.

**Figure 4.** *TP53* **alterations drive a** *relapse phenotype* **through MYC signaling and metabolism deregulation. A.** Supervised hierarchical clustering of the top fifty most differentially expressed genes between Dx and Rel T-ALL samples. Heatmap representation of the most variables genes at Dx and Rel in 4 isogenic PDX pairs (primografts). *TP53* expression is highlighted in red. Independent triplicates are

analysed for each isogenic PDX pair. **B-C.** GSEA of enriched pathways in *TP53*alt T-

ALL relapse samples. **B.** Comparative Gene Set Enrichment Analysis (GSEA) of

differentially expressed genes in Rel and Dx samples from 4 isogenic PDX pairs.

Representation of selected Hallmark pathways significantly upregulated at relapse,

defining a *relapse expression signature* in all isogenic Dx-Rel PDX pairs pooled

together (1st column "All_PDXs") and in each individual PDX pair (2nd to 5th columns;

respectively PDX11, PDX30, PDX42 and PDX43) (*P* adj <0.05). **C.** GSEA of the

upregulated (upper panel) and downregulated (lower panel) *relapse expression*

*signature* obtained from PDX samples (panel B) tested in the original T-ALL patients

samples. UP_PDX, upregulated transcriptomic signature in Rel PDXs; DOWN_PDX,

downregulated transcriptomic signature in Rel PDXs. FDR, false discovery rate; NES,

normalized enrichment score. **D-G.** Direct evaluation of mitochondrial respiration in

leukemic cells from isogenic Dx-Rel PDXs using Seahorse assays. **D.** Real-time

measurements of oxygen consumption rate (OCR) in isogenic Dx-Rel leukemic cells

from 4 T-ALL cases. **E.** Quantification of maximal respiration in leukemic cells from 4

isogenic Dx-Rel PDX pairs evaluated in 3 independent experiments. **F.** Real-time

measurements of OCR in *TP53*-silenced (*TP53* KD) and matched control leukemic

cells from diagnosis PDX samples (PDX42D and PDX43D). **G.** Quantification of

maximal respiration in *TP53* KD and control leukemic cells from 3 independent

diagnosis PDX samples (PDX30D, PDX42D and PDX43D) evaluated in 2 independent

experiments

Statistical data in this figure are represented as mean $\pm$ s.d. Significance is indicated

by *P* values; E and G: Mann-Whitney test. ns, not significant; **, *p* < 0.01; ***, *p* <

0.001; ****, *p* < 0.0001.

**Figure 5. Single-cell profiling of isogenix Dx-Rel T-ALLs can unveil a minor pre-**

**relapse subclone at diagnosis. A.** UMAP visualizations of isogenic Dx-Rel primary

samples from 4 T-ALL patients and their derived PDXs analysed by short read single-

cell RNA (scRNA) sequencing. UMAP plots of 117.119 single cells coloured by T-ALL

samples (n=16, primary and PDXs samples from four T-ALL cases at diagnosis and

relapse; left panel), leukemic cell identity (Diagnosis versus Relapse; middle panel)

and by leukemic cell origin (Primary versus PDX; right panel). **B-C.** GSEA of biological

pathways enriched at relapse in primary and PDX T-ALL samples. GSEA from pseudo

bulk analysis of differentially expressed genes between relapse (Rel) and diagnosis

(Dx) samples from isogenic patient (**B**) and PDX (**C**) T-ALL pairs retrieves the *relapse*

*expression signature* defined in Figure 4B. Representation of selected Hallmark

pathways significantly deregulated at relapse (*P* adj <0.05) in all isogenic pairs pooled

together (1st column ") and in each individual pair (2nd to 5th columns). **D.** UMAP

visualizations of Dx and Rel primary samples from TALL42 case analysed by short

read scRNA sequencing. UMAP plots of 14.049 single leukemic cells coloured by cell

identity (Diagnosis vs Relapse; upper panel) and cluster-based by transcriptomic data

(fourteen different clusters; lower panel). The red dotted ellipse highlights cluster 6 in

TALL42D sample. **E.** Dot plot visualization of biomarker genes average expression

selected from the unsupervised differential expression analysis between Rel and Dx

samples from TALL42 case. Average expression and proportion of expressing cells for

selected biomarkers are represented in each cluster from Dx and Rel samples. The

red dotted line highlights the "pre-relapse" clone. **F-G.** GSEA of positively and

negatively enriched pathways at relapse and "pre-relapse" stages in TALL42 samples.

**F.** Representation of selected Hallmark pathways significantly deregulated in relapse

(1st column Rel vs Dx; *P* adj <0.05) and "pre-relapse" (2nd column Sub vs Dx, red dotted

line; *P* adj <0.05) clusters as compared to matched diagnosis counterparts. **G.**

Representation of specific gene sets from the MSigDB C1 and C2 collections

significantly upregulated (in red) or downregulated (in blue) at relapse (1st column Rel

vs Dx; *P* adj <0.05) and "pre-relapse" (2nd column Sub vs Dx; *P* adj <0.05) as compared

to matched primary diagnosis counterparts. Color code indicates the cell proliferation,

invasiveness, stemness and oxidative phosphorylation signatures. **H.** Single-cell

scores for several leukemia stem cell (LSC) signatures in TALL42 diagnosis (Dx), pre-

relapse (Sub) and relapse (Rel) clusters. The box includes the median, hinges mark

the 25th and 75th percentiles. *P* values from a two-sided paired t-test are shown. ns,

not significant; ****, *p* < 0.0001 **I.** CNV inference analysis of cluster 6 ('Sub'; "pre-

relapse" subpopulation) from TALL42D and all the clusters from TALL42R (clusters 0,

3, 4 and 9; 'Rel') interrogating copy number alterations on each chromosome (lower

part) as compared to the remaining diagnosis sample (clusters 1, 2, 5, 7, 10 and 11;

'Dx') as a reference (upper part). The red circle highlights the CNV status for

chromosome 17p in the pre-relapse clone (cluster 6) from TALL42D.

# Figure 1

**A**

Primary Diagnosis

T-ALL patient

Primary *TP53*-altered Relapse

Xenotransplantation NSG mice

PDX Diagnosis

PDX *TP53*-altered Relapse

Isogenic PDX pairs

**C**

**B**

# Figure 2

# Figure 3

**A**

**B**

Primary injection
PDX30D

■ sh CTL mCherry (n=6)
▲ sh *TP53* GFP (n=6)

**C**

Primary injection
PDX30D

– – sh CTL mCherry (n=6; 150.5 days)
– – sh *TP53* GFP (n=6; 128 days)

*p* = 0.0061

**D**

Secondary injection
PDX30D

■ sh CTL mCherry (n=6)
▲ sh *TP53* GFP (n=6)

**E**

Secondary injection
PDX30D

– – sh CTL mCherry (n=5; 150 days)
– – sh *TP53* GFP (n=5; 102 days)

*p* = 0.0018

**F**

Secondary injection
PDX30D

**G**

# Figure 4

# Figure 5