***In vivo* isogenic modelling unveils *TP53*-associated relapse trajectories in T-cell**

acute lymphoblastic leukemia

Stéphanie Gachet[1,2,3,#], Samuel Quentin[1,2,4*], Lucie Hernandez[1,2*], Loïc Maillard[1,2],

Marie Passet[1,2,4], Rathana Kim[1,2,3,4], Hugo Bergugnat[1,2], Melha Benlebna[1,2], Rémy

Jelin[5], Marine Aglave[5], Maxime Boy[1,2], Véronique Parietti[1,2], Pierre Fenaux[6], André

Baruchel[7], Hervé Dombret[1,4,6], Nicolas Boissel[1,2,3,8], François Sigaux[1,2,3,4], Hugues de

Thé[1,2,3,4], Emmanuelle Clappier[1,2,3,4] and Jean Soulier[1,2,3,4].

[1]Institut de Recherche Saint-Louis (IRSL), Université Paris Cité, 75010 Paris, France;

[2]INSERM UMR1342 / CNRS EMR8000, Paris, France;

[3]Institut de la Leucémie, Hôpital Saint-Louis, APHP, Université Paris Cité, Paris,

France;

[4]Hôpital Saint-Louis, Laboratoire d'Hématologie, APHP, Paris, France;

[5]Gustave Roussy, Département de la recherche, F-94805, Villejuif, France;

[6]Hôpital Saint-Louis, Département Hématologie Senior, APHP, Paris, France;

[7]Hôpital Robert Debré, APHP, Université Paris Cité, Paris, France;

[8]Hôpital Saint-Louis, Service d'Hématologie Adolescents et Jeunes Adultes, APHP,

Paris, France.

[*]These co-authors contributed equally

Corresponding authors:

Drs. Stéphanie Gachet, PhD: stephanie.gachet@inserm.fr and Jean Soulier, MD, PhD:

jean.soulier@aphp.fr

[#]Present address: Laboratoire des cellules Souches Hématopoïétiques et des

Leucémies, Equipe Niche, Cancer et Radiations dans l'Hématopoïèse, Inserm U1274,

Commissariat à l'Energie Atomique et aux Energies Alternatives (CEA), Université

Paris-Saclay.

Running title:

TP53-altered relapse trajectories in T-ALL

Keywords: T-cell acute lymphoblastic leukemia, somatic *TP53* alterations, relapse, leukemia initiating cells, metabolic adaptation

Word counts:

Abstract, 150; Text, 3382

Number of Figures: 5

Footnotes:

Supplementary informations for this article (Supplementary Figures, 5 and Supplementary Tables, 3) are available online.

ABSTRACT

Many patients with T-cell acute lymphoblastic leukemia (T-ALL) relapse into a treatment-resistant disease. The mechanisms driving relapse remain largely elusive, in part due to the lack of faithful experimental models. Here, we leveraged patient-derived xenograft (PDX) pairs generated from diagnosis and relapse T-ALLs to functionally address the cellular mechanisms driving *TP53*-altered relapse. Beyond inter-T-ALL variability, comparative analyses revealed a unique, cell-intrinsic relapse phenotype that includes greater leukemia-initiating capacity and that can be conferred to diagnosis cells by *TP53* silencing. Transcriptomic profiling linked the relapse phenotype to deregulated OXPHOS metabolism and MYC signaling. Integration of single-cell profilings uncovered *TP53*-wildtype cell populations at diagnosis expressing a relapse profile, possibly reflecting a pre-existing modulation of *TP53* signaling. These cells sequentially evolved towards biallelic *TP53* inactivation at relapse. Collectively, our findings support a model in which T-ALL relapses emerge from a selected pre-existing transcriptional state characterized by deregulated metabolism that favors subsequent *TP53* inactivation.

INTRODUCTION

T-cell acute lymphoblastic leukemia (T-ALL) is an aggressive hematologic malignancy that represents 15% of pediatric and 25% of all adult ALL cases[1,2]. Multistep deregulation of key transcription factors and genes controlling self-renewal, senescence, apoptosis, differentiation or proliferation in a T-cell precursor leads to clonal transformation and expansion of blast cells in the lymphoid organs and the bone marrow[1,3–7]. Although most T-ALL patients at diagnosis achieve remission with intensive chemotherapy and glucocorticoids, 15% to 40% of the patients subsequently experience relapse into a resistant leukemia with dismal prognosis[8–10]. Therefore, comprehensive understanding of the functional mechanisms underlying relapse and resistance is a major unmet need to develop more efficient therapeutic strategies. While exhaustive genetic and immunophenotypic landscape studies have enabled to identify a large variety of T-ALL subtypes[2,7,11,12], these last only partially correlate with the onset of relapse. It rather appears that resistance initiates upon treatment by the selection of residual T-ALL cells with cellular features driving cell survival despite therapy and allowing re-expansion[13–19]. Selection of *NT5C2* somatic mutations leading

to cell resistance to 6-mercaptopurine is a well-established, although rare, example of

resistance-driven mechanism in T- and B-ALL patients[20–24], but alternative

mechanisms remain elusive.

Here, we investigated *TP53* alterations (*TP53*[alt], i.e. *TP53* somatic mutations and/or

deletions) as a factor of resistance acquisition in T-ALL. While rarely detected at

diagnosis (2-3%), *TP53*[alt] are frequently found at relapse (up to 25%) and are

associated to poor clinical outcomes[25–30]. *TP53*[alt] cells are genetically trackable,

allowing to detect and characterize them at various stages of T-ALL progression

(diagnosis, remission or relapse). Using *in vivo* isogenic patient-derived xenografts

(PDX) of diagnosis and relapse T-ALL pairs, we unveiled a unique relapse phenotype

that included increased leukemia initiating activity and modulation of key signaling and

metabolic pathways such as MYC and OXPHOS. Integration of multi-omics and

functional data, including single-cell analyses, allowed us to delineate the relapse

trajectory of rare, therapy-selected cells pre-existing at diagnosis that subsequently

expanded sequentially and drove relapse.

RESULTS

Isogenic, patient-derived xenografts (PDXs) faithfully capture T-ALL diagnosis

and *TP53*[alt]-driven relapse genomic patterns

We aimed at uncovering functional mechanisms involved in leukemia resistance and

relapse. We focused on *TP53*-altered T-ALL since *TP53* alterations are well-

recognized, genetically-trackable, drivers of relapse but are still poorly mechanistically

understood in T-ALL[25–29,31]. We selected four cases of T-ALL with *TP53* alterations at

relapse only for which cryopreserved diagnosis and relapse patient samples were

available. To overcome the limitations of *ex vivo* functional studies on primary samples

or cell lines, we established PDXs of paired diagnosis (Dx) and relapse (Rel) leukemia

in immunodeficient NOD.Cg-*Prkdc$_{scid}$ IL2rg$_{tm1Wjl}$*/SzJ (NSG) mice (**Figure 1A**).

Genomic analyses of the resulting isogenic, Dx-Rel PDX pairs showed that these

PDXs largely captured the genomic pattern of the patient cells, both at diagnosis and

at relapse, respectively (**Figure 1B** and **Supplementary Tables 1-2**). *TP53* alterations

in patient T-ALL relapse cells were missense mutations in the *TP53* DNA-binding

domain within mutational hotspot residues R175 and R248 (cases TALL42; and

TALL11 and TALL43, respectively) and a less classical but already reported

pathogenic mutation H214 (case TALL30) (**Figure 1C**). The other allele was lost by 17p deletion in 3 cases (TALL30, TALL42 and TALL43) or by 17p uniparental disomy of the mutated allele in 1 case (TALL11) (**Figure 1B**). The resulting PDXs carried the original *TP53* missense mutations in association with the loss of the remaining wild-type (WT) allele, as in primary patient samples, or in 1 case only the deleted allele (PDX43R).

Interestingly, other somatic copy number variants (CNVs) and point mutations (i.e., in addition to *TP53*[alt]) were largely similar in both the diagnosis and relapse samples in the T-ALL pairs, suggesting that *TP53*-driven T-ALL relapse did not acquire stronger genomic instability (**Figure 1B**). Therefore, these T-ALL PDX models faithfully captured the diagnosis and the relapse patient samples, respectively, enabling further functional comparison experiments using this unlimited ressources of isogenic, human leukemic cells.

Relapse cells display a cell-intrinsic, stable, relapse phenotype compared to their isogenic diagnosis cells

To uncover functional mechanisms involved in leukemia resistance and relapse, we compared the leukemia propagation ability of the diagnosis and relapse T-ALL pairs. First, primary leukemic cells from paired Dx-Rel patient samples were transplanted in parallel at equivalent cell doses in NSG mice, and leukemia onset and mice survival were monitored. Despite strong inter-patient T-ALL heterogeneity, we found in each pair that the relapse cells propagated leukemia more efficiently than their diagnosis counterparts (**Supplementary Figure 1A**), which translated into a significantly shorter mice survival (**Figure 2A** and **Supplementary Figure 1B**). This increased leukemia propagating activity at relapse was observed in all PDXs established from the 4 pairs. We performed multiple iterative transplantations using decreasing cell doses (**Figures 2B-C** and **Supplementary Figures 1C-F**) and found that the propagation phenotype was stable upon serial reinjection, without additional acceleration (**Supplementary Figures 1D,1F**). Limiting dilution *in vivo* assays (injection of 10^6 to 1 leukemic cell/mouse) showed that the leukemia initiating cell (LIC) frequencies were increased in the Rel PDXs compared to Dx counterparts (**Figure 2D** and **Supplementary Table 3**). Consistently, *in vitro* long-term culture-initiating cell assays in coculture with MS5-DL1 stromal cells[32] showed a significantly increased clonogenicity in Rel PDXs from all T-ALL pairs, demonstrating that increased LIC activity is a cell-intrinsic feature of

relapse cells (**Supplementary Figure 1G** and **Supplementary Table 3**). Real-time
and end-point *in vitro* proliferation using Bromodeoxyuridine (BrdU) pulse labelling and
apoptosis experiments, in particular cleaved caspase 3 staining, showed increased
proliferation and decreased apoptosis in Rel versus Dx PDX cells (**Figures 2E-F** and
Supplementary Figures 1H-I). Moreover, homing of the leukemic cells in the bone
marrow, spleen and liver at 24 hrs was increased in PDXs from Rel for the 4 T-ALL
pairs (**Supplementary Figures 1J-K**). However, when homing was bypassed by direct
injection of leukemic cells into the femoral bone marrow cavity, leukemic cells from Rel
PDXs still propagated leukemia more efficiently than Dx PDXs (**Supplementary**
Figures 1L-M).
Collectively, our data indicate that despite inter-T-ALL heterogeneity, $TP53^{alt}$ mediate
a unique, cell-intrinsic, relapse phenotype that is associated with increased
proliferation, cell survival, and cell homing and post-homing advantage, resulting in
higher LIC activity.
162
Experimental *TP53* silencing confers the cell-intrinsic relapse phenotype to
diagnosis leukemic cells
To model *TP53* alterations as driver of the relapse phenotype, we inactivated *TP53*
using a lentiviral shRNA approach in Dx PDX leukemic cells followed by transplantation
in NSG mice (**Figure 3A** and **Supplementary Figure 2A**). *TP53* downregulation
increased the leukemia propagation capacities of Dx PDX cells when transplanted
under parallel and competitive settings (**Figure 3B** and **Supplementary Figures 2B,**
2D-E), and resulted in enhanced leukemia spreading and shorter survival of the
recipient mice (**Figure 3C** and **Supplementary Figure 2C**). This phenotype was
retained in secondary transplants, both in parallel and competitive settings (**Figures**
3D-F and **Supplementary Figures 2F-H**).
In vitro long term culture-initiating cell assays and proliferation experiments on the
TP53-silenced Dx PDX cells confirmed the relapse phenotype, indicating a *TP53*-
dependent increased clonogenic activity (**Figure 3G**, **Supplementary Table 3** and
Supplementary Figure 2I). Finally, *in vivo TP53*-silenced Dx cells also displayed
increased homing and post homing capacities (**Supplementary Figures 2J-M**).
Altogether, these results demonstrate that *TP53* silencing suffices to confer functional
features of the relapse phenotype to patient diagnosis cells.
181

182 ***TP53*** **alterations deregulate MYC signaling and metabolism pathways in T-ALL**

183 **cells and directly contribute to the** ***relapse phenotype***

184 To dig into the biological pathways deregulated by $TP53^{alt}$ in T-ALL relapse, we

185 compared transcriptional profiles of isogenic Dx and Rel T-ALL PDX samples. While

186 91 genes were upregulated at relapse, 181 genes were downregulated including *TP53*

187 itself (except in PDX11R where *TP53* was point-mutated on both alleles) (**Figure 4A**

188 and **Supplementary Figures 3A-B**). Geneset enrichment analysis (GSEA) identified

189 the MYC, E2F, and G2M checkpoint pathways as enriched in Rel cells along with

190 several cell metabolism pathways, including oxidative phosphorylation (OXPHOS),

191 glycolysis and cholesterol homeostasis, defining a relapse expression signature

192 (**Figure 4B**). Moreover, we interrogated genesets linked to cell proliferation,

193 invasiveness and stemness, and evidenced their upregulation at relapse

194 (**Supplementary Figure 3C**). Analysis in secondary transplant PDXs showed

195 consistent expression patterns, demonstrating functional stability (**Supplementary**

196 **Figures 3D-G**). The relapse expression signature was also retrieved from the

197 corresponding primary T-ALL samples, further outlining that our Dx-Rel PDX models

198 faithfully captured the features of the patient cells (**Figure 4C**).

199 Considering the role of TP53 in regulating MYC signaling and metabolism

200 pathways[33,34], we focused on OXPHOS and glycolysis and performed real-time

201 measurements of oxygen consumption rate (OCR) and extra cellular acidification rate

202 (ECAR) in isogenic Dx-Rel PDX pairs. We found increased basal and maximal

203 mitochondrial respiration, along with increased glycolysis, in the Rel vs Dx T-ALL cells

204 (**Figures 4D-E** and **Supplementary Figures 3H-K**). Consistently, experimental

205 silencing of *TP53* in the diagnosis PDX cells conferred a relapse mitochondrial and

206 glycolytic profile, demonstrating a driver role of *TP53* inactivation into deregulation of

207 metabolic functions (**Figures 4F-G** and **Supplementary Figures 3L-N**).

208

209 **Backtracking pre-relapse** ***TP53*^{alt} **clones at diagnosis**

210 As T-ALL relapse may emerge from minor pre-existing subclones at

211 diagnosis[13,17,19,22,35,36], we next aimed to backtrack rare *TP53*-mutated cells that would

212 constitute a genetically defined pre-relapse reservoir. We performed ultra-sensitive,

213 quantitative, droplet digital PCR (ddPCR) approach on the isogenic Dx-Rel primary

214 and PDX samples. We retrieved the expected *TP53*-associated relapse mutation in

215 the primary and/or PDX sample from diagnosis in 3 out of 4 cases (all but TALL11),

with very low variant allele frequencies (VAF) ranging from 0.01 to 0.04% that could be extrapolated to approximately 1/1250 to 1/5000 total cells, i.e., below the usual medical diagnosis detection thresholds of dominant genetic subclones). These results indicate the presence at diagnosis of a very minor *TP53*-mutant clone that was further selected and expanded in the post-treatment relapse sample. Interestingly, a cryopreserved remission sample was also available in case TALL11, in which we detected the expected *TP53*-R248Q mutation with a low frequency (VAF, 0.02%; approximately 1/2500 total cells), demonstrating the existence of the mutant cells reservoir during treatment. When this remission sample was injected into NSG mice, a full-blown leukemia developed from these rare cells. All the resulting leukemic cells carried the *TP53*-R248Q missense mutation. Interestingly, this leukemia retained a wildtype allele (*TP53*-R248Q allele frequency, 46% in sorted leukemic cells), contrasting with the relapse cells that demonstrated copy-neutral loss of the wildtype allele (*TP53*-R248Q allele frequency, 92%). This highlights the selective advantage upon treatment conferred by the monoallelic point mutation to a minor clone at remission, which further expanded in the PDX. The relapse cells displayed a complete loss of the wildtype *TP53* allele and function with duplication of the *TP53*-R248Q allele through acquisition of an uniparental disomy.

Those data demonstrate that relapse emerged from a pre-existing ultra-minor *TP53*-mutant clone that expanded during treatment through selective growth advantage, and subsequent acquisition of bi-allelic *TP53* loss at full relapse.

Combined single-cell profiling uncovers pre-relapse, reservoir cells at diagnosis that express features of the relapse phenotype

To further evaluate whether the rare *TP53*-mutant population from diagnosis expressed a relapse-like transcriptional state, we performed single-cell RNAseq analysis on the 4 T-ALL pairs. The 16 resulting samples (8 primary and 8 from PDXs) did not globally cluster by their diagnosis or relapse nature (i.e., all diagnosis *versus* all relapse samples), nor by their origin (i.e., all primary *versus* all PDX samples), or by cell cycle repartition, but rather by individual T-ALL case (**Figure 5A** and **Supplementary Figures 4A-B**), underlying prominent inter T-ALL biological heterogeneity. GSEA confirmed oxidative phosphorylation signature as the most differentially expressed pathway in all relapse samples along with cell proliferation, cell

signaling and cell metabolism pathways (**Supplementary Figures 4C-F** and **Figures 5B-C**).

Comparison of the diagnosis and relapse samples uncovered a minor cell population in two diagnosis samples (TALL42 and TALL30) that clustered nearby their paired relapse counterparts, suggesting the existence of a "pre-relapse" cells transcriptional state at diagnosis (**Supplementary Figures 4C, 4E**). We further integrated the combined analyses of mutation, copy number and TCR rearrangement to reconstitute leukemia architecture before and after treatment along with single-cell expression profiling in these two cases (**Figure 5D**).

In case TALL42, the small cluster of cells represented approximately 10% of the total leukemic cells at diagnosis (9.6% and 15% of the primary and PDX samples, respectively) (**Figures 5E-F** and **Supplementary Figures 4G-Q**). This minor cell population displayed both a distinct TCRB rearrangement (TCRB#2) and a distinct second-allele 9p/*CDKN2A* deletion from the dominant clone at diagnosis, and no 17p deletion, while the relapse population displayed the same subclonal TCRB#2 but yet another 9p/*CDKN2A* deletion (specific to relapse) and both the *TP53*-R175H and del17p/*TP53* (those last being not detected by NGS and inference analysis in the 10% subclone at diagnosis) (**Figure 5D** and **Supplementary Figures 4I, 4K, 4P, 4R**). Interestingly, the minor subpopulation at diagnosis was enriched in most of the relapse expression signature including OXPHOS, MTORC1 signalling, MYC targets V1, E2F targets, cholesterol homeostasis and DNA repair, these pathways being even more upregulated in the matched relapse cells (**Figure 5G** and **Supplementary Figure 4S**). We interrogated additional geneset collections and found a significant enrichment of stemness, invasiveness and OXPHOS signatures combined with downregulation of cell proliferation pathways, in line with the properties attributed to a quiescent, LIC-activity enriched, pre-relapse leukemia reservoir (**Figure 5H** and **Supplementary Figure 4T**). Specifically, interrogation of several leukemia stem cell (LSC) signatures[37–40] showed that this cluster had increased LSC signature scores compared to diagnosis cells, as in relapse cells (**Figure 5I**).

In the second case, TALL30, a minor cell population representing approximately 5% at diagnosis, trackable by the same TCRB rearrangement as relapse cells but distinct from the dominant diagnostic population, also demonstrated the deregulation of the relapse signature (**Supplementary Figure 5**).

Clearly, the very low frequency of the *TP53* mutation in the two cases (VAF <0.02%
as found by ddPCR) contrasted with the 5-10% frequency of the minor cell population
expressing the relapse signature. In case TALL42, the clonal architecture showed that
both the 5-10% minor clone population and the *TP53^alt* mutation derived from a
common ancestor bearing the same TCRB rearrangement from which they diverged
by acquiring a distinct 9p/*CDKN2A* second-allele deletion in addition to the *TP53*
mutation in the relapse clone (**Figure 5D**). In case TALL30, the *TP53*-mutated cells
may have emerged from the minor cell population or from an ancestor pre-relapse
reservoir. Although somewhat counterintuitive, these data suggest that relapse may
be primed in a pre-existing reservoir population expressing the relapse expression
signature, the *TP53* mutation occurring in a fraction of those cells that further expand
upon treatment. To investigate this hypothesis, we performed deep-sequencing, long-
read Nanopore scRNAseq on the 10X cDNAs to track the *TP53* mutations at the single-
cell resolution, allowing to directly assess the transcriptional profile of the very rare
TP53-mutant cells. In TALL42, we obtained a final mean depth of 110X at diagnosis
and 40X at relapse, in 8.861 cells and 6.022 cells respectively. The expected *TP53*-
R175H missense mutation was retrieved in all (35/35) covered reads from the del17p
relapse sample, but also in rare (6/193) reads from the diagnosis sample (**Figures 5J-
L** and **Supplementary Figures 4U-V**). While not directly numerically comparable to
ddPCR since they operate at different biological and technical levels (allele level
quantification versus transcriptional profiling), these data confirmed the enrichment of
the relapse expression signature in the rare *TP53*-mutated cells as compared to the
vast majority of *TP53 WT* cells from diagnosis (**Figure 5M**). Similar results were
obtained in the second TALL30 case in which the relapse expression signature was
also enriched in the rare *TP53*-mutant cells detected at diagnosis (**Supplementary
Figures 5P-U**).

Collectively, our data show that *TP53*-mutated cells pre-existing at diagnosis
constituted a pre-relapse reservoir population expressing the relapse signature, that
expanded upon treatment along with sequential inactivation of the second *TP53* allele.

311
312

313　**DISCUSSION**

314　Treatment failure and subsequent patient death for ALL are largely related to the
315　emergence, within the first 2-years, of a relapse much less responsive to therapies

than the initial diagnosis leukemia. Comprehensive longitudinal genomic studies on patient cohorts showed that rather than being subtype dependent, resistance initiates upon treatment by the selection of residual, ancestor T-ALL cells surviving therapy and re-expanding[13–19]. However, no clear genomic driver of relapse were detected in most cases and the underlying mechanisms, possibly non-genetic, remain elusive. Here, we leveraged *TP53*[alt] T-ALL relapse cases, i.e. leukemia with a clear, genetically-trackable driver of resistance to develop faithfull *in vivo* models, based on isogenic diagnosis-relapse paired xenografts, and analysed cellular features driving cell survival and re-expansion. Contrasting with many other T-ALL cases where we previously shown that the PDX resulting from diagnosis more closely resembled relapse samples than bulk diagnosis samples[13], diagnosis and relapse features were faithfully captured in these isogenic PDX pairs resulting from *TP53*[alt] T-ALL. This substantial difference may be due to the *TP53* lesions nature confering an *in vivo* advantage upon treatment only, as seen in other premalignant conditions[18,41]. Importantly, these isogenic PDX models enable to overcome fundamental limitations of *ex vivo* functional studies on primary samples or cell lines, i.e. rarity and practical workability for primary samples; highly rearranged and drifted cells for cell lines; and, more generally, a lack of microenvironnement support. In addition, isogenic PDXs also allow to avoid bias related to inter-patient heterogeneity, such as the markedly distinct proliferation range between our four T-ALL cases, faithfully captured by the isogenic PDX pairs (**Supplementary Figure 1B**).

Interestingly, we found that the *TP53*[alt]-driven T-ALL relapse cells did not display strong genomic instability, suggesting a direct oncogenic role of *TP53* alterations in those leukemias as "gatekeeper" rather than its commonly recognized "caretaker" role. This was confirmed by *TP53* silencing experiments evidencing a cell-intrinsic gain of malignancy in relapse cells largely driven by the *TP53* inactivation itself. The relapse phenotype we identified includes a cell-intrinsic functional gain of LIC and proliferation, along with an expression profile that involved cell metabolism, signaling, MYC pathways, and stemness, consistently with a role of *TP53* in the regulation of normal and pathologic stem cell function in hematopoiesis[42]. We especially evidenced functionally that increased *TP53*[alt]-dependent OXPHOS was crucial to the relapse phenotype, as may be seen in other acute leukemia subtypes such as acute myeloid leukemia or various solid cancers[43,44].

By combining single-cell transcriptomics, genomics and ultra-sensitive specific
backtracking by ddPCR at diagnosis, we could reconstitute the clonal architecture at
diagnosis and relapse in two cases (**Figure 5D**). We detected the expected *TP53*
mutations at diagnosis at ultra-low frequencies, showing that the relapse originated
from a pre-existing ultra-rare reservoir. Interestingly, a remission sample from one case
could be injected and expanded towards an overt T-ALL PDX that was *TP53-*
mutatedbut not yet 17p deleted, demonstrating the sequential bi-allelic acquisition
through leukemia progression. Intriguingly, a transcriptional cluster representing a cell
population of approximately 5-10% of cells was detected in two cases at diagnosis,
although the vast majority of those cells were *TP53*^WT. In one of these two cases
(TALL42), clonal architecture demonstrated that a rare *TP53*-mutated clone and the
minor (5-10%) cell population issued from the same ancestor, distinct from the
dominant clone at diagnosis. In the second case, the *TP53*-mutation may have
emerged directly from the minor cell population expressing the relapse signature or
from an ancestor. These data raise the possibility that the *TP53* mutation would
stabilize a pre-existing transcriptional state, possibly non genetically-driven. Taken the
similarity of the relapse signature and the effects of *TP53* inactivation, this minor
population may exhibit transcriptional features associated with the relapse because of
a functional, rather than genetic, inactivation of TP53 signaling. Beyond the *TP53-*
associated relapses we explored in this study, many T-ALL relapses do not display
any additional "relapse-associated genes"[17], suggesting that leukemia resistance and
relapse might be driven by non-genetic mechanisms. In line with this observation,
several studies have raised a spectrum of key epigenetic, transcriptionnal remodeling,
cell plasticity and microenvironmental mechanisms that may contribute to
relapse[17,35,37,40,48,49]. It will thus be of interest to study whether the *TP53*^alt-driven
relapse phenotype we identified here can also be relevant in T-ALL cases lacking *TP53*
genetic alterations at relapse, suggesting non-genetic driver mechanisms.
Like in the vast majority of cancers, *TP53*^alt T-ALL relapses are associated with very
poor clinical outcomes and limited therapeutic options, representing a major clinical
challenge. Detecting very rare pre-existing *TP53*-mutant cells at diagnosis, i.e., below
usual thresholds, potentially combined with single-cell transcriptomic detecting high-
risk cellular state, and *TP53*^alt bi-allelic deletion tracking, may be of broad interest for
patient monitoring and early intervention with targeted therapies or hematopoietic stem
cell transplantation.

METHODS

Patient cohort, collection of biological material and annotations. Primary samples from 42 T-ALL patients who have relapsed were obtained from blood or bone marrow (BM) aspirates/biopsies at initial diagnosis, remission (non tumoral, considered as germline reference) and first relapse. The cohort included children and young adults (age, 1 to 42 years old). Samples were stored in the Tumor Bank of Saint-Louis Hospital, Paris (BiRTH, Biobank for Research in Translational Hematology) accordingly to the French law, including informed consent. Annotations of T-ALL mutations were obtained using the NGS sequencing and analysis pipeline from Saint-Louis Hospital, using a VAF threshold of 2%[50]. Seven of the 42 patients (16.7%) had *TP53* alterations at relapse, including 4 for which live cells from both diagnosis and relapse could be recovered and injected to mice. IRB approval from the Inserm was given to this study under the number #12–078.

Animal studies. Mice experiments described in this study comply with the current ethical regulations. Authorizations for animal experimentation were obtained from the *Ministère de l'enseignement supérieur et de la recherche* with the No. APAFIS#19958-20190326150746255v2 and APAFIS#47448-2024020909144467v3. Mice were housed under a dark-light cycle at ambient temperature ($\approx$20-24°C) with a humidity of 40-60% and handled in a pathogen-free animal facility in accordance with the guidelines of the Animal Care and Use Committee (IRSL, Université Paris Cité). Animals were excluded from experiments if any sign of stress were observed without clinical signs of full-blown leukemia development (i.e. absence of human leukemic cells in blood, bone marrow and spleen).

Patient derived xenografts (PDXs) of isogenic diagnosis-relapse T-ALL samples. Isogenic diagnosis (Dx) and relapse (Rel) T-ALL patient samples were thawed, qualified and injected intravenously (unless otherwise stated) at equivalent cell doses into sublethally irradiated immunodeficient NOD.Cg-Prkdc$_{scid}$ IL2rg$_{tm1Wjl}$/SzJ (NSG)

mice from the Jackson Laboratory that are bred locally, as described previously[50].
Leukemia development was monitored monthly by flow cytometry (FACS) analysis of
blood or BM aspirates using hCD45 and hCD7-based staining protocol detailed in the
'Flow cytometry' section to confirm disease engraftment.

Human T-ALL cells from both patients and derived PDXs (BM of fully leukemic mice
with >95% blasts) were used for genomics, molecular and functional experiments. To
evaluate the stability of the phenotype from PDX models, leukemic cells from PDXs at
first passage were serially transplanted into NSG recipient mice by intravenous
injection (unless otherwise stated) at equivalent cell doses. In some circumstances to
evaluate post homing involvement, leukemic cells from isogenic Dx-Rel PDXs were
injected into sublethally irradiated mice by intrafemoral route. Recipient mice were
monitored daily by visual and clinical observations and monthly for biological
parameters, i.e. signs of stress (weakness, weight loss, ruffled coat, lethargy or ataxia)
and the presence of human leukemic cells in blood and/or BM. To monitor survival,
mice were sacrificed at or before reaching ethical endpoints. Pathology and flow
cytometry analyses confirmed full-blown T-ALL disease with marrow, spleen and liver
infiltration by leukemic cells.

Flow cytometry. Leukemic cells obtained from peripheral blood after Ficoll separation
and from BM aspirates were washed and resuspended in PBS (*Gibco*) supplemented
with 2% FBS (*Gibco*) before staining. Human leukemic cells infiltration in NSG recipient
mice was measured in legs, spleen and liver that were crushed or mashed. Leukemic
cells suspensions were washed and resuspended in PBS supplemented with 2% FBS
for staining. Human leukemic cells were then stained for 30 min at 4°C with anti-human
CD45 (APC or VioBlue) and CD7 (PE or PE-Vio770) antibodies (*Miltenyi Biotec*) (1:100
dilution). Cells were then washed and analysed on a FACS Canto II or LSR Fortessa
analyzer (*BD Biosciences*). Flow cytometry analyses were performed using Flowjo
10.10 (*TreeStar*).

Whole Exome Sequencing Analysis of primary and PDX T-ALL samples. Primary
samples from matched tumoral (initial diagnosis and first relapse) and non-tumoral
(remission) obtained from T-ALL patients and the corresponding isogenic Dx-Rel PDXs
were subjected to deep whole-exome sequencing (WES). DNA libraries were prepared
using SureSelect XT Target Enrichment according to the manufacturer's instructions

(*Agilent Technologies*, Santa Clara, CA, USA). Exon capture was performed with Agilent SureSelect Human All Exon V5 kit. The samples were sequenced on an Illumina NextSeq 500 platform (*Illumina*, San Diego, CA, USA) in a 2 × 100 bp paired-end sequencing run. The average sequencing depth was 110X. Raw data were first submitted to quality check using Trim_Galore (v0.6.6) to perform quality and adapter trimming for paired-end reads. The paired end clean reads were aligned to the human reference genome (hg19) using BWA (v0.7.17-r1188). Duplicates reads were flagged using Picard-tools (v2.3.0) and local realignment around InDels and base quality score recalibration was performed using Genome Analysis ToolKit (GATK v4.2.5.0).

Somatic mutations were called by VarScan somatic (v2.4.2) and annotated by ANNOVAR (version 2022-04-13) before filtering. Somatic copy number variants (CNVs) were detected by facets (v0.16.0-1) and by an in-house R script. The final retained abnormalities were validated by visual inspection considering a number of markers ≥2 and log2ratio with respect to the individual background noise.

Exon capture was performed with Agilent SureSelect Human All Exon V6 UTR kit for the isogenic Dx-Rel PDX pairs. The same procedure as above was used to analyse PDX samples (n= 8) using the corresponding primary non-tumour sample as reference for somatic calling variant step. The overlap between V5 and V6_UTR capture was performed for the upstream analysis.

***In vivo* limiting dilution experiments.** Leukemic cells from paired Dx-Rel PDXs (primografts) were intravenously injected under limiting dilution conditions from 10^6 to 1 cell per sublethally irradiated mouse (up to 6 mice per condition). Leukemia development in recipient NSG mice was monitored monthly by blood or BM aspirates. All mice were sacrificed at or before reaching ethical endpoints. Blast infiltration was confirmed as detailed above. Pathology analysis confirmed full-blown T-ALL diagnosis, with BM, spleen and liver infiltration by lymphoblasts expressing human CD45 and CD7 markers. Leukemia initiating cell (LIC) frequencies of isogenic Dx-Rel T-ALL PDXs were calculated from the number of engrafted mice for each cell dose in each condition using the WEHI bioinformatic ELDA (Extreme Limiting Dilution Analysis) software[51]. *P* values were calculated with overall tests for differences in LIC frequencies between any of the group tested.

In vitro long term culture-initiating cell **experiment.** Leukemic cells from isogenic Dx-Rel T-ALL PDX samples (primografts) and from paired control mCherry[+] and *TP53* KD GFP[+] PDX samples at diagnosis were sorted by flow cytometry based on CD45[+]CD7[+] (+/- GFP[+] or mCherry[+]) and seeded on irradiated (30 Gy) confluent stromal cells under limiting dilution conditions from 78.125 to 1 cell per well (24 wells per condition). Leukemic cells were then cocultured on stromal feeder cells for 2 to 4 weeks under hypoxic condition (3% O_2) as previously described[50]. Cell medium was partially changed twice a week. Positive wells for leukemic cells outgrowth were visually scored at regular time intervals. Leukemic cells detection was confirmed by flow cytometry analysis of recovered wells after 2 to 4 weeks of culture upon hypoxia as detailed above to confirm the presence of leukemic cells. LIC frequencies were calculated from the number of positive wells for leukemic cells outgrowth for each cell dose in each condition using the WEHI bioinformatic ELDA software[51]. *P* values were calculated with overall tests for differences in LIC frequencies between any of the group tested.

Real-time measurement of leukemic cells proliferation and apoptosis with the IncuCyte® Live-Cell Analysis System. Real-time measurement of *in vitro* leukemic cell proliferation and apoptosis was performed using the IncuCyte® Live-Cell Analysis System (*Sartorius*). Human CD45[+]CD7[+] leukemic cells from isogenic Dx-Rel T-ALL PDX samples (primografts) were sorted by flow cytometry and 100.000 cells were seeded on 96-well flat bottom plate (4 wells per condition) previously coated with 0.4μg of purified human Delta4-Fc fusion protein (*PX'Therapeutics*) according to manufacturer's recommendation. Leukemic cells were cultured as previously described[50] for up to 96 hours under normoxia in the IncuCyte® system (5% CO_2 and 37 °C). Real-time *in vitro* leukemic cell growth was monitored using live-cell time-lapse imaging by capturing phase contrast images every 2 hours and analysed using the integrated confluence algorithm provided. Cell proliferation was represented as the proportion of phase objects confluence over time, without labels. Leukemic cell apoptosis was detected using the IncuCyte® Caspase-3/7 green reagent for apoptosis (*Sartorius*). Briefly, leukemic cells were cultured as previously described in the presence of 5μM Caspase-3/7 green reagent. Real-time live-cell imaging of apoptosis was monitored by capturing phase contrast and green fluorescence images every 2 hours and analysed using the integrated algorithm provided. Leukemic cell apoptosis

was represented as the proprotion of green fluorescently labelled nuclei after activated caspase-3/7 cleavage of inert Incucyte® Caspase-3/7 dye.

Apoptosis and cell proliferation end-point assays. Leukemic cells from isogenic Dx-Rel PDX T-ALL samples were co-cultured on stromal cells[50] for 48 hours under normoxia in a humidified 5% CO_2 incubator at 37°C. To perform apoptosis assay, leukemic cells from co-cultures were then collected, washed and stained as detailed above. Leukemic cells were then fixed and stained with PE or Biotin anti-Cleaved Caspase 3 antibody (*BD Biosciences*) as recommended by the manufacturer's recommendations, washed and then analyzed by flow cytometry on FACS Canto II or a LSR Fortessa analyzers (*BD Biosciences*). Cell cycle assays were performed using the APC BrdU Flow kit (*BD Pharmigen*) as recommended by the manufacturer's recommendations. Briefly, leukemic cells co-cultured on stromal cells for 48 hours were pulse-labeled for one hour at 37°C with 1mM Bromodeoxyuridine (BrdU). Leukemic cells from the recovered co-cultures were stained with cell surface antibodies as above. Leukemic cells were then fixed, permeabilized, incubated with DNase and stained with APC-conjugated anti-BrdU antibody (1:100 dilution) for 20 min at room temperature and incubated with Vybrant violet (1:1000 dilution, *Invitrogen*) for 30 min at 37°C before FACS analyses.

***In vivo* homing assay.** Leukemic cells from isogenic Dx-Rel PDXs or from matched *TP53*KD and controls Dx PDXs were injected intravenously ($2\text{-}6\times10^6$ cells/mouse) into sublethally irradiated immunodeficient NSG recipient mice (n = 2 to 4 mice per group). Homing capacities of human leukemic cells were evaluated 24 hours after intravenous injection by FACS analysis of BM, spleen and liver.

***TP53* gene silencing in human leukemic cells and *in vivo* transplantations.** The Hu ShRNA anti-*TP53* was previously described[52]. The human *TP53* cDNA was cloned under the H1 promoter in the backbone pRRL.SIN.cPPT.PGK.GFP lentiviral vector (*Addgene* plasmid #22038). The pRRL_H1shCTL_PGK_mCherry lentivector was constructed in house from the pRRL_H1shLacZ_PGK_eGFP mentioned above. Lentiviral particles were produced as described[53] with jetPRIME reagents (*Polyplus transfection*) by co-transfection of viral production plasmids, according to the manufacturer's protocol. Lentiviral supernatants titration was performed using the

qPCR Lentivirus Titer Kit (*Applied Biological Materials*) according to the manufacturer's

recommendation.

Leukemic cells from primary PDX samples at diagnosis were cultured on 24-wells

plates previously coated with 2.5µg of purified human Delta4-Fc fusion protein

(*PX'Therapeutics*) and transduced for 48-72 hours with the lentiviral supernatants

including controls (at MOI 20-50) in the presence of 4ug/ml Protamine sulfate (*Sigma*).

Transduced leukemic cells were then washed , GFP- and mCherry-FACS sorted and

immediately intravenously injected into sub-lethally irradiated NSG mice, at equivalent

dose in parallel settings. For competitive experiments, leukemic cells transduced with

a shRNA CTL mCherry or shRNA *TP53*-GFP lentivirus were FACS-sorted, and mixed

in equivalent number before intravenous injection into sub-lethally irradiated NSG

mice. The starting GFP$^+$/mCherry$^+$ ratio was re-checked by flow cytometry before IV-

injection. Gene silencing efficiency was checked by immunoblot (human p53 D0-1,

Santa Cruz; Actin,*Santa Cruz*).

Large-scale gene expression analysis of primary and PDX T-ALL samples by

bulk RNA sequencing analysis. Total RNAs were extracted from frozen pellets of

leukemic blasts at initial diagnosis and first relapse (n= 8 samples) and from BM of fully

leukemic NSG mice for the corresponding derived PDXs (n= 24 PDX samples, first

injection) using Qiagen RNA extraction kit (*Qiagen,* miRNeasy Tissue cell advance)

according to the manufacturer's recommendations. cDNA libraries were prepared

using the TruSeq Stranded mRNA kit (*Illumina*, San Diego, CA, USA). Alignment to

the hg19 (GRCh37) reference genome was performed using STAR aligner (Version

2.7.8a) with quantMode GeneCounts parameter to build raw count matrix for gene-

level summarisation based on the in-built hg19 RefSeq-based annotation.

Normalisation by the method of trimmed mean of M-values (TMM) was performed

using the calcNormFactors function in edgeR. A linear model fitted in limma package

was performed to identify differential gene expression between relapse and diagnosis

samples in basal conditions. Differentially expressed genes were identified based on

the adjusted *p*-value < 0.05 and filtered with a fold change absolute value > 0.5. The

relationship between samples was visualized by hierarchical clustering and principal

component analysis (PCA). Over Representation Analysis (ORA) and Gene set

enrichment analyses (GSEA) were then performed to identify significant - hallmark,

positional (C1), curated (C2), and ontology (C5) - genes sets using the fgsea and

msigdbr packages. Additional analyses on PDX samples from secondary transplants were performed using the ClariomTM D Assay plateform (*Thermo Fisher Scientific*) using the manufacturer's recommendations.

Mitochondrial Respiration and Glycolytic Function Measurements. Oxygen consumption rate (OCR) and extracellular acidification rate (ECAR) were respectively measured with the Seahorse XFp Cell Mito Stress and XFp Glycolysis Stress Test kits on the Seahorse XFp Analyser (*Agilent, Seahorse Bioscience*) in accordance with the manufacturer's recommendations. Leukemic cells recovered from 24h co-culture on stromal cells were seeded at 4×10^5 cells per well (optimal cell seeding density was determined in preliminary experiments) in an XFp-well miniplate previously coated with Cell-Tak (*Corning, Fisher Scientific*) in 180µl of XF DMEM Medium (pH 7.4), supplemented with 2 mM L-glutamine (*Gibco*) (for OCR and ECAR), 1 mM fresh sodium pyruvate (*Gibco*) (for OCR only) and 10 mM D-glucose (*Sigma*) (for OCR only) on the day of the assay. For OCR measurements, sequential compound injections of Oligomycin (1µM), FCCP (1µM), Rotenone (0.5 µM) and Antimycin A (0.5µM) enabled measurements of basal respiration, ATP production, proton leak, maximal respiration, spare respiratory capacity, and non-mitochondrial respiration. For ECAR measurements, sequential compound injections of Glucose (10mM), Oligomycin (1µM) and 2-Deoxy-glucose (2-DG, 50mM) enabled measurements of glycolysis, glycolytic capacity, glycolytic reserve and non-glycolytic acidification. Key parameters of mitochondrial respiration and glycolytic functions were recorded for 90 minutes after cells were placed into the analyzer. OCR (pmol/min) and ECAR (mpH/min) were measured simultaneously in all wells, three times at each step, by optical fluorescent O2 sensors, and a minimum of three replicates were analyzed per condition in any given experiment. Unless otherwise stated, all compounds and material were obtained from *Agilent Seahorse Bioscience*.

Cell lines and human T-ALL cells culture. HEK-293T cell line was purchased from ATCC (CRL-3216). HEK-293T cells were grown in DMEM medium (*Gibco*) supplemented with 2mM L-glutamine (*Gibco*), 1% penicillin/streptomycin (*Gibco*), and 10% heat-inactivated fetal bovine serum (FBS) (*Gibco*). Mouse MS5-Delta-like1[13,54] and human hTERT-immortalized primary bone marrow MSCs[55] derived stromal cells were maintained in αMEM medium (*Gibco*) supplemented with 2mM L-glutamine

(*Gibco*), 1% penicillin/streptomycin (*Gibco*), 1μM Hydrocortisone (*Sigma*, for MSCs

only) and 10% heat-inactivated FBS (*Gibco*). All cell lines were grown in a humidified

5% CO_2 incubator at 37 °C and tested monthly negative for Mycoplasma using

MycoAlert PLUS Mycoplasma Detection Kit (*Lonza,*) or MycoStrip Mycoplasma

Detection Kit (*Invivogen*).

Primary and PDX human T-ALL leukemic cells were cocultured on stromal cells (MS5-

DL1 or hMSC cells) in αMEM medium (*Gibco*) supplemented with 2mM L-glutamine

(*Gibco*), 1% penicillin/streptomycin (*Gibco*), 10% heat-inactivated FBS (*Stem Cell*

Technologies), 10% heat-inactivated human AB serum (*Institut de Biotechnologies*

Jacques Boy), 20nM Human Recombinant Insuline (*SAFC Sigma*), 50ng/ml Human

Recombinant SCF (*Miltenyi Biotec*), 10ng/ml Human Recombinant IL7 (*Miltenyi*

Biotec), 20ng/ml Human Recombinant Flt3 ligand (*Miltenyi Biotec*) as already

described[13,50].

***TP53*-mutant cells backtracking in isogenic primary and PDX T-ALL samples by**

droplet-digital PCR.

Droplet-digital PCR (ddPCR) reactions were performed following Bio-Rad

recommendation's with 2X ddPCR SuperMix for probes No dUTP and commercially

designed ddPCR Mutation Assays for each *TP53* mutation (*Bio-Rad*). Mutation assays

are a mix of two fluorescent probes in competition for the annealing on the same target

sequence: one specific probe for the wild-type sequence (FAM probe) and a HEX

probe, specific of the mutated sequence. Two wet-lab validated ddPCR MUT FAMHEX

Assay were used for detection of patient's specific mutations (*TP53*-R175H (Assay ID

dHsaMDV2010105) and *TP53*-R248Q (Assay ID dHsaMDV2010127)) and a non-wet-

lab validated assay IS ddPCR MUT FAMHEX Assay was used for *TP53*-H214R

mutation detection (Assay ID dHsaMDS2510824). Experiments were performed on

genomic DNA from isogenic primary and PDX T-ALL samples. Several wells were

loaded with DNA from each sample to test enough material. Droplets were generated

using QX200 Droplet Generator (*Bio-Rad*). PCR reactions were performed on an

Eppendorf Nexus Gradient thermocycler following cycling conditions after validation of

optimum annealing/extension temperature: 95°C for 10 min, 40 cycles of two steps

PCR 94°C for 30 sec, 55°C for 1 min with a 2°C/sec ramping, an enzyme denaturation

step at 98°C for 10 min and a final hold at 4°C. FAM and HEX fluorescence's of each

droplet were then read on the QX200 Droplet reader. Visualization, analysis and VAFs

calculation were performed with QXOne Manager 1.2 Standard edition software (*Bio-*

Rad). Thresholds for each assay were determined aiming maximum specificity, despite

sensitivity and quantification accuracy by only incorporated double-positive droplets of

a well when at least a single positive-droplet was spotted to avoid potential false

double-negative droplets (none as "rain droplets") in the analysis. Non-template

controls containing all reagents and the corresponding amount of carrier DNA to be

equivalent to the tested samples were included in the analysis. VAFs were calculated

by considering all the wells analysed.

High-throughput short read single-cell RNA sequencing analysis of paired Dx-

Rel primary and PDX T-ALL samples. Single-cell RNA-sequencing (scRNAseq)

studies were performed on viable leukemic cells prepared from cryopreserved BM

samples from T-ALL patients and the corresponding derived PDXs. Libraries from

10.000 leukemic cells were prepared using the Chromium Next GEM Single Cell 3'

v3.1 reagent kits (*10x Genomics*) and subjected to short read sequencing on the

Illumina NovaSeq 6000 platform (*Illumina*, San Diego, CA, USA) in a 28-8-91 paired-

end sequencing run (50.000 reads per cell) allowing the captured expression of ~2000

genes per cell. FASTQ files were aligned to the human reference sequence (hg38)

using CellRanger count pipeline (*10× Genomics*, v3.1.0). The output gene-barcode

matrices were imported into the Seurat R package (v4.3.0) for scRNAseq data analysis

and visualization. Cells with nFeature_RNA <500 or >20% mitochondrial genes

content were filtered out. After removing of low-quality cells, data were normalized

using the Normalized-Data function with "LogNormalize" and a scale factor of 10,000.

Highly variable features were identified using the FindVariableFeatures function with

selection method "vst". Data were scaled using the ScaleData function and S.Score

and G2M.Score obtained from CellCycleScoring function were regressed out. Cells

were clustered using FindNeighbors (dims =1:50) with the FindClusters function

(resolution =0.6) based on the shared nearest neighbour. Datasets were visualized

through non-linear dimensional reduction using the Uniform Manifold Approximation

and Projection (UMAP). To identify a list of differentially expressed genes, gene

expression comparisons between clusters and/or samples were performed using the

fast Wilcoxon rank-sum test and area under the receiver operating characteristic curve

(auROC) analysis implemented in the R package Presto (v1.0.0). Gene signature

scores were computed using the AddModuleScore_UCell function from the UCell R

package (v2.6.2), which applies a rank-based scoring method for robust and scalable
analysis. To detect large-scale copy number variations in tumor cells, we utilized the
inferCNV package (v1.20.0)[56,57]. Gene expression profiles of all relapse cells, or of a
defined subset representing the pre-relapse subclone, were contrasted against
diagnostic cells to infer chromosomal variations in tumor cell clusters by analyzing
expression intensities across genomic locations and thereby explore potential
chromosomal alterations.
695
3' single-cell long read sequencing on isogenic Dx-Rel primary T-ALL samples.
Single-cell suspension was converted into barcoded single cell double strand cDNAs
using the Next GEM Single Cell 3' Kit (*10X Genomics*, V3.1), aiming for 10,000 cells
following the manufacturer's instructions. 10 ng of cDNA amplicons prepared using
Next GEM Single Cell 3'Kits (*10X Genomics*, V3.1) were tagged with biotin using PCR
amplification following the Oxford Nanopore Technologies (ONT) manufacturer's
instructions (SST_9198_v114_revK_29Jan2025) with custom-ordered oligos.
[Btn]Fwd_3580_partial_read1_defined_for_3'_cDNA at 10 µM:
5'/5Biosg/CAGCACTTGCCTGTCGCTCTATCTTCCTACACGACGCTCTTCCGATCT-
3'
Rev_PR2_partial_TSO_defined_for_3'_cDNA at 10 µM:
5'-CAGCTTTCTGTTGGTGCTGATATTGCAAGCAGTGGTATCAACGCAGAG-3' and
LongAmp Hot Start Taq 2X Master Mix (*NEB*).
Pull-down of the amplicons on M280 streptavidin beads (10 µg/µl, *Invitrogen*) was
performed before a second PCR using the PCR Primers (PRM) included in the PCR
Expansion kit (EXP-PCA001) and LongAmp Hot Start Taq 2X Master Mix (*NEB*). A
standard Ligation Sequencing Kit V14 library preparation (SQK-LSK114) was
completed to prepare the cDNA ends for sequencing on a PromethION device. 200
fmol of purified cDNA (Agencourt AMPure XP beads, *Beckman Coulter*™) from the
average fragment size identified were quantified using an Agilent Fragment Analyzer
and ligated. Only flow cells with at least 5,000 pores at the initial scan were used for
sequencing. Prepared library was loaded onto R10.4.1 flow cells (FLO-PRO114M,
ONT) on a PromethION P-48 device (*ONT*) (*Gustave Roussy Genomic Facility, Villejuif*
France) with MinKNOW (v6.0.14) and super high-accuracy base-calling mode.
720
3' single-cell long read ONT sequencing analysis.

Real-time base calling was run using MinKNOW (v7.4.14) with the Dorado basecaller
and the "Super-accurate basecalling v4.3.0" model. Alignments have been made using
the single-cell (v2.0.3) workflow from epi2me-labs (*https://github.com/epi2me-labs/wf-*
single-cell). After identification of the 10X adapters (from 3' V3.1 kit, *10X Genomics*)
and extraction of the cells barcodes and UMIs, the alignment on the human reference
(GRCh38-2024-A, *10X Genomics*) was performed using minimap2 (v2.24) as detailed
in the code, following parameters "-ax splice -uf --secondary=no". Genes and
transcripts assignment was performed using an in-house script from epi2me-labs
similar to FLAMES.

Cell barcode by symbol count table were loaded in R (version 4.1.0). To call real cells
from empty droplets, we used the emptyDrops() function from the dropletUtils package
(version 1.12.2), which assesses whether the RNA content associated with a cell
barcode is significantly distinct from the ambient background RNA present within each
sample. Barcodes with *p*-value < 0.001 (Benjamini-Hochberg-corrected) were
considered as legitimate cells for further analysis.

The count matrix was filtered to exclude genes detected in less than 5 cells, cells with
less than 1000 UMIs or less than 200 detected genes, as well as cells with
mitochondrial transcripts proportion >20%.The proportion of ribosomal gene counts
and the proportion of mechanical stress-response gene counts were also estimated
but not used to filter cells. Barcodes corresponding to doublet cells were identified and
discarded using the union of two methods: scDblFinder (version 1.6.0) and scds
(version 1.8.0) with its hybrid method with default parameters for both. We manually
verified that the cells identified as doublets did not systematically correspond to cells
in G2M phase.

Seurat (version 4.0.4) was applied for further data processing. The SCTransform
normalization method (Hafemeister & Satija Genome Biol. 2019) was used to
normalize, scale, select 3000 Highly Variable Genes and regress out bias factors
(nCount_RNA, the number of detected genes). Person residuals from this regression
were used for dimension reduction by Principal Component Analysis (PCA). The
number of PCA dimensions to keep for further analysis was evaluated by assessing a
range of reduced PCA spaces using 3 to 49 dimensions, with a step of 2. For each
generated PCA space, Louvain clustering of cells was performed using a range of
values for the resolution parameter from 0.1 to 1.2 with a step of 0.1. The optimal space
was manually evaluated as the one combination of kept dimensions and clustering

resolution resolving the best structure (clusters homogeneity and compacity) in a
UMAP space. Additionally, we used the clustree method (version 0.4.3) to assess if
the selected optimal space corresponded to a relatively stable position in the clustering
results tested for these dimensions / resolution combinations.
An automatic annotation of cell types was performed by SingleR (version 1.6.1) (with
fine-tuning step) and ClustifyR (version 1.5.1), using packages built-in references. It
labels clusters (or cells) from a dataset based on similarity (Spearman correlation
score) to a reference dataset with known labels. The labels with a correlation score
greater than 0.25 for SingleR or greater than 0.35 for ClustifyR were kept. The
annotation was made with CelliD (version 1.0.0) with genes signatures from panglao
database. Marker genes for Louvain clusters were identified through a «one
versus others» differential analysis using the Wilcoxon test through the
FindAllMarkers() function (Seurat), considering only genes with a minimum log fold-
change of 0.5 in at least 75% of cells from one of the groups compared, and FDR-
adjusted p-values <0.05 (Benjaminin-Hochberg method). SNP were called by using
bam-readcount on individual BAMs per barcodes.
772
Statistical analyses. Statistical analyses were performed with GraphPad Prism
(version 10.3; *GraphPad Software Inc.*) or R packages using Mann-Whitney, Anova,
Wilcoxon and Kruskal-Wallis tests, as indicated in figure legends. Unless otherwise
stated, the data are expressed as mean±s.d. of *n*=3 or more determinations.
For xenotransplantation analysis, Kaplan–Meier survival curves were performed and
the statistical significance was determined with the Log-rank (Mantel-Cox) test. Overall
survival was defined as the time from cell injection to euthanasia or death at predefined
endpoints. Animals were censored if the cause of death/sacrifice was not leukemia
related. Number of mice and median survival are indicated into brackets.
For limiting dilution assays, statistical analysis was performed using the WEHI
bioinformatic ELDA (Extreme Limiting Dilution Analysis) software[51]). Differences were
considered statistically significant at *p*<0.05 (*), *p*<0.01 (**), *p*<0.001 (***) or *p*<0.0001
(****).
786

Authors' Contributions

Conception and design of the study: SG, EC and JS

Development of methodology: SG, SQ, LH, RJ, MA and JS

Acquisition of data (provided materials and animals, acquired and managed patients, provided facilities): SG, SQ, LH, LM, MP, RK, HB, MB, VP, PF, AB, HD, NB, EC and JS

Analysis and interpretation of data (biostatistics, computational analysis, statistical analysis): SG, SQ, LH, LM, MB, RJ, MA, FS, EC and JS

Writing of the manuscript - original draft preparation, review and editing: SG, HdT, EC and JS

Grants acquisition and study supervision: SG and JS

All authors critically reviewed the manuscript.

Disclosure of potential conflicts of interest

The authors declare no potential conflicts of interest.

Acknowledgments

The authors thank Niclas Setterblad, Christelle Doliger, Antonio Alberdi, Sophie Duchez and Claire Maillard from the Technological core facility of the Saint-Louis Research Institut, Inserm UMR1342 / CNRS EMR8000; Nathalie Droin and Maëla Francillette from the Genomic core facility of Gustave Roussy; the ICGex NGS platform of the Curie Institut for high-throughput sequencing of *10X* cDNA librairies.

Fundings

This work was supported by grants to Jean Soulier from the ERC St Grant Consolidator #311660, the team FRM 2022 program from the Fondation pour la Recherche Médicale #00107939, the Ligue Contre le Cancer IDF, the Fondation Leucémie Espoir, and by grants to Stéphanie Gachet from the Fondation de France, the Fondation ARC pour la Recherche sur le Cancer (PJA 20191209702), and the Paris Saint-Louis Leukemia Institute program (a grant managed by the French National Research Agency under the France 2030 program, ANR-23-IAHU-0005, Institut de la Leucémie Paris Saint-Louis). The ICGex platform was supported by the grants from the French National Research Agency, the Canceropole Ile-de-France and by the SiRIC-Curie program.

Data-sharing statement

The data that support the findings of this study are publically available at EBI (https://www.ebi.ac.uk/biostudies) under the accession numbers BioStudies S-BSST2554 and S-BSST2808, or in the Supplementary materials. The custom code used for the analyses will be available on the published version of the article.

REFERENCES

1. Van Vlierberghe, P. & Ferrando, A. The molecular basis of T cell acute lymphoblastic leukemia. *J Clin Invest* **122**, 3398–406 (2012).

2. Buckley, M., Yeung, D. T., White, D. L. & Eadie, L. N. T-cell acute lymphoblastic leukaemia: subtype prevalence, clinical outcome, and emerging targeted treatments. *Leukemia* **39**, 1294–1310 (2025).

3. Weng, A. P. *et al.* Activating mutations of NOTCH1 in human T cell acute lymphoblastic leukemia. *Science* **306**, 269–71 (2004).

4. Ferrando, A. A. *et al.* Gene expression signatures define novel oncogenic pathways in T cell acute lymphoblastic leukemia. *Cancer Cell* **1**, 75–87 (2002).

5. Soulier, J. *et al.* HOXA genes are included in genetic and biologic networks defining human acute T-cell leukemia (T-ALL). *Blood* **106**, 274–86 (2005).

6. Girardi, T., Vicente, C., Cools, J. & De Keersmaecker, K. The genetics and molecular biology of T-ALL. *Blood* **129**, 1113–1123 (2017).

7. Pölönen, P. *et al.* The genomic basis of childhood T-lineage acute lymphoblastic leukaemia. *Nature* **632**, 1082–1091 (2024).

8. Desjonqueres, A. *et al.* Acute lymphoblastic leukemia relapsing after first-line pediatric-inspired therapy: a retrospective GRAALL study. *Blood Cancer J* **6**, e504 (2016).

9. Heikamp, E. B. & Pui, C. H. Next-Generation Evaluation and Treatment of Pediatric Acute Lymphoblastic Leukemia. *J Pediatr* **203**, 14-24 e2 (2018).

10. Pagliaro, L. *et al.* Acute lymphoblastic leukaemia. *Nat. Rev. Dis. Primer* **10**, 41 (2024).

11. Brady, S. W. *et al.* The genomic landscape of pediatric acute lymphoblastic leukemia. *Nat. Genet.* **54**, 1376–1389 (2022).

12. Pölönen, P., Mullighan, C. G. & Teachey, D. T. Classification and risk stratification in T-lineage acute lymphoblastic leukemia. *Blood* **145**, 1464–1474 (2025).

13. Clappier, E. *et al.* Clonal selection in xenografted human T cell acute lymphoblastic leukemia recapitulates gain of malignancy at relapse. *J Exp Med* **208**, 653–61 (2011).

14. Mullighan, C. G. *et al.* Genomic analysis of the clonal origins of relapsed acute lymphoblastic leukemia. *Science* **322**, 1377–80 (2008).

15. Tosello, V. *et al.* WT1 mutations in T-ALL. *Blood* **114**, 1038–1045 (2009).

16. Anderson, K. *et al.* Genetic variegation of clonal architecture and propagating cells in leukaemia. *Nature* **469**, 356–61 (2011).

17. Kunz, J. B. *et al.* Pediatric T-cell lymphoblastic leukemia evolves into relapse by clonal selection, acquisition of mutations and promoter hypomethylation. *Haematologica* **100**, 1442–50 (2015).

18. Wong, T. N. *et al.* Role of TP53 mutations in the origin and evolution of therapy-related acute myeloid leukaemia. *Nature* **518**, 552–555 (2015).

19. Oshima, K. *et al.* Mutational landscape, clonal evolution patterns, and role of RAS mutations in relapsed acute lymphoblastic leukemia. *Proc Natl Acad Sci U A* **113**, 11306–11311 (2016).

20. Meyer, J. A. *et al.* Relapse-specific mutations in NT5C2 in childhood acute lymphoblastic leukemia. *Nat. Genet.* **45**, 290–294 (2013).

21. Tzoneva, G. *et al.* Activating mutations in the NT5C2 nucleotidase gene drive chemotherapy resistance in relapsed ALL. *Nat Med* **19**, 368–71 (2013).

22. Tzoneva, G. *et al.* Clonal evolution mechanisms in NT5C2 mutant-relapsed acute lymphoblastic leukaemia. *Nature* **553**, 511–514 (2018).

23. Dieck, C. L. *et al.* Structure and Mechanisms of NT5C2 Mutations Driving Thiopurine Resistance in Relapsed Lymphoblastic Leukemia. *Cancer Cell* **34**, 136-147.e6 (2018).

24. Barz, M. J. *et al.* Subclonal NT5C2 mutations are associated with poor outcomes after relapse of pediatric acute lymphoblastic leukemia. *Blood* **135**, 921–933 (2020).

25. Diccianni, M. B. *et al.* Clinical significance of p53 mutations in relapsed T-cell acute lymphoblastic
leukemia. *Blood* **84**, 3105–12 (1994).

26. Hof, J. *et al.* Mutations and deletions of the TP53 gene predict nonresponse to treatment and poor
outcome in first relapse of childhood acute lymphoblastic leukemia. *J Clin Oncol* **29**, 3185–93
(2011).

27. Hof, J. *et al.* NOTCH1 mutation, TP53 alteration and myeloid antigen expression predict outcome
heterogeneity in children with first relapse of T-cell acute lymphoblastic leukemia. *Haematologica*
102, e249–e252 (2017).

28. Richter-Pechańska, P. *et al.* Identification of a genetically defined ultra-high-risk group in relapsed
pediatric T-lymphoblastic leukemia. *Blood Cancer J.* **7**, e523 (2017).

29. Simonin, M. *et al.* Prognostic value and oncogenic landscape of TP53 alterations in adult and
pediatric T-ALL. *Blood* **141**, 1353–1358 (2023).

30. Kempter, T. *et al.* Subclonal TP53 and KRAS variants combined with poor treatment response
identify ultrahigh-risk pediatric patients with T-ALL. *Blood Adv.* **9**, 1267–1279 (2025).

31. Li, B. *et al.* Therapy-induced mutations drive the genomic landscape of relapsed acute
lymphoblastic leukemia. *Blood* https://doi.org/10.1182/blood.2019002220 (2020)
doi:10.1182/blood.2019002220.

32. Armstrong, F. *et al.* NOTCH is a key regulator of human T-cell acute leukemia initiating cell activity.
Blood **113**, 1730–1740 (2009).

33. Liu, Y., Su, Z., Tavana, O. & Gu, W. Understanding the complexity of p53 in a new era of tumor
suppression. *Cancer Cell* **42**, 946–967 (2024).

34. Dong, Y., Tu, R., Liu, H. & Qing, G. Regulation of cancer cell metabolism: oncogenic MYC in the
driver's seat. *Signal Transduct. Target. Ther.* **5**, 124 (2020).

35. Richter-Pechańska, P. *et al.* Pediatric T-ALL type-1 and type-2 relapses develop along distinct
pathways of clonal evolution. *Leukemia* **36**, 1759–1768 (2022).

36. Albertí-Servera, L. *et al.* Single-cell DNA amplicon sequencing reveals clonal heterogeneity and
evolution in T-cell acute lymphoblastic leukemia. *Blood* **137**, 801–811 (2021).

37. Calvo, J. *et al.* High CD44 expression and enhanced E-selectin binding identified as biomarkers of
chemoresistant leukemic cells in human T-ALL. *Leukemia* **39**, 323–336 (2025).

38. Xu, J. *et al.* A multiomic atlas identifies a treatment-resistant, bone marrow progenitor-like cell
population in T cell acute lymphoblastic leukemia. *Nat. Cancer* **6**, 102–122 (2025).

39. Ebinger, S. *et al.* Characterization of Rare, Dormant, and Therapy-Resistant Cells in Acute
Lymphoblastic Leukemia. *Cancer Cell* **30**, 849–862 (2016).

40. Costea, J. *et al.* Role of stem-like cells in chemotherapy resistance and relapse in pediatric T-cell
acute lymphoblastic leukemia. *Nat. Commun.* **16**, 5413 (2025).

41. Hsu, J. I. *et al.* PPM1D Mutations Drive Clonal Hematopoiesis in Response to Cytotoxic
Chemotherapy. *Cell Stem Cell* **23**, 700-713.e6 (2018).

42. Chen, S. *et al.* Mutant p53 drives clonal hematopoiesis through modulating epigenetic pathway.
Nat. Commun. **10**, 5649 (2019).

43. Liu, Y., Su, Z., Tavana, O. & Gu, W. Understanding the complexity of p53 in a new era of tumor
suppression. *Cancer Cell* **42**, 946–967 (2024).

44. Chomczyk, M. *et al.* Impact of p53-associated acute myeloid leukemia hallmarks on metabolism
and the immune environment. *Front. Pharmacol.* **15**, 1409210 (2024).

45. Chiu, P. P. L., Jiang, H. & Dick, J. E. Leukemia-initiating cells in human T-lymphoblastic leukemia
exhibit glucocorticoid resistance. *Blood* **116**, 5268–5279 (2010).

46. Belmonte, M., Hoofd, C., Weng, A. P. & Giambra, V. Targeting leukemia stem cells: which pathways
drive self-renewal activity in T-cell acute lymphoblastic leukemia? *Curr Oncol* **23**, 34–41 (2016).

47. Gerby, B. *et al.* High-throughput screening in niche-based assay identifies compounds to target
preleukemic stem cells. *J. Clin. Invest.* **126**, 4569–4584 (2016).

48. Yadav, B. D. *et al.* Heterogeneity in mechanisms of emergent resistance in pediatric T-cell acute
lymphoblastic leukemia. *Oncotarget* **7**, 58728–42 (2016).

49. Ariës, I. M. *et al.* PRC2 loss induces chemoresistance by repressing apoptosis in T cell acute
lymphoblastic leukemia. *J. Exp. Med.* **215**, 3094–3114 (2018).

50. Gachet, S. *et al.* Deletion 6q Drives T-cell Leukemia Progression by Ribosome Modulation. *Cancer*
Discov **8**, 1614–1631 (2018).

51. Hu, Y. & Smyth, G. K. ELDA: extreme limiting dilution analysis for comparing depleted and enriched
populations in stem cell and other assays. *J. Immunol. Methods* **347**, 70–78 (2009).

52. Ceccaldi, R. *et al.* Bone marrow failure in Fanconi anemia is triggered by an exacerbated p53/p21
DNA damage response that impairs hematopoietic stem and progenitor cells. *Cell Stem Cell* **11**,
36–49 (2012).

53. Sebert, M. *et al.* Clonal hematopoiesis driven by chromosome 1q/MDM4 trisomy defines a canonical
route toward leukemia in Fanconi anemia. *Cell Stem Cell* **30**, 153-170.e9 (2023).

54. Gerby, B. *et al.* Optimized gene transfer into human primary leukemic T cell with NOD-
SCID/leukemia-initiating cell activity. *Leukemia* **24**, 646–9 (2010).

55. Mihara, K. *et al.* Development and functional characterization of human bone marrow mesenchymal
cells immortalized by enforced expression of telomerase. *Br. J. Haematol.* **120**, 846–849 (2003).

56. Patel, A. P. *et al.* Single-cell RNA-seq highlights intratumoral heterogeneity in primary glioblastoma.
Science **344**, 1396–1401 (2014).

57. Zhu, Q. *et al.* Single cell multi-omics reveal intra-cell-line heterogeneity across human cancer cell
lines. *Nat. Commun.* **14**, 8170 (2023).

945
946

FIGURE LEGENDS

Figure 1. *In vivo* **isogenic modelling of diagnosis and relapse** *TP53***-altered T-ALL. A.** Experimental design (Created in BioRender. Gachet, S. (2026) https://BioRender.com/zczr8er). Isogenic Dx-Rel PDX pairs were established after injection into immunodeficient NSG mice of paired primary diagnosis and *TP53*-altered (*TP53*^{alt}) relapse samples from the same T-ALL patients. **B.** Comparative Whole Exome Sequencing (WES) analysis of the isogenic Dx-Rel PDX samples established from 4 T-ALL patients who relapsed with *17p/TP53* alterations. In each case, PDX samples (PDXD and PDXR) are compared side-by-side to the corresponding primary samples (TALLD and TALLR, respectively). **C.** Lollipop plot of *TP53* mutations observed in 4 primary relapse samples (TALL11R, TALL30R, TALL42R and TALL43R).

Figure 2. *TP53*^{alt} **T-ALL relapse cells display a cell-intrinsic gain of malignancy phenotype. A.** Kaplan-Meier survival curves of PDX mice engrafted with primary isogenic Dx-Rel leukemic cells from 4 T-ALL patients (TALL11, TALL30, TALL42 and TALL43). N=4-5 recipient mice injected at equivalent cell doses per sample. **B-C.** Parallel serial transplantations of paired Dx-Rel T-ALL PDX samples. Kaplan-Meier survival curves of mice injected with 3 cell doses (10^6, 10^5 and 10^4 cells) of isogenic Dx-Rel PDX pairs (n = 4 TALL cases) raised from primary (**B**) and secondary (**C**) xenografts. **D.** LIC frequencies of isogenic Dx-Rel PDX sample pairs calculated from *in vivo* limiting dilutions. **E-F.** Real-time *in vitro* cell proliferation and apoptosis of leukemic cells from isogenic Dx-Rel PDX pairs, measured with the Incucyte® Live-Cell Analysis System. Proliferation was quantified by measuring the proportion of phase objects confluence over time, without labels **(E)**, and apoptosis by the quantification of green fluorescently labelled nuclei after activated caspase-3/7 cleavage of inert Incucyte® Caspase-3/7 dyes **(F)**.

Statistical data in this figure are represented as mean ± s.d. Significance is indicated by *P* values; A-C: Log-rank test (Mantel-Cox); D: overall tests for differences using ELDA software; E-F: two-way Anova test. Number of recipient mice and median survival are indicated into brackets on Kaplan-Meier survival curves. ns, not significant; *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$; ****, $p < 0.0001$.

Figure 3. *TP53* silencing confers a relapse phenotype to diagnosis leukemic

cells. A. Experimental design of *TP53* silencing in diagnosis cells from PDX samples

using a lentiviral shRNA approach (Created in BioRender. Gachet, S. (2026)

https://BioRender.com/usb6t9c). **B-C.** Primary transplantation of paired control

mCherry⁺ (shCTL mCherry) and *TP53* KD GFP⁺ (sh *TP53* GFP) leukemic cells from

PDX30D injected in parallel settings into recipient mice. **B.** Engraftment kinetics in the

bone marrow (BM) of recipient mice (n=6) injected with paired control and *TP53* KD

leukemic cells from PDX30D. **C.** Kaplan-Meier survival curves of mice injected with

paired control and *TP53* KD leukemic cells from PDX30D. **D-F.** Secondary transplant

of paired control mCherry⁺ (shCTL mCherry) and *TP53* KD GFP⁺ (sh *TP53* GFP)

leukemic cells from PDX30D injected either in parallel (panels D-E) or competitive

(panel F) settings into recipient mice. **D.** Engraftment kinetics in the BM of recipient

mice (n=5) injected in parallel settings with cells from control and *TP53* KD conditions.

E. Kaplan-Meier survival curves of mice injected in parallel settings with sorted control

or *TP53* KD leukemic cells from PDX30D. **F.** Engraftment kinetics in the BM of recipient

mice (n=4) injected with sorted cells from both control and *TP53* KD in competitive

settings (ratio 1:1). Data are represented as ratios of *TP53* KD GFP⁺ over control

mCherry⁺ leukemic cells within BM of the same recipient mouse and compared to the

input. **G.** LIC frequencies of paired control-*TP53* KD samples calculated from *in vitro*

long-term culture-initiating cell assays of sorted GFP⁺ or mCherry⁺ CD45⁺CD7⁺

leukemic cell from diagnosis PDX samples (PDX30D and PDX42D). LIC frequencies

are estimated based on the proportions of positive wells for GFP⁺/mCherry⁺

CD45⁺CD7⁺ leukemic cells evaluated after 2 and 4 weeks of culture.

Statistical data in this figure are represented as mean ± s.d. Significance is indicated

by *P* values; B and D: Mann-Whitney test; C and E: Log-rank test (Mantel-Cox); F:

Kruskal-Wallis test; G: overall tests for differences using ELDA software. Number of

recipient mice and median survival are indicated into brackets on Kaplan-Meier

survival curves. ns, not significant; *, $p < 0.05$; **, $p < 0.01$; ****, $p < 0.0001$.

1008

1009 **Figure 4. *TP53* alterations drive a relapse phenotype through MYC signaling and**

1010 **metabolism deregulation. A.** Supervised hierarchical clustering of the top fifty most

1011 differentially expressed genes between Dx and Rel T-ALL samples. Heatmap

1012 representation of the most variables genes at Dx and Rel in 4 isogenic PDX pairs

1013 (primografts). *TP53* expression is highlighted in red. Independent triplicates are

analysed for each isogenic PDX pair. **B-C.** GSEA of enriched pathways in *TP53*[alt] T-ALL relapse samples. **B.** Comparative Gene Set Enrichment Analysis (GSEA) of differentially expressed genes in Rel and Dx samples from 4 isogenic PDX pairs. Representation of selected Hallmark pathways significantly upregulated at relapse, defining a relapse expression signature in all isogenic Dx-Rel PDX pairs pooled together (1st column "All_PDXs") and in each individual PDX pair (2nd to 5th columns; respectively PDX11, PDX30, PDX42 and PDX43) (*P* adj <0.05). **C.** GSEA of the upregulated (upper panel) and downregulated (lower panel) relapse expression signature obtained from PDX samples (panel B) tested in the original T-ALL patients samples. UP_PDX, upregulated transcriptomic signature in Rel PDXs; DOWN_PDX, downregulated transcriptomic signature in Rel PDXs. FDR, false discovery rate; NES, normalized enrichment score. **D-G.** Direct evaluation of mitochondrial respiration in leukemic cells from isogenic Dx-Rel PDXs using Seahorse assays. **D.** Real-time measurements of oxygen consumption rate (OCR) in isogenic Dx-Rel leukemic cells from 4 T-ALL cases. **E.** Quantification of maximal respiration in leukemic cells from 4 isogenic Dx-Rel PDX pairs evaluated in 3 independent experiments. **F.** Real-time measurements of OCR in *TP53*-silenced (*TP53* KD) and matched control leukemic cells from diagnosis PDX samples (PDX42D and PDX43D). **G.** Quantification of maximal respiration in *TP53* KD and control leukemic cells from 3 independent diagnosis PDX samples (PDX30D, PDX42D and PDX43D) evaluated in 2 independent experiments

Statistical data in this figure are represented as mean $\pm$ s.d. Significance is indicated by *P* values; E and G: Mann-Whitney test. ns, not significant; **, $p < 0.01$; ***, $p < 0.001$; ****, $p < 0.0001$.

Figure 5. Single-cell profiling of isogenix Dx-Rel T-ALLs can unveil a pre-relapse cells reservoir at diagnosis. A. UMAP visualizations of isogenic Dx-Rel primary samples from 4 T-ALL patients and their derived PDXs analysed by short read single-cell RNA (scRNA) sequencing. UMAP plots of 117.119 single cells coloured by T-ALL samples (n=16, primary and PDXs samples from 4 T-ALL cases at diagnosis and relapse; left panel), leukemic cell identity (Diagnosis versus Relapse; middle panel) and by leukemic cell origin (Primary versus PDX; right panel). **B-C.** GSEA of biological pathways enriched at relapse in primary and PDX T-ALL samples. GSEA from pseudo bulk analysis of differentially expressed genes between relapse (Rel) and diagnosis

(Dx) samples from isogenic patient (**B**) and PDX (**C**) T-ALL pairs retrieves the relapse

expression signature defined in Figure 4B. Representation of selected Hallmark

pathways significantly deregulated at relapse (*P* adj <0.05) in all isogenic pairs pooled

together (1st column ") and in each individual pair (2nd to 5th columns). Note that TALL30

case eventually diverged from the other cases by the late acquisition of a *NT5C2*

mutation, known to drive resistance to maintenance therapy. **D.** Oncogenetic

trajectories for diagnosis and relapse TALL42 and TALL30 samples. **E.** UMAP

visualizations of Dx and Rel primary samples from TALL42 case analysed by short

read scRNA sequencing. UMAP plots of 14.049 single leukemic cells coloured by cell

identity (Diagnosis vs Relapse; upper panel) and cluster-based by transcriptomic data

(fourteen different clusters; lower panel). The red dotted ellipse highlights cluster 6 in

TALL42D sample. **F.** Dot plot visualization of biomarker genes average expression

selected from the unsupervised differential expression analysis between Rel and Dx

samples from TALL42 case. Average expression and proportion of expressing cells for

selected biomarkers are represented in each cluster from Dx and Rel samples. The

red dotted line highlights the pre-relapse cells reservoir. **G-H.** GSEA of positively and

negatively enriched pathways at relapse and pre-relapse stages in TALL42 samples.

G. Representation of selected Hallmark pathways significantly deregulated in relapse

(1st column Rel vs Dx; *P* adj <0.05) and pre-relapse (2nd column Sub vs Dx, red dotted

line; *P* adj <0.05) clusters as compared to matched diagnosis counterparts. **H.**

Representation of specific gene sets from the MSigDB C1 and C2 collections

significantly upregulated (in red) or downregulated (in blue) at relapse (1st column Rel

vs Dx; *P* adj <0.05) and pre-relapse (2nd column Sub vs Dx; *P* adj <0.05) as compared

to matched primary diagnosis counterparts. Color code indicates the cell proliferation,

invasiveness, stemness and oxidative phosphorylation signatures. **I** Single-cell scores

for several leukemia stem cell (LSC) signatures in TALL42 diagnosis (Dx), pre-relapse

(Sub) and relapse (Rel) clusters. The box includes the median, hinges mark the 25th

and 75th percentiles. *P* values from a two-sided paired t-test are shown. ns, not

significant; ****, *p* < 0.0001 **J.** UMAP visualizations of Dx and Rel primary samples

from TALL42 case analysed by long read single-cell RNA sequencing. UMAP plots of

14.883 single leukemic cells (8.861 cells from diagnosis and 6.022 cells from relapse)

coloured by cell identity (Diagnosis vs Relapse; left panel) and cluster-based by

transcriptomic data (9 different clusters corresponding to leukemic cells; right panel).

K. Projection of *TP53* WT and mutant (*TP53*-R175H*)* reads at chromosomal location

chr17:7'675'088 (hg38) from TALL42 Dx and Rel samples. **L.** Quantification of *TP53*

mutant reads retrived in isogenic Dx (6/193) and Rel (35/35) TALL42 samples. **M.**

Single-cell signature scores for relevant hallmark pathways of the relapse expression

signature (OXPHOS, MYC target V1 and MTORC1 signaling) in *TP53* WT and mutant

cells. The box includes the median, hinges mark the 25th and 75th percentiles.

1087

Figure 1

Figure 2

Figure 3

A

B

Primary injection
PDX30D

ns   ns   *   ns

% transduced leukemic cells / total leucocytes (BM)

Weeks post engraftment   4   8   12   16

■ sh CTL mCherry (n=6)
▲ sh *TP53* GFP (n=6)

C

Primary injection
PDX30D

Percent survival

Elapsed days   0   50   100   150   200

** *p* = 0.0061

- - sh CTL mCherry (n=6; 150.5 days)
- - sh *TP53* GFP (n=6; 128 days)

D

Secondary injection
PDX30D

**   ns   *

% transduced leukemic cells / total leucocytes (BM)

Weeks post engraftment   4   8   12

■ sh CTL mCherry (n=6)
▲ sh *TP53* GFP (n=6)

E

Secondary injection
PDX30D

Percent survival

Elapsed days   0   50   100   150   200

** *p* = 0.0018

- - sh CTL mCherry (n=5; 150 days)
- - sh *TP53* GFP (n=5; 102 days)

F

Secondary injection
PDX30D

ns   ns   *

Ratio GFP+ / mCherry+ leukemic cells (BM)

100
10
1
0.1

Weeks post engraftment   0   4   8   12

G

PDX30D   PDX42D

****   ****

LIC frequency (95% CI)

1
1/10
1/10^2
1/10^3
1/10^4

sh CTL mCherry
sh *TP53* GFP
sh CTL mCherry
sh *TP53* GFP

Figure 4

Figure 5