Phylogenetic analysis of enteroviruses from non-human primates reveals two new species

within the genus _Enterovirus_ and inter-species recombination

Corentin Aubé[1], Paulina Cruz de Casas[1], Matthieu Prot[2], Artem Baidaliuk[2], Marie-Claire

Endegue Zanga[3], Etienne Simon-Lorière[2], Nolwenn Jouvenet[1], Serge A. Sadeuh-Mba[3,4], Maël

Bessaud[1, 5]

[1] Institut Pasteur, Université de Paris Cité, CNRS UMR 3569, Virus sensing and signaling Unit.

25-28, rue du Dr Roux 75015 Paris. France

[2] Institut Pasteur, Université Paris Cité, CNRS UMR2000, Evolutionary Genomics of RNA

Viruses Unit. 25-28, rue du Dr Roux 75015 Paris. France

[3] Centre Pasteur du Cameroun, Virology Laboratory, _451, avenue 2005 - Yaoundé 2. Cameroon_

[4] Salisbury Animal Health Laboratory, Maryland Department of Agriculture, 27722 Nanticoke

Rd, Salisbury, MD 21801, United States of America

[5] Laboratoire associé au Centre national de référence pour les entérovirus & paréchovirus -

25-28, rue du Dr Roux 75015 Paris. France

Abstract

To date, 15 species have been described within the genus *Enterovirus*. Previous studies suggested the existence of another species comprising strains isolated from the stool specimens of non-human primates (NHPs) in Central Africa. Moreover, numerous full-length or partial genomic sequences of NHP enteroviruses (EVs) can be found in GenBank without being properly classified. To our knowledge, no comprehensive synthesis of NHP EV data exists, leaving genetic relationships between strains across independent studies unclear. To address these gaps, we sequenced the complete genome of four NHP EVs from our stool collection and conducted an extensive search of NHP EV sequences in GenBank to perform a comprehensive phylogenetic analysis. Our analyses revealed two new species tentatively named *Enterovirus mbel* and *Enterovirus noa*, which contain at least 6 and 2 virus types, respectively. We also identified new virus types within the known species EV-J.

Phylogenetic analyses strongly suggest interspecies recombination events between NHP EVs in the non-structural region of the genome, challenging the long-held view that recombination is confined within narrowly defined subsets of EVs belonging to the same species. We also performed the first comprehensive comparative analysis of full length human and NHP EV genomes, focusing on GC content, dinucleotide frequencies and codon-usage bias. GC content emerged as the most robust host-associated marker: all NHP-associated virus types within species *E. alphacoxsackie* and *E. betacoxsackie* displayed GC % below 47 %, whereas human-derived virus types exhibited GC % above 47 %. Dinucleotide frequency, Effective Number of Codons (ENC) and Relative Synonymous Codon Usage highlighted distinct codon-bias clusters that mirror the phylogenetic relationships between EVs but only partially correlate with their respective hosts of origin.

This work enhances our understanding of EVs circulating in NHPs and paves the way

for future research aiming at understanding the mechanisms underlying host-adaptation among

EVs.

Keywords

Enterovirus, *Picornaviridae*, Recombination, non-human primates, GC content, host-

adaptation

INTRODUCTION

Enteroviruses (EVs) are small, non-enveloped viruses with an icosahedral capsid. Their genome is a single-stranded and positive-sense RNA molecule of approximately 7,200-8,500 nucleotides (nt) in length. It contains two untranslated regions (5'UTR and 3'UTR) flanking a large open reading frame (ORF) that encodes a polyprotein, which is subsequently cleaved into three precursors (P1, P2 and P3). P1 gives rise to the four capsid proteins (VP1 to VP4) while P2 and P3 are cleaved into different proteins involved in viral replication. A short additional upstream ORF (uORF) encoding a single protein has been described in the genome of some EVs (Lulla et al., 2019).

The classification of EVs has undergone significant revisions since their discovery. Initially, EVs were classified into different groups based on their phenotypic properties (e.g. tropism and pathogenicity in mice) and physicochemical characteristics (density of the virus particles, sensibility to acid pHs) (Bessaud et al., 2018). Within each group, virus isolates were grouped into serotypes, based on *in vitro* cross-neutralization assays. In the 2000s, genetic criteria replaced phenotypic and serological criteria as the basis for classification (M. S. Oberste et al., 1999). Enteroviruses are now classified into species and virus types, depending on their phylogenetic relationships. According to the International Committee for Taxonomy of Viruses guidelines (ICTV), EVs belong to the same species if they share at least 70% amino acid (aa) identity in the complete ORF and in the 2C+3CD genes, and at least 60% aa identity in the capsid (P1 region). The *Enterovirus* genus currently comprises 15 species: *Enterovirus alphacoxsackie* (EV-A), *Enterovirus betacoxsackie* (EV-B), *Enterovirus coxsackiepol* (EV-C), *Enterovirus deconjuncti* (EV-D) *Enterovirus eibovi* (EV-E), *Enterovirus fitauri* (EV-F) *Enterovirus geswini* (EV-G), *Enterovirus hesimi* (EV-H), *Enterovirus idromi* (EV-I), *Enterovirus jesimi* (EV-J), *Enterovirus krodeni* (EV-K), *Enterovirus lesimi* (EV-L), *Enterovirus alpharhino* (RV-A), *Enterovirus betarhino* (RV-B) and *Enterovirus cerhino* (RV-C), each of

which comprise multiple virus types based on the VP1-encoding sequence. Virus types within

a species are defined by >75% nt identity and >85%-88% aa identity in this region (Brown et

al., 2009; M. Steven Oberste et al., 1999).

EVs infect a wide range of mammals, including humans, non-human primates (NHPs),

small ruminants, pigs, cattle, camels and rodents. While each EV appears adapted to a limited

set of host species that constitutes its natural reservoirs, host barriers can be overcome, as

demonstrated by numerous documented cross-species transmissions between humans and

animals, bidirectionally, or among different animal species (Doté et al., 2024; Fieldhouse et al.,

2018; Modiyinji et al., 2025). Notably, EVs do not exhibit clear host-associated phylogenetic

clustering, as some species include virus types circulating in different hosts. For instance, some

EV-As, EV-Bs and EV-Ds circulate in humans while others circulate in NHPs. Similarly, some

EV-Gs circulate in pigs while others circulate in small ruminants.

The first identified NHP EVs were detected in simian cell cultures and laboratory

animals in the mid-20[th] century (Oberste et al., 2002). Since then, new EV types have been

discovered in captive, wild, healthy and sick NHPs. Over the past 20 years, next-generation

sequencing has increased the number of publicly available NHP EV sequences, however, data

remain sparse. Furthermore, many sequences span diverse sub-genomic regions and remain to

be properly classified. Moreover, no comprehensive synthesis of all available NHP EV

molecular data exists, leaving the genetic relationships between NHP EVs across independent

studies fragmented. Additionally, sequences of NHP EVs sampled in Central Africa have

suggested the existence of at least one additional EV species (Sadeuh-Mba et al., 2014), but it

has never been formally described. To address these gaps, we conducted an extensive search of

NHP EV sequences in GenBank and performed a comprehensive phylogenic analysis. To make

this analysis as complete as possible, we also sequenced the complete genome of four NHP EVs

from our NHP stool collection and samples from our previous work (Sadeuh-Mba et al., 2014).

Furthermore, we compared genetic characteristics, such as GC percentage (GC%), dinucleotide

ratios or codon usage (effective number of codons, ENC and relative synonymous codon usage,

RSCU) between NHP and human EVs. Our analysis identified two previously uncharacterized

EV species and revealed that specific genomic signatures align with phylogeny and may be

associated with host-adaptation.

METHODS

Amplicon-based sequencing

EV-B114 strain Z057 was grown in RD cells and RNA was extracted from 250 µL of

clarified cell culture supernatant using the High Pure Viral RNA kit (Roche Diagnostics,

Meylan, France) following the manufacturer's instructions. Genome amplification was

performed by RT-PCR with primers targeting previously described conserved regions of the

EV genome (Joffret et al., 2018). Sequencing was performed by Illumina technology and de

novo assembly conducted with CLC Genomics Workbench as previously described (Bessaud

et al., 2016). As EV-J121 strain CHE20 could not be propagated in cell culture (Sadeuh-Mba

et al., 2014), RNA was extracted directly from 140 µL of clarified 20% (Weight/Volume) stool

suspension using the QIAamp Viral RNA Mini Kit (Qiagen, France). Genome amplification

was performed by RT-PCR with primers targeting the sub-genomic regions previously

sequenced (Sadeuh-Mba et al., 2014), as well as a primer targeting the 3' terminal poly A

sequence. Sequencing was performed by the Sanger technique.

Metatranscriptomic next-generation sequencing

Two strains, RCMH3 and RCMH6, that could not be grown in cell culture were

sequenced using a protocol previously described (Gámbaro et al., 2021). After extraction from

140 µl of fecal suspension using the QIAamp Viral RNA Mini Kit (Qiagen) followed by Turbo

DNAse treatment (Ambion), RNA was purifed with Agencourt RNAClean XP beads. Host

rRNA were depleted using custom probes and RNAse H treatment (Matranga et al., 2014).

After purification using AMPure RNA clean beads (Beckman Coulter Genomics) and elution

in 10 µl of RNase-free water, RNA was retro transcribed using random primers and SuperScript

IV (Invitrogen). Second-strand DNA was synthesized using *E. coli* DNA ligase, RNAse H and

DNA polymerase (New England Biolabs) and purified using Agencourt AMPure XP beads

(Beckman Coulter). Libraries were then prepared using the Nextera XT DNA Library Prep Kit

(Illumina), quality controlled using a Bioanalyzer high sensitivity DNA analysis kit (Agilent)

and sequenced on an Illumina NextSeq500 (2×75 cycles).

Adapters and low-quality reads were removed using Trimmomatic v0.39 (Bolger et al.,

2014). Trimmed reads were assembled using megahit v1.2.9 (Li et al., 2016) with default

parameters. The contigs were queried against the NCBI non-redundant protein database using

DIAMOND v2.0.4 (Buchfink et al., 2015), leading to the identification of novel EV sequences.

The EV genomes were then reconstructed by iterative mapping of the trimmed reads, using clc-

assembly-cell v5.1.0. We used Samtools v1.10 (Li et al., 2009) to generate alignment statistics,

and ivar v1.0 (Grubaugh et al., 2019) to generate consensus sequences, using a minimum of $3\times$

read depth coverage. The mapping data was visually checked to confirm the accuracy of the

obtained genomes using Geneious Prime 2023 (www.geneious.com).

Phylogenetic analysis

Multiple sequence alignments were performed with CLC Main Workbench 24.0.2

software (CLC bio, Aarhus, Denmark). Phylogenetic trees were inferred by the neighbour-

joining (NJ) or maximum likelihood (ML) methods implemented in MEGA X (Kumar et al.,

2018). Distance-based phylogenetic trees were reconstructed by the NJ method with the

Tajima-Nei model for computing evolutionary distances (Tajima and Nei, 1984). The ML

phylogenetic trees were reconstructed by the Tajima-Nei model and Nearest-Neighbor-

Interchange method for ML heuristic method. The reliability of tree topologies was estimated

by bootstrap analysis with 1,000 pseudoreplicates for NJ trees and 500 pseudoreplicates for ML

trees due to computational constraints. Tree representations were generated using MEGA X

(version 10.2.6) or FigTree (version 1.4.4) (https://tree.bio.ed.ac.uk/software/figtree/). Links to

Nextstrain interactive representations of trees from Figures 2 and 6, Supplementary Figure 1,

and tanglegrams are available on GitHub:

https://github.com/CorentinAube/NHP_EV_genomic_analysis.

Genomic comparison

Prototype genomes of human and NHP EVs were retrieved from the Picornaviridae

database (http://www.picornaviridae.com), while more recent sequences were retrieved from

the BV-BRC database (Olson et al., 2023). Other partial NHP EV sequences were retrieved

from the scientific literature indexed in PubMed. Protein sequence translation, MSA, nucleotide

and amino acid identity matrices were calculated with CLC Main Workbench. The GC content,

dinucleotide ratios, ENC and RSCU values were calculated using custom Python (version

3.12.12) scripts available on GitHub with also the list of all accession numbers of sequences

used (https://github.com/CorentinAube/NHP_EV_genomic_analysis). The formulas for

dinucleotide ratio, ENC and RSCU calculations were obtained from Zeng and colleagues (Zeng

et al., 2022). The determination of GC3 bias was based on the method published by Butt and

colleagues (Butt et al., 2016).

Data availability

189    Nucleotide sequences generated in this study were deposited on the GenBank database

190 (accession numbers PX682287-PX682290). An isolate of the EV-B114 strain Z057 has been

191 shared via the EVAg repository (Romette et al., 2018).

193 **RESULTS**

194 **Whole-genome sequencing of four NHP EVs from our stool collection**

195    Four EVs that had only been previously partially sequenced (Sadeuh-Mba et al., 2014),

196 were subjected to whole genome sequencing. Contigs covering nearly the entire viral genome

197 were obtained for three viruses: Z057, CHE20, and RCMH3 (Figure 1A). For RCMH6,

198 approximately 150 nt could not successfully be sequenced at the 5' extremity. All 4 viruses

199 featured the canonical EV genomic organization, with a large ORF flanked by two UTRs.

200 Comparing these sequences with annotated reference EV genomes, canonical polyprotein

201 cleavage sites were identified, suggesting that the polyprotein processing in these viruses is

202 similar to that of other EVs (Supplementary Table 1). Additionally, the genomes of Z057,

203 RCMH3, and RCMH6 contained a putative uORF. This putative uORF overlaps with the main

204 ORF with a frameshift, and its putative start codon is located 133 or 136 nt upstream of the

205 main ORF (Figure 1B). In contrast, the EV-J121 CHE20 genome lacked a uORF, as previously

206 reported for other members of the EV-J species (Lulla et al., 2019). All four genomes also

207 contained a cis-replication element (CRE) within the 2C gene, matching the consensus

208 sequence 5'-GXXXAAAXXXXXXA-3' (Figure 1C) (Cordey et al., 2008).

210 **Comprehensive phylogeny of non-human primate enteroviruses**

211    The search for NHP EV sequences in GenBank led to the identification of 336

212 sequences, of which 24 were complete genomes, 91 were VP1 sequences, 103 were VP4/2

sequences, 19 were VP2 sequences, 26 were 3D sequences and 29 were sequences covering

various regions of the genome, referred to as partial (Table 1).

A phylogenetic tree based on the complete VP1 gene was generated using all available

NHP EV sequences spanning this region along with representatives of all known human derived

EV types (Figure 2 and Supplementary Figure 1). This VP1-based tree clearly delineates the

EV species. Non-human primate derived EVs were found in species EV-A, -B, -D, -H, -J and -

L. The data for EV-L are very sparse with only one known representative. Species EV-H

includes only 6 sequences, of which 4 are epidemiologically related to each other. To determine

whether these sequences belong to the same virus type, the 75% nt and 85% aa identity rule

(Brown et al., 2009) was applied to their VP1 sequences. As previously described (Nguyen et

al., 2014), these sequences belong to a single virus type (EV-H1), sharing 84.09%-100% nt

identity and 96.17%-100% aa identity (Supplementary Figure 2). In contrast to the EV-H and

EV-L species, distinct EV-J virus types have been previously described. The capsid-based

phylogram accurately delineated the EV-J103, EV-J108 and EV-J122 virus types and confirmed

the EV-J121 type, of which CHE20 is the first known representative that was previously

characterized based on partial sequence (Sadeuh-Mba et al., 2014) (Figure 3A). In the VP4/VP2

region, three strains (CPML8127, CPOV3340 and CPOV3342) were closely related to CHE20

(Figure 3C), thus suggesting that they could also be assigned to the EV-J121 virus type. Identity

matrices were calculated to assign several untyped EV-J sequences (Figure 3A and 3B,

Supplementary Figure 3). NOLA1 and NOLA3 share over 75.41% and 85.31% at nt and aa

level, respectively, with an EV-J103 strain and were thus assigned to this virus type. Similarly,

the sequence comparison of the partial VP1 sequence of cg5275 with EV-J prototype strains

supported its assignment to the EV-J103 type (>77.36% and >85.43 nt and aa identity,

respectively, with other EV-J103 viruses, Supplementary Figure 4). As previously reported

(Wang et al., 2016) the Sev-nJ strain represent a novel virus type, sharing less than 74.48% and

84.27% with other EV-J sequences (Supplementary Figure 3). The CHN-BJ-LXL1 sequence, published 5 years later than the Sev-nJ strain, belonged to the same EV-J type, as it shares 82.09% and 92.63% nt and aa identity with Sev-nJ, respectively (Supplementary Figure 3). We propose to assign these two strains to a novel virus type, EV-J123, following current taxonomic nomenclature. Additionally, several partial VP1 and VP4/2 sequences remained to be assigned (Harvala et al., 2014). Based on partial VP1 sequences, the CPOV3336 strain is related to EV-J121 strain CHE20 (Figure 3B), but their nt and aa identities (66.04% and 75.81%, Supplementary Figure 4) indicating that CPOV3336 belongs to a putative distinct virus type (tentatively named EV-J124). A set of short VP4/VP2 sequences (CPOV3337, DJ4139, CPGR7921, CPLM8102 and CPLM8125) were very close that of CPOV3336 (Figure 3C), suggesting that they belong to the newly proposed EV-J124 type.

Some viruses previously identified as EV-Js (CHN-BJ-2J, -1J, -8J, -10J and -9J) form a specific group (Figure 3). To determine whether these viruses defined a separated additional new species, we applied the ICTV criteria. The CHN-BJ-2J, 1J, 8J, 10J and 9J sequences share 69.41% to 72.74% identity in the ORF and 80.72% to 84.50% in the 2C+3CD genes with other EV-J types (Supplementary Figure 5A and B). However, they share only between 55.36% to 59.25% aa identity in the capsid gene (Supplementary Figure 5C), as previously reported (Li et al., 2020). These results support the existence of a separated new species, which we tentatively named *Enterovirus nao* (EV-N), after the Chinese word náo (猱) meaning "monkey" (Couvreur, 1896; Loewe, 1993). As previously suggested (Li et al., 2020), CHN-BJ-1J, 8J, 9J, 10J and CHN-BJ-1-2J form two distinct EV types within this species, as they share less than 72.94% nt and 80.99% aa identity (Supplementary Figure 3). We propose naming these types EV-N1 (for the CHN-BJ-1J, 8J, 9J, 10J strains) and EV-N2 (for the CHN-BJ-2J strain).

Three EV species, EV-A, -B and -D, include some EVs that originated from humans while others were recovered from NHPs. Within the EV-B species, NHP EVs segregate from

human derived EVs and define a specific branch (Figure 2). The unique exception to the host association of EV-B types is the case of GAB98 strain, isolated from wild chimpanzees, which is an EV-B107 known to circulate in humans in Asia (Maan et al., 2019; Yamashita et al., 2010). The isolation of strain GAB98 in a chimpanzee was explained as a sporadic human-to-NHP transmission, potentially facilitated by the adjacent or overlapping humans and chimpanzee habitats (Mombo et al., 2017). As previously reported, 4 distinct virus types exist in the NHP specific branch of the EV-B species. Strain Z057, of which a complete genome sequence was obtained in this work (Figure 1), belongs to the EV-B114 virus type (Figure 4), that was first isolated in 1963 from NHP derived cell cultures (Malherbe and Harwin, 1963). Since its discovery, only two EV-B114 EVs have been reported: CPOV3344 (Nguyen et al., 2014) and Z057 (Sadeuh-Mba et al., 2014). They share 82.09%-84.58% nt and 92.88%-93.95% aa identities between each other in the VP1 sequence; thus confirming that they belong to the same virus type (Figure 4). In contrast to EV-B species, the phylogenetic boundaries between NHP and human EVs are less distinct in species EV-A and EV-D (Figure 2). As previously reported, members of the species EV-A fall into four main phylogroups. Clades 1 and 2 consist of EV types that circulate in humans worldwide (Clade 1: EV-A120, EV-A71, CVA7, CVA14 and CVA16; Clade 2: EV-A114, CVA2, CVA3, CVA4, CVA5, CVA6, CVA8, CVA10 and CVA12). Clade 3 includes EVs detected both in human and NHPs (Clade 3: EV-A76, EV-A89, EV-A90, EV-A91, EV-A119, EV-A121 and EV-A125). Finally, clade 4 contains only NHP EV strains (Clade 4: EV-A122, EV-A123 and EV-A124). Similarly, within EV-D, some types are human-specific (EV-D70, EV-D68 and EV-D94), or NHP-specific (EV-D120), while EV-D111 has been detected in both hosts (Sadeuh-Mba et al., 2019). No unidentified or misidentified NHP derived EV-A -B or -D were found in public databases.

Notably, 7 sequences spanning the entire main ORF (RCMH3, RCMH6 -both sequenced in this study-, CPML3961, CPML8109, GR2815 and NGR_NHP_1) clustered together in the

VP1-based phylogenetic tree and did not align with any known species (Supplementary Figure 1). To determine whether these sequences belong to a single new species, we calculated their pairwise nt and aa identities (Supplementary Figure 6). Our analysis showed that the corresponding viruses met the ICTV criteria for classification as the same species: they share over 70% aa identity in the ORF (85.69%-95.02%) and in the 2C+3CD sequences (97.03%-99.69%), as well as over 60% aa identity in the capsid sequences (69.27%-89.82%). Therefore, we propose the creation of a new species, tentatively named *Enterovirus mbel* (EV-M), after the name of *Pterocarpus soyauxii*, a tree native to the tropical forest in Central African (Albreht et al., 2025), known as *mbel* in various Bantu languages spoken in Central and Southern Cameroon, where the first representatives of this species were sampled (Galley, 1964; Tsala, 1956). Phylogenetic analyses of the VP1 gene revealed several clusters within this new EV-M species (Figure 5). Based on the established thresholds for discriminating EV types (>75% nt identity and >85% aa identity in the VP1 gene) (Brown et al., 2009), we identified 6 distinct virus types (Supplementary Figure 7): EV-M1 (prototype: RCMH6), EV-M2 (prototype: RCMH3), EV-M3 (prototype: CPML3961), EV-M4 (prototype: CPML8109), EV-M5 (prototype: GR2815) and EV-M6 (prototype: NGR_NHP_1). Identity matrices showed that numerous VP1 sequences deposited in GenBank belong to these newly defined EV-M virus types (Supplementary Figure 8 and Figure 5). EV-M1, EV-M2, EV-M4 and EV-M5 virus types gathered sequences that displayed >78.57% nt and >90.62% aa identities between each other. Within the EV-M3 type, all sequences except two (LM1962 and CPOV3334) displayed a nt identity >77.04% with the prototype strain CPML3961, exceeding the 75% threshold. The nt identities of LM1962 and CPOV3334 compared to CPML396 fell slightly below this threshold (73.22% and 74.14%, respectively), but both shared an aa identity >86.0% with the prototype, supporting their classification as EV-M3. Besides the prototype NGR_NHP_1, no additional members of virus type EV-M6 were identified.

Shorter VP1 or VP2 sequences, averaging 350 nt in length, have also been published

and annotated as belonging to the species EV-J (Mombo et al., 2023, 2021, 2017). However,

dendrograms based on these regions revealed that they actually cluster within the EV-M species

(Supplementary Figure 9A and 9B). The corresponding sequences are too short to allow

accurate typing of the corresponding viruses.

Collectively, these results reveal a complex landscape of NHP EV diversity, including

two new species (EV-M and EV-N) and two new EV-J types (EV-J123 and EV-J124). This

refined classification will facilitate the proper assignation of NHP EV strains to their respective

species and virus type.

Phylogenetic relationships in the non-structural genomic regions

Recombination events are frequent among co-circulating EVs, generating new viruses

with chimeric genomes that may outcompete parental strains if they confer a fitness advantage.

Because recombination constantly reshuffles EV genomes, different genomic regions are not

subjected to the same extend of selective pressure and do not follow the same evolutionary

trajectories. After clarifying the phylogenetic relationships among the NHP derived EVs in the

capsid-encoding region, we investigated their evolutionary history by constructing

phylogenetic trees considering 2A2B, 2C, 3A3B, 3C, and 3D regions, using all available full-

length ORF sequences (Figure 6). As previously reported (Mombo et al., 2017; Oberste et al.,

2007; Sadeuh-Mba et al., 2023), striking intra-species phylogenetic incongruences were

observed between trees generated from structural (capsid) and non-structural (2A2B, 2C, 3A3B

and 3D) regions. Within the EV-B species, the NHP and human derived EVs formed two distinct

clades, consistent with the VP1 phylogeny (Figure 6). Of note, all known EV-Ms clustered with

the NHP EV-B clade across all non-structural regions. Consequently, EV-Ms and NHP EV-Bs

are phylogenetically indistinguishable from each other across the non-structural part of the

genome. Similarly, as previously reported (Oberste et al., 2007; Sadeuh-Mba et al., 2023), EV-

A125 strain BA13 did not cluster with other EV-As in the non-structural regions downstream the capsid. Our analyses further revealed that BA13 strain clustered with EV-J108 strains in the 2A2B, 2C and 3A3B regions, with the EV-J103 strains in the 3C gene and with the members of the new species EV-N in the 3D gene (Figure 6). Finally, the species EV-D, EV-H, and EV-L exhibited complex phylogenetic relationships. Indeed, they form a single group in the 2A2B region, but segregated into three distinct branches in the 3A3B and 3C based trees. The other regions showed specific pairwise associations: EV-D grouped with EV-H and EV-L in the 2C region and the 3D region, respectively.

Comparison of genomic signatures of human and NHP derived EVs

To our knowledge, no comparative genomic analysis of human and NHP derived EV species has been published so far. A previous study, limited to EV-A species, revealed that clade 3 viruses displayed distinct relative dinucleotide ratios and codon usage biases compared to human EV-As of clades 1 and 2 (Zeng et al., 2022), suggesting that specific genomic signatures could be associated with the virus' natural history and, particularly its host range. To determine whether NHP derived EVs from different host species shared unique genomic traits, we compared GC content (GC%), dinucleotide ratio, and codon usage (ENC and RSCU analysis) between human and NHP derived EVs. To minimize potential bias in these analyses, we included, where possible, at least the prototype strain and two additional genomes sampled from different countries for each virus type.

The GC% ranged from 41% to 49% across all species (Figure 7A). Interestingly, the GC% strongly correlated with the host origin rather than EV species classification, with all NHP virus types featuring a GC% <47%. In particular, NHP-specific EV species (EV-J, EV-H, EV-L, EV-M and EV-N) display low GC%, ranging from 43% to 46.4%, as did the few known EV-

D virus types (41.0%-43.5%). Strikingly, virus types segregated with respect to their host of origin within the EV-A and EV-B species: all EV-A virus types of clades 1 and 2, originating from humans, displayed GC%>47%, while NHP-associated clades 3 and 4 had GC%<47% (Figure 7B). A similar clustering was found within EV-B species, with all human derived EV types featuring a GC% >47%, while all NHP derived EV types had a GC% <45% (Figure 7C). Notably, the GC % of EV-C virus types, which are restricted to human hosts, mirrored their phylogenetic relationships: respiratory virus types (EV-C104, EV-C105, EV-C109, EV-C117 and EV-C118) exhibited a high GC % (48–48.5%), cultivable enteric virus types (PV1, PV2, PV3, CVA11, CVA13, CVA17, CVA20, CVA21, CVA24, EV-C95, EV-C96, EV-C99 and EV-C102) displayed an intermediate GC % (45–46.5 %), while non-cultivable viruses (CVA1, CVA19, CVA22, EV-C113, EV-C116 and EV-C119) showed a low GC % (43-44 %) (Figure 7D).

A previous study revealed that EV-As clade 3 viruses display lower CpG and UpA dinucleotide frequencies compared to clades 1 and 2 (Zeng et al., 2022). To determine whether NHP derived EV-As of clade 4 share this trait, we analyzed dinucleotide ratios of EV-A strains representative of the different clades (Figure 8). Our analysis revealed that EV-A viruses belonging to clade 4 had the same low CpG ratio as clade 3 EV-As (Figure 8B). The same observation was made within EV-B species, with NHP derived EV types displaying a CpG ratio substantially lower than that of human derived EV types (NHP EVs<0.42 vs Human EVs>0.56) (Figure 8C). With the exception of EV-H, a low CpG ratio seems to be a common characteristic of NHP derived EVs. Indeed, members of the EV-J, EV-L, EV-M and EV-N species featured CpG ratios<0.5 and comparable to those of NHP derived EV-A and EV-B species (Figure 8A). It is noteworthy that all EV-D virus types had CpG ratio<0.35, irrespective of their host of origin.

We also investigated codon usage bias in human and NHP derived EVs. The effective

number of codons (ENC), which ranges from 20 (strong bias, with one codon used for each

amino-acid) to 61 (no bias, even use of each codon) (Wright, 1990), revealed distinct patterns.

Human EV-As (clades 1 and 2), EV-Bs, respiratory EV-Cs, and NHP derived EV-Hs had ENC

greater than 54 (Figure 9A), while clade 3, NHP derived EV-As, NHP derived EV-Bs, cultivable

and non-cultivable EV-Cs, EV-Ds, EV-Js and EV-Ms featured ENC values between 50 and 54.

The sole known EV-L had an ENC value of 48. Since the GC% at the third codon position

(GC3) could introduce a bias in the ENC analysis, we compared the ENC values for each EV

species with the expected ENC based on the respective GC3 percentage (Butt et al., 2016)

(Supplementary Figure 10). All calculated ENC values fell below the expected curve,

confirming that GC3 content did not account for the variations across virus group.

We also analyzed the Relative Synonymous Codon Usage (RSCU), which quantifies

how frequently each synonymous codon is used in a gene or genome relative to the expected

usage. The results, summarized in a cladogram, grouped EV types into three main clusters

(Figure 9B). Group A contains only the EV-H1 type. Group B includes all other NHP-specific

EV types (except EV-A123), whichever their species, all human cultivable EV-Cs, all EV-Ds

and EV-A clades 3 and 4. Group C contains all other human EV-Cs and EV-A123. These results

suggest that codon-usage bias varies markedly among EV groups, yet it only partially mirrors

the underlying phylogenetic relationships among them.

Collectively, these results reveal that dinucleotide ratios, ENC and RSCU values vary

among NHP EV species. Remarkably, GC% mirrors phylogenetic relationships between EVs

and could serve as a predictive marker of human versus NHP origin of EVs.

DISCUSSION

For decades, no standardized nomenclature existed for NHP derived EVs. In the 1980s, it was proposed to designate these viruses under the common name, *Simian enterovirus*, followed by a number corresponding to a given serotype (Kalter, 1982). For a long time, NHP derived EVs were not assigned to specific species. In the 2000s and 2010s, the generalization of genetic sequencing brought about major changes in EV classification, as phylogenetic relationships replaced phenotypic criteria to delineate EV types and species. Based on their genetic relationships, NHP derived EVs were classified into different species, some of which also included human derived EVs. By considering all NHP EV genome sequences available in public databases and integrating our two new full-genome sequences, our analyses confirmed the existence of a new species (Figure 5 and Supplementary Figure 6) that had been previously suggested by two independent studies (Nguyen et al., 2014; Sadeuh-Mba et al., 2014). We found that this new EV species gathered viruses sampled in three countries, Gabon, Cameroon, and Nigeria (Mombo et al., 2023, 2021, 2017; Nguyen et al., 2014; Sadeuh-Mba et al., 2023). This geographic distribution justifies the name proposed for this species, *Enterovirus mbel*, which refers to an endemic tree of this region, called African padauk in English (Albreht et al., 2025). We also identified another new EV species, comprising a few viruses sampled in China previously misidentified as EV-Js (Li et al., 2020) (Figure 3 and Supplementary Figure 5). Since this species has only been documented in China, we propose naming it *Enterovirus nao,* after the Chinese word referring to a monkey in a poem written 2,500-3,000 years ago (Couvreur, 1896; Loewe, 1993). However, it is important to note that the genuine geographic area of circulation of these two new EV species remain to be documented, especially for *E. nao* which was detected in captive animals.

The discovery of the uORF in the genome of some EVs a few years ago constituted a drastic change in our understanding of their biology. This uORF was experimentally shown to encode a non-essential protein that enhances virus replication in gut cells (Lulla et al., 2019).

Interestingly, the putative uORFs identified in the genomes of EV-B114 Z057, EV-M1 RCMH6, and EV-M3 RCMH3 are respectively located 134 nt, 133 nt, and 136 nt upstream of the AUG start codon of the main ORF. These distances fall below the cutoff set by Lulla et al. for screening for uORF of EV genomic sequences (Lulla et al., 2019). Although functional validation of these candidate uORFs is still lacking, our findings suggest that the prevalence of uORFs across EVs may have been underestimated and warrants re-evaluation using a more permissive distance threshold.

The phylogenetic analyses performed on the non-structural part of the genome further confirmed the close genetic relationships between NHP EVs recovered from different species, as previously documented (Figure 6) (Mombo et al., 2017; Nguyen et al., 2014; Oberste et al., 2007; Sadeuh-Mba et al., 2014). This close genetic relatedness could be explained by convergent evolution, which may have driven EV genomes from different species to become phylogenetically more similar as a consequence of their replication within NHP hosts. However, it is unlikely that convergent evolution could uniformly affect the entire non-structural region of the genome, as it would be the case for NHP derived EV-Bs and EV-Ms that cluster together across this region. Furthermore, convergent evolution among NHP derived EVs is not supported by the fact that most known NHP derived EV-As do not cluster with other NHP derived EVs, except for EV-A125 strain BA13, which is phylogenetically close to EV-Js downstream the capsid. If convergent evolution was shaping the evolution of EV-A125, it would be expected to similarly affect other NHP derived EV-As, which seems unlikely. Therefore, these phylogenetic incongruences found in this study are probably due to recombinations, a well-established driver of EV evolution (Muslin et al., 2019).

While interspecies recombination has been observed in the 5'UTR (Boros et al., 2012; Muslin et al., 2015; Santti et al., 1999), it has never been documented downstream of the capsid-encoding region among human derived EVs. Even intraspecies recombination is limited to

specific subsets of EV types that can recombine with one another but not with members of other subsets. For instance, EV-As of clade 3 do not exchange genetic sequences with EV-As of clades 1 and 2 (Oberste et al., 2005; Sadeuh-Mba et al., 2014). Similarly, respiratory EV-Cs and non-cultivable EV-Cs do not recombine with each other or with cultivable EV-Cs (Brouwer et al., 2020). Several factors could explain the apparent inability of some EVs to generate recombinant viruses, such as functional incompatibilities between genomic regions from distant phylogenetic groups, lack of fitness advantages conferred to the recombinant viruses, or low frequency of co-infection due to different host and tissue tropisms. Our analysis provided substantial evidence to the existence of interspecies recombination between some NHP derived EV species, particularly between EV-Ms and EV-Bs. These findings change our understanding of recombination between EVs, which appear to be capable of producing larger leaps than previously believed. This raise fundamental questions about the past and future evolutionary path of these viruses. For instance, it is unknown whether NHP derived EV-Bs acqired non-structural genes from EV-Ms, or vice-versa, unless both inherited these genes from a yet-unidentified common source. The ability of EV-Ms to recombine with human derived EV-Bs and the potential for such inter-species macro-evolutionary events to produce new human-adapted EV-Ms are also difficult to predict.

Previous studies have shown that genomes of clade 3 EV-As, which were detected in both NHPs and humans, had a lower CpG frequency than human specific EV-As and featured a specific codon usage bias that impairs their expression of these EV-A genomes in human cells (Zeng et al., 2022). These observations led us to hypothesize that some genetic characteristics shared by NHP derived EVs may play a role in host adaptation. Nucleotide composition has been previously used to predict the host of novel picorna-like viruses (Kapoor et al., 2010). While our analysis of dinucleotide frequency and codon usage bias across the various EV species circulating in humans and in NHPs did not reveal a clear segregation between human

and NHP derived viruses (Figure 9), our results identified a relatively low GC% appears as a

signature shared by virus types circulating among NHPs, regardless of their virus species

(Figure 7). Unfortunately, the lack of data precluded robust statistical analysis, as some EV

types are represented by hundreds of sequences while others, such as EV-L species, are

represented by only one or two sequences. Based on genetic and epidemiological data, we and

others have previously postulated that EV-Ds known in humans may have recently emerged

from zoonotic sources, presumably NHPs (Kono et al., 1981; Miyamura et al., 1986; Sadeuh-

Mba et al., 2019). The low GC% values displayed by all EV-D virus types, irrespective of their

tissue tropism or host of origin, are consistent with that hypothesis. The relative relatedness of

EV-Ds and NHP-specific EV species in the non-structural part of the genome further supporting

the hypothesis of a zoonotic origin of EV-Ds. However, host adaptation is not the only driver

of GC%, as illustrated by EV-C virus types, which display a broad range of GC% values despite

all circulating in humans (Figure 7). In addition, the GC% of human and NHP EV strains falls

within the broad range observed in the human transcriptome (average 48%, varying from <40%

to >60% among mRNAs) (Courel et al., 2019; Piovesan et al., 2019). Unfortunately, the GC%

of transcriptomes of NHP infected with EVs remains uncharacterized to our knowledge. The

GC% overlap between EVs and human transcriptome is insufficient to conclude host-species

adaptation, and the lack of data prevents comparative analysis that could elucidate potential

host-driven GC% adaptation in NHP EV strains. The genomic signature result from a dynamic

equilibrium influenced by many factors, of which polymerase fidelity, RNA secondary structure

requirements, host translational preferences, immune pressure, and genetic recombination.

Furthermore, identifying genomic signatures that reliably distinguish NHP derived EVs from

human derived EVs is currently limited by the paucity of fully sequenced strains, our limited

knowledge of the natural history of these viruses, and particularly the uncertainty surrounding

their host tropisms.

ACKNOWLEDGEMENT

The authors are indebted to Dr. WEI Yu, who shared her knowledge of ancient Chinese poetry, and Dr. Nick J. Knowles, for his feedback regarding enterovirus phylogeny. This work has used the computational and storage services (Maestro cluster) provided by the IT department at Institut Pasteur, Paris.

AUTHORS CONTRIBUTIONS

MB and SASM conceived the study. SASM, PCdC, MP, MLJ, MCEZ, AB and ESL generated the sequencing data. CA, ESL, and MB performed the analyses. CA, NJ and MB wrote the article, which was approved by all co-authors.

FUNDING

This project was funded by the Institut Pasteur through a program of incentive actions (*Actions concertée inter-pasteuriennes*, ACIP-162). CA's salary is funded by a grant of the Gates Foundation (INV-067045). The E.S.-L. laboratory is funded by Institut Pasteur, the INCEPTION program (Investissements d'Avenir grant ANR-16-CONV-0005), the Ixcore foundation for research, the French Government's Investissement d'Avenir programme, Laboratoire d'Excellence 'Integrative Biology of Emerging Infectious Diseases' (grant no. ANR-10-LABX-62-IBEID), the HERA Projects DURABLE (grant no 101102733) and LEAPS (grant no 101094685). The funders of this study had no role in study design, data collection, analysis and interpretation, or writing of the article.

REFERENCES

Albreht, L., Bourland, N., Klitgård, B., Schley, R., Janssens, S.B., Hardy, O.J., 2025. Species delimitation and phylogenomics of economically important African Pterocarpus trees, with an implication for the development of DNA-based species identification tools. Mol Phylogenet Evol 204, 108277. https://doi.org/10.1016/j.ympev.2024.108277

Bessaud, M., Blondel, B., Joffret, M.-L., Mac Kain, A., Delpeyroux, F., 2018. Nomenclature et classification des entérovirus : quelle histoire ! Virologie (Montrouge) 22, 289–303. https://doi.org/10.1684/vir.2018.0756

Bessaud, M., Sadeuh-Mba, S.A., Joffret, M.-L., Razafindratsimandresy, R., Polston, P., Volle, R., Rakoto-Andrianarivelo, M., Blondel, B., Njouom, R., Delpeyroux, F., 2016. Whole Genome Sequencing of Enterovirus species C Isolates by High-Throughput Sequencing: Development of Generic Primers. Front Microbiol 7, 1294. https://doi.org/10.3389/fmicb.2016.01294

Bolger, A.M., Lohse, M., Usadel, B., 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. Bioinformatics 30, 2114–2120. https://doi.org/10.1093/bioinformatics/btu170

Boros, Á., Pankovics, P., Knowles, N.J., Reuter, G., 2012. Natural interspecies recombinant bovine/porcine enterovirus in sheep. J Gen Virol 93, 1941–1951. https://doi.org/10.1099/vir.0.041335-0

Brouwer, L., Benschop, K.S.M., Nguyen, D., Kamau, E., Pajkrt, D., Simmonds, P., Wolthers, K.C., 2020. Recombination Analysis of Non-Poliovirus Members of the Enterovirus C Species; Restriction of Recombination Events to Members of the Same 3DPol Cluster. Viruses 12, 706. https://doi.org/10.3390/v12070706

Brown, B.A., Maher, K., Flemister, M.R., Naraghi-Arani, P., Uddin, M., Oberste, M.S., Pallansch, M.A., 2009. Resolving ambiguities in genetic typing of human enterovirus species C clinical isolates and identification of enterovirus 96, 99 and 102. J Gen Virol 90, 1713–1723. https://doi.org/10.1099/vir.0.008540-0

Buchfink, B., Xie, C., Huson, D.H., 2015. Fast and sensitive protein alignment using DIAMOND. Nat Methods 12, 59–60. https://doi.org/10.1038/nmeth.3176

Butt, A.M., Nasrullah, I., Qamar, R., Tong, Y., 2016. Evolution of codon usage in Zika virus genomes is host and vector specific. Emerg Microbes Infect 5, e107. https://doi.org/10.1038/emi.2016.106

Cordey, S., Gerlach, D., Junier, T., Zdobnov, E.M., Kaiser, L., Tapparel, C., 2008. The cis-acting replication elements define human enterovirus and rhinovirus species. RNA 14, 1568–1578. https://doi.org/10.1261/rna.1031408

Courel, M., Clément, Y., Bossevain, C., Foretek, D., Vidal Cruchez, O., Yi, Z., Bénard, M., Benassy, M.-N., Kress, M., Vindry, C., Ernoult-Lange, M., Antoniewski, C., Morillon, A., Brest, P., Hubstenberger, A., Roest Crollius, H., Standart, N., Weil, D., 2019. GC content shapes mRNA storage and decay in human cells. Elife 8, e49708. https://doi.org/10.7554/eLife.49708

Couvreur, S., 1896. Chant IX. Kio Koung, in: Cheu King. Texte Chinois Avec Une Double Traduction En Français et En Latin, Une Introduction et Un Vocabulaire. Imprimerie de la Mission catholique, Ho-kien fou, pp. 302–304.

Doté, J., Joffret, M.-L., Beta, B.N., Ait-Ahmed, M., Banga-Mingo, V., Knowles, N.J., Jouvenet, N., MBaïkoua, M.-N., Gouandjika-Vasilache, I., Bessaud, M., 2024.

579   Characterization of enteroviruses circulating among farm animals and children in
580   Central African Republic. Emerg Microbes Infect 13, 2368212.
581   https://doi.org/10.1080/22221751.2024.2368212
582 Fieldhouse, J.K., Wang, X., Mallinson, K.A., Tsao, R.W., Gray, G.C., 2018. A systematic
583   review of evidence that enteroviruses may be zoonotic. Emerging Microbes &
584   Infections 7, 1–09. https://doi.org/10.1038/s41426-018-0159-1
585 Galley, S., 1964. Dictionnaire Fang -Français et Français - Fang. Éditions Henri
586   Messeiller, Neuchatel, Suisse.
587 Gámbaro, F., Pérez, A.B., Agüera, E., Prot, M., Martínez-Martínez, L., Cabrerizo, M.,
588   Simon-Loriere, E., Fernandez-Garcia, M.D., 2021. Genomic surveillance of
589   enterovirus associated with aseptic meningitis cases in southern Spain, 2015–
590   2018. Sci Rep 11, 21523. https://doi.org/10.1038/s41598-021-01053-4
591 Grubaugh, N.D., Gangavarapu, K., Quick, J., Matteson, N.L., De Jesus, J.G., Main, B.J.,
592   Tan, A.L., Paul, L.M., Brackney, D.E., Grewal, S., Gurfield, N., Van Rompay, K.K.A.,
593   Isern, S., Michael, S.F., Coffey, L.L., Loman, N.J., Andersen, K.G., 2019. An
594   amplicon-based sequencing framework for accurately measuring intrahost virus
595   diversity using PrimalSeq and iVar. Genome Biol 20, 8.
596   https://doi.org/10.1186/s13059-018-1618-7
597 Harvala, H., Van Nguyen, D., McIntyre, C., Ahuka-Mundeke, S., Ngole, E.M., Delaporte,
598   E., Peeters, M., Simmonds, P., 2014. Co-circulation of enteroviruses between
599   apes and humans. J Gen Virol 95, 403–407. https://doi.org/10.1099/vir.0.059048-
600   0
601 Joffret, M.-L., Polston, P.M., Razafindratsimandresy, R., Bessaud, M., Heraud, J.-M.,
602   Delpeyroux, F., 2018. Whole Genome Sequencing of Enteroviruses Species A to D
603   by High-Throughput Sequencing: Application for Viral Mixtures. Front Microbiol 9,
604   2339. https://doi.org/10.3389/fmicb.2018.02339
605 Kalter, S.S., 1982. Enteric viruses of nonhuman primates. Vet Pathol Suppl 7, 33–43.
606 Kapoor, A., Simmonds, P., Lipkin, W.I., Zaidi, S., Delwart, E., 2010. Use of nucleotide
607   composition analysis to infer hosts for three novel picorna-like viruses. J Virol 84,
608   10322–10328. https://doi.org/10.1128/JVI.00601-10
609 Kono, R., Sasagawa, A., Yamazaki, S., Nakazono, N., Minami, K., Otatsume, S., Robin, Y.,
610   Renaudet, J., Cornet, M., Afoakwa, S.N., Mingle, J.A., Obinim, J.K., Huros, A.,
611   1981. Seroepidemiologic studies of acute hemorrhagic conjunctivitis virus
612   (enterovirus type 70) in West Africa. III. Studies with animal sera from Ghana and
613   Senegal. Am J Epidemiol 114, 362–368.
614   https://doi.org/10.1093/oxfordjournals.aje.a113203
615 Kumar, S., Stecher, G., Li, M., Knyaz, C., Tamura, K., 2018. MEGA X: Molecular
616   Evolutionary Genetics Analysis across Computing Platforms. Mol Biol Evol 35,
617   1547–1549. https://doi.org/10.1093/molbev/msy096
618 Li, D., Luo, R., Liu, C.-M., Leung, C.-M., Ting, H.-F., Sadakane, K., Yamashita, H., Lam, T.-
619   W., 2016. MEGAHIT v1.0: A fast and scalable metagenome assembler driven by
620   advanced methodologies and community practices. Methods 102, 3–11.
621   https://doi.org/10.1016/j.ymeth.2016.02.020
622 Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis,
623   G., Durbin, R., 1000 Genome Project Data Processing Subgroup, 2009. The
624   Sequence Alignment/Map format and SAMtools. Bioinformatics 25, 2078–2079.
625   https://doi.org/10.1093/bioinformatics/btp352

Li, W., Qiang, X., Qin, S., Huang, Y., Hu, Y., Bai, B., Hou, J., Gao, R., Zhang, X., Mi, Z., Fan, H., Ye, H., Tong, Y., Mao, P., 2020. Virome diversity analysis reveals novel enteroviruses and a human picobirnavirus in stool samples from African green monkeys with diarrhea. Infect Genet Evol 82, 104279. https://doi.org/10.1016/j.meegid.2020.104279

Loewe, M., 1993. Shih ching, in: Early Chinese Texts, Early China Special Monographs Series. The Society for the Study of Early China and The Institute of East Asian Studies, University of California, Berkeley, pp. 415–423.

Lulla, V., Dinan, A.M., Hosmillo, M., Chaudhry, Y., Sherry, L., Irigoyen, N., Nayak, K.M., Stonehouse, N.J., Zilbauer, M., Goodfellow, I., Firth, A.E., 2019. An upstream protein-coding region in enteroviruses modulates virus infection in gut epithelial cells. Nat Microbiol 4, 280–292. https://doi.org/10.1038/s41564-018-0297-1

Maan, H.S., Dhole, T.N., Chowdhary, R., 2019. Identification and characterization of nonpolio enterovirus associated with nonpolio-acute flaccid paralysis in polio endemic state of Uttar Pradesh, Northern India. PLoS One 14, e0208902. https://doi.org/10.1371/journal.pone.0208902

Malherbe, H., Harwin, R., 1963. The cytopathic effects of vervet monkey viruses. S Afr Med J 37, 407–411.

Matranga, C.B., Andersen, K.G., Winnicki, S., Busby, M., Gladden, A.D., Tewhey, R., Stremlau, M., Berlin, A., Gire, S.K., England, E., Moses, L.M., Mikkelsen, T.S., Odia, I., Ehiane, P.E., Folarin, O., Goba, A., Kahn, S.H., Grant, D.S., Honko, A., Hensley, L., Happi, C., Garry, R.F., Malboeuf, C.M., Birren, B.W., Gnirke, A., Levin, J.Z., Sabeti, P.C., 2014. Enhanced methods for unbiased deep sequencing of Lassa and Ebola RNA viruses from clinical and biological samples. Genome Biol 15, 519. https://doi.org/10.1186/PREACCEPT-1698056557139770

Miyamura, K., Tanimura, M., Takeda, N., Kono, R., Yamazaki, S., 1986. Evolution of enterovirus 70 in nature: All isolates were recently derived from a common ancestor. Archives of Virology 89, 1–14. https://doi.org/10.1007/BF01309875

Modiyinji, A.F., Joffret, M.-L., Nombot-Yazenguet, M.P. de M., Endengue Zanga, M.-C., Sadeuh-Mba, S., Njouom, R., Bessaud, M., 2025. Molecular characterization of enteroviruses circulating among pigs and goats in two Central African countries, Cameroon and the Central African Republic. Access Microbiol 7, 000886.v3. https://doi.org/10.1099/acmi.0.000886.v3

Mombo, I.M., Boundenga, L., Suquet, E., Ngoubangoye, B., Maganga, G.D., Leroy, E.M., Charpentier, M.J., Rougeron, V., 2021. Natural infection of free-ranging mandrills (Mandrillus sphinx) by enteroviruses and astroviruses in southern Gabon. Microb Pathog 150, 104659. https://doi.org/10.1016/j.micpath.2020.104659

Mombo, I.M., Lukashev, A.N., Bleicker, T., Brünink, S., Berthet, N., Maganga, G.D., Durand, P., Arnathau, C., Boundenga, L., Ngoubangoye, B., Boué, V., Liégeois, F., Ollomo, B., Prugnolle, F., Drexler, J.F., Drosten, C., Renaud, F., Rougeron, V., Leroy, E., 2017. African Non-Human Primates Host Diverse Enteroviruses. PLOS ONE 12, e0169067. https://doi.org/10.1371/journal.pone.0169067

Mombo, I.M., Ngoubangoye, B., Boundenga, L., Mebaley, T.G.N., Assoghe, A.A., Tsoumbou, T., Bisseye, C., Maganga, G., Leroy, E., 2023. Natural Non-Polio-Enteroviruses Detected in Captive Mandrills (Mandrillus sphinx) and Chimpanzees (Pan troglodytes troglodytes) Housed in a Primate Center in

672   Franceville, Gabon. Int J Primatol 44, 847–851. https://doi.org/10.1007/s10764-
673   023-00361-y
674 Muslin, C., Joffret, M.-L., Pelletier, I., Blondel, B., Delpeyroux, F., 2015. Evolution and
675   Emergence of Enteroviruses through Intra- and Inter-species Recombination:
676   Plasticity and Phenotypic Impact of Modular Genetic Exchanges in the 5'
677   Untranslated Region. PLoS Pathog 11, e1005266.
678   https://doi.org/10.1371/journal.ppat.1005266
679 Muslin, C., Kain, A.M., Bessaud, M., Blondel, B., Delpeyroux, F., 2019. Recombination in
680   Enteroviruses, a Multi-Step Modular Evolutionary Process. Viruses 11, 859.
681   https://doi.org/10.3390/v11090859
682 Nguyen, D.V., Harvala, H., Ngole, E.M., Delaporte, E., Woolhouse, M.E.J., Peeters, M.,
683   Simmonds, P., 2014. High Rates of Infection with Novel Enterovirus Variants in
684   Wild Populations of Mandrills and Other Old World Monkey Species. Journal of
685   Virology 88, 5967–5976. https://doi.org/10.1128/jvi.00088-14
686 Oberste, M. Steven, Maher, K., Kilpatrick, D.R., Flemister, M.R., Brown, B.A., Pallansch,
687   M.A., 1999. Typing of Human Enteroviruses by Partial Sequencing of VP1. Journal
688   of Clinical Microbiology 37, 1288–1293. https://doi.org/10.1128/jcm.37.5.1288-
689   1293.1999
690 Oberste, M. S., Maher, K., Kilpatrick, D.R., Pallansch, M.A., 1999. Molecular evolution of
691   the human enteroviruses: correlation of serotype with VP1 sequence and
692   application to picornavirus classification. J Virol 73, 1941–1948.
693   https://doi.org/10.1128/JVI.73.3.1941-1948.1999
694 Oberste, M.S., Maher, K., Michele, S.M., Belliot, G., Uddin, M., Pallansch, M.A., 2005.
695   Enteroviruses 76, 89, 90 and 91 represent a novel group within the species
696   Human enterovirus A. J Gen Virol 86, 445–451.
697   https://doi.org/10.1099/vir.0.80475-0
698 Oberste, M.S., Maher, K., Pallansch, M.A., 2007. Complete genome sequences for nine
699   simian enteroviruses. J Gen Virol 88, 3360–3372.
700   https://doi.org/10.1099/vir.0.83124-0
701 Oberste, M.S., Maher, K., Pallansch, M.A., 2002. Molecular Phylogeny and Proposed
702   Classification of the Simian Picornaviruses. Journal of Virology 76, 1244–1251.
703   https://doi.org/10.1128/jvi.76.3.1244-1251.2002
704 Olson, R.D., Assaf, R., Brettin, T., Conrad, N., Cucinell, C., Davis, J.J., Dempsey, D.M.,
705   Dickerman, A., Dietrich, E.M., Kenyon, R.W., Kuscuoglu, M., Lefkowitz, E.J., Lu, J.,
706   Machi, D., Macken, C., Mao, C., Niewiadomska, A., Nguyen, M., Olsen, G.J.,
707   Overbeek, J.C., Parrello, B., Parrello, V., Porter, J.S., Pusch, G.D., Shukla, M.,
708   Singh, I., Stewart, L., Tan, G., Thomas, C., VanOeffelen, M., Vonstein, V., Wallace,
709   Z.S., Warren, A.S., Wattam, A.R., Xia, F., Yoo, H., Zhang, Y., Zmasek, C.M.,
710   Scheuermann, R.H., Stevens, R.L., 2023. Introducing the Bacterial and Viral
711   Bioinformatics Resource Center (BV-BRC): a resource combining PATRIC,
712   IRD and ViPR. Nucleic Acids Res 51, D678–D689.
713   https://doi.org/10.1093/nar/gkac1003
714 Piovesan, A., Pelleri, M.C., Antonaros, F., Strippoli, P., Caracausi, M., Vitale, L., 2019. On
715   the length, weight and GC content of the human genome. BMC Res Notes 12,
716   106. https://doi.org/10.1186/s13104-019-4137-z
717 Romette, J.L., Prat, C.M., Gould, E.A., de Lamballerie, X., Charrel, R., Coutard, B., Fooks,
718   A.R., Bardsley, M., Carroll, M., Drosten, C., Drexler, J.F., Günther, S., Klempa, B.,

Pinschewer, D., Klimkait, T., Avsic-Zupanc, T., Capobianchi, M.R., Dicaro, A.,
Ippolito, G., Nitsche, A., Koopmans, M., Reusken, C., Gorbalenya, A., Raoul, H.,
Bourhy, H., Mettenleiter, T., Reiche, S., Batten, C., Sabeta, C., Paweska, J.T.,
Eropkin, M., Zverev, V., Hu, Z., Mac Cullough, S., Mirazimi, A., Pradel, F., Lieutaud,
P., 2018. The European Virus Archive goes global: A growing resource for
research. Antiviral Res 158, 127–134.
https://doi.org/10.1016/j.antiviral.2018.07.017
Sadeuh-Mba, S., Doté, J., Joffret, M.-L., Chesnais, M., Filhol, T., Gouandjika-Vasilache, I.,
Jouvenet, N., Bessaud, M., 2023. Animal enteroviruses: a glimpse of a wide
evolutionary landscape. Virologie (Montrouge) 27, 22–34.
https://doi.org/10.1684/vir.2023.1000
Sadeuh-Mba, S.A., Bessaud, M., Joffret, M.-L., Zanga, M.-C.E., Balanant, J., Ngole, E.M.,
Njouom, R., Reynes, J.-M., Delpeyroux, F., Rousset, D., 2014. Characterization of
Enteroviruses from Non-Human Primates in Cameroon Revealed Virus Types
Widespread in Humans along with Candidate New Types and Species. PLOS
Neglected Tropical Diseases 8, e3052.
https://doi.org/10.1371/journal.pntd.0003052
Sadeuh-Mba, S.A., Joffret, M.-L., Mazitchi, A., Endegue-Zanga, M.-C., Njouom, R.,
Delpeyroux, F., Gouandjika-Vasilache, I., Bessaud, M., 2019. Genetic and
phenotypic characterization of recently discovered enterovirus D type 111. PLoS
Negl Trop Dis 13, e0007797. https://doi.org/10.1371/journal.pntd.0007797
Santti, J., Hyypiä, T., Kinnunen, L., Salminen, M., 1999. Evidence of recombination
among enteroviruses. J Virol 73, 8741–8749.
https://doi.org/10.1128/JVI.73.10.8741-8749.1999
Tajima, F., Nei, M., 1984. Estimation of evolutionary distance between nucleotide
sequences. Mol Biol Evol 1, 269–285.
https://doi.org/10.1093/oxfordjournals.molbev.a040317
Tsala, T., 1956. Dictionnaire Ewondo - Français. Lyon.
Wang, X., Shao, S., Wang, H., Shen, Q., Yang, S., Zhang, W., 2016. An enterovirus from a
captive primate in China. Springerplus 5, 1281. https://doi.org/10.1186/s40064-
016-2966-y
Wright, F., 1990. The "effective number of codons" used in a gene. Gene 87, 23–29.
https://doi.org/10.1016/0378-1119(90)90491-9
Yamashita, T., Ito, M., Tsuzuki, H., Sakae, K., Minagawa, H., 2010. Molecular
identification of enteroviruses including two new types (EV-98 and EV-107)
isolated from Japanese travellers from Asian countries. J Gen Virol 91, 1063–
1066. https://doi.org/10.1099/vir.0.016014-0
Zeng, L., Chen, M., Wang, M., Zhu, L., Yan, J., Zhang, X., Xu, J., Zhang, S., 2022.
Enterovirus A Shows Unique Patterns of Codon Usage Bias in Conventional
Versus Unconventional Clade. Front. Cell. Infect. Microbiol. 12.
https://doi.org/10.3389/fcimb.2022.941325
760
761
762
763

Legends of figures and table

Figure 1. Genome organization of CHE20, Z057, RCMH6, and RCMH3 strains. (A) Schematic representation of the genomic organization of the four newly sequenced genomes. The uORF is shown in brown (Lulla et al., 2019), the P1 region in purple, the P2 region in green, and the P3 region in blue. **(B)** Alignment of nucleotide sequences of the potential uORF compared to the E7 Wallace strain (highlighted by brown boxes). The start codons of the putative uORF and the main ORF, as well as, the stop codons of the uORF are highlighted. **(C)** Alignment of the CRE sequence compared to the PV3 Leon strain (Cordey et al., 2008). The sequence alignment was performed using the CLC Main Workbench v24 software.

Figure 2. Phylogenetic analysis of complete VP1 sequences from non-human primate and representative human enteroviruses. The Neighbor-Joining tree was reconstructed from a full-length VP1 nucleotide sequence alignment. Branches are coloured according to EV species: EV-A (light green), EV-B (dark blue), EV-C (red), EV-D (dark green), EV-H (light blue), EV-J (pink), EV-L (purple), EV-M (brown) and EV-N (orange). Circles indicate NHP derived EV strains. The arrows indicate new genomes sequenced in our study (Figure 1). The tree is rooted with the PV1 Mahoney strain (V01148). Bootstrap values greater than 70% (based on 1,000 pseudoreplicates) are shown at the corresponding nodes. The sequence alignment was performed using the CLC Main Workbench v24, the phylogenetic tree was constructed with MEGA X and the visualization was generated with FigTree. The tree is drawn to scale, with branch lengths measured in the number of substitutions per site.

Figure 3. Phylogenetic analysis of enterovirus J species. The maximum-likelihood trees were constructed with all available full capsid (length of alignment 2571 nt) **(A)**, partial VP1

(length of alignment 754 nt) **(B)** and partial VP4/2 (length of alignment 429 nt) **(C)** sequences.

The trees were rooted with the EV-A124 OM22 strain. The symbol * indicates proposed new

type classifications. The colours of the strain names indicate their host origins: *Mandrillus*

(green), *Chlorocebus* (orange), *Cercocebus* (brown), *Papio* (blue), *Macaca* (purple) and *Pan*

(red). Bootstrap values greater than 70% (based on 500 pseudoreplicates) are shown at the

corresponding nodes. The sequence alignments were performed using the CLC Main

Workbench v24, the phylogenetic trees were constructed with MEGA X and the visualization

was generated with FigTree. The trees are drawn to scale, with branch lengths measured in the

number of substitutions per site.

Figure 4. VP1-based phylogenetic tree of enterovirus B species. The maximum-likelihood

tree was constructed with all available full NHP EV-B VP1 sequences. The tree is rooted with

the EV-B3 SSM strain. The colours of the strain names indicate their host origin: *Mandrillus*

(green), *Homo* (black), *Cercopithecus* (grey), and *Pan* (red). Bootstrap values greater than 70%

(based on 500 pseudoreplicates) are shown at the corresponding nodes. Percentages of

nucleotide (Nt) and amino acid (Aa) identities between EV-B114 strains are indicated. The

sequence alignment was performed using the CLC Main Workbench v24, and the phylogenetic

tree was constructed with MEGA X. The tree is drawn to scale, with branch lengths measured

in the number of substitutions per site.

Figure 5. VP1-based phylogenetic tree of the enterovirus M species. The maximum-

likelihood tree was constructed with all available full VP1 sequences. VP1 sequences derived

from complete genomes are in bold. The colours of the strain names indicate their host origin:

Mandrillus (green), *Cercocebus* (brown), *Macaca* (purple) and *Pan* (red). The tree is rooted

with the EV-J121 CHE20 strain. Bootstrap values greater than 70% (based on 500

pseudoreplicates) are shown at the corresponding nodes. The sequence alignment was

performed using the CLC Main Workbench v24, and the phylogenetic tree was constructed with

MEGA X. The tree is drawn to scale, with branch lengths measured in the number of

substitutions per site.

Figure 6. Phylogenetic analysis of non-structural genes derived from complete genomes of

NHP and representative human EVs. The maximum-likelihood trees were constructed from

the 2A2B, 2C, 3A3B, 3C and 3D regions of all complete NHP and representative human EV

genomes. Branches are coloured according to EV species: EV-A (light green), EV-B (dark blue),

EV-C (red), EV-D (dark green), EV-H (light blue), EV-J (pink), EV-L (purple), EV-M (brown)

and EV-N (orange). The colours of the strain names indicate their host origin: *Macaca* (purple),

Pan (red), *Papio* (light blue), *Chlorocebus* (orange), *Mandrillus* (green), *Cercocebus* (brown)

and *Homo* (black). The trees are rooted with the PV1 Mahoney strain (V01148). Bootstrap

values greater than 70% (based on 500 pseudoreplicates) are shown at the corresponding nodes.

The sequence alignments were performed using the CLC Main Workbench v24, the

phylogenetic trees were constructed with MEGA X, and the visualizations were generated with

FigTree. The trees are drawn to scale, with branch lengths measured in the number of

substitutions per site.

Figure 7. Distribution of GC content across enterovirus species. GC content distribution for

all EV species **(A)**, EV-A species **(B)**, EV-B species **(C)**, and EV-C species **(D)** are plotted. Blue

boxes indicate clusters of human derived EV while brown boxes indicate clusters of mainly

NHP derived EVs. When possible, three sequences per type were included in the analysis and

standard deviations are plotted. Abbreviations: Cult, cultivable; NC, non-cultivable; Resp, respiratory; Hs, *Homo sapiens*; NHP, non-human primate.

Figure 8. Dinucleotide ratio analysis of human and non-human primate enterovirus genomes. (A) All dinucleotide ratios are shown. **(B)** The CpG dinucleotide ratio is specifically highlighted for each EV species. Abbreviations: Cult, cultivable; NC, non-cultivable; Resp, respiratory; Hs, *Homo sapiens*; NHP, non-human primate.

Figure 9. Codon usage bias analysis of enterovirus species. (A) Effective number of codons (ENC) values for all NHP EVs and representative human EVs are plotted. When possible, three sequences per type were included in the analysis. **(B)** Relative synonymous codon usage (RSCU) values for all NHP EVs and representative human EVs were summarized in a cladogram. The three main groups separated by the cladogram are labeled A, B and C. Abbreviations: Hs, *Homo sapiens*; NHP, non-human primate.

Table 1. Summary of included studies and number of available enterovirus sequences. For each study, the first author, year of publication, and the number of sequences available are listed. Data were obtained from: (Ao et al., 2016; Han et al., 2021; Handley et al., 2012; Harvala et al., 2011, 2014; Kapusinszky et al., 2017; Li et al., 2020; Mombo et al., 2017, 2021, 2023; Negrey et al., 2022; Nguyen et al., 2014; Sadeuh-Mba et al., 2014; Wang et al., 2016; Zhao et al., 2019).

Figure 1. Genome organization of CHE20, Z057, RCMH6, and RCMH3 strains

Figure 2. Phylogenetic analysis of complete VP1 sequences from non-human primate and representative human enteroviruses

Figure 3. Phylogenetic analysis of enterovirus J species

Figure 4. VP1-based phylogenetic tree of enterovirus B species

Figure 5. VP1-based phylogenetic tree of the enterovirus M species

Figure 6. Phylogenetic analysis of non-structural genes derived from complete genomes of NHP and representative human EVs

Figure 7. Distribution of GC content across enterovirus species

Figure 8. Dinucleotide ratio analysis of human and non-human primate enterovirus genomes

Figure 9. Codon usage bias analysis of enterovirus species

First author	Year	5'UTR	VP4	VP4/2	VP2	VP1	3D	full genome	partial	Total
Harvala	2011	3		3		3	3			12
Handley	2012								3	3
Sadeuh-Mba	2014	36				27	23			86
Nguyen	2014			86		36		3		125
Harvala	2014			14		6				20
Ao	2016							1		1
Wang	2016							1		1
Mombo	2017				19	9		3		31
Kapusinszky	2017								22	22
Zhao	2019							4	1	5
George (unpublished)	2019								1	1
Li	2020							5	1	6
Mombo	2021					10				10
Han	2021							7		7
Negrey	2022								1	1
Mombo	2023				5					5
Total		39	0	103	19	91	26	24	29	336

Table 1. Summary of included studies and number of available enterovirus sequences

Supplementary Table 1. Predicted polyprotein cleavage sites of enterovirus Z057, CHE20, RCMH3, and RCMH6. Cleavage site predictions are based on multiple sequence alignment with the annotated poliovirus type 1 Mahoney reference strain (V01148.1). The analysis was performed using CLC Main Workbench v24 software.

Strains	VP4/VP2	VP2/VP3	VP3/VP1	VP1/2A	2A/2B	2B/2C	2C/3A	3A/3B	3B/3C	3C/3D
CHE20	LK/SP	TQ/GL	LQ/SP	TA/GA	EQ/GI	KQ/AD	FQ/GP	FQ/GA	VQ/GP	EQ/GE
Z057	LN/SP	PQ/GL	LQ/SP	TY/GA	EQ/GV	RQ/AD	FQ/GP	FQ/GA	VQ/GP	EQ/GE
RCMH3	LK/SP	TQ/GL	LQ/GD	NT/GA	EQ/GV	RQ/AD	FQ/GP	FQ/GA	VQ/GP	EQ/GE
RCMH6	LK/SP	AQ/GL	LQ/GD	TT/GA	EQ/GV	RQ/AD	FQ/GP	FQ/GA	VQ/GP	EQ/GE

Supplementary Figure 1. Phylogenetic analysis of complete VP1 sequences from non-human primate and representative human enteroviruses. This phylogenetic tree is the same as in the Figure 2, with the name of each EV type indicated. For NHP EV, the following nomenclature is used: Species-Type-Strain name-Accession number-Country-Year-Host species-Condition of life of host. For prototypic EVs, the following nomenclature is used: Species-Type-Strain name-Accession Number. Abbreviations: CAM, Cameroon; CHN, China; USA, United States of America; GAB, Gabon; NGR, Nigeria; Msph, *Mandrillus sphinx*; Mleu, *Mandrillus leucophaeus*; Ctor, *Cercocebus torquatus*; Caet, *Chlorocebus aethiops*; Ptro, *Pan troglodytes*; Ggor, *Gorilla gorilla*; Mmul, *Macaca mulatta*; W, wild; C, captive. The arrows indicate new genomes sequenced in our study (Figure 1). The tree is drawn to scale, with branch lengths measured in the number of substitutions per site.

		1	2	3	4	5	6	
EV-H DJ4141 KJ420733	1		100.00	99.88	99.88	85.6	84.09	
EV-H DJ4144 KJ420736	2	100.00		99.88	99.88	85.6	84.09	
EV-H DJ4143 KJ420735	3	99.65	99.65		100.00	85.71	84.2	Percentage of
EV-H DJ4142 KJ420734	4	99.65	99.65	100.00		85.71	84.2	nucleotide identity
EV-H1 SV4-1715 UWB AF326759	5	97.56	97.56	97.91	97.91		84.44	
EV-H1 A-2 plaque AF201894	6	95.82	95.82	96.17	96.17	96.86		

Percentage of amino acid identity

Supplementary Figure 2. Type characterization of the enterovirus H species. Nucleotide and amino acid identity matrices of EV-H complete VP1 sequences were calculated using CLC Main Workbench v24.

A

Group	Strain	#	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
J121	EV-J12J CJIE20 KF541639	1		60.89	62.39	61.80	62.27	61.29	60.48	63.33	57.35	55.87	50.45	50.23	50.17	50.63	49.71	49.60
J103	EV-J103 POo1 FJ007373	2	60.89		82.23	75.29	76.81	76.25	71.61	71.71	59.43	57.98	54.98	52.93	50.68	50.68	51.03	51.37
	EV-J103 NOLA1 MZ312496	3	62.39	82.23		75.41	76.31	77.33	71.89	73.89	59.91	58.70	54.57	52.70	51.94	51.71	52.05	52.05
	EV-J103 NOLA3 MZ312494	4	61.80	75.29	75.41		77.82	77.33	72.64	72.55	60.57	60.51	56.00	52.70	52.05	51.71	51.48	51.71
	EV-J103 cg3697 KT961657	5	62.27	76.81	76.31	77.82		87.02	74.48	73.17	61.63	59.84	56.18	53.39	52.85	52.97	53.54	53.77
	EV-J103 cg8227 KT961656	6	61.29	76.25	77.33	77.33	87.02		72.31	73.95	60.30	59.31	55.42	52.82	52.51	52.17	53.88	54.00
J123*	EV-J123 Sev-nJ KT581857	7	60.48	71.61	71.89	72.64	74.48	72.31		82.09	62.53	59.93	57.32	51.78	51.48	51.94	52.17	52.40
	EV-J123 CHN-BJ-LXL1 OK397296	8	63.33	71.71	73.89	72.55	73.17	73.95	82.09		62.87	60.51	56.86	53.86	52.17	52.51	51.71	51.94
J108	EV-J108 N125 NC010384	9	57.35	59.43	59.91	60.57	61.63	60.30	62.53	62.87		75.44	54.84	50.58	50.58	50.35	51.04	51.38
	EV-J108 N203 NC013695	10	55.87	57.98	58.70	60.51	59.84	59.31	59.93	60.51	75.44		54.61	50.12	49.08	49.54	48.96	49.42
J122	EV-J122 SV6 AF326766.2	11	50.45	54.98	54.57	56.00	56.18	55.42	57.32	56.86	54.84	54.61		52.63	53.97	53.29	52.61	52.72
N2*	EV-N2 CHN-BJ-2J MN865119	12	50.23	52.93	52.70	52.70	53.39	52.82	51.78	53.86	50.58	50.12	52.63		72.24	71.78	72.71	72.94
N1*	EV-N1 CHN-BJ-1J MN427526	13	50.17	50.68	51.94	52.05	52.85	52.51	51.48	52.17	50.58	49.08	53.97	72.24		98.12	92.37	92.25
	EV-N1 CHN-BJ-8J MN427527	14	50.63	50.68	51.71	51.71	52.97	52.17	51.94	52.51	50.35	49.54	53.29	71.78	98.12		91.20	91.08
	EV-N1 CHN-BJ-10J MN865120	15	49.71	51.03	52.05	51.48	53.54	53.88	52.17	51.71	51.04	48.96	52.61	72.71	92.37	91.20		99.18
	EV-N1 CHN-BJ-9J MN427528	16	49.60	51.37	52.05	51.71	53.77	54.00	52.40	51.94	51.38	49.42	52.72	72.94	92.25	91.08	99.18	

B

Group	Strain	#	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
J103	EV-J103 POo1 FJ007373	1		92.31	87.41	89.16	85.31	82.11	83.16	66.10	60.14	62.20	52.90	47.26	46.92	47.60	47.26	46.58
	EV-J103 NOLA1 MZ312496	2	92.31		87.06	88.81	85.66	81.82	83.92	65.87	60.82	62.54	52.22	48.63	48.29	48.97	48.97	47.95
	EV-J103 cg3697 KT961657	3	87.41	87.06		94.76	86.36	84.27	84.62	68.26	61.86	61.86	51.54	48.63	48.63	48.97	48.97	49.32
	EV-J103 cg8227 KT961656	4	89.16	88.81	94.76		87.41	84.97	85.66	66.89	62.89	62.89	50.85	48.97	48.97	49.66	49.66	48.97
	EV-J103 NOLA3 MZ312494	5	85.31	85.66	86.36	87.41		82.17	82.87	65.53	60.82	61.86	50.85	48.63	48.63	47.95	47.95	47.60
J123*	EV-J123 Sev-nJ KT581857	6	82.11	81.82	84.27	84.97	82.17		92.63	67.47	61.86	62.20	52.90	48.63	48.29	48.97	48.97	46.58
	EV-J123 CHN-BJ-LXL1 OK397296	7	83.16	83.92	84.62	85.66	82.87	92.63		68.15	62.89	63.92	54.27	48.97	48.63	48.63	48.63	47.26
J121	EV-J12J CJIE20 KF541639	8	66.10	65.87	68.26	66.89	65.53	67.47	68.15		55.07	55.07	47.99	46.28	46.28	46.28	46.28	45.27
J108	EV-J108 N125 NC010384	9	60.14	60.82	61.86	61.51	60.82	61.86	62.89	55.07		89.72	47.78	44.48	44.48	43.79	44.14	44.83
	EV-J108 N203 NC013695	10	62.20	62.54	61.86	62.89	61.86	62.20	63.92	55.07	89.72		51.19	44.83	44.83	44.48	44.48	44.83
J122	EV-J122 SV6 AF326766.2	11	52.90	52.22	51.54	50.85	50.85	52.90	54.27	47.99	47.78	51.19		46.78	46.44	46.44	46.44	46.10
N1*	EV-N1 CHN-BJ-1J MN427526	12	47.26	48.63	48.63	48.97	48.63	48.63	48.97	46.28	44.48	44.83	46.78		99.65	94.37	95.07	79.58
	EV-N1 CHN-BJ-8J MN427527	13	46.92	48.29	48.63	48.97	48.63	48.29	48.63	46.28	44.48	44.83	46.44	99.65		94.01	94.72	79.58
	EV-N1 CHN-BJ-10J MN865120	14	47.60	48.97	48.97	49.66	47.95	48.97	48.63	46.28	43.79	44.48	46.44	94.37	94.01		98.59	79.93
	EV-N1 CHN-BJ-9J MN427528	15	47.26	48.97	48.97	49.66	47.95	48.97	48.63	46.28	44.14	44.48	46.44	95.07	94.72	98.59		80.99
N2*	EV-N2 CHN-BJ-2J MN865119	16	46.58	47.95	49.32	48.97	47.60	46.58	47.26	45.27	44.83	44.83	46.10	79.58	79.58	79.93	80.99	

Supplementary Figure 3. Type characterization of enterovirus J strains based on complete VP1 sequences. Nucleotide **(A)** and amino acid **(B)** identity matrices of VP1 sequences are shown. Translation of nucleotide sequences, multiple sequence alignments, and identity matrices were performed using CLC Main Workbench v24. Putative novel type assignments are indicated by an asterisk (*).

Supplementary Figure 4. Type characterization of enterovirus J strains based on partial VP1 sequences. Nucleotide **(A)** and amino acid **(B)** identity matrices of partial VP1 sequences are shown. Translation of nucleotide sequences, multiple sequence alignments, and identity matrices were performed using CLC Main Workbench 24. Putative novel type assignments are indicated by an asterisk (*).

A — ORF

		1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
J121	EV-J121 CHE20 KF541639	1		78.44	83.67	82.70	83.03	82.34	82.34	83.71	80.43	80.57	71.82	71.53	71.53	70.85	71.53
J122	EV-J122 SV6 AF326766.2	2	78.44		83.84	83.24	83.84	83.24	83.15	84.48	75.26	74.93	71.93	72.25	72.30	71.84	72.34
J103	EV-J103 cg3697 KT961657	3	83.67	83.84		94.59	95.23	94.13	91.42	92.85	79.62	79.39	72.65	72.51	72.51	72.28	72.74
	EV-J103 POo1 FJ007373	4	82.70	83.24	94.59		96.19	93.21	90.09	91.70	78.88	79.07	71.83	71.83	71.78	71.37	71.87
	EV-J103 NOLA1 MZ312496	5	83.03	83.84	95.23	96.19		94.36	91.15	92.85	79.30	79.48	72.28	72.15	72.05	71.74	72.28
	EV-J103 NOLA3 MZ312494	6	82.34	83.24	94.13	93.21	94.36		90.27	91.38	78.92	78.66	72.15	72.24	72.05	71.51	72.05
J123*	EV-J123 Sev-nJ KT581857	7	82.34	83.15	91.42	90.09	91.15	90.27		95.69	79.06	78.39	71.55	71.83	71.69	71.14	71.74
	EV-J123 CHN-BJ-LXL1 OK397296	8	83.71	84.48	92.85	91.70	92.85	91.38	95.69		79.84	79.43	72.01	72.28	72.15	71.51	72.15
J108	EV-J108 N125 NC010384	9	80.43	75.26	79.62	78.88	79.30	78.92	79.06	79.84		92.73	70.85	70.65	70.52	69.79	70.70
	EV-J108 N203 NC013695	10	80.57	74.93	79.39	79.07	79.48	78.66	78.39	79.43	92.73		70.32	70.14	70.18	69.41	70.05
N2*	EV-N2 CHN-BJ-2J MN865119	11	71.82	71.93	72.65	71.83	72.28	72.15	71.55	72.01	70.85	70.32		92.91	92.72	92.31	93.23
N1*	EV-N1 CHN-BJ-1J MN427526	12	71.53	72.25	72.51	71.83	72.15	72.24	71.83	72.28	70.65	70.14	92.91		99.40	96.45	97.74
	EV-N1 CHN-BJ-8J MN427527	13	71.53	72.30	72.51	71.78	72.05	72.05	71.69	72.15	70.52	70.18	92.72	99.40		96.17	97.46
	EV-N1 CHN-BJ-10J MN865120	14	70.85	71.84	72.28	71.37	71.74	71.51	71.14	71.51	69.79	69.41	92.31	96.45	96.17		98.43
	EV-N1 CHN-BJ-9J MN427528	15	71.53	72.34	72.74	71.87	72.28	72.05	71.74	72.15	70.70	70.05	93.23	97.74	97.46	98.43	

B — 2C+3CD

		1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
J121	EV-J121 CHE20 KF541639	1		88.60	90.04	90.45	90.14	90.14	90.97	91.17	89.85	89.74	84.50	84.19	84.19	84.39	82.85
J123*	EV-J123 Sev-nJ KT581857	2	88.60		93.12	94.15	94.35	94.25	94.46	96.41	84.92	84.62	82.34	82.24	82.24	82.34	81.01
J103	EV-J103 POo1 FJ007373	3	90.04	93.12		96.92	97.43	97.33	94.97	95.28	86.46	86.46	83.37	83.26	83.26	83.37	82.34
	EV-J103 cg3697 KT961657	4	90.45	94.15	96.92		97.95	97.95	96.00	96.30	86.56	86.46	83.26	83.26	83.16	83.37	82.34
	EV-J103 NOLA3 MZ312494	5	90.14	94.35	97.43	97.95		99.28	96.71	96.51	86.67	86.77	83.26	83.16	83.16	83.26	82.03
	EV-J103 NOLA1 MZ312496	6	90.14	94.25	97.33	97.95	99.28		96.51	96.41	86.56	86.67	83.47	83.26	83.26	83.37	82.14
J122	EV-J122 SV6 AF326766.2	7	90.97	94.46	94.97	96.00	96.71	96.51		96.92	86.97	86.46	82.96	82.85	82.85	82.96	81.83
J123*	EV-J123 CHN-BJ-LXL1 OK397296	8	91.17	96.41	95.28	96.30	96.51	96.41	96.92		86.67	86.67	83.16	83.16	83.06	83.16	81.72
J108	EV-J108 N125 NC010384	9	89.85	84.92	86.46	86.56	86.67	86.56	86.97	86.67		94.97	83.18	83.28	83.28	83.49	81.74
	EV-J108 N203 NC013695	10	89.74	84.62	86.36	86.46	86.77	86.67	86.46	86.67	94.97		82.15	81.85	81.85	82.05	80.72
N2*	EV-N2 CHN-BJ-2J MN865119	11	84.50	82.34	83.37	83.26	83.26	83.47	82.96	83.16	83.18	82.15		98.56	98.56	98.77	96.82
N1*	EV-N1 CHN-BJ-1J MN427526	12	84.19	82.24	83.26	83.26	83.16	83.26	82.85	83.16	83.28	81.85	98.56		99.79	99.38	96.71
	EV-N1 CHN-BJ-8J MN427527	13	84.19	82.24	83.26	83.16	83.16	83.26	82.85	83.06	83.28	81.85	98.56	99.79		99.38	96.71
	EV-N1 CHN-BJ-9J MN427528	14	84.39	82.34	83.37	83.37	83.26	83.37	82.96	83.16	83.49	82.05	98.77	99.38	99.38		97.33
	EV-N1 CHN-BJ-10J MN865120	15	82.85	81.01	82.34	82.34	82.03	82.14	81.83	81.72	81.74	80.72	96.82	96.71	96.71	97.33	

C — Capsid

		1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
J121	EV-J121 CHE20 KF541639	1		74.74	75.06	74.74	73.42	73.92	72.05	68.30	69.66	61.42	55.79	55.97	56.09	55.62	55.74
J123*	EV-J123 Sev-nJ KT581857	2	74.74		95.05	87.97	86.44	87.26	85.38	71.20	71.40	65.65	57.63	58.97	58.86	58.51	58.62
	EV-J123 CHN-BJ-LXL1 OK397296	3	75.06	95.05		88.56	87.60	88.68	85.51	70.85	71.74	65.89	57.63	58.97	58.86	58.39	58.51
J103	EV-J103 cg3697 KT961657	4	74.74	87.97	88.56		90.80	91.39	89.28	70.30	70.96	64.95	58.44	58.74	58.86	58.97	59.09
	EV-J103 POo1 FJ007373	5	73.42	86.44	87.60	90.80		94.46	87.51	69.45	70.58	65.77	57.51	58.39	58.39	58.16	58.28
	EV-J103 NOLA1 MZ312496	6	73.92	87.26	88.68	91.39	94.46		87.40	69.72	69.72	65.06	57.86	58.51	58.39	58.51	58.62
	EV-J103 NOLA3 MZ312494	7	72.05	85.38	85.51	89.28	87.51	87.40		68.99	69.53	63.58	57.97	59.21	58.86	58.39	58.51
J108	EV-J108 N125 NC010384	8	68.30	71.20	70.85	70.30	69.45	69.72	68.99		91.29	59.56	56.29	55.71	55.48	55.36	55.71
	EV-J108 N203 NC013695	9	69.66	71.40	71.74	70.96	70.58	69.72	69.53	91.29		60.21	56.83	56.94	56.94	56.36	56.48
J122	EV-J122 SV6 AF326766.2	10	61.42	65.65	65.89	64.95	65.77	65.06	63.58	59.56	60.21		57.33	59.14	59.25	59.14	59.25
N2*	EV-N2 CHN-BJ-2J MN865119	11	55.79	57.63	57.63	58.44	57.51	57.86	57.97	56.29	56.83	57.33		84.95	84.59	85.07	85.30
N1*	EV-N1 CHN-BJ-1J MN427526	12	55.97	58.97	58.97	58.74	58.39	58.51	59.21	55.71	56.94	59.14	84.95		99.04	95.33	95.57
	EV-N1 CHN-BJ-8J MN427527	13	56.09	58.86	58.86	58.86	58.39	58.39	58.86	55.48	56.94	59.25	84.59	99.04		94.73	94.97
	EV-N1 CHN-BJ-10J MN865120	14	55.62	58.51	58.39	58.97	58.16	58.51	58.39	55.36	56.36	59.14	85.07	95.33	94.73		99.28
	EV-N1 CHN-BJ-9J MN427528	15	55.74	58.62	58.51	59.09	58.28	58.62	58.51	55.71	56.48	59.25	85.30	95.57	94.97	99.28	

Supplementary Figure 5. Species characterization of enterovirus J strains. Amino acid identity matrices of **(A)** the complete ORF, **(B)** the 2C+3CD and **(C)** the capsid sequences are shown. Translation of nucleotide sequences, multiple sequence alignments, and identity matrices were performed using CLC Main Workbench v24. Putative novel type assignments are indicated by an asterisk (*).

A ORF

	1	2	3	4	5	6	7	8	9	10	11	12	13	14
EV-M3 CPML3961 KJ420748 (1)		95,02	94,06	89,69	88,50	86,67	77,78	73,85	71,13	69,44	76,19	72,11	71,49	64,97
EV-M1 RCMH6 (2)	95,02		94,56	89,78	88,36	86,94	78,14	74,16	71,40	70,30	76,46	72,60	71,94	65,20
EV-M2 RCMH3 (3)	94,06	94,56		90,69	89,18	86,81	78,28	74,12	71,22	69,94	76,73	72,46	71,71	65,25
EV-M4 CPML8109 KJ420749 (4)	89,69	89,78	90,69		93,17	87,06	78,40	74,23	72,14	70,13	76,80	72,75	71,63	65,20
EV-M5 GR2815 KJ420747 (5)	88,50	88,36	89,18	93,17		85,69	77,54	73,96	71,05	70,00	76,40	72,60	71,36	64,61
EV-M6 NGR_NHP_1 MN265403 (6)	86,67	86,94	86,81	87,06	85,69		77,91	74,60	71,03	69,71	76,01	72,47	71,55	63,85
EV-B114 Z057 (7)	77,78	78,14	78,28	78,40	77,54	77,91		91,48	85,50	81,98	87,92	67,65	67,63	58,76
EV-B114 B165 AF326751 (8)	73,85	74,16	74,12	74,23	73,96	74,60	91,48		81,20	82,07	87,65	65,52	65,60	57,90
EV-B113 GAB653 KJ701249 (9)	71,13	71,40	71,22	72,14	71,05	71,03	85,50	81,20		85,24	80,22	61,09	60,81	53,66
EV-B110 K-UT-18-Q3 MW876511* (10)	69,44	70,30	69,94	70,13	70,00	69,71	81,98	82,07	85,24		87,21	61,50	61,49	53,48
EV-B112 GAB130 KJ418244 (11)	76,19	76,46	76,73	76,80	76,40	76,01	87,92	87,65	80,22	87,21		66,74	66,50	57,90
EV-J121 CHE20 (12)	72,11	72,60	72,46	72,75	72,60	72,47	67,65	65,52	61,09	61,50	66,74		80,64	63,29
EV-J08 N125 NC010384 (13)	71,49	71,94	71,71	71,63	71,36	71,55	67,63	65,60	60,81	61,49	66,50	80,64		61,42
EV-A122 M19 AF326754 (14)	64,97	65,20	65,25	65,20	64,61	63,85	58,76	57,90	53,66	53,48	57,90	63,29	61,42	

B 2C+3CD

	1	2	3	4	5	6	7	8	9	10	11	12	13	14
EV-M4 CPML8109 KJ420749 (1)		99,69	99,59	99,28	98,77	98,46	98,56	97,85	94,15	84,51	81,95	83,50	75,21	65,20
EV-M5 GR2815 KJ420747 (2)	99,69		99,38	99,18	98,87	98,26	98,36	97,64	93,95	84,31	81,95	83,30	75,02	65,20
EV-M2 RCMH3 (3)	99,59	99,38		98,97	98,46	98,26	98,36	97,64	94,05	84,31	81,74	83,30	75,02	65,40
EV-M1 RCMH6 (4)	99,28	99,18	98,97		98,56	98,05	98,26	97,54	94,15	84,41	81,64	83,09	75,12	65,10
EV-M3 CPML3961 KJ420748 (5)	98,77	98,87	98,46	98,56		97,54	98,15	97,03	93,54	83,79	81,85	82,89	74,75	64,99
EV-B114 Z057 (6)	98,46	98,26	98,26	98,05	97,54		97,54	97,33	93,54	84,31	81,74	83,20	75,02	64,99
EV-B112 GAB130 KJ418244 (7)	98,56	98,36	98,36	98,26	98,15	97,54		97,64	93,74	84,38	81,91	82,99	75,02	64,69
EV-M6 NGR_NHP_1 MN265403 (8)	97,85	97,64	97,64	97,54	97,03	97,33	97,64		94,15	84,31	80,92	83,61	73,27	65,40
EV-B114 B165 AF326751 (9)	94,15	93,95	94,05	94,15	93,54	93,54	93,74	94,15		80,41	77,44	80,94	73,27	65,40
EV-B110 K-UT-18-Q3 MW876511* (10)	84,51	84,31	84,31	84,41	83,79	84,31	84,38	84,31	80,41		94,18	71,00	63,96	55,27
EV-B113 GAB653 KJ701249 (11)	81,95	81,95	81,74	81,64	81,85	81,74	81,91	80,92	77,44	94,18		67,42	61,01	53,22
EV-J121 CHE20 (12)	83,50	83,30	83,30	83,09	82,89	83,20	82,99	83,61	80,94	71,00	67,42		80,81	65,51
EV-J08 N125 NC010384 (13)	75,21	75,02	75,02	75,12	74,75	75,02	75,02	73,27	73,27	63,96	61,01	80,81		57,42
EV-A122 M19 AF326754 (14)	65,20	65,20	65,40	65,10	64,99	64,99	64,69	65,40	65,40	55,27	53,22	65,51	57,42	

C Capsid

	1	2	3	4	5	6	7	8	9	10	11	12	13	14
EV-M3 CPML3961 KJ420748 (1)		89,82	87,49	75,93	74,18	71,84	63,51	57,21	56,40	47,56	47,79	48,01	47,73	47,84
EV-M1 RCMH6 (2)	89,82		87,94	75,35	73,60	71,96	64,51	58,78	57,32	48,07	48,30	49,20	48,92	48,35
EV-M2 RCMH3 (3)	87,49	87,94		77,34	75,35	71,61	64,05	57,97	56,86	47,84	47,84	48,30	48,58	49,03
EV-M4 CPML8109 KJ420749 (4)	75,93	75,35	77,34		85,14	71,94	63,93	58,56	56,66	48,63	48,97	50,68	48,80	48,69
EV-M5 GR2815 KJ420747 (5)	74,18	73,60	75,35	85,14		69,27	62,77	59,21	56,89	48,23	48,57	49,71	49,31	48,05
EV-M6 NGR_NHP_1 MN265403 (6)	71,84	71,96	71,61	71,94	69,27		60,07	56,89	55,79	49,72	49,72	50,28	49,37	48,92
EV-A122 M19 AF326754 (7)	63,51	64,51	64,05	63,93	62,77	60,07		59,44	56,78	47,61	47,84	48,41	47,55	47,78
EV-J121 CHE20 (8)	57,21	58,78	57,97	58,56	59,21	56,89	59,44		69,04	47,43	47,89	47,89	48,28	47,60
EV-J08 N125 NC010384 (9)	56,40	57,32	56,86	56,66	56,89	55,79	56,78	69,04		47,61	47,49	47,38	47,43	47,21
EV-B114 Z057 (10)	47,56	48,07	47,84	48,63	48,23	49,72	47,61	47,43	47,61		94,35	85,90	79,55	79,55
EV-B114 B165 AF326751 (11)	47,79	48,30	47,84	48,97	48,57	49,72	47,84	47,89	47,49	94,35		86,02	80,14	78,85
EV-B113 GAB653 KJ701249 (12)	48,01	49,20	48,30	50,68	49,71	50,28	48,41	47,89	47,38	85,90	86,02		78,26	77,32
EV-B112 GAB130 KJ418244 (13)	47,73	48,92	48,58	48,80	49,31	49,37	47,55	48,28	47,43	79,55	80,14	78,26		88,92
EV-B110 K-UT-18-Q3 MW876511* (14)	47,84	48,35	49,03	48,69	48,05	48,92	47,78	47,60	47,21	79,55	78,85	77,32	88,92	

Supplementary Figure 6. Species characterization of enterovirus M strains. Amino acid identity matrices of **(A)** the complete ORF, **(B)** the 2C+3CD and **(C)** the capsid sequences are shown. Translation of nucleotide sequences, multiple sequence alignments, and identity matrices were performed using CLC Main Workbench v24. The asterisk (*) indicates the sequence where the last 399 nt are missing in the 3D gene.

		1	2	3	4	5	6
EV-M2 RCMH3	1		72.59	67.59	68.4	65.63	59.01
EV-M1 RCMH6	2	86.81		64.59	64.13	64.13	57.77
EV-M3 CPML3961 KJ420748	3	84.72	87.5		74.45	73.36	58.22
EV-M4 CPML8109 KJ420749	4	72.66	70.24	71.97		73.82	62.5
EV-M5 GR2815 KJ420747	5	69.2	69.9	69.9	80.84		60.02
EV-M6 NGR_NHP_1 MN265403	6	60.81	60.47	64.19	59.46	58.45	

Percentage of nucleotide identity

Percentage of amino acid identity

Supplementary Figure 7. Type characterization of enterovirus M strains based on complete VP1 sequences. Nucleotide and amino acid identity matrices of EV-M VP1 sequences derived from complete genomes were calculated using CLC Main Workbench v24.

Supplementary Figure 8. Type characterization of enterovirus M strains based on all available VP1 sequences. Nucleotide **(A)** and amino acid **(B)** identity matrices of VP1 sequences are shown. Translation of nucleotide sequences, multiple sequence alignments, and identity matrices were performed using CLC Main Workbench v24.

Supplementary Figure 9. Phylogenetic analysis of enterovirus M strains based on partial VP1 and VP2 sequences. Maximum likelihood phylogenetic trees based on **(A)** VP1 (average length: 354 nucleotides) and **(B)** VP2 (average length: 320 nucleotides) partial sequences are shown. Sequence alignments were performed using CLC Main Workbench v24 and phylogenetic trees were constructed with MEGA X (version 10.2.6). The trees are drawn to scale, with branch lengths measured in the number of substitutions per site. Complete genomes are in bold and symbols indicate references of published sequences: ^, Mombo et al. 2023; *, Mombo et al. 2017; °, Mombo et al. 2021.

Supplementary Figure 10. ENC–GC3 plots to determine bias mediated by GC3 position. The blue curve indicates the expected ENC values if GC3 composition alone drives codon usage bias. **(A)** EV-A clade 1 species. **(B)** EV-A clade 2 species. **(C)** EV-A clade 3 species. **(D)** EV-A clade 4 species. **(E)** Human EV-B. **(F)** NHP EV-B.

Supplementary Figure 10. ENC–GC3 plots to determine bias mediated by GC3 position. The blue curve indicates the expected ENC values if GC3 composition alone drives codon usage bias. **(G)** Non-cultivable EV-C. **(H)** Respiratory EV-C. **(I)** Cultivable EV-C. **(J)** EV-D. **(K)** EV-L. **(L)** EV-H. **(M)** EV-M. **(N)** EV-N. Resp, Respiratory; NC, Non-cultivable; Cult: cultivable.