Information-Theoretic Origins of Metabolic Scaling

Andrea Tabi

[1]University of Amsterdam, The Netherlands

KEYWORDS: universality | information neutrality | Kullback-Leibler divergence | coarse-graining | stochastic processes

Abstract

Metabolic rate scales with body size, however its universality remains debated and
unresolved. We show that such universal scaling may arise from information neutrality
in stochastic cell dynamics. Using a stochastic ontogenetic growth model of cellular
dynamics, we identify an optimal microscopic noise structure where organism level
metabolic fluctuations are least sensitive to the underlying microscopic cellular noise
and have maximal dependence on organism size. At this point, the macroscopic scaling
exponent collapses to a universal value across species size close to Kleiber's law. These
results reveal a noncritical RG-like behavior, suggesting that universality emerges here
from an information-theoretic optimum of stochastic metabolic fluctuations.

1   Introduction

Power-law distribution is prevalent in natural complex systems. This distribution is one of the
signatures of phase transitions which emerge when a control parameter is tuned to a critical
threshold [1]. This underlies the concept of self-organized criticality, i.e. when natural complex
systems evolve towards a critical threshold in order to achieve maximal information processing [2].
Since "life at the edge of chaos" argument has gained traction in biology over the past decades,
many biological processes have been observed to reside close to phase transitions [3, 4]. However,
various mechanisms can generate power laws, such as aggregation [5], preferential attachment [6],
or stochastic processes [7, 8] without criticality [9]. Thus identifying the generative process remains
crucial for understanding the biological significance and interpretation of such scaling laws.

In ecology, metabolic scaling has been a long-standing candidate for a unifying principle across
scales [10], which describes the sublinear scaling between organisms' body size and metabolic rate
rooted in an empirical observation called the Kleiber's law [11]. Since body size being a master trait,
metabolic scaling belongs to a broader class of biological allometries [12] with life-history traits,
population growth rates, and energy fluxes [13] to name a few. This suggests that living systems

must obey internal constraints as a consequence of historical contingencies and functional and structural rules [14, 15]. The universality principle of this relationship has been up for debate for a long time due to largely varying observations [16, 17, 18], however the most popular explanations include noncritical phenomena such as fractal-like structures of distribution networks [19], surface-to-volume ratio of the organisms [20, 21] or life-history optimisation [22]. In biological contexts, such scale invariance implies efficient information propagation across biological hierarchy under noisy environment [23]. Coarse-grained variables such as metabolic rate thus one of the most informative macroscopic descriptors that captures the system behavior as a whole.

A recent work has shown that metabolic fluctuations can successfully reproduce the observed variability of the metabolic scaling exponents across and within taxa [8]. In ecological systems, demographic noise arises from discrete birth-death processes, and is a fundamental source of intrinsic stochasticity [24, 25]. It has been shown to generate macroscopic population fluctuations even without environmental forcings [26]. Thus, demographic stochasticity constitutes size-dependent fluctuations that have been shown to modulate metabolic scaling relations [8]. This raises a question under what conditions do macroscopic patterns, in this case the allometric exponents, become universal given the underlying microscopic noise structure. In this study, we present an information-theoretic model to show that optimal cellular fluctuations can give rise to universal metabolic scaling across species. The universality presented here is rooted in the information neutrality between stochastic cellular processes and organism-level metabolic fluctuations.

2 Methods

2.1 Stochastic metabolic model of cellular bursts

The ontogenetic growth is modeled at the cellular level. First, we model cell dynamics which is assumed to be a bursty stochastic process. The cell dynamics regulates the metabolic rate and the body mass of the organism. Cell dynamics is modeled as a stochastic jump process where the jumps J(t) are approximated as a symmetric exponential jumps and sampled from a Laplace distribution [27]. The cell population evolves according to a stochastic logistic model [8]:

$$\frac{dN_c}{dt} = aN_c\left(1 - \frac{N_c}{N}\right) + J(t), \tag{1}$$

where a is the cellular growth rate, N is the total number of cells at maturity, $J(t) \sim$ Laplace$(0, b)$, is the jump process and the magnitude throughout the life cycle can be written as:

$$b = N_c^{\alpha/2}\left[\phi + (1 - \phi)\left(1 - \frac{N_c}{N}\right)^{\theta}\right]. \tag{2}$$

where α is the correlation exponent, ϕ is the fluctuation coefficient. It is assumed that the

magnitude of fluctuations are higher in the growth phase pf the organism compared to the adult phase, which gradually decays over time. This decay is modulated by θ. The magnitude of the jump process is proportional to organism size and at the adult stage it simplifies to:

$$b = \phi N^{\alpha/2} \tag{3}$$

The exponent $\alpha/2$ derives from the assumption that the variance of the Laplace distribution is proportional to the system size, i.e. $Var(J) = 2b^2$, where $Var(J) \propto N_c^{\alpha}$. This allows for a tunable correlation strength of cell bursts, where $\alpha = 1$ represents independent cells, $\alpha = 2$ represents fully correlated cells and $1 < \alpha < 2$ represents partially correlated cell population. Values of $\alpha > 1$ lead to sublinear allometric scaling, whereas $\alpha < 1$ can result in superlinear allometric scaling exponents β (Fig. S1).

Then, we estimate organism-level metabolism as the sum of maintenance costs of each individual cell (E_M), the growth costs (E_G) and an overhead cost of stochastic cellular dynamics (H) [8].:

$$B(t) = B_M(t) + B_G(t) + H(t). \tag{4}$$

The maintenance component is proportional to the number of cells:

$$B_M(t) = N_c(t)B_c, \tag{5}$$

where B_c is the maintenance power expenditure per cell ($B_c = 1.08e - 7$ W [28]). The growth component accounts for the energetic cost of creating new cells:

$$B_G = \begin{cases} \dfrac{E_c}{\eta} \dfrac{dN_c}{dt}, & \text{if } \dfrac{dN_c}{dt} \geq 0, \\[3mm] 0, & \text{if } \dfrac{dN_c}{dt} < 0 \end{cases} \tag{6}$$

where E_c is the energy required for cell birth ($E_c = 3.7e - 7J$ [29]), and η is the growth efficiency ($\eta = 0.7$). The stochastic inefficiency component H represents the overhead costs of cell bursts [30, 31] and is proportional to the magnitude of the jump term:

$$H = \begin{cases} (1 - \eta)E_c\dfrac{dN_c}{dt} + \gamma\left|J(t)\right|, & \text{if } \dfrac{dN_c}{dt} \geq 0, \\[3mm] \kappa E_c\left|\dfrac{dN_c}{dt}\right| + \gamma\left|J(t)\right|, & \text{if } \dfrac{dN_c}{dt} < 0 \end{cases} \tag{7}$$

where γ is a stochastic overhead cost that converts the stochastic cellular burst into energy dissipation ($\gamma = 1$ J/cell) and κ is the apoptosis cost set to 0.3. The total body mass evolves

over time as the net energy available for biomass production, after subtracting maintenance and stochastic inefficiency [32]:

$$\frac{dm}{dt} = \frac{m_c}{E_c}(B - H) - \frac{mB_c}{E_c}. \tag{8}$$

where m_c is the average mass of a cell ($m_c = 1e - 12kg$ [33]). This final form describes the dynamics of body mass based on energy input, metabolic noise, and maintenance cost.

2.2 Estimation of species-level metabolic fluctuations

We derive analytical formulas for the expected species-level metabolic rate, body mass and the allometric scaling exponent in the adult phase (steady state). The metabolic rate in the adult phase simplifies to:

$$B(t) = NB_c + c_+ J_+ + c_- J_-, \tag{9}$$

$$X \sim \text{Laplace}(0, s), \qquad s = \phi N^{\alpha/2} \sqrt{\frac{\Delta t}{2}}. \tag{10}$$

where $J_+ = \max(X, 0)$ and $J_- = \max(-X, 0)$ are the positive and negative parts of the jump term (J). The noise amplitude ϕ controls the microscopic variability, while α governs how fluctuations scale with system size. Positive and negative bursts are weighted by

$$c_+ = E_c\left(\frac{1}{2\eta} + \frac{1 - \eta + \kappa}{2}\right) + \gamma, \tag{11}$$

$$c_- = \kappa E_c + \gamma, \tag{12}$$

The metabolic fluctuations r driven by the microscopic noise structure (ϕ, α) is defined as:

$$r(t) = \log\left(\frac{B_{t+1}}{B_t}\right) \tag{13}$$

where the only difference between B_t and B_{t+1} arises from a stochastic jump in cell number. In this study, we focus on the noise structure, quantified by stochasticity magnitude ϕ, and the correlation exponent α and all other model parameters are kept fixed. For each $\alpha \in [0.9, 2]$, $\phi \in [10^{-6}, 10^{-2}]$ and $N \in [10^{10}, 10^{16}]$ grid, we compute the allometric scaling exponent β_S. We simulate $n_{\text{sim}} = 10^5$ stochastic trajectories for each (N, α, ϕ) parameter combination using Monte Carlo simulations. For each parameter set we draw $X_1, X_2 \overset{\text{iid}}{\sim} \text{Laplace}(0, b)$, to construct B_t and B_{t+1} and compute the sample of r. This procedure yields unbiased estimates that match the simulated time-series statistics for all species sizes.

2.3 Estimation of the allometric exponent β_S at steady state

At adult steady state the expected value of the metabolic rate is

$$\mathbb{E}[B_{\text{adult}}] = A\,N + C\,N^{\alpha/2}, \qquad A = B_c, \qquad C = \phi\sqrt{\frac{dt}{2}}\left[E_c\left(\frac{1}{2\eta} + \frac{1-\eta+\kappa}{2}\right) + \gamma\right]. \tag{14}$$

Similarly, the expected adult mass is

$$\mathbb{E}[m_{\text{adult}}] = m_c N + m_c E N^{\alpha/2}, \qquad E = \phi\sqrt{\frac{dt}{2}}\left(\frac{E_c}{2\eta B_c}\right), \tag{15}$$

where the second term is negligible for large N. Then, we define the metabolic scaling exponent β_S as the fraction of the derivatives of the expected values of adult metabolism and mass.

$$\beta_S = \frac{d\ln\mathbb{E}[B_{\text{adult}}]}{d\ln\mathbb{E}[m_{\text{adult}}]} \tag{16}$$

After differentiating each quantity with respect to N, the steady-state allometric exponent can be calculated as:

$$\beta_S = \frac{AN + \dfrac{\alpha}{2}CN^{\alpha/2}}{AN + CN^{\alpha/2}} \cdot \frac{N + EN^{\alpha/2}}{N + \dfrac{\alpha}{2}EN^{\alpha/2}} \tag{17}$$

The maintenance term scales linearly with cell number, $B_M \propto N_c$, while the stochastic term scales as $H \propto N_c^{\alpha/2}$, producing an effective nonlinear allometric relationship between total metabolism and body mass. The exponent β_S interpolates continuously between sublinear and superlinear regimes as a mixture of $aN + bN^{\alpha/2}$. This simple mechanism generates scaling naturally from the interplay between deterministic processes and stochastic fluctuations and reproduces known biological scaling laws and can be generalized to other stochastic systems such as economic and informational systems.

2.4 Information-theoretic analysis

In order to find the optimal noise structure and the corresponding allometric exponent, first we measure for each α the Kullback-Leibler (KL) divergence between the conditional fluctuation distribution of the mesoscopic fluctuations r and α and the pooled distribution of r across α:

$$D_{\text{KL}}(\alpha, r) = D_{KL}(p(r|\alpha) \,\|\, p(r)) = \sum_{i=1}^{n_{\text{bins}}} p(r_i \mid \alpha)\,\log\frac{p(r_i \mid \alpha)}{p(r_i)}. \tag{18}$$

For each species size (i.e., each N), we evaluate $D_{\text{KL}}(\alpha, r)$ across all α values between $[0.9, 2]$. Note that here we define optimal as maximal robustness of macroscopic fluctuations to microscopic noise. Then, we find the minimal information neutrality point, where $p(r|\alpha)$ is most similar to the pooled distribution $p(r)$:

$$\alpha_{\text{min}}(N, \phi) = \arg\min_{\alpha} D_{\text{KL}}(\alpha, r) \tag{19}$$

At the information neutrality point, α has minimal effect on the distribution of r. We define the information-neutral valley as the region in parameter space where the distribution of macroscopic fluctuations is least sensitive to variations in the microscopic correlation exponent α, quantified by a minimum in the $D_{KL}(\alpha, r)$. This minimum defines the most robust correlation exponent α_{min}. To reduce Monte Carlo noise, we smoothed $(\alpha, D_{\mathrm{KL}})$ trajectories before locating the minimum. Then, we calculate the corresponding allometric exponents $\beta(\alpha_{min})$ for each body size across all ϕ values.

The goal is to determine whether there exists a species–independent optimal noise scale ϕ^* at which the allometric scaling exponent is universal. To do so, we calculate the overlap of the species-specific $\alpha_{min} - \phi$ curve representing the universal crossover. This value or interval represent the point where all species exhibit maximal robustness to microscopic noise and at the same time maximally correlated microscopic noise structure. For each species we identify the noise values ϕ at which robustness is maximal, i.e. when $\alpha_{\min}(\phi, N)$ achieves its maximum over all possible ϕ values:

$$\phi^*(N) = \max_{\phi'} \alpha_{\min}(\phi, N) \tag{20}$$

which defines the structural ridge or crossover region. Then, for each ϕ^*, we compute the corresponding allometric scaling exponent β^*.

To check for universality across species trajectories, we apply a min-max normalization ($x' = \frac{x - x_{\min}}{x_{\max} - x_{\min}}$) of α and β, performed independently for each species size (N). This procedure is consistent with the linear reparametrization of coupling space under the renormalization group [34, 35], and produces a dimensionless coordinate in which the trajectories collapse onto a universal curve. Although the Renormalization Group (RG) terminology used here is phenomenological, we explicitly test the stability of the information-neutral landscape under blocking transformations (see for more details in Supplementary Material).

As a complementary analysis, we quantify the sensitivity of the mesoscopic observable r to the scaling exponent α using the Fisher Information:

$$I_r(\alpha) = \sum_{i=1}^{n_{\mathrm{bins}}} \frac{\left(\partial_\alpha p(r_i \mid \alpha)\right)^2}{p(r_i \mid \alpha)}. \tag{21}$$

$$\partial_\alpha p(r_i \mid \alpha) \approx \frac{p(r_i \mid \alpha + \delta) - p(r_i \mid \alpha - \delta)}{2\delta}, \tag{22}$$

where is set to $\delta = 0.02$. Then, calculating $I_r(\alpha_{min})$ gives the sensitivity of the most robust correlation exponent to microscopic fluctuations.

Figure 1: **Schematic representation of the analytical concept.** (A) Metabolic fluctuations generated from uncorrelated (top) and correlated (bottom) bursty cell dynamics. (B) In order to identify the optimal noise structure across species sizes, first we minimise the KL divergence between the correlation exponent and the metabolic fluctuations, then we maximise the identified α_{min} across the fluctuation coefficients ϕ. (C) This results in a *crossover* region representing an optimal noise structure, which leads to identify a universal scaling exponent β^*.

3   Results

Our results reveal how information mapping from microscopic bursty cell dynamics to organism-level fluctuations shapes the species-level allometric exponent. First, we computed the KL divergence between the α and the organism-level metabolic fluctuations r across a α-ϕ grid for each species sizes (Fig.2A). In each parameter combination, a correlation exponent α_{min} can be identified, where the KL divergence reached its minimum (information-neutral valley). These minimum correlation exponents are in the range of $\sim$ 1.4–1.8 across species sizes. At α_{min} the metabolic fluctuations $r(t)$ retain the least sensitivity to the microscopic parameter α, indicating that the coarse-grained description is maximally insensitive to microscopic details.

To test for universality, we use the normalised trajectories of the exponents $\beta'(\alpha_{min})$ and α', which closely collapse across species, revealing a universal scaling law emerging at the point of minimal dependence on α (Fig.2B). This is consistent with the renormalization group theory, i.e. values of the control parameter at which the coarse-grained statistics become invariant under changes in microscopic parameters. In this framework, species size (N) plays the role of a system-size, where increasing N corresponds to moving toward the macroscopic limit, where microscopic details become irrelevant. These invariant structures are identified through the behavior of the KL divergence, therefore reflects an information-theoretic signature of universality.

As a next step, we calculated the crossover region where the minimum correlation exponents

Figure 2: **Sensitivity of stochastic metabolic fluctuations across species sizes.** (A) KL divergence as a function of the microscopic correlation exponent α for a representative species size and noise amplitude ϕ, revealing an information-neutral valley. (B) Each curve corresponds to a trajectory parameterized by ϕ. After normalization, trajectories collapse across species sizes. (C) The maximised $\alpha_{min}(\phi)$ reveal the structural ridge, and the corresponding $\beta*$ exponent. (D) The sensitivity measured as the Fisher Information of the metabolic fluctuations to the scaling exponents α_{min}.

reach their maximum, which identifies the optimal fluctuation coefficient ϕ^* (Fig.2C). This represents the structural ridge where the correlation exponents are minimal and maximally size-dependent and correlated. This minimax criterion identifies the most robust noise structure. Within the crossover region, the metabolic scaling exponent is narrowly distributed around $\beta^* = 0.77$ with a standard deviation of 0.02. This indicates that, once coarse-grained to the crossover regime, species spanning six orders of magnitude in maximum cell number share an effectively universal metabolic scaling exponent that is in the vicinity of the 3/4 Kleiber's law. The tight clustering of β^* values across species in the universal region provides direct evidence for universality that does not rely on criticality.

Furthermore, we analysed the sensitivity of r with respect to α quantified by the Fisher Information. On the joint information plane where the KL divergence $D_{\mathrm{KL}}[\alpha]$ has its minimum, the Fisher information is not divergent and of intermediate magnitude (Fig.2D). For large ϕ, the dynamics amplify noise, the Fisher and KL divergence decrease and the system becomes nearly insensitive to microscopic noise leading to diminishing allometric exponents.

4 Discussion

The presented analysis confirms that universality can arise from a simple information neutrality criterion in biological systems without phase transition. Contrary to the Edge of Chaos assumption [2], our results show that metabolic scaling does not operate at a point of maximal information transmission or maximal sensitivity. Instead, universality emerges where the information transmission is minimal. The identified information-neutral valley corresponds to a noncritical crossover point or region where Fisher information is low to moderate and where organism-level metabolic fluctuations are maximally robust to microscopic stochasticity and species converge onto a universal scaling exponent. This indicates that universal metabolic scaling is the consequence of evolution tuning itself rather towards a point or plateau of maximal robustness then to critical boundary. Interestingly, the canonical 3/4 scaling exponent [11] emerges close to this robust crossover region.

These results correspond to a phenomenological RG, which emerges from coarse-graining of microscopic cellular bursts into mesoscopic metabolic fluctuations and consequently to macroscopic scaling laws. Importantly, the universality observed, i.e. the collapse of normalised α' and β' across species sizes, reflects a non-critical RG-like behavior. Note that here RG-like is used in a phenomenological sense, referring to the collapse of normalized trajectories under coarse-graining, rather than to a formally defined renormalization transformation. Such noncritical universal behaviors are increasingly recognized in biology. They appear in developmental scaling [36], neural population dynamics [37], biochemical reaction networks, protein aggregation, and scaling laws in ecosystems [38]. In these systems, robustness and universality emerge not from being near a phase transition, but rather from the natural elimination of microscopic detail as biological units are aggregated into larger scales [5]. These findings are similar to a previous model, where universal metabolic scaling arises from life-history optimisation [22].

The large variance in observed empirical diversity is more likely a result of a locally optimised trait space rather than global optimisation. Evolution can be formalised as a stochastic adaptive walk on rugged landscapes [39, 40] constrained by historical contingencies and developmental architecture [14, 41, 15], which prevent natural selection from globally optimizing traits. The average scaling exponent of ~ 0.77 recovered in this analysis therefore represents a purely theoretical optimum closely coinciding with the observed Kleiber's law. Note that a local minimum appears around $\alpha = 1$, which corresponds to uncorrelated noise structure.

This study offers a new theoretical perspective on how information neutrality can give rise to universal scaling laws. More broadly, the framework developed here suggests that information neutrality principles may underlie the universal scaling behavior of noisy processes. This model is generalizable to other complex systems.

References

[1] T. Mora, W. Bialek, Are Biological Systems Poised at Criticality? *Journal of Statistical Physics* **144**, 268 (2011).

[2] C. G. Langton, Computation at the edge of chaos: Phase transitions and emergent computation. *Physica D: Nonlinear Phenomena* **42**, 12 (1990).

[3] M. A. Munoz, Colloquium: Criticality and dynamical scaling in living systems. *Reviews of Modern Physics* **90**, 031001 (2018). Publisher: APS.

[4] E. F. W. Heffern, H. Huelskamp, S. Bahar, R. F. Inglis, Phase transitions in biology: from bird flocks to population dynamics. *Proceedings of the Royal Society B: Biological Sciences* **288**, 20211111 (2021). Publisher: Royal Society.

[5] P. A. Marquet, *et al.*, Scaling and power-laws in ecological systems. *Journal of Experimental Biology* **208**, 1749 (2005). Publisher: Company of Biologists.

[6] A.-L. Barabási, R. Albert, Emergence of Scaling in Random Networks. *Science* **286**, 509 (1999). Publisher: American Association for the Advancement of Science.

[7] D. Naug, Metabolic scaling as an emergent outcome of variation in metabolic rate. *Philosophical Transactions of the Royal Society B: Biological Sciences* **379**, 20220495 (2024). Publisher: Royal Society.

[8] A. Tabi, W. Merbis, F. Santos, R. Sole, Metabolic fluctuations explain allometric scaling diversity. (2025). ISSN: 2692-8205 Pages: 2025.11.06.686202 Section: New Results.

[9] J. P. Sethna, J. P. Sethna, *Statistical Mechanics: Entropy, Order Parameters, and Complexity*, Oxford Master Series in Physics (Oxford University Press, Oxford, New York, 2021), second edition, second edition edn.

[10] J. H. Brown, J. F. Gillooly, A. P. Allen, V. M. Savage, G. B. West, Toward a Metabolic Theory of Ecology. *Ecology* **85**, 1771 (2004). _eprint: https://onlinelibrary.wiley.com/doi/pdf/10.1890/03-9000.

[11] M. Kleiber, Body size and metabolism. *Hilgardia* **6**, 315 (1932).

[12] K. Schmidt-Nielsen, Scaling in biology: The consequences of size. *Journal of Experimental Zoology* **194**, 287 (1975). _eprint: https://onlinelibrary.wiley.com/doi/pdf/10.1002/jez.1401940120.

[13] W. A. Calder, *Size, Function, and Life History* (Harvard University Press, 1984). Google-Books-ID: teSzzQEACAAJ.

[14] P. Alberch, The logic of monsters: Evidence for internal constraint in development and evolution. *Geobios* **22**, 21 (1989).

[15] R. Solé, *et al.*, Fundamental constraints to the logic of living systems. *Interface Focus* **14**, 20240010 (2024). Publisher: Royal Society.

[16] J. P. DeLong, J. G. Okie, M. E. Moses, R. M. Sibly, J. H. Brown, Shifts in metabolic scaling, production, and efficiency across major evolutionary transitions of life. *Proceedings of the National Academy of Sciences* **107**, 12941 (2010). Publisher: Proceedings of the National Academy of Sciences.

[17] A. Clarke, P. Rothery, N. J. B. Isaac, Scaling of basal metabolic rate with body mass and temperature in mammals. *Journal of Animal Ecology* **79**, 610 (2010). _eprint: https://onlinelibrary.wiley.com/doi/pdf/10.1111/j.1365-2656.2010.01672.x.

[18] T. M. Hoehler, *et al.*, The metabolic rate of the biosphere and its components. *Proceedings of the National Academy of Sciences* **120**, e2303764120 (2023).

[19] G. B. West, J. H. Brown, B. J. Enquist, A General Model for the Origin of Allometric Scaling Laws in Biology. *Science* **276**, 122 (1997). Publisher: American Association for the Advancement of Science.

[20] M. Rubner, Über den Einfluss der Körpergrösse auf Stoff- und Kraftwechsel. (1883). Pages: 536 Publication Title: Zeitschrift für Biologie Volume: 19.

[21] A. A. Heusner, Energy metabolism and body size. I. Is the 0.75 mass exponent of Kleiber's equation a statistical artifact? *Respiration Physiology* **48**, 1 (1982).

[22] C. R. White, L. A. Alton, C. L. Bywater, E. J. Lombardi, D. J. Marshall, Metabolic scaling is the product of life-history optimization. *Science* **377**, 834 (2022).

[23] G. Tkačik, W. Bialek, Information Processing in Living Systems. *Annual Review of Condensed Matter Physics* **7**, 89 (2016). Publisher: Annual Reviews.

[24] R. Lande, *et al.*, *Stochastic Population Dynamics in Ecology and Conservation*, Oxford Series in Ecology and Evolution (Oxford University Press, Oxford, New York, 2003).

[25] R. M. Nisbet, W. Gurney, *Modelling fluctuating populations : reprint of first Edition (1982)* (Blackburn Press, 2003).

[26] A. J. McKane, T. J. Newman, Predator-Prey Cycles from Resonant Amplification of Demographic Stochasticity. *Physical Review Letters* **94**, 218102 (2005). Publisher: American Physical Society.

[27] F. A. Labra, P. A. Marquet, F. Bozinovic, Scaling metabolic rate fluctuations. *Proceedings of the National Academy of Sciences* **104**, 10900 (2007). Publisher: Proceedings of the National Academy of Sciences.

[28] G. B. West, W. H. Woodruff, J. H. Brown, Allometric scaling of metabolic rate from molecules and mitochondria to cells and mammals. *Proceedings of the National Academy of Sciences of the United States of America* **99**, 2473 (2002).

[29] E. Ortega-Arzola, P. M. Higgins, C. S. Cockell, The minimum energy required to build a cell. *Scientific Reports* **14**, 5267 (2024).

[30] M. Esposito, C. Van den Broeck, Three Detailed Fluctuation Theorems. *Physical Review Letters* **104**, 090601 (2010). Publisher: American Physical Society.

[31] U. Seifert, Stochastic thermodynamics, fluctuation theorems and molecular machines. *Reports on Progress in Physics* **75**, 126001 (2012). Publisher: IOP Publishing.

[32] G. B. West, J. H. Brown, B. J. Enquist, A general model for ontogenetic growth. *Nature* **413**, 628 (2001). Publisher: Nature Publishing Group.

[33] A. M. Makarieva, *et al.*, Mean mass-specific metabolic rates are strikingly similar across life's major domains: Evidence for life's metabolic optimum. *Proceedings of the National Academy of Sciences* **105**, 16994 (2008). Publisher: Proceedings of the National Academy of Sciences.

[34] K. G. Wilson, J. Kogut, The renormalization group and the expansion. *Physics Reports* **12**, 75 (1974).

[35] J. Cardy, *Scaling and Renormalization in Statistical Physics* (1996). Publication Title: Scaling and Renormalization in Statistical Physics ADS Bibcode: 1996scph.book.....C.

[36] V. M. Lauschke, C. D. Tsiairis, P. François, A. Aulehla, Scaling of embryonic patterning based on phase-gradient encoding. *Nature* **493**, 101 (2013). Publisher: Nature Publishing Group.

[37] L. Meshulam, J. L. Gauthier, C. D. Brody, D. W. Tank, W. Bialek, Coarse Graining, Fixed Points, and Scaling in a Large Population of Neurons. *Physical Review Letters* **123**, 178103 (2019). Publisher: American Physical Society.

[38] J. R. Banavar, J. Damuth, A. Maritan, A. Rinaldo, Scaling in Ecosystems and the Linkage of Macroecological Laws. *Physical Review Letters* **98**, 068104 (2007). Publisher: American Physical Society.

[39] S. Kauffman, S. Levin, Towards a general theory of adaptive walks on rugged landscapes. *Journal of Theoretical Biology* **128**, 11 (1987).

[40] T. Lenormand, D. Roze, F. Rousset, Stochasticity in evolution. *Trends in Ecology & Evolution* **24**, 157 (2009). Publisher: Elsevier.

[41] S. J. Gould, R. C. Lewontin, The spandrels of San Marco and the Panglossian paradigm: a critique of the adaptationist programme. *Proceedings of the Royal Society of London. Series B. Biological Sciences* **205**, 581 (1997). Publisher: Royal Society.

Supplementary Materials for:
Information-Theoretic Origins of Metabolic Scaling

S1 Coarse-graining of mesoscopic fluctuations

To assess whether the information-neutral regime is stable under coarse-graining, we performed a blocking analysis of the mesoscopic fluctuation observable. Note that this analysis is intended only as a diagnostic test of RG-like phenomenology rather than as a formal renormalization-group analysis.

First, we define the coarse-grained fluctuation of the k^{th} step by block summation,

$$R_k(t) = \sum_{i=0}^{k-1} r(t+i) = \log \frac{B(t+k)}{B(t)}.$$

(1)

This corresponds to aggregating fluctuations over k consecutive steps and is analogous to blocking transformations used in coarse-graining analyses.

For independent or weakly correlated increments, one would generically expect the variance of R_k to grow with k. However, across the explored parameter ranges we find that the standard deviation of R_k is constant over all block sizes. This happens because $r(t)$ is defined as a relative increment of a stationary process, such that aggregation does not lead to systematic broadening of the distribution. Therefore, no rescaling of R_k (e.g. z-score or $\sqrt{k}$ normalisation) is required to compare distributions across different coarse-graining levels.

Using the coarse-grained variable R_k, we computed the $D_{KL}(\alpha, r)$,

$$D_{\mathrm{KL}}\big(p(R_k \mid \alpha) \,\|\, p(R_k)\big),$$

(2)

where $p(R_k)$ denotes the distribution pooled across α. We find that the KL landscape as a function of α is invariant under coarse-graining: both the shape of the landscape and the location of its minimum are unchanged as k is increased (Figure S2). This invariance indicates that the information-neutral exponent $\alpha_{\min}$ identified is stable under coarse-graining of mesoscopic fluctuations and therefore supports the phenomenological RG interpretation.

Figure S1: **Species allometric exponents by organism size across different noise structures (α and ϕ).**

Figure S2: **Coarse-grained fluctuations.**

Figure S3: **Example of Fisher Information across** α. The dashed lines correspond to the α_{min} values, i.e. the minimum of D_{KL} curve with $\phi = 10^{-5.3}$. At that point, the Fisher Information metrics take an intermediate value.