Title: **Opposite Effects of Alpha Oscillations on Mind-wandering
With Eyes Open and Closed**

Abbreviated Title: Mind-wandering, Alpha Oscillations, and Eye State

Esther M. Thielking[1,2], Hannah Ramsey[1], Jason Samaha[3], Thomas Andrillon[4] & Luca Iemi[1,2]

Affiliations:
1. *Neuroscience & Behavior Department, Barnard College, New York, NY, 10027, USA*
2. *Reid Hall, Columbia University, 75006, Paris, France*
3. *Department of Psychology, University of California, Santa Cruz, CA, 95064, USA*
4. *Sorbonne Université, Institut du Cerveau - Paris Brain Institute - ICM, Inserm, CNRS, APHP, Hôpital de la Pitié Salpêtrière, 75013, Paris, France*

Corresponding author: Esther M. Thielking; esther.thielking@gmail.com

Pages: 49
Figures: 5
Tables: 2
Word count: Abstract: 245; Introduction: 645; Discussion: 1499

Acknowledgements:
We thank Barnard College students Emma Kornberg, Hiya Jain, Jena Mamdani, and Sophia Faisal for assistance with the EEG recording; Peter Balsam for providing resources, and Esperanza Jubera-García, Niko A. Busch, and Saskia Haegens for helpful feedback.

Competing Interests:
The authors have no competing financial interests to declare.

Abstract

Spontaneous fluctuations in brain activity underlie mind-wandering, when mental

experience sometimes becomes unrelated to what we are currently doing. Mind-wandering

research has investigated ongoing neural oscillations in the alpha frequency band (7-14 Hz),

thought to reflect changes in arousal and attention. However, studies disagree whether alpha

power increases or decreases while mind-wandering compared to focused attention. We

hypothesized that these opposite effects arise from differences in eye state across studies, as

eye closure increases alpha power and can reverse its relationship to sleepiness, which is itself

related to mind-wandering. To test this, we recorded EEG while male and female participants with

eyes either open or closed were probed to report whether they were focused on the auditory

attention task or mind-wandering, as well as how sleepy they were. Consistent with our

hypothesis, we found that increased alpha power made both mind-wandering and sleepiness

reports more likely in the eyes-open group but less likely in the eyes-closed group. A systematic

review of past literature largely replicated these opposite relationships across eye states in

different paradigms. We propose that these results reflect an inverted-U relationship between

alpha power and mind-wandering, formed by a positive relationship at low alpha power values

typical of eyes-open states and a negative relationship at high alpha power values typical of eyes-

closed states. By sampling alpha power across eye states in the same paradigm, this work

reconciles contradictions in prior literature and clarifies how ongoing neural oscillations reflect the

stream of mental experiences.

Significance Statement

Brain signals tracked with electroencephalography (EEG) can provide a read-out of a

person's current mental experience. Increases in a brain signal called alpha oscillations usually

indicate sleepiness when recorded with eyes open. Surprisingly, the read-out can be reversed

when eyes are closed: increases in alpha oscillations indicate alertness. Here, we found that this

reversal explains the puzzling relationship between alpha oscillations and mind-wandering, when

mental experience becomes unrelated to the task-at-hand. In moments of increased alpha,

participants with eyes open reported more mind-wandering and sleepiness, whereas participants

with eyes closed reported more on-task focus and alertness. One intriguing possibility is that

decreases in arousal reflected by changes in alpha oscillations trigger episodes of mind-

wandering.

Introduction

Our mental experience can spontaneously become unrelated to what we are currently

doing and perceiving: this is the common experience of mind-wandering. At the same time, the

state of the brain changes from moment to moment due to ongoing, internally generated neural

processes occurring independently of task demands or sensory input. How do these spontaneous

fluctuations of mental experience and brain state relate to each other? To address this question,

numerous studies have recorded a prominent type of spontaneous brain activity, alpha

oscillations (7–14 Hz), while participants sustained their attention on repetitive tasks and

intermittently reported mind-wandering (Kam et al., 2022). These studies show that moment-to-

moment fluctuations in alpha power are correlated with mind-wandering reports. However, the

direction of this relationship has been inconsistent: some studies report positive relationships

(stronger alpha power while mind-wandering; Compton et al., 2019; Arnau et al., 2020; Nakatani

et al., 2024), whereas others report negative relationships (weaker alpha power while mind-

wandering; Braboszcz and Delorme, 2011; van Son et al., 2019a; Rodriguez-Larios and Alaerts,

2021; Rout et al., 2024).

These findings have been described as "conflicting," "inconsistent," or "complex"

(Compton et al., 2019; Arnau et al., 2020; Kam et al., 2022; Nakatani et al., 2024), and although

methodological differences have been proposed as an explanation, the debate remains

unresolved. We reasoned that these inconsistencies reflect differences in eye state across

studies. Closing the eyes increases the power of alpha oscillations (Berger, 1929) and can

reverse the direction of its relationship with arousal (e.g., subjective sleepiness: Strijkstra et al.,

2003; Kaida et al., 2006), which is thought to influence mind-wandering (Andrillon et al., 2019;

Jubera-García et al., 2021). Accordingly, we propose an underlying inverted-U function relating

mind-wandering with alpha power across eye states. Here, mind-wandering and alpha power are

positively correlated when alpha oscillations fluctuate within a low power range as during eyes-

open states, corresponding to the left side of the inverted-U (Fig. 1). By contrast, mind-wandering

and alpha power are negatively correlated when alpha oscillations fluctuate within a high power

range as during eyes-closed states, corresponding to the right side of the inverted-U (Fig. 1). In

this way, the seemingly contradictory positive and negative relationships observed in prior

literature can be understood as reflecting different sides of an underlying non-linear function.

Figure 1. Inverted-U model.
Schematic illustration of the inverted-U relationship between alpha (7-14 Hz) power and mind-
wandering. The x-axis represents the range of alpha power values observed across eye states.
The left y-axis shows the distribution density of alpha power values. The red distribution
represents eyes-open states corresponding to the low alpha power range, while the blue
distribution represents eyes-closed states corresponding to the high alpha power range. The right
y-axis shows mind-wandering rate, with the gray curve depicting the inverted-U relationship
between alpha power and mind-wandering rate: the left side spanning the low alpha power range
shows a positive relationship, and the right side spanning the high alpha power range shows a
negative relationship.

To test this, we recorded EEG during eyes-open and eyes-closed resting states followed

by a sustained attention-to-response task (SART; auditory go/no-go) with intermittent experience-

sampling probes and subjective sleepiness ratings. This paradigm allowed us to quantify

fluctuations in mind-wandering and sleepiness through subjective reports, as well as reaction

times (RTs), known to co-vary with mind-wandering (Leszczynski et al., 2017; Andrillon et al.,

2021). To sample different sides of the inverted-U function, we manipulated the range of alpha

power values by varying eye state between participants: one group performed the task with eyes

open (EO group), while another group performed it with eyes closed (EC group). By comparing

alpha power between the task and the resting states, we confirmed that eye-closed states boosted

alpha power and thus produced a high power range relative to eyes-open states (Barry et al.,

2007). For each group, we then assessed how ongoing alpha power was correlated with

momentary fluctuations of subjective experience and RTs. We found that alpha power showed a

positive correlation with subjective reports of mind-wandering and sleepiness in the EO group,

and a negative correlation in the EC group, consistent with the inverted-U model.

Further, a systematic literature review dividing the studies by eye state largely replicated

this pattern of positive and negative relationships, further supporting the inverted-U model. RTs

did not co-vary with mind-wandering and were negatively correlated with alpha and beta power

only in the EC group, replicating previous results obtained in the same eye state. Taken together,

the results of this study reconcile inconsistencies in the literature and help specify the link between

spontaneous neural oscillations and ongoing mental experience.

Materials and Methods

Participants

Sixty healthy adults participated in this study. All provided written informed consent prior to data collection. Participants were randomly assigned to perform the main task with eyes either closed (EC group; N=30) or open (EO group; N=30). Before pre-processing, participants were excluded due to EEG acquisition issues (N=5) or insufficient probe response variability (i.e., percentage mind-wandering reports below 10% or above 90%; N=5). Groups did not differ in sex (EO: 4M 22F, EC: 1M 23F; Fisher's exact test, p=0.351) or age (EO: mean 19.8 ± standard error of the mean (SEM) 0.23 years; EC: mean 19.7 ± SEM 0.19 years; t_{48}=0.574, p=0.569). All analyses included 24 EC and 26 EO subjects. Experimental procedures were approved by the Institutional Review Board (IRB) at Barnard College.

Behavioral Paradigm

Participants completed a resting-state and a task session (Fig. 2). The resting-state session consisted of two 2-min blocks (eyes-open and eyes-closed) presented in randomized order and separated by a short self-paced break. This was used to establish each participant's low (eyes-open) and high (eyes-closed) ranges of alpha power values. Participants were instructed to stay still, fixate on a central mark (in eyes-open rest), and avoid thinking of anything in particular. Following the resting state session, participants completed six 5-min task blocks of an auditory sustained attention to response task (SART), each separated by a self-paced break. For the duration of each task block, participants in the EO group were instructed to maintain fixation on a cross, whereas participants in the EC group were instructed to maintain eye closure.

In the main task, participants performed an auditory SART that required them to respond as quickly and as accurately as possible via button press to frequent 'go' stimuli and to withhold responses to infrequent 'no-go' stimuli (i.e., auditory go/no-go task; Fig. 2). Task performance

was measured as misses (no button press in go trials), false alarms (button press in no-go trials),

and RTs (analyzed on correct go trials only). At pseudorandom intervals, participants were probed

to report whether they were mind-wandering or on-task just before the probe. Participants were

instructed that mind-wandering consisted of attending to task-unrelated thoughts and that on-task

consisted of attending to the auditory SART (Smallwood and Schooler, 2015; Arnau et al., 2020;

Compton et al., 2024). Participants were also informed that we consider mind-wandering to be a

fundamental part of human cognition and were encouraged to respond honestly based on their

subjective experience.

After each resting and task block, participants rated their subjective sleepiness on the 9-

point Likert-type Karolinska Sleepiness Scale (KSS; 1= "extremely alert", 9= "very sleepy, great

effort to keep awake"; Kaida et al., 2006) and reported whether their eyes were open or closed to

confirm compliance with the task instructions.

Figure 2. Experimental paradigm.
Participants (N=60) completed an eyes-open and an eyes-closed resting state block, with each
block lasting 2 min and the order randomized across participants, followed by an auditory
sustained attention to response task (SART). Participants were randomly assigned to perform the
SART with eyes either open or closed. The SART comprised 6 blocks, with each lasting 5 min. A
spoken digit was presented on each SART trial, with an inter-stimulus interval (ISI) of 2 s:
participants responded to frequent go digits and withheld responses to infrequent no-go digits.
With a pseudorandom inter-probe interval (IPI) of 50 to 70 s, participants were probed to report
whether they had just been on-task or mind-wandering. Sleepiness ratings (Karolinska Sleepiness
Scale, KSS) were collected after each rest and task block. EEG was recorded from 8 occipito-
parietal electrodes.

Stimuli

The task was implemented in MATLAB (R2022b; The MathWorks; RRID:SCR_001622) using Psychtoolbox-3 (RRID:SCR_002881; Brainard, 1997). Auditory stimuli were synthesized with Google Cloud Text-to-Speech (*Cloud Text-to-Speech API v1*, n.d.). Spoken digits (0.499 s duration) were SART stimuli (go and no-go), and the spoken word "pause" was the experience-sampling probe (0.545 s duration). Files were sampled at 48 kHz and delivered via in-earbuds at a fixed volume. There were 9 spoken digits, ranging from "one" to "nine;" "three" and "seven" were no-go stimuli, while the others were go stimuli. Each task block included 158 trials (34 no-go, 119 go, 5 probes). The stimulus sequence was pseudorandomized for each block: 17 independent random permutations of the spoken digits were concatenated to form the base digit list (length 153) from which 5 probe insertion indices were drawn separated by a uniformly random step of 25–35 trials (mean 30.19 ± SEM 0.04 trials), corresponding to a 50-70 second inter-probe interval (IPI) comparable to previous probe-caught experience-sampling paradigms (Compton et al., 2019; Schubert et al., 2020; Andrillon et al., 2021; Rodriguez-Larios and Alaerts, 2021; Kucyi et al., 2024). The inter-stimulus-interval (ISI) was 2 s (Bastian and Sackur, 2013; Leszczynski et al., 2017).

EEG Recording and Preprocessing

An Enobio-8 system (Neuroelectrics) was used to record continuous EEG sampled at 500 Hz. We used 8 gel channels over occipital and parietal areas of interest (P3, Pz, P4, PO3, PO4, PO7, PO8, Oz; 10-10 system), corresponding to the scalp location of the most prominent alpha rhythm in the human EEG (Tenke et al., 2015; Hohaia et al., 2022). EEG data were referenced to an ear-clip electrode. The FieldTrip toolbox (version 20180801; Oostenveld et al., 2011) running on MATLAB (R2024b; The MathWorks; RRID:SCR_001622) was used to process and analyze the data. Continuous data were epoched into 1000 ms epochs consecutively for resting-state blocks or from -2000 to 500 ms relative to stimulus onset for task blocks. Epoched data were

visually inspected to remove major artifacts. One task block was removed from a participant in

the EO group, who reported closing their eyes during the block. A mean of 20.94 ± SEM 5.55

epochs were rejected per participant. In three participants, one channel was excluded; in one

participant, two channels were excluded.

Spectral analysis

Milliseconds-scale spectral analysis: We computed time–frequency representations

(TFRs) of power from the epoched data using an adaptive sliding-time window of three cycles in

length ($\Delta t = 3/f$), with a Hanning taper applied before obtaining power via a fast Fourier transform

(FFT) approach. The TFRs were computed at 100 ms intervals across each epoch, and spectral

power was estimated from 5 to 30 Hz in 0.5 Hz steps. This milliseconds-scale TFR was used in

the analysis of the relationship between spectral power and RTs.

Seconds-scale spectral analysis: We extracted 1000 ms consecutively for resting epochs

and from –1000 to 0 ms relative to stimulus onset on each trial for task epochs, then estimated

spectral power using a Hanning-tapered fast FFT (5–30 Hz, 0.5 Hz steps). The second half of the

2s ISI were analyzed to measure "ongoing" power, as this window minimizes contamination from

evoked or induced activity related to stimulus presentation or motor responses from the preceding

trial. We then constructed a seconds-scale time–frequency representation (TFR) by stacking the

power spectra from the five 1-s epochs preceding each probe. This created a temporal axis of

sequential pre-probe periods (e.g., –9 to –8 s, –7 to –6 s, –5 to –4 s, –3 to –2 s, and –1 to 0 s

relative to probe onset) consistent with the time range analyzed in prior studies (Baldwin et al.,

2017; Compton et al., 2019; Jin et al., 2019; Schubert et al., 2020; Andrillon et al., 2021; Kucyi

et al., 2024). This seconds-scale TFR was used for the analysis of the relationship between

spectral power and mind-wandering. We averaged the seconds-scale data within each block for

the analysis of the relationship between spectral power and sleepiness.

Epoch-wise power values were averaged across channels for all analyses. Because we

were primarily interested in brain activity before experimental events (pre-probe and prestimulus),

we did not apply baseline correction in either the milliseconds-scale or seconds-scale spectral

analyses.

Testing the Inverted-U Model

By "inverted-U" we specify that the relationship between alpha power and subjective

reports changes in direction (sign) along the range of alpha power values observed across eye

states from positive sign in the low range of alpha power to negative sign in the high range of

alpha power. The eye state manipulation in our study allowed us to sample low (EO group) and

high (EC group) ranges of alpha power values, corresponding to the left and right sides of the

hypothesized inverted-U. To test this, we fit separate linear relationships within the EO and EC

groups and tested whether they have individually significant slopes with opposite signs ("two-

lines" test: Simonsohn, 2018). We chose this method over fitting a single quadratic function

because it is compatible with our between-subject design and because quadratic models can both

make unjustified assumptions about the function's shape and inflate false-positive rates

(Simonsohn, 2018). Evidence for an inverted-U is obtained if two criteria are met: (i) eyes-open

and eyes-closed states exhibit non-overlapping alpha power ranges (see *Analysis of Spectral*

Power and Eye State); (ii) the EO group shows a significantly positive relationship and the EC

group shows a significantly negative relationship, between alpha power and subjective reports

(see *Analysis of Behavioral Responses and Their Relationship With Spectral Power*).

Analysis of Spectral Power and Eye State

To demonstrate that eyes-open and eyes-closed states produce low and high ranges of

alpha power values, respectively, we performed two analyses. First, we computed the difference

in spectral power between (i) eyes-open and eyes-closed epochs from resting blocks across all

participants and (ii) epochs from resting blocks (either eyes-open or eyes-closed) and an equal

number of randomly selected epochs from task blocks of the opposite eye state, separately for

each participant group. Second, we examined whether eyes-open and eyes-closed states were

distinguished by partially non-overlapping ranges of power distributions. For each participant and

frequency we computed the percentage of epochs whose power values are observed in one eye

state but never in the other. The eyes-open non-overlap was defined as the proportion of eyes-

open epochs with power values below the minimum power value observed in the eyes-closed

distribution. The eyes-closed non-overlap was defined as the proportion of eyes-closed epochs

with power values above the maximum power value observed in the eyes-open distribution. We

computed these within-participant non-overlap metrics in two ways: (i) by comparing eyes-open

and eyes-closed epochs from resting blocks across participants, and (ii) by comparing all epochs

from resting blocks (either eyes-open or eyes-closed) and an equal number of randomly selected

epochs in task blocks of the opposite eye state, separately for each participant group. At the group

level, we assessed the significance of eyes-open and eyes-closed non-overlap metrics using

separate cluster-based permutation tests across frequencies (see *Statistical Analyses*).

Analysis of Behavioral Responses and Their Relationship With Spectral Power

For each participant, we used separate generalized linear models (GLMs) to examine first

(i) the contribution of sleepiness ratings in explaining block-average mind-wandering probability,

and second the contributions of EEG spectral power during the task in explaining (ii)

mind-wandering reports, (iii) sleepiness ratings, and (iv) RTs.

For GLMs (i), (iii), and (iv), we fit standard linear regression models of the form:

$$Y = \beta_0 + \beta_1 X + \varepsilon$$

where X is the z-scored predictor (sleepiness rating or spectral power, the latter normalized by

$\log_{10}$-transformation: Smulders et al., 2018), Y is the corresponding z-scored behavioral measure

(mind-wandering probability, sleepiness rating, or RTs, the latter $\log_{10}$-transformed), β_0 is the

intercept, β_1 is the slope coefficient, and ε is the residual error. From these models, we use β_1 to

quantify the contribution of X to explaining variation in Y.

For GLM (ii), we fit a logistic regression model of the form:

241                        $$\text{logit}(P(Y = 1|X)) = \beta_0 + \beta_1 X$$

where X is EEG spectral power ($\log_{10}$-transformed and z-scored), Y is the binary report to the

experience-sampling probe (Y=1 for a mind-wandering report, Y=0 for an on-task report), and

$P(Y = 1|X)$ the probability of reporting mind-wandering given X. From this model, we use β_1 to

quantify the contribution of EEG spectral power to explaining the log-odds of reporting

mind-wandering.

Statistical Analyses

A two-stage statistical approach was applied to test the significance of the observed

statistics described in the preceding sections. At the subject level, we converted the observed

statistics to z-scores using a permutation-based null distribution. This process involved randomly

shuffling either the eye state (differences and non-overlap scores) or the mapping between the

predictor and the behavioral measure (GLM β_1) for 1000 iterations and recomputing the statistic

for each permutation to create a null distribution. The observed statistic was then standardized

by subtracting the mean and dividing by the standard deviation of this null distribution, resulting

in one z-score for each subject, and for each frequency and time point (if applicable) (Samaha et

al., 2017; Iemi and Busch, 2018).

At the group level, we tested whether the subject-level z-scores were significant across

participants (i.e., whether their signs were consistent). For analyses without multiple comparisons

(behavioral GLM), significance was assessed using a two-tailed t-test against zero on the subject-

level z-scores. For other analyses (EEG GLMs), significance was assessed using a

nonparametric cluster-based permutation test, which addresses multiple comparisons across

frequencies and time points (if applicable) (Maris and Oostenveld, 2007). We obtained a

distribution of z-scores under the null hypothesis by randomly permuting their signs 1000 times.

On each iteration, we tested the resulting z-scores with a two-tailed t-test against zero and

assessed the sum of the t values within the largest contiguous cluster of significant time-frequency

points (or frequency points) (cluster alpha: p<0.05), resulting in a distribution of t sums expected

under the null hypothesis. A final p-value was calculated as the proportion of t sums under the

null hypothesis larger than the sum of t values within clusters in the observed data. Thus, p-values

smaller than 0.05 (final alpha) indicate that the observed coefficients were significantly different

from zero.

We quantified evidence for the null hypothesis using Jeffreys–Zellner–Siow (JZS) Bayes

factors (BF_{10}) (MATLAB BayesFactor toolbox; Krekelberg, 2021) with the default Cauchy prior on

standardized effect sizes with scale r = 0.707 (Rouder et al., 2009). We interpreted BF as follows:

BF > 3 indicates evidence favoring the alternative, BF < 1/3 indicates evidence favoring the null,

and values between 1/3 and 3 were treated as inconclusive. For analyses with multiple

comparisons that yielded a significant cluster in only one group, we extracted the mask of

significant time-frequency points from that group and computed the BF at each corresponding

point in the non-significant group. We then reported the percentage of time-frequency points

showing evidence for the null, or inconclusive evidence (Iemi et al., 2017).

Systematic literature review

We conducted a systematic literature review using the search engine PubMed. Our aim

was to find previous studies that examined the relationship between ongoing alpha power and

subjective reports of attentional state (mind-wandering) in neurotypical humans. We searched in

the abstracts and titles for the following key words: (spontaneous OR pre-stimulus OR prestimulus

OR ongoing OR preprobe OR pre-probe) AND (Alpha OR α OR 10 Hz OR 8–12 Hz OR 7-14 Hz)

AND (variability OR oscillations OR fluctuations OR activity OR rhythmic OR power OR amplitude)

AND (subjective attention OR mind-wandering OR mind wandering OR task-focused OR lapses

OR misses). Inclusion required that studies analyzed neurotypical participants with no specialized

training (e.g., meditators were excluded), ongoing alpha power in relation to mind-wandering

reports (i.e., pre-probe window for probe-caught studies), and immediate (not retrospective)

reports of attentional state. Further, we excluded studies that quantified alpha activity using a p-

episode approach (e.g., Boudewyn and Carter, 2018) or that applied baseline correction to alpha

power values (e.g., Hua et al., 2022).

From 29 eligible studies, we extracted (1) eye state during experience sampling: eyes

open or eyes closed, (2) direction of attention: exteroceptive (attending to external stimuli) or

interoceptive (attending to bodily stimuli, e.g., breath counting), with sensory modality coded for

exteroceptive tasks as visual (V), auditory (A), audiovisual (AV), or (A,V) for studies with separate

tasks in auditory and visual modalities, and interoceptive tasks coded as breath counting (Breath),

(3) sampling method: probe-caught (experimentally-initiated report about current attentional state)

or self-caught (participant-initiated reports upon recognizing mind-wandering), (4) relationship

between alpha power and mind-wandering: expressed as the correlation that is expected based

on the results reported by each study. In probe-caught studies, this was measured as a contrast

between ongoing alpha power preceding probes where participants reported mind-wandering and

on-task states; in self-caught studies, this was measured as a contrast between ongoing alpha

power in the period before the button press (interpreted as mind-wandering) and after the button

press (interpreted as on-task). Findings of significantly stronger alpha power before mind-

wandering compared to on-task states are expressed as a positive correlation; findings of

significantly weaker ongoing alpha power before mind-wandering compared to on-task states are

expressed as a negative correlation; findings of no significant difference in ongoing alpha power

between mind-wandering and on-task states are expressed as a "null" correlation.

Code Accessibility

The code used for this study will be publicly released when the study is published.

                                                        Results

314   **Behavior**

315   We probed participants about once a minute to self-report whether they were focused on

performing the auditory SART or engaged in task-unrelated mental experiences (i.e., mind-

wandering), and about once every five minutes to report their sleepiness levels (Fig. 2). Eye state

was manipulated between groups: the EO group performed the task with eyes open, and the EC

group with eyes closed (Fig. 2). Participants reported frequent mind-wandering (EC 67.28±3.67%;

EO 52.14±3.60%; mean ± SEM) and "some signs of sleepiness" during task performance (EO

6.083 ± 0.266; EC 6.182 ± 0.340). Mind-wandering was more likely in blocks when participants

reported feeling sleepier (EO t_{25}=2.236, p=0.035; EC t_{23}=4.104, p<0.001), suggesting that mind-

wandering increased with sleepiness. Additionally, mind-wandering reports were preceded by

more false alarms (EO t_{25}=2.845, p=0.009; EC t_{23}=3.845, p<0.001) and more misses over the 5

trials pre-probe in the EC group (EO t_{25}=1.449, p=0.160, BF=0.524; EC t_{23}=2.114, p=0.046),

suggesting that mind-wandering had a detrimental effect on task accuracy. By contrast, mind-

wandering didn't affect average reaction times (RTs), demonstrated by no difference in mean RT

between correct trials preceding on-task and mind-wandering reports (5 pre-probe trials: both

groups p>0.517, BF<0.279; 1 pre-probe trial: both groups p>0.190, BF<0.52).

330   Comparing EC and EO groups allowed us to investigate the effects of eye state on

participants' behavior (Table 1). The EC group reported more instances of mind-wandering

compared to the EO group, suggesting that eye closure increased mind-wandering. The effects

on task performance were either null (RTs, misses) or inconclusive (false alarms, mind-wandering

and sleepiness correlation). There was a greater increase in sleepiness during the task relative

to rest in the EC group than in the EO group. Interestingly, across participants, eyes-closed rest

was associated with higher sleepiness ratings (5.520 ± 0.220) than eyes-open rest (4.900 ± 0.259;

$t_{49}=3.313$, p=0.002), suggesting that eye closure increased sleepiness not only during the task

but also during rest.

Taken together, these behavioral results suggest that, in this study, mind-wandering is

associated with increased sleepiness and worse task accuracy (but no change in average RT),

and that eye closure results in increased mind-wandering and sleepiness.

Behavioral Metrics	EC (mean ± SEM)	EO (mean±SEM)	t_{48}	p	BF
Mind-wandering (%)	67.28 ± 3.67	52.14 ± 3.60	2.944	0.005	8.384
Δ Sleepiness (%)	42.810 ± 11.474	11.389 ± 5.400	2.541	0.014	3.653
MW-S correlation z(β)	0.795 ± 0.194	0.369 ± 0.165	1.685	0.099	0.893
RTs (s)	0.675 ± 0.021	0.690 ± 0.027	-0.415	0.680	0.304
Misses (%)	1.44 ± 0.57	1.38 ± 0.38	0.090	0.929	0.284
False alarms (%)	19.80 ± 2.60	27.38 ± 3.56	-1.697	0.096	0.907

Table 1. Behavioral effects of eye state.
Relative to the eyes-open (EO) group, the eyes-closed (EC) group showed significantly more
mind-wandering and a significant increase in sleepiness during the task relative to rest. No
significant between-group differences were found in the correlation (z-scored GLM β coefficient;
z(β)) between mind-wandering and sleepiness (MW-S), or in reaction times (RTs), misses, or
false alarms. Comparisons were made with unpaired t-tests, and Bayes factor (BF) analysis was
used to determine whether non-significant results reflected evidence for the null (BF<1/3) or
inconclusive evidence (1/3<BF<3).

Eyes-open and eyes-closed states produce partially non-overlapping ranges of alpha power values

In this study, we used a manipulation of eye state to sample left and right sides of the proposed inverted-U function relating alpha oscillations and mind-wandering (Fig. 1). Based on prior research (Berger, 1929), we expect eyes-open states to sample the low range of alpha power values or the left side of the inverted-U, and eyes-closed states to sample the high range of alpha power values or the right side of the inverted-U (Fig. 1). Thus, we assessed criterion (i) of the inverted-U model: whether eye closure increased mean alpha power and shifted its range to include values that are not observed during eyes-open states (i.e., a different side of the inverted-U).

We first compared ongoing oscillatory power between eyes-open rest and eyes-closed rest. Cluster permutation testing combining participants from both groups revealed that eye closure during rest increased power in the alpha and surrounding bands (5–23.5 Hz, p=0.001, peak t_{49}=10.869 at 10 Hz; data not visualized). We then compared ongoing oscillatory power during task engagement with that observed during rest in the other eye state (for EO and EC groups separately). Cluster-based permutation testing revealed a significant increase in power in the alpha and surrounding bands during task blocks in the EC group relative to eyes-open rest (5–24 Hz, p=0.001, peak t_{23}=7.940 at 11.5 Hz; Fig. 3b), and a significant decrease in power in a similar frequency range during task blocks in the EO group relative to eyes-closed rest (6-22.5 Hz, p=0.006, peak t_{25}=-6.602 at 10 Hz; Fig. 3a). Taken together, these findings confirm the well-documented "alpha blocking" effect (Berger, 1929; Barry et al., 2007; Tenke et al., 2015; ElShafei et al., 2022; Hohaia et al., 2022; Petro et al., 2022) and show that alpha power increases when the eyes are closed regardless of whether oscillatory activity is recorded during rest or task-engaged states.

We hypothesized that manipulating eye state enables sampling alpha power in left and

right sides of the inverted-U. To assess this, we measured the extent of non-overlap between the

eyes-open and eyes-closed epoch-wise power distributions: the proportion of eyes-open values

lower than the eyes-closed minimum (EO non-overlap) and the proportion of eyes-closed values

higher than the eyes-open maximum (EC non-overlap). In resting-state data combining

participants from both groups, cluster permutation testing revealed significant eyes-open non-

overlap spanning the alpha band and surrounding frequencies (5.00–24.50 Hz, p<0.001, peak

t_{49}=4.973 at 10.5 Hz; 25.50–27.00 Hz, p=0.014, peak t_{49}=3.608 at 26 Hz; data not visualized) as

well as significant eyes-closed non-overlap spanning the alpha band and surrounding frequencies

(5.00–13.50 Hz, p<0.001, peak t_{49}=6.066 at 10.5 Hz and 16.00–22.00 Hz, p<0.001, peak

t_{49}=3.841 at 18.0 Hz; data not visualized). We next tested whether alpha power fluctuations during

the task showed non-overlapping values with those observed during rest in the other eye state

for each group separately. Cluster permutation testing revealed significant eyes-open non-overlap

in the alpha and beta bands in the EO group (6.5-7 Hz, p=0.034; 8.0–12.0 Hz, p=0.001, peak

t_{25}=3.273 at 10 Hz; 15.5–19.0 Hz, p=0.002; 29.5-30 Hz, p=0.033; Fig. 3c) and significant eyes-

closed non-overlap in the alpha and surrounding bands in the EC group (6–19 Hz, p=0.001, peak

t_{23}=4.96 at 12.5 Hz; 21–22.5 Hz, p=0.044, and 27.5–29.5 Hz, p=0.037; Fig. 3d).

Taken together, these results confirm criterion (i) of the inverted-U model: the eye-state

manipulation produced non-overlapping alpha-power ranges corresponding to the left (eyes

open) and right (eyes closed) sides of the hypothesized inverted-U function. Critically, establishing

this non-overlap allows us to evaluate the relationship between alpha power and mind-wandering

on each side of the inverted-U.

Figure 3. Eye-state effects on spectral power.
(a,b) Group-average power spectra (mean ± shaded SEM) during task and eyes-closed rest in the eyes-open (EO) group (a), and during task and eyes-open rest in the eyes-closed (EC) group (b). (c,d) Group-average frequency representation of EO non-overlap: proportion of EO task epochs with power values below the minimum value observed in eyes–closed rest (c), and EC non-overlap: proportion of EC task epochs with power values above the maximum value observed in eyes-open rest (d). Significant frequency clusters are indicated with black horizontal lines (cluster-based permutation test, p<0.05). Each eye state is characterized by distinct, partially non-overlapping alpha power ranges: specifically, eyes-closed states increase alpha power, shifting it to values never observed during eyes-open states.

Alpha power shows an inverted-U relationship with mind-wandering and sleepiness

We tested a model proposing that mind-wandering follows an inverted-U shaped function across the range of alpha power values observed in eyes-open and eyes-closed states (Fig. 1). The left side of the inverted-U corresponds to the lower range of alpha power values (sampled here during eyes-open states) where a positive correlation with mind-wandering probability is predicted; the right side of the inverted-U corresponds to the higher range of alpha power values (sampled here during eyes-closed states) where a negative correlation with mind-wandering probability is predicted. Thus, we assessed criterion (ii) of the inverted-U model by fitting separate

GLMs for the EO and EC groups to examine the pre-probe, trial-wise correlation between

oscillatory power and mind-wandering reports.

Cluster-based permutation testing revealed in the EO group, a significant positive

correlation between mind-wandering probability and alpha power over the 5 seconds preceding

the probe (11.5–13.5 Hz, –5 to 0 s, $p<0.001$, peak $t_{25}=3.904$ at 12.5 Hz and 0 s; Fig. 4 a,c), and

in the EC group, a significant negative correlation between mind-wandering probability and alpha

power over the 7 seconds preceding the probe (9–12.5 Hz, –7 to 0 s, $p<0.001$, peak $t_{23}=-3.825$

at 9.5 Hz and –7 s; Fig. 4 b,c), consistent with the inverted-U model.

Note that alpha power in the pre-probe window may be influenced by events in the

preceding trial. For example, a motor response in trial t may induce alpha-band desynchronization

in trial t + 1 (Pfurtscheller and Lopes Da Silva, 1999; Nikulin et al., 2007; Mazaheri and Jensen,

2008). Because mind-wandering reports are preceded by more button presses due to increased

false alarm rates, changes in pre-probe alpha power attributed to mind-wandering might instead

reflect differences in motor-related desynchronization, rather than endogenous fluctuations. To

rule out motor confounds, we repeated the GLM analysis including only pre-probe epochs

following correct go responses, all of which involved a button press in the preceding trial. This

control analysis replicated the positive correlation in the EO group ($p=0.021$) and the negative

correlation in the EC group ($p<0.001$), ruling out motor confounds.

The inverted-U model of mind-wandering was motivated by prior findings suggesting an

inverted-U relationship between alpha power and sleepiness, a state closely related to mind-

wandering (Jubera-García et al., 2021) characterized by positive and negative correlations during

eyes-open and eyes-closed states, respectively (Kaida et al., 2006). To replicate these findings,

we applied the same approach used for the mind-wandering analysis, fitting separate GLMs in

the EO and EC groups to test the block-wise relationships between ongoing oscillatory power and

sleepiness ratings. Cluster-based permutation testing revealed a significant positive correlation

between sleepiness and alpha power in the EO group (9.5–12 Hz, p=0.013, peak t_{25}=4.298 at

11.5 Hz; Fig. 4 d,f) and a significant negative correlation in the EC group (9–11.5 Hz, p=0.011,

peak t_{23}=-4.633, at 10 Hz; Fig. 4 e,f), consistent with previous results. The restricted GLMs

controlling for motor confounds replicated the positive correlation in EO group (p=0.021) and

negative correlation in EC group (p=0.006).

Taken together, these results confirm criterion (ii) of the inverted-U model: alpha power

shows a positive correlation with mind-wandering and sleepiness at low power ranges (EO group)

and a negative correlation at high power ranges (EC group), providing evidence for an inverted-

U relationship between alpha power and subjective reports.

Figure 4. Relationship between power and subjective reports of mind-wandering and
sleepiness.
Participants were probed to report whether they were on-task or mind-wandering during each task
block and to rate their sleepiness after each block. We used generalized linear models (GLMs) to
correlate fluctuations in ongoing spectral power with each of these subjective reports separately
in the eyes-open (EO) and eyes-closed (EC) groups. (a, b) Group-level t-statistics map the
correlation between pre-probe power and the probability of reporting mind-wandering for the EO
(a) and EC (b) groups, respectively. Blue indicates that mind-wandering became more likely with

decreasing pre-probe power; yellow indicates that mind-wandering became more likely with
increasing pre-probe power. Maps are masked to show cluster-corrected significant effects
(p<0.05). Time 0 s denotes probe onset. **(d, e)** *Group-level frequency-domain representations of*
the correlation between power and sleepiness ratings for the EO (d) and EC (e) groups. Significant
frequency clusters are indicated with black horizontal lines (cluster-based permutation test,
p<0.05). **(c, f)** *Descriptive visualization of the inverted-U across EO and EC groups. For each*
participant, power was first averaged across time-frequency (c) or frequency (f) points identified
as significant in either group by the cluster-based permutation tests shown in (a)/(b) for mind-
wandering ((c): 9–13.5 Hz, –7 to 0 s), and in (d)/(e) for sleepiness ((f): 9–12 Hz). Trials (mind-
wandering) or blocks (sleepiness) were median split into low and high power bins, and mind-
wandering probability (c) or mean KSS rating (f) was computed for each bin. To represent EO
and EC groups on a common axis, power and behavioral measures were expressed as percent
change relative to each participant's resting baseline (mean power across EO and EC rest) and
overall mean report, respectively. Error bars show group SEM (x: power; y: behavior). Panels (c)
and (f) are descriptive summaries of the results in panels (a)/(b), and (d)/(e), respectively, and
were not independently tested.

Alpha and beta power show a negative relationship with reaction times in the eyes-closed

group

Then, we examined the relationship between prestimulus alpha power and RTs to

replicate previous findings (e.g., ElShafei et al., 2018). We applied the same approach as for the

analysis of subjective reports, fitting separate GLMs for the EO and EC groups. Cluster-based

permutation testing in the EC group showed a significant negative correlation between RTs and

power in the alpha and beta bands within approximately one second before stimulus onset (6.5–

25 Hz, –1.1 to –0.3 s, p=0.018, peak t_{23}=–3.593 at 19 Hz and –0.9 s; Fig. 5 b,c), indicating that

stronger oscillations in alpha and beta bands over occipital and parietal scalp regions preceded

faster responses to auditory stimuli. By contrast, cluster-based permutation testing in the EO

group revealed no significant clusters (Fig. 5 a,c). Bayesian analysis at each time-frequency point

within the EC significant cluster showed that, in the EO group, 33% of points favored the null

hypothesis and 64% were inconclusive.

To rule out motor confounds, we repeated the GLM analysis using two subsets of epochs:

(i) epochs following correct go responses, which were always preceded by a button press, and

(ii) epochs following correct no-go responses, which were not preceded by a button press. These

control analyses replicated the inconclusive correlation in the EO group (no significant correlation

in the alpha band) and the negative correlation spanning alpha and beta bands in the EC group

(i: p=0.028; ii: p=0.007).

Note that average RT was not affected by mind-wandering (see *Behavioral Results*),

therefore these results cannot provide evidence for or against the inverted-U model of mind-

wandering. Instead, the results in the EC group replicate some previous findings showing that

RTs are negatively correlated with prestimulus alpha power from task-unrelated areas.

Figure 5. Relationship between prestimulus power and reaction times.
Group-level t statistics map the correlation coefficient reflecting the relationship between EEG
power and reaction times (RTs) in the (a) eyes-open (EO) and (b) eyes-closed (EC) groups. Blue
and yellow values indicate a negative and positive correlation between ongoing power and RT,
respectively. The maps are masked to show cluster-corrected significant effects (p<0.05). Time 0
s denotes stimulus onset. (c) Descriptive visualization of the EO and EC group comparison across
resting-state normalized data. For each participant, power was first averaged across the
significant points from the cluster-based permutation test of the EC group ((b): 6.5–25 Hz, –1.1
to –0.3 s). Trials were then median split into low vs high power bins, and average RT was
computed for each bin. To place EO and EC groups on a common axis, power and RT measures
were expressed as percent change relative to each participant's resting baseline (mean power
across EO and EC rest) and overall mean RT, respectively. Error bars show group SEM (x: power;
y: RT). (c) is a descriptive summary of the results in panels A and B and was not independently
tested.

Systematic Literature Review Supports an Inverted-U Relationship Between Alpha Power and Mind-Wandering

We conducted a systematic literature review of studies analyzing the relationship between the power of alpha oscillations and immediate mind-wandering reports in neurotypical individuals. We categorized studies based on whether the task was performed with eyes open (Table 2, top) or eyes closed (Table 2, bottom). We found that 17/24 eyes-open studies reported a positive relationship, 3/24 reported a negative relationship, and 4/24 reported no significant relationship. This shows that prior eyes-open studies are predominantly consistent with our results in the EO group and with the left side of the inverted-U. Our study replicated this positive relationship using an auditory task, whereas most eyes-open studies (16/24) used visual tasks. Notably, although the only eyes-open study that used a self-caught methodology reported a negative relationship, alpha power in that study was higher in a distant pre-report window labeled "mind-wandering" (–10 to –5 s) than in the immediately pre-report window labeled "awareness of mind-wandering" (–5 to 0 s) (Shinagawa et al., 2025). If meta-awareness reflects an on-task state, this would constitute a positive relationship consistent with most eyes-open studies, and with the left side of the inverted-U. All 5/5 eyes-closed studies reported a negative relationship, consistent with our results in the EC group and with the right side of the inverted-U. We additionally found that 23/24 prior eyes-open studies used exteroceptive tasks and probe-caught mind-wandering reports, whereas all eyes-closed studies used interoceptive tasks and most used self-caught mind-wandering reports. Accordingly, prior literature alone could not have determined whether the opposite relationships in eyes-open and eyes-closed studies reflect different sides of the inverted-U (compatible with our model), or different mechanisms related to the type of sustained attention task or reporting method used.

Reference	Relationship between alpha power and mind-wandering	Sustained attention task (modality)	Mind-wandering report method
Eyes-open paradigm			
Arnau et al. (2020)	Positive	Exteroceptive (V)	Probe
Baird et al. (2014)	Negative	Exteroceptive (V)	Probe
Baldwin et al. (2017)	Positive	Exteroceptive (V)	Probe
Broadway et al. (2015)	Null	Exteroceptive (V)	Probe
Compton et al. (2019)	Positive	Exteroceptive (V)	Probe
Compton et al. (2024)	Positive	Exteroceptive (V)	Probe
Conrad and Newman (2021)	Positive	Exteroceptive (AV)	Probe
Dhindsa et al. (2019)	Null	Exteroceptive (AV)	Probe
Dias Da Silva et al. (2022)	Positive	Exteroceptive (V)	Probe
Esposito et al. (2022)	Positive	Exteroceptive (V)	Probe
Gouraud et al. (2021)	Positive	Exteroceptive (AV)	Probe
Groot et al. (2021)	Positive	Exteroceptive (V)	Probe
Ibáñez-Molina and Iglesias-Parro (2014)	Null	Exteroceptive (AV)	Probe
Jin et al. (2019)	Positive	Exteroceptive (V)	Probe
Kam et al. (2021)	Positive	Exteroceptive (V)	Probe

Kam et al. (2025)	Positive	Exteroceptive (AV, A,V)	Probe
Kirschner et al. (2012)	Positive	Exteroceptive (V)	Probe
Macdonald et al. (2011)	Positive	Exteroceptive (V)	Probe
Polychroni et al. (2022)	Positive	Exteroceptive (A)	Probe
Prieto et al. (2021)	Negative	Exteroceptive (AV)	Probe
Qin et al. (2011)	Null	Exteroceptive (V)	Probe
Shinagawa et al. (2025)	Negative	Exteroceptive (A)	Self (*)
	Negative	Interoceptive (Breath)	Self
Wamsley and Summer (2020)	Positive	Exteroceptive (V)	Probe
Yang et al. (2022)	Positive	Exteroceptive (V)	Probe
Eyes-closed paradigm			
Braboszcz and Delorme (2011)	Negative	Interoceptive (Breath)	Self
Rodriguez-Larios and Alaerts (2021)	Negative	Interoceptive (Breath)	Probe
Rout et al. (2024)	Negative	Interoceptive (Breath)	Self
van Son et al. (2019a)	Negative	Interoceptive (Breath)	Self
van Son et al. (2019b)	Negative	Interoceptive (Breath)	Self

Table 2. Review of prior studies examining the relationship between alpha power and mind-wandering.

Studies using eyes-open and eyes-closed paradigms are categorized by sustained-attention task type, mind-wandering report method, and the direction of the observed relationship: "positive" means alpha power increases in mind-wandering relative to on-task states, "negative" means it decreases, and "null" means no reliable relationship. Exteroceptive tasks used visual (V), auditory (A), audiovisual (AV), or auditory and visual (A,V) modalities separately, while all interoceptive tasks involved breath counting (Breath). Probe-caught studies contrast activity preceding probes classified as mind-wandering versus on-task. Self-caught studies contrasted alpha power

Discussion

Alpha oscillations and mind-wandering

The literature on mind-wandering faces a paradox: mind-wandering is associated with stronger alpha power in some studies but weaker alpha power in others. We reasoned that an inverted-U relationship between alpha power and mind-wandering could dissipate this paradox, with studies sampling different alpha power ranges (sides of the inverted-U) reporting opposite effects. We further hypothesized that whether participants' eyes were open or closed could vary the range of alpha power explored by different studies. Accordingly, we found that mind-wandering was positively correlated with alpha power in the EO group, consistent with previous eyes-open studies using similar methods (Table 2, top) and the left side of the inverted-U. Eye closure (EC group) shifted alpha power into a higher range and reversed its correlation with mind-wandering, consistent with the right side of the inverted-U (Fig. 1).

Although previous eyes-closed studies reported a negative correlation (Table 2, bottom), whether it reflects the right side of the inverted-U or other methodological differences remains unclear. First, alpha power may show opposite effects in interoceptive and exteroceptive attention tasks (Cooper et al., 2003; Villena-González et al., 2017), though Shinagawa et al. (2025) found evidence against this. Second, eyes-closed studies assumed self-caught mind-wandering reports to be preceded by mind-wandering and followed by on-task states (Rout et al., 2024). However, if participants report mind-wandering only after becoming aware of it and returning on-task (Shinagawa et al., 2025), pre-report periods are mislabeled as "mind-wandering." If participants immediately resume mind-wandering post-report, these periods are mislabeled as "on-task" (Rout

et al., 2024). This mislabeling could reverse the sign of the observed correlation and limit

interpretability of self-caught studies.

By using the same exteroceptive task and probe-caught experience-sampling in both eye

states, our study demonstrates that the reversal of the correlation in the EC group arises from eye

closure. Future research might test if this reversal replicates independent of eye closure by

manipulating alpha power via perceptual deprivation (Ganzfeld: Miskovic et al., 2019),

neurofeedback (Hanslmayr et al., 2005; Brickwedde et al., 2019), and neurostimulation (Vossen

et al., 2015).

Taken together, our findings demonstrate that the paradoxical positive and negative

correlations between alpha oscillations and mind-wandering reported previously, reflect two sides

of an inverted-U relationship spanning alpha power ranging across eye states.

Alpha oscillations and sleepiness

We first tested if subjective sleepiness was related to mind-wandering (Jubera-García et

al., 2021). Participants mind-wandered more when they felt sleepier, and those with eyes closed

showed both higher sleepiness and more frequent mind-wandering than those with eyes open.

This is consistent with prior research showing that mind-wandering is more likely during states of

low arousal reflected in subjective reports (sleepiness: Stawarczyk and D'Argembeau, 2016;

Carciofo et al., 2014) and physiological metrics (reduced baseline pupil diameter: Unsworth and

Robison, 2018; Stawarczyk et al., 2020; Andrillon et al., 2021; but see Smallwood et al., 2012;

Groot et al., 2021; reduced stimulus-locked pupil dilation: Jubera-García et al., 2021; and strong

EEG slow waves: Andrillon et al., 2021). Further, increasing sleepiness via sleep deprivation

triggers mind-wandering (Poh et al., 2016; Walker and Trick, 2018; Yang et al., 2025) and, via

pharmacology, increases attentional lapses (Pinggal et al., 2022), suggesting a causal link.

We next examined whether subjective sleepiness showed an inverted-U relationship with

alpha power, as suggested by prior research. In the EO group, alpha power was positively

correlated with sleepiness ratings, corresponding to the left side of the inverted-U. This aligns with

previous eyes-open studies showing that alpha power increases with sleepiness measured

directly through subjective ratings (Stampi et al., 1995; Cajochen et al., 1996; Kaida et al., 2006;

Iemi et al., 2019), or indirectly through time-on-task (Benwell et al., 2019; Kopčanová et al., 2025).

By contrast, in the EC group, alpha power was negatively correlated with sleepiness, consistent

with the right side of the inverted-U. This replicates prior eyes-closed findings showing that alpha

power decreases with increasing subjective sleepiness (Stampi et al., 1995; Strijkstra et al., 2003;

Kaida et al., 2006; Snipes et al., 2024) and during sleep onset ("alpha dropout;" De Gennaro et

al., 2001; Lacaux et al., 2024).

Taken together, these findings show that subjective reports of sleepiness and mind-

wandering are behaviorally correlated and both exhibit an inverted-U relationship with alpha

power.

Alpha/Beta oscillations and reaction times

In our study, mean reaction times (RTs) didn't differ between mind-wandering and on-task

states in either group, inconsistent with our hypothesis (Leszczynski et al., 2017; Andrillon et al.,

2021; but see Kam et al., 2011; Arnau et al., 2020). Note that our study did not distinguish between

task-unrelated states with (i.e., mind-wandering) and without content (i.e., mind-blanking;

Andrillon et al., 2024). These states may produce opposite effects on RTs (Andrillon et al., 2021;

Munoz-Musat et al., 2025), so conflating them could yield null results.

We thus examined the relationship between prestimulus alpha power and RTs to replicate

previous findings. In the EC group, we found that stronger alpha/beta power (in occipito-parietal

electrodes) preceded faster (auditory) RTs. This is consistent with prior studies estimating

oscillations from parietal and visual regions, and RTs in non-visual tasks (e.g. somatosensory:

Linkenkaer-Hansen et al., 2004; ElShafei et al., 2022; auditory: Bollimunta et al., 2008; ElShafei

et al., 2018; Brickwedde et al., 2025). However, the EO group showed mostly inconclusive

evidence. This contrasts one study using source-level MEG, which likely boosts the signal-to-

noise ratio of alpha oscillations, showing a similar relationship in both eye states (ElShafei et al.,

2022). Thus, the effects on RTs, unlike subjective reports, are likely incompatible with an inverted-

U across eye states. Importantly, because mean RTs didn't track mind-wandering in our study,

these results cannot provide evidence for or against the inverted-U model of mind-wandering.

Mechanistic Interpretations

While this study doesn't directly address why the inverted-U exists, by clarifying existing

literature it provides a necessary first step toward understanding underlying mechanisms. We

consider two mechanistic interpretations post-hoc.

First, alpha fluctuations during mind-wandering may reflect changes in arousal (see *Alpha*

oscillations and sleepiness). Accordingly, the inverted-U we observed is compatible with mind-

wandering occurring in states of decreased arousal indexed by (i) higher eyes-open alpha power

(left side of the inverted-U) and (ii) lower eyes-closed alpha power (right side of the inverted-U).

Decreased arousal is thought to induce a state of perceptual decoupling, reflected in dampened

physiological responses to external sensory stimuli: for example, reduced event-related potentials

at sleep onset (Harsh et al., 1994; Ogilvie, 2001; Andrillon and Kouider, 2020) and during mind-

wandering (Baird et al., 2014), as well as reduced stimulus-locked pupil responses during mind-

wandering (Jubera-García et al., 2020). Importantly, unlike selective attention, perceptual

decoupling is thought to attenuate processing at all sensory sources, affecting task-related and

task-unrelated information equally (Barron et al., 2011; Schooler et al., 2011). At the experiential

level, this attenuation may yield mental states with either no content at all (mind-blanking) or with

stimulus-independent content (i.e., unrelated to the immediate sensory environment; Boulakis et

al., 2023).

Second, alpha fluctuations during mind-wandering may reflect shifts in selective attention

between task-related and task-unrelated sensory information (Mazaheri et al., 2014; Morrow et

al., 2023). According to functional inhibition, alpha oscillations modulate the excitability of

neurons, locally enhancing or suppressing information processing within those neurons (Klimesch

et al., 2007; Haegens et al., 2011; Iemi et al., 2022). Thus, the inverted-U may be compatible with

(i) eyes-open alpha power inhibiting task-related sensory sources (likely reflecting mixed

sources), shifting attention to non-auditory stimuli, and thereby increasing mind-wandering (left

side of the inverted-U) and (ii) eyes-closed alpha power (likely reflecting a visual source) inhibiting

task-unrelated sensory sources, shifting attention to auditory stimuli, and thereby decreasing

mind-wandering (right side of the inverted-U).

An outstanding question is how to reconcile the inverted-U relationship between alpha

power and arousal with the functional inhibition account. Strong alpha power in eyes-open states

(left side of the inverted-U) reflects decreased arousal and plausibly induces perceptual

decoupling through its known inhibitory effects (Iemi et al., 2019, 2022). However, the right side

of the inverted-U challenges this view: in eyes-closed states, strong alpha power reflects

increased arousal (Kaida et al., 2006) and excitability (de Graaf et al., 2017). Future work is

needed to clarify the underlying neurophysiological mechanisms.

This study cannot adjudicate between these interpretations due to some limitations. First,

the limited spatial resolution of our EEG recordings prevents us from determining whether eyes-

open and eyes-closed alpha oscillations arise from different sensory sources (interpretation 2) or

a global/common signal (interpretation 1). Second, because we did not probe content, we cannot

determine whether changes in alpha power relate to mind-wandering about stimuli represented

in the corresponding task-unrelated sensory sources (interpretation 2) or a mixture between mind-

blanking and stimulus-independent mind-wandering (interpretation 1). Future studies could

address this with higher-resolution recordings and more fine-grained experience-sampling

methods.

In conclusion, these results demonstrate a relationship between mind-wandering and

alpha oscillations that follows an inverted-U across eye states, with eye closure shifting alpha

power into a higher range and reversing its correlation with mind-wandering. By considering eye

state, this study clarifies contradictions in prior literature and specifies how ongoing neural

oscillations shape subjective experience.

References

Andrillon, T., Windt, J., Silk, T., Drummond, S. P. A., Bellgrove, M. A., & Tsuchiya, N.

(2019). Does the Mind Wander When the Brain Takes a Break? Local Sleep in

Wakefulness, Attentional Lapses and Mind-Wandering. *Frontiers in Neuroscience*, *13*,

949. https://doi.org/10.3389/fnins.2019.00949

Andrillon, T., and Kouider, S. (2020). The vigilant sleeper: Neural mechanisms of sensory

(de)coupling during sleep. *Current Opinion in Physiology*, *15*, 47–59.

https://doi.org/10.1016/j.cophys.2019.12.002

Andrillon, T., Burns, A., Mackay, T., Windt, J., & Tsuchiya, N. (2021). Predicting lapses of

attention with sleep-like slow waves. *Nature Communications*, *12*(1), 3657.

https://doi.org/10.1038/s41467-021-23890-7

Andrillon, T., Lutz, A., Windt, J., & Demertzi, A. (2024). *Where is my Mind?: A*

Neurocognitive Investigation of Mind Blanking. PsyArXiv.

https://doi.org/10.31234/osf.io/xmtga

Arnau, S., Löffler, C., Rummel, J., Hagemann, D., Wascher, E., & Schubert, A.-L. (2020).

Inter-trial alpha power indicates mind wandering. *Psychophysiology*, *57*(6), e13581.

https://doi.org/10.1111/psyp.13581

Baird, B., Smallwood, J., Lutz, A., & Schooler, J. W. (2014). The Decoupled Mind: Mind-

wandering Disrupts Cortical Phase-locking to Perceptual Events. *Journal of Cognitive*

Neuroscience, *26*(11), 2596–2607. https://doi.org/10.1162/jocn_a_00656

Baldwin, C. L., Roberts, D. M., Barragan, D., Lee, J. D., Lerner, N., & Higgins, J. S. (2017).

Detecting and Quantifying Mind Wandering during Simulated Driving. *Frontiers in*

Human Neuroscience, *11*, 406. https://doi.org/10.3389/fnhum.2017.00406

Barron, E., Riby, L. M., Greer, J., & Smallwood, J. (2011). Absorbed in Thought: The Effect

of Mind Wandering on the Processing of Relevant and Irrelevant Events. *Psychological*

Science, *22*(5), 596–601. https://doi.org/10.1177/0956797611404083

Barry, R. J., Clarke, A. R., Johnstone, S. J., Magee, C. A., & Rushby, J. A. (2007). EEG

differences between eyes-closed and eyes-open resting conditions. *Clinical*

Neurophysiology, *118*(12), 2765–2773. https://doi.org/10.1016/j.clinph.2007.07.028

Bastian, M., and Sackur, J. (2013). Mind wandering at the fingertips: Automatic parsing of

subjective states based on response time variability. *Frontiers in Psychology*, *4*.

https://doi.org/10.3389/fpsyg.2013.00573

Benwell, C. S. Y., London, R. E., Tagliabue, C. F., Veniero, D., Gross, J., Keitel, C., & Thut,

G. (2019). Frequency and power of human alpha oscillations drift systematically with

time-on-task. *NeuroImage*, *192*, 101–114.

https://doi.org/10.1016/j.neuroimage.2019.02.067

Berger, H. (1929) Über das elektrenkephalogramm des menschen. European Archives of

Psychiatry and Clinical Neuroscience, 87, 527-570.

https://doi.org/10.1007/BF01797193

Bollimunta, A., Chen, Y., Schroeder, C. E., & Ding, M. (2008). Neuronal Mechanisms of

Cortical Alpha Oscillations in Awake-Behaving Macaques. *The Journal of*

Neuroscience, *28*(40), 9976–9988. https://doi.org/10.1523/JNEUROSCI.2699-08.2008

Boudewyn, M. A., and Carter, C. S. (2018). I must have missed that: Alpha-band

oscillations track attention to spoken language. Neuropsychologia, 117, 148–155.

https://doi.org/10.1016/j.neuropsychologia.2018.05.024

Boulakis, P. A., Mortaheb, S., Calster, L. van, Majerus, S., & Demertzi, A. (2023). Whole-

Brain Deactivations Precede Uninduced Mind-Blanking Reports. *Journal of*

Neuroscience, *43*(40), 6807–6815. https://doi.org/10.1523/JNEUROSCI.0696-23.2023

Braboszcz, C., and Delorme, A. (2011). Lost in thoughts: Neural markers of low alertness

during mind wandering. *NeuroImage*, *54*(4), 3040–3047.

https://doi.org/10.1016/j.neuroimage.2010.10.008

Brainard, D. H. (1997). The Psychophysics Toolbox. *Spatial Vision*, *10*(4), 433–436.

https://doi.org/10.1163/156856897X00357

Brickwedde, M., Krüger, M. C., & Dinse, H. R. (2019). Somatosensory alpha oscillations

gate perceptual learning efficiency. *Nature Communications*, *10*(1), 263.

https://doi.org/10.1038/s41467-018-08012-0

Brickwedde, M., Limachya, R., Markiewicz, R., Sutton, E., Postzich, C., Shapiro, K.,

Jensen, O., & Mazaheri, A. (2025). *Cross-modal interaction of Alpha Activity does not*

reflect inhibition of early sensory processing: A frequency tagging study using EEG and

MEG. https://doi.org/10.7554/eLife.106050.3

Broadway, J. M., Franklin, M. S., & Schooler, J. W. (2015). Early event-related brain

potentials and hemispheric asymmetries reveal mind-wandering while reading and

predict comprehension. *Biological Psychology*, *107*, 31–43.

https://doi.org/10.1016/j.biopsycho.2015.02.009

Cajochen, C., Kräuchi, K., Von Arx, M.-A., Möri, D., Graw, P., & Wirz-Justice, A. (1996).

Daytime melatonin administration enhances sleepiness and theta/alpha activity in the

waking EEG. *Neuroscience Letters*, *207*(3), 209–213. https://doi.org/10.1016/0304-

3940(96)12517-9

Carciofo, R., Du, F., Song, N., & Zhang, K. (2014). Mind Wandering, Sleep Quality, Affect

and Chronotype: An Exploratory Study. *PLoS ONE*, *9*(3), e91285.

https://doi.org/10.1371/journal.pone.0091285

Cloud Text-to-Speech API V1. (n.d.). Google Cloud Documentation. Retrieved November 11, 2024, from https://docs.cloud.google.com/text-to-speech/docs/reference/rest

Compton, R. J., Gearinger, D., & Wild, H. (2019). The wandering mind oscillates: EEG alpha power is enhanced during moments of mind-wandering. *Cognitive, Affective, & Behavioral Neuroscience*, *19*(5), 1184–1191. https://doi.org/10.3758/s13415-019-00745-9

Compton, R. J., Shudrenko, D., Mann, K., Turdukulov, E., Ng, E., & Miller, L. (2024). Effects of task context on EEG correlates of mind-wandering. *Cognitive, Affective, & Behavioral Neuroscience*, *24*(1), 72–86. https://doi.org/10.3758/s13415-023-01138-9

Conrad, C., and Newman, A. (2021). Measuring Mind Wandering During Online Lectures Assessed With EEG. *Frontiers in Human Neuroscience*, *15*. https://doi.org/10.3389/fnhum.2021.697532

Cooper, N. R., Croft, R. J., Dominey, S. J. J., Burgess, A. P., & Gruzelier, J. H. (2003). Paradox lost? Exploring the role of alpha oscillations during externally vs. internally directed attention and the implications for idling and inhibition hypotheses. *International Journal of Psychophysiology*, *47*(1), 65–74. https://doi.org/10.1016/S0167-8760(02)00107-1

De Gennaro, L., Ferrara, M., & Bertini, M. (2001). The boundary between wakefulness and sleep: Quantitative electroencephalographic changes during the sleep onset period. *Neuroscience*, *107*(1), 1–11. https://doi.org/10.1016/S0306-4522(01)00309-8

De Graaf, T. A., Duecker, F., Stankevich, Y., Ten Oever, S., & Sack, A. T. (2017). Seeing in the dark: Phosphene thresholds with eyes open versus closed in the absence of visual inputs. *Brain Stimulation*, *10*(4), 828–835. https://doi.org/10.1016/j.brs.2017.04.127

Dhindsa, K., Acai, A., Wagner, N., Bosynak, D., Kelly, S., Bhandari, M., Petrisor, B., & Sonnadara, R. R. (2019). Individualized pattern recognition for detecting mind wandering from EEG during live lectures. *PLOS ONE, 14*(9), e0222276. https://doi.org/10.1371/journal.pone.0222276

Dias Da Silva, M. R., Gonçalves, Ó. F., Branco, D., & Postma, M. (2022). Revisiting consciousness: Distinguishing between states of conscious focused attention and mind wandering with EEG. *Consciousness and Cognition, 101*, 103332. https://doi.org/10.1016/j.concog.2022.103332

ElShafei, H. A., Bouet, R., Bertrand, O., & Bidet-Caulet, A. (2018). Two Sides of the Same Coin: Distinct Sub-Bands in the α Rhythm Reflect Facilitation and Suppression Mechanisms during Auditory Anticipatory Attention. *Eneuro, 5*(4), ENEURO.0141-18.2018. https://doi.org/10.1523/ENEURO.0141-18.2018

ElShafei, H. A., Orlemann, C., & Haegens, S. (2022). The Impact of Eye Closure on Anticipatory α Activity in a Tactile Discrimination Task. *Eneuro, 9*(1), ENEURO.0412-21.2021. https://doi.org/10.1523/ENEURO.0412-21.2021

Esposito, A., Braccili, E., Sgro, F., Chiarantano, E., D'Ippolito, M., Pisotta, I., Bigioni, A., Guerrieri, A., Mattia, D., & Cincotti, F. (2022). Attention, Boredom and Mind Wandering during a Vigilance Task: EEG and Ocular Markers. *2022 IEEE International Conference on Metrology for Extended Reality, Artificial Intelligence and Neural Engineering (MetroXRAINE)*, 477–482. https://doi.org/10.1109/MetroXRAINE54828.2022.9967678

Gouraud, J., Delorme, A., & Berberian, B. (2021). Mind Wandering Influences EEG Signal in Complex Multimodal Environments. *Frontiers in Neuroergonomics, 2*, 625343. https://doi.org/10.3389/fnrgo.2021.625343

Groot, J. M., Boayue, N. M., Csifcsák, G., Boekel, W., Huster, R., Forstmann, B. U., &

Mittner, M. (2021). Probing the neural signature of mind wandering with simultaneous

fMRI-EEG and pupillometry. NeuroImage, 224, 117412.

https://doi.org/10.1016/j.neuroimage.2020.117412

Haegens, S., Nácher, V., Luna, R., Romo, R., & Jensen, O. (2011). α-Oscillations in the

monkey sensorimotor network influence discrimination performance by rhythmical

inhibition of neuronal spiking. *Proceedings of the National Academy of Sciences of the*

United States of America, *108*(48), 19377–19382.

https://doi.org/10.1073/pnas.1117190108

Hanslmayr, S., Sauseng, P., Doppelmayr, M., Schabus, M., & Klimesch, W. (2005).

Increasing Individual Upper Alpha Power by Neurofeedback Improves Cognitive

Performance in Human Subjects. *Applied Psychophysiology and Biofeedback*, *30*(1),

1–10. https://doi.org/10.1007/s10484-005-2169-8

Harsh, J., Voss, U., Hull, J., Schrepfer, S., & Badia, P. (1994). ERP and behavioral

changes during the wake/sleep transition. *Psychophysiology*, *31*(3), 244–252.

https://doi.org/10.1111/j.1469-8986.1994.tb02213.x

Hohaia, W., Saurels, B. W., Johnston, A., Yarrow, K., & Arnold, D. H. (2022). Occipital

alpha-band brain waves when the eyes are closed are shaped by ongoing visual

processes. *Scientific Reports*, *12*(1), 1194. https://doi.org/10.1038/s41598-022-05289-

6

Hua, J., Wolff, A., Zhang, J., Yao, L., Zang, Y., Luo, J., Ge, X., Liu, C., & Northoff, G.

(2022). Alpha and theta peak frequency track on- and off-thoughts. Communications

Biology, 5(1), 209. https://doi.org/10.1038/s42003-022-03146-w

Ibáñez-Molina, A. J., and Iglesias-Parro, S. (2014). Fractal characterization of internally and

externally generated conscious experiences. *Brain and Cognition*, *87*, 69–75.

https://doi.org/10.1016/j.bandc.2014.03.002

Iemi, L., Chaumon, M., Crouzet, S. M., & Busch, N. A. (2017). Spontaneous Neural

Oscillations Bias Perception by Modulating Baseline Excitability. *The Journal of*

Neuroscience, *37*(4), 807–819. https://doi.org/10.1523/JNEUROSCI.1432-16.2016

Iemi, L., and Busch, N. A. (2018). Moment-to-Moment Fluctuations in Neuronal Excitability

Bias Subjective Perception Rather than Strategic Decision-Making. *Eneuro*, *5*(3),

ENEURO.0430-17.2018. https://doi.org/10.1523/ENEURO.0430-17.2018

Iemi, L., Busch, N. A., Laudini, A., Haegens, S., Samaha, J., Villringer, A., & Nikulin, V. V.

(2019). Multiple mechanisms link prestimulus neural oscillations to sensory responses.

eLife, *8*, e43620. https://doi.org/10.7554/eLife.43620

Iemi, L., Gwilliams, L., Samaha, J., Auksztulewicz, R., Cycowicz, Y. M., King, J.-R., Nikulin,

V. V., Thesen, T., Doyle, W., Devinsky, O., Schroeder, C. E., Melloni, L., & Haegens,

S. (2022). Ongoing neural oscillations influence behavior and sensory representations

by suppressing neuronal excitability. *NeuroImage*, *247*, 118746.

https://doi.org/10.1016/j.neuroimage.2021.118746

Jin, C. Y., Borst, J. P., & van Vugt, M. K. (2019). Predicting task-general mind-wandering

with EEG. *Cognitive, Affective, & Behavioral Neuroscience*, *19*(4), 1059–1073.

https://doi.org/10.3758/s13415-019-00707-1

Jubera-García, E., Gevers, W., & Van Opstal, F. (2020). Influence of content and intensity

of thought on behavioral and pupil changes during active mind-wandering, off-focus,

and on-task states. *Attention, Perception, & Psychophysics*, *82*(3), 1125–1135.

https://doi.org/10.3758/s13414-019-01865-7

Jubera-García, E., Gevers, W., & Van Opstal, F. (2021). Local build-up of sleep pressure

could trigger mind wandering: Evidence from sleep, circadian and mind wandering

research. *Biochemical Pharmacology*, *191*, 114478.

https://doi.org/10.1016/j.bcp.2021.114478

Kaida, K., Takahashi, M., Åkerstedt, T., Nakata, A., Otsuka, Y., Haratani, T., & Fukasawa,

K. (2006). Validation of the Karolinska sleepiness scale against performance and EEG

variables. *Clinical Neurophysiology*, *117*(7), 1574–1581.

https://doi.org/10.1016/j.clinph.2006.03.011

Kam, J. W. Y., Dao, E., Farley, J., Fitzpatrick, K., Smallwood, J., Schooler, J. W., & Handy,

T. C. (2011). Slow Fluctuations in Attentional Control of Sensory Cortex. *Journal of*

Cognitive Neuroscience, *23*(2), 460–470. https://doi.org/10.1162/jocn.2010.21443

Kam, J. W. Y., Irving, Z. C., Mills, C., Patel, S., Gopnik, A., & Knight, R. T. (2021). Distinct

electrophysiological signatures of task-unrelated and dynamic thoughts. *Proceedings*

of the National Academy of Sciences, *118*(4), e2011796118.

https://doi.org/10.1073/pnas.2011796118

Kam, J. W. Y., Rahnuma, T., Park, Y. E., & Hart, C. M. (2022). Electrophysiological

markers of mind wandering: A systematic review. *NeuroImage*, *258*, 119372.

https://doi.org/10.1016/j.neuroimage.2022.119372

Kam, J. W. Y., Rahnuma, T., Nanjappan Jothiraj, S., Ouellette-Zuk, A. A., & Knight, R. T.

(2025). Electrophysiological signatures of ongoing thoughts during naturalistic

behavior. *Imaging Neuroscience*, *3*, IMAG.a.20. https://doi.org/10.1162/IMAG.a.20

Kirschner, A., Kam, J. W. Y., Handy, T. C., & Ward, L. M. (2012). Differential

Synchronization in Default and Task-Specific Networks of the Human Brain. Frontiers

in Human Neuroscience, 6. https://doi.org/10.3389/fnhum.2012.00139

Klimesch, W., Sauseng, P., & Hanslmayr, S. (2007). EEG alpha oscillations: The inhibition-

860   timing hypothesis. *Brain Research Reviews*, *53*(1), 63–88.

861   https://doi.org/10.1016/j.brainresrev.2006.06.003

862   Kopčanová, M., Thut, G., Benwell, C. S. Y., & Keitel, C. (2025). Characterising time-on-task

863   effects on oscillatory and aperiodic EEG components and their co-variation with visual

864   task performance. *Imaging Neuroscience*, *3*, imag_a_00566.

865   https://doi.org/10.1162/imag_a_00566

866   Krekelberg, B. (2021). *klabhub/bayesFactor: Ttest updates* [Computer software]. Zenodo.

867   https://doi.org/10.5281/zenodo.5707551

868   Kucyi, A., Anderson, N., Bounyarith, T., Braun, D., Shareef-Trudeau, L., Treves, I., Braga,

869   R. M., Hsieh, P.-J., & Hung, S.-M. (2024). Individual variability in neural

870   representations of mind-wandering. *Network Neuroscience*, *8*(3), 808–836.

871   https://doi.org/10.1162/netn_a_00387

872   Lacaux, C., Strauss, M., Bekinschtein, T. A., & Oudiette, D. (2024). Embracing sleep-onset

873   complexity. *Trends in Neurosciences*, *47*(4), 273–288.

874   https://doi.org/10.1016/j.tins.2024.02.002

875   Leszczynski, M., Chaieb, L., Reber, T. P., Derner, M., Axmacher, N., & Fell, J. (2017). Mind

876   wandering simultaneously prolongs reactions and promotes creative incubation.

877   *Scientific Reports*, *7*(1), 10197. https://doi.org/10.1038/s41598-017-10616-3

878   Linkenkaer-Hansen, K., Nikulin, V. V., Palva, S., Ilmoniemi, R. J., & Palva, J. M. (2004).

879   Prestimulus Oscillations Enhance Psychophysical Performance in Humans. *The*

880   *Journal of Neuroscience*, *24*(45), 10186–10190.

881   https://doi.org/10.1523/JNEUROSCI.2584-04.2004

Macdonald, J. S. P., Mathan, S., & Yeung, N. (2011). Trial-by-Trial Variations in Subjective

Attentional State are Reflected in Ongoing Prestimulus EEG Alpha Oscillations.

Frontiers in Psychology, *2*. https://doi.org/10.3389/fpsyg.2011.00082

Maris, E., and Oostenveld, R. (2007). Nonparametric statistical testing of EEG- and MEG-

data. *Journal of Neuroscience Methods*, *164*(1), 177–190.

https://doi.org/10.1016/j.jneumeth.2007.03.024

Mazaheri, A., and Jensen, O. (2008). Asymmetric Amplitude Modulations of Brain

Oscillations Generate Slow Evoked Responses. *Journal of Neuroscience*, *28*(31),

7781–7787. https://doi.org/10.1523/JNEUROSCI.1631-08.2008

Mazaheri, A., Van Schouwenburg, M. R., Dimitrijevic, A., Denys, D., Cools, R., & Jensen,

O. (2014). Region-specific modulations in oscillatory alpha activity serve to facilitate

processing in the visual and auditory modalities. *NeuroImage*, *87*, 356–362.

https://doi.org/10.1016/j.neuroimage.2013.10.052

Morrow, A., Elias, M., & Samaha, J. (2023). Evaluating the Evidence for the Functional

Inhibition Account of Alpha-band Oscillations during Preparatory Attention. *Journal of*

Cognitive Neuroscience, *35*(8), 1195–1211. https://doi.org/10.1162/jocn_a_02009

Munoz-Musat, E., Le Coz, A., Corcoran, A. W., Belloli, L., Naccache, L., & Andrillon, T.

(2025). Behavioral, experiential, and physiological signatures of mind blanking.

Proceedings of the National Academy of Sciences, 122(52), e2510262122.

https://doi.org/10.1073/pnas.2510262122

Miskovic, V., Bagg, J. O., Ríos, M., & Pouliot, J. J. (2019). Electrophysiological and

phenomenological effects of short-term immersion in an altered sensory environment.

Consciousness and Cognition, 70, 39–49.

https://doi.org/10.1016/j.concog.2019.02.003

Nakatani, C., Bernhard, H., & Van Leeuwen, C. (2024). Prior EEG marks focused and

mind-wandering mental states across trials. *Cerebral Cortex*, *34*(10), bhae403.

https://doi.org/10.1093/cercor/bhae403

Neuroelectrics. (n.d.) Enobio 8. [EEG Device]. Barcelona, Spain.

https://www.neuroelectrics.com/products/research/enobio/enobio8

Nikulin, V. V., Linkenkaer-Hansen, K., Nolte, G., Lemm, S., Müller, K. R., Ilmoniemi, R. J., &

Curio, G. (2007). A novel mechanism for evoked responses in the human brain.

European Journal of Neuroscience, *25*(10), 3146–3154. https://doi.org/10.1111/j.1460-

9568.2007.05553.x

Ogilvie, R. D. (2001). The process of falling asleep. *Sleep Medicine Reviews*, *5*(3), 247–

270. https://doi.org/10.1053/smrv.2001.0145

Oostenveld, R., Fries, P., Maris, E., & Schoffelen, J.-M. (2011). FieldTrip: Open Source

Software for Advanced Analysis of MEG, EEG, and Invasive Electrophysiological Data.

Computational Intelligence and Neuroscience, *2011*(1), 156869.

https://doi.org/10.1155/2011/156869

Petro, N. M., Ott, L. R., Penhale, S. H., Rempe, M. P., Embury, C. M., Picci, G., Wang, Y.-

P., Stephen, J. M., Calhoun, V. D., & Wilson, T. W. (2022). Eyes-closed versus eyes-

open differences in spontaneous neural dynamics during development. *NeuroImage*,

258, 119337. https://doi.org/10.1016/j.neuroimage.2022.119337

Pfurtscheller, G., and Lopes Da Silva, F. H. (1999). Event-related EEG/MEG

synchronization and desynchronization: Basic principles. *Clinical Neurophysiology*,

110(11), 1842–1857. https://doi.org/10.1016/S1388-2457(99)00141-8

Pinggal, E., Dockree, P. M., O'Connell, R. G., Bellgrove, M. A., & Andrillon, T. (2022).

Pharmacological Manipulations of Physiological Arousal and Sleep-Like Slow Waves

930    Modulate Sustained Attention. *The Journal of Neuroscience, 42*(43), 8113–8124.

931    https://doi.org/10.1523/JNEUROSCI.0836-22.2022

932 Poh, J.-H., Chong, P. L. H., & Chee, M. W. L. (2016). Sleepless night, restless mind:

933    Effects of sleep deprivation on mind wandering. Journal of Experimental Psychology:

934    General, 145(10), 1312–1318. https://doi.org/10.1037/xge0000207

935 Polychroni, N., Herrojo Ruiz, M., & Terhune, D. B. (2022). Introspection confidence predicts

936    EEG decoding of self-generated thoughts and meta-awareness. *Human Brain*

937    *Mapping, 43*(7), 2311–2327. https://doi.org/10.1002/hbm.25789

938 Prieto, M., Iglesias-Parro, S., Soriano, M. F., & Ibáñez-Molina, A. (2021). Local Oscillatory

939    Brain Dynamics of Mind Wandering in Schizophrenia. *Brain Sciences, 11*(7), 910.

940    https://doi.org/10.3390/brainsci11070910

941 Qin, J., Perdoni, C., & He, B. (2011). Dissociation of Subjectively Reported and

942    Behaviorally Indexed Mind Wandering by EEG Rhythmic Activity. *PLOS ONE, 6*(9),

943    e23124. https://doi.org/10.1371/journal.pone.0023124

944 Rodriguez-Larios, J., and Alaerts, K. (2021). EEG alpha–theta dynamics during mind

945    wandering in the context of breath focus meditation: An experience sampling approach

946    with novice meditation practitioners. *European Journal of Neuroscience, 53*(6), 1855–

947    1868. https://doi.org/10.1111/ejn.15073

948 Rouder, J. N., Speckman, P. L., Sun, D., Morey, R. D., & Iverson, G. (2009). Bayesian t

949    tests for accepting and rejecting the null hypothesis. *Psychonomic Bulletin & Review*,

950    *16*(2), 225–237. https://doi.org/10.3758/PBR.16.2.225

951 Rout, T. W., Duda, A. T., De Blasio, F. M., Clarke, A. R., & Barry, R. J. (2024). Global alpha

952    power fluctuations in a self-caught mind-wandering paradigm are independent of

953    arousal. *International Journal of Psychophysiology, 205*, 112428.

954    https://doi.org/10.1016/j.ijpsycho.2024.112428

Samaha, J., Iemi, L., & Postle, B. R. (2017). Prestimulus alpha-band power biases visual

discrimination confidence, but not accuracy. *Consciousness and Cognition*, *54*, 47–55.

https://doi.org/10.1016/j.concog.2017.02.005

Schooler, J. W., Smallwood, J., Christoff, K., Handy, T. C., Reichle, E. D., & Sayette, M. A.

(2011). Meta-awareness, perceptual decoupling and the wandering mind. Trends in

Cognitive Sciences, 15(7), 319–326. https://doi.org/10.1016/j.tics.2011.05.006

Schubert, A.-L., Frischkorn, G. T., & Rummel, J. (2020). The validity of the online thought-

probing procedure of mind wandering is not threatened by variations of probe rate and

probe framing. *Psychological Research*, *84*(7), 1846–1856.

https://doi.org/10.1007/s00426-019-01194-2

Shinagawa, K., Tanaka, Y., Terasawa, Y., & Umeda, S. (2025). Brain–body interactions

associated with the transition from mind wandering to awareness of its occurrence.

Neuroscience of Consciousness, 2025(1), niaf059. https://doi.org/10.1093/nc/niaf059

Simonsohn, U. (2018). Two Lines: A Valid Alternative to the Invalid Testing of U-Shaped

Relationships With Quadratic Regressions. *Advances in Methods and Practices in*

Psychological Science, *1*(4), 538–555. https://doi.org/10.1177/2515245918805755

Smallwood, J., Brown, K. S., Baird, B., Mrazek, M. D., Franklin, M. S., & Schooler, J. W.

(2012). Insulation for Daydreams: A Role for Tonic Norepinephrine in the Facilitation of

Internally Guided Thought. *PLOS ONE*, *7*(4), e33706.

https://doi.org/10.1371/journal.pone.0033706

Smallwood, J., and Schooler, J. W. (2015). The Science of Mind Wandering: Empirically

Navigating the Stream of Consciousness. *Annual Review of Psychology*, *66*(1), 487–

518. https://doi.org/10.1146/annurev-psych-010814-015331

Smulders, F. T. Y., ten Oever, S., Donkers, F. C. L., Quaedflieg, C. W. E. M., & van de Ven,

V. (2018). Single-trial log transformation is optimal in frequency analysis of resting

EEG alpha. *European Journal of Neuroscience, 48*(7), 2585–2598.

https://doi.org/10.1111/ejn.13854

Snipes, S., Meier, E., Accascina, S., & Huber, R. (2024). Extended wakefulness alters the

relationship between EEG oscillations and performance in a sustained attention task.

Journal of Sleep Research, 33(6), e14230. https://doi.org/10.1111/jsr.14230

Stampi, C., Stone, P., & Michimori, A. (1995). A new quantitative method for assessing

sleepiness: The alpha attenuation test. *Work & Stress, 9*(2–3), 368–376.

https://doi.org/10.1080/02678379508256574

Stawarczyk, D., and D'Argembeau, A. (2016). Conjoint influence of mind-wandering and

sleepiness on task performance. Journal of Experimental Psychology: Human

Perception and Performance, 42(10), 1587–1600. https://doi.org/10.1037/xhp0000254

Stawarczyk, D., François, C., Wertz, J., & D'Argembeau, A. (2020). Drowsiness or mind-

wandering? Fluctuations in ocular parameters during attentional lapses. Biological

Psychology, 156, 107950. https://doi.org/10.1016/j.biopsycho.2020.107950

Strijkstra, A. M., Beersma, D. G. M., Drayer, B., Halbesma, N., & Daan, S. (2003).

Subjective sleepiness correlates negatively with global alpha (8–12 Hz) and positively

with central frontal theta (4–8 Hz) frequencies in the human resting awake

electroencephalogram. *Neuroscience Letters, 340*(1), 17–20.

https://doi.org/10.1016/S0304-3940(03)00033-8

Tenke, C. E., Kayser, J., Abraham, K., Alvarenga, J. E., & Bruder, G. E. (2015). Posterior

EEG alpha at rest and during task performance: Comparison of current source density

and field potential measures. *International Journal of Psychophysiology, 97*(3), 299–

309. https://doi.org/10.1016/j.ijpsycho.2015.05.011

Unsworth, N., and Robison, M. K. (2018). Tracking arousal state and mind wandering with

pupillometry. *Cognitive, Affective, & Behavioral Neuroscience*, *18*(4), 638–664.

https://doi.org/10.3758/s13415-018-0594-4

van Son, D., De Blasio, F. M., Fogarty, J. S., Angelidis, A., Barry, R. J., & Putman, P.

(2019a). Frontal EEG theta/beta ratio during mind wandering episodes. *Biological*

Psychology, *140*, 19–27. https://doi.org/10.1016/j.biopsycho.2018.11.003

van Son, D., de Rover, M., De Blasio, F. M., van der Does, W., Barry, R. J., & Putman, P.

(2019b). Electroencephalography theta/beta ratio covaries with mind wandering and

functional connectivity in the executive control network. *Annals of the New York*

Academy of Sciences, *1452*(1), 52–64. https://doi.org/10.1111/nyas.14180

Villena-González, M., Moënne-Loccoz, C., Lagos, R. A., Alliende, L. M., Billeke, P., Aboitiz,

F., López, V., & Cosmelli, D. (2017). Attending to the heart is associated with posterior

alpha band increase and a reduction in sensitivity to concurrent visual stimuli.

Psychophysiology, *54*(10), 1483–1497. https://doi.org/10.1111/psyp.12894

Vossen, A., Gross, J., & Thut, G. (2015). Alpha Power Increase After Transcranial

Alternating Current Stimulation at Alpha Frequency (α-tACS) Reflects Plastic Changes

Rather Than Entrainment. *Brain Stimulation*, *8*(3), 499–508.

https://doi.org/10.1016/j.brs.2014.12.004

Walker, H. E. K., and Trick, L. M. (2018). Mind-wandering while driving: The impact of

fatigue, task length, and sustained attention abilities. *Transportation Research Part F:*

Traffic Psychology and Behaviour, *59*, 81–97. https://doi.org/10.1016/j.trf.2018.08.009

Wamsley, E. J., and Summer, T. (2020). Spontaneous Entry into an "Offline" State during

Wakefulness: A Mechanism of Memory Consolidation? *Journal of Cognitive*

Neuroscience, *32*(9), 1714–1734. https://doi.org/10.1162/jocn_a_01587

Yang, H., Paller, K. A., & Van Vugt, M. (2022). The steady state visual evoked potential

(SSVEP) tracks "sticky" thinking, but not more general mind-wandering. Frontiers in

Human Neuroscience, 16, 892863. https://doi.org/10.3389/fnhum.2022.892863

Yang, Z., Williams, S. D., Beldzik, E., Anakwe, S., Schimmelpfennig, E., & Lewis, L. D.

(2025). Attentional failures after sleep deprivation are locked to joint neurovascular,

pupil and cerebrospinal fluid flow dynamics. *Nature Neuroscience*, 1–11.

https://doi.org/10.1038/s41593-025-02098-8