Federated single-cell QTL meta-analysis reveals novel disease mechanisms

Daniel Kaptijn[1,2,*], Lieke Michielsen[3,4,*], Drew Neavin[5,6,7,*], Aida Ripoll-Cladellas[8,9,*], José Alquicira-Hernández[5,10,$], Maryna Korshevniuk[1,2,$], Jimmy Tsz Hang Lee[11,$], Roy Oelen[1,2,$], Martijn Vochteloo[1,2,$], Robert Warmerdam[1,2,$], Yoshinari Ando[12], Maria Ban[13], Odmaa Bayaraa[14], Marijn Berg[15,16], Irene van Blokland[1,17], Daniel Considine[11,18], Mame M. Dieng[14], Ryuya Edahiro[19,20,21], M. Grace Gordon[22,23,24,25], Hilde E. Groot[17], Pim van der Harst[17], Matthias Heinig[26,27], Chung-Chau Hon[12], Youssef Idaghdour[14,28], Pooja Kathail[29], Niek de Klein[11], Wenchao Li[30], Yang Li[30,31], Corinna Losert[26,27], Vinu Manikanda[14,28], Jonathan Moody[12], Haroon Naeem[32], Younes Mokrab[32,33,34], Martijn C. Nawijn[15,16], Mihai Netea[31], Jelmer Niewold[1,2], Yukinori Okada[19,35,36,37], Stephen Sawcer[13], Issiaka Soulama[38], Oliver Stegle[11,39,40], Yakov Tsepilov[11,18], Woong-Yang Park[41], Deepa Rajagopalan[42], Tala Shahin[14], Jay W. Shin[42], Gosia Trynka[11,18], Urmo Võsa[43], Harm-Jan Westra[1,2], Seyhan Yazar[5,44,45], Jimmie Ye[23,24,46,47,48,49], zhenhua zhang[30], eQTLGen consortium[50], Martin Hemberg[51,&], Ahmed Mahfouz[3,4,&], Marta Melé[8,&], Joseph E. Powell[5,6,52,&], Marc Jan Bonder[1,2,#,⌀], Lude Franke[1,2,#,⌀], Monique G.P. van der Wijst[1,2,#,⌀]

1. Department of Genetics, University of Groningen, University Medical Center Groningen.
2. Oncode Institute, Groningen, the Netherlands.
3. Department of Human Genetics, Leiden University Medical Center, Leiden, The Netherlands.
4. Delft Bioinformatics Laboratory, Delft University of Technology, Delft, The Netherlands
5. Translational Genomics Program, Garvan Institute of Medical Research, Darlinghurst, Sydney, NSW, Australia
6. University of New South Wales, Kensington, Sydney, NSW, Australia
7. Current affiliation: Institute for Molecular Bioscience, University of Queensland, Brisbane, QLD, Australia
8. Life Sciences Department, Barcelona Supercomputing Center, Barcelona, Catalonia, Spain
9. Universitat de Barcelona, Barcelona, Spain
10. Computational Genomics, Institute for Molecular Bioscience, University of Queensland, Brisbane, Australia
11. Wellcome Sanger Institute, Wellcome Genome Campus, Cambridge, UK.
12. Laboratory for Genome Information Analysis, RIKEN Center for Integrative Medical Sciences, Japan.
13. University of Cambridge, Department of Clinical Neurosciences, Cambridge Biomedical Campus, Cambridge, CB2 0QQ, UK
14. Biology Program, New York University Abu Dhabi, United Arab Emirates.
15. Department of Pathology and Medical Biology, University of Groningen, University Medical Center Groningen, Groningen, the Netherlands.
16. GRIAC research institute, University Medical Center Groningen, Groningen, the Netherlands.
17. Department of Cardiology, University of Groningen, University Medical Center Groningen, Groningen, the Netherlands.
18. Open Targets, Wellcome Genome Campus, Hinxton, Cambridgeshire
19. Department of Statistical Genetics, Graduate School of Medicine, The University of Osaka, Suita, Japan.
20. Laboratory for Systems Genetics, RIKEN Center for Integrative Medical Sciences, Yokohama, Japan.
21. Department of Respiratory Medicine and Clinical Immunology, Graduate School of Medicine, The University of Osaka, Suita, Japan
22. Biological and Medical Informatics Graduate Program, University of California San Francisco, San Francisco, USA.
23. UCSF Division of Rheumatology, Department of Medicine, University of California San Francisco, San Francisco, CA, USA.
24. Institute for Human Genetics, University of California San Francisco, USA.
25. Department of Bioengineering and Therapeutic Sciences, University of California San Francisco, San Francisco, USA.
26. Institute of Computational Biology, German Research Center for Environmental Health, Helmholtz Zentrum München, Neuherberg, Germany
27. Department of Computer Science, TUM School of Computation, Information and Technology, Technical University of Munich, Garching, Germany 5. Open Targets, Wellcome Genome Campus, Hinxton, Cambridges
28. Public Health Research Center, New York University Abu Dhabi Institute, New York University Abu Dhabi, United Arab Emirates.

29. Department of Electrical Engineering and Computer Science, Center for Computational Biology, University of California Berkeley, Berkeley, CA, USA

30. Centre for Individualised Infection Medicine (CiiM) & TWINCORE, joint ventures between the Helmholtz-Centre for Infection Research (HZI) and the Hannover Medical School (MHH), Hannover, Germany

31. Department of Internal Medicine and Radboud Center for Infectious Diseases, Radboud University Nijmegen Medical Center, Nijmegen, Netherlands

32. Sidra Medicine, Doha, Qatar.

33. Department of Biomedical Science, College of Health Sciences, Qatar University, Doha, Qatar.

34. Department of Genetic Medicine, Weill Cornell Medicine, Qatar.

35. Department of Genome Informatics, Graduate School of Medicine, the University of Tokyo, Tokyo, Japan.

36. Laboratory of Statistical Immunology, Immunology Frontier Research Center (WPI-IFReC), The University of Osaka, Suita, Japan

37. Premium Research Institute for Human Metaverse Medicine (WPI-PRIMe), The University of Osaka, Suita, Japan

38. Institut de Recherche en Sciences de la Santé, Ouagadougou, Burkina Faso

39. Division of Computational Genomics and Systems Genetics, German Cancer Research Center (DKFZ), Heidelberg, Germany.

40. European Molecular Biology Laboratory (EMBL), Genome Biology Unit, Heidelberg, Germany.

41. Samsung Genome Institute, Samsung Medical Center, Seoul, Korea.

42. A*STAR Genome Institute of Singapore, Singapore.

43. Estonian Genome Centre, Institute of Genomics, University of Tartu, Tartu, Estonia

44. School of Clinical Medicine, UNSW Medicine and Health, Sydney, NSW, Australia

45. Precision Immunology Program, Garvan Institute of Medical Research, Darlinghurst, Sydney, NSW, Australia

46. Bakar Computational Health Sciences Institute, University of California San Francisco, San Francisco, CA, USA.

47. Department of Epidemiology and Biostatistics, University of California San Francisco, San Francisco, CA, USA.

48. Parker Institute for Cancer Immunotherapy, San Francisco, CA, USA.

49. Chan Zuckerberg Biohub, San Francisco, CA, USA

50. Banner author, see **Note S1**

51. The Gene Lay Institute of Immunology and Inflammation, Brigham and Women's Hospital, Massachusetts General Hospital, and Harvard Medical School, Boston, USA

52. UNSW Cellular Genomics Futures Institute, University of New South Wales, Sydney, Australia.

*, $, &, # These authors have contributed equally.

⬚ Correspondence: Marc Jan Bonder (m.j.bonder@umcg.nl), Lude Franke (l.h.franke@umcg.nl), Monique van der Wijst (m.g.p.van.der.wijst@umcg.nl)

91 **Abstract**

Genetic effects on gene expression are often cell type-specific and obscured in bulk analyses. To
resolve this context-dependent regulation, we performed a federated *cis*-eQTL meta-analysis across
12 PBMC datasets (2,032 individuals, 2.5 million cells). Across six immune cell types, we identified *cis*-
eQTLs for 6,592 genes and fine-mapped 14,985 independent loci. Notably, the 42% of eQTLs that
were undetected in a bulk eQTL study on 43,301 whole blood samples also showed stronger
enrichment for disease GWAS loci. We further identified three genome-wide significant and 65
suggestive loci affecting the abundance of (rare) immune cell types and validated these using
previously reported hematological GWAS and bulk-derived *trans*-eQTLs. Integrating single-cell *cis*-
eQTLs with bulk *trans*-eQTLs enabled us to anchor 6,382 *trans*-eGenes (37.2% novel) to upstream
regulators and reconstruct directed gene regulatory relationships. For example, a hemorrhoidal
disease-associated variant showed a CD4+ T cell-specific *cis*-eQTL on *BACH1* that colocalized with 45
immune and metabolic *trans*-eGenes. These results demonstrate the power of single-cell QTL meta-
analysis in interpreting complex trait genetics.

Introduction

Genome-wide association studies (GWAS) have now revealed over 200,000 genetic risk factors, mostly single nucleotide polymorphisms (SNPs), for more than 200 different traits[1]. Deciphering how these genetic variants contribute to disease is fundamental for advancing therapeutic strategies. However, as the majority of these GWAS SNPs are in the non-coding part of the genome, it has been thus far challenging to pinpoint the molecular mechanisms underlying how these SNPs confer disease risk.

Linking GWAS SNPs to cell type abundance or specific molecular data layers through quantitative trait locus (QTL) analyses has already increased our understanding of these underlying mechanisms. Previously, the largest effort, involving 563,085 individuals, linked 7,122 genetic variants to cell type abundance QTLs (i.e., ccQTLs) of predefined blood cell types, measured using flow cytometry[2,3]. Importantly, these ccQTLs often colocalize with disease[2–4], providing a mechanism how cell type composition changes may be involved in disease pathogenesis. Moreover, the largest molecular QTL analyses to-date have been conducted using gene expression data, i.e., expression QTL (eQTL) mapping. Such efforts span broad multi-tissue initiatives, as done in the Genotype-Tissue Expression (GTEx) consortium that mapped eQTLs in 838 donors across 49 tissues[5], as well as large single tissue studies, exemplified by the eQTLGen consortium that mapped eQTLs in whole blood of 43,301 individuals[6]. Despite these large-scale efforts, no potential underlying molecular mechanism has been identified for most disease-associated variants. For example, the GTEx consortium found that only 43% of disease-associated loci colocalize with a known local (*cis*) eQTL effect in at least one tissue[2]. Similarly, the eQTLGen consortium (phase 2) found a *cis* or distal (*trans*) eQTL effect for 45.9% of the 1,848 independent immune disease-associated SNPs tested[6]. Different explanations have been provided for this relatively small overlap. First, many genetic variants may act only in a specific cellular or environmental context[7,8] that is not adequately captured by tissue-level studies such as GTEx[5] or eQTLGen[6]. Second, molecular readouts other than gene expression levels (e.g., splicing, chromatin accessibility) may be needed to capture additional overlap with disease-associated variants[9]. Finally, despite the enormous sample sizes of current eQTL consortia, statistical power to detect genome-wide eQTL effects is still limited for tissues other than blood. Yet it is these context-dependent and distal eQTL effects that may be most relevant for our disease understanding[10,11].

Single-cell RNA-sequencing (scRNA-seq) data provide a unique opportunity to unbiasedly detect high resolution ccQTLs and context-dependent eQTL effects. However, currently available individual single-cell datasets are still limited in sample size compared to previous bulk QTL consortia efforts[8,12,13]. To increase the power of sc-QTL studies, we initiated the sc-eQTLGen consortium in which the scRNA-seq and genotype data from individual cohorts are processed with standardized computational pipelines so that their ccQTL or eQTL summary statistics can ultimately be combined in a federated QTL meta-analysis[14]. In the current study, we unify 12 single-cell peripheral blood mononuclear cell (PBMC) datasets in a federated QTL meta-analysis and map cell type abundance and cell type-specific regulatory variants across 2,032 donors. We then fine-map these effects, integrate them with a large collection of GWAS studies and with *cis*- and *trans*-eQTLs from the eQTLGen consortium, and demonstrate their value for interpreting immune disease risk (**Figure 1**). This information helps pinpoint how and in which cell type these GWAS risk variants act, thereby providing insights that may inform risk stratification, therapeutic development, and prevention strategies.

Results

Building harmonized pipelines to perform QTL meta-analysis across 12 cohorts

We performed ccQTL and eQTL meta-analyses in the sc-eQTLGen consortium, combining scRNA-seq data from 2,032 individuals and ~2.5M PBMCs (**Figure 1a**, and **Table S1**). We harmonized data from 12 cohorts (**Note S2**), generated by different single-cell technologies and chemistries (e.g., full-length

Smart-seq2, 3'- and 5'-end 10X Genomics) and from individuals from multiple ancestries, different
ages, and both sexes (**Table S1**). Specifically, we included the OneK1K study[13] (1,017, European
ancestry), CD4+ T cells from the Trynka study[15] (117, European ancestry), CD4+ T cells from the ARMS
study[16] (45, European ancestry), the 300BCG study[17] (38, European ancestry), the OASIS study[18,19] (91,
East Asian Ancestry), cerebrospinal fluid (CSF) data[20] from the Sawcer lab (73, European ancestry),
the Burkina Faso malaria study[21] (178, African Ancestry), the Ye SLE study[12] (170, European, East
Asian and Southern Asian ancestries), the van der Wijst study[22] (40, European ancestry), unpublished
single nucleus dataset (118, European ancestry), and unstimulated samples from the 1M-scBloodNL
project[8] (120, European ancestry) and the van Blokland STEMI study[23] (38, European ancestry) **(Note**
S2).
Because QTL analyses involve privacy-sensitive genotype information, raw data could not be
centralized. Instead, standardized analysis pipelines were run within each cohort, and only summary
statistics for the federated meta-analyses were shared. To facilitate this setup, we established several
working groups (WG) to harmonize: WG1. genotype imputation and scRNA-seq data preprocessing,
quality control (QC), and donor demultiplexing, WG2. cell type assignment, and WG3. ccQTL and
eQTL mapping (**Figure 1b**, see **Methods**). These pipelines are publicly available on Github **(Code**
Availability), and several parts of these pipeline elements have been described in detail in separate
papers (sample demultiplexing[24], cell type assignment[25–27], and eQTL mapping[28]).
In short, we removed cells that were assigned ambiguously or as doublets using a
combination of tools that together capture both heterogenic doublets (from different donors) and
homogenic doublets (from the same donor)[24]. Additionally, we removed cells that expressed fewer
unique molecular identifiers (UMIs) or RNA molecules than the median minus twice the median
absolute deviation (MAD) of the entire dataset or whose mitochondrial gene content percentage
exceeded the median plus three times the MAD. Next, we annotated the QC-passing cells to seven
major cell types, which were further resolved into 27 cell subtypes: B cells (B intermediate, B
memory, B naïve, plasmablast), CD4+ T cells (CD4 cytotoxic T, CD4 naïve, CD4 proliferating, CD4
central memory (TCM), CD4 effector memory (TEM), regulatory T cell), CD8+ T cells (CD8 naïve, CD8
proliferating, CD8 TCM, CD8 TEM), monocytes (classical CD14+ monocyte, non-classical CD16+
monocyte), natural killer cells (NKdim, NKbright, NK proliferating) and dendritic cells (AXL+ DC,
conventional DC1 (cDC1), cDC2, plasmacytoid DC (pDC)), ILCs (innate lymphoid cells) and three other
T cell subtypes (MAIT, dnT, gdT) (**Table S2**). Cell type annotation was performed using hierarchical
scPred[25,27] and Azimuth[26]. We only retained cells that were concordantly classified by both tools at
the major cell type resolution. The annotation reference for these tools was trained on joint scRNA-
seq and sc-surface protein data (CITE-seq) of human PBMCs as defined previously[26]. These QCed and
annotated cells were then used for downstream QTL analyses **(Figure 1c)**.

190
191
Figure 1. Overview of the single-cell eQTLGen consortium. a. We initiated the sc-eQTLGen
consortium (Phase 1), which currently includes meta-analysis QTL summary statistics derived from 12
cohorts for 2,032 donors and each of the cell types found in PBMCs. **b.** Standardized computational
pipelines were developed to harmonize (WG1) quality control, demultiplexing, (WG2) cell type
annotation, and (WG3) QTL mapping across cohorts, enabling centralized meta-analysis of QTL
summary statistics. **c.** We conducted several downstream analyses: 1. ccQTL mapping on 63 different
cell type composition estimates (on both major PBMC cell types and their subtypes), identifying 68
unique genetic signals associated with differences in cell type proportion; 2. cell type-specific eQTL
analysis, which identified 6,592 eGenes and 14,985 independent eQTL effects; 3. systematic
colocalization of eQTLs with over 7,000 different GWASs identified colocalization to 5,004 GWASs; 4.
systematic colocalization of *trans*-eQTL signals from the bulk blood eQTLGen study (n = 43,301
samples) with our cell type-specific *cis*-eQTL signals, which identified 21,212 *cis-trans* gene pairs.
204

Single cell–derived ccQTLs capture bulk-derived *trans*-eQTLs

Flow cytometry-based analyses have previously shown the value of ccQTLs to interpret disease-associated variants[2,3]. To achieve higher cellular resolution in characterizing these ccQTLs, we therefore explored the usage of scRNA-seq data. We assessed the effect of genetics on cell type composition by conducting a ccQTL meta-analysis at different cellular resolutions. Per donor we calculated the ratio of cells belonging to: the lymphoid (B, CD4+ T, CD8+ T, NK, ILC) versus myeloid (monocyte, DC) lineage, each of the seven major cell types versus total PBMCs, each of the 27 cell subtypes versus total PBMCs, and cell subtypes versus the major cell type they belong to, totaling 63 phenotypes. As some of our ccQTLs represent cell type fractions within total PBMCs and ccQTLs typically have effect sizes smaller than eQTLs[29], we limited our meta-analysis to those datasets that were most similar (7 out of 12 datasets). Specifically, we excluded datasets that provided only partial PBMC representation rather than the full PBMC spectrum (the cerebrospinal fluid dataset (Sawcer CSF[20]) and two datasets that contained only a single major cell type (Trynka[15] and ARMS[16])), one dataset with fewer than 40 donors (300BCG[17]), and one dataset from multiple ancestries (Ye SLE[12]). Furthermore, we limited our analysis to donors who had at least 350 total PBMCs, resulting in the inclusion of 1,710 donors. We defined Gaussian-normalized cell type ratios as phenotypes and applied a linear mixed model to test associations with genetic variants, testing variants with MAF >5% and Hardy-Weinberg Equilibrium (HWE) P-value > 1×10^{-4} (see **Methods**). The resulting summary statistics were subsequently combined using an inverse-variance weighted meta-analysis, keeping only effects tested in all datasets. To be conservative, given the remaining differences between datasets, we removed variants with a high association heterogeneity ($I^2 > 40\%$).

We noted an interdependency across many of the tested cell type proportions. For instance, the proportions of lymphocytes to PBMCs were negatively correlated with those for myeloid cells to PBMCs. ccQTL analysis on these proportions would therefore likely yield the same genetic association signals, albeit with opposite effect directions. Therefore, to more accurately count the number of associated genetic loci, we first defined association loci using "Locus breaker"[30] (see **Methods and Fig SX**). In these loci we performed fine-mapping on each of the ccQTL summary statistics sets, using SuSiE[31] with the multiple causal variant assumption (see **Methods**). These fine-mapped loci we termed independent loci. We subsequently performed colocalization of the independent loci with coloc[32] to identify sharedness of genetic signal among cell composition traits. If fine-mapping was not successful, we deemed the loci to be the same if the association regions overlapped, and counted non-overlapping regions as a single independent locus.

After applying fine-mapping and colocalization, we identified three genome-wide significant independent loci ($P < 5.0 \times 10^{-8}$), associated with cell subtype composition in monocytes, CD8+ T cells, and NK cells (**Figure 2a** and **Table S3**). At a suggestive threshold ($P < 5.0 \times 10^{-6}$), we identified an additional 65 independent loci, including loci influencing the proportion of B cells, CD4+ T cells, and DCs (**Table S3**). Thus, in all, we found 68 independent genetic signals associated with cell type composition derived from PBMC single-cell data. Most of these signals corresponded to known regulators of immune cell abundance: 63 of the 68 independent loci (92.6%) overlapped with cell count traits in the UK Biobank (UKB) cell composition GWAS[3] (leukocyte count, monocyte count, lymphocyte percentage of leukocytes, and monocyte percentage of leukocytes) (**Table S3**). For example, the UKB 'monocyte count' locus rs67122419 ($P = 3.2 \times 10^{-151}$) aligned with our genome-wide signal of CD14+ versus CD16+ monocyte proportions. Likewise, a genome-wide association for CD8+ TEM in total PBMC counts colocalized with the UKB 'leukocyte count' locus rs2523579 ($P = 4.8 \times 10^{-191}$). Moreover, a suggestive CD4+ T cell in total PBMC counts signal corresponded to the UKB 'lymphocyte percentage of leukocytes' rs4648050 ($P = 6.2 \times 10^{-44}$). However, we note that, while nearly all of the independent loci have been reported before[2,3], the traits do not precisely match, as the UKB measurements were derived from flow cytometry–based technologies. Therefore, some differences between the association signals may exist due to differences in cell count resolution and the use of different measures to quantify cell count traits. By leveraging our single-cell data, we have added resolution to the existing associations; however, as resolutions differ, we cannot directly replicate effect directions.

As an alternative way to validate the ccQTLs at the appropriate resolution, we reasoned we
could use the *trans*-eQTLs identified in bulk data. These *trans*-eQTLs are likely a mixture of genetic
effects on cell type composition and intracellular expression regulation. For instance, if a gene is
predominantly expressed in a specific cell subtype, this can lead to an apparent *trans*-eQTL effect in
bulk RNA-seq data if the associated variant changes the proportion of that cell subtype, rather than
changing gene expression. Therefore, we colocalized our ccQTLs with *trans*-eQTLs identified using
bulk RNA-seq data from 43,301 whole blood samples in eQTLGen (phase 2)[6]. Then, using one of the
representative scRNA-seq datasets (UMCG v3[8,23]) in sc-eQTLGen, we used UCell[33] to compute a rank-
based expression score indicating how highly expressed the ccQTL-associated trans-eGenes are in a
single cell (**Methods**). Scores were averaged per cell (sub)type per donor. We next tested whether
UCell scores differed between the relevant cell type and the other cells contributing to the fraction
underlying the ccQTL phenotype, using a paired Wilcoxon test. This approach allowed us to assess
whether the bulk-derived *trans*-eGenes that were associated with the same genetic variant as the
ccQTL were differentially expressed in the cell type implicated in the ccQTL. At the same time, it
enabled us to distinguish *trans*-eQTL signals driven by changes in cell type composition from those
reflecting intracellular regulatory effects.
We could apply this approach to four independent ccQTL loci, as we required the ccQTL to
colocalize with at least two *trans*-eGenes: two out of three genome-wide loci (monocyte and NK
effects) and two suggestive loci (CD4+ T in PBMC and CD8+ TEM in CD8+ T cells). All four ccQTL loci
were validated using this approach: the module scores were significantly different between the cell
(sub)types and were highest in the cell type whose abundance increased in association with the
specific allele (monocyte locus: Wilcox $P = 4.1 \times 10^{-17}$; NK locus: Wilcox $P = 1.9 \times 10^{-17}$; CD8+ TEM locus:
Wilcox $P = 1.2 \times 10^{-12}$; CD4+ T locus: Wilcox $P = 0.02$) (**Figure 2b–f**). While we expect that our list of
ccQTLs and *trans*-eQTLs are incomplete, the concordant replication and consistent effect directions
provide additional support for the robustness of our ccQTL findings.
Next, we assessed whether the identified ccQTL loci contribute to disease, as was previously
reported for flow cytometry-based ccQTLs[34]. To this end, we examined the lambda (λ) inflation to
evaluate whether our ccQTLs show an enriched association signal in external GWASs. We confirmed
that our ccQTL variants showed a larger λ-inflation for hematological measurement traits as
compared to other GWAS (Wilcox $P = 0.012$, median λ-inflation difference = 0.02, **Figure 2g**),
indicating enrichment for genetic determinants of blood cell composition. In contrast, we observed
no λ-inflation for disease or immune system disease in general, or any specific immune disease GWAS
—likely reflecting the limited number of significant ccQTL and disease loci. The absence of immune
disease-related enrichment compared to previous flow cytometry-based measurements[34] may reflect
the substantially smaller sample size of our study (up to 224-fold fewer individuals) or that disease-
relevant effects arise in cell types not represented in PBMCs, such as neutrophils or eosinophils.
293

Figure 2. Overview of the ccQTL results. a. Combined Manhattan plot of the three genome-wide significant (P < 5x10^{-8}) and 65 suggestive (P < 5x10^{-6}) cell count GWAS traits. These association signals are colored by the cell type whose proportions are affected by the variant. b. Subset of the UMAP for the monocytes. The two cell subtypes are circled (CD16+ on top, CD14+ monocytes bottom). Colors indicate the UCell scores per cell based on the *trans*-eGenes linked to the ccQTL SNP. c-f. Violin plots of donor-aggregated averages of the UCell score in the monocyte (c), NK (d), CD8+ T (e) and CD4+ T (f) cell (sub)types. g. Violin plots of the λ-inflation of the ccQTL variants versus GWAS-associated variants for hematological traits (left) or other GWAS (right) (Methods).

Single-cell data detects novel eQTLs that are more often lineage-specific than bulk-derived eQTLs

Many ccQTLs likely act through *cis*-regulatory mechanisms: a genetic variant first influences the expression level of a specific gene (*cis*-eQTL) involved in, for example, lineage differentiation, after which cell type composition is affected. However, genetic variants can also affect gene expression levels without altering cell type composition. Therefore, to systematically capture such regulatory effects, we conducted a *cis*-eQTL meta-analysis across all 12 cohorts, assessing all QC-passing variants within a 1 megabase (Mb) distance around the gene (see **Methods**). We limited the eQTL analysis to the six most abundant major cell types (CD4+ T, CD8+ T, NK, B, Monocyte and DC), to have sufficient

cell numbers in each dataset for reliable pseudobulks. Across all cell types, we detected eQTL effects
for 6,592 genes (eGenes), covering 56.4% of the expressed genes (see **Methods**) (**Figure 3a**).
Compared to the largest sc-eQTL study within our meta-analysis (OneK1K, n = 1,018 donors),
doubling the number of donors through meta-analysis increased the number of detected eGenes by
1.3x (for CD4+ T cells) to 6.7x (for DCs) (**Figure 3a**). The number of detected eGenes per cell type was
closely related to the average number of donors and cells per donor available for that cell type
(**Figure 3b, Figure SX**): we identified the highest number of eGenes in the cell type for which most
donors and cells per donor were available (CD4+ T: 4,988 eGenes, 39.2% of all expressed genes), and
the fewest for the cell type with least donors and fewest cells per donor (DCs: 375 eGenes, 3.8% of all
expressed genes).
Multiple genetic variants within a cell type or across cell types may regulate the same eGene.
First, to identify the independent loci that regulate the same eGene within a cell type, we conducted
fine-mapping using SuSiE[31]. In total, we identified 14,985 independent eQTLs (**Table S4**), increasing
the total number of eQTL signals by 20.4⬚36.2% (**Figure 3a**). We observed the highest number of
independent effects in CD4+ T cells, in line with the higher numbers of donors and cells for that cell
type. Most independent eQTL effects were identified within 100 kb around the transcription start site
of the gene they are regulating. In line with previous observations[35], secondary eQTL effects were, on
average, further from the gene than the primary eQTL effect (**Figure S1**).
We then studied the cell type-specificity of eQTL effects by comparing the effect of significant
eQTLs across cell types using two complementary strategies. In the first, we assessed pair-wise
similarity across cell types between expressed genes (**Figure 3c**) and between eQTL effect sizes
(**Figure 3d**) using the r_b metric[36]. In the second strategy, we employed systematic colocalization of the
fine-mapped loci across the 6 major cell types and over genes (**Figure 3e**). While the r_b metric
evaluates whether shared genes are similarly regulated by genetics, the colocalization formally tests
whether the genetic signal is truly overlapping, but only for signals that are strong enough to be fine-
mappable. The first strategy showed that whether a gene is expressed (**Figure 3c**) and the gene's
genetic regulation (**Figure 3d**) are more similar within cells from the same lineage than across
lineages. For instance, CD4+ and CD8+ T cells showed high concordance of eQTL effect sizes ($r_b = 0.9$),
whereas the CD4+ T cells and monocytes showed substantially lower concordance ($r_b = 0.6$). The
colocalization strategy supported these observations. We identified 6,434 independent loci across the
six major cell types, allowing for loci to affect multiple genes (**Figure S2**). Of these, 82.0% were
associated with a single gene, although some, particularly in the HLA locus, affected up to 66 genes.
Overall, most loci (62.2%) were specific to a single cell type. This is likely in part explained by the
relatively large number of *cis*-eGenes uniquely detected in CD4+ T cells due to higher statistical
power in this cell type. However, when restricting our analysis to the 431 genes tested in at least five
of the six cell types, the results were more similar to those of the r_b-based approach (**Figure S3**: 91%
of these genes were shared within rather than across lineages (85%).
Next, we assessed how many of our independent eQTL loci were also found in eQTLGen
(phase 2)[6] (**Methods**). We observed comparable results for each of the six cell types (**Fig. 3e**). Overall,
57.8% of the independent loci were found in both the sc- and bulk-derived datasets and shared at
least one eGene, providing confidence in our results while also allowing annotation of the cellular
context in which these genetic variants act. The remaining loci either did not have any overlapping
eGene assigned (17.5%) or were not identified in eQTLGen (24.7%). As expected, loci shared across
single-cell and bulk data were more often shared across lineages (on average 39.6% cross-lineage
effects for bulk-shared effects and 21.0% for single-cell-specific effects, **Figure 3e**). Together, these
results highlight the added value of single-cell data for detecting novel, often lineage-specific, eQTLs
and for providing the context in which an eQTL acts.
360

Figure 3. eQTL meta-analysis results. a. Comparison of the number of eGenes in sc-eQTLGen (light
blue) and the number of eQTLs in the largest individual dataset (OneK1K[13], gray). Additionally
showing the number of independent loci after fine-mapping (darker shade of blue). **b.** Scatter plot
showing, per cell type, the average number of cells per donor versus the number of detected eGenes.
c. Heatmap showing the number of shared expressed genes between the six cell types. Columns are
the discovery cell type. Rows are the replication cell types. **d.** Heatmap as in **(c)** but now showing the
r_b between the eQTL effect sizes of the shared expressed genes. **e.** Bar chart depicting the number of
independent loci observed in the six cell types annotated by their replicability in bulk eQTLGen[6] (blue
= bulk-shared, yellow= sc-specific). The bulk-shared and sc-specific loci are further subdivided by their
replicability across blood lineages (darker shade = lineage-specific, lighter = lineage-agnostic).

Single-cell QTLs shed new light on disease
Next, we determined whether and how single-cell QTLs provided new disease insights. As we
assumed that all ccQTLs relevant to disease will act through eQTLs, we related all our comparisons to
eQTLs first. For this, we colocalized the eQTL loci to GWAS loci by leveraging the OpenTargets
Genetics database[37] of more than 7,000 GWASs. We found 2,271 independent loci (6,434 eQTL,
Figure 4a) were linked to 5,004 different GWASs (Table S5). Among those GWASs, immune system

disease loci were most often colocalizing with eQTL loci (median ratio 0.40) and least often with
hematological measurement loci (median ratio 0.30) **(Figure 4b)**. Interestingly, other diseases than
immune system diseases also showed higher colocalization ratios than hematological measurements
(median ratio = 0.34), likely due to some immune involvement in those diseases as well **(Figure 4c)**.
To better understand which eQTL loci are more likely to colocalize to GWAS loci, we used
logistic regression (see **Methods**). This revealed that eQTL loci affecting a larger number of genes (P =
7.2×10^{-14}), detected across multiple cell types (P= 2.8×10^{-5}), or identified in both lymphoid and
myeloid cells (P= 2.0×10^{-4}), were significantly more likely to colocalize with a GWAS signal **(Figure 4d)**.
Interestingly, these same analyses showed that eQTL loci shared between single-cell and bulk
eQTLGen are more likely to colocalize to GWAS traits (P= 2.6×10^{-4}) **(Figure 4d)**. More specifically,
when assessing characteristics of shared loci we found that these are significantly enriched in
hematological measurements (P= 2.4×10^{-3}), anthropometric traits (P= 0.02), other (non-disease-
associated) traits (P= 5.0×10^{-3}) **(Figure 4e)**. These colocalization-based observations are in line with
SNP-based enrichments (considering linkage disequilibrium (LD)), in which both the immune system
disease GWAS (EFO0000540, cross cell type Fisher OR: 8.7, P= 1.6×10^{-48}) and all disease trait GWAS
sets (EFO0000408, cross cell type Fisher OR 4.1, P= 8.1×10^{-104}) were enriched in our sc-eQTLs over all
other SNPs assessed **(Table S6)**. Notably, we observed that the sc-eQTLs are significantly more
enriched for disease GWAS than bulk eQTLGen eQTLs (Z-test OR 2.9, P= 4×10^{-4}, **Table S6**),
underscoring the added value of sc-eQTLs for disease interpretation. Together, these analyses show
that effects unique to single-cell data are more relevant to the interpretation of disease GWAS loci,
and shared effects are more likely to capture cell (sub)type composition related loci.
As an example of a disease association uniquely identified using sc-eQTLs, we considered
two eQTL loci that both colocalized with 'eczema or dermatitis' (GCST90038680)[38]. The
first locus, containing risk variant rs12712145 (Chr2:102,392,150, GWAS P = 2.2×10^{-10}), was present
in NK and CD4+ T cells, but different genes were implicated depending on the cell type. The C risk
allele increased expression of *IL18R1* in NK cells (eQTL Z = 14.6) but lowered expression of *IL18RAP* in
CD4+ T cells (eQTL Z = -0.45, **Figure S4**). The second locus was found in B cells, where the disease risk
allele A of variant rs28618323 (Chr14:35,106,586, GWAS P = 8.2×10^{-10}) decreased expression
of *FAM177A1* (eQTL Z = -6.1). Together, these genes are all involved in the same signaling pathway.
IL18R1 and IL18RAP form the IL18 receptor complex that activates downstream NF-κB signaling upon
binding of IL18[39]. FAM177A1 competitively binds TRAF6, preventing its downstream activation of NF-
κB signaling[40]. This implies that the elevated IL-18 signaling in atopic dermatitis patients, which also
correlates with disease severity[41], may be facilitated through increased signaling in NK cells and
reduced suppression in B cells **(Figure 4h)**.
Another locus uniquely identified in single-cell data as both ccQTL and eQTL and associated
with disease (early onset of asthma, FINNGEN_R12_ASTHMA_CHILD_EXMORE, GWAS P = 5.5×10^{-9}),
was the risk variant rs391934 (Chr2:8,304,364, **Figure 4i**). The T risk allele increased the proportion of
CD8+ TEM to total CD8+ T cells (ccQTL Z-score = 4.8) and increased the expression of the long-non-
coding RNA *LINC00299* in NK cells (eQTL Z-score = 4.9, **Figure S5**). Although CD8+ TEM have not been
directly associated with asthma before, CD8+ TEMRA have been[42]. This terminally-differentiated
subset of CD8+ T cells can arise from CD8+ TEM that re-express CD45RA[43]. Because our eQTL meta-
analyses were conducted primarily on 3'-end scRNA-seq data, we could not readily distinguish the
CD8+ TEM and CD8+ TEMRA subtypes, which are defined by their expression of different CD45
protein isoforms (CD45RO/CD45RA) encoded by the *PTPRC* gene. We therefore suspect that this
ccQTL may have instead captured a change in the abundance of CD8+ TEMRA cells. Further research
is needed to define whether the regulation of *LINC00299* expression in NK cells contributes to the
abundance change of the CD8+ TEMRA subset, or that the eQTL effect may have been present in the
CD8+ TEM subset as well as a consequence of the ccQTL. Notably, the latter possibility is supported
by the high expression of *LINC00299* within the CD8+ TEM subset.
429

430
431

Figure 4. QTL to GWAS trait colocalization. a. Systematic colocalization of independent eQTL effects
with GWAS traits. For colocalizing signals, sharing of eQTLs between single-cell and bulk eQTLGen was
evaluated. **b.** Per GWAS trait category the proportion of genetic signals that colocalized with eQTLs. **c.**
The other disease category (from **b**) split into more precise disease categories, demonstrating that
immune⬚derived eQTLs inform traits beyond immune diseases. **d.** SNP features that increase chances
(log OR) that a sc-eQTL colocalizes with GWAS traits. **e.** Likelihood (log OR) that eQTL loci shared
between single-cell and bulk eQTLGen, colocalize with different GWAS traits (black: not significant,
red: significant). **f. & g.** Example locus plots for loci that are uniquely associated to sc-eQTL effects
colocalizing to the GWAS on "Eczema or dermatitis". **f.** Locus plot for the colocalization between the
GWAS signal (rs12712145) and *IL18R1* in NK cells. **g.** Locus plot for the colocalization between the
GWAS signal (rs28618323) and *FAM177A1* in B cells. **h.** Illustration of the hypothetical mechanism by
which IL-18 signaling to NF-kB plays a role in the disease pathogenesis of Eczema and Dermatitis. **i.**
Example of disease association (early onset of asthma) identified by ccQTL effect in sc-eQTLGen.
445
Expanding disease-relevant *cis-trans* gene pairs identified in bulk by leveraging single-cell
resolution
Finally, we assessed whether the *cis*-eQTLs unique to single-cell data could also help refine gene
regulatory networks relevant to disease. Ideally, this would involve mapping these *cis*-eQTLs and
trans-eQTLs in our single-cell data, such that *cis*-regulatory variants can be directly linked to their
downstream *trans* effects. However, the sample size of our single-cell meta-analysis is underpowered
for genome-wide *trans*-eQTL discovery. In contrast, eQTLGen (phase 2)[6] provides adequate power for
such analyses, with a sample size >20-fold larger. Within eQTLGen, 48,037 *trans*-eQTLs were
identified (defined as effects >5Mb away from the eSNP, reflecting 9,199 unique trans-eGenes), with
23,248 (52.8%) linked to at least one *cis*-eGene (**Table S7**). These *cis-trans* associations enabled
inference of directional gene-gene relationships, offering additional insights into disease mechanisms.
To leverage both the resolution of the single-cell data and the statistical power of bulk data, we
integrated these approaches to improve identification of true regulatory *cis-trans* gene pairs
associated with disease.
First, we examined how many sc-derived *cis*-eQTLs could be colocalized with bulk-derived
trans-eQTLs: 14,683 *trans*-eQTLs (5,563 *trans*-eGenes). Most variants could not be colocalized as
there was no *cis*-eQTL effect in the single-cell data. In total, we identified 6,382 *trans*-eQTLs that
colocalized with at least one sc-derived *cis*-eQTL, corresponding to 3,706 *trans*-eGenes. 37.2% of
these colocalizing *trans*-eQTLs (**Figure 5a**) were newly annotated to at least one *cis*-eQTL using the
single-cell data and most of these links (58%) were associated with disease (annotation based on
eQTLGen[6]). In total these colocalizing effects reflected 21,212 unique *cis-trans* gene pairs (linking on
average 5.7 *cis*-eGenes to a *trans*-eGene), including 15,563 (73.4%) that were not observed in
eQTLGen.
A particularly noteworthy single-cell-specific effect was a *trans*-eQTL locus linked to a *cis*-
eQTL locus for *BACH1*. Two independent *cis*-eQTLs were found in single-cell data: one in CD4+ and
CD8+ T cells (rs2832300) and the other in monocytes and CD8+ T cells (rs66723060) (**Figure 5b**).
Interestingly, only the T cell-specific locus was linked to the *trans*-eQTL effects, corresponding to 45
trans-eGenes. These genes were significantly enriched for *BACH1*[43] P = 0.01 (corrected for the
number of assessed sets, **Methods**)), supporting the validity of the observed *trans*-eQTL effects. The
cis-locus was further implicated in a GWAS for hemorrhoidal disease[44] (PP H4: 0.88), a condition
influenced by genetic, immunological, and metabolic factors[30]. Consistent with this, pathways related
to the immune system (Reactome:R-HSA-168256, g:Profiler P = 5.4×10^{-5}) and metabolism
(Reactome:R-HSA-1430728, g:Profiler P = 2.7×10^{-5}) were enriched among the *trans*-eGenes after
accounting for genes expressed in eQTLGen. The T cell-specific locus was not detected in the bulk
data, likely due to the averaging of the two sc-eQTL loci in bulk data (**Figure 5b**). This was supported
by the higher LD between bulk (rs10563577) and single-cell lead variants (R^2 = 0.29 bulk-monocyte
effect (rs66723060); R^2 = 0.37 bulk-CD4+ T cell effect (rs2832300)) than between the two single-cell
effects (R^2 = 0.12 monocyte-CD4+ T cell).

Moreover, even for the *trans*-eGenes previously annotated using bulk data, single-cell *cis*-
eQTLs provide additional disease insights. For example, in bulk eQTLGen data, the *IKZF3 cis*-eQTL
colocalized with 48 *trans*-eGenes, whereas leveraging sc-eQTLGen data allowed us to annotate the
IKZF3 cis-eQTL as lymphocyte-specific and link the SNP to 255 *trans*-eGenes. Using also these
additional *trans*-eGenes, we found *IKZF* motifs to be significantly enriched among the linked *trans*-
eGenes (g:Profiler P = 3.0×10^{-3}, precision = 87%), providing additional support for *IKZF3* being the
key regulator of these genes. Interestingly, the *IKZF3 cis*-eQTL locus colocalized with several
autoimmune and allergic disease GWASs, including asthma (PP H4: 0.97), Crohn's disease (PP H4:
0.99), and rheumatoid arthritis (PP H4: 0.97). Additional evidence for the role of the 255 *trans*-
eGenes in these diseases was provided by a subset of 22 rare variant-implicated disease genes that
were also previously reported in Orphanet, ClinVar, or gene-burden analyses, as collected within the
Open Targets genetics platform[37] **(Figure 5c)**. Importantly, 15 of these genes were only linked through
single-cell data, further strengthening the connection with *IKZF3* and supporting the role of this locus
in regulation of genes critical for disease in general. These findings highlight the value of integrating
single-cell and bulk eQTLs to identify putatively causal *cis-to-trans* eQTL pairs that can help uncover
disease-relevant gene regulatory networks.

501
502

Figure 5. Distal disease networks uncovered via single-cell eQTL mapping. a. The number of bulk eQTLGen[3] *trans*-eQTLs that colocalize to single-cell or bulk eQTLGen *cis*-eQTLs, and the number of directional *cis-trans* gene pairs that could be inferred from that. **b.** Single-cell-specific *cis-trans* gene pairs identified for the T-cell-specific, but not monocyte *cis*-eQTL locus for *BACH1*. Only the T-cell-specific *BACH1 cis*-eQTL locus colocalizes with hemorrhoidal disease, providing mechanistic insights into how this locus contributes to the disease pathogenesis. Importantly, these effects could not be identified in bulk as the bulk-averaged *cis*-eQTL signal did not colocalize to either the *trans*-eQTLs or the disease GWAS. **c.** *Cis-trans* gene pairs identified for the bulk and single-cell *IKZF3 cis*-eQTL locus. Many additional *trans*-eGenes were linked to the *IKZF3 cis*-eGene using single-cell eQTL data. These additional *trans*-eGenes were enriched for *IKZF3* regulation, and a subset was also involved in rare disease through previously reported rare variants.

Discussion

We performed a ccQTL and eQTL meta-analysis in scRNA-seq data from up to 2,032 individuals in 2.5M peripheral immune cells. This meta-analysis setup increased the donor and cell sample size by more than two-fold compared to the largest study within the consortium[13], while also covering a wider range of different ancestries (African, European, East Asian and South Asian). Our ccQTL analyses identified three genome-wide significant and 65 suggestive CSs that affected the cell subtype composition of blood. Although most identified loci had previously been implicated with blood composition effects[2,3], our single-cell framework resolves these effects at substantial higher cellular resolution, revealing which specific immune populations are genetically perturbed. In addition, we could validate our findings by overlaying our ccQTL results with whole blood bulk-derived *trans*-eQTLs from over 43 thousand individuals[6]. The module scores of the *trans*-eGenes that share genetic signal with our sc-derived ccQTLs differed most in the cell type affected by the ccQTL, indicating correct assignment of a composition effect. This also shows that these specific *trans*-eQTLs, i.e., those linked to ccQTLs, likely reflect cell composition effects rather than intracellular expression effects. These results showcase how combining sc- and bulk-derived QTL results can synergize for interpretation of ccQTL and *trans*-eQTL effects. Moreover, the large number of replicated effects we observe, including the suggestive loci, indicates that many more reliable ccQTLs are expected to be discovered with increasing sample size.

Besides genetic effects on blood cell proportions, our sc-eQTL meta-analysis substantially expands the catalog of variants known to change gene expression in specific immune cell types. Compared to the largest sc-eQTL study included in our meta-analysis[13], which included about half the donors and one-third of the total number of cells, we discovered between 25-363% more *cis*-eQTLs, showcasing the value of our meta-analysis approach. We observed that 68% of these eQTLs act in a lineage-specific manner, and it was this category that was enriched among the 42% of eQTLs undetected in bulk blood-based eQTL meta-analyses efforts, even in over 20-fold more individuals[6]. This underscores a key limitation of bulk analyses: averaging across heterogeneous cell populations dilutes regulatory effects that are confined to specific immune lineages or activation states. As it is exactly these cell- and context-specific eQTLs that are more strongly enriched for disease[44], this explains why stronger enrichment for GWAS loci is observed among our single-cell-specific eQTLs as opposed to those shared with bulk data.

A further conceptual advance of this work lies in integrating single-cell *cis*-eQTLs with bulk *trans*-eQTLs to reconstruct disease-relevant regulatory cascades. While *trans*-eQTL discovery remains underpowered in single-cell data, the large-scale eQTLGen resource[6] provides a complementary opportunity to map downstream regulatory effects. By linking bulk *trans*-eQTLs to upstream regulators (i.e., *cis*-eQTLs) identified at single-cell resolution, we uncovered thousands of *cis*–*trans* gene pairs that were previously unannotated in bulk data. This integration not only increased the number of inferred regulatory relationships but also resolved their cellular context, as illustrated by the *BACH1* and *IKZF3* loci, where single-cell *cis*-eQTLs revealed lineage-specific regulatory hubs linked to immune and metabolic disease pathways.

Altogether, our study provides a valuable resource for anyone interested in understanding how genetic variation may impact disease, by identifying new *cis*-eGenes linked to disease, connecting these *cis*-eGenes with *trans*-eGenes to identify the regulatory intermediates that tie them together, and uncovering the cell types in which these variants act. However, several limitations should be considered. First, QTL discovery power and the accuracy of causal variant fine-mapping differ among the cell types, as the total number of cells and cells per donor used for QTL mapping differed widely among the 6 major cell types assessed. These differences across cell types will likely impact our ability to correctly assign cell type-specificity of an eQTL. Therefore, to overcome this issue, instead we assessed the sharedness of eQTL effects at lineage rather than cell type level, as we expected that eQTL effects are most likely shared within the same lineage (lymphoid or myeloid)[45]. Second, technical heterogeneity across cohorts (such as differences in PBMC isolation, preservation, and

preparation) likely introduced variability in cell type composition that may exceed the effect size of
individual genetic variants. While this variability could have reduced power for ccQTL detection, the
federated meta-analysis design and stringent heterogeneity filtering make spurious associations
unlikely. Third, some ccQTLs may indirectly affect expression, resulting in an eQTL effect entirely
driven by changes in cell (sub)type composition[29]. While this is mainly a risk for bulk eQTL studies,
which use samples composed of many different cell types[6,46], our sc-eQTL meta-analyses may have
been impacted by similar cell subtype changes. As a potential example, we described the ccQTL on
CD8+ TEM over total CD8+ T cells that also resulted in an eQTL effect on *LINC00299*, a long non-
coding RNA primarily expressed in the CD8+ TEM subtype. To address this, future efforts of the
consortium will further expand the donor and cell sample sizes to facilitate sufficient discovery power
to map eQTLs at higher cell subtype resolution. Finally, while the current study mapped eQTLs at cell
type resolution, several recent sc-QTL studies have shown that additional eQTLs can be identified
when leveraging cell subtypes or dynamic cell state annotations instead[29,47]. This is a unique aspect of
single-cell data that could provide additional resolution over previous bulk efforts, and it would
greatly benefit from the large amount of single-cell data available within the sc-eQTLGen consortium.
However, this is an aspect we would like to address in future consortium efforts, as the federated
meta-analysis setup complicates uniform assignment of cells along a dynamic trajectory.

In summary, this work highlights the benefits of integrating many sc-eQTL studies within a unified
meta-analysis framework. Such framework not only improves the QTL discovery power, but also
eliminates the need to share privacy-sensitive data or to rerun analyses whenever new datasets are
included. We expect that this work will benefit from further increases in sample size in the near
future and from combining the discovery power of bulk eQTL data with the cellular and context-
specific resolution of sc-eQTL data through the combined efforts of the eQTLGen[6] and sc-eQTLGen
consortia[14]. This will help us interpret disease-associated genetic variation and aid in the
implementation of personalized disease prevention and treatment.
591

Methods

Datasets

Data from 12 cohorts was meta-analyzed using harmonized computational pipelines (see below). The data derived from these cohorts is generated using diverse single-cell technologies and chemistries—including full-length Smart-seq2, as well as 3ʹ- and 5ʹ-end 10x Genomics scRNA-seq and sc-multiome platforms—and spanning individuals of multiple ancestries, age groups, and both sexes (**Table S1**). Detailed information on the data generation of the scRNA-seq and genotype information is provided in the **Supplementary Information**. Notably, scRNA-seq alignment to the reference genome was not harmonized across the studies: all the 10X studies followed (different versions of) the 10X Genomics CellRanger pipeline, while details of alignment of the Smart-seq2 data of the ARMS cohort can be found in the original publication[16].

Genotype imputation

Genotype imputation was carried out for each dataset separately on GRCh38 using a Snakemake (v6.4.1)[48] pipeline. Briefly, sample quality and consistency were ensured by removing samples missing a large amount of genotype calls (call rate >0.9) and ensuring consistency between genotype-determined and self-reported sex (when provided). If necessary, genotypes were lifted over to build 38, using CrossMap v0.4.1[49]. Next, donor genomes were identified to be most similar to African, Native American, European, East Asian or South Asian ancestries using 1000 Genomes samples as a reference[49]. Samples were grouped by genetic ancestry and imputed in each group separately, leveraging the 30X WGS 1000G reference panel[49]. Post imputation genotype quality was ensured by removing variants with a call rate < 0.99, and a MACH R2 quality score < 0.6. For QTL mapping we removed multiallelic variants, and per study we removed SNPs that had a MAF < 1%, had a HWE P < 1×10^{-4} and ensured that every genotype was observed in every study at least once. At the meta-analysis stage, all SNPs with a MAF < 5% over the whole study were removed, leveraging information calculated using Genotype-IO (https://github.com/molgenis/systemsgenetics/wiki/Genotype-IO). Sample and genotype filtering were carried out using plink 1.9 (v1.90b6.24) and plink 2 (v2.00a3LM)[50], bcftools (v1.10.2)[51], vcftools (v0.1.17)[52], Genotype Harmonizer (v1.4.23)[53], and R (v3.6.3). Eagle (v2.4.1)[54] was used to phase the genotypes before imputation with Minimac4 (v1.0.2)[55] using the 1000 Genomes 30x WGS dataset as imputation reference.

Demultiplexing and Doublet Removal

Souporcell (v2)[56] and Demuxlet[57] were used to demultiplex pools that contained multiple unrelated donors and identify heterotypic doublets. The transcription-based doublet detecting methods DoubletDetection (v3.0)[58], scds (v1.1.2)[59], and scrublet (v0.2.1)[60] were additionally used to identify homotypic doublets (those from the same donors). Droplets were classified as singlets if all software packages identified the droplet as a singlet and both Demuxlet and Souporcell annotated the droplet to the same donor.

Cell type assignment

We predicted the cell type label of the cells using two different methods: Azimuth[26] and HierscPred (v1.9.2)[25,27]. For Azimuth, we followed the multimodal reference mapping vignette from Seurat (v4.0.5) (https://satijalab.org/seurat/articles/multimodal_reference_mapping). HierscPred is the hierarchical version of scPred. It trains a scPred classifier for every parent node in the hierarchy. During the prediction step, HierscPred starts at the root node and uses the corresponding scPred model to predict which child node the cell belongs to. If the probability of the predicted child node is higher than the threshold (default = 0, so no rejection option), we continued with making predictions using the child node's scPred classifier. We continued until we reach a leaf node or the prediction probability fell below the threshold. In the latter case, the assigned label is the label of the cell type we last predicted with sufficient confidence. Both methods use the Human-PBMC reference from Hao

et al. as the reference/training dataset[14]. This dataset is annotated at three resolutions (celltype.L1,
celltype.L2, and celltype.L3). Using both methods, we annotated the cells at the celltype.L2
resolution. Then, based on this L2 resolution, we manually assigned cells to their respective L1
resolution. This L1 resolution was used as input for the eQTL analysis in 6 major cell types: B cells (B
intermediate, B memory, B naïve, plasmablast), CD4+ T cells (CD4 CTL, CD4 naïve, CD4 proliferating,
CD4 TCM, CD4 TEM, Treg), CD8+ T cells (CD8 naïve, CD8 proliferating, CD8 TCM, CD8 TEM),
monocytes (CD14 mono, CD16 mono), NK cells (NKdim, NKbright, NK proliferating), and dendritic
cells (ASDC, cDC1, cDC2, pDC). Since both methods are subject to methodological limitations, we only
used cells for downstream analyses when the predictions were concordant at this L1 resolution. For
the ccQTL mapping we took the L2 annotations as derived from Azimuth as the L2 label, and
additionally included ILCs (L1), as well as the three other T cell groups (MAIT, gdT, dnT). As most
datasets were sequenced in batches, we tried to ensure that batch effects did not affect the results
by annotating the cells per batch. A singularity image to annotate the cells using both methods is
available here: https://github.com/sc-eQTLgen-consortium/WG2-pipeline-classification/tree/main.
657
scRNA-seq data preprocessing and quality control
We removed those cells that expressed fewer UMIs or RNA molecules than the median minus 2 times
the MAD of the entire dataset or had mitochondrial gene content percentage was larger than the
median plus 3 times the MAD. We also removed cells that were assigned ambiguously or as doublet
using a combination of tools, see **Demultiplexing and Doublet Removal**. Finally, we kept those cells
that were concordantly annotated with both hierarchical scPred[25,27] and Azimuth[26] trained on human
PBMC reference data, as defined previously[26], see **Cell type assignment**. After QC, we leveraged the
PFlog1PF normalization[61] to normalize the scRNA-seq data per cell type.
666
Linkage disequilibrium calculations
LD was calculated for each dataset separately using the Python (v3.8.12) implementation of
LDSTORE2 v2.0_x86_64[62]. For this, we generated the following files per chromosome: bgen, bgi, z,
sample, master, and bdose, as described in the documentation for LDSTORE2. We then calculated LD
in windows to reduce the overlap in the LD matrices, with the window size determined by the
distance between genes. We used a minimum window size of 5M bases, with the start of the window
1M bases before the start of a gene and the end of the window 1M bases after a gene. If there were
other genes located within a window, the end of the window was determined by the gene that is
furthest along in the window.
To store and share LD efficiently, we performed a principal component analysis (PCA) on the
LD matrix to reduce the dimensions of the matrix from variants x variants to samples-1 x variants.
This reduced matrix, as well as the components and means, were saved to allow for reconstruction of
the LD matrix at the central analysis site, where we selected the number of components to be n
samples minus one as this allowed for a near perfect reconstruction, at a fraction of the space of the
original LD matrices.
682
Cell type composition QTL mapping
For ccQTL analyses, we included all cells that were concordantly annotated to the L1 level, and did
not perform any additional QC on the cells. Moreover, we removed donors based on a PCA on the
genotype information, excluding genetic outliers based on 2x SD within the respective study. We
assessed the effect of genetics on different compositional resolutions: 1. lymphoid versus myeloid
lineage, 2. major cell type compared to total PBMCs, 3. minor cell type compared to total PMBCs, and
4. compared to the major cell type it belongs to. We limited our meta-analysis to 7 out of the 12
datasets, as we excluded the cerebrospinal fluid dataset (Sawcer CSF), one dataset with fewer than 40
donors (300BCG), datasets from multiple ancestries (Ye), and those that contained only a single cell
type (Trynka and ARMS). In the remaining datasets, we limited our analyses to donors that had at
least 350 PBMCs. We mapped ccQTLs by running a linear mixed effect model, implemented in LIMIX
QTL[28], on cell count ratios forced to a normal distribution (Gaussian-normalized), which we corrected

for age, sex, and total number of cells per donor (n_cells) by including these as fixed effects in the
linear model, and accounted for possible population substructure and repeated observations from
the same donor by including a PLINK[50] king kinship matrix (u_K) as a random effect as follows:
698
699 Cell type proportions ~ SNP + age + sex + n_cells + u_K + ε
700
eQTL mapping
All 12 datasets were used for eQTL mapping. Pseudobulk mean averaged expression data per cell
type (L1 resolution) and per donor was used as input for the eQTL mapping[28]. We removed donors
whose gene expression (Gaussian-normalized) deviated more than two times the SD from the mean
of both PC1 and PC2 within the respective study. For the Ye dataset, we deviated from this as this
dataset had samples from two ancestries (European and Asian). For this dataset, we included the first
two genetic PCs as additional covariates when mapping eQTLs. eQTL mapping was performed using
LIMIX QTL on Gaussian-normalized expression data, we included 10 expression PCs as fixed effects
and added the PLINK king kinship matrix (u_K) as a random effect.
710
711 LIMIX eQTL model:
712 Cell type composition ~ SNP + expression PC[1-10] + u_K + ε
713 For the Ye dataset:
714 Cell type composition ~ SNP + expression PC[1-10] + genetic PC[1-2] + u_K + ε
715
Each dataset performed 100 permutations for the eQTL mapping, after which the permuted and eQTL
data were separately meta-analyzed at the central analysis site. To efficiently share the permutation
information we calculated Z-scores of the associations (beta/beta_se) for the permuted associations,
and fitted a normal distribution over these per SNP. The mean and variance of each association were
shared to allow further SNP selection at the central analysis site. At the central site the permutation
where combined. Specifically, we sampled from the distributions per association, leveraging the in
sample LD, to get back to a full permutation matrix per study. These were meta-anlyzed as outlined in
the meta-analysis section. Per gene, per permutation round we selected the top associated variant,
and leveraged the beta-distribution to increase resolution when calculating the permuted P value per
association as outlined in the FastQTL paper[63].
726
Meta-analysis
Both the ccQTLs and the eQTLs were meta-analyzed at the central analysis site. Here, final SNP and
trait selection was performed over all datasets at once (see **genotype QC**). For ccQTL meta-analysis
no further filtering was needed as all traits were shared over the datasets. However, for eQTL meta-
analysis, we first subsetted our results to genes expressed in all 12 datasets. For the ccQTL meta-
analysis, we used an inverse variance based meta-analysis, the top performing meta-analysis in our
benchmarks[64]. In contrast, for the eQTL meta-analysis a sample-size-based weighted meta-analysis[64]
was used instead, as this significantly reduced the amount of data that had to be shared between the
sites (inverse variance weighting requires sharing of standard errors per test). To filter associations
that are heterogeneous across the datasets we applied the I2 variation filter[65] after ccQTL mapping,
removing effects with I2 over 40 percent.
738
Fine-mapping and colocalization analysis
SuSiE was used for fine-mapping[31], setting the maximum number of credible sets per eGene to 5,
leaving the other values at default. For traits without a specific genomic location associated to it, i.e.
GWAS traits or ccQTLs, we leveraged the Locus Breaker methodology[30] to define loci that are used as
input for fine-mapping. Throughout the manuscript we refer to these credible sets as independent
loci. When SuSiE failed to converge, we defaulted back to reporting a single independent locus per
eGene and these effects were not taken forward for colocalizations. We used in sample LD for this
procedure, to derive this we meta-analyzed the dataset-specific LD by calculating the sample size-

weighted mean LD over the datasets. For the eQTLGen (phase 2)[6] *cis*-eQTLs we leveraged the same
procedure, where we subsetted the eQTLGen summary statistics down to the variants considered in
sc-eQTLGen, and used the in sample LD of sc-eQTLGen. This limited differences between the
finemapping results for the downstream colocalization. For *trans*-eQTL colocalization, we leveraged
the results as taken from eQTLGen to match the disease colocalization in eQTLGen and limit the
computational burden due to re-finemapping. To colocalize *cis*-eQTLs to each other, as well as for
colocalizing *cis-trans* effects we used the coloc package[32], in which the log bayes factor (LBF)
information from SuSiE was used as input, specifically we used the function: "coloc.bf_bf" (from
coloc), leveraging default parameters.
To colocalize the sc-eQTLGen *cis*-eQTLs to the Open Targets data we converted the SuSiE[31]
output into StudyLocus format from gentropy v2.0 (https://github.com/opentargets/gentropy) to be
consistent with the GWAS credible data format in Open Targets. These independent loci were then
overlapped with all GWAS independent loci in the Open Targets ecosystem (March 2025 data
release). We considered overlap when two independent loci (GWAS vs eQTL) share at least one
variant. The LBFs for each variant were then used to calculate coloc posterior probabilities, where we
infer an LBF of 0 for variants which do not exist in one of the two independent loci of an overlap. All
analyses were performed on Google Cloud Platform via Airflow using our pipeline orchestration. The
mathematics in the coloc implementation is the same as the original coloc R package[32], and is
previously benchmarked against that to ensure it produces the same output. Independent loci were
filtered such that the minimal LBF was 2, or >= 0.8686 in log10BF scale, and the independent loci had
at least one association at $P \leq 5.0 \times 10^{-3}$. For all colocalization analyses we defined traits to be
colocalized if the sum of PP H3 and PP H4 was higher than 0.9 and the ratio between PPH4 and PPH3
was higher than 3.
770
ccQTL validation
We validated the ccQTLs by looking up the variants within the ccQTL CSs with UK Biobank cell count
GWASes[3] and whole blood *trans*-eQTLs from eQTLGen (phase 2)[6]. For the latter, we calculated a
UCell[33] module activity score per cell (in the UMCG V3 dataset as representative example) for each
independent ccQTL locus. First we colocalized (see **finemapping and colocalization**) each ccQTL locus
with *trans*-eQTLs derived from eQTLGen. Next we split up the colocalizing *trans*-eQTLs based on the
effect sign, and derived two UCell module scores[33], one for the positive *trans*-eQTLs and one for the
negative *trans*-eQTLs. We subtracted the module score of negatively associated *trans*-eGenes from
that of positively associated ones, to get to the final UCell score. For this calculation we required at
least two colocalizing trans-eQTL effects per ccQTL loci. We then used a Wilcoxon Signed-Rank test to
define whether this module activity score was significantly different in the cells that changed
composition relative to all other cells over which the ccQTL was calculated (i.e. versus all PBMCs, or
versus the other cells within the same major group depending on the effect), and we assessed
whether the direction of effect was the same as the ccQTL effect.
785
786 **ccQTL inflation**
787 To quantify the relation between ccQTLs and GWAS variants, we calculated the λ-inflations.
788 Specifically, we used the fine-mapped variants from the disease GWAS as taken from the Open
789 Targets fine-mapping, and calculated the λ-inflation[66] for the ccQTL variants that overlap with GWAS
790 variants.
791
Annotation of GWAS-linked independent loci
To assess which features of our independent sc-eQTLs are shared with loci associated to GWAS or
associated to bulk *cis*-eQTLs from eQTLGen, we leveraged two logistic regression models. One
predicting independent loci with disease GWAS colocalization (yes = 1, no = 0), and another one
predicting loci that are shared with bulk *cis*-eQTLs from eQTLGen effect (yes = 1, no = 0). As predictors
we included: the disease EFO category, and the number of cell types and genes an independent locus
is identified in; whether the locus is shared among the two major linages (lineage mixing) and shared

with bulk *cis*-eQTLs from eQTLGen. The logistic regression models were fitted with the glm function in
R 4.2.1.
801
Variant-based GWAS enrichment of eQTLs
All fine-mapped eQTL loci were used for variant-based GWAS enrichment. We annotated these
variants with GWAS variants with genome-wide significant ($P < 5 \times 10^{-8}$) associations (v1.0.2)
downloaded from the GWAS Catalog, while considering LD. We used EFO terms to group variants into
different trait groups. We mapped the GWAS traits to EFO-IDs[10] and corresponding parent terms from
the GWAS catalog[67] using gwasrapidd v0.99.17. The different trait categories were defined as follow:
- All traits: no filtering; all traits in GWAS catalog
- Disease GWAS: only kept traits assigned to these EFO parent terms: 'Cancer (EFO_0000616)',
  'Immune system disorder (EFO_0000540)', 'Other disease (EFO_0000408)', 'Cardiovascular
  disease (EFO_0000319)', 'Neorological disorder (EFO_0000618)', 'Metabolic disorder
  ('EFO_0000589')
- Immune disease GWAS: only kept traits assigned to these EFO parent terms:'Immune system
  disorder (EFO_0000540)'
To analyze the enrichment across all cell types ('All' category) we used only one cell type observation
per eGene. If an eGene had an eQTL effect in multiple cell types, the observation of the cell type with
the lowest p-value of the eQTL effect was used.
In sample LD information was used to filter the eQTL and complete set of assessed variants
(R2 ≥ 0.1) as well as determining whether the GWAS variants are in LD (R2 ≥ 0.8). The eSNPs were
filtered such that the most significant variant in high LD was kept. eSNPs are determined to be GWAS
variants either if they are present in the GWAS catalog set of GWAS variants or if they are in LD with
any of the GWAS variants.
823
eGene enrichment analysis
To gain insight into the potential upstream regulator that may be associated to the *trans*-eGenes that
were linked to the same *cis*-eGenes, we performed gene set enrichments using the R package
g:profiler[68]. We tested only *trans*-eGenes that formed *cis-trans* gene-gene pairs in which the same
cis-eGene was linked to at least 9 other *trans*-eGenes. All settings were left at defaults provided by
the tool developers, other than the background sets. The background gene set consisted of all genes
that showed at least one *trans*-eQTLs in eQTLGen.
Additionally, to get higher resolution on the relevance of TFs we prepared specific test information by
parsing the information from the ReMap project[69]. Specifically, we added three extra information
sources, we added the *cis*-regulatory modules (CRMs) as defined by ReMap, the non-redundant TF
information over all cell types and the non-redundant TF information for blood relevant cell types[70].
These information sources were added by providing custom data to g:profiler[68].
836

837 Code availability
838 The pipelines for (WG1) genotype imputation and scRNA-seq data preprocessing, QC and donor
839 demultiplexing; (WG2) cell type assignment; (WG3) eQTL mapping and meta-analysis and
840 downstream analyses, can be found on Github (https://github.com/sc-eQTLgen-consortium). The
841 pipelines are established in Snakemake[48], and all software is provided in a Singularity[71] or Docker[72]
842 image for consistency across datasets.
843

844 Data availability
845 Full eQTL summary statistics will be made available after publication the sc-eQTLGen consortium
846 webpage: https://www.eqtlgen.org/sc/. eQTLGen summary statistics will be publicly available upon
847 acceptance of the respective paper.
848

849 Ethical compliance

All cohorts included in this study enrolled participants with informed consent and collected and
analyzed data in accordance with ethical and institutional regulations. The information about
individual institutional review board approvals is available in the original publications for each cohort.
All procedures performed in studies involving human participants were in accordance with the ethical
standards of the institutional and/or national research committee and with the 1964 Helsinki
declaration and its later amendments or comparable ethical standards.

Acknowledgments

We thank all the children who participated in the Burkina Faso study and their families. We thank the
staff of the Centre National de Recherche et de Formation sur le Paludisme and of the Groupe de
Recherche Action en Santé for facilitating sample collection and clinical work. We acknowledge
support from the National Institute for Health Research (NIHR) Cambridge Biomedical Research
Centre, United Kingdom. The images in Figure 4f are created using Servier Medical Art
(https://smart.servier.com), licensed under CC BY 4.0. We thank Katherine McIntyre for the English
editing of our manuscript. The authors wish to acknowledge the contributions of all participating
cohorts and study participants.

Funding

L.F., D.K., M.J.B. are supported by a grant from the Dutch Research Council (ZonMW-VICI
09150182010019 to L.F. ZonMW LongCOVID grant 10430302110002), European Union's Horizon
Europe Research and Innovation Program grant 101057553 (LongCovid) and through a Senior
Investigator Grant from the Oncode Institute and a grant from Saxum Volutum (Pericode). M.J.B. is
supported by a grant from the Dutch Research Council (ZonMW-VIDI 09150172310068). M.W. is
supported by a grant from the Dutch Research Council (ENW-VIDI 223.041). The Sawcer CSF study
was supported by funding from the Multiple Sclerosis Society in the UK, grant reference 43 to M.Ban
and S.S. The Burkina Faso malaria study is supported by NYUAD Grant AD105 to Y.I. and the resources
to run the sc-eQTL analysis by Tamkeen under the Research Institute, NYUAD Grant G1206 to Y.I. G.T.
and N.K. were supported by the Wellcome Trust 220540/Z/20/A and Open Targets (OTAR2064). Y.M.
and H.N. are supported in part by the Qatar National Research Fund (QNRF Awards PPM1-1122-
150008) and internal funds from Sidra Medicine. L.M and A.M. were supported by an NWO
Gravitation grant 024.004.012. This research was also supported by the Chan Zuckerberg Initiative
through grant 2021-239720 (5022) to M.M. This work was supported by the Estonian Research
Council grant (PSG1230) to U.V.

The eQTLGen consortium is supported in part by NWO ZonMW-VENI Grant (no. 9150161910057 and
09150162310207), the Estonian Research Council (MOB3ERC115), the German Federal Ministry of
Education and Research (BMBF) within the framework of the e:Med research and funding concept
(grant # 01ZX1906B & 01ZX2206B), the Deutsche Forschungsgemeinschaft (DFG, German Research
Foundation) – 542489987, the Innovative Medicines Initiative grant 3TR (GA#831434), the Innovative
Medicines Initiative grant 3TR (GA#831434), the Intramural Research Program of the National Cancer
Institute, National Institutes of Health, US Department of Health and Human Services, the core
funding from the Wellcome Trust [220540/Z/20/A], the Innovative Medicines Initiative Joint
Undertaking (grant 115565), the Spanish Ministry of Science and Education (PID2022-13929208-I00),
the Academy of Finland (Grant number: 338395), a R01 AG075884 grant, the European Research
Council under the European Union's Horizon 2020 research and innovation programme (grant
agreement n° 772376 and 101198904 and 101017802), the Diabetes Fonds (Junior Fellowship
2023.81.006), the Medical Research Council (MR/V002503/1), Wellcome Trust Investigator Award
(204969/Z/16/Z), Wellcome Trust Grants (090532/Z/09/Z and 203141/Z/16/Z), the Chinese Academy
of Medical Sciences Innovation Fund for Medical Science (grant number: 2018-I2M-2-002), the Swiss
National Science Foundation grant 10003484 and Novo Nordisk and the Fondation pour la recherche
sur le diabete, the JSPS KAKENHI (25H01057), and AMED (JP24km0405217, JP24ek0109594,

JP24ek0410113, JP24kk0305022, JP223fa627001, JP223fa627002, JP223fa627010, JP223fa627011, JP22zf0127008, JP24tm0524002, JP24wm0625504, JP24gm1810011), JST Moonshot R&D (JPMJMS2021, JPMJMS2024), Takeda Science Foundation, Ono Pharmaceutical Foundation for Oncology, Immunology, and Neurology, Bioinformatics Initiative of Osaka University Graduate School of Medicine, Institute for Open and Transdisciplinary Research Initiatives, Center for Infectious Disease Education and Research, and Center for Advanced Modality and DDS, Osaka University, RIKEN TRIP initiative, the Innovative Medicines Initiative Joint Undertaking (grant 115565), the core funding from the British Heart Foundation (RG/F/23/110103), NIHR Cambridge Biomedical Research Centre (NIHR203312) [*], BHF Chair Award (CH/12/2/29428), Cambridge BHF Centre of Research Excellence (RE/24/130011), and by Health Data Research UK, which is funded by the UK Medical Research Council, Engineering and Physical Sciences Research Council, Economic and Social Research Council, Department of Health and Social Care (England), Chief Scientist Office of the Scottish Government Health and Social Care Directorates, Health and Social Care Research and Development Division (Welsh Government), Public Health Agency (Northern Ireland), British Heart Foundation and the Wellcome trust, the Intramural Research Program of the National Institutes of Health and NIH grant R35GM139580 [#], the KNAW Academy Professor Award (PAH/6635), a R01 MH115676 and R01 MH115676 grant, a U01 HG012009 grant, the Estonian Research Council (MOB3ERC115, PRG1911 and TT17), the Swiss National Science Foundation (315230_219587), the Ministry of Education and Research Centres of Excellence grant TK214 name of CoE, the European Regional Development Fund, the programme Mobilitas Pluss (MOBTP108).

*The views expressed are those of the authors and not necessarily those of the NIHR or the Department of Health and Social Care.
The contributions of the NIH author(s) were made as part of their official duties as NIH federal employees, are in compliance with agency policy requirements, and are considered Works of the United States Government. However, the findings and conclusions presented in this paper are those of the author(s) and do not necessarily reflect the views of the NIH or the U.S. Department of Health and Human Services.

Author contributions

Conceived and designed the computational framework: D.K., L.M., D.N., A.R., O.S., M.Hemberg, A.M., M.M., J.P., M.J.B., L.F., M.W.

Analyzed the data: D.K., L.M., D.N., A.R., J.A., M.K., J.T.H.L., R.O., M.V., R.W., Y.A., M.Ban, O.B., M.Berg, I.B., M.D., R.E., M.G.G., H.G., M.Heinig, P.H., C.H., Y.I., P.K., N.K., W.L., Y.L., C.L., V.M., J.M., M.C.N., M.N., J.N., Y.O., S.S., I.S., W.P., D.R., T.S., J.S., G.T., U.V., S.Y., J.Y., Z.Z., eQTLGen consortium, M.J.B., M.W.

Performed statistical analysis: D.K., L.M., D.N., A.R., D.C., M.Heinig, C.L., Y.T., G.T.

Contributed analysis tools: D.K., L.M., D.N., A.R., J.A., M.K., J.T.H.L., R.O., M.V., R.W., D.C., Y.T., G.T.

Contributed dataset: D.K., M.K., R.O., Y.A., M.Ban, O.B., M.Berg, I.B., M.D., R.E., M.G.G., H.G., P.H., C.H., Y.I., P.K., N.K., W.L., Y.L., V.M., J.M., H.N., Y.M., M.C.N., M.N., J.N., Y.O., S.S., I.S., W.P., D.R., T.S., J.S., G.T., U.V., S.Y., J.Y., Z.Z., eQTLGen, J.P. M.J.B., L.F., M.W.

Jointly supervised research: M.J.B., L.F., M.W.

Wrote the paper: M.J.B., M.W.

Competing interests

946 M.V. is currently a full-time employee of Illumina and owns stock in the company; these interests
947 began after M.V.'s contribution to this work was completed. L.F. has ongoing contract-based research
948 with Biogen and Roche, not related to this work.
949

Figures

Tables

Supplementary Figures

Supplementary Tables

References

1. Sollis, E. *et al.* The NHGRI-EBI GWAS Catalog: knowledgebase and deposition resource. *Nucleic*
Acids Res. **51**, D977–D985 (2023).
2. Akbari, P. *et al.* A genome-wide association study of blood cell morphology identifies cellular
proteins implicated in disease aetiology. *Nature Communications* **14**, 5023 (2023).
3. Vuckovic, D. *et al.* The Polygenic and Monogenic Basis of Blood Traits and Diseases. *Cell* **182**,
1214-1231.e11 (2020).
4. Orrù, V. *et al.* Complex genetic signatures in immune cells underlie autoimmunity and inform
therapy. *Nature Genetics* **52**, 1036–1045 (2020).
5. The GTEx Consortium *et al.* The GTEx Consortium atlas of genetic regulatory effects across
human tissues. *Science* **369**, 1318–1330 (2024).
6. Warmerdam, R., eQTLGen consortium, Franke, L. & Võsa, U. Trans-eQTLs reveal the architecture
of human gene regulatory networks. *biorXiv* (2026).
7. Natri, H. M. *et al.* Cell-type-specific and disease-associated expression quantitative trait loci in the
human lung. *Nat.Genet.* **56**, 595–604 (2024).
8. Oelen, R. *et al.* Single-cell RNA-sequencing of peripheral blood mononuclear cells reveals
widespread, context-specific gene expression regulation upon pathogenic exposure. *Nature*
Communications **13**, 3267 (2022).
9. Hormozdiari, F. *et al.* Leveraging molecular quantitative trait loci to understand the genetic
architecture of diseases and complex traits. *Nat.Genet.* **50**, 1041–1047 (2018).
10. Mostafavi, H., Spence, J. P., Naqvi, S. & Pritchard, J. K. Systematic differences in discovery of
genetic effects on gene expression and complex traits. *Nat.Genet.* **55**, 1866–1875 (2023).
11. Umans, B. D., Battle, A. & Gilad, Y. Where Are the Disease-Associated eQTLs? *Trends Genet.* **37**,
109–124 (2021).
12. Perez, R. K. *et al.* Single-cell RNA-seq reveals cell type–specific molecular and genetic
associations to lupus. *Science* **376**, eabf1970 (2022).
13. Yazar, S. *et al.* Single-cell eQTL mapping identifies cell type–specific genetic control of
autoimmune disease. *Science* **376**, eabf3041 (2022).
14. van der Wijst, M. *et al.* The single-cell eQTLGen consortium. *Elife* **9**, 10.7554/eLife.52155 (2020).
15. Soskic, B. *et al.* Immune disease risk variants regulate gene expression dynamics during CD4+ T
cell activation. *Nat.Genet.* **54**, 817–826 (2022).
16. Vieira Braga, F. A. *et al.* A cellular census of human lungs identifies novel cell states in health and
in asthma. *Nat.Med.* **25**, 1153–1163 (2019).
17. Li, W. *et al.* A single-cell view on host immune transcriptional response to in vivo BCG-induced
trained immunity. *Cell.Rep.* **42**, 112487 (2023).
18. Edahiro, R. *et al.* Single-cell analyses and host genetics highlight the role of innate immune cells in
COVID-19 severity. *Nat.Genet.* **55**, 753–767 (2023).
19. Namkoong, H. *et al.* DOCK2 is involved in the host genetics and biology of severe COVID-19.
Nature **609**, 754–760 (2022).
20. Ban, M. *et al.* Expression profiling of cerebrospinal fluid identifies dysregulated antiviral
mechanisms in multiple sclerosis. *Brain* **147**, 554–565 (2024).
21. Shahin, T. *et al.* Single-cell transcriptomics reveals inter-ethnic variation in immune response to
Falciparum malaria. *The American Journal of Human Genetics* **112**, 709–723 (2025).
22. van der Wijst, M. G. P. *et al.* Single-cell RNA sequencing identifies celltype-specific cis-eQTLs and
co-expression QTLs. *Nat.Genet.* **50**, 493–497 (2018).

23. van Blokland, I. V. *et al.* Single-Cell Dissection of the Immune Response After Acute Myocardial
Infarction. *Circ.Genom.Precis Med.* **17**, e004374 (2024).
24. Neavin, D. *et al.* Demuxafy: improvement in droplet assignment by integrating multiple single-cell
demultiplexing and doublet detection methods. *Genome Biol.* **25**, 94 (2024).
25. Alquicira-Hernandez, J., Sathe, A., Ji, H. P., Nguyen, Q. & Powell, J. E. scPred: accurate
supervised method for cell-type classification from single-cell RNA-seq data. *Genome Biol.* **20**, 264
(2019).
26. Hao, Y. *et al.* Integrated analysis of multimodal single-cell data. *Cell* **184**, 3573-3587.e29 (2021).
27. Michielsen, L., Reinders, M. J. T. & Mahfouz, A. Hierarchical progressive learning of cell identities
in single-cell data. *Nature Communications* **12**, 2799 (2021).
28. Cuomo, A. S. E. *et al.* Optimizing expression quantitative trait locus mapping workflows for single-
cell studies. *Genome Biol.* **22**, 188 (2021).
29. Rumker, L. *et al.* Identifying genetic variants that influence the abundance of cell states in single-
cell data. *Nature Genetics* **56**, 2068–2077 (2024).
30. Pirastu, N. *et al.* GWAS for male-pattern baldness identifies 71 susceptibility loci explaining 38% of
the risk. *Nature Communications* **8**, 1584 (2017).
31. Wang, G., Sarkar, A., Carbonetto, P. & Stephens, M. A Simple New Approach to Variable Selection
in Regression, with Application to Genetic Fine Mapping. *J.R.Stat.Soc.Ser.B.Stat.Methodol.* **82**,
1273–1300 (2020).
32. Pullin, J. M. & Wallace, C. Variant-specific priors clarify colocalisation analysis. *PLOS Genetics* **21**,
1–23 (2025).
33. Andreatta, M. & Carmona, S. J. UCell: Robust and scalable single-cell gene signature scoring.
Computational and Structural Biotechnology Journal **19**, 3796–3798 (2021).
34. Sobczyk, M. K. & Gaunt, T. R. Integrative Mendelian Randomization approaches for therapeutic
target prioritisation in immune-mediated diseases. *medRxiv* 2024.04.27.24306475 (2024)
doi:10.1101/2024.04.27.24306475.
35. Zhang, Z. E., Kim, A., Suboc, N., Mancuso, N. & Gazal, S. Efficient count-based models improve
power and robustness for large-scale single-cell eQTL mapping. *medRxiv* 2025.01.18.25320755
(2025) doi:10.1101/2025.01.18.25320755.
36. Qi, T. *et al.* Identifying gene targets for brain-related traits using transcriptomic and methylomic
data from blood. *Nature Communications* **9**, 2282 (2018).
37. Ghoussaini, M. *et al.* Open Targets Genetics: systematic identification of trait-associated genes
using large-scale genetics and functional genomics. *Nucleic Acids Res.* **49**, D1311–D1320 (2020).
38. Dönertaş, H. M., Fabian, D. K., Valenzuela, M. F., Partridge, L. & Thornton, J. M. Common genetic
associations between age-related diseases. *Nat Aging* **1**, 400–412 (2021).
39. Rex, D. A. B. *et al.* A comprehensive pathway map of IL-18-mediated signalling. *Journal of Cell
Communication and Signaling* **14**, 257–266 (2020).
40. Liao, B.-W. *et al.* FAM177A1 Inhibits IL-1β–Induced Signaling by Impairing TRAF6–Ubc13
Association. *The Journal of Immunology* **207**, 3090–3097 (2021).
41. Trzeciak, M. *et al.* Relationship between serum levels of interleukin-18, IgE and disease severity in
patients with atopic dermatitis. *Clinical and Experimental Dermatology* **36**, 728–732 (2011).
42. Ramonell, R. P. *et al.* CD8+ TEMRAs in severe asthma associate with asthma symptom duration
and escape proliferation arrest. *JCI Insight* **10**, (2025).
43. McGuire, D. J. *et al.* Regulation of CD45 isoforms during human effector and memory CD8 T cell
differentiation: Implications for T cell nomenclature. *Proc Natl Acad Sci U S A* **122**, e2322982122
(2025).
44. Kim-Hellmuth, S. *et al.* Cell type-specific genetic regulation of gene expression across human
tissues. *Science* **369**, (2020).
45. Peters, J. E. *et al.* Insight into Genotype-Phenotype Associations through eQTL Mapping in
Multiple Cell Types in Health and Immune-Mediated Disease. *PLoS Genet.* **12**, e1005908 (2016).
46. Võsa, U. *et al.* Large-scale cis- and trans-eQTL analyses identify thousands of genetic loci and
polygenic scores that regulate blood gene expression. *Nat.Genet.* **53**, 1300–1310 (2021).
47. Popp, J. M. *et al.* Cell type and dynamic state govern genetic regulation of gene expression in
heterogeneous differentiating cultures. *Cell Genom* **4**, 100701 (2024).
48. Mölder, F. *et al.* Sustainable data analysis with Snakemake. *F1000Res* **10**, 33 (2021).
49. Byrska-Bishop, M. *et al.* High-coverage whole-genome sequencing of the expanded 1000
Genomes Project cohort including 602 trios. *Cell* **185**, 3426-3440.e19 (2022).
50. Chang, C. C. *et al.* Second-generation PLINK: rising to the challenge of larger and richer datasets.
Gigascience **4**, 7-8. eCollection 2015 (2015).

51. Danecek, P. *et al.* Twelve years of SAMtools and BCFtools. *Gigascience* **10**, giab008. doi:
10.1093/gigascience/giab008 (2021).
52. Danecek, P. *et al.* The variant call format and VCFtools. *Bioinformatics* **27**, 2156–2158 (2011).
53. Deelen, P. *et al.* Genotype harmonizer: automatic strand alignment and format conversion for
genotype data integration. *BMC Res.Notes* **7**, 901–901 (2014).
54. Loh, P.-R. *et al.* Reference-based phasing using the Haplotype Reference Consortium panel.
Nat.Genet. **48**, 1443–1448 (2016).
55. Das, S. *et al.* Next-generation genotype imputation service and methods. *Nat.Genet.* **48**, 1284–
1287 (2016).
56. Heaton, H. *et al.* Souporcell: robust clustering of single-cell RNA-seq data by genotype without
reference genotypes. *Nat.Methods* **17**, 615–620 (2020).
57. Kang, H. M. *et al.* Multiplexed droplet single-cell RNA-sequencing using natural genetic variation.
Nat.Biotechnol. **36**, 89–94 (2018).
58. Gayoso, A., Shor, J., Carr, A. J., Sharma, R. & Pe'er, D. GitHub: DoubletDetection.
https://doi.org/10.5281/zenodo.2678041 (2020) doi:http://doi.org/10.5281/zenodo.2678041.
59. Bais, A. S. & Kostka, D. scds: computational annotation of doublets in single-cell RNA sequencing
data. *Bioinformatics* **36**, 1150–1158 (2020).
60. Wolock, S. L., Lopez, R. & Klein, A. M. Scrublet: Computational Identification of Cell Doublets in
Single-Cell Transcriptomic Data. *Cell.Syst.* **8**, 281-291.e9 (2019).
61. Booeshaghi, A. S., Hallgrímsdóttir, I. B., Gálvez-Merchán, Á. & Pachter, L. Depth normalization for
single-cell genomics count data. *bioRxiv* 2022.05.06.490859 (2022)
doi:10.1101/2022.05.06.490859.
62. Benner, C. *et al.* Prospects of Fine-Mapping Trait-Associated Genomic Regions by Using
Summary Statistics from Genome-wide Association Studies. *Am.J.Hum.Genet.* **101**, 539–551
(2017).
63. Ongen, H., Buil, A., Brown, A. A., Dermitzakis, E. T. & Delaneau, O. Fast and efficient QTL mapper
for thousands of molecular phenotypes. *Bioinformatics* **32**, 1479–1485 (2016).
64. Korshevniuk, M. *et al.* Optimized summary-statistic-based single-cell eQTL meta-analysis.
Scientific Reports **15**, 28407 (2025).
65. Higgins, J. P. T. & Thompson, S. G. Quantifying heterogeneity in a meta-analysis. *Stat Med* **21**,
1539–1558 (2002).
66. Devlin, B. & Roeder, K. Genomic control for association studies. *Biometrics* **55**, 997–1004 (1999).
67. Cerezo, M. *et al.* The NHGRI-EBI GWAS Catalog: standards for reusability, sustainability and
diversity. *Nucleic Acids Research* **53**, D998–D1005 (2025).
68. Raudvere, U. *et al.* g:Profiler: a web server for functional enrichment analysis and conversions of
gene lists (2019 update). *Nucleic Acids Research* **47**, W191–W198 (2019).
69. Hammal, F., de Langen, P., Bergon, A., Lopez, F. & Ballester, B. ReMap 2022: a database of
Human, Mouse, Drosophila and Arabidopsis regulatory regions from an integrative analysis of
DNA-binding sequencing experiments. *Nucleic Acids Res* **50**, D316–D325 (2022).
70. Kaptijn, D. *et al.* Disease-associated variants are enriched for altering cell-type-specific gene co-
expression relationships. *bioRxiv* 2025.09.06.674678 (2025) doi:10.1101/2025.09.06.674678.
71. Kurtzer, G. M., Sochat, V. & Bauer, M. W. Singularity: Scientific containers for mobility of compute.
PLoS One **12**, e0177459 (2017).
72. Merkel, D. Docker: lightweight Linux containers for consistent development and deployment. *Linux*
J. **2014**, (2014).