The attachment rate of *Vibrio anguillarum* strains to microplastics strongly varies with abiotic and biotic factors

Authors: Massot Méril[1], Wimmer Lukas[2,3], Mhamad Aly Moussawi[4,5], Hamet Jeanne[1,6], Tatjana N. Parac-Vogt[4], Dailey Lea Ann[2], Callens Martijn[7], Bedhomme Stéphanie[1]

Affiliations

[1] CEFE, CNRS, Univ Montpellier, EPHE, IRD, Montpellier, France

[2] Department of Pharmaceutical Sciences, University of Vienna, Vienna, Austria

[3] Vienna Doctoral School of Pharmaceutical, Nutritional and Sport Sciences (PhaNuSpo), University of Vienna, Vienna, Austria

[4] Department of Chemistry, KU Leuven, Celestijnenlaan 200F, 3001 Leuven, Belgium

[5] School of Chemistry, The University of Glasgow, Glasgow G11 6EW, United Kingdom

[6] HydroSciences Montpellier, University of Montpellier, CNRS, IRD, Montpellier, France

[7] Animal Science Unit, Flanders Research Institute for Agriculture, Fisheries and Food (ILVO), Oostende, Belgium

Abstract

Microplastics (MPs), resulting from plastic objects and waste degradation, are increasingly abundant, particularly in marine environments. They exhibit a hydrophobic surface on which biofilms form easily. Metagenomic analyses of these biofilms have revealed that they often contain bacterial species potentially pathogenic to humans or animals. For this reason, MPs are suspected to present a risk for public health by acting as a vector for pathogenic bacteria species. To better understand this hazard, we studied different factors potentially affecting the bacterial attachment rate to MPs. Focusing on the fish pathogen *Vibrio anguillarum,* a collection of 16 strains was assembled and GFP-labelled. Their attachment rates were measured using fluorescence microscopy on three types of MPs (milled polypropylene and polyethylene terephthalate particles and commercially available polystyrene beads). A strong effect of the particle type was found, likely linked to both the chemical composition of the particles and the surface characteristics, with higher attachment rates on rough particles. Our results also revealed a strong intra-specific variation in attachment rate, highlighting the need of testing several strains of the same species to assess attachment rate and related hazards.

34 Finally, it was observed that when a biofilm already formed on the MPs (by field-incubation of
35 the MPs along the Mediterranean French coast), differences in attachment rates between
36 particle types were erased. It was concluded that the attachment rate of *V. anguillarum* depends
37 on a combination of biotic and abiotic factors, which makes risk assessment of MPs as vectors
38 of pathogenic bacteria species a very complex task.
39
40 **Keywords**: microplastics, biofilm, marine pollution, Vibrio pathogen, plastisphere, intraspecific
41 variation, *Vibrio anguillarum*
42

Introduction

44
Approximately 9 billion tons of plastic objects have been manufactured since 1950, of
which more than half have been produced since the early 2000s [1]. Of all the discarded plastics
so far, some 14% have been incinerated and less than 10% recycled [1], whereas the vast
majority ends up in various ecosystems. As a result, plastic material can already be found
everywhere on the planet [2], [3], [4], [5], and is the largest source of debris produced on the
continents and ending up in the oceans [6].

Microplastics (fragments between 1 μm and 5 mm) and nanoplastics (fragments smaller
than 1 μm), are considered to be a concerning element of plastic pollution. Studies indicate they
can reach virtually all organs of the human body, e.g. the blood [7], the brain [8], the kidneys [8],
the placenta [9], and the digestive system [8], [10]. A growing body of literature raises concerns
regarding their cytotoxicity [11], [12] and broader health implications [13], [14], [15]. In addition
to the negative effects on animal and human health through physical damage and chemical
interactions, microplastics can present a risk through the molecules and living organisms that
adhere on them. Indeed, when immersed in seawater or freshwater, microplastics are rapidly
covered by a layer of proteins and other biomolecules, called the "ecocorona" [16]. This nutrient-
rich ecocorona attracts bacteria which can also adhere to the microplastics and form a biofilm.
The term 'plastisphere' has been coined to describe communities of microbes associated to
microplastics [17]. Several studies have detected an enrichment of potential pathogens in
biofilms formed on microplastics collected from seawater, compared to the water column [18],
[19], [20], [21]. However, many knowledge gaps have been identified and should be investigated
to confirm or disprove the assumption of microplastics acting as vectors for pathogens [22].

A first potential issue is that numerous *in situ* studies have detected potential pathogenic
bacteria within the plastisphere *via* 16S rRNA gene sequencing [23], [24], [25], [26], [27], [28].
This technique is essential for studying microbial communities, but lacks consistent taxonomic
granularity that allows identification at the species level, preventing conclusions about the actual
presence of pathogenic species [29]. Even when pathogenic species were detected *via* other
techniques, their viability or their virulence was not tested [19], [30], [31]. Moreover, strain
identity and diversity within pathogenic species of the plastisphere remain overlooked although
there can be a large variability of pathogenicity (ability to cause a disease) and virulence
(severity of the disease) within a bacterial species. Intraspecific variability in gene content
involved in motility and biofilm production has been described in several bacterial taxa [32], [33],
[34]. These functions partially determine the expression of virulence towards a host, as they
enable adhesion to its epithelia. Hence, genes involved in the adhesion to surfaces and
virulence often belong to the accessory genome and that there is a strong intra-specific
variability for the expression of these traits. Although studying intraspecific variability is key to
understand bacterial attachment to solid surfaces and biofilm formation, this has received less
attention than interspecific variability.

A second potential issue is that the term "microplastics" covers a huge diversity of
particles in terms of physical and chemical properties like polymer type, density, shape,
additives, surface regularity and chemistry (e.g. hydrophobicity), or fragmentation process [35].
The influence of these properties, alone and in combination, on the potential of microplastics as
a vector of pathogenic bacteria has been largely overlooked in previous studies. Furthermore,
the majority of microplastics studied to date were collected in marine environments [21], [36],
and consequently, these properties were not controlled.

Many reports on enrichment of the plastisphere with potential pathogens concern the
bacterial genus *Vibrio*. It contains several species causing severe diseases in humans and
marine animals, including the well-known causative agent of cholera, *Vibrio cholerae*. This
genus is usually found in the early stages of biofilm formation on microplastic surfaces, where it
becomes a minor member of mature biofilms [25], [26], [27]. However, a certain inter-study
variability remains.

The objective of this study was to investigate some of the factors driving the attachment
of pathogenic bacteria to microplastics. *Vibrio anguillarum* was chosen as the model species*,* as
it is lethal to fish and crustaceans and is responsible for substantial losses in aquaculture. More
specifically, four questions were asked: (1) Is there an intraspecific variation of the attachment

rate to microplastics in *V. anguillarum* and, if so, can we identify the genomic features behind
this variation? (2) Does the attachment rate vary between microplastic types (polymer and
shape)? (3) Does the presence of a natural biofilm, formed by in-field incubation, change the
attachment rate of *V. anguillarum*? (4) Does the attachment to microplastics increase the
resistance of *V. anguillarum* to bleach, a common water sanitizing treatment?

Material and methods

1. Bacterial strain and genomic analyses

1.1. Selection of *V. anguillarum* strains

108 To explore the intraspecific variability of attachment rates on MPs, 16 *V. anguillarum*
109 strains were obtained from the Belgian Coordinated Collection of Microorganisms (Table S1).
110 These strains were selected to represent a wide geographic range and diverse ecological
111 origins (isolated from infections in various host species or from environmental samples) (Figure
112 1A).

1.2. Fluorescent labelling of *V. anguillarum* strains

The pMRB plasmid [37] containing a green fluorescent protein (GFP) coding gene and a
chloramphenicol resistance gene as selection marker was introduced by conjugation in all *V.*
anguillarum strains (supplementary material). Briefly, 200 µL of *Escherichia coli* ß3914
(diaminopimelate-auxotrophic donor) containing pMRB-GFP plasmid was mixed with 800 µL of
recipient culture, centrifuged and spotted on top of a sterile cellulose acetate filter (0.45 µm pore
size). Filters were placed on LBA + NaCl 0.5M supplemented with diaminopimelate (0.3mM).
Plates were incubated overnight at 28 °C. Bacterial cells were resuspended in LB + NaCl 0.5M,
and *V. anguillarum* transconjugants were selected on LBA + NaCl 0.5M + chloramphenicol (5
µg/ml). For each *V. anguillarum* strain, single colonies were isolated, and their GFP expression
was assessed, using the fluorescence microscope Leica DM2000 LED (Leica microsystem,
Wetzlar, Germany).

1.3. Genome sequencing and phylogenetic reconstruction

Sequencing

We acquired high-quality genome sequences using a hybrid approach that combined long-read data from Oxford Nanopore Technology (ONT, Oxford, United-Kingdom) with high-accuracy short reads from Illumina (San Diego, CA, USA). Genomic DNA was extracted using the QIAGEN Blood and tissue DNA extraction kit (Hilden, Germany) following manufacturer's instructions on overnight cultures in LB + NaCl 0.5M at 28 °C.

Illumina libraries were prepared using the Nextera XT Flex DNA prep kit, and ONT library with the Native barcoding kit V14 SQK-NBD114, both according to the manufacturer's instructions (supplementary material).

Read processing, assembly and annotation

The quality of raw Illumina reads was controlled using FastQC [38] (Phred score threshold= 30) and adapters and regions with low quality-scores were cleaned with Cutadapt (version 4.0) [39]. ONT reads with quality score lower than 15 were discarded using NanoPack2 [40], [41].

Hybrid *de novo* assembly was done using Hybracter (version 0.11.2) [42] on the filtered reads. Genomes were annotated using PROKKA (version 1.14.6) [43].

Phylogenetic reconstruction

All the available *V. anguillarum* genomes were retrieved from the curated collection NCBI RefSeq. Subsequently, we assessed the genome integrity using CheckM (version 1.1.3), focusing on completeness and contamination [44]. The analysis was conducted in December 2024 and yielded 140 genomes in total.

Roary (version 3.13.0) [45] was used to determine the core genome of the 157 strains (140 genomes from RefSeq, our 16 sequenced strains, and the *V. ordalii* FS-238 strain as an outgroup). A phylogeny based on single nucleotide polymorphisms (SNPs) in the core genome was built. All core genes were aligned using MAFFT (version 7.505) with the 'auto' option [46]. A core gene nucleotide alignment supermatrix was produced by concatenating all aligned core genes. A SNP distance matrix was obtained from the alignment with SNP-dists (version 0.7.0) [47]. A maximum-likelihood phylogenetic tree was constructed with IQ-Tree 2.1.4-beta [48], and a GTR+F+R9 model of substitution chosen according to Bayesian information criterion obtained with ModelFinder [49].

2. Microplastic and non-plastic control materials

Attachment experiments were conducted with irregularly shaped polypropylene (PP) and polyethylene terephthalate (PET) fragments (30-50 µm) produced by milling (supplementary material). The milling procedure described by [50] was modified to obtain smaller PP microplastics. A different previously published method [51] was optimized for the production of PET microplastics. MPs passing through a 50 µm sieve and retained by a 38 µm sieve were collected for subsequent analyses. The particle size distributions of the PP and PET fractions were measured *via* laser diffraction, confirming that the majority of particles fell within the range 30-50 µm (Figure S1). Monodisperse polystyrene (PS) spheres a diameter of 30 µm (Sigma-Aldrich, MO, USA) were utilized as comparator plastic material. The surface roughness of plastic particles was assessed using electron microscopy. Scanning electron microscopy (SEM) micrographs were recorded using a JEOL-6010LV SEM. Before imaging, the samples were coated with a palladium layer using a JEOL JFC-1300 autofine coater under Ar plasma. PS particles exhibited smooth surfaces, whereas PET and PP particles showed rough surfaces with irregular cavities, which were absent in PS beads (Figure 2A).

Glass spheres with a diameter > 108 µm and lignin fragments were used as non-plastic controls (Sigma-Aldrich, MO, USA).

3. Experiment 1: *V. anguillarum* attachment on pristine microplastics

3. 1. Preparation of bacteria and microplastic mixtures

V. anguillarum strains carrying the pMRB-GFP plasmid were grown overnight in LB + NaCl 0.5M supplemented with chloramphenicol (10 µg/mL) at 28 °C, with constant agitation at 200 rpm. Concentrations of the bacteria cultures were determined by flow cytometry absolute counting (supplementary material) and mixtures at a 1:10 ratio of microplastics to bacteria were prepared using microplastic suspensions at an approximate concentration of $5*10^5$ particles per mL. A volume of 1.8 mL of the resulting mixtures were incubated at 30 °C for 24 hours with constant agitation at 200 rpm in 2 mL Eppendorf tubes. Density differences of the polymeric materials

compared to the used media caused either flotation (PP) or sedimentation (PET, PS) of the
particles. To account for this, the tubes were placed horizontally to maximize the likelihood of
interaction between planktonic cells and particles. Each combination of the 16 strains and three
plastic polymers (PET, PS and PP) was tested in triplicate.

3. 2. Fluorescence microscopy and imaging

After 24h of incubation, 10 µl of each mixture were dropped onto a Malassez cell, which
was placed under the fluorescence microscope Leica DM2000 LED after a sedimentation period
of 3 minutes. Multiple pictures were captured of 10 fields forming a row or a column of squares
(each square representing a volume of 0.01 µl) using a Flexacam C3 (Leica microsystem,
Wetzlar, Germany) connected to the microscope. The specific row or column was selected to
ensure that between 40% and 60% of the area was covered by MPs, where feasible, and before
any fluorescence detection. A brightfield (BF) picture of the MPs and the grid of the Malassez
cell were taken, before a green fluorescence filter was employed to detect the GFP signal. Ten
pictures (GF pictures) of each field were captured under fluorescence excitation light at intervals
of 20 µm in height, thus covering the total distance between the Malassez slide and the
coverslip. This series of pictures allowed for comprehensive assessment of all planktonic
bacteria and bacteria attached to the MPs within a volume of 0.1 µl. All the pictures were taken
at a magnification of x200 and saved as TIFF.
205

3. 3. Image treatment

To achieve high-quality and reproducible image treatment from multiple raw microscopy captures, we developed a pipeline using Fiji (ImageJ version 2.14.0/1.54i) [52]. An overview of the pipeline is depicted in Figure S2. Parameter specifications can be found in Table S2, and green fluorescent images of bacteria attached to microplastics are shown in Figure S3A. Briefly, for each sample, we generated one brightfield megapicture encompassing the 10 fields, and 10 green fluorescent megapictures (one per depth). From that, we segmented them to get a mask for both microparticles and green fluorescent pixels emitted by fluorescent bacteria.

Raw image processing and stitching

Raw BF images were converted to an 8-bit grayscale. For PET and PS microparticle images, a bandpass filter was applied to isolate structures in the microparticle size range. For PP images, a variance filter was applied to highlight regions with local variability, and a Gaussian blur was applied to smooth the images. The transformed BF images of the 10 fields were stitched *via* the MIST plugin [53], [54], with pre-stitching for optimal alignment (Table S2, supplementary material). Cropping was performed to center images, normalize pixel count, and remove edge artifacts.

Pixel segmentation

Segmentation of MPs employed trained models *via* Fiji's trainable Weka segmentation plugin [55], [56], a machine learning-based image analysis. The segmentation resulted in binary pictures capturing the position of the MPs on the stitched grayscale transformed BF picture. Each mask produced by its utilization was visually verified by overlaying it onto the original image. Manual corrections were made in cases of local mis-segmentation (Table S2, supplementary material). The GF pictures were split into the three color channels, and only the green channel was retained for further analyses. Fluorescent bacteria were segmented using different methods depending on the polymer type, including machine learning or differential intensity threshold approaches to account for particle autofluorescence (Table S2, supplementary material).

 Fluorescent pixel classification

Fluorescent pixels were categorized into different groups based on their overlap with
microplastic masks. Fluorescent pixels overlapping those of the MPs and present at the same
depth (from 200 to 140 µm for PS beads; from 200 to 120 µm for PET) were classified as
originating from bacteria attached to the microplastics. For the PP particles, we observed two
behaviors in the water column. In some samples, the particles floated at a depth between 120
and 40 µm, while in others, the particles settled at the bottom of the column, between 200 and
120 µm in depth. This difference in behavior between floating and sinking PP was considered
during pixel classification. Fluorescent pixels overlapping the microplastics but at shallower
depths were treated as noise, as we could not distinguish their origin between a clump of
bacteria attached below or a planktonic bacterium at a shallower depth. Fluorescent pixels not
overlapping with the MPs were classified as originating from planktonic bacteria, regardless of
depth.

Strain attachment to supports was evaluated according to Equation 1, where $n_{overlap}$ was
the number of fluorescent pixels overlapping MPs and n_{mask} was the number of pixels in the MP
mask).

$$Attachment\ (\%) = \frac{100\ \times n_{overlap}}{n_{mask}} \qquad\qquad \text{(Equ. 1)}$$

3.4 Exploration of genomic features involved in microplastic attachment

We formulated several hypotheses regarding the potential association of the intraspecific
variability in attachment to microplastics with genomic features: attachment efficiency is
associated with either (1) intra-species phylogenetic or gene content distances, (2) virulence
gene content, or (3) the capacity of bacteria to use plastic polymers as a source of carbon.

First, we performed a phylogenetic reconstruction on the 16 tested strains only, using
the same methods as for the complete collection of *V. anguillarum* genomes (see above
'phylogenetic analyses') and we computed the phylogenetic variance–covariance matrix
obtained from the phylogenetic tree using the vcv.phylo function of the 'ape' package [57]. This
matrix was later introduced as a random effect in a mixed effect model to account for a
phylogenetic influence on attachment to microplastics (see below 'statistical analyses' section).
Roary was also used to obtain the accessory gene content of each strain. This was later used to

test the association between the attachment rate and the presence/absence of genes in the accessory genome (see below 'statistical analyses' section).

Second, to determine whether attachment efficiency was associated with virulome content, we constructed a database compiling all experimentally validated genes involved in the virulence of *V. anguillarum*. Briefly, we looked for '*Vibrio anguillarum*' in Pubmed, and read title and abstract of all the articles published between January 1965 and January 2025 to identify relevant studies. Accession numbers corresponding to virulence genes were retrieved from publicly available genomic repositories NCBI [58] and DDBJ [59]. Gene sequences were curated and organized according to their virulence system into a structured database of 308 sequences for further comparative analyses. The dataset used in this study is publicly available at Zenodo (DOI: 10.5281/zenodo.18151786). Then, the 16 *V. anguillarum* genome scaffolds were screened for virulence genes using Abricate (version 1.0.1, minimum percent sequence identity: 80%, minimum percent coverage: 90%) [60] against our custom database, as well as the available virulence database 'vfdb' (last update: January 2025) [61], [62].

Third, scaffolds were screened for hydrocarbon degradation protein sequences using Proteinortho (version 6.3.4) [63] against the HADEG database (version 'HADEG_protein_database_231119.faa') [64], to assess whether certain strains could use plastic polymers as a source of carbon.

4. Experiment 2: *V. anguillarum* attachment on field-colonized microplastics

4.1. On-field experimental set-up

Incubation sites

Three localities were chosen along the French Western Mediterranean coast: a semi-enclosed lagoon (bassin de Thau, GPS coordinates of the experimental set-up: 43.37897°N, 3.57164°E), a marina (Carnon's harbor, GPS coordinates: 43.54609°N, 3.97173°E), and a protected area (the Marine Reserve of Cerbères-Banyuls, GPS coordinates: 42°29300 N – 03°08700 E). Each location was selected due to differences in the type and intensity of human activities: the Bassin de Thau is known for its oyster farming and recreational nautical activities, Carnon's harbor is a marina. Both are highly frequented during summer, whereas access to the marine reserve is restricted all year long.

 Field-colonized microplastic incubation

We prepared the sampling device as described in [65]. For each incubation, four nylon bags
containing 50 mg of PP, PS, PET or glass microparticles were hung with Ty-raps within an
oyster cage (Nodus factory, France). The cage was loaded with four 1 kg diving weights and
immersed one meter below the surface. The incubation period was four weeks.

At the end of the incubation period, the bags containing MPs were processed in the lab as
described in [65]. Briefly, the bags were cut open and the content was resuspended in a sterile
artificial seawater (Sea salt, InstantOcean, VA, USA) by aspersion. The obtained suspensions
were transferred onto a 20 μm mesh filter to collect MPs and discard the planktonic material.
The field-colonized MPs were resuspended in 2 mL of artificial seawater, of which 400 μl were
dedicated to *V. anguillarum* attachment rate measurement. Brightfield images of natural biofilms
formed around microplastics are shown in Figure S3B.

4.2. Incubation of field-colonized particles with fluorescently labelled *V. anguillarum* strains

V. anguillarum strain selection

For attachment experiments with microplastics colonized by marine biofilms, strains NB10,
VIB8, and VIB205 were chosen based on (1) their genetic distance, each strain being distinct
from each other by more than 30,000 SNPs in their core genome (average SNP number among
pairs of the 16 strains = 30,443), and (2) their Jaccard distance calculated on their accessory
gene content, which was above 0.30 for each pair (average Jaccard distance among pairs of
the 16 strains= 0.26).

In vitro assay and image processing

The number of resuspended microplastics was estimated for each sample by counting the
particles under a microscope using a Malassez counting chamber. The incubation protocol and
image capturing scheme were similar to the ones described for the first experiment. Briefly, 100
μl of the field-colonized MP suspension was mixed with the required volume of bacteria
suspension to achieve a ratio 1:10. The mixture was incubated for 24 hours at 30 °C, with
constant agitation at 200 rpm. Each combination of the three *V. anguillarum* strains, three
locations, three plastic polymers, and the non-plastic control (PET, PS, PP, and glass) was

tested in triplicate. For each combination, we also incubated virgin microparticles in parallel as a
paired control under the same conditions.

The same image processing procedure as for the first experiment was employed using our
custom Fiji pipelines for PET, PP and PS microparticles. For glass beads, a modified version of
the PS microparticle pipeline was developed (Table S2, supplementary material). The outcome
variable of this experiment was the difference in the proportion of fluorescent pixels overlapping
with particles between field-colonized particles and their paired control.

5.   Experiment 3: Effect of microplastics on *V. anguillarum* survival to bleach treatment

We tested the effect of biofilm on the survival of *V. anguillarum* to bleach treatment. We
used PP and PET microparticles, fragments of lignin (Sigma-Aldrich, MO, USA) as non-plastic
control, and VIB205 as focal strain. MPs were mixed with the required volume of bacteria
suspension to achieve a ratio 1:10. The mixture was incubated in pre-filtered (0.45 µm) and
autoclaved seawater, which was sampled from Carnon's harbor (GPS coordinates:
43.546231°N, 3.971839°E). Each combination of the focal strain and material (PET, PP, and
lignin) was tested in triplicate. We also incubated the focal strain in absence of MPs under the
same conditions as a 'non-biofilm support' control.

Incubation was conducted in a total volume of 100 mL at 28 °C with constant agitation at
150 rpm to enable *V. anguillarum* cells to attach to MPs. After 5 days of incubation, 2.7 µL of
bleach at 37 g L^{-1} were added to each sample (final concentration of 1 mg L^{-1}). After 30 minutes
of exposure to bleach, each sample was poured on PluriStrainer Mini cell strainer (PluriSelect,
Germany) with a mesh size of 20 µm to retain colonized MPs. Biofilms were separated from
MPs as described in [66]. According to the track plate method, 10 µL of undiluted resuspension
and dilutions from 10^{-1} to 10^{-3} were spotted on TCBS plates and incubated at 28 °C. Survival to
bleach treatment was evaluated based on the number of colonies formed after 48 hours.

6.   Statistical analyses

For the first experiment, attachment was evaluated based on the proportion of fluorescent pixels covering the MP surface. To analyze the effect of polymer type, a generalized linear mixed-effect model (GLMM) was employed. The Beta distribution was used to model the proportion data. Along with polymer type, we also assessed the effect of *V. anguillarum* strains and their interaction. The day of the experiment was included as a random effect to account for variability between replicates.

Model parameters were estimated in a Bayesian framework, using a Markov Chain Monte Carlo (MCMC) algorithm, implemented with the R package brms [67], [68], [69] (supplementary material). The models were compared using the Watanabe-Akaike Information Criterion (WAIC). Among the models with the smallest WAIC values and a difference of less than 4 units, we selected the most parsimonious model. Once the best model was selected, we included the phylogenetic variance–covariance matrix as a 'random effect'. We tested whether the proportion of variance attributed to the phylogenetic structure significantly contributed to the total variance in the model, and in which proportion.

Next, we tested the association between the attachment rate and the presence/absence of genes in the accessory genome of the 16 strains using Scoary2 (version 0.0.15, Benjamini-Hochberg correction of p-values with a cut-off of 0.05) [70]. To enhance statistical power, we selected candidate genes present in at least 4 strains and no more than 12 strains.

For the second experiment, the difference in the proportions of fluorescent pixels overlapping with particles between field-colonized particles and their paired control was used. Model parameters were estimated in the same Bayesian framework, as described above. The effects of polymer type, location of particle field-colonization, and *V. anguillarum* strains were analyzed using a generalized linear model. The Beta distribution was used to model the data after applying an affine transformation, which mapped the data from the interval [-1, 1] to [0, 1]. The day of the experiment was included as a random effect to account for variability between replicates.

All statistical analyses were conducted using R version 4.4.0 [71].

Results

1. Phylogenetic analyses

We gathered a total of 157 WGS, consisting of 156 *V. anguillarum* strains and one *V. ordalii* strain, which was placed as an outgroup of the phylogenetic tree (Figure 1B). Among these, the genomes (composed of two chromosomes in *V. anguillarum* species) of the sixteen tested strains for MP attachment were assembled *de novo* using hybrid Illumina–ONT sequencing. The average genome size was 4,189,500 bp, with an average GC content of 44.5%. The assemblies consisted of 2 to 10 contigs each, with fourteen genomes resolved into complete circular chromosomes. The hybrid assemblies showed high coverage depth (average >500X), supporting accurate reconstruction of chromosomal structures. The core genome comprised 2,135 genes, with a total of 14,364 genes detected across all genomes, resulting in an average of 3,739 genes per strain. A phylogeny based on single nucleotide polymorphisms (SNPs) in the core genome was built using the *V. anguillarum* genome assemblies from RefSeq, the newly sequenced genomes, and the *V. ordalii* FS-238 strain. The tested strains were distributed throughout the phylogenetic tree, with three clusters of closely related strains observed (< 1,000 SNP): NB10, VIB1, VIB271, and VIB273; VIB197, VIB200, and VIB220; VIB205, and VIB206 (Figure 1B). The average number of SNPs among pairs of tested strains was 30,443.

2. Experiment 1: *V. anguillarum* attachment on pristine microplastics

Attachment depends on plastic polymer and V. anguillarum *strain*

Technical issues resulted in missing brightfield and green fluorescence raw images for 10 out of 144 samples. Nevertheless, all affected samples, including those with PS (2 samples), PP (5 samples), and PET (3 samples), were retained in the final analysis (supplementary results, Table S4). After cropping, we compared the size of the reconstructed megapictures, expressed as a number of pixels, to ensure comparability between polymers. No significant difference was detected between polymers (anova, p-value= 0.06, Figure S4).

The mean proportion of fluorescent pixels covering the microplastic surface for PET, PP
and PS was 72.6% (± 19.6%), 50.0% (± 21.0%), and 15.7% (± 7.2%), respectively, with a wide
distribution for PET and PP (ranging from 35.3% to 99.6% and 10.9% to 93.8%, respectively,
Figure 2). For PS, this low proportion combined with very few fluorescent pixels in absolute
number was consistent with the almost complete absence of fluorescent bacteria attached to
MPs observed by visual inspection under the microscope. Within each polymer, a continuum of
attachment was observed among the strains, rather than distinct groups (Figure 2).

Using a generalized linear mixed-effects model, we assessed whether the proportion of
fluorescent pixels covering the microplastic surface differed between plastic polymers. The
model including polymer type and *V. anguillarum* strain as explanatory variables, with no
interaction terms, had the lowest WAIC (Table 1) and was thus considered the best model.
Based on the Bayesian model comparison, we found substantial evidence supporting a strong
effect of polymers and strains (Table 2, Bayes Factor$_{model\ null-\ model\ 1}$= 0.0006 and BF$_{model\ 1-\ model\ 2}$
$<10^{-5}$).

The absence of significant interaction between strain and polymer in the model retained
suggests that strains exhibited similar behavior in their attachment across the three polymers.
Despite an important inter-replicate variability, strains with a high attachment phenotype on PET
also exhibited a high attachment phenotype on PP. These similarities in ranking between PET
and PP were not observed in the presence of PS, where very few attached bacteria were
observed.

Exploration of genomic features involved in microplastic attachment

To test the hypothesis that differences in attachment efficiency was explained by
phylogenetic distances, the variance-covariance matrix derived from the phylogenetic tree of the
core genome was introduced as a random effect in the mixed effect model. The proportion of
variance attributable to phylogenetic relatedness was 0.03 [0.00; 0.09]$_{CI95\%}$, indicating no
phylogenetic signal. No significant association was found between any gene of the accessory
genome and the attachment gradient to PET or PP.

Among the 307 Vibrio anguillarum-specific gene sequences encompassing 23 virulence
systems gathered in our custom database, we detected between 220 and 277 virulence
genes (median: 230), with the majority localized on the first chromosome (median number on
the first and second chromosome being 167 and 64, respectively). The number of virulence
genes was not correlated with their attachment efficiency (rho= -0.13, p= 0.6, Spearman's

correlation test). There was no correlation either when focusing on genes involved in adhesion
and quorum sensing (rho= -0.07, p= 0.8, Spearman's correlation test).

To assess whether the attachment rates of *V. anguillarum* strains to MPs could be linked
to the presence of genes encoding plastic degrading enzymes, the annotated genomes were
screened for the presence of genes of the HADEG database. The outer membrane lipoprotein
Blc (UniProt accession number Q08790) and the protein-tyrosine-phosphatase Wzb (UniProt
accession number P0AAB2) were the only sequences detected in 15 and 10 strains,
respectively. No association was found for the attachment gradient, since strains without these
sequences did not exhibit an outlier phenotype.

3. Experiment 2: *V. anguillarum* attachment on field-colonized microplastics

Technical issues resulted in missing brightfield and green fluorescence images for 7
samples out of 216 samples. Nevertheless, all affected samples, including those with PS (2
samples), PET (1 sample), PP (2 samples), and GL (2 samples), were retained in the final
analysis (supplementary material). After cropping, there was no significant difference between
the size of pictures between naturally-colonized particles and their control, nor between strains
and locations. The surface of BF pictures covered by PS beads was significantly lower than for
the other substrates (posthoc Tukey test, p-value < 10^{-5} for the three comparisons, Figure S5),
as for Thau compared to Carnon incubation sites (posthoc Tukey test, p-value= 0.001).

As a preliminary step, we confirmed that strain attachment to pristine MP particles was
consistent with the results of the first experiment (Figure S6). Attachment was highest on PET
(81% ± 15.7% on average), intermediate on PP (73.4% ± 20.1%), and lowest on PS (15.9% ±
12.1%). For each of the three strains, the proportion of fluorescent pixels overlapping particles
in experiment 2 was in line with that observed in experiment 1, with the largest difference in
mean observed for VIB205 on PET (70.7% ± 27.5% on average in experiment 1, 90.6% in
experiment 2 ± 8.4%).

For each combination of strain, field location and polymer type, we computed the
difference in fluorescent pixel overlap between field-colonized and pristine particles (Figure S7).
Positive differences reflected increased attachment of GFP-labelled *V. anguillarum* to
microplastics in the presence of naturally formed biofilm, while negative differences reflected
decreased attachment.

The difference in fluorescent pixel overlap was negative for PET and PP (PET= -22.3 ±
23.0, PP= -26.5 ± 28.3, Figure 3, Table S5), while positive for PS beads (22.0 ± 27.5). The
difference in proportion of fluorescent pixels covering the microplastic surface between naturally
colonized microparticles and their paired control was similar among incubation sites (-4.8 ±
22.3, -5.1 ± 36.3, and -7.4 ± 33.9 for Banyuls, Carnon and Thau, respectively, Table S5) and
among tested strains (-12.5 ± 31.1, -5.8 ± 33.8, and 0.9 ± 27.7 for NB10, VIB8, and VIB205,
respectively, Table S5).

The model with polymer and *V. anguillarum* strain variables and with an interaction term
exhibited the lowest WAIC, while the simpler model that included only the polymer variable
differed by less than 4 points (Table 3). Given the principle of parsimony, we selected the model
including only the polymer type for further analyses.

In the presence of naturally formed biofilms, *V. anguillarum* showed significantly reduced
attachment to PP and PET, whereas attachment was significantly enhanced on PS, compared
with the non-plastic support (Table 4, -0.53 [-0.82; -0.26]$_{CI95\%}$, -0.62 [-0.89; -0.37]$_{CI95\%}$, 0.40
[0.11; 0.68]$_{CI95\%}$, for PET, PP and PS, respectively).

4.   Experiment 3: Effect of microplastics on *V.*

anguillarum survival after bleach treatment

We tested the effect of attachment to microplastics on the survival of *V. anguillarum* after
30 minutes of bleach exposure at a concentration of 1 mg L^{-1}. As expected, the no-support
control conditions showed no *V. anguillarum* growth after bleach treatment, whereas viable
colonies formed on TCBS following incubation with PET and lignin in two out of three replicates
(Table 5), suggesting that attachment to PET or lignin reduces the efficacy of bleach treatment.
Colony counts were higher in lignin samples than in PET: lignin replicates yielded 5.3*10□ and
1.5*10^{4} CFU/ml, compared with 900 and 100 CFU/ml in PET replicates. No colony was found
from PP.

Discussion

502
The number of scientific studies that show plastics can be major negative disruptors of
marine ecosystems keep increasing [15], [72], [73], [74], [75]. One aspect is that plastic debris,

and in particular MPs, can be vectors or vehicles for pathogenic bacteria species in marine
environments [16]. Here, we explored some of the driving factors for attachment of the fish-
pathogen species *Vibrio anguillarum* to MPs. We measured the *V. anguillarum* attachment on
PS, PP, and PET microparticles in lab-controlled conditions. This experimental approach was
repeated with 16 strains, isolated in both an infection or non-infection context, representing the
within-species diversity. We further investigated if the *V. anguillarum* attachment was influenced
by a field-acquired biofilm on the MP surface, and whether the effect of attachment to an MP
support influences strain survival when exposed to bleach.

Our experimental approach revealed a strong effect of the particle type on the
attachment rate of the 16 *V. anguillarum* strains tested (Figure 2B). All three particle types, *i.e.*
milled PP MPs, milled PET MPs and commercially manufactured PS beads, were of similar
median size (30 µm) but differed in their shape, size range, chemical composition and surface
characteristics, e.g. roughness. Furthermore, while the in-house produced PP and PET particles
contain no additives, the PS beads might also contain undisclosed additives (PS-beads
commercially available but no list of additives was provided). Physically, PS beads have a very
smooth surface whereas PP and PET particles obtained by milling have a rough surface with
numerous crevices (Figure 2A). The higher attachment rates on PP and PET particles suggest
that the physical differences (*i.e.* surface roughness) may explain the "particle type" effect,
although a fully crossed design would be necessary to disentangle the effects of roughness and
chemical composition on the attachment rate. The positive effect of rugosity on attachment rates
has been found in other microplastic and non-microplastic systems. For example, Foulon and
collaborators showed that *Vibrio crassostreae* can remain attached for several days on PS
microparticles, with longer attachment observed on particles exhibiting greater rugosity [76].
Additionally, Hossain and collaborators showed in a freshwater microcosm that colonization of
microplastics by three non-Vibrio species was also correlated to rugosity [77]. The studies
analyzing the link between roughness and bacteria attachment rates have been reviewed [78]
and there is either a positive relationship between roughness and attachment rate or no
relationship. These results outline the importance of the choice of the MP models when
designing experiments on bacteria attachment rate to evaluate the risk of pathogen transport by
MPs. The use of "realistic" MPs (*i.e.* with similar shape and chemical composition to
environmentally aged materials) is key and in this regard, the MPs obtained by milling are more
similar to MPs produced by plastic object degradation than PS beads.

We next sought to establish whether there were differences in attachment rates between
strains. The strains used in the present study were chosen to be diverse in terms of sampling
sites and dates. Furthermore, the obtained whole genome sequences confirmed that they are
also spread within the phylogenetic tree of the species, and that they differ greatly in their
accessory gene content. Our experimental approach revealed a strong intraspecific variation,
with strain-specific attachment rates ranging from 35 to 100% on PET and from 10 to 94% on
PP (Figure 2). The intraspecific variability in attachment rates has rarely been tested but
measurements performed on other *Vibrio* species [79] revealed important differences between
strains involved in biofilm formation on plastic surfaces. The strong intra-specific variability in
attachment rates found here highlights that determining the attachment rate of a species is not
straightforward and cannot be based on measuring a single "representative" strain. It further
means that assessing the risk for a particular MP type as a vehicle of a particular pathogenic
species is, in itself, a complex task.

The strong inter-strain variability and the similarity in attachment rates for PET and PP
particles suggests that there are genetic determinants of the attachment rate. We tried to
identify them by testing associations between the attachment rate measurements and the
genomic information contained in the whole genome sequences. However, no significant
phylogenetic signal was found and no genomic determinant could be identified. Despite
combining a general approach with a more targeted screening of *V. anguillarum* virulence genes
from an ad-hoc database (Table S3, 10.5281/zenodo.17525113), we found no evidence that the
presence of specific genes, including virulence-associated ones, could account for the
differences observed among strains. Our decision to focus on virulence genes was based on
the fact that some life-history traits involved in interactions with the environment, particularly
adhesion to surfaces, have been shown to depend on virulence factors [80]. This led us to
hypothesize that strains with certain virulence genes could have a higher attachment rate, which
could, in turn, increase the frequency of virulent or pathogenic strains on MPs. The lack of
significant association between genetic composition and attachment rate might be due to
insufficient statistical power – given the number of strains used – limited by the tractability of the
experimental approach, and image analysis. Another explanation could be non-genetic
differences between strains (and even between replicates) in the level of expression of genes
involved in quorum sensing, motility or exopolysaccharide synthesis [81], as reported by Foulon
et al. [76] in a laboratory experiment or Kesy *et al.* [82] in field conditions.

We found that one-month field-colonization in a marine environment altered the

attachment rates of three tested *V. anguillarum* strains, such that the effect of the particle type

described on pristine particles was mitigated. Attachment rates on PP and PET particles were

lower for field-colonized particles than for pristine particles, whereas attachment rates on PS

particles were higher for field-colonized particles than for pristine particles (Figure 3). This

suggests that the presence of a field-formed biofilm homogenizes surface properties, by

masking physical and chemical surface properties. Such patterns could also or additionally

evolve due to ecological interactions including both attraction and repulsion cues from the

microorganisms of the field-formed biofilms [83]. The dynamics of the plastisphere assembly

may largely depend on a combination of stochastic encounters between microplastics and

marine bacteria, differences in composition of the ecocorona between polymers [84], differential

opportunistic colonization strategies mediated by environmental conditions [79], as well as

differential capacities to adhere to surfaces of varying rugosity. The abundance of early

colonizers could therefore be determined by the order of attachment resulting from the sum of

these forces, whereby species that settle first may gain a competitive advantage or facilitate the

subsequent attraction of other species. Such variability could explain long-term changes that are

often observed in plastisphere composition dynamics [85], [86].

Biofilm formation is known to increase the phenotypic resistance of bacteria to biocides

such as antimicrobials [87] and disinfectants [88]. Since MPs provide surfaces for biofilm

development, their presence in aquatic environments may reduce the effectiveness of

sanitization measures. Among these, we selected bleach treatment as a widely employed

strategy for sewage disinfection and water potabilization. We examined whether MP pollution

could reduce its efficacy by facilitating the persistence of pathogenic bacteria. We found that

PET microparticles enabled survival of *V. anguillarum* during 10 minutes bleach treatment, as

did lignin particles. Further validation of these results will require repeating the experiment with

a larger number of replicates, as well as using older and more naturally developed biofilms.

Despite a large number of zero values resulting from bacterial mortality caused by the

treatment, they stress the need to explore the potential role of MPs as shelter during bleach

treatment. To our knowledge, this is the first study that aims to quantify the additional risk of

MPs over bacterial survival to bleach treatment.

In conclusion, our experimental approach revealed that the adherence of *V. anguillarum*

to MPs, and as a consequence the risk for MPs being a vector of pathogenic *V. anguillarum*

strains depends on many factors, including physico-chemical particle surface properties, the

presence of other microorganisms, and also the identity of the strain. This combination of

factors, and of potential others not studied here, makes the evaluation of the risk a very complex

task.

Concerns have also been raised regarding the enrichment of antibiotic-resistant bacteria

in biofilms associated to microplastics compared to free-living bacteria [89]. However, studies on

non-plastic suspended matter in freshwaters have likewise reported an enrichment of mobile

genetic elements linked to antibiotic resistance [90]. This further underscores the importance of

elucidating the colonization dynamics of pathogenic species on these surfaces. Pathogenic

species that ride plastic litter may disrupt fragile ecosystems, already pressured by overfishing

and other types of pollutions. While it is relatively rare for a non-native species to successfully

survive in a new environment, rafting on ubiquitous plastic debris could potentially increase the

frequency of such events.

Acknowledgments

We thank Frédérique Le Roux, Delphine Destoumieux-Garzon, Guillaume Charrière,

and François Delavat for providing conjugation protocol and plasmids.

We are grateful to the ImpTox project's members, in particular Andreja Rajkovic and

Elsa Gadoin for useful discussions. We also thank Tim Janicke for useful discussions,

and Vahiniaina Andriamanga and Marie-Pierre Dubois for their assistance in ONT sequencing.

We also thank MARBEC unit (Univ Montpellier, CNRS, IFREMER, IRD, Montpellier,

France) and Ifremer institute (Unité observation et écologie de la restauration des écosystèmes

littoraux, Laboratoire Environnement Ressources Occitanie, Sète, France) for giving us

authorization to install the incubation device on Ifremer's experimental oyster farm in Bassin de

Thau, in particular Delphine Serais, Hervé Violette. We thank the scientific council of the marine

reserve of Cerbères-Banyuls (Réserve Naturelle Marine de Cerbère-Banyuls, Département des

Pyrénées-Orientales, France), for providing authorizations to incubate the device on site, in

particular Virginie Hartmann. We also thank the Banyuls Observation Sea Service (BOSS

FR3724, Observatoire Océanologique de Banyuls, Banyuls-sur-mer, France), in particular

Laurent Zudaire. We thank Morgann Michel and the harbor's office for providing authorizations

to incubate the device in Carnon's harbor as well.

All authors were supported by Horizon 2020 grant (grant number: 965173). We report no

conflict of interest. IA tools were used to help improve English. Genome assemblies have been

deposited in the European Nucleotide Archive (ENA) under BioSample accession numbers

ERS28308090 to ERS28308105, with corresponding assembly accession numbers ERZ28769549 to ERZ28769564. All scripts used for the analyses have been deposited on Codeberg and are openly accessible at the following link: https://codeberg.org/merimas/ImpTox_project_Vibrio_anguillarum_attachment_to_microplastics

The virulence gene database and the fasta file used as reference database are openly available at Zenodo (DOI: 10.5281/zenodo.17525113, DOI: 10.5281/zenodo.18151786).

Declaration of generative AI and AI-assisted technologies in the manuscript preparation process

During the preparation of this work the authors used ChatGPT in order to improve language. After using this tool, the authors reviewed and edited the content as needed and take full responsibility for the content of the published article.

CRediT author statement

Méril Massot: Data curation, Formal analyses, Investigation, Software, Validation, Writing- Original draft preparation

Lukas Wimmer: Data curation, Investigation, Resources, Validation, Visualization, Writing – review & editing

Mhamad Aly Moussawi: Resources, Visualization, Writing – review & editing

Jeanne Hamet: Data curation, Resources, Validation, Writing – review & editing

Tatjana N. Parac-Vogt: Funding acquisition, Methodology, Project administration, Supervision, Writing – review & editing

Lea-Ann Dailey: Funding acquisition, Methodology, Project administration, Supervision, Writing – review & editing

Martijn Callens: Conceptualization, Funding acquisition, Methodology, Resources, Writing – review & editing

Stéphanie Bedhomme: Conceptualization, Funding acquisition, Methodology, Project administration, Resources, Supervision, Writing – original draft, Writing – review & editing,

664 **Table 1.** Model selection for the first experiment. Model in bold correspond to the model
665 selected.
666

Model	Covariables	WAIC
null	null	-133.7
1	Plastic polymer	-133.3
2	**Plastic polymer** **Strains**	**-162.1**
3	Plastic polymer Strains Interaction(plastic polymer; strain)	-137.1

667
668
669 **Table 2.** Median and 95% credible intervals of posterior distributions for the estimates of the
670 proportion of fluorescent pixels covering microplastics. Levels used as reference were "PS" and
671 "VIB1" for polymer and *V. anguillarum* strains, respectively. Estimates in bold correspond to
672 distributions for which the confidence interval does not contain zero.

	Estimate	95% CI
FIXED EFFECTS		
Intercept	**-2.81**	**[-3.59 ; -2.07]**
Polymer		
PET	**2.75**	**[1.96 ; 3.51]**
PP	**1.69**	**[0.92 ; 2.46]**
Strains		
VIB10	**2.07**	**[1.36 ; 2.76]**
VIB8	**1.86**	**[1.16 ; 2.59]**
VIB273	**1.55**	**[0.88 ; 2.24]**
VIB72	**1.50**	**[0.78 ; 2.25]**
VIB5	**1.48**	**[0.83 ; 2.19]**
NB10	**1.32**	**[0.64 ; 2.10]**
VIB197	**1.23**	**[0.54 ; 1.93]**
VIB205	**1.13**	**[0.51 ; 1.91]**
VIB206	**1.04**	**[0.40 ; 1.78]**
VIB271	**1.02**	**[0.35 ; 1.75]**
VIB220	**1.03**	**[0.30 ; 1.74]**
VIB200	**0.93**	**[0.28 ; 1.66]**
VIB203	**0.88**	**[0.13 ; 1.59]**
VIB7	0.64	[-0.08 ; 1.43]
VIB71	0.61	[-0.12 ; 1.33]
RANDOM EFFECTS		
'day of experiment' variance	0.39	[0.14 ; 0.82]
residual variance	9.61	[7.37 ; 12.10]

673
674

675 **Table 3.** WAIC of models run for the second experiment. Model in bold correspond to the model
676 selected.
677

Model	Covariables	No interaction	Interaction
null	null	-87.5	-
1	**Plastic polymer**	**-136.4**	-
2	Plastic polymer Strain	-138.3	-135.5
2	Plastic polymer Location	-132.4	-140.0
3	Plastic polymer Strains Interaction(plastic polymer; strain)	-135.5	-118.9

678
679
680
681 **Table 4.** Median and 95% credible intervals of posterior distributions for the estimates of the
682 difference of proportions of fluorescent pixels covering microplastics between field-colonized
683 particles and pristine. Level used as reference was "GL". Estimates in bold correspond to
684 distributions for which the confidence interval does not contain zero.
685

	Estimate	95% CI
FIXED EFFECTS		
Intercept	0.06	[-0.40 ; 0.44]
Polymer		
PET	**-0.53**	**[-0.82 ; -0.26]**
PP	**-0.62**	**[-0.89 ; -0.37]**
PS	**0.40**	**[0.11 ; 0.68]**
RANDOM EFFECTS		
'day of experiment' variance	0.23	[0.01 ; 0.68]
residual variance	14.40	[10.90 ; 18.52]

698
699
700 **Table 5.** Number of colonies found on TCBS plates after bleach treatment. Numbers in bold
701 were used to estimate the concentration of bacteria after survival.
702

Material	PET			PP			Lignin			No support		
Replicate	R1	R2	R3	R1	R2	R3	R1	R2	R3	R1	R2	R3
No dilution	**0**	**9**	**1**	**0**	**0**	**0**	>100	48	**0**	**0**	**0**	**0**
10^{-1}	0	3	0	0	0	0	**53**	**15**	0	0	0	0
10^{-2}	0	0	0	0	0	0	12	2	0	0	0	0
10^{-3}	0	0	0	0	0	0	0	0	0	0	0	0
CFU/ml estimated	0	900	100	0	0	0	$5.3*10^4$	$1.5*10^4$	0	0	0	0

Figure 1. Phylogenetic tree and geographic origin of the tested *V. anguillarum* strains. Panel A represents the geographical origin of each strain. Colors refer to the identity of each strain on the phylogenetic tree. Countries of origin are colored in gray. Panel B represents the maximum likelihood phylogenetic tree of the 156 *V. anguillarum* strains. The tree was rooted on the *V. ordalii* FS-32 strain. It was built using IQ-Tree 2.1.4-beta, and is based on the 2,135 genes that composed the core genome. For clarity purpose, the tree presents only the 16 strains tested in this analysis.

Figure 2. Scanning Electron Microscopy (SEM) images showing microplastic surface morphology, and proportion of fluorescent pixels covering microplastics for the 16 *V. anguillarum* strains tested. Panel A shows SEM images of PET, PP and PS particle surfaces, shown from left to right, respectively. Scale and magnifications are provided at the bottom of each image. Panel B displays the proportion of fluorescent pixels covering the microplastics. Each graph represents attachment data for microplastics from different polymers (from left to right: PET, PP, and PS). For each strain, gray points show the observed data for three replicates, while colored diamonds indicate the mean proportion. The color code corresponds to the strain, and is consistent across all graphs. Strains are organized in descending order based on their mean values for PET.

Figure 3. Change in proportion of fluorescent pixels overlap between field-colonized microparticles and control, according to the nature of substrate. For each strain, gray points show the difference between paired samples for each combination of {substrate - location - *V. anguillarum* strain}, while red diamonds indicate the mean difference for each substrate. A difference below zero indicates lower colonization of field-colonized substrate compared to the control by the focal *V. anguillarum* strain.

Table S1. Information on the origin of *V. anguillarum* strains used for experiments. Information was retrieved from the Belgian Coordinated Collection of Microorganisms catalogue.

Strain	LMG name	Serotype	Sample origin	Country of origin	Year indicated in BCCM
NB10	LMG 32357	O1	rainbow trout (*Oncorhynchus mykiss*)	Sweden	1990
VIB 1	LMG 10861	O1	rainbow trout (*Oncorhynchus mykiss*)	Denmark	1990
VIB 10	LMG 10870	O10	cod (*Gadus morhua*)	Denmark	1990
VIB 197	LMG 13224	VaNT7	rotifer	Greece	1991
VIB 5	LMG 10865	O5	cod (*Gadus morhua*)	Denmark	1990
VIB 7	LMG 10867	O7	eel (*Anguilla anguilla*)	Denmark	1990
VIB 71	LMG 4411	O2a	Young sea trout (*Salmo trutta*)	Scotland	1960
VIB 72	LMG 4437	O2a	cod (*Gadus morhua*), ulcerous lesion	Norway	1956
VIB 8	LMG 10868	O8	cod (*Gadus morhua*)	Denmark	1990
VIB200	LMG 13227	VaNT7	sea bream (*Sparus aurata*), larvae	Greece	1991
VIB203	LMG 13230	O6	sea bream (*Sparus aurata*), larvae	Spain	1991
VIB205	LMG 13232	O6	water	Spain	1991
VIB206	LMG 13233	O6	rotifer	Spain	1990
VIB220	LMG 13247	unknown	water	Greece	1991
VIB271	LMG 13583	O1	chinook salmon (*Oncorhynchus tschawytscha*)	Oregon, United States	1974
VIB273	LMG 13585	O1	rainbow trout (*Oncorhynchus mykiss*)	Arizona, United States	1968

769 **Table S2. Fiji's operation details.** Each step is presented with specific parameters for each
770 type of microparticle. Different strategies were applied due to heterogeneity in microparticle
771 density and opacity.

Analysis step	PET	PP	PS	GL
1. Raw image processing - BF	1. converted to 8-bit grayscale 2. bandpass filter to filter out structures larger than 100 pixels and smaller than 20 pixels, with a tolerance of 5	1. converted to 8-bit grayscale two copies of the image 2. variance filter with a radius of 1 pixel applied to copy 1 3. Gaussian blur with a sigma value of 5 applied to copy 2 4. Copies 1 & 2 merged	1. converted to 8-bit grayscale 2. bandpass filter to filter out structures larger than 100 pixels and smaller than 20 pixels, with a tolerance of 5	1. converted to 8-bit grayscale 2. bandpass filter to filter out structures larger than 100 pixels and smaller than 20 pixels, with a tolerance of 5
1. Raw image processing - GF	1. split into the three color channels 2. green channel retained			
2. Stitching	1. Each raw image was copied and converted to an 8-bit grayscale 2. Variance filter applied with a radius of 3 pixels to enhance regions with high pixel variability. 3. Threshold applied to refine the segmentation: pixels with intensity values between 10 and 255 considered as foreground, while those below 10 were designated as background. 4. Images converted to binary masks. The background was set to black. 5. MIST plug-in and generation of a text file containing the relative coordinates of images was generated and saved for later assembly 6. Stitching of the 10 images transformed for later segmentation using the MIST plug-in feeded with the text file containing the relative coordinates of images			
3. Cropping	Horizontal megapictures: rectangular region for cropping starting at position {x=0, y= 110}. This rectangle uses the full image width, and adjusted height to 890 pixels Vertical megapictures: rectangular region for cropping starting at position {x=300, y= 0}. This rectangle uses the full image height, and adjusted width to 1,200 pixels			
4. Segmentation - BF	1. Trainable Weka segmentation classifiers n = 3 2. Series of Dilate and Erode to fill masks	1. Trainable Weka segmentation classifier n = 1 2. Series of Dilate and Erode to fill masks	1. Trainable Weka segmentation classifier n = 1 2. Series of Dilate and Erode to fill masks	1. Trainable Weka segmentation classifier n = 2 2. Series of Dilate and Erode to fill masks
4. Segmentation - GF	Vibrio cells attached to microplastic: Setting intensity threshold to 90 Planktonic Vibrio cells: removing microparticle areas with brightfield mask, then setting intensity threshold to 82	Setting intensity threshold to 82	Trainable Weka segmentation classifier n = 1	Trainable Weka segmentation classifier n = 1
5. Image operations	Apply AND operation between microparticle mask and fluorescent pixel masks originating from 120-200 µm depths' pictures	Apply AND operation between microparticle mask and fluorescent pixel masks originating from: -120-200 µm depths' pictures (sinking particles) - 40-120 µm depths' pictures (floating particles)	Apply AND operation between microparticle mask and fluorescent pixel masks originating from 140-200 µm depths' pictures	Apply AND operation between microparticle mask and fluorescent pixel masks originating from 100-200 µm depths' pictures

774 **Table S3. Database accession identifiers of *V. anguillarum* virulence genes.** Database
775 compiling experimentally validated genes involved in the virulence of *V. anguillarum*, updated as
776 of January 2025.
777
778 **Table S4. Summary of samples that were not meeting requirements after image pipeline
779 analysis.** Each sheet corresponds to a separate experiment and batches of samples are
780 separated by an empty line for clarity purposes. Deviations from the pipeline description are
781 indicated for each sample.
782
783 **Table S5. Descriptive statistics of fluorescent pixel overlap on microparticles:
784 comparison between field-colonized and control particles, by nature of particles, *V.
785 anguillarum* strain, and origin of the field-colonized particles.** Negative values indicate
786 lower colonization by the focal *V. anguillarum* strain of field-colonized substrate compared to the
787 control.
788

Variables		mean	standard deviation	min	max
nature of particle	PET	-22.3	23.0	-60.3	20.9
	PP	-26.5	28.3	-78.7	34.2
	PS	22.0	27.5	-48.2	84.1
	GL	3.6	17.2	-20.8	62.2
strain	NB10	-12.5	31.1	-78.7	56.2
	VIB8	-5.8	33.8	-60.2	62.2
	VIB205	0.9	27.7	-53.3	84.1
location	BA	-4.8	22.3	-50.3	37.3
	CA	-5.1	36.3	-60.2	84.1
	TH	-7.4	33.9	-78.7	72.8

808
Figure S1. Particle size distribution curve for PET and PP microparticles used in this work.
Individual lines represent the particle size distribution of one batch.

Figure S2. Pipeline depicting megapicture reconstruction and pixel classification.
Panel A represents image capture protocol of MP/Vibrio cells suspension. Ten microliters were placed on a Malassez cell and a brightfield image of the field was taken to capture microparticle positions at a magnification of x200. A green fluorescence filter was applied to capture GFP signal, and 10 pictures of the field were taken at different heights in 20 μm intervals, encompassing the total volume of the mixture between the slide and the coverslip. Multiple pictures were captured of 10 fields, representing a grand volume of 0.01 μl for each sample. Panels B, C, D, and E represent the different steps using Fiji to generate one brightfield megapicture encompassing the 10 fields, and 10 green fluorescent megapictures (one per depth). From that, we segmented them to get a mask for both microparticles and green fluorescent pixels emitted by fluorescent bacteria. Panel F represents step during which all masks were manually checked and corrected in cases of local mis-segmentation. Panel G represents fluorescent pixel classification, based on their position compared to microplastic masks: (1) those overlapping and at the same depth were classified as emitted by attached bacteria, (2) those not overlapping were classified as planktonic bacteria, (3) those overlapping but at shallower and deeper depths were considered noise. Attachment was evaluated based on the proportion of fluorescent pixels covering the microplastic area.

Figure S3. Biofilm colonization of microplastic particles observed by light microscopy.
Horizontal bars represent a scale of 10 μm in all images. Panel A shows green fluorescent microscopy images of PS and PET particles after 24 hours if incubation with GFP-labelled *V. anguillarum* in experiment 1. Panel B shows brightfield microscopy images of microplastics after one month incubation along the Mediterranean coast, in experiment 2. Arrows indicate the boundaries of biofilms associated with particles. 'MP'= microplastic particle

Figure S4. Number of pixels analyzed in experiment 1 in reconstructed megapictures based on the polymer (*V. anguillarum* attachment on pristine microplastics). Left panel shows the size of reconstructed megapictures as a function of the polymer, expressed as a number of pixels. The middle panel shows the surface of microparticles covering pictures, while the right panel shows the total number of fluorescent pixels as a function of polymer.

Figure S5. Number of pixels analyzed in experiment 2 in reconstructed megapictures, based on the polymer (*V. anguillarum* attachment on field-colonized microplastics). Left panel shows the size of reconstructed megapictures as a function of the polymer, expressed as

a number of pixels. The middle panel shows the surface of microparticles covering pictures, while the right panel shows the total number of fluorescent pixels as a function of polymer.

Figure S6. Comparisons of proportions of fluorescent pixels covering pristine microplastics between experiments. Each graph represents attachment data for microplastics from different polymers (from left to right: PET, PP, and PS) and strains tested in both experiments (from top to bottom: VIB8, NB10, and VIB205). For each strain, gray points show the observed data, while colored diamonds indicate the mean proportion. Nine measurements per strain and polymer were obtained in experiment 1, and 27 measurements in experiment 2.

Figure S7. Change in proportion of fluorescent pixels overlap between field-colonized microparticles and control, according to the tested *V. anguillarum* strain, nature of substrate (vertical panels), and origin of the field-colonized particles (horizontal panels). A difference below zero indicates lower colonization by the focal *V. anguillarum* strain of field-colonized substrate compared to the control. For each strain, gray points show the observed data for three replicates, while colored diamonds indicate the mean proportion. The color code corresponds to the strain, and is consistent across all graphs.

Bibliography

[1] R. Geyer, J. R. Jambeck, et K. L. Law, « Production, use, and fate of all plastics ever made », *Sci. Adv.*, vol. 3, n^o 7, p. e1700782, 2017, doi: 10.1126/sciadv.1700782.

[2] S. M. Abel, F. Wu, S. Primpke, G. Gerdts, et A. Brandt, « Journey to the deep: plastic pollution in the hadal of deep-sea trenches. », *Environ. Pollut.*, vol. 333, p. 122078, sept. 2023, doi: 10.1016/j.envpol.2023.122078.

[3] S. Allen *et al.*, « Atmospheric transport and deposition of microplastics in a remote mountain catchment », *Nat. Geosci.*, vol. 12, n^o 5, p. 339‑344, mai 2019, doi: 10.1038/s41561-019-0335-5.

[4] A. N. Meyer, B. Lutz, et M. Bergmann, « Where does Arctic beach debris come from? Analyzing debris composition and provenance on Svalbard aided by citizen scientists », *Front. Mar. Sci.*, vol. 10, févr. 2023, doi: 10.3389/fmars.2023.1092939.

[5] M. Palazot *et al.*, « First national reference of microplastic contamination of French soils », *Sci. Total Environ.*, vol. 918, p. 170564, mars 2024, doi: 10.1016/j.scitotenv.2024.170564.

[6] J. R. Jambeck *et al.*, « Marine pollution. Plastic waste inputs from land into the ocean », *Science*, vol. 347, n^o 6223, p. 768‑771, févr. 2015, doi: 10.1126/science.1260352.

[7] H. A. Leslie, M. J. M. van Velzen, S. H. Brandsma, A. D. Vethaak, J. J. Garcia-Vallejo, et M. H. Lamoree, « Discovery and quantification of plastic particle pollution in human blood », *Environ. Int.*, vol. 163, p. 107199, mai 2022, doi: 10.1016/j.envint.2022.107199.

[8] A. J. Nihart *et al.*, « Bioaccumulation of microplastics in decedent human brains », *Nat. Med.*, vol. 31, n^o 4, p. 1114‑1119, avr. 2025, doi: 10.1038/s41591-024-03453-1.

[9] M. A. Garcia *et al.*, « Quantitation and identification of microplastics accumulation in human placental specimens using pyrolysis gas chromatography mass spectrometry », *Toxicol. Sci.*, vol. 199, n^o 1, p. 81‑88, mai 2024, doi: 10.1093/toxsci/kfae021.

[10] J. Zhang, L. Wang, L. Trasande, et K. Kannan, « Occurrence of polyethylene terephthalate and polycarbonate microplastics in infant and adult feces », *Environ. Sci. Technol. Lett.*, vol. 8, n^o 11, p. 989‑994, nov. 2021, doi: 10.1021/acs.estlett.1c00559.

[11] M. Peng *et al.*, « Cellular and bioenergetic effects of polystyrene microplastic in function of cell type, differentiation status and post-exposure time », *Environ. Pollut.*, vol. 337, p. 122550, nov. 2023, doi: 10.1016/j.envpol.2023.122550.

[12] M. Peng *et al.*, « Probing long-term impacts: low-dose polystyrene nanoplastics exacerbate mitochondrial health and evoke secondary glycolysis via repeated and single dosing », *Environ. Sci. Technol.*, vol. 58, n^o 23, p. 9967‑9979, juin 2024, doi: 10.1021/acs.est.3c10868.

[13] R. Marfella *et al.*, « Microplastics and nanoplastics in atheromas and cardiovascular Events », *N. Engl. J. Med.*, vol. 390, n^o 10, p. 900‑910, mars 2024, doi: 10.1056/NEJMoa2309822.

[14] Y. Deng, Y. Zhang, B. Lemos, et H. Ren, « Tissue accumulation of microplastics in mice and biomarker responses suggest widespread health risks of exposure », *Sci. Rep.*, vol. 7, n^o 1, p. 46687, avr. 2017, doi: 10.1038/srep46687.

[15] H. S. Charlton-Howard, A. L. Bond, J. Rivers-Auty, et J. L. Lavers, « 'Plasticosis': Characterising macro- and microplastic-associated fibrosis in seabird tissues », *J. Hazard. Mater.*, vol. 450, p. 131090, mai 2023, doi: 10.1016/j.jhazmat.2023.131090.

[16] J. Bowley, C. Baker-Austin, A. Porter, R. Hartnell, et C. Lewis, « Oceanic Hitchhikers – Assessing Pathogen Risks from Marine Microplastic », *Trends Microbiol.*, vol. 29, n^o 2, p. 107‑116, févr. 2021, doi: 10.1016/j.tim.2020.06.011.

[17] E. R. Zettler, T. J. Mincer, et L. A. Amaral-Zettler, « Life in the "plastisphere": microbial
communities on plastic marine debris », *Environ. Sci. Technol.*, vol. 47, n⁰ 13, p.
7137⬚7146, juill. 2013, doi: 10.1021/es401288x.
[18] D. Debroas, A. Mone, et A. Ter Halle, « Plastics in the North Atlantic garbage patch: A
boat-microbe for hitchhikers and plastic degraders », *Sci. Total Environ.*, vol. 599⬚600, p.
1222⬚1232, déc. 2017, doi: 10.1016/j.scitotenv.2017.05.059.
[19] I. V. Kirstein *et al.*, « Dangerous hitchhikers? Evidence for potentially pathogenic *Vibrio*
spp. on microplastic particles », *Mar. Environ. Res.*, vol. 120, p. 1⬚8, sept. 2016, doi:
10.1016/j.marenvres.2016.07.004.
[20] A. L. Laverty, S. Primpke, C. Lorenz, G. Gerdts, et F. C. Dobbs, « Bacterial biofilms
colonizing plastics in estuarine waters, with an emphasis on Vibrio spp. and their
antibacterial resistance », *PLOS ONE*, vol. 15, n⁰ 8, p. e0237704, août 2020, doi:
10.1371/journal.pone.0237704.
[21] M. L. Pedrotti, A. L. e Figueiredo Lacerda, S. Petit, J. F. Ghiglione, et G. Gorsky, « Vibrio
spp and other potential pathogenic bacteria associated to microfibers in the North-Western
Mediterranean Sea », *PLOS ONE*, vol. 17, n⁰ 11, p. e0275284, nov. 2022, doi:
10.1371/journal.pone.0275284.
[22] R. Tavelli, M. Callens, C. Grootaert, M. F. Abdallah, et A. Rajkovic, « Foodborne
pathogens in the plastisphere: Can microplastics in the food chain threaten microbial food
safety? », *Trends Food Sci. Technol.*, vol. 129, p. 1⬚10, nov. 2022, doi:
10.1016/j.tifs.2022.08.021.
[23] A. K. Coons, K. Busch, M. Lenz, U. Hentschel, et E. Borchert, « Biogeography rather than
substrate type determines bacterial colonization dynamics of marine plastics », *PeerJ*, vol.
9, p. e12135, 2021, doi: 10.7717/peerj.12135.
[24] X. Jiang *et al.*, « Dissemination of antibiotic resistance genes from antibiotic producers to
pathogens », *Nat. Commun.*, vol. 8, n⁰ 1, p. 15784, juin 2017, doi: 10.1038/ncomms15784.
[25] K. Kesy, S. Oberbeckmann, B. Kreikemeyer, et M. Labrenz, « Spatial environmental
heterogeneity determines young biofilm assemblages on microplastics in Baltic Sea
mesocosms », *Front. Microbiol.*, vol. 10, 2019, Consulté le: 22 mars 2022. [En ligne].
Disponible sur: https://www.frontiersin.org/article/10.3389/fmicb.2019.01665
[26] X. Sun *et al.*, « Impact of mariculture-derived microplastics on bacterial biofilm formation
and their potential threat to mariculture: A case in situ study on the Sungo Bay, China »,
Environ. Pollut., vol. 262, p. 114336, juill. 2020, doi: 10.1016/j.envpol.2020.114336.
[27] T. J. Sushmitha, M. Rajeev, P. Sriyutha Murthy, S. Ganesh, S. R. Toleti, et S. Karutha
Pandian, « Bacterial community structure of early-stage biofilms is dictated by temporal
succession rather than substrate types in the southern coastal seawater of India », *PloS*
One, vol. 16, n⁰ 9, p. e0257961, 2021, doi: 10.1371/journal.pone.0257961.
[28] R. J. Wright, M. G. I. Langille, et T. R. Walker, « Food or just a free ride? A meta-analysis
reveals the global diversity of the plastisphere », *ISME J.*, vol. 15, n⁰ 3, Art. n⁰ 3, mars
2021, doi: 10.1038/s41396-020-00814-9.
[29] J. M. Janda et S. L. Abbott, « 16S rRNA gene sequencing for bacterial identification in the
diagnostic laboratory: pluses, perils, and pitfalls », *J. Clin. Microbiol.*, vol. 45, n⁰ 9, p.
2761⬚2764, sept. 2007, doi: 10.1128/jcm.01228-07.
[30] H. Liang *et al.*, « Detection of faecal bacteria and antibiotic resistance genes in biofilms
attached to plastics from human-impacted coastal areas », *Environ. Pollut. Barking Essex*
1987, vol. 319, p. 120983, févr. 2023, doi: 10.1016/j.envpol.2022.120983.
[31] V. Radisic, P. S. Nimje, A. M. Bienfait, et N. P. Marathe, « Marine plastics from Norwegian
West Coast carry potentially virulent fish pathogens and opportunistic human pathogens
harboring new variants of antibiotic resistance genes », *Microorganisms*, vol. 8, n⁰ 8, p.
1200, août 2020, doi: 10.3390/microorganisms8081200.

[32] A. Lianou et K. P. Koutsoumanis, « Strain variability of the biofilm-forming ability of
Salmonella enterica under various environmental conditions », *Int. J. Food Microbiol.*, vol.
160, n° 2, p. 171‑178, nov. 2012, doi: 10.1016/j.ijfoodmicro.2012.10.002.
[33] J. Garita-Cambronero, A. Palacio-Bielsa, M. M. López, et J. Cubero, « Comparative
genomic and phenotypic characterization of pathogenic and non-pathogenic strains of
Xanthomonas arboricola reveals insights into the infection process of bacterial spot
disease of stone fruits », *PLOS ONE*, vol. 11, n° 8, p. e0161977, août 2016, doi:
10.1371/journal.pone.0161977.
[34] F. Oliveira, C. A. Lima, S. Brás, Â. França, et N. Cerca, « Evidence for inter- and
intraspecies biofilm formation variability among a small group of coagulase-negative
staphylococci », *FEMS Microbiol. Lett.*, vol. 362, n° 20, p. fnv175, oct. 2015, doi:
10.1093/femsle/fnv175.
[35] S. Zhao *et al.*, « The distribution of subsurface microplastics in the ocean », *Nature*, vol.
641, n° 8061, p. 51‑61, mai 2025, doi: 10.1038/s41586-025-08818-1.
[36] P. Jiang, S. Zhao, L. Zhu, et D. Li, « Microplastic-associated bacterial assemblages in the
intertidal zone of the Yangtze Estuary », *Sci. Total Environ.*, vol. 624, p. 48‑54, mai 2018,
doi: 10.1016/j.scitotenv.2017.12.105.
[37] F. Le Roux, B. M. Davis, et M. K. Waldor, « Conserved small RNAs govern replication and
incompatibility of a diverse new plasmid family from marine bacteria », *Nucleic Acids Res.*,
vol. 39, n° 3, p. 1004‑1013, févr. 2011, doi: 10.1093/nar/gkq852.
[38] S. Andrews, « FastQC: a quality control tool for high throughput sequence data ». 2010.
[En ligne]. Disponible sur: http://www.bioinformatics.babraham.ac.uk/projects/fastqc/
[39] M. Martin, « Cutadapt removes adapter sequences from high-throughput sequencing
reads », *EMBnet.journal*, vol. 17, n° 1, Art. n° 1, mai 2011, doi: 10.14806/ej.17.1.200.
[40] W. De Coster, S. D'Hert, D. T. Schultz, M. Cruts, et C. Van Broeckhoven, « NanoPack:
visualizing and processing long-read sequencing data », *Bioinformatics*, vol. 34, n° 15, p.
2666‑2669, août 2018, doi: 10.1093/bioinformatics/bty149.
[41] W. De Coster et R. Rademakers, « NanoPack2: population-scale evaluation of long-read
sequencing data », *Bioinformatics*, vol. 39, n° 5, p. btad311, mai 2023, doi:
10.1093/bioinformatics/btad311.
[42] G. Bouras *et al.*, « Hybracter: enabling scalable, automated, complete and accurate
bacterial genome assemblies », *Microb. Genomics*, vol. 10, n° 5, p. 001244, 2024, doi:
10.1099/mgen.0.001244.
[43] T. Seemann, *Prokka: rapid prokaryotic genome annotation*. (28 février 2022). Perl.
Consulté le: 8 mars 2022. [En ligne]. Disponible sur: https://github.com/tseemann/prokka
[44] D. H. Parks, M. Imelfort, C. T. Skennerton, P. Hugenholtz, et G. W. Tyson, « CheckM:
assessing the quality of microbial genomes recovered from isolates, single cells, and
metagenomes », *Genome Res.*, vol. 25, n° 7, p. 1043‑1055, janv. 2015, doi:
10.1101/gr.186072.114.
[45] A. J. Page *et al.*, « Roary: rapid large-scale prokaryote pan genome analysis »,
Bioinformatics, vol. 31, n° 22, p. 3691‑3693, nov. 2015, doi:
10.1093/bioinformatics/btv421.
[46] K. Katoh, K. Misawa, K. Kuma, et T. Miyata, « MAFFT: a novel method for rapid multiple
sequence alignment based on fast Fourier transform », *Nucleic Acids Res.*, vol. 30, n° 14,
p. 3059‑3066, juill. 2002.
[47] T. Seemann, *tseemann/snp-dists*. (21 mars 2025). C. Consulté le: 21 mars 2025. [En
ligne]. Disponible sur: https://github.com/tseemann/snp-dists
[48] B. Q. Minh *et al.*, « IQ-TREE 2: new models and efficient methods for phylogenetic
inference in the genomic era », *Mol. Biol. Evol.*, vol. 37, n° 5, p. 1530‑1534, mai 2020, doi:
10.1093/molbev/msaa015.

[49]   S. Kalyaanamoorthy, B. Q. Minh, T. K. Wong, A. von Haeseler, et L. S. Jermiin,
1027   « ModelFinder: fast model selection for accurate phylogenetic estimates », *Nat. Methods*,
1028   vol. 14, n° 6, p. 587‑589, juin 2017, doi: 10.1038/nmeth.4285.
1029   [50]   T. Kaseke *et al.*, « Polypropylene micro- and nanoplastics affect the digestion of cow's milk
1030   proteins in infant model of gastric digestion », *Environ. Pollut.*, vol. 383, p. 126803, oct.
1031   2025, doi: 10.1016/j.envpol.2025.126803.
1032   [51]   S. Ducoli *et al.*, « A different protein corona cloaks "true-to-life" nanoplastics with respect to
1033   synthetic polystyrene nanobeads », *Environ. Sci. Nano*, vol. 9, n° 4, p. 1414‑1426, avr.
1034   2022, doi: 10.1039/D1EN01016F.
1035   [52]   J. Schindelin *et al.*, « Fiji: an open-source platform for biological-image analysis », *Nat.*
1036   *Methods*, vol. 9, n° 7, p. 676‑682, juill. 2012, doi: 10.1038/nmeth.2019.
1037   [53]   J. Chalfoun *et al.*, « MIST: accurate and scalable microscopy image stitching tool with
1038   stage modeling and error minimization », *Sci. Rep.*, vol. 7, n° 1, p. 4988, juill. 2017, doi:
1039   10.1038/s41598-017-04567-y.
1040   [54]   T. Blattner, W. Keyrouz, J. Chalfoun, B. Stivalet, M. Brady, et S. Zhou, « A hybrid CPU-
1041   GPU system for stitching large scale optical microscopy images », in *2014 43rd*
1042   *International Conference on Parallel Processing*, sept. 2014, p. 1‑9. doi:
1043   10.1109/ICPP.2014.9.
1044   [55]   I. Arganda-Carreras *et al.*, « Trainable Weka Segmentation: a machine learning tool for
1045   microscopy pixel classification », *Bioinformatics*, vol. 33, n° 15, p. 2424‑2426, août 2017,
1046   doi: 10.1093/bioinformatics/btx180.
1047   [56]   I. Arganda-Carreras, V. Kaynig, C. Rueden, J. Schindelin, A. Cardona, et H. S. Seung,
1048   *Trainable_Segmentation: Release v3.1.2*. (3 août 2016). Zenodo. doi:
1049   10.5281/zenodo.59290.
1050   [57]   E. Paradis et K. Schliep, « ape 5.0: an environment for modern phylogenetics and
1051   evolutionary analyses in R », *Bioinformatics*, vol. 35, n° 3, p. 526‑528, févr. 2019, doi:
1052   10.1093/bioinformatics/bty633.
1053   [58]   E. W. Sayers *et al.*, « Database resources of the National Center for Biotechnology
1054   Information », *Nucleic Acids Res.*, vol. 52, n° D1, p. D33‑D43, nov. 2023, doi:
1055   10.1093/nar/gkad1044.
1056   [59]   A. Fukuda, Y. Kodama, J. Mashima, T. Fujisawa, et O. Ogasawara, « DDBJ update:
1057   streamlining submission and access of human data », *Nucleic Acids Res.*, vol. 49, n° D1,
1058   p. D71‑D75, janv. 2021, doi: 10.1093/nar/gkaa982.
1059   [60]   T. Seemann, *ABRicate*. (7 mars 2022). Perl. Consulté le: 8 mars 2022. [En ligne].
1060   Disponible sur: https://github.com/tseemann/abricate
1061   [61]   L. Chen *et al.*, « VFDB: a reference database for bacterial virulence factors », *Nucleic*
1062   *Acids Res.*, vol. 33, n° Database issue, p. D325-328, janv. 2005, doi: 10.1093/nar/gki008.
1063   [62]   L. Chen, D. Zheng, B. Liu, J. Yang, et Q. Jin, « VFDB 2016: hierarchical and refined
1064   dataset for big data analysis--10 years on », *Nucleic Acids Res.*, vol. 44, n° D1, p. D694-
1065   697, janv. 2016, doi: 10.1093/nar/gkv1239.
1066   [63]   M. Lechner, S. Findeiß, L. Steiner, M. Marz, P. F. Stadler, et S. J. Prohaska,
1067   « Proteinortho: Detection of (Co-)orthologs in large-scale analysis », *BMC Bioinformatics*,
1068   vol. 12, n° 1, p. 124, avr. 2011, doi: 10.1186/1471-2105-12-124.
1069   [64]   J. Rojas-Vargas, H. G. Castelán-Sánchez, et L. Pardo-López, « HADEG: A curated
1070   hydrocarbon aerobic degradation enzymes and genes database », *Comput. Biol. Chem.*,
1071   vol. 107, p. 107966, déc. 2023, doi: 10.1016/j.compbiolchem.2023.107966.
1072   [65]   M. Massot, E. Gadoin, A. Rajkovic, et S. Bedhomme, « Microplastics in marine
1073   environments: protocol for isolating natural biofilms from seawater-incubated particles. »,
1074   protocols.io. Consulté le: 16 septembre 2024. [En ligne]. Disponible sur:
1075   dx.doi.org/10.17504/protocols.io.kqdg3pzjel25/v1

[66] M. Arias-Andres, U. Klümper, K. Rojas-Jimenez, et H.-P. Grossart, « Microplastic pollution
increases gene exchange in aquatic ecosystems », *Environ. Pollut.*, vol. 237, p. 253‑261,
juin 2018, doi: 10.1016/j.envpol.2018.02.058.

[67] P.-C. Bürkner, « brms: an R package for Bayesian multilevel models using Stan », *J. Stat.*
Softw., vol. 80, p. 1‑28, août 2017, doi: 10.18637/jss.v080.i01.

[68] P.-C. Bürkner, « Bayesian item response modeling in R with brms and Stan », *J. Stat.*
Softw., vol. 100, p. 1‑54, nov. 2021, doi: 10.18637/jss.v100.i05.

[69] B. Carpenter *et al.*, « Stan: A Probabilistic Programming Language », *J. Stat. Softw.*, vol.
76, p. 1‑32, janv. 2017, doi: 10.18637/jss.v076.i01.

[70] T. Roder *et al.*, « Scoary2: rapid association of phenotypic multi-omics data with microbial
pan-genomes », *Genome Biol.*, vol. 25, p. 93, avr. 2024, doi: 10.1186/s13059-024-03233-
7.

[71] R core team, *R: A language and environment for statistical computing.* (2019). R
Foundation for Statistical Computing, Vienna, Austria. [En ligne]. Disponible sur:
https://www.r-project.org/

[72] J. B. Lamb *et al.*, « Plastic waste associated with disease on coral reefs », *Science*, vol.
359, n^o 6374, p. 460‑462, janv. 2018, doi: 10.1126/science.aar3320.

[73] null Suyadi et C. Y. Manullang, « Distribution of plastic debris pollution and it is
implications on mangrove vegetation », *Mar. Pollut. Bull.*, vol. 160, p. 111642, nov. 2020,
doi: 10.1016/j.marpolbul.2020.111642.

[74] C. E. J. van Bijsterveldt *et al.*, « Does plastic waste kill mangroves? A field experiment to
assess the impact of macro plastics on mangrove growth, stress response and survival »,
Sci. Total Environ., vol. 756, p. 143826, févr. 2021, doi: 10.1016/j.scitotenv.2020.143826.

[75] S. Rech, M. Thiel, Y. J. Borrell Pichs, et E. García-Vazquez, « Travelling light: Fouling
biota on macroplastics arriving on beaches of remote Rapa Nui (Easter Island) in the
South Pacific Subtropical Gyre », *Mar. Pollut. Bull.*, vol. 137, p. 119‑128, déc. 2018, doi:
10.1016/j.marpolbul.2018.10.015.

[76] V. Foulon, F. Le Roux, C. Lambert, A. Huvet, P. Soudant, et I. Paul-Pont, « Colonization of
polystyrene microparticles by *Vibrio crassostreae*: light and electron microscopic
investigation », *Environ. Sci. Technol.*, vol. 50, n^o 20, p. 10988‑10996, oct. 2016, doi:
10.1021/acs.est.6b02720.

[77] M. R. Hossain, M. Jiang, Q. Wei, et L. G. Leff, « Microplastic surface properties affect
bacterial colonization in freshwater », *J. Basic Microbiol.*, vol. 59, n^o 1, p. 54‑61, 2019, doi:
10.1002/jobm.201800174.

[78] Y. Cheng, G. Feng, et C. I. Moraru, « Micro- and nanotopography sensitive bacterial
attachment mechanisms: a review », *Front. Microbiol.*, vol. 10, févr. 2019, doi:
10.3389/fmicb.2019.00191.

[79] R. E. Leighton *et al.*, « *Vibrio parahaemolyticus* and *Vibrio vulnificus in vitro* colonization on
plastics influenced by temperature and strain variability », *Front. Microbiol.*, vol. 13, janv.
2023, doi: 10.3389/fmicb.2022.1099502.

[80] E. Robino *et al.*, « Resistance of the oyster pathogen *Vibrio tasmaniensis* LGP32 against
grazing by Vannella sp. marine amoeba involves Vsm and CopA virulence factors »,
Environ. Microbiol., vol. 22, n^o 10, p. 4183‑4197, 2020, doi: 10.1111/1462-2920.14770.

[81] A. Morot, F. Delavat, A. Bazire, C. Paillard, A. Dufour, et S. Rodrigues, « Genetic insights
into biofilm formation by a pathogenic strain of *Vibrio harveyi* », *Microorganisms*, vol. 12, n^o
1, p. 186, janv. 2024, doi: 10.3390/microorganisms12010186.

[82] K. Kesy, M. Labrenz, B. S. Scales, B. Kreikemeyer, et S. Oberbeckmann, « Vibrio
colonization is highly dynamic in early microplastic-associated biofilms as well as on field-
collected microplastics », *Microorganisms*, vol. 9, n^o 1, p. 76, janv. 2021, doi:
10.3390/microorganisms9010076.

[83] M. S. Datta, E. Sliwerska, J. Gore, M. F. Polz, et O. X. Cordero, « Microbial interactions
lead to rapid micro-scale successions on model marine particles », *Nat. Commun.*, vol. 7,
n^o 1, p. 11965, juin 2016, doi: 10.1038/ncomms11965.
[84] H. Deng *et al.*, « Microplastic-associated biofilm in an intensive mariculture pond: Temporal
dynamics of microbial communities, extracellular polymeric substances and impacts on
microplastics properties », *J. Clean. Prod.*, vol. 319, p. 128774, oct. 2021, doi:
10.1016/j.jclepro.2021.128774.
[85] X. Xu *et al.*, « Marine microplastic-associated bacterial community succession in response
to geography, exposure time, and plastic type in China's coastal seawaters », *Mar. Pollut.*
Bull., vol. 145, p. 278⬛286, août 2019, doi: 10.1016/j.marpolbul.2019.05.036.
[86] C. Lemonnier *et al.*, « Time-series incubations in a coastal environment illuminates the
importance of early colonizers and the complexity of bacterial biofilm dynamics on marine
plastics », *Environ. Pollut.*, vol. 312, p. 119994, nov. 2022, doi:
10.1016/j.envpol.2022.119994.
[87] O. Ciofu, C. Moser, P. Ø. Jensen, et N. Høiby, « Tolerance and resistance of microbial
biofilms », *Nat. Rev. Microbiol.*, vol. 20, n^o 10, p. 621⬛635, oct. 2022, doi: 10.1038/s41579-
022-00682-4.
[88] J.-Y. Maillard et I. Centeleghe, « How biofilm changes our understanding of cleaning and
disinfection », *Antimicrob. Resist. Infect. Control*, vol. 12, p. 95, sept. 2023, doi:
10.1186/s13756-023-01290-4.
[89] E. M. Stevenson, O. Rushby-Jones, A. Buckling, M. Cole, P. K. Lindeque, et A. K. Murray,
« Selective colonization of microplastics, wood and glass by antimicrobial-resistant and
pathogenic bacteria », *Microbiology*, vol. 170, n^o 10, p. 001506, 2024, doi:
10.1099/mic.0.001506.
[90] M. De la Cruz Barrón, C. Merlin, H. Guilloteau, E. Montargès-Pelletier, et X. Bellanger,
« Suspended materials in river waters differentially enrich Class 1 Integron- and IncP-1
plasmid-carrying bacteria in sediments », *Front. Microbiol.*, vol. 9, juill. 2018, doi:
10.3389/fmicb.2018.01443.

1154
1155
1156