On the correctness of gene tree tagging and the consistency of ASTRAL-pro under a unified model of gene duplication, loss, and coalescence

Rachel Parsons[1], Yunzhuo Liu[1], Parth Dua[1], Alexey Markin[2], and Erin K. Molloy[1,3,*]

[1] *Department of Computer Science, University of Maryland, College Park, 20742, USA*

[2] *Virus and Prion Research Unit, National Animal Disease Center, Agricultural Research Service,*
United States Department of Agriculture, Ames, IA, 50010

[3] *University of Maryland Institute for Advanced Computer Studies, College Park, MD 20740*

*Email: ekmolloy@umd.edu

January 21, 2026

Abstract

Motivation: ASTRAL-pro is the leading method for reconstructing species trees under complex evolutionary scenarios involving gene duplication, loss, and coalescence. A major open question is whether ASTRAL-pro is statistically consistent under a unified model of these processes, called DLCoal. This question is challenging to address because ASTRAL-pro seeks a species tree that maximizes the number of four-taxon trees (called quartets) also displayed by the input (multi-copy) gene trees, *excluding* those induced by duplications and *agglomerating* those that are homeomorphic up to duplications. Critically, there is no notion of correctness when tagging gene tree vertices as duplication or speciation events in the context of deep coalescence.

Results: Here, we propose that a gene tree vertex is correctly tagged as a duplication if it is the most recent common ancestor of *at least one* pair of gene copies related via a duplication event. Under our definition, deep coalescence propagates duplication tags across gene tree vertices, sometimes resulting in the exclusion of quartets on orthologous gene copies. Nevertheless, we show that A-pro is statistically consistent under the DLCoal model for an exclusion-only version of its objective function, assuming the input gene trees are correctly rooted and tagged. To empirically evaluate this modification, we exclude "duplication quartets" in the related method TREE-QMC and find that it achieves similar accuracy to A-pro on simulated data under varying rates of deep coalescence, duplication and loss, and gene tree estimation error, as well as on a plant data set.

Availability and Implementation: TREE-QMC-pro is available on Github: https://github.com/molloy-lab/TREE-QMC/tree/tqmc-pro.

Keywords: Quartets, Multispecies Coalescent, Gene Duplication and Loss, Gene Tree Estimation Error

1   Introduction

The availability of genome-scale data sets has increased dramatically over the last decade, thanks to technological advances and large-scale initiatives, like One Thousand Plant Transcriptomes [Leebens-Mack et al., 2019] and Bird 10K [Feng et al., 2020]. Fully leveraging these data for species tree reconstruction requires accounting for gene tree heterogeneity (GTH), which broadly refers to variation in the evolutionary histories across different genomic regions, called genes [Maddison, 1997]. Perhaps the most pervasive source of GTH is incomplete lineage sorting (ILS) [Edwards, 2009], where gene trees disagree with the species tree due to deep coalescence, a population-genetics phenomenon, modeled by the Multi-Species

Coalescent (MSC) [Pamilo and Nei, 1988, Rannala and Yang, 2003]. Under the MSC, the most probable gene tree agrees with the species tree for unrooted four-taxon trees, called quartets [Allman et al., 2011]. This result motivates the most popular species tree estimation method, ASTRAL, which seeks a species tree that maximizes the number of quartets also induced by the input gene trees, estimated from molecular sequences [Mirarab et al., 2014].

ASTRAL's popularity may be attributed to its practical properties (speed and accuracy) as well as its statistical guarantee; specifically, ASTRAL is *consistent* under the MSC, meaning that it returns the true unrooted species tree topology with high probability, as the number of gene trees generated under the MSC tends to infinity. An underlying assumption is that genes evolve without duplications. Thus, multi-copy gene families must be restricted to orthologs or else excluded from species tree reconstruction. The former can present computational challenges (see Majidian et al., 2025 for a recent review), and the latter can dramatically reduce the amount of data available for species tree reconstruction (consider that Wickett et al., 2014 curated more than 9,000 gene families but just 424 were single-copy). To address this issue, some methods reconstruct species trees directly from multi-copy gene trees. However, the accuracy of traditional approaches based on gene tree parsimony (e.g., Parsons and Bansal, 2025) or Robinson-Foulds supertrees (e.g., Molloy and Warnow, 2020) is challenged by deep coalescence as well as gene tree estimation error (Fig. 2b in Zhang et al., 2020).

The current leading methods are based on quartets. ASTRAL-multi (called A-multi for brevity) extends ASTRAL's objective function to multi-copy genes in the natural way [Rabiee et al., 2019]. Although A-multi is statistically consistent under a unified duplication-loss-coalescent model, called DLCoal [Markin and Eulenstein, 2021, Hill et al., 2022], it is less accurate than a related method, called A-pro, in simulations (Figs. 2b and 4 in Zhang et al. [2020]). Unlike A-multi, A-pro internally roots and *tags* gene trees so that each internal vertex is labeled as duplication or speciation. This information is used to *exclude* quartets induced by duplications, called "duplication quartets", and *agglomerate* the remaining quartets that are homeomorphic up to duplications. Tagging these events is natural when genes evolve with gene duplication and loss (GDL) but not with deep coalescence. Perhaps as a result, the consistency of A-pro under the DLCoal model has remained a major open question since its introduction in 2020.

To address this challenge, we propose a definition of correct tagging that can be applied when genes evolve with deep coalescence, in addition to duplication and loss. Specifically, we argue that a gene tree vertex is correctly tagged as a duplication if it is the most recent common ancestor of *at least one* pair of gene copies related via a duplication event. As we will show, under our definition, deep coalescence propagates duplication tags across gene tree vertices, sometimes resulting in the exclusion of quartets on orthologous gene copies. Nevertheless, we show that A-pro is consistent under the DLCoal model for an exclusion-only version of its scoring function, assuming the input gene trees are correctly rooted and tagged; we leave agglomeration to future work.

To empirically evaluate the modified objective function, we exclude "duplication quartets" within TREE-QMC [Han and Molloy, 2023], which is a heuristic for the same optimization problem as ASTRAL. We also implement functionality for rooting/tagging trees as well as providing rooted/tagged trees directly as input. These features enable us to systematically evaluate how different rates of deep coalescence, GDL, and estimation error impact the accuracy of gene tree tagging and, in turn, species tree reconstruction under different quartet-based objective functions. We find that our new method TQMC-pro achieves similar results to A-pro on these simulated data as well as on a plant data set. Moreover, both A-pro and TQMC-pro outperform methods that do not exclude "duplication quartets," especially for small numbers of genes. Collectively, our contributions mark critical steps forward in understanding the performance of A-pro and related techniques in theory and in practice.

2   Background

To introduce our theoretical results, we review terminology and background for phylogenetic methods and the DLCoal model.

2.1   Terminology

A *phylogenetic tree* T is a tuple $(g, \mathcal{X}, \phi)$, where g is a tree in the graph-theoretic sense, $\mathcal{X}$ is a set of species labels (also called taxa), and ϕ is a map from leaves of g to species labels in $\mathcal{X}$. We say that T is *singly-labeled* if ϕ is a bijection; otherwise, we say that T is *multi-labeled*. For simplicity, we use lower case letters to denote the leaves of gene and locus trees and upper case letters to denote their respective

species labels (i.e., $\phi(a) = A$). Sometimes we use subscripts (e.g., $a_0, a_1, \dots$) to indicate different leaves with the same species label. We let $L(T)$ denote the leaf set of T.

A phylogenetic tree T is *rooted* if there exists a directed path from a special vertex with in-degree zero, called the root, to each leaf; it is *unrooted* if all edges are undirected. Non-terminal edges are called internal edges, and non-leaf vertices are called internal vertices. Trees are assumed to be *binary*, meaning that every internal vertex except the root has degree 3 (note that the root vertex has degree 2, otherwise degree 2 vertices are prohibited).

A **quartet** is an unrooted phylogenetic tree with four uniquely labeled leaves. There are three possible quartet topologies indicated by the one internal edge that splits the species set in half; for example, $q_1 = A, B|C, D$, $q_2 = A, C|B, D$, and $q_3 = A, D|B, C$ are the three quartet topologies on species set $\{A, B, C, D\}$. A quartet can be rooted at its internal branch, which yields a **balanced tree**, or at one of its four terminal branches, which yields **pectinate tree** (sometimes called a caterpillar tree). A larger tree T *induces* a collection of quartets that can be computed by restricting T to every subset of four uniquely labeled leaves (i.e., deleting all other leaves and suppressing vertices of degree 2). Singly-labeled trees can be restricted according to their species labels as the outcome is unambiguous. If T is rooted, the induced subtree on A, B, C, D, denoted $T|_{A,B,C,D}$, inherits the rooting, although it is ignored when convenient and clear from context.

2.2 DLCoal Model

The DLCoal model, introduced by Rasmussen and Kellis [2012], is parameterized by a **species tree** T on taxon set $\mathcal{X}$. Species trees must be rooted, binary, singly-labeled. Each branch e of T is annotated with length $t(e)$, indicating the number of generations, as well as width $N_e(e)$, indicating the effective population size. Additionally, duplication and loss rates, λ and μ respectively, must be specified (rates correspond to the number of events per generation per lineage). Gene trees evolve within a species tree under duplication, loss, and coalescent processes in two steps, which we summarize next (see Hill et al., 2022 for details).

First, a **locus tree** tree ℓ evolves within the species tree T according to the stochastic GDL model of Arvestad et al. [2009] (Fig. S1A–B). A single lineage starts at the root of T and grows forward in time. Lineages are duplicated and lost on the branches of T with rate λ and μ, respectively. A **duplication event** instantaneously splits the impacted lineage, with one side being arbitrarily designated as the original (i.e., mother) copy and the other being designated as the duplicated (i.e., daughter) copy. A loss event prohibits the impacted lineage from growing forward (it is later pruned from ℓ). When a lineage arrives at an internal node v of T, it undergoes a **speciation event**, instantaneously splitting so that one side enters the branch $v \mapsto v.l$ and the other side enters the branch $v \mapsto v.r$, where $v.l$ and $v.r$ indicate the left and right child of v, respectively. Once a lineage arrives at a leaf vertex of T, it inherits the associated taxon label; likewise, branches in ℓ inherit lengths and widths from T.

Next, a **gene tree** g evolves within the locus tree ℓ under the Multilocus Coalesecent (MLC) (Fig. S1C–D), which is closely related to the MSC. Lineages sampled at leaves of ℓ trace their ancestry backward in time. Multiple lineages entering same branch of ℓ can coalesce with equal probability; the resulting internal vertices represent **coalescent events**. The *bounded coalescent* is used to force gene copies to coalesce prior to the duplication event from which they originate (e.g., gene copies a_2 and b_2 in Fig. S1C–D). Lineages entering the branch above the root of ℓ will eventually coalesce into a single lineage [Rannala and Yang, 2003].

To summarize, the DLCoal model, given species tree T on taxon set $\mathcal{X}$, generates a gene-locus tree pair, denoted $\ell, g \sim DLCoal(T, \mathcal{X})$, although only g is observed. Because of this generation process, ℓ, g are rooted and binary and can be multi-labeled. One lineage is sampled per leaf in ℓ, so there is a bijection between the leaf sets of ℓ and g (so we reference both of them with the variables, typically a, b, c, d, which are labeled A, B, C, D, respectively). Any collection of gene trees generated under DLCoal is assumed to be independent and identically distributed.

2.3 Maximum Quartet Support Species Trees (MQSST)

ASTRAL is a heuristic for the NP-hard MQSST problem [Lafond and Scornavacca, 2019], defined by the following input/output:

- **Input**: a set $\mathcal{T}$ of k unrooted, binary trees, each with leaves labeled by species set X
- **Output**: an unrooted, binary tree with leaves bijectively labeled by X that maximizes the number of quartets also displayed by the input gene trees

Likewise, A-multi is a heuristic for MQSST but extended to multi-labeled input trees in the natural way. Let $N^t(T|_{A,B,C,D})$ be <u>number</u> of quartets in a multi-labeled input tree t that agree with some singly-labeled tree T restricted to A, B, C, D, that is

$$N^t(T|_{A,B,C,D}) = \sum_{\substack{a \in L(t): \\ \phi(a)=A}} \sum_{\substack{b \in L(t): \\ \phi(b)=B}} \sum_{\substack{c \in L(t): \\ \phi(c)=C}} \sum_{\substack{d \in L(t): \\ \phi(d)=D}} \delta(T|_{A,B,C,D} \cong t|_{a,b,c,d}) \tag{1}$$

where $\delta(\cdot)$ returns 1 if the two quartets are isomorphic and 0 otherwise. Then, A-multi's objective function is

$$Z^{\mathcal{T}}(T) = \sum_{t \in \mathcal{T}} \sum_{\substack{A,B,C,D \in X: \\ A \neq B \neq C \neq D}} N^t(T|_{A,B,C,D}). \tag{2}$$

It is easy to see that A-multi's objective function is equivalent to ASTRAL's when the input trees are singly-labeled.

As previously mentioned, the leading method A-pro is similar to A-multi but (1) *excludes* "duplication quartets" and (2) *agglomerates* "speciation quartets" that are homeomorphic up to duplications. A quartet is classified as a **speciation quartet** (SQ) if for every selection of three leaves, their most recent common ancestor (MRCA) is a speciation vertex (Fig. S1E–F); otherwise, the quartet is classified as a **duplication quartet** (DQ) (Fig. S1G–H). Thus, a balanced quartet is an SQ if the root (i.e., the MRCA of its leaves) is a speciation vertex; in contrast, a pectinate quartet is an SQ if both the root and the MRCA of its three non-outgroup leaves are speciation vertices. To classify quartets as SQs or DQs, the input trees must be rooted and **tagged** so that their internal vertices are labeled as duplications or speciations. For convenience, A-pro implements its own tagging algorithm and roots each input tree such that the GDL score associated with tagging is minimized.

The motivation behind A-pro is that (1) quartets whose topologies are driven by duplication vertices do not represent information about speciation events and thus should be excluded and (2) isomorphic quartets whose topologies are driven by the same speciation vertices (but whose leaves differ due to duplications) do not provide new information and thus should be counted as a single unit [Zhang et al., 2020]. Henceforth, we focus on a version of A-pro's objective function that counts SQs only, denoted by subscript sq (e.g., δ_{sq}, N_{sq}, and Z_{sq}); we leave agglomeration of homeomorphic SQs to future work.

3 Theoretical Results and Algorithms

3.1 Correctness of Gene Tree Tagging under DLCoal

As previously mentioned, when genes evolve with GDL only, there is a natural definition of correct tagging. Take Figure S1A–B, in which locus tree ℓ evolves within species tree T, as an example. Consider the internal vertex v in locus tree ℓ associated with the duplication event at the MRCA of A, B in T. The leaves below $v.l$ (i.e., a_0, b_0) are descendants of the original (i.e., mother) copy, and those below $v.r$ (i.e., a_2, b_2) are descendants of the duplicated (i.e., daughter) copy. Any pair of leaves with MRCA v descend from the same duplication event and thus are **paralogs**.

This argument breaks down with deep coalescence. Take Figure S1C–D, in which gene tree g evolves within locus tree ℓ, as an example. Consider the internal vertex u in gene tree g with leaves b_0, a_2, b_2 descending from $u.l$ and leaves a_0, c_0, d_0 descending from $u.r$. Looking at the collection of leaf pairs with MRCA u, some are related via a duplication event (e.g., a_2 and a_0) but others are not (e.g., b_0 and a_0). The latter occurs because of deep coalescence (e.g., both a_0 and c_0 fail to coalesce until they reach a deeper ancestral population, after which they coalesce with each other). To summarize, whether or not u should be tagged as a duplication vertex depends on the gene copies being examined; however, a single tag must be designated. We take the conservative approach.

Definition 1 (Correct Tagging for Gene Trees). *Let ℓ, g be a locus-gene tree pair, and let u be the internal node in g. We say that u is correctly tagged as a duplication if there exists one leaf x below $u.l$ and one leaf y below $u.r$ such that x, y are paralogs (i.e., the MRCA of x, y in ℓ is a duplication vertex).*

Our definition of correct tagging has two advantages. First, it is backwards compatible with locus trees (i.e., applying our definition of correct tagging to a locus tree successfully captures the duplication events). Second, it agrees with A-pro's tagging algorithm, which would designate u as a duplication vertex because it is the MRCA of two gene copies a_0 and a_1 in the same species A so they must be related via a duplication event. It is worth noting that A-pro's algorithm can fail to correctly tag gene trees, and even locus trees, because duplications and losses can produce locus subtrees that appear to

Figure S1: **A)** Locus tree evolving within species tree, with duplications (no losses shown). **B)** Correctly tagged locus tree with green circles indicating duplication events. **C)** Gene tree evolving within locus tree, with deep coalescence (branch lengths/widths of locus tree are not drawn to scale). Black circles represent coalescent events. **D)** Correctly tagged gene tree with green circles indicating tagged duplication vertices. **E)** There are $3 \cdot 3 \cdot 2 \cdot 2 = 36$ ways to select four gene copies (i.e., leaves) uniquely labeled by A, B, C, D. Five induce SQs in the locus tree, corresponding to the ways of selecting four gene copies that descend from the same locus lineage at R, shown in black and blue. The SQ on a_2, b_2, c_0, c_0 is <u>not</u> shown. **F)** Because of deep coalescence, just one of the five SQs in the locus tree is also an SQ in the gene tree. **G)** and **H)** Example DQs induced by the locus tree and gene tree, respectively.

be single-copy but are not (Fig. 2D in Molloy and Warnow, 2020). Similar issues can arise due to deep coalescence when tagging gene trees.

An interesting phenomenon arises from our definition of correct tagging, which will later be important in our proof of consistency under the DLCoal model. Observe that quartets on **orthologs** can be correctly tagged as duplications because of deep coalescence (e.g., a_0, b_0, c_0, d_0 in Fig. S1H). Although this may seem odd, recall that SQs are not restricted to orthologs, even for locus trees (e.g., the quartet on paralogs a_2, b_0, c_0, d_0 is an SQ in Fig. S1E).

3.2   Consistency of A-pro under DLCoal

We are now ready to prove our main result:

Theorem 1. *An optimal solution under an exclusion-only version of A-pro's objective function is a consistent estimator of the (unrooted) species tree (topology) under the DLCoal model, assuming the input gene trees are correctly tagged (we implicitly assume no estimation errors or errors in rooting).*

Proof. Let $\mathcal{P}$ be a collection of gene trees generated under the DLCoal model given species tree T on species set $\mathcal{X}$. We want to show that the optimal solution to an exclusion-only scoring function T^* is isomorphic to T with high probability, as $|\mathcal{P}|$ tends to ∞. Our proof follows the structure of Theorem 3 in Legried et al. [2021]. Let $\hat{T}$ be an arbitrary species tree on $\mathcal{X}$. By gene tree independence and the law of large numbers, $Z_{sq}^{\mathcal{P}}(\hat{T})/|\mathcal{P}|$ converges almost surely to its expectation simultaneously for all species trees on X. Thus, its expectation can be simplified as

$$\mathbb{E}\Big[\frac{1}{|\mathcal{P}|}Z_{sq}^{\mathcal{P}}(\hat{T})\Big] = \sum_{\substack{A,B,C,D \in X: \\ A \neq B \neq C \neq D}} \mathbb{E}\big[N_{sq}^{g}(\hat{T}|_{A,B,C,D})\big] \qquad (3)$$

where g is an arbitrary tree in $\mathcal{P}$, that is $g \sim DLCoal(T, \mathcal{X})$, and $N^g_{sq}(\hat{T}|_{A,B,C,D})$ is a random variable for the <u>number</u> of SQs induced by gene tree g that are isomorphic to $\hat{T}|_{A,B,C,D}$ (Equation 1). Then, it is easy to see that $Z^{\mathcal{P}}_{sq}(\hat{T})$ will be maximized by the true species tree T if for an arbitrary subset of four species A, B, C, D in $\mathcal{X}$, the following inequality holds

$$\mathbb{E}\big[N^g_{sq}(q_1)\big] > \max\big\{\mathbb{E}\big[N^g_{sq}(q_2)\big], \mathbb{E}\big[N^g_{sq}(q_3)\big]\big\}$$

where $T|_{A,B,C,D} = q_1 = A, B|C, D$, without loss of generality, and q_2, q_3 are the alternative topologies. We prove that this inequality holds in Theorem 2. □

To prove Theorem 2, we consider how gene copies descend from different lineages in the locus tree ℓ, called **locus lineages**, at different vertices of the species tree T, similar to Markin and Eulenstein [2021] and Hill et al. [2022]. Let l_X denote the set of locus lineages that exist at vertex X in T, and let i^x_X denote the locus lineage in l_X from which gene copy x descends. If there are multiple locus lineages in l_X, they must descend from duplication events above X, so going backward in time, they join at duplication events, eventually becoming one lineage.

Take Figure S1A, in which a locus tree ℓ evolves within a balanced species tree T, as an example. Let R denote the MRCA of A, B, C, D in T. There are two locus lineages at R colored black and blue. Gene copies $a_0, a_2, b_0, b_2, c_0, d_0$, which descend from the black locus lineage, correspond to the original locus, whereas gene copies a_1, b_1, c_1, d_1, which descend from the blue locus lineage, correspond to the duplicated locus. For any four gene copies a, b, c, d, we can consider the relationship among their locus lineages at R. As an example, the locus lineage relationship at R for a_0, b_2, c_1, d_1 is $i^{a_0}_R = i^{b_2}_R \neq i^{c_1}_R = i^{d_1}_R$, also denoted ab, cd. Locus lineage relationships, also called root lineage scenarios, are enumerated by Markin and Eulenstein [2021] and Hill et al. [2022] and then used to partition the computation of $N^g(\hat{T}|_{A,B|C,D})$ (Equation 1). As we will show, locus lineage relationships are also useful for classifying gene tree induced quartets as SQs and DQs.

We now analyze locus lineage relationships, moving down the species tree T. To begin, consider locus lineages from **duplication events above the MRCA of A, B, C, D, denoted R.**

Lemma 1. *Assume $\ell, g \sim DLCoal(T, \mathcal{X})$ are correctly tagged. For an arbitrary subset of four taxa $\{A, B, C, D\} \subseteq \mathcal{X}$, if gene copies a, b, c, d do <u>not</u> descend from the same locus lineage at R, then $\ell|_{a,b,c,d}$ and $g|_{a,b,c,d}$ are DQs.*

Proof. Let $l'_R = \{i^a_R, i^b_R, i^c_R, i^d_R\} \subseteq l_R$. If $|l'_R| > 1$, the locus lineages in l'_R must descend from duplication events above R. Therefore, the MRCA of a, b, c, d in ℓ is a duplication event above R, and $\ell|_{a,b,c,d}$ is a DQ. Now consider the MRCA of a, b, c, d in g, denoted u. If $|l'_R| > 1$, it is always possible to select one gene copy $x \in L(u.l) \cap \{a, b, c, d\}$ and one gene copy $y \in L(u.r) \cap \{a, b, c, d\}$ such that x, y descend from different locus lineages in l'_R Then, by our definition of correct tagging, u is a duplication vertex; it follows that $g|_{a,b,c,d}$ is a DQ. □

By Lemma 1, only gene copies that descend from the same locus lineage at R can be SQs, so we make this assumption going forward. Continuing down the species tree T, our analyses depend on the rooting of $T|_{A,B,C,D}$. Its rooted topology is either the balanced tree $((A, B), (C, D))$ or one of four pectinate trees; we assume $(((A, B), C), D)$, without loss of generality. Now consider locus lineages from **duplication events above the MRCA of A, B, C, denoted P, but below R for the pectinate tree.**

Lemma 2. *Assume $T|_{A,B,C,D} = (((A, B), C), D)$. If gene copies a, b, c, d descend from the same locus lineage at R but a, b, c do <u>not</u> descend from the same locus lineage at P, then $\ell|_{a,b,c,d}$ and $g|_{a,b,c,d}$ are DQs.*

Proof. Suppose gene copies a, b, c, d descend from the same locus lineage r at R. Locus lineage r instantaneously splits at R in a speciation event that (1) corresponds to the MRCA of a, b, c, d in ℓ and (2) forces $\ell|_{a,b,c,d}$ to be pectinate with outgroup d. For $\ell|_{a,b,c,d}$ to be a DQ, the MRCA of a, b, c must be a duplication vertex. Let $l'_P = \{i^a_P, i^b_P, i^c_P\} \subseteq l_P$. If $|l'_P| > 1$, the locus lineages in l'_P must descend from duplication events above P but below R, as they must join together prior to the speciation event at R. This gives us our result. Due to deep coalescence, we cannot make assumptions about the rooted topology of $g|_{a,b,c,d}$. There are three cases to consider (1) balanced, (2) pectinate with an outgroup from a, b, c, or (3) pectinate with outgroup d. For cases (1) and (2), consider the MRCA of a, b, c, d in g, denoted u. If $|l'_P| > 1$, it is always possible to select one gene copy $x \in L(u.l) \cap \{a, b, c\}$ and one gene copy $y \in L(u.r) \cap \{a, b, c\}$ such that x, y descend from different locus lineages in l'_P. Then, by our definition of correct tagging, u is a duplication vertex; it follows that $g|_{a,b,c,d}$ is a DQ for cases (1) and (2). The same logic holds in case (3) but setting u to be the MRCA of a, b, c in g. □

By Lemma 2, only gene copies that descend from the same locus lineage at P can be SQs for the pectinate tree, so we additionally make this assumption going forward. We are now ready to proceed with proving Theorem 2.

Theorem 2. *Assume $\ell, g \sim DLCoal(T, \mathcal{X})$ are correctly tagged. For an arbitrary subset of four species $A, B, C, D \subseteq \mathcal{X}$, the following inequality holds*

$$\mathbb{E}\left[N_{sq}^g(q_1)\right] > \max\left\{\mathbb{E}\left[N_{sq}^g(q_2)\right], \mathbb{E}\left[N_{sq}^g(q_3)\right]\right\} \tag{4}$$

where $T|_{A,B,C,D} = q_1 = A, B|C, D$, without loss of generality, and q_2, q_3 are the alternative topologies.

Proof. Our proof follows the structure of Theorem 4 in Hill et al. [2022]. To begin, we break our proof into two cases depending on whether $T|_{A,B,C,D}$ is balanced or pectinate.

Balanced species tree ((A,B),(C,D)). By Lemma 1, we have

$$\mathbb{E}\left[N_{sq}^g(T|_{A,B,C,D})\right] = \sum_{r \in l_R} \mathbb{E}_r\left[N_{sq}^g(T|_{A,B,C,D})\right] \tag{5}$$

where subscript r denotes conditioning the probability space and the expected values on having gene copies from A, B, C, D descending from the same locus lineage r at R. Locus lineage r instantaneously splits at R in a speciation event, which is the MRCA of a, b, c, d in ℓ, denoted R'. Therefore, we can also condition based on one of two events: (1) no coalescence among a, b, c, d below R', denoted $\mathcal{NC}$, and (2) its complement $\mathcal{NC}^C$.

Coalescence below R'. For event $\mathcal{NC}^C$, either a, b coalesce below R', c, d coalesce below R', or both. Regardless, $g|_{a,b,c,d}$ is isomorphic to $q_1 = A, B|C, D$, so $\mathbb{P}_r\left[N_{sq}^g(q_2) = 0|NC^C\right] = \mathbb{P}_r\left[N_{sq}^g(q_3) = 0|\mathcal{NC}^C\right] = 1$. Although the unrooted topology of $g|_{a,b,c,d}$ is fully determined by our conditioning, it could be either an SQ or a DQ, depending on g (because complex scenarios involving duplications and deep coalescence, even among gene copies outside a, b, c, d, can trigger duplication vertices across g; see Fig. S1). To ensure $\mathbb{E}_r\left[N_{sq}^g(q_1)|\mathcal{NC}^C\right] > 0$, consider that (1) there is a strictly positive probability that $g \sim MLC(\ell)$ is isomorphic to ℓ and (2) $\ell|_{a,b,c,d}$ must be an SQ by our assumption that all gene copies descend from r, which splits at R in a speciation event that corresponds to the MRCA of a, b, c, d and forces $\ell|_{a,b,c,d}$ to have rooted topology $((a, b), (c, d))$; also see Zhang et al. [2020]. This gives us $\mathbb{P}_r\left[N_{sq}^g(q_1) \geq 1|\mathcal{NC}^C\right] > 0$. Then, by Markov's Inequality, $\mathbb{E}_r\left[N_{sq}^g(q_1)|\mathcal{NC}^C\right] > 0$. Putting everything together, we have

$$\mathbb{E}_r\left[N_{sq}^g(q_1)|\mathcal{NC}^C\right] > 0 = \mathbb{E}_r\left[N_{sq}^g(q_2)|\mathcal{NC}^C\right] = \mathbb{E}_r\left[N_{sq}^g(q_3)|\mathcal{NC}^C\right]. \tag{6}$$

No coalescence below R'. For event $\mathcal{NC}$, gene copies a, b, c, d fail to coalesce and enter above R', so all three quartet topologies are possible and occur with equal probability, by exchangeability [Allman et al., 2011]. Additional constraints imposed by the bounded coalescent do not prevent the exchangeability of a, b, c, d labels, since we conditioned on them descending from the same locus lineage r, and therefore they have the same coalescent constraints above R'. Moreover, the three quartet topologies are SQs with equal probability, by exchangeability. Consider an arbitrary gene tree t that (1) induces SQ $a, b|c, d$ and (2) meets our conditions (i.e., a, b, c, d descend from r and fail to coalescence before R'). Then, swapping labels b, c (or a, c) in t produces a gene tree t' with the same probability as t under the MLC model, but t' induces SQ $a, c|b, d$ (or $a, d|b, c$) instead (and continuing swaps in this fashion). It follows that

$$\mathbb{E}_r\left[N_{sq}^g(q_1)|\mathcal{NC}\right] = \mathbb{E}_r\left[N_{sq}^g(q_2)|\mathcal{NC}\right] = \mathbb{E}_r\left[N_{sq}^g(q_3)|\mathcal{NC}\right]. \tag{7}$$

Summary. Combining Equations 5–7 gives Equation 4, completing our proof for the balanced species tree.

Pectinate Species Tree (((A,B),C),D). By Lemmas 1 and 2, we have

$$\mathbb{E}\left[N_{sq}^g(T|_{A,B,C,D})\right] = \sum_{r \in l_R} \sum_{p \in l_P} \mathbb{E}_{r,p}\left[N_{sq}^g(T|_{A,B,C,D})\right]$$

where subscript r, p denotes conditioning the probability space and expected values on having gene copies from A, B, C, D descending from the same locus lineage r at R with those from A, B, C descending from the same locus lineage p at P. Locus lineage p instantaneously splits at P in a speciation event, which is the MRCA of a, b, c in ℓ, denoted P'. The remainder of the proof is the same as for the balanced species tree but replacing the speciation event R' with P' and excluding d from leaf swaps because only a, b, c have an opportunity to coalesce above P' but below R'; see Allman et al., 2011 for details. $\qquad\square$

Lastly, it is worth noting that our proofs do not make use of the duplication rate λ or loss rate μ, so they hold for generic GDL only models, as discussed by Molloy and Warnow [2020].

3.3 TREE-QMC-pro

We have shown that an optimal solution to an exclusion-only version of A-pro's objective function is a consistent estimator of the species tree under the DLCoal model. To empirically evaluate this modified objective function, we exclude DQs within TREE-QMC [Han and Molloy, 2023]. TREE-QMC (or TQMC for brevity) is a heuristic for the same NP-hard optimization problem as ASTRAL but is instead based on the Quartet Max Quartet (QMC) framework of Snir and Rao [2010].

QMC reconstructs the species tree in a divide-and-conquer fashion. At each step in the divide phase, the goal is to partition the taxon set in half, producing an internal branch of the output species tree. Each (bi)partition is identified by constructing a graph in which vertices are the taxon set for the given subproblem and edges are weighted according to the input quartets. Each quartet $A, B|C, D$ contributes two bad edges (i.e., (A, B) and (C, D)) and four "good" edges (i.e., (A, C), (A, D), (B, C), and (B, D)). Cutting the bad edges is undesirable because it breaks apart siblings, whereas cutting the good edges is desirable; this motivates seeking a maximum cut; see Snir and Rao [2010].

The main contribution of TREE-QMC is that it constructs the quartet graph in an efficient manner directly from gene trees, without enumerating all quartets. The latest version enables quartets to be weighted based on gene tree branch lengths and support values to improve robustness to gene tree estimation error [Han and Molloy, 2025]. To build the weighted graph efficiently, TREE-QMC uses "auxiliary values" that store information about how taxa below any given node can be combined to produce good or bad edges. Auxiliary values are computed in either preorder or postorder traversals of the input gene tree; after, good/bad edges are computed in postorder traversals. Our contribution is modifying the recurrences for auxiliary values and good/bad edges to exclude contributions from DQs (see Section 1 of the Supplement for details and equations). To our knowledge, our new method, called TQMC-pro, is the first to enable both the exclusion of DQs and the weighting of quartets based on gene tree branch information. Including the agglomeration in this framework is challenging because homeomorphic SQs can have different weights.

Lastly, the other contribution of TREE-QMC is that it **normalizes** the quartet graph to account for "artificial taxa," which are introduced so that solutions on subproblems can be combined during the divide phase (this framework naturally accommodates multi-labeled trees as input). To normalize the graph efficiently, TQMC requires the number of leaves in a gene tree labeled by a taxon to be indicative of the number of quartets that taxon participates in. This relationship breaks down when excluding DQs. Therefore, TQMC-pro does not support graph normalization; however, it did not impact accuracy in our study.

4 Empirical Study

Next, we describe an empirical study evaluating the exclusion-only objective function implemented within TQMC-pro. Software versions and exact commands are available in the Supplement.

Simulation Study. To evaluate TQMC-pro, data were simulated under the DLCoal model with 25 taxa (and an outgroup) and 2000 gene trees using the same protocol as the A-pro study [Zhang et al., 2020]. Re-simulation was necessary to save the true duplication/loss events, which were needed to evaluate tagging accuracy. Full simulation parameters are provided in Table S1 of the Supplement and summarized below.

The first experiment varied the gene duplication rate as well as the effective population size, which in turn **varied the ILS level** (loss rate of zero). Data were simulated with four duplication levels (0, 1, 2, 5) and four ILS levels (0, 0.2, 0.5, 0.7), where duplication level was evaluated as the mean number of gene copies per species minus one and ILS level was evaluated as the mean RF distance between true locus and gene tree pairs for the dup=0 condition. In total, there were 16 model conditions, each with 25 replicates.

The second experiment varied **gene tree estimation error (GTEE)**. Molecular sequences of 500bp were simulated under the GTR model for the first 10 replicates of each condition described above. After, gene trees were estimated from the first 100bp and 500bp under the GTR model using IQ-TREE [Minh et al., 2020]. Gene tree estimation error (GTEE) was evaluated as the mean RF distance between the true and estimated gene trees. For 100bp and 500bp, GTEE was around 0.5 and 0.2, respectively.

The third experiment varied the duplication rate λ as well as **loss/duplication rate ratio** μ/λ (effective population size was set to the same value as the 0.5 ILS level). Data were simulated with four duplication levels (0, 1, 2, 5) and three ratios (0, 0.5, 1) (the simulator did not allow $\mu > \lambda$; see the Supplement for details). In total, there were 9 model conditions, each with 25 replicates.

Evaluation Metrics. The accuracy of gene tree tagging was evaluated for each pair of gene copies (i.e., leaves). Specifically, TPs are paralogs in both the true and A-pro taggings, FPs are orthologs in the true tagging and paralogs in the A-pro tagging, and FNs are paralogs in the true tagging but orthologs in the A-pro tagging. Precision and recall were computed for each model condition taking the first 100 gene trees of the 10 replicates. Species tree error was evaluated as the normalized RF distance between the true and estimated species trees; it corresponds to the fraction of FN/FP branches, as both trees are binary.

Methods. TQMC-pro was evaluated against A-pro and standard quartet-based species tree methods: A-multi and TQMC, which is similar to A-multi. All methods were executed given the first 250, 500, 1000, or 2000 gene trees per replicate data set. We ran TQMC in default mode, which uses n2 graph normalization, as well as without normalization (n0) to match TQMC-pro. For each data set, the true duplication events were used to tag the true rooted gene trees based on our definition of correct tagging. Additionally, true and estimated gene trees were rooted and tagged using A-pro. These two sets of trees were given as inputs to TQMC-pro. To our knowledge, A-pro does not take tagged gene trees as input and thus performed rooting and tagging internally.

Plant re-analysis. Lastly, we re-analyzed the 1kp plant data set [Wickett et al., 2014] with 83 taxa and 9,237 gene family trees using A-pro, TQMC-pro (with A-pro tagged gene trees), and A-multi. Trees were compared based on their quartet scores computed with both A-pro and A-multi. We also evaluated differences against the ASTRAL tree (80 taxa) estimated from single-copy genes, reporting support (local posterior probability) computed with either A-pro or A-multi on differing branches.

5   Empirical Results and Discussion

We now describe the results of simulations and plant re-analysis.

Varying Duplication and ILS Levels We first looked at interactions between varying the duplication and ILS levels, holding the loss rate at 0. The number of duplication vertices in the gene trees increases with ILS level, jumping from $\sim$23 to $\sim$31 for the highest duplication level (5), as suggested by our example (Fig. S1). Tagging accuracy increased with duplication level (although so did the number of pairwise comparisons) but decreased with ILS level. Nevertheless, precision was greater than 0.75 and recall was greater than 0.8, across all model conditions (Fig. S2A). Perhaps as a result, there was little to no difference in species tree error between TQMC-pro given the true tagging or A-pro's rooting/tagging (Fig. S2B). Likewise, TQMC-pro and A-pro achieved similar error rates across all model conditions and outperformed the "multi" methods, with differences most pronounced for small numbers of genes, higher duplication levels, and high ILS levels. The performance of "pro" methods tended to improve with increasing duplication level, whereas the performance of "multi" methods deteriorated. As an example, at the highest ILS level (0.7) and smallest number of genes (250), the pro method error *decreased* from $\sim$0.1 to <0.05 when going from duplication level 0 to 5, whereas the multi method error *increased* from $\sim$0.1 to >0.15 (A-multi and TQMC-n0) or stayed the same (TQMC-n2). One possible explanation is that duplications produce more conflicting quartets for the multi methods but not the pro methods (additionally, quartets impacted by deep coalescence are more likely to be duplication quartets). It is also worth noting that the ILS level decreased from 0.71 for dup=0 to 0.62 for dup=5 in this example, perhaps due to the added constraints of the bounded coalescent, so the model condition became less challenging (Table S2).

Varying GTEE Next, we evaluated the impact of gene tree estimation error. Tagging precision decreased as GTEE increased, whereas recall stayed largely the same. This suggests an increase in FPs (Fig. S2A,C,D), and relatedly, the number of vertices tagged as duplications by A-pro increased with GTEE (Table S2). Likewise, species tree error typically increased as GTEE increased, when holding model conditions constant, especially for A-multi and TQMC-n0 (Figs. S2E and S1), although there were exceptions at the highest ILS levels (Figs. S2–S3). In the context of GTEE, increasing the duplication level sometimes decreased species tree error, even for the multi-methods (especially for duplication levels 1-2 and small numbers of gene trees). One possible explanation is that the greater number of quartets available from duplications has an advantage when there is noise.

Figure S2: **A, B, and C** Precision and recall of A-Pro's tagging algorithm on true gene trees and gene trees estimated 500bp and 100bp, respectively. **D)** Species tree error varying duplication and ILS levels (loss rate=0). **E)** Species tree error varying GTEE (ILS level=0.2). Dots are averages over 25 replicates for true gene trees and 10 replicates for estimated gene trees.

Varying Loss/Duplication Rate Ratio   Next, we evaluated the impact of varying the loss/duplication rate ratio, holding the effective population size constant (ILS level of 0.5). Losses had little impact on either tagging accuracy (Fig. S4) or species tree error (Fig. S5). However, increasing the loss rate also lowered the amount of ILS, especially as the duplication rate increased (Table S3), likely related to the bounded coalescent.

Plant Re-analysis   Lastly, we re-analyzed the 1kp plant data set. Both ASTRAL-pro and TQMC-pro produced similar trees to the published ASTRAL tree on single-copy genes, differing by just 5 and 4 branches out of 77, respectively, and recovering all major clades (Fig. S3 and Fig. S6). They differed from each other by just 4 branches, often with lower branch support. In contrast, A-multi differed from the ASTRAL tree by 58 branches and failed to recover the major clades. The inter-mixing of these clades often led to branch support values of 0. To rule out the possibility that A-multi had just failed to find a good solution to its optimization problem, we computed the A-multi score for both the A-pro and TQMC-pro trees. We found that the A-multi tree was indeed the highest scoring, suggesting that the problem is due to its objective function not excluding duplication quartets.

6   Conclusions

In this work, we presented a definition of correct gene tree tagging and used it to provide the first proof that A-pro is statistically consistent under the DLCoal model for an exclusion-only version of its objective function. Our proof does not consider SQ agglomeration; however, we show empirically that DQ exclusion within TQMC (called TQMC-pro) achieves similar performance to A-pro on real and

Figure S3: Species trees estimated by A-Pro, TQMC-pro, and A-Multi on the multi-copy 1kp plant data set. All trees were rooted at $(Uronema, (Volvox, Chlamydomonas))$, not shown. Dashed lines indicate the species was missing from the single-copy ASTRAL tree. The A-pro tree and TQMC-pro tree differed from the single-copy ASTRAL tree by 5 and 4 branches, respectively, shown in red; they differed from each other by four branches; shown in blue. Branch support (range 0 to 1) computed with A-pro is indicated on these branches as well as on the clades for Monocots and Eudicots+Magnolids, which are split in the A-multi tree. Branch support computed by A-multi is shown for select branches where these clades intermix. Quartet scores are shown at the bottom.

simulated data sets. Our empirical results also highlight that excluding DQs may lead to better sample complexity than A-multi. Currently, there are no sample complexity results for these methods, and we expect formal proofs to be intricate based on related results under the DLCoal model [Hill et al., 2022]. Our proof also assumes correct rooting and tagging; it may be possible to relax these assumptions, as rooting/tagging with A-pro's algorithm yielded good empirical results, both in terms of tagging accuracy and species tree accuracy. Relatedly, we evaluated tagging accuracy in a pairwise fashion, as it was computationally efficient and applicable to gene trees with estimation errors; however, pairwise errors do not necessarily translate to errors in the DQ/SQ classification (as only a subset of pairwise relationships need to be inferred correctly to reach the correct classification). Lastly, it would be interesting to extend this work to more complex unified models of duplication-loss-coalescence, such as the MLMSC [Li et al., 2020].

7   Competing interests

No competing interests to declare.

8   Author contributions statement

RP, PD, YL, EKM developed the definition of correct gene tree tagging. RP, AM, EKM proved statistical consistency. RP designed TREE-QMC-pro and implemented it. RP, YL, EKM designed the experiments. YL, RP generated the results. All authors analyzed the results. RP, EKM drafted the manuscript. All authors reviewed and edited the manuscript. EKM conceptualized and supervised the study.

9   Acknowledgments

The authors thank Yunheng Han for helpful discussions about the TREE-QMC code base. This material is based upon work supported by the U. S. National Science Foundation (NSF) under Grant Number 2441458 (to EKM) as well as the State of Maryland. AM was supported in part by the USDA-ARS (ARS project number 5030-32000-231-000D) and USDA-APHIS (ARS project number 5030-32000-231-111-I). This work was initiated "Theory, Methods, and Applications of Quantitative Phylogenomics" program held at the NSF-funded Institute for Computational and Experimental Research in Mathematics. The funders had no role in study design, data collection, and interpretation, or the decision to submit the work for publication. Mention of trade names or commercial products in this article is solely for the purpose of providing specific information and does not imply recommendation or endorsement by the USDA. USDA is an equal opportunity provider and employer.

References

E. S. Allman, J. H. Degnan, and J. A. Rhodes. Identifying the rooted species tree from the distribution of unrooted gene trees under the coalescent. *J Math Biol*, 62(6):833–862, 2011.

L. Arvestad, J. Lagergren, and B. Sennblad. The Gene Evolution Model and Computing Its Associated Probabilities. *J ACM*, 56(2):7, 2009.

S. V. Edwards. Is a new and general theory of molecular systematics emerging? *Evolution*, 63(1):1–19, 2009.

S. Feng et al. Dense sampling of bird diversity increases power of comparative genomics. *Nature*, 587 (7833):252–257, 2020.

Y. Han and E. K. Molloy. Improving quartet graph construction for scalable and accurate species tree estimation from gene trees. *Genome Res*, 33(7):1042–1105, 2023.

Y. Han and E. K. Molloy. Improved robustness to gene tree incompleteness, estimation errors, and systematic homology errors with weighted TREE-QMC. *Syst Biol*, 74(5):758–777, 2025.

M. Hill, B. Legried, and S. Roch. Species tree estimation under joint modeling of coalescence and duplication: Sample complexity of quartet methods. *The Annals of Applied Probability*, 32, 2022.

M. Lafond and C. Scornavacca. On the weighted quartet consensus problem. *Theoretical Computer Science*, 769:1–17, 2019.

J. H. Leebens-Mack et al. One thousand plant transcriptomes and the phylogenomics of green plants. *Nature*, 574(7780):679–685, 2019.

B. Legried, E. K. Molloy, T. Warnow, and S. Roch. Polynomial-time statistical estimation of species trees under gene duplication and loss. *J Comput Biol*, 28(5):452–468, 2021.

Q. Li, C. Scornavacca, N. Galtier, and Y.-B. Chan. The multilocus multispecies coalescent: A flexible new model of gene family evolution. *Systematic Biol*, 70(4):822–837, 2020.

W. P. Maddison. Gene trees in species trees. *Syst Biol*, 46(3):523–536, 1997.

S. Majidian et al. Quest for Orthologs in the era of data deluge and ai: Challenges and innovations in orthology prediction and data integration. *J Mol Evol*, 93(6):702–719, 2025.

A. Markin and O. Eulenstein. Quartet-based inference is statistically consistent under the unified duplication-loss-coalescence model. *Bioinformatics*, 37(22):4064–4074, 2021.

B. Q. Minh, H. A. Schmidt, O. Chernomor, D. Schrempf, M. D. Woodhams, A. von Haeseler, and R. Lanfear. IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference in the Genomic Era. *Molecular Biology and Evolution*, 37(5):1530–1534, 2020.

S. Mirarab, R. Reaz, M. S. Bayzid, T. Zimmermann, M. S. Swenson, and T. Warnow. ASTRAL: genome-scale coalescent-based species tree estimation. *Bioinformatics*, 30(17):i541–i548, 2014.

E. K. Molloy and T. Warnow. FastMulRFS: fast and accurate species tree estimation under generic gene duplication and loss models. *Bioinformatics*, 36(Supplement_1):i57–i65, 2020.

P. Pamilo and M. Nei. Relationships between gene trees and species trees. *Mol Biol Evol*, 5(5):568–583, 1988.

R. A. Parsons and M. S. Bansal. DupLoss-2: Improved phylogenomic species tree inference under gene duplication and loss. *Syst Biol*, page syaf073, 2025.

M. Rabiee, E. Sayyari, and S. Mirarab. Multi-allele species reconstruction using ASTRAL. *Mol Phylogenet Evol*, 130:286–296, 2019.

B. Rannala and Z. Yang. Bayes estimation of species divergence times and ancestral population sizes using DNA sequences from multiple loci. *Genetics*, 164(4):1645–1656, 2003.

M. D. Rasmussen and M. Kellis. Unified modeling of gene duplication, loss, and coalescence using a locus tree. *Genome Res*, 22(4):755–765, 2012.

S. Snir and S. Rao. Quartets MaxCut: A divide and conquer quartets algorithm. *IEEE/ACM Trans Comput Biol Bioinform*, 7(4):704–718, 2010.

N. J. Wickett et al. Phylotranscriptomic analysis of the origin and early diversification of land plants. *Proc Natl Acad Sci USA*, 111(45):E4859–E4868, 2014.

C. Zhang, C. Scornavacca, E. K. Molloy, and S. Mirarab. ASTRAL-Pro: Quartet-based species-tree inference despite paralogy. *Mol Biol Evol*, 37(11):3292–3307, 2020.