Title: Mutualistic rhizobia harbor genetic variation for traits related to parasite infection

Teaser text

Nutritional mutualisms are everywhere. These mutualisms are defined by the exchange of

resources between partners, but many have secondary effects on resistance to parasites,

pathogens, or herbivores. Here, we show that a textbook nutritional mutualist (nitrogen-fixing

rhizobia) harbors genetic variation for the host's response to parasite infection. Our results imply

that nutritional mutualists can impact the evolutionary potential of infection-related traits in their

hosts, suggesting that these mutualisms may be overlooked drivers of defense evolution.

Abstract

Nutritional mutualisms are defined by resource exchange, but growing evidence suggests that

these interactions also shape resistance to parasites and pathogens. The evolutionary significance

of this phenomenon is unclear, primarily because research has focused on mutualist-mediated

plasticity rather than on mutualist contributions to heritable variation in the outcome of infection.

To test whether nutritional mutualists harbor genetic variation for infection-related traits, we

performed a quantitative genetic experiment in the nutritional mutualism between legumes and

nitrogen-fixing rhizobia. We paired 10 mutualistic *Sinorhizobium meliloti* rhizobia strains with

20 *Medicago truncatula* plant genotypes in an incomplete factorial design, and experimentally

infected plants with parasitic root-knot nematodes (*Meloidogyne hapla*). We used this design to

estimate plant and rhizobia contributions to genetic variation in four infection-related traits:

resistance, tolerance, virulence, and mutualism robustness to infection. Rhizobia contributed

directly to genetic variation in virulence, and to parasite resistance and mutualism robustness via

genotype-by-genotype interactions. Rhizobia did not contribute to genetic variation in tolerance.

The effect of rhizobia on parasite resistance was partially explained by their effect on root

growth. Our results raise the possibility that some nutritional mutualists play a role comparable

to defensive mutualists in shaping the evolutionary potential of host defense traits.

Keywords

Genetic variation, parasitism, mutualism, host-microbe, root-knot nematode, symbiosis

Introduction

Mutualisms are traditionally categorized by the benefits they provide. Nutritional mutualists

exchange resources, defensive mutualists provide protection, and transportation mutualists

facilitate movement (Bronstein, 2015). Yet mutualist impacts on their hosts can extend well

beyond the primary benefit of the interaction (Batstone et al., 2022; Friesen et al., 2011; Fronk &

Sachs, 2022; Ma et al., 2020; O'Brien et al., 2021). Nutritional mutualisms, for example,

frequently affect resistance to parasites, pathogens, and herbivores (Buxton-Martin et al., 2025;

Koch & Schmid-Hempel, 2011; O'Keeffe et al., 2022). This observation indicates that effects on

interactions with enemies are not limited to defensive mutualisms, and raises an important

question: do non-defensive mutualisms also influence the evolution of defense traits?

There are now many examples of how nutritional mutualisms affect their host's

resistance to parasites, pathogens, or herbivores (reviewed in Wood et al., *in review*). This

phenomenon is especially well studied in mycorrhizal fungi, which usually increase the host

plant's resistance to infection (Jung et al., 2012; Liu et al., 2023), but can decrease resistance as

well (Eck et al., 2022; Miozzi et al., 2019). Nitrogen-fixing rhizobia increase their host plant's

vulnerability to a parasitic nematode (Buxton-Martin et al., 2025; Wood et al., 2018), as well as

to herbivores (Simonsen & Stinchcombe, 2014). Similar effects on interactions with enemies

have been documented in animals as well (Salem et al., 2015; Stevens et al., 2021). From these

examples, it is clear that nutritional mutualists can have a profound impact on their host's

interactions with enemies. Furthermore, unlike defensive mutualists, nutritional mutualists are

not always protective, implying that nutritional and defensive mutualisms may have markedly

different consequences for the evolution of host traits related to defense.

In defensive mutualisms, symbionts contribute directly to the evolution of defense via

heritable variation in host traits that resides in symbiont populations (reviewed in Vorburger &

Perlman, 2018). In other words, defense is jointly determined by host and symbiont genomes.

For example, *Drosophila* fruit flies inoculated with different *Wolbachia* strains differ in

resistance to viral infection, indicating that *Wolbachia* harbors genetic variation for antiviral

resistance (Martinez et al., 2014, 2017). In a landmark study, (Oliver et al., 2005) found that

parasite resistance was explained by the genotype of the *Hamiltonella defensa* symbiont, not the

aphid host. Subsequent studies, primarily in pea aphids (*Acyrthosiphon pisum*), established that

differences in protection among defensive symbiont strains are common (Hafer-Hahmann &

Vorburger, 2020; McLean et al., 2020; Parker et al., 2017; Rouchet & Vorburger, 2014; Schmid

et al., 2012). In parallel, experimental (co)evolution, primarily in *Caenorhabditis elegans*

nematodes, has directly linked defensive symbionts to the evolution of resistance and tolerance,

as well as pathogen virulence (Ford et al., 2016, 2017; Hoang et al., 2024; King & Bonsall, 2017;

Rafaluk-Mohr et al., 2022; Smith et al., 2025).

By contrast, the evolutionary significance of *non-defensive* mutualist effects on

interactions with enemies remains largely untested. This is primarily because research in non-

defensive mutualisms has emphasized mutualist-mediated *plasticity* over mutualist effects on

heritable variation in infection-related traits. A few studies have reported differences in enemy

interactions between hosts inoculated with different nutritional mutualist strains (Buxton-Martin

et al., 2025; Simonsen & Stinchcombe, 2014). For example, *Medicago* plants inoculated with

two different rhizobia strains differed in resistance to parasitic nematodes, as well as in their

transcriptomic response to nematode infection (Buxton-Martin et al., 2025). However, two-strain

experiments like this cannot directly estimate the mutualist's contribution to genetic variation in

infection-related traits, because genetic variance is a population-level property that requires

many strains to measure accurately. Furthermore, benchmarking mutualist genetic variation

against host genetic variation is essential to gauge the mutualist's relative contribution to the

evolutionary potential of a host trait. We're aware of only one study in any mutualism that

directly compared host and mutualist contributions to genetic variance in infection-related traits

(Parker et al., 2017), in an aphid defensive symbiosis.

Here, we tested whether nutritional mutualists influence the evolutionary potential of host

traits related to parasite infection. We asked three questions: First, do nutritional mutualists

harbor genetic variation for infection-related host traits? Second, how much genetic variation do

mutualists contribute to these traits, relative to hosts? Finally, do nutritional mutualists contribute

equally to parasite resistance, tolerance, and virulence? We addressed these questions in the

tripartite interaction between legumes, nitrogen-fixing rhizobia, and plant-parasitic root knot

nematodes. *M. truncatula* is an annual legume native to the Mediterranean (Cook, 1999) and

forms a mutualistic symbiosis with nitrogen-fixing rhizobia, such as *S. meliloti*. Rhizobia are

housed within nodules that develop from the roots of their host plant where they trade with the

host plant for carbon-containing compounds. The northern root-knot nematode (*Meloiogyne*

hapla) is an obligate endoparasite that forms galls on plant roots, where it siphons nutrients from

the host (M. G. K. Jones & Goto, 2011; Koltai et al., 2001). A previous study found that *M.*

truncatula plants inoculated with two different *S. meliloti* strains differed in their resistance to

nematode infection (Buxton-Martin et al., 2025).

We paired 20 *Medicago truncatula* plant accessions and 10 *Sinorhizobium meliloti*

rhizobia strains in an incomplete factorial design, and infected half of the plants with parasitic

nematodes (Figure 1A). We then partitioned total genetic variation in four infection-related traits

(described below) into genetic variation in the host (direct genetic variance; G_{Host}), genetic

variation in the mutualism (indirect genetic variance; $G_{Rhizobia}$), and genotype-by-genotype

interactions between host and mutualist (intergenomic epistasis; $G_{Host \times Rhizobia}$) (De Lisle et al.,

2022; Heath & Tiffin, 2007; Wade, 2007). Genotype-by-genotype interactions are pervasive in

both nutritional and defensive mutualisms (Heath, 2010; Heath & Tiffin, 2007; Parker et al.,

2017; Schmid et al., 2012), so it is crucial to account for genotype dependence when estimating

genetic variation in joint host-symbiont traits.

We focused on four infection-related traits: resistance, tolerance, virulence, and

mutualism robustness to infection (Figure 1B) (Bush et al., 1997). Resistance (the inverse of

susceptibility) is the host's ability to prevent parasite infection, and is measured by infection

intensity (the total number of parasites on a host). Tolerance is the fitness cost per parasite,

measured as the slope of the relationship between infection intensity and fitness. Virulence is the

total fitness cost of infection, as measured as the difference in fitness between infected and

uninfected hosts. All three of these traits are typically determined jointly by both the host's and

parasite's genotype and differ in their implications for host-parasite interactions (Bruns et al.,

2012; Jutzeler et al., 2024; Salvaudon et al., 2007). In addition to these three classic infection-

related traits, we also measured mutualism robustness to parasite infection, defined as the impact

of parasite infection on mutualist colonization (Figure 1B). This captures an often-overlooked

cost of parasite infection: disrupted interactions with mutualists. Finally, we evaluated whether

the rhizobial effect on infection intensity was related to their effect on host growth (i.e.,

nutritional mutualists increase plant size, and bigger plants host more parasites).

We found that rhizobia contributed directly to genetic variation in parasite virulence

($G_{Rhizobia}$), and to parasite resistance and mutualism robustness via genotype-by-genotype

interactions ($G_{Host \times Rhizobia}$). A path analysis revealed that rhizobia primarily influence resistance

indirectly via their effect on plant size, raising the possibility that mutualist effects on resistance

are common in growth-promoting resource mutualisms. Our results demonstrate that non-

defensive mutualists can influence the evolutionary potential of infection-related traits in their

hosts. However, if the genotype-by-genotype interactions for infection are common, as we found

in this study, the contribution of nutritional mutualisms to heritable variation in infection-related

traits will depend on how hosts and mutualists pair up in wild populations.

Methods

Study system

Medicago truncatula Gaertn. is an annual legume native to the Mediterranean (Cook, 1999). *M.*

truncatula plants form association with nitrogen-fixing rhizobia in the soil in the genus

Sinorhizobium, through the successful exchange of compatible chemical cues and the migration

of rhizobia into root tissues. Rhizobia are then housed within nodules that develop from the roots

of their host plant. Within these nodules rhizobia fix nitrogen that is traded with the host plant for

carbon-containing compounds. The parasitic root knot nematode *Meloidogyne hapla* is an

obligate endoparasite with an extremely large host range (Opperman et al., 2008). Female *M.*

hapla burrow through intracellular spaces of host roots until they reach a suitable infection site in

the vascular cylinder. They then pierce plant cells with their stylet mouthpiece and inject effector

proteins that suppress host immune responses and induce cell differentiation producing giant

cells from which the nematode siphons nutrients. A gall forms around these infection sites (M.

G. K. Jones & Goto, 2011). Previous studies showed that plant maternal families (hereafter,

genotypes) and plants inoculated with different rhizobia strains differ in resistance to this

parasite (Buxton-Martin et al., 2025; Wood et al., 2018).

We chose 20 accessions of *M. truncatula* from the French National Institute for

Agriculture Research (INRAE) germplasm library. We haphazardly chose these accessions to

span the range of standing genetic variation in *M. truncatula*, including representative diversity

across structured populations as well as resistance to nematode infection and nodulation with

rhizobia reported in Wood *et al.* (2018) (see Supplemental for details). We chose 10 strains of *S.*

meliloti that were isolated from *M. truncatula* (STable 2). Using FASTANI (Jain et al., 2018),

these strains showed a minimum average nucleotide identity of no less than 98% (SFigure 3),

indicating all strains belong to a single species. The sources of all *S. meliloti* strains can be found

in the Supplementary Materials. We sourced *M. hapla* line LM from the Department of

Entomology and Plant Pathology at North Carolina State University, which was originally

collected in France (Chen & Roberts, 2003).

Experimental design and data collection

Following the protocol of Garcia *et al.* (2006), we grew *M. truncatula* in 66 mL Cone-tainer pots

(Stuewe & Sons, Inc., Tangent, OR, USA) within a growth chamber at the University of

Pennsylvania. Plants were assigned to treatments such that plant genotypes encountered 2-4

rhizobia strains, and each rhizobia strain encountered 5 or 6 host genotypes (Figure 1A). We

replicated this experimental design across nematode-infected and nematode-uninfected

treatments. Plants were inoculated with nematodes and rhizobia 14 days after planting. Each

treatment (plant genotype-rhizobia strain combination) initially included 7-8 replicates, to ensure

that at least 4-5 replicates survived to harvesting. See the Supplementary Materials for details

regarding inoculation, growth conditions, and sample preparation.

To prevent rhizobia or nematode contamination, each rack of plants contained a single

rhizobia treatment and nematode treatment. We distributed rhizobia strain and nematode

treatments across different shelves and within the growth chamber in an incomplete block design

and randomized plant genotypes within racks. We included shelf location within growth

chambers as a fixed-effect blocking variable in our statistical models.

Plants were harvested 52-56 days after planting. Approximately 5-40 nodules were

dissected from plant roots during harvesting for a separate experiment and the number of nodules

removed was recorded. To harvest, roots were washed in water, and roots and shoots were

separated with scissors. Shoots were dried for a minimum of 48 hours in a drying oven at 60 °C

and root systems were frozen in -20 °C for later processing.

After drying, shoots were weighed to the nearest tenth of a milligram. To quantify

parasitic nematode infection, we dyed the root systems of nematode-infected plants with red food

coloring following the methods outlined in Kandouh *et al.* (2019), which stains mature nematode

egg sacs. Roots were scanned on an Epson Perfection V850 Pro scanner at 1200 dpi with high

contrast background correction following methods outlined in York (2023). Nodules, galls, and

nematode egg sacs were manually counted from these images using ImageJ (Schneider et al.,

2012). After scanning, roots were dried for >48 hours in a drying oven and weighed to the

nearest 0.0001 gram.

Variance component analyses: Quantifying rhizobia contributions to genetic variation in

infection-related traits

To estimate the contribution of rhizobia strains to variation in resistance, virulence,

tolerance, and mutualism robustness, we used variance component analyses (Figure 1B). We fit

two models for each trait except virulence: one with (Figures 2, 4, 5) and one without (SFigures

5-7) root biomass as a covariate, to capture rhizobia strain effects on infection intensity measured

as parasite density (i.e. number of galls per root biomass) and total infection intensity (i.e.

number of galls per individual plant). We primarily present models with parasite density but

include results that do not correct for root biomass in the Supplementary Materials (SFigures 5-

7). Results were qualitatively similar for most analyses.

To test whether rhizobia strains contributed to variation in resistance to nematode

infection, we fit mixed-effects models with number of galls as the response variable and rhizobia

strain, host genotype, and the genotype-by-strain interaction as random effects. This analysis

only included nematode-infected plants. In this analysis, a significant effect of rhizobia strain or

significant genotype-by-strain interaction indicates that rhizobia contribute to genetic variation in

resistance via indirect genetic effects ($G_{Rhizobia}$) or intergenomic epistasis ($G_{Host \times Rhizobia}$)

respectively. Experimental block was included as a fixed effect, and we fit a negative binomial

error distribution (Lindén & Mäntyniemi, 2011).

Virulence is measured as the difference in fitness between infected and uninfected hosts.

To test whether rhizobia strains contributed to variation in virulence, we fit a mixed-effect model

with shoot biomass, our host fitness proxy, as the response variable and rhizobia strain, host

genotype, genotype-by-strain, and all three of these effects interacting with nematode infection

status as random effects. Experimental block and nematode infection status were included as

fixed effects. In this analysis, a significant effect of infection-by-strain indicates rhizobia strain

effects on virulence and a significant effect of infection-by-strain-by-genotype indicates that

rhizobia contribute to virulence via indirect genetic effects ($G_{Rhizobia}$) or intergenomic epistasis

($G_{Host \times Rhizobia}$) respectively.

Tolerance is the relationship between infection intensity and host fitness. To test whether

rhizobia strains contributed to variation in tolerance, we fit mixed-effects models with shoot

biomass as the response variable and rhizobia strain, host genotype, and both of these effects

interacting with the number of galls as random effects. Unlike our other analyses, because each

genotype-by-strain combination had 4-8 replicates, we did not have the statistical power to

confidently estimate slopes for specific host genotype-rhizobia strain combinations and therefore

did not include genotype-by-strain or infection-by-genotype-by-strain effects in our model. For

random effects interacting with the number of galls, we included random slopes and no random

intercept for the number of galls. We included the number of galls and experimental block as

fixed effects. In this analysis, a significant interaction between number of galls and rhizobia

strain indicates that rhizobia strain contributes to genetic variation in tolerance via indirect

genetic effects ($G_{Rhizobia}$).

We defined mutualism robustness as the difference in nodulation between infected and

uninfected hosts. To test whether rhizobia strains contributed to variation in mutualism

robustness, we fit mixed-effects models with the number of nodules as the response variable and

rhizobia strain, host genotype, genotype-by-strain, and all three of these effects interacting with

nematode infection status as random effects. Experimental block and nematode infection status

were included as fixed effects. In this analysis, a significant effect of infection-by-strain or

infection-by-strain-by-genotype indicates rhizobia strain effects on mutualism robustness via

indirect genetic effects ($G_{Rhizobia}$) or intergenomic epistasis ($G_{Host \times Rhizobia}$) respectively. We used

a negative binomial distribution error (Lindén & Mäntyniemi, 2011).

We implemented our data analysis in R. We fit mixed-effects models using the

glmmTMB package (Brooks et al., 2017), and used the Sattherthwaite's degrees of freedom

method implemented in the lmerTest package (Kuznetsova et al., 2017) to estimate p-values and

infer statistical significance. We used the DHARMa package (Florian, 2016) to check that model

residuals were normally distributed and homoscedastic. We tested significance of random effects

with likelihood ratio tests. We used the MuMIn package (Bartoń, 2025) to calculate the

conditional and marginal R^2 values for each model and used the trigamma method when possible

per the package author's suggestion.

Path analysis: Distinguishing direct and indirect effects of rhizobia on infection intensity

Resource mutualists like rhizobia could affect the intensity of parasite infection (i.e., the number

of parasites on a host) directly, or indirectly via their effects on host growth, which impacts the

amount of host tissue available to infect. We therefore used path analysis to ask whether the

effect of rhizobia on resistance was mediated by root biomass. Our path analysis used the

following models:

Outcome model: Number of galls ~ Root biomass + Host genotype + Rhizobia strain

Mediating model: Root biomass ~ Host genotype + Rhizobia strain

To test for significant effects in the overall path analysis, we used a negative binomial GLM

implemented with the glm.nb function from the MASS package in R (Venables & Ripley, 2002)

to fit our outcome models that had gall number as the response variable (Lindén & Mäntyniemi,

2011) and a Gaussian linear model implemented in the lm function from the STAT package

(Bolar, 2019) to fit the mediating model. To estimate standardized path coefficients we used a

Gaussian distribution for both models fit with the lm function. We used the piecewiseSEM

package (Lefcheck et al., 2025) to test and structure our path analysis. We used sum-to-zero

contrasts in all our models. To test for the significance of direct and indirect effects in the per

strain path analyses, we used the boot function in the boot package (Canty & Ripley, 2025) to

perform bootstrapping analysis with 10,000 resamples to acquire confidence intervals for the

product of path coefficients and the proportion of meditation using the Gaussian models.

Results

Resistance

Our resistance model (Figure 2) explained 41.4% of the total variation in the number of galls

(R^2_C). Of the total variation in the number of galls, 10.28% was explained by fixed effects (R^2_M)

leaving 31.12% explained by our random effects. Of the variation in the number of galls

explained by random effects, our variance component analysis attributed 86% to host genotype

($p < 0.0001$), 14% to the genotype-by-strain interaction ($p = 0.0610$), and no significant variation

to rhizobia strain ($p = 0.1730$). These results indicate that host genotype was the largest

determinant of resistance but that genotype-by-strain interaction does contribute a significant

variation to resistance. Our variance component analysis that did not correct for root biomass

indicated that only host genotype contributed to variation in gall numbers.

Virulence

Our virulence model (Figure 3) explained 92.6% of the total variance in shoot biomass (host

fitness) (R^2_C). Of the total variance in shoot biomass, 27.51% was explained by fixed effects

(R^2_M) leaving approximately 65.09% explained by our random effects. Of the variation in shoot

biomass explained by random effects, 5% was attributed to the infection-by-strain interaction (p

$= 0.048$). The infection-by-genotype interaction and infection-by-genotype-by-rhizoba-strain

interaction explained 3% and 8% of the variance, respectively, but neither was statistically

significant ($p = 0.133$ and $p = 0.222$ respectively). These results indicate that rhizobia strains

contribute to variation in the fitness consequences of nematode infection, but that their effect

depends on the genotype of their host plant.

Tolerance

Our tolerance model that corrected for root biomass (Figure 4) explained 77.18% of the total

variation in shoot biomass (host fitness) (R^2_C). Of the total variation in shoot biomass, 69.57%

was explained by fixed effects (R^2_M) leaving approximately 7.61% explained by our random

effects. Of the variation in shoot biomass explained by random effects, our variance component

analysis attributed no variance to effects interacting with nematode infection severity (number of

galls). Instead, all variation in host fitness was attributed to host genotype (100%, $p < 0.0001$).

Qualitatively similar results were found from our tolerance model that did not correct for root

biomass ($R^2_C = 0.2253$; $R^2_M = 0.0697$). These results indicate that rhizobia strains do not

contribute to variation in tolerance.

Mutualist robustness

Our mutualism robustness model that corrected for root biomass (Figure 5) explained 46.86% of

the total variation in nodule counts (R^2_M). Of the total variation in nodule counts, 11.71% was

explained by fixed effects (R^2_C) leaving approximately 35.15% explained by random effects. Of

the variation in nodule counts explained by random effects, 12% was attributed to the infection-

by-genotype interaction ($p = < 0.0001$) and 16% to the infection-by-genotype-by-strain

interaction ($p = 0.0480$). No variance was attributed to the infection-by-strain interaction ($p =$

0.5409). Our mutualism robustness model that did not correct for root biomass showed

qualitatively similar results. These results indicate that individual host genotype and rhizobia

strain combinations contribute to variation in mutualism robustness to parasite infection, but that

the rhizobia effect depends on the genotype of its host.

Rhizobia effects on nematode infection intensity are primarily mediated by their effect on plant

size

In our path analysis (Figure 6), we found that rhizobia strains varied in their influence on root

biomass (p<0.0001) but not on gall numbers (p=0.1381) (Figure 2). Gall numbers increased with

root biomass (β=0.3151, p<0.0001). These results indicate that after correcting for the

differences in host genotype, larger roots are associated with greater number of galls, and

rhizobia differ in their effect on root growth. These path analysis results are consistent with the

difference between variance component analyses that did (Figure 2) and did not (SFigure 5)

correct for root biomass and indicate that their effect on plant growth is one mechanism through

which rhizobia strains influence resistance to nematodes.

Discussion

Here we report that nitrogen-fixing rhizobia contribute to genetic variation in infection-related

traits in their host plant. These effects manifested as differences in parasite virulence among

plants inoculated with different rhizobia strains (i.e., $G_{Rhizobia}$), as well as intergenomic epistasis

for resistance (i.e., $G_{Host \times Rhizobia}$) (Table 1). To our surprise, we detected no significant host

genetic variance for parasite virulence, only rhizobia strain and host-genotype-by-rhizobia-strain

effects.The rhizobia effect on resistance is partially explained by a core property of this

nutritional mutualist: its effect on plant growth. In addition to these canonical infection-related

traits, genotype-by-genotype interactions between plants and rhizobia also contributed to genetic

variation in mutualism robustness to parasite infection. These results add to the growing

literature on the ecological and evolutionary significance of symbiotic extended phenotypes, and

suggests that non-defensive mutualists may be overlooked contributors to the evolutionary

potential of defense traits in their hosts.

Rhizobia contributed to genetic variation in resistance and virulence

Rhizobia contributed significantly to genetic variation in two of the three canonical

infection-related traits in our study: resistance (total parasite load) and virulence (the difference

in fitness between infected and uninfected hosts). The contribution of rhizobia to genetic

variation in resistance was small relative to the host, and completely tied up in intergenomic

epistasis ($G_{Host \times Rhizobia}$). Overall, variation in resistance was primarily governed by the host

genome (G_{Host} = 86% of total genetic variance; Figure 2). By contrast, rhizobia contributed

substantially to genetic variation in virulence. Our variance component analyses attributed 5% of

variation in virulence to rhizobia (the nematode infection-by-rhizobia strain interaction for plant

biomass) and an additional 18% of variation to the genotype-by-genotype interaction (GxG)

between plants and rhizobia (the infection-by-genotype-by-strain interaction for plant biomass)

(although this term was not statistically significant; Figure 3). To our surprise, only 3% of

variation in virulence was attributed to the host alone (infection-by-plant genotype interaction for

plant biomass). In other words, rhizobia contributed to nearly 90% of total genetic variation in

virulence. These results show that rhizobial strain identity is a primary determinant of the

virulence of nematode infections in *M. truncatula*.

Significant genotype-by-genotype interactions (GxG) for resistance are consistent with

the established prevalence of intergenomic epistasis in both nutritional and defensive mutualisms

(Batstone et al., 2022; Heath, 2010; Heath & Tiffin, 2007; Parker et al., 2017; Schmid et al.,

2012). Intergenomic epistasis is hypothesized to contribute to the maintenance of genetic

variation in mutualisms, because it creates a situation where the mutualist strain favored by

selection differs across host genetic backgrounds (and vice versa) (Heath & Stinchcombe, 2014).

Ultimately, though, the evolutionary ramifications of GxG depend on how hosts and symbionts

partner in wild populations (De Lisle et al., 2022). If plants and rhizobia pair non-randomly,

intergenomic epistasis could contribute to additive genetic variance in parasite resistance,

because a rhizobia strain's effect on resistance will only occur in a subset of host genetic

backgrounds. Non-random pairing between host and symbiont could arise from partner choice,

which is common in the legume-rhizobia mutualism (Simms & Taylor, 2002). However, to the

best of our knowledge, the covariance between host and symbiont genotypes has not been

directly measured in a facultative mutualism (De Lisle et al., 2022).

Finally, we observed no rhizobia strain effects on tolerance. Tolerance and resistance are

two strategies that hosts evolve in response to parasites, each with differing evolutionary

consequences. Increased resistance tends to select for increased virulence, potentially leading to

an evolutionary arms race whereas tolerance does not (Vorburger & Perlman, 2018). Examples of

symbionts that protect their hosts via increasing tolerance appear to be rare or underrepresented

in the literature (Vorburger & Perlman, 2018) and so the lack of rhizobial effects on host

tolerance may support this notion that microbial effects on host tolerance may be rare. However,

resource availability can influence tolerance (Budischak & Cressler, 2018; Yi et al., 2025),

making this lack of rhizobial effect on tolerance somewhat surprising.

Rhizobia contribute to genetic variation in mutualism robustness to parasite infection

Previous research found that, in addition to stealing resources from their host, parasitic

nematodes decrease nodulation with rhizobia. The magnitude of this effect varied across plant

genetic backgrounds, indicating that mutualism robustness is a heritable plant trait (Wood et al.,

2018). In the present study, we replicated this result, finding significant host genetic variation for

mutualism robustness (infection-by-host genotype interaction for nodulation; Figure 5).

Furthermore, here we found that mutualism robustness is also influenced by a genotype-by-

genotype interaction between plants and rhizobia (infection-by-genotype-by-strain interaction for

nodulation; Figure 5). Rhizobia thus contain genetic variation for mutualism robustness in the

face of nematode infection, indicating that rhizobia evolution may influence mutualism

robustness. This result is consistent with the established importance of context dependence in

mutualisms (Carlson et al., 2025; Chamberlain et al., 2014; Heath & Tiffin, 2007; Parker et al.,

2017). It suggests that parasite infection may be an important environmental factor shaping

mutualism expression and evolution, and suggests that the mutualism disruption of mutualisms

may be an overlooked cost of infection

The genetic and physiological basis of strain effects on parasite infection

Mutualist effects on parasite infection could arise from two underlying mechanisms:

immune crosstalk between mutualism and parasitism, or resource competition or facilitation

(Carlson et al., 2025). While our experiment cannot directly discriminate between immune- and

resource-mediated mechanisms, our path analysis (Figure 6) provides a basis for hypothesizing

about the biological processes that may mediate the effects we observed. We used this analysis to

evaluate whether the rhizobia effect on nematode resistance is mediated by the rhizobia effect on

plant growth (root biomass), consistent with a resource-mediated link between rhizobia and

parasite resistance. Our results are consistent with this hypothesis: rhizobia differed in their

effect on root biomass, and root biomass was associated with parasite load, suggesting that the

rhizobia effect on parasite load is associated with the resources they provide. This suggests that

strain effects on parasite resistance may arise from genotype-by-genotype interactions between

plants and rhizobia for resource- or growth-related biological processes.

By contrast, the direct path linking rhizobia and parasite load was not significant. One

possible interpretation of this result is that immune crosstalk between rhizobia and nematodes

did not substantially influence nematode resistance in this study. However, in our variance

component analyses that corrected for plant size (i.e., analyses that included root biomass as a

covariate), we still detected a significant rhizobia contribution to genetic variation in resistance,

via GxG between rhizobia and host. Together, these observations suggest that, at least for some

strains on some plant genetic backgrounds, the rhizobia effect on parasite resistance is not fully

explained by the resources they provide. This inference is also supported by our recent

transcriptomic study showing that rhizobia strains promote differential expression of genes

enriched for immune functions in nematode galls (Buxton-Martin et al., 2025).

Ultimately, deciphering the biological basis involved in rhizobial strain effects on

infection-related traits is important because these mechanisms differ in their implications for

host-parasite evolution. In defensive mutualisms, the mechanisms by which defensive symbionts

affect virulence (Armitage et al., 2022; Vorburger & Perlman, 2018) and the costs associated

with harboring defensive mutualists (King & Bonsall, 2017) influence the evolutionary

consequences of those protective effects. Competition between parasites and mutualists for

resources is expected to in most cases increase virulence; interference competition is expected to

decrease virulence, and effects mediated by immune responses may vary significantly in their

evolutionary consequences for virulence (Armitage et al., 2022; Vorburger & Perlman, 2018).

Our path analysis also highlights an important consideration for researchers studying

interactions between parasites and resource mutualists: is absolute parasite load or parasite

density (i.e., body size-adjusted parasite load) a more appropriate metric of infection severity?

While absolute parasite load may provide ecologically relevant information about parasite

populations, parasite density may be a better indicator of the fitness cost of parasite infection. If

larger hosts typically harbor more parasites, resource mutualists may often increase absolute

parasite load because they facilitate host growth. However, their effects on parasite density and

the net fitness cost of infection may be less predictable. With this in mind we recommend

considering carefully how infection-related traits – especially infection intensity – is quantified

when in systems in which resource mutualists impact host growth.

Ecological and evolutionary implications of nutritional mutualist effects on heritable variation in

infection-related traits

Our study shows that mutualist contributions to genetic variation in infection-related

traits are not restricted to defensive mutualisms, i.e., interactions where the primary benefit is

defense. This is consistent with pervasive effects of microbial symbionts on a wide range of host

traits (Friesen et al., 2011). Studies in defensive mutualisms suggest that these effects are

evolutionarily significant: defensive mutualists contribute substantially to heritable variation in

infection related traits (Martinez et al., 2017; Parker et al., 2017) and they can drive the rapid

evolution and coevolution of resistance, tolerance, and virulence (Ford et al., 2016, 2017; May &

Nelson, 2014; Rafaluk-Mohr et al., 2018, 2022; Vorburger & Perlman, 2018). Our study adds to

the growing body of studies showing that these effects may not be restricted to defensive

mutualisms (Wood et al., *in review*). Nitrogen-fixing rhizobia are textbook nutritional mutualists.

Yet our study and several others show that they can affect their host's interactions with enemies

(Irmer et al., 2015; Prévitali et al., 2025; Simonsen & Stinchcombe, 2014; Wood et al., 2018).

Future studies should directly test whether genetic variation in non-defensive mutualists

contributes to the evolution of defense traits in their hosts, as it does in defensive mutualisms.

Non-defensive mutualists could accelerate defense evolution by increasing genetic variation

available to selection in hosts, or directly via the rapid evolution of microbial extended

phenotypes (De Lisle et al., 2022; Henry et al., 2021; O'Brien et al., 2021; Wood et al., *in*

review). However, if the underlying microbial genes are subject to conflicting selection (for

example, divergent selection between the free-living and symbiotic life stages), then joint trait

evolution will be difficult to predict without understanding how selection operates in both the

host and the microbe. Therefore, understanding how selection acts on symbiotic extended

phenotypes could be key to predicting the evolution of joint traits, including those related to

parasite infection. Furthermore, pervasive GxG between hosts and mutualists means that total

genetic variation in a joint trait like defense depends on how hosts and symbionts partner in wild

populations. Rhizobia could actually *decrease* total genetic variance in the plants response to

infection if GxG masks heritable differences among host genotypes (Heath, 2010; Wood et al., *in*

review).

Finally, our study raises several additional questions. First, are non-defensive mutualists

more likely to influence resistance against specific types of antagonists? Understanding the

immune- and resource-mediated mechanisms underlying nutritional mutualist effects on defense

traits might allow us to predict when mutualists will affect interactions with antagonists, and if

so, which ones. Microbial mutualists may be especially likely to influence defense against

antagonists governed by similar immune pathways, or defenses whose cost overlaps with the

resources provided or consumed by the mutualist. For example, in our system, nematodes and

rhizobia have significant overlapping pathways by which they interact with their host (Koltai et

al., 2001). Finally, our results highlighted the importance of intergenomic epistasis between hosts

and symbiotic mutualists, but we did not consider GxG between the symbionts themselves. Our

experiment involved single-strain inoculations, yet plants are can be colonized by multiple

rhizobia strains and the benefit that the rhizobia provide depends on the strain(s) they are co-

inoculated with (Rahman et al., 2023; Westhoek et al., 2021). This suggests that other symbiotic

extended phenotypes, including rhizobia effects on infection-related traits, may also be

influenced by GxG between rhizobia strains. These lines of inquiry parallel the emerging focus

within parasitology and phytopathology on concepts like the pathobiome (Vayssier-Taussat et

al., 2014) and disease tetrahedron (Leveau, 2024), which incorporate ecological realism into

host-parasite interactions.

References

Armitage, S. A., Genersch, E., McMahon, D. P., Rafaluk-Mohr, C., & Rolff, J. (2022). Tripartite interactions: How immunity, microbiota and pathogens interact and affect pathogen virulence evolution. *Current Opinion in Insect Science*, *50*, 100871. https://doi.org/10.1016/j.cois.2021.12.011

Bartoń, K. (2025). *MuMIn: Multi-Model Inference* (Version 1.48.11) [Computer software]. https://cran.r-project.org/web/packages/MuMIn/index.html

Batstone, R. T., Lindgren, H., Allsup, C. M., Goralka, L. A., Riley, A. B., Grillo, M. A., Marshall-Colon, A., & Heath, K. D. (2022). Genome-Wide Association Studies across Environmental and Genetic Contexts Reveal Complex Genetic Architecture of Symbiotic Extended Phenotypes. *mBio*, *13*(6), e01823-22. https://doi.org/10.1128/mbio.01823-22

Bolar, K. (2019). *STAT: Interactive Document for Working with Basic Statistical Analysis* (Version 0.1.0) [Computer software]. https://cran.r-project.org/web/packages/STAT/index.html

Bronstein, J. L. (Ed.). (2015). *Mutualism*. Oxford University Press. https://doi.org/10.1093/acprof:oso/9780199675654.001.0001

Brooks, M., Kristensen, K., Benthem, K. van, Magnusson, A., Berg, C., Nielsen, A., Skaug, H., Mächler, M., & Bolker, B. (2017). glmmTMB Balances Speed and Flexibility Among Packages for Zero-inflated Generalized Linear Mixed Modeling. *The R Journal*. https://digitalcommons.unl.edu/r-journal/675

Bruns, E., Carson, M., & May, G. (2012). Pathogen and host genotype differently affect pathogen fitness through their effects on different life-history stages. *BMC Evolutionary Biology*, *12*, 135. https://doi.org/10.1186/1471-2148-12-135

Budischak, S. A., & Cressler, C. E. (2018). Fueling Defense: Effects of Resources on the Ecology and Evolution of Tolerance to Parasite Infection. *Frontiers in Immunology*, *9*. https://doi.org/10.3389/fimmu.2018.02453

Bush, A. O., Lafferty, K. D., Lotz, J. M., & Shostak, A. W. (1997). Parasitology Meets Ecology on Its Own Terms: Margolis et al. Revisited. *The Journal of Parasitology*, *83*(4), 575. https://doi.org/10.2307/3284227

Buxton-Martin, A. D., Gluck-Thaler, E., Yi, E., Stinchcombe, J. R., & Wood, C. W. (2025). Variation in a microbial mutualist has transcriptional and phenotypic consequences for host–parasite interactions. *Functional Ecology*, *39*(10), 2656–2673. https://doi.org/10.1111/1365-2435.70143

Canty, A., & Ripley, B. D. (2025). *boot: Bootstrap Functions* (Version 1.3-32) [Computer software]. https://cran.r-project.org/web/packages/boot/index.html

Carlson, C. I., Frederickson, M. E., & Osmond, M. M. (2025). *How genotype-by-environment interactions can maintain variation in mutualisms* (p. 2024.07.19.604331). bioRxiv. https://doi.org/10.1101/2024.07.19.604331

Chamberlain, S. A., Bronstein, J. L., & Rudgers, J. A. (2014). How context dependent are species interactions? *Ecology Letters*, *17*(7), 881–890. https://doi.org/10.1111/ele.12279

Chen, P., & Roberts, P. A. (2003). Genetic analysis of (a)virulence in Meloidogyne hapla to resistance in bean (Phaseolus vulgaris). *Nematology*, *5*(5), 687–697. https://doi.org/10.1163/156854103322746869

Cook, D. R. (1999). *Medicago truncatula*—A model in the making! *Current Opinion in Plant Biology*, *2*(4), 301–304. https://doi.org/10.1016/S1369-5266(99)80053-3

De Lisle, S. P., Bolnick, D. I., Brodie, E. D., III, Moore, A. J., & McGlothlin, J. W. (2022). Interacting phenotypes and the coevolutionary process: Interspecific indirect genetic effects alter coevolutionary dynamics. *Evolution; International Journal of Organic Evolution*, *76*(3), 429–444. https://doi.org/10.1111/evo.14427

Eck, J. L., Kytöviita, M.-M., & Laine, A.-L. (2022). Arbuscular mycorrhizal fungi influence host infection during epidemics in a wild plant pathosystem. *New Phytologist*, *236*(5), 1922–1935. https://doi.org/10.1111/nph.18481

Florian, H. (2016). DHARMa: Residual Diagnostics for Hierarchical (Multi-Level / Mixed) Regression Models. *CRAN: Contributed Packages*. https://doi.org/10.32614/cran.package.dharma

Ford, S. A., Kao, D., Williams, D., & King, K. C. (2016). Microbe-mediated host defence drives the evolution of reduced pathogen virulence. *Nature Communications*, *7*(1), 13430. https://doi.org/10.1038/ncomms13430

Ford, S. A., Williams, D., Paterson, S., & King, K. C. (2017). Co-evolutionary dynamics between a defensive microbe and a pathogen driven by fluctuating selection. *Molecular Ecology*, *26*(7), 1778–1789. https://doi.org/10.1111/mec.13906

Friesen, M. L., Porter, S. S., Stark, S. C., Wettberg, E. J. von, Sachs, J. L., & Martinez-Romero, E. (2011). Microbially Mediated Plant Functional Traits. *Annual Review of Ecology, Evolution, and Systematics*, *42*(Volume 42, 2011), 23–46. https://doi.org/10.1146/annurev-ecolsys-102710-145039

Fronk, D. C., & Sachs, J. L. (2022). Symbiotic organs: The nexus of host–microbe evolution. *Trends in Ecology & Evolution, Special Issue: Symbiosis*, *37*(7), 599–610. https://doi.org/10.1016/j.tree.2022.02.014

Garcia, J., Barker, D. G., & Journet, E.-P. (2006). Seed storage and germination. In *Medicago truncatula Handbook* (p. 9). Noble Institute.

Hafer-Hahmann, N., & Vorburger, C. (2020). Parasitoids as drivers of symbiont diversity in an insect host. *Ecology Letters*, *23*(8), 1232–1241. https://doi.org/10.1111/ele.13526

Heath, K. D. (2010). Intergenomic epistasis and coevolutionary constrain in plants and rhizobia. *Evolution*, *64*(5), 1446–1458. https://doi.org/10.1111/j.1558-5646.2009.00913.x

Heath, K. D., & Stinchcombe, J. R. (2014). Explaining mutualism variation: A new evolutionary paradox? *Evolution*, *68*(2), 309–317. https://doi.org/10.1111/evo.12292

Heath, K. D., & Tiffin, P. (2007). Context dependence in the coevolution of plant and rhizobial mutualists. *Proceedings of the Royal Society B: Biological Sciences*, *274*(1620), 1905–1912. https://doi.org/10.1098/rspb.2007.0495

Henry, L. P., Bruijning, M., Forsberg, S. K. G., & Ayroles, J. F. (2021). The microbiome extends host evolutionary potential. *Nature Communications*, *12*(1), 5141. https://doi.org/10.1038/s41467-021-25315-x

Hoang, K. L., Read, T. D., & King, K. C. (2024). Incomplete immunity in a natural animal-microbiota interaction selects for higher pathogen virulence. *Current Biology*, *34*(6), 1357-1363.e3. https://doi.org/10.1016/j.cub.2024.02.015

Irmer, S., Podzun, N., Langel, D., Heidemann, F., Kaltenegger, E., Schemmerling, B., Geilfus, C.-M., Zörb, C., & Ober, D. (2015). New aspect of plant–rhizobia interaction: Alkaloid biosynthesis in Crotalaria depends on nodulation. *Proceedings of the National Academy of Sciences*, *112*(13), 4164–4169. https://doi.org/10.1073/pnas.1423457112

Jain, C., Rodriguez-R, L. M., Phillippy, A. M., Konstantinidis, K. T., & Aluru, S. (2018). High throughput ANI analysis of 90K prokaryotic genomes reveals clear species boundaries. *Nature Communications*, *9*(1), 5114. https://doi.org/10.1038/s41467-018-07641-9

Jones, M. G. K., & Goto, D. B. (2011). Root-knot Nematodes and Giant Cells. In J. Jones, G. Gheysen, & C. Fenoll (Eds.), *Genomics and Molecular Genetics of Plant-Nematode Interactions* (pp. 83–100). Springer Netherlands. https://doi.org/10.1007/978-94-007-0434-3_5

Jung, S. C., Martinez-Medina, A., Lopez-Raez, J. A., & Pozo, M. J. (2012). Mycorrhiza-Induced Resistance and Priming of Plant Defenses. *Journal of Chemical Ecology*, *38*(6), 651–664. https://doi.org/10.1007/s10886-012-0134-6

Jutzeler, K. S., Le Clec'h, W., Chevalier, F. D., & Anderson, T. J. C. (2024). Contribution of parasite and host genotype to immunopathology of schistosome infections. *Parasites & Vectors*, *17*(1), 203. https://doi.org/10.1186/s13071-024-06286-6

Kandouh, B., Hasan, A., & Al-Hakeem, A. M. (2019). A complete safe and cost effective method for staining root-knot nematodes meloidogyne spp. *Plant Archives*, *19*(2), 3885–3888.

King, K. C., & Bonsall, M. B. (2017). The evolutionary and coevolutionary consequences of defensive microbes for host-parasite interactions. *BMC Evolutionary Biology*, *17*(1), 190. https://doi.org/10.1186/s12862-017-1030-z

Koch, H., & Schmid-Hempel, P. (2011). Socially transmitted gut microbiota protect bumble bees against an intestinal parasite. *Proceedings of the National Academy of Sciences*, *108*(48), 19288–19292. https://doi.org/10.1073/pnas.1110474108

Koltai, H., Dhandaydham, M., Opperman, C., Thomas, J., & Bird, D. (2001). Overlapping Plant Signal Transduction Pathways Induced by a Parasitic Nematode and a Rhizobial Endosymbiont. *Molecular Plant-Microbe Interactions®*, *14*(10), 1168–1177. https://doi.org/10.1094/MPMI.2001.14.10.1168

Kuznetsova, A., Brockhoff, P. B., & Christensen, R. H. B. (2017). lmerTest Package: Tests in Linear Mixed Effects Models. *Journal of Statistical Software*, *82*, 1–26. https://doi.org/10.18637/jss.v082.i13

Lefcheck, J., Byrnes, J., & Grace, J. (2025). *piecewiseSEM: Piecewise Structural Equation Modeling* (Version 2.3.1) [Computer software]. https://cran.r-project.org/web/packages/piecewiseSEM/index.html

Leveau, J. H. J. (2024). Re-Envisioning the Plant Disease Triangle: Full Integration of the Host Microbiota and a Focal Pivot to Health Outcomes. *Annual Review of Phytopathology*, *62*(Volume 62, 2024), 31–47. https://doi.org/10.1146/annurev-phyto-121423-042021

Lindén, A., & Mäntyniemi, S. (2011). Using the negative binomial distribution to model overdispersion in ecological count data. *Ecology*, *92*(7), 1414–1421. https://doi.org/10.1890/10-1831.1

Liu, X., Matsumoto, H., Lv, T., Zhan, C., Fang, H., Pan, Q., Xu, H., Fan, X., Chu, T., Chen, S., Qiao, K., Ma, Y., Sun, L., Wang, Q., & Wang, M. (2023). Phyllosphere microbiome induces host metabolic defence against rice false-smut disease. *Nature Microbiology*, 1–15. https://doi.org/10.1038/s41564-023-01379-x

Ma, Y., Dias, M. C., & Freitas, H. (2020). Drought and Salinity Stress Responses and Microbe-Induced Tolerance in Plants. *Frontiers in Plant Science*, *11*. https://doi.org/10.3389/fpls.2020.591911

Martinez, J., Longdon, B., Bauer, S., Chan, Y.-S., Miller, W. J., Bourtzis, K., Teixeira, L., &
Jiggins, F. M. (2014). Symbionts Commonly Provide Broad Spectrum Resistance to
Viruses in Insects: A Comparative Analysis of Wolbachia Strains. *PLOS Pathogens*,
10(9), e1004369. https://doi.org/10.1371/journal.ppat.1004369

Martinez, J., Tolosana, I., Ok, S., Smith, S., Snoeck, K., Day, J. P., & Jiggins, F. M. (2017).
Symbiont strain is the main determinant of variation in Wolbachia-mediated protection
against viruses across Drosophila species. *Molecular Ecology*, *26*(15), 4072–4084.
https://doi.org/10.1111/mec.14164

May, G., & Nelson, P. (2014). Defensive mutualisms: Do microbial interactions within hosts
drive the evolution of defensive traits? *Functional Ecology*, *28*(2), 356–363.
https://doi.org/10.1111/1365-2435.12166

McLean, A. H. C., Hrček, J., Parker, B. J., Mathé-Hubert, H., Kaech, H., Paine, C., & Godfray,
H. C. J. (2020). Multiple phenotypes conferred by a single insect symbiont are
independent. *Proceedings of the Royal Society B: Biological Sciences*, *287*(1929),
20200562. https://doi.org/10.1098/rspb.2020.0562

Miozzi, L., Vaira, A. M., Catoni, M., Fiorilli, V., Accotto, G. P., & Lanfranco, L. (2019).
Arbuscular Mycorrhizal Symbiosis: Plant Friend or Foe in the Fight Against Viruses?
Frontiers in Microbiology, *10*. https://doi.org/10.3389/fmicb.2019.01238

O'Brien, A. M., Ginnan, N. A., Rebolleda-Gómez, M., & Wagner, M. R. (2021). Microbial
effects on plant phenology and fitness. *American Journal of Botany*, *108*(10), 1824–1837.
https://doi.org/10.1002/ajb2.1743

O'Keeffe, K. R., Wheeler, B. T., & Mitchell, C. E. (2022). A Microbial Mutualist Within Host
Individuals Increases Parasite Transmission Between Host Individuals: Evidence From a
Field Mesocosm Experiment. *Frontiers in Microbiology*, *13*.
https://doi.org/10.3389/fmicb.2022.824211

Oliver, K. M., Moran, N. A., & Hunter, M. S. (2005). Variation in resistance to parasitism in
aphids is due to symbionts not host genotype. *Proceedings of the National Academy of
Sciences*, *102*(36), 12795–12800. https://doi.org/10.1073/pnas.0506131102

Opperman, C. H., Bird, D. M., Williamson, V. M., Rokhsar, D. S., Burke, M., Cohn, J., Cromer,
J., Diener, S., Gajan, J., Graham, S., Houfek, T. D., Liu, Q., Mitros, T., Schaff, J.,
Schaffer, R., Scholl, E., Sosinski, B. R., Thomas, V. P., & Windham, E. (2008).
Sequence and genetic map of Meloidogyne hapla: A compact nematode genome for plant
parasitism. *Proceedings of the National Academy of Sciences of the United States of
America*, *105*(39), 14802–14807. https://doi.org/10.1073/pnas.0805946105

Parker, B. J., Hrček, J., McLean, A. H. C., & Godfray, H. C. J. (2017). Genotype specificity
among hosts, pathogens, and beneficial microbes influences the strength of symbiont-
mediated protection. *Evolution*, *71*(5), 1222–1231. https://doi.org/10.1111/evo.13216

Prévitali, T., Rouault, M., Pichereaux, C., & Gourion, B. (2025). Lotus resistance against
Ralstonia is enhanced by Mesorhizobium and does not impair mutualism. *New
Phytologist*, *245*(3), 1249–1262. https://doi.org/10.1111/nph.20276

Rafaluk-Mohr, C., Ashby, B., Dahan, D. A., & King, K. C. (2018). Mutual fitness benefits arise
during coevolution in a nematode-defensive microbe model. *Evolution Letters*, *2*(3), 246–
256. https://doi.org/10.1002/evl3.58

Rafaluk-Mohr, C., Gerth, M., Sealey, J. E., Ekroth, A. K. E., Aboobaker, A. A., Kloock, A., &
King, K. C. (2022). Microbial protection favors parasite tolerance and alters host-parasite

coevolutionary dynamics. *Current Biology*, *32*(7), 1593-1598.e3. https://doi.org/10.1016/j.cub.2022.01.063

Rahman, A., Manci, M., Nadon, C., Perez, I. A., Farsamin, W. F., Lampe, M. T., Le, T. H., Martínez, L. T., Weisberg, A. J., Chang, J. H., & Sachs, J. L. (2023). Competitive interference among rhizobia reduces benefits to hosts. *Current Biology*, *33*(14), 2988-3001.e4. https://doi.org/10.1016/j.cub.2023.06.081

Rouchet, R., & Vorburger, C. (2014). Experimental Evolution of Parasitoid Infectivity on Symbiont-Protected Hosts Leads to the Emergence of Genotype Specificity. *Evolution*, *68*(6), 1607–1616. https://doi.org/10.1111/evo.12377

Salem, H., Onchuru, T. O., Bauer, E., & Kaltenpoth, M. (2015). Symbiont transmission entails the risk of parasite infection. *Biology Letters*, *11*(12), 20150840. https://doi.org/10.1098/rsbl.2015.0840

Salvaudon, L., Héraudet, V., & Shykoff, J. A. (2007). Genotype-specific interactions and the trade-off between host and parasite fitness. *BMC Evolutionary Biology*, *7*, 189. https://doi.org/10.1186/1471-2148-7-189

Schmid, M., Sieber, R., Zimmermann, Y.-S., & Vorburger, C. (2012). Development, specificity and sublethal effects of symbiont-conferred resistance to parasitoids in aphids. *Functional Ecology*, *26*(1), 207–215. https://doi.org/10.1111/j.1365-2435.2011.01904.x

Schneider, C. A., Rasband, W. S., & Eliceiri, K. W. (2012). NIH Image to ImageJ: 25 years of image analysis. *Nature Methods*, *9*(7), 671–675. https://doi.org/10.1038/nmeth.2089

Simms, E. L., & Taylor, D. L. (2002). Partner Choice in Nitrogen-Fixation Mutualisms of Legumes and Rhizobia. *Integrative and Comparative Biology*, *42*(2), 369–380. https://doi.org/10.1093/icb/42.2.369

Simonsen, A. K., & Stinchcombe, J. R. (2014). Herbivory eliminates fitness costs of mutualism exploiters. *New Phytologist*, *202*(2), 651–661. https://doi.org/10.1111/nph.12668

Smith, C. A., Renegado, S., & Ashby, B. (2025). The evolution of parasite virulence in the presence of resistance-conferring defensive symbionts. *Journal of Evolutionary Biology*, *38*(8), 1050–1058. https://doi.org/10.1093/jeb/voaf049

Stevens, E. J., Bates, K. A., & King, K. C. (2021). Host microbiota can facilitate pathogen infection. *PLOS Pathogens*, *17*(5), e1009514. https://doi.org/10.1371/journal.ppat.1009514

Vayssier-Taussat, M., Albina, E., Citti, C., Cosson, J. F., Jacques, M.-A., Lebrun, M.-H., Le Loir, Y., Ogliastro, M., Petit, M.-A., Roumagnac, P., & Candresse, T. (2014). Shifting the paradigm from pathogens to pathobiome: New concepts in the light of meta-omics. *Frontiers in Cellular and Infection Microbiology*, *4*. https://doi.org/10.3389/fcimb.2014.00029

Venables, W. N., & Ripley, B. D. (2002). *Modern Applied Statistics with S* (4th ed.). Springer. https://www.stats.ox.ac.uk/pub/MASS4/

Vorburger, C., & Perlman, S. J. (2018). The role of defensive symbionts in host–parasite coevolution. *Biological Reviews*, *93*(4), 1747–1764. https://doi.org/10.1111/brv.12417

Wade, M. J. (2007). The co-evolutionary genetics of ecological communities. *Nature Reviews Genetics*, *8*(3), 185–195. https://doi.org/10.1038/nrg2031

Westhoek, A., Clark, L. J., Culbert, M., Dalchau, N., Griffiths, M., Jorrin, B., Karunakaran, R., Ledermann, R., Tkacz, A., Webb, I., James, E. K., Poole, P. S., & Turnbull, L. A. (2021). Conditional sanctioning in a legume– *Rhizobium* mutualism. *Proceedings of the National Academy of Sciences*, *118*(19), e2025760118. https://doi.org/10.1073/pnas.2025760118

Wood, C. W., Howard, M. M., Yi, E., Heath, K. D., O'Keeffe, K. R., & Lau, J. A. (n.d.).
Integrating microbial mutualists into the evolution of defense. *Trends in Ecology &*
Evolution (In Review).
Wood, C. W., Pilkington, B. L., Vaidya, P., Biel, C., & Stinchcombe, J. R. (2018). Genetic
conflict with a parasitic nematode disrupts the legume–rhizobia mutualism. *Evolution*
Letters, *2*(3), 233–245. https://doi.org/10.1002/evl3.51
Yi, E., Meng, N., & Wood, C. W. (2025). *Mutualist-Provided Resources Increase Susceptibility*
to Parasites (p. 2023.10.15.562412). bioRxiv. https://doi.org/10.1101/2023.10.15.562412
York, L. (2023). Root scanning using a flatbed scanner with transparency unit. *Zenodo*.
https://doi.org/10.5281/zenodo.7926796
704

Tables

	G_{Host}	$G_{Rhizobia}$	$G_{Host \times Rhizobia}$
Resistance	✓✓	N.S.	(✓✓)
Virulence	N.S.	✓✓	N.S.
Tolerance	✓✓	N.S.	Not tested
Mutualism robustness	✓✓	N.S.	✓✓

Table 1. A summative table showing whether genetic variances attributed to host genotype, rhizobia strain, or genotype-by-strain were found for each infection-related traits tested. Check marks indicate a significant effect ($p < 0.05$), check marks within parentheses indicate a marginally significant ($p < 0.1$) effect, and N.S. indicates no significant effect ($p > 0.1$).

 Figure Legends

Figure 1. (A) A conceptual diagram showing our experimental design. (B) Conceptual diagrams

that illustrate the absence (top row) and presence (bottom row) of rhizobia genetic variation for

the four infection-related traits we measured. Rhizobia genetic variation for resistance is

indicated by differences in infection intensity across rhizobia strains. Rhizobia genetic variation

for virulence is indicated by a strain-dependent decrease in fitness in parasite-infected plants.

Rhizobia genetic variation for tolerance is indicated by differences in the slope of the

relationship between host fitness and infection intensity for different rhizobia strains. Rhizobia

genetic variation for mutualism robustness is indicated by a strain-dependent decrease in

mutualist colonization in parasite-infected plants.

Figure 2. Genetic variation in **resistance** to nematode infection. (A) Variance component

analysis of resistance, partitioning variance into three sources: host genotype, rhizobia strain, and

the genotype-by-strain interaction. (B-D) Visualizing G_{Host} (B), $G_{Rhizobia}$ (C), and $G_{Host \times Rhizobia}$

(D) as variation among host genotypes, rhizobia strains, and a subset of genotype-strain

combinations, respectively. Genotype and strain means are the best linear unbiased predictors

(BLUPs) for number of galls, extracted from our mixed model (see Methods). Because our

experiment was an incomplete factorial design, we only show a subset of plant-rhizobia

combinations for the genotype-by-strain interaction (D); see SFigure 2 for subclusters.

Figure 3. Genetic variation in **parasite virulence**. (A) Variance component analysis of

virulence, partitioning variance into three sources: host genotype (the infection x genotype

interaction), rhizobia strain (the infection x strain interaction), and the genotype-by-strain

interaction (the three-way interaction). (B-D) Visualizing G_{Host} (B), $G_{Rhizobia}$ (C), and $G_{Host \times}$

$_{Rhizobia}$ (D) as variation among host genotypes, rhizobia strains, and genotype-strain

combinations, respectively, in the fitness cost of infection. Genotype and strain means are the

best linear unbiased predictors (BLUPs) extracted from our mixed model (see Methods).

Because our experiment was an incomplete factorial design, we only show a subset of plant-

rhizobia combinations for the genotype-by-strain interaction (D); see SFigure 2 for subclusters.

Figure 4. Genetic variation in **tolerance**. (A) Variance component analysis of virulence,

partitioning variance in tolerance into three sources: host genotype (the galls x genotype

interaction) and rhizobia strain (galls x strain interaction). We did not test for a genotype-by-

strain interaction for tolerance. (B-C) Visualizing G_{Host} (B) and $G_{Rhizobia}$ (C) as variation among

host genotypes and rhizobia strains, respectively, in the relationship between parasite load and

fitness. No significant infection-by-strain interactions were observed, indicating no effect of

rhizobia strain on host tolerance.

Figure 5. Genetic variation in **mutualism robustness to infection**. (A) Variance component

analysis of virulence, partitioning variance into three sources: host genotype (the infection x

genotype interaction), rhizobia strain (the infection x strain interaction), and the genotype-by-

strain interaction (the three-way interaction). (B-D) Visualizing G_{Host} (B), $G_{Rhizobia}$ (C), and G_{Host}

$_{\times Rhizobia}$ (D) as variation among host genotypes, rhizobia strains, and genotype-strain

combinations, respectively, in the impact of parasite infection on nodulation. Genotype and

strain means are the best linear unbiased predictors (BLUPs) extracted from our mixed model

(see Methods). Because our experiment was an incomplete factorial design, we only show a

subset of plant-rhizobia combinations for the genotype-by-strain interaction (D); see SFigure 5

for subclusters.

Figure 6: Path analysis results showing significant effect of rhizobia strain on root biomass ($p <$

0.0001) and a significant effect of root biomass on the number of galls ($\beta=0.3151$, $p < 0.0001$)

but no significant direct effect of rhizobia strains on the number of galls ($p = 0.1381$). This path

analysis corrected for variation in these traits among host genotypes (see Methods).

Figure 1. (A) A conceptual diagram showing our experimental design. (B) Conceptual diagrams that illustrate the absence (top row) and presence (bottom row) of rhizobia genetic variation for the four infection-related traits we measured. Rhizobia genetic variation for resistance is indicated by differences in infection intensity across rhizobia strains. Rhizobia genetic variation for virulence is indicated by a strain-dependent decrease in fitness in parasite-infected plants. Rhizobia genetic variation for tolerance is indicated by differences in the slope of the relationship between host fitness and infection intensity for different rhizobia strains. Rhizobia genetic variation for mutualism robustness is indicated by a strain-dependent decrease in mutualist colonization in parasite-infected plants.

Resistance

A. Variance explained: number of galls

0%　　　　　　25%　　　　　　50%　　　　　　75%　　　　　　100%

B. Host genotype (86%, p < 0.001)

Number of Galls BLUP

0.4

0.0

-0.4

-0.8

Host genotype: HM001 HM011 HM012 HM041 HM044 HM048 HM049 HM052 HM065 HM071 HM084 HM086 HM097 HM112 HM138 HM145 HM148 HM167 HM168 HM170

C. Rhizobia strain (0%, p = 0.173)

Number of Galls BLUP

0.4

0.0

-0.4

-0.8

Rhizobia strain: KH35c KH48e MAG154 MAG177 MAG282 MAG358 MAG533 MAG540 USDA1021 WSM1022

D. Host genotype by rhizobia strain (14%, p = 0.061)

Number of Galls BLUP

0.4

0.0

-0.8

| Subcluster 1 | Subcluster 2 | Subcluster 3 | Subcluster 4 | Subcluster 5 |

Subcluster 1: HM049 HM138
Subcluster 2: HM012 HM052
Subcluster 3: HM011 HM071
Subcluster 4: HM012 HM138
Subcluster 5: HM071 HM084

Host genotype

Rhizo: KH35c KH48e MAG177 MAG282 MAG358 MAG540 USDA1021

Figure 2. Genetic variation in **resistance** to nematode infection. (A) Variance component analysis of resistance, partitioning variance into three sources: host genotype, rhizobia strain, and the genotype-by-strain interaction. (B-D) Visualizing G_{Host} (B), $G_{Rhizobia}$ (C), and $G_{Host \times Rhizobia}$ (D) as variation among host genotypes, rhizobia strains, and a subset of genotype-strain combinations, respectively. Genotype and strain means are the best linear unbiased predictors (BLUPs) for number of galls, extracted from our mixed model (see Methods). Because our experiment was an incomplete factorial design, we only show a subset of plant-rhizobia combinations for the genotype-by-strain interaction (D); see SFigure 2 for subclusters.

Parasite virulence

A. Variance explained: host shoot biomass

Host genotype
(39%, p < 0.001)

Host genotype by rhizobia strain
(35%, p < 0.001)

Rhizobia strain (0%, p = 0.071)

B. Infection by host genotype (3%, p = 0.133)

C. Infection by rhizobia strain (5%, p = 0.048)

D. Infection by host geotype by rhizobia strain (18%, p = 0.222)

Subcluster 1   Subcluster 2   Subcluster 3   Subcluster 4   Subcluster 5

Host Shoot Biomass BLUP

Nematode Infection Status

Infected   Uninfected

Genotype

HM011   HM049   HM071   HM138
HM012   HM052   HM084

Rhizo

KH35c   MAG177   MAG358   USDA1021
KH48e   MAG282   MAG540

Figure 3. Genetic variation in **parasite virulence**. (A) Variance component analysis of virulence, partitioning variance into three sources: host genotype (the infection x genotype interaction), rhizobia strain (the infection x strain interaction), and the genotype-by-strain interaction (the three-way interaction). (B-D) Visualizing G_{Host} (B), $G_{Rhizobia}$ (C), and $G_{Host \times Rhizobia}$ (D) as variation among host genotypes, rhizobia strains, and genotype-strain combinations, respectively, in the fitness cost of infection. Genotype and strain means are the best linear unbiased predictors (BLUPs) extracted from our mixed model (see Methods). Because our experiment was an incomplete factorial design, we only show a subset of plant-rhizobia combinations for the genotype-by-strain interaction (D); see SFigure 2 for subclusters.

Host tolerance

Figure 4. Genetic variation in **tolerance**. (A) Variance component analysis of virulence, partitioning variance in tolerance into three sources: host genotype (the galls x genotype interaction) and rhizobia strain (galls x strain interaction). We did not test for a genotype-by-strain interaction for tolerance. (B-C) Visualizing G_{Host} (B) and $G_{Rhizobia}$ (C) as variation among host genotypes and rhizobia strains,

respectively, in the relationship between parasite load and fitness. No significant infection-by-strain interactions were observed, indicating no effect of rhizobia strain on host tolerance.

Mutualism robustness

A. Variance explained: number of nodules

Host genotype (40%, p < 0.001)

Host genotype by rhizobia strain (28%, p < 0.001)

Rhizobia strain (5%, p < 0.001)

B. Infection by host genotype (12%, p < 0.001)

C. Infection by rhizobia strain (0%, p = 0.541)

D. Infection by host geotype by rhizobia strain (16%, p = 0.048)

Subcluster 1 | Subcluster 2 | Subcluster 3 | Subcluster 4 | Subcluster 5

Genotype: HM011, HM049, HM071, HM138, HM012, HM052, HM084

Rhizo: KH35c, MAG177, MAG358, USDA1021, KH48e, MAG282, MAG540

Figure 5. Genetic variation in **mutualism robustness to infection**. (A) Variance component analysis of virulence, partitioning variance into three sources: host genotype (the infection x genotype interaction), rhizobia strain (the infection x strain interaction), and the genotype-by-strain interaction (the three-way interaction). (B-D) Visualizing G_{Host} (B), $G_{Rhizobia}$ (C), and $G_{Host \times Rhizobia}$ (D) as variation among host genotypes, rhizobia strains, and genotype-strain combinations, respectively, in the impact of parasite infection on nodulation. Genotype and strain means are the best linear unbiased predictors (BLUPs) extracted from our mixed model (see Methods). Because our experiment was an incomplete factorial design, we only show a subset of plant-rhizobia combinations for the genotype-by-strain interaction (D); see SFigure 5 for subclusters.

Figure 6: Path analysis results showing significant effect of rhizobia strain on root biomass (p < 0.0001) and a significant effect of root biomass on the number of galls (β=0.3151, p < 0.0001) but no significant direct effect of rhizobia strains on the number of galls (p = 0.1381). This path analysis corrected for variation in these traits among host genotypes (see Methods).

Summary table of significant sources of genetic variances for infection-related traits

	G$_{Host}$	G$_{Rhizobia}$	G$_{Host \times Rhizobia}$
Resistance	✓✓	N.S.	(✓✓)
Virulence	N.S.	✓✓	N.S.
Tolerance	✓✓	N.S.	Not tested
Mutualism robustness	✓✓	N.S.	✓✓

Table 1: A summative table showing whether genetic variances attributed to host genotype, rhizobia strain, or genotype-by-strain were found for each infection-related traits tested. Check marks indicate a significant effect (p<0.05), check marks within parentheses indicate a marginally significant (p<0.1) effect, and N.S. indicates no significant effect (p>0.1).