Comparing DNA Extraction Protocols for Freshwater Prokaryotic Communities: Impacts on Yield and Microbial Profiling

Arthur SZYLIT[1], Ludwig JARDILLIER[2], Maria-Cristina CIOBANU[2], Léa CABROL[3,4], Maialen BARRET[5], Urania CHRISTAKI[1]

[1] Université Littoral Côte d'Opale, Université de Lille, UMR CNRS 8187 LOG, Wimereux, France

[2] Université Paris-Saclay, CNRS, AgroParisTech, Ecologie Société Evolution, 91190, Gif-sur-Yvette, France

[3] Aix Marseille University, Univ Toulon, CNRS, IRD, Mediterranean Institute of Oceanography (MIO), Marseille, France

[4] Millenium Institute Biodiversity of Antarctic and Subantarctic Ecosystems (BASE), Santiago, Chile

[5] Université de Toulouse, Toulouse INP, CNRS, IRD, CRBE, Toulouse, France

corresponding author: SZYLIT Arthur

arthur.szylit@univ-littoral.fr

32 Av. du Maréchal Foch, 62930 Wimereux, France

Abstract:

DNA extraction from aquatic samples is a critical process that influences the quantity and purity of the DNA obtained. This can have profound effects on the accuracy of the community depiction. In this study, DNA extraction workflows of seven commercial DNA extraction kits, along with several modifications of manufacturers' protocols were tested; accounting for a total of 18 different protocols. For each protocol, DNA yield (quantity, replicability and quality), richness and compositional reproducibility based on 16S rRNA gene sequencing, as well as processing time and cost were assessed. With the exception of one kit, the standard protocols recommended by the manufacturer of the kits showed comparable DNA yield results. Shared ASVs between all protocols accounted for >90% of the reads and were mostly abundant ASVs, indicating consistent detection of dominant taxa across all protocols. Adding supplementary lysis and elution steps to the manufacturer's protocols yielded up to ~4× more DNA. However, total read counts and ASV richness were lower as total DNA increased. Manufacturer's protocols therefore showed higher values than their modified versions, although these effects were not significant on community composition. We conclude that the choice of a protocol is the balance between recovering sufficient DNA of good quality versus potential effects on downstream sequencing output (reads and ASVs).

Key words: DNA extraction, Protocols comparison, Prokaryotic community, Freshwater sample, Metabarcoding, 16S rRNA gene

Introduction

No less than three centuries separate Leuwenhoek's pioneering observations of microorganisms and the direct enumeration of aquatic bacteria [1]. It is well known that less than 1% of natural bacteria can be cultured and studied under standard laboratory conditions, a phenomenon called "the great plate count anomaly" [2–4]. Our knowledge of the composition and functions of microbial communities from a variety of habitats on Earth has expanded greatly over the past decades, driven primarily by progress in DNA sequencing technologies [5,6] to a point that environmental science has moved into the 'age of the microbes' [7]. Accurately deciphering community composition and function is essential for addressing key ecological questions, such as understanding how microbial communities are structured in space and time, how this structure affects their functional roles, and ultimately their contribution to biogeochemical cycles. The reduction in both cost and processing time of high-throughput sequencing platforms facilitated their widespread adoption [8,9]. Yet the accuracy of sequencing-based analyses depends strongly on upstream steps, particularly the isolation of high-quality DNA from environmental samples and, for metabarcoding approaches, the choice of primer sets [10]. DNA extraction from aquatic samples is a critical but often delicate process influencing DNA quantity and purity that can have profound effects on the accuracy of the community composition assessment [11]. Despite initiatives for standardization in the framework of international projects such as the Earth Microbiome Project (https://earthmicrobiome.org/protocols-and-standards/), Ocean Sampling Day Initiative [12], NEON [13] and Biogeoscapes (https://biogeoscapes.org/si/), no consensus method has been adopted so far. Differences in sample origin, targeted organisms, research goals, and available resources often lead laboratories to adopt protocols best suited to their constraints and objectives. Historically, DNA extraction relied on phenol-chloroform protocols, combining enzymatic or chemical lysis with ethanol precipitation [14]. Although effective, these methods used hazardous chemicals, were time-consuming, and difficult to standardise. The growing interest in microbial research has led to the development of numerous commercial kits. While the exact composition of these kits is generally not disclosed for commercial reasons, they typically follow similar steps: cell lysis, DNA binding, washing, and elution. Despite these common steps, several studies have reported significant differences between manufactured kits in terms of DNA yield and the downstream analysis of prokaryotic

communities across diverse sample types including, for example, soils [15],
permafrost [16,17], bioreactors [18], marine sediments [19], plant tissues [20], human
microbiomes [21,22], ancient DNA [23], and freshwaters [24].
In aquatic ecosystems, analysing prokaryotic communities typically involves a filtration
step to concentrate biomass onto a membrane. While this filtration step, through filter
porosity and subsequent treatments, significantly influences community differentiation
[25], high levels of suspended particulate matter —independently of prokaryote
abundance— can quickly clog filters, thereby reducing microbial biomass recovery and
impairing sample quality. Such conditions are common in estuarine and coastal marine
environments, and also occur in most freshwater ecosystems. Limited size and depth
make freshwater ecosystems particularly sensitive to allochthonous inputs such as
erosion runoff, wind and rainfall, which can load particles or resuspend previously
settled particles. In this context, the DNA extraction step of water samples remains
critical, as it aims to recover as much DNA as possible from a potentially low biomass.
The objective of the present study was to provide practical recommendations for
selecting and optimizing DNA extraction workflows for prokaryotic community studies
in freshwaters that, by extension, could also apply to microbial eukaryotes and marine
environments. Seven commercial DNA extraction kits were tested. In addition to
manufacturers' protocols, we benchmarked targeted lysis and elution modifications,
leading to 18 protocols that we compared. DNA recovery efficiency was evaluated in
terms of DNA yield, richness and compositional reproducibility using 16S rRNA gene
sequencing, as well as processing time and cost.

Material and Methods

<u>Sample collection</u>: An artificial lake located near Paris, France (48.715506° N,
2.157887° E) was sampled in 2023. This 330 m² waterbody was formed in 2017. Water
samples were collected from the shoreline using a clean plastic beaker attached to a
3 m rod and transferred into a sterile 10 L container. The collected water was
prefiltered sequentially through 100-µm and 30-µm polycarbonate membranes
(Milipore Isopore) to remove large particles. The prefiltered water was filtered onto a
0.2-µm pore-size and 47 mm diameter polycarbonate membrane (Whatman
Nucleopore). The same volume (200 mL) was filtered for all samples. A total of 57
filters were collected across two independent sampling events: Experiment K (Kit
comparison, 45 filters) and Experiment L (Lysis optimization, 12 filters). All membranes

were immediately placed in sterile Petri dishes and stored at –80 °C until further
processing.

<u>DNA extraction protocols</u>: We chose to test some of the most commonly used
manufacturer DNA extraction kits. The kits and associated protocols that were tested
are summarised in Table 1. Originally designed for soil matrices, these kits include
more stringent purification steps than standard water protocols, and are therefore
more appropriate for inhibitor-rich samples such as those usually collected in
freshwaters. For all tests, membranes were cut with sterile scissors under aseptic
conditions prior to DNA extraction to enhance the efficiency of bead-based mechanical
lysis. All subsequent extraction procedures were performed on ice in triplicate.
Experiment K (Kit comparison) compared 15 extraction protocols derived from seven
commercial kits (E1 to E7) produced by four different manufacturers (Table 1). For
each kit, the manufacturer protocol was followed (indicated by suffix "A", Table 1). For
the NucleoSpin Soil kit (Macherey-Nagel), E5A1 and E5A2 correspond to the
manufacturer's protocol using lysis Buffers SL1 and SL2, respectively (Table 1).
Modified protocols for Qiagen kits (E1D, E2B, E3B, E4B) incorporated the following
variations (Table 1): membranes were first incubated overnight at 5°C with agitation
using a HulaMixer (Thermo Scientific™ 15920D) in PowerBead Solution (or CD1 for
DNeasy PowerSoil Pro). For the PowerLyser kit, two simplified protocols (E1C and
E1B) omitted this overnight incubation step. Following this, samples underwent four
freeze-thaw cycles between liquid nitrogen (until completely frozen) and a 65°C water
bath (until fully thawed). For PowerSoil, PowerMax, and PowerLyser PowerSoil kits,
Solution C1 was then added and samples were incubated at 65°C for 10 min prior to
mechanical lysis. For the DNeasy PowerSoil Pro (which does not use Solution C1),
protocols were tested with (E2C) and without (E2B) the addition of 80 µL of 10% SDS
(final concentration 1%) before the 65°C incubation. Mechanical lysis was performed
by vortexing for 25 min. Elution modifications included extended column drying by
placing tubes on ice with loose caps for 5 min following centrifugation. Elution was
performed in two sequential steps using preheated elution buffer at 65°C (room
temperature for E1B): 50 µL was added, incubated for 5 min, and centrifuged; this step
was repeated with an additional 50 µL. For the PowerMax kit, the manufacturer's
protocol (E4A) uses a single 5 mL elution, whereas the modified protocol (E4B)
employed three sequential elution steps (500 µL, 500 µL, and 2500 µL; Table 1)

Furthermore, experiment L (Lysis optimization) compared four extraction protocols
using the PowerLyser kit, all starting from the E1D protocol. E1D was retained as a
control, while three modified versions (E1E, E1F, E1G) replaced the overnight
incubation, freeze-thaw cycles, and vortex-based mechanical lysis with FastPrep-24
bead-beating (MP Biomedicals). FastPrep conditions varied in cycle number, duration,
and speed as specified in Table 1. In protocols with multiple cycles (E1E and E1G), a
30-second pause was included between cycles to reduce heat buildup. All other steps
(Solution C1 incubation, elution modifications) remained identical to E1D.

DNA quality assessment was measured for all extractions from both experiments,
where DNA fragment size was assessed on a 1% agarose gel, and absorbance ratios
(A260/280 nm and A260/230 nm) were measured using a NanoDrop 2000
spectrophotometer (Thermo Scientific). DNA quantity was determined using an Qubit
3 Fluorometer (Invitrogen).

Table1: Summary of DNA extraction kits and protocols tested during the two experiments. Protocols E1–E7A correspond to the manufacturers' protocols, while

other codes indicate modified versions. Checkmarks (✓) indicate applied modifications; dashes (—) indicate omissions. The "+" denotes SDS addition during

65 °C incubation.

Brand	Kit	Protocol	Lysis Modifications			Mechanical lysis				Elution Modifications			
			Agit. O/N	Freeze-thaw 4×	Inc. 65°C (+ SDS)	Instrument	Cycles	Duration	Speed	5mn Air-dry	Buffer 65°C	5mn incubation	Multi-step
Experiment K (45 samples)													
Qiagen	DNeasy PowerLyzer PowerSoil	E1A	—	—	—	MS1 MiniShaker	1	25 mn	1000 rpm	—	—	—	—
		E1B	—	✓	✓					✓	—	✓	2× 50 µl
		E1C	—	✓	✓					✓	✓	✓	2× 50 µl
		E1D	✓	✓	✓					✓	✓	✓	2× 50 µl
	DNeasy PowerSoil Pro	E2A	—	—	—					—	—	—	—
		E2B	✓	✓	✓					✓	✓	✓	2× 50 µl
		E2C	✓	✓	✓+					✓	✓	✓	2× 50 µl
	DNeasy PowerSoil	E3A	—	—	—					—	—	—	—
		E3B	✓	✓	✓					✓	✓	✓	2× 50 µl
	DNeasy PowerMax Soil	E4A	—	—	—					—	—	—	—
		E4B	✓	✓	✓					✓	✓	✓	2x 500 µl + 1 x 2500 µl
Macherey-Nagel	Nucleospin Soil	E5A1 (Buffer SL1)	—	—	—					—	—	—	—
		E5A2 (Buffer SL2)	—	—	—					—	—	—	—
Zymo	ZymoBIOMICS DNA	E6A	—	—	—					—	—	—	—
MP Biomedicals	FastDNA SPIN	E7A	—	—	—					—	—	—	—

Brand	Kit	Protocol	Lysis Modifications			Mechanical lysis				Elution Modifications			
			Agit. O/N	Freeze-thaw 4×	Inc. 65°C (+ SDS)	Instrument	Cycles	Duration	Speed	5mn Air-dry	Buffer 65°C	5mn incubation	Multi-step
Experiment L (12 samples)													
Qiagen	DNeasy PowerLyzer PowerSoil	E1D	✓	✓	✓	MS1 MiniShaker	1	25 mn	1000 rpm	✓	✓	✓	2× 50 µl
		E1E	—	—	✓	FastPrep 24	4	30 s	4 m/s	✓	✓	✓	2× 50 µl
		E1F	—	—	✓		1	45 s	6 m/s	✓	✓	✓	2× 50 µl
		E1G	—	—	✓		4	30 s	6 m/s	✓	✓	✓	2× 50 µl

16S rDNA amplification and sequencing: Prokaryotic community composition of all
samples (57 samples) was determined by high-throughput sequencing of the gene
encoding the small subunit of 16S rRNA in prokaryotes (bacteria and archaea), using
primers U515F and Pk926R (411 bp) targeting the V4–V5 region [26], following the
Earth Microbiome Project protocol (https://earthmicrobiome.org/protocols-and-
standards/16s/). PCR amplification was carried out in a 25 µl reaction volume using
Platinum Taq DNA Polymerase (Invitrogen, Thermo Fisher Scientific, Waltham, MA,
USA), following the manufacturer's recommended reaction mix (Table S1),
supplemented with DTT from the PCR Decontamination Kit (Enzo Life Sciences) as
per manufacturer's instructions. The thermocycling conditions followed the
manufacturer's protocol, with an annealing temperature of 95°C. Amplicon length was
verified by electrophoresis on a 1% agarose gel. For each sample, products from 3 to
5 PCR replicates were combined and purified using the QIAquick PCR Purification Kit
(Qiagen, Germany). Amplicon concentrations were normalised to 20 ng.µL^{-1} prior to
sequencing. All libraries were sequenced in a single MiSeq 2 × 250 run (GENEWIZ,
Germany) to avoid inter-run bias.

Bioinformatic pipeline: A total of 2 806 613 raw paired-end sequences were processed
using the DADA2 pipeline [27] implemented in R (version 4.3.3). Reads were filtered
and trimmed using the *filterAndTrim()* function with the following parameters: forward
and reverse reads were truncated at 240 and 200 bases, respectively, based on
quality profiles generated with QIIME 2 version 2023.9 [28], which showed a median
Phred score of 38 at both positions. The first 19 and 20 bases were removed to
eliminate primers, ambiguous reads were discarded (maxN = 0), and low-quality tails
were trimmed at a quality threshold of 2 (truncQ = 2). Error rates were learned with
the *learnErrors()* function, and amplicon sequence variants (ASVs) were inferred using
dada(). Read pairs were merged using the *mergePairs()* function with a minimum
overlap of 15 bp and a maximum of 1 mismatch. Only merged sequences between
364 and 382 bp in length were retained. Chimeric sequences were removed using the
removeBimeraDenovo() function with the consensus method. ASV taxonomic
assignment was performed with the *assignTaxonomy()* function, using the naive
Bayesian classifier trained on the SILVA SSU reference database version 138.1 [29],
with reverse complement detection enabled (tryRC = TRUE). Before analysis,
unclassified ASVs at the phylum level, or affiliated with Mitochondria or Chloroplast

were removed. Singleton ASVs and those representing less than 0.005% of total reads

across all samples were excluded [30], reducing the dataset from 3 709 to 371 ASVs

(about 10% retained) while preserving 98.4% of the reads (from 1 856 016 to 1 826

234 reads). The sequence dataset was then rarefied to the minimum sequencing

depth observed across samples (19 485 reads), resulting in a final dataset of 57

samples, 1 110 645 sequences, and 371 ASVs. All raw sequences have been

deposited in the European Nucleotide Archive under project accession number

PRJEB94317.

<u>Data Analysis</u>: All statistical analyses were performed using R version 4.3.3 [31]. To

compare DNA quantities between extraction conditions, a one-way ANOVA was

performed after verifying the assumption of homogeneity of variances using Levene's

test (car package, Fox and Weisberg, 2018) and the normality of residuals through the

Shapiro–Wilk test and visual inspection of residual distributions.

The prokaryotic dataset was processed using the phyloseq package version 1.46.0

[33]. Data manipulation was performed using the dplyr package version 1.1.4 [34] and

tidyr version 1.3.1 [35]. Graphs were created using the ggplot2 package version 3.5.1

[36]. Alpha diversity indices (Richness, and Evenness) were calculated from the

rarefied prokaryote dataset using the *estimate_richness* function within the phyloseq

package for each sample. The core community was defined as the set of ASVs

common to all samples (100% prevalence), with a relative abundance of at least

0.001%.

A phylogenetic tree was constructed to compute Weighted UniFrac distances.

Reference sequences were aligned using DECIPHER [37], and a Neighbor-Joining

tree was built from maximum likelihood distances with phangorn [38]. The tree was

then optimised under a GTR+I+Γ model using nearest-neighbor interchange (NNI)

rearrangements. Weighted Unifrac distance [39] was used to study beta diversity in

the rarefied prokaryotic community. A Principal Coordinate Analysis (PCoA) was

performed to visualise the structure of the prokaryotic communities between samples.

Differences in prokaryotic community composition between extraction methods were

assessed using a permutational multivariate analysis of variance (PERMANOVA) [40]

with the *adonis2()* function from the vegan package (version 2.6.4) [41], after verifying

the homogeneity of multivariate dispersions using the *betadisper()* function (vegan

package). Differential abundance analysis was performed at the genus level using

ANCOM-BC [42] to identify genera that significantly differed in abundance according
to the extraction protocol (independently for each sampling campaign). P-values were
adjusted for multiple testing using false discovery rate (FDR) correction. Only genera
with an adjusted p-value below 0.05 and an absolute $\log_2$ fold change greater than 1
were considered significantly differentially abundant.

Results

<u>DNA yield (quantity and replicability)</u>. In Experiment K, DNA yield ranged from 58.6 ±
24 ng for standard Nucleospin (E5A2) to 1745.7 ± 119.2 ng for modified DNeasy
Powerlyser (E1D; Fig 1A). Nucleospin kit (E5A1) showed significantly lower yield
compared to other unmodified manufacturer protocols (ANOVA, p<0.03), while no
significant differences were observed between the remaining 6 unmodified protocols
(E1A, E2A, E3A, E4A, E6A, E7A, ANOVA, p=0.6 ; Fig 1A).
For the four Qiagen kits, only the DNeasy PowerLyzer showed a significant difference
(ANOVA, p<0.001) between the manufacturer's protocol (E1A) and the tested
modifications (E1D). The three modified protocols of the DNeasy PowerLyzer (E1B,
C, D) were also significantly different between them (Tukey post-hoc test, p<0.01) (i.e,
preheating the elution buffer increased yield, and adding an overnight incubation,
Table 1) . For the DNeasy PowerSoil Pro, the SDS addition in E2C did not increase
yield relative to E2B. Overall, the protocol E1D combined the highest DNA yield
(approximately 4-fold higher than the manufacturer's protocol) with a low coefficient of
variation (RSD=6.8%), compared with other modified protocols E2B and E2C
(RSD=33.5% and 24.5%), indicating more consistent performance across replicates
(Fig 1A).
Consequently, the DNeasy PowerLyzer E1D protocol (Table 1) was tested with
stronger mechanical lysis in Experiment L. DNA yields did not then differ significantly
among protocols E1D, E1E, E1F, and E1G (ANOVA, p = 0.75; Fig 1B). E1D showed
a coefficient of variation (CV) of 6.5%, while replicate variability increased with lysis
intensity across the FastPrep conditions, with CVs of 4.9%, 20.8%, and 37.3% for E1E,
E1F, and E1G, respectively.

Figure 1: DNA yield across extraction protocols for Experiment K (A) and Experiment L (B). ANOVAs were performed within each experiment. Protocols in bold correspond to manufacturers' protocols. Asterisks (*) indicate manufacturer protocols that differed significantly from other manufacturer protocols.

DNA quality: In Experiment K, absorbance ratios varied among protocols (Fig 2A). For A260nm/230nm, only PowerLyzer protocols (E1) showed significant differences (ANOVA, p = 0.006), with ratios of 0.9 ± 0.2 for the manufacturer protocol (E1A) and 1.6 ± 0.1 for the modified protocol (E1D) (Fig 2A). For A260nm/280nm, only PowerSoil Pro based protocols (E2) showed a difference (Kruskal-Wallis, p = 0.01), with ratios of 2.1 ± 0.1 for E2A and 1.9 ± 0.05 for E2C (Fig 2A). In Experiment L, A260nm/230nm ratios differed significantly among protocols (ANOVA, p = 0.006), with FastPrep protocols (E1E/F/G) showing higher ratios (1.32 ± 0.17) compared to E1D (0.81 ± 0.12; Fig 2B). A260nm/280nm ratios did not differ significantly (ANOVA, p = 0.65). Gel electrophoresis revealed differences in DNA fragment size between lysis methods (Fig S1): protocols without FastPrep-based lysis (E1D) produced DNA fragments of approximately 48.5 kb, whereas FastPrep protocols (E1E, E1F, E1G) yielded fragments ranging from 10 to 15 kb.

Figure 2: Absorbance ratios A260/230 and A260/280 across extraction protocols for Experiment K (A) and Experiment L (B). Protocols in bold correspond to manufacturers' protocols. ND indicates out-of-range data due to low DNA concentration and contamination leading to unreliable ratios.

Alpha diversity: In Experiment K, ASV richness ranged from 173 ± 7 (E2B) to 192 ± 7 (E1A) (Fig 3A). As for DNA yield described above, among all kits tested, only the DNeasy PowerLyzer protocols (E1A, E1B, E1C, E1D) differed significantly in ASV richness (ANOVA, p = 0.004). Also, within the PowerLyzer, the standard protocol (E1A) showed significantly higher richness (Tukey, p < 0.02, Fig 2A) than its modified versions (E1B-C-D). Read counts mirrored this pattern, decreasing from 32 980 ± 3 189 for E1A to 28 683 ± 1 533 for E1D (Fig S2). Evenness differed significantly among manufacturer protocols (ANOVA, p = 0.03), with Macherey-Nagel kits (E5A1 and E5A2) having the highest values (0.830 ± 0.004). For all Qiagen kits, manufacturer and modified protocols did not differ significantly for this index (Fig 3A). In Experiment L, ASV richness did not differ among E1D, E1E, E1F, and E1G (Fig 3B). Within E1F, two of the three replicates had lower ASV abundance (118 and 136 ASVs), yet the mean of the replicate values did not differ from the others (ANOVA, p=0.1). Evenness differed significantly among the four protocols (ANOVA, p = 0.0003, Fig 3B). E1G had

the lowest values (0.54 ± 0.02) and differed significantly from all other protocols

(Tukey, $p < 0.05$). E1E also had significantly lower values than E1D (Tukey, $p = 0.005$).

Figure 3: Alpha diversity indices (ASV richness and Evenness) of prokaryotic

communities across extraction protocols for Experiment K (A) and Experiment L (B).

Protocols in bold correspond to the manufacturers' protocols.

Overall, taking in account all protocols and replicates, DNA yield was negatively

correlated with ASV richness and reads number in experiment K (Pearson correlation,

p< 0.001, Fig 4A-B).

Figure 4: Relationship between DNA yield and sequencing output across extraction protocols in Experiment K. Correlation between DNA quantity and ASV richness (A) and between DNA quantity and the number of sequencing reads (B). Colors and shapes indicate the extraction protocol. Pearson correlation coefficients and p-values are shown in each panel.

Community structure: In Experiment K, 109 ASVs (representing 42% of the total 261 ASVs) formed the core community (Fig 5A). On average, these ASVs accounted for $93.7 \pm 1\%$ of the reads per tested protocol. Similar results were obtained in Experiment L, where the core community comprised 96 ASVs representing 55% of the total 176 ASVs and accounted for $96.7 \pm 0.83\%$ of reads across protocols (Fig 5B). In Experiment K, most abundant core genera *Limnohabitans*, "hgcI clade" and "CL500-29" accounted for about 21 % of the total reads. In Experiment L, the order Synechococcales, exclusively composed of the genus Cyanobium, was dominant in the core community across all protocols ($38.1 \pm 8.2\%$). Its relative abundance varied among protocols: from 28.5% (E1D) to 35.2% (E1F), 40.9% (E1E), and 47.7% (E1G) (Fig S3). In contrast to the core community, the number of rare ASVs varied among protocols (Fig S4A). In Experiment K, rare ASVs ranged from 14 (E6) to 25 (E4B). Manufacturer protocols generally detected more rare ASVs than their modified counterparts, except for PowerSoil (E3), which detected 18 in both versions. In Experiment L (Fig S4B), a single, intense FastPrep cycle (E1F) detected only 3 rare ASVs, compared to 7–8 in protocols using multiple cycles (E1E, E1G) or vortex-based lysis (E1D).

PCoA based on all ASVs showed significant differences among protocols (PERMANOVA, p = 0.001 for both experiments) and among manufacturers (PERMANOVA, p = 0.003 for Experiment K; Fig S5A-B). However, pairwise comparisons did not identify significant differences between specific manufacturers and protocols, and no clear phylogenetic pattern emerged. In Experiment K, differential abundance analysis revealed that several genera were enriched in manufacturer's protocols compared to their modified counterparts (Fig S6A). Specifically, *Clostridium* was most enriched in E1A (vs E1D), *Fictibacillus* in E2A (vs E2B) and E3A (vs E3B), and *Tumebacillus* in E4A (vs E4B). In Experiment L, differential abundance analysis showed that genera from the Proteobacteria phylum were enriched in E1E and E1G relative to E1D (Fig S6B). While 13 genera differed in abundance between E1D/E1E and E1D/E1G, only one genus differed significantly between E1E and E1G (Fig S6B).

Figure 5: Core prokaryotic community across extraction protocols for Experiment K (A) and Experiment L (B). The core was defined using a prevalence threshold of 100% and a detection threshold of 0.001%. For each donut chart, the center shows the number and percentage of core ASVs and their corresponding reads, while the outer ring displays the proportion of reads assigned to core ASVs for each extraction protocol.

Discussion

On a global scale, aquatic ecosystems contribute greatly to the functioning of the biosphere through their role in the regulation of climate and carbon flows, for example [43,44]. Elucidating the processes driving these ecosystems, particularly biogeochemical fluxes, is essential to appreciate their temporal dynamics and their responses to disturbances. Microbial communities drive all biogeochemical cycles and therefore are key players in the functioning of aquatic ecosystems. The development of molecular approaches and sequencing technologies has rapidly expanded our knowledge of the tree of life, mostly for microbial diversity. This advent has revealed the tremendous microbial phylogenetic diversity and its associated metabolism diversity. It is therefore of primary importance to be able to depict accurately how microbial community composition changes with their environment and, in consequence, to link any change in composition with changes in functions and ultimately on biogeochemical cycles. A large fraction of the microbial diversity consists of low-abundance taxa. Although traditional paradigms attributed the main roles in biogeochemical cycles to abundant taxa, numerous studies have emphasised the significant importance of rare taxa [45,46]. Accurately deciphering spatio-temporal changes in these taxa and their effects on biogeochemical cycles is essential. This requires careful evaluation of all analytical steps. While the effects of primer sets and bioinformatic pipelines have been extensively assessed [47,48], earlier steps, particularly DNA extraction, have received less attention.

Compared to previous studies, the present investigation assessed the differences in the DNA yield and quality as well as in the community depiction between a larger number of manufactured kits and protocol modifications (Table 2). As in previous works realised on freshwater ecosystems, most of the compared protocols involved Qiagen kits (Table 2). Qiagen kits compared here are routinely used for water, while Zymo and MP Biomedicals are more occasionally used [49,50]. The Qiagen DNeasy PowerSoil kit (E3, Table 1) has often been cited for high-quality DNA from water [24] and soil [16] and was later discontinued and replaced by the DNeasy PowerSoil Pro kit (E2). In the present study increasing DNA yield had a negative effect on community richness (Fig 2, Table 3). Although these effects were moderate, they demonstrate—as in previous studies (Table 2)—that protocol selection requires careful consideration of trade-offs based on study-specific objectives.

365 **Table 2**: Comparative studies of DNA extraction protocols from aquatic samples (2005-2023)

Study	Target organism	Matrix	Location	Methods compared	Manufacturers tested	modifications comparison	DNA Measurement	target gene	Sequencing Approach	Main finding
[73]	Bacteria Pathogens	Wastewater	Canada	commercial kit (1) phenol-chloroform variants (6) Other chemical methods (3)	MoBio	none	Yield Quality	16S rRNA	targeted	Few of ten procedures were adequate for bacterial pathogens; bead-beating/ammonium acetate was optimal for microarray detection
[74]	Eukaryotes	Freshwater (waste ponds)	UK	commercials kit (3)	Qiagen MoBio*	none	Yield Quality	18S rRNA	NA (PCR)	QIAGEN DNeasy Blood & Tissue yielded the highest DNA yield and quality for freshwater microalgae, while diversity profiles were comparable across kits
[75]	Bacteria	Freshwater (River)	China	phenol-chloroform variants (3)	NA	Lysis Purification	Yield quality	16S rNA	NA (PCR)	Modified protocol was most efficient, yielding high-quality DNA suitable for molecular applications
[76]	Bacteria Eukaryotes	Freshwater (Lake,River)	Switzerland	commercial kits (2) phenol–chloroform (1)	Qiagen MoBio	none	Yield	16S rRNA COI	metabarcoding targeted	Extraction protocol choice alters biodiversity detection, affecting DNA yield, sequence recovery, and community composition
[24]	Bacteria	Freshwater (River) Wastewater	USA	commercial kits (4)	Qiagen MoBio	none	NA	16S rRNA	metabarcoding	DNA Mini Kit and PowerSoil Kit yielded the most consistent sequencing results across water and wastewater samples
[77]	Prokaryotes Phytoplankton Vertebrates	Seawater (Pacific)	USA	commercial kit (2) phenol-chloroform (1)	Qiagen MoBio	none	Yield	16S rRNA 18S rRNA 12S rRNA	metabarcoding	DNeasy and MoBio kits produce comparable results for multi-trophic eDNA, while phenol-chloroform significantly alters community composition
[78]	Vertebrates	Freshwater (Stream and aquaria) Tap water	Australia	commercial kit (2)	Qiagen MoBio	none	Yield	12S rRNA	NA (qPCR)	Extraction with the Qiagen DNeasy kit provided the highest DNA yield and best cost-efficiency, especially when paired with cellulose nitrate filtration.
[79]	Bacteria Fungi, Plant Vertebrates, Animal	Freshwater (Stream)	New Zealand	commercial kits (6)	Qiagen MoBio ZymoBIOMICS	none	Yield Quality	16S rRNA ITS2, 12S rRNA COI	metabarcoding	DNeasy PowerSoil performs best across taxa and sample types, but no single method is optimal for all organisms
[80]	Bacteria	Freshwater (Lake)	Brazil	commercial kit (1) phenol–chloroform (1)	MoBio	none	NA	16S rRNA	metabarcoding	Phenol–chloroform and kit extractions yielded microbial communities too different to compare.
[81]	Bacteria Eukaryotes	Freshwater (Reservoir)	China	commercial kits (5)	Qiagen MoBio MP Biomedicals	none	Yield Quality	16S rRNA 18S rRNA	metabarcoding	Extraction methods bias rare taxa more than abundant ones; some protocols favor diversity, others reproducibility
[82]	Eukaryotes	Seawater (Atlantic)	Spain	commercial kit (3) Other chemical methods (2)	Qiagen Omega Bio-tek Epicentre	none	Yield Quality	18S rRNA	metabarcoding	Extraction method choice critically affects detection of difficult-to-identify nano- and picoplankton eukaryotes
[83]	Prokaryotes	Seawater (Adriatic Sea)	Croatia	commercials kit (2) phenol-chloroform variants (3) direct PCR (1)	Qiagen	Lysis	Yield	16S rRNA	metabarcoding	All methods showed similar biases; methodological consistency matters more than method choice
This study	Prokaryotes	Freshwater (Lake)	France	commercial kits (18)	Qiagen MoBio Macherey-Nagel Zymo MP Biomedicals	Lysis Elution	Yield	16S rRNA	metabarcoding	Modified commercial kit increased DNA yield, but higher yield negatively correlated with read count and ASV richness

DNA yield (quantity and replicability) and community composition: In contrast to large
between-kit differences in other matrices such as soil or marine sediments [15,19], our
results showed that with the exception of the NucleoSpin Soil kit, all DNA yields were
broadly comparable across manufacturer protocols (Fig 1A, Table 3). The lower yield
obtained with the NucleoSpin Soil kit may be explained by several factors, including
weaker mechanical or chemical lysis, buffer chemistry that promotes inhibitor
carryover or reduces DNA binding, or lower binding/elution efficiency. However,
because the exact buffer compositions and lysis chemistries are proprietary and not
disclosed, we were unable to distinguish the underlying mechanisms in this study.
Interestingly the lower DNA yield of the NucleoSpin Soil kit (E5A1-2, Table 3) was
accompanied by higher Evenness, suggesting reduced dominance rather than
increased richness (Fig 2A).

Among the Qiagen kits tested, the DNeasy PowerLyzer modified protocol (E1D)
yielded up to ~4× more DNA than the standard protocol (E1A) (Fig 1A, Table 3). These
modifications likely acted through complementary mechanisms: freeze-thaw cycles
disrupt cells through heat shock [51,52]; incubation at 65 °C enhances chemical lysis
and destabilises cell walls and membranes [53–55]; and complete ethanol removal
improves DNA desorption from the silica membrane and prevents PCR inhibition. A
two-step elution further increases recovery by rehydrating the matrix twice and
collecting DNA that remains after the first pass [53,56]. Although the modifications
were applied as a bundle, preventing isolation of individual effects, their combined
impact enhanced DNA yield. In addition, two other modifications, tested
independently, also increased DNA recovery: overnight incubation with agitation in the
PowerBead Solution and heating the elution buffer to 65 °C. Overnight agitation likely
promoted cell detachment from particulate matter and improved sample
homogenization, thereby enhancing the efficiency of bead-beating [57]. Likewise,
preheating the elution buffer increased recovery [58], probably by improving
membrane rehydration, increasing DNA solubility, and weakening DNA–silica
interactions. The improved A260nm/230nm ratios observed with protocol
modifications confirm that additional lysis and purification steps effectively reduce
contaminants such as carbohydrates and phenolic compounds, which are common in
freshwater samples with high particulate matter. However, these modifications also
reduced rare ASV detection, reinforcing the trade-off between DNA purity and
preservation of low-abundance community members. Improving contaminant removal

could degrade DNA from rare ASV, compromising the completeness of community
characterization despite higher overall DNA yields.

Within the PowerLyzer-based protocols, all modifications were associated with fewer
observed ASVs despite higher DNA yields (Fig 3, Table 3). ASV counts and read depth
tended to decrease as total DNA increased, in line with previous observations [59].
This negative correlation can be partly explained by co-extraction of non-bacterial
DNA. Additionally, DNA shearing, resulting in smaller DNA fragments, represents a
critical consequence of intensive mechanical lysis that compromises PCR efficiency
and reduces effective amplicon recovery after quality filtering. Intensive lysis benefits
are outweighed by the loss of DNA integrity through fragmentation. Multiple cycles
(E1E, E1G) were more effective at enriching dominant taxa (e.g. Cyanobium) than a
single, more intense cycle (E1F), suggesting that repeated mechanical stress, rather
than instantaneous lysis intensity alone, drives shifts toward dominance. Conversely,
this single intense cycle detected fewer rare ASV, suggesting over-aggressive lysis
disproportionately degrades low-abundance DNA and yielded fewer reads and ASVs
overall. These patterns reflect a trade-off: stronger lysis raises the total amount of
extracted DNA but can degrade sequencing output, requiring an intermediate
optimum.

While protocol modifications reduced rare taxa detection, shared ASVs (common to
all compared protocols) accounted for > 93% of total reads, suggesting consistent
detection of abundant taxa across protocols, with differences concentrated in rare
ASVs as shown in previous study [60]. Although the PCoA of the communities of all
samples tested did not show any clear pattern in phylogenetic structure, community
composition differed significantly among protocols and manufacturers, consistent with
previous findings that extraction methods influence the observed prokaryotic
community [61,62]. The distinct clustering of PowerMax (E4) samples (Fig S5) likely
reflects a mismatch between kit design and sample type: PowerMax is designed to
process up to 10 g of soil, whereas other kits handle up to 0.25 g. Water membrane
samples represent far lower biomass than these capacities, potentially underloading
the PowerMax columns and reducing DNA binding efficiency, which may have altered
the representation of certain taxa.

Cell wall and membrane architecture can influence the efficiency of DNA extraction
[63,64]. Previous studies highlighted that some protocols preferentially recover Gram-

negative taxa [61], whereas others favor Gram-positive taxa [65]. We hypothesised
that adding lysis steps would increase the recovery of Gram+ bacteria because their
thicker peptidoglycan layers require harsher disruption or other microbes with robust
membranes. Contrary to this expectation, several Gram+ genera—*Clostridium*,
Mycobacterium, *Fictibacillus*, *Tumebacillus*, and *Romboutsia*—were relatively more
abundant under the manufacturer protocols than under the modified ones (Fig S6A)
[66–70]. A plausible explanation is a relative abundance artefact caused by
preferential lysis of Gram- cells. Intensified mechanical and thermal disruption may
disproportionately increase the recovery of Gram- DNA, thereby lowering the relative
abundance of Gram+ taxa, even if their absolute recovery also increases. In other
words, more aggressive lysis may improve Gram+ lysis, but the concurrent gain for
Gram- taxa can mask that effect in relative abundance data.

In addition to differences in DNA yield and quality, the time and cost of all 18 protocols
were systematically evaluated given their practical relevance (Table 3; Table S2). For
some pairs, total processing time varied by roughly 1.8-fold; for example, E1F required
about 2h, whereas E1D last for about 4h, excluding the overnight incubation. E4B was
the clear outlier, with approximately 6 h of processing plus an overnight incubation and
a much higher price, about 300% higher than the other kits. Among the protocols,
DNeasy PowerMax Soil (E4) had the highest cost per sample (Table 2-3). This kit has
been used for water samples [71] and has been reported as effective for soils [72], but
it is designed for large input masses. This requires larger buffer volumes and higher-
capacity columns, which increases cost. In addition, E4 is sold only in boxes of 10
extractions, whereas other kits are available in larger formats (up to 250), further
increasing the per-sample cost. Among the standard and modified versions of DNeasy
PowerMax Soil, the modified protocol (E4B) had a longer processing time, mainly
because the freeze-thaw phase required more time at each step.

Table 3: Summary of DNA extraction protocols and performance metrics. Cells are coded as "+" (more favorable) or "−" (less favorable). Handling time was
classified as more than 3 hours plus overnight incubation ("--"), more than 3 hours ("-"), approximately 2 hours 30 minutes ("o"), or less than 2 hours 30 minutes
("+"). Cost per extraction was classified as more than €40 ("--"), between €6 and €15 ("o"), or less than €6 ("+"). Contamination was assessed by absorbance
ratios A260/230 and A260/280 ("+" indicates at least one ratio close to optimal values: ~2.0 for A260/230 and ~1.85 for A260/280; "-" indicates both ratios were
not optimal), and fragment length was assessed by gel electrophoresis ("+" indicates high molecular weight DNA ~48.5 kb, "-" indicates fragmented DNA 10–
15 kb). NA indicates out-of-range data due to low DNA concentration and contamination leading to unreliable ratios. For DNA yield (quantity and reproducibility),
ASV richness, and intra-replicate dissimilarity (wUniFrac), categories were defined by tertiles from the first sampling (n = 45). Lower wUniFrac indicates higher
compositional reproducibility. For the second sampling (n = 12), categories for E1E, E1F, and E1G were assigned relative to E1D and its tertile from the first
sampling.

Description			Logistic		DNA Yields		DNA quality		Community Structure	
Brand	Kit	Protocol	Handling time	Cost	Quantity	Reproductibility	Contamination	Length	Richness (ASV)	Intra-replicate dissimilarity
Experiment K (45 samples)										
Qiagen	DNeasy PowerLyzer PowerSoil	E1A	o	o	-	+	-	+	+	o
		E1B	-		o	+	-	+	o	o
		E1C	-		+	+	+	+	-	+
		E1D	- -		+	+	+	+	-	+
	DNeasy PowerSoil Pro	E2A	o	o	o	o	-	+	o	-
		E2B	- -		+	o	-	+	-	o
		E2C	- -		+	o	+	+	-	+
	DNeasy PowerSoil	E3A	o	NA	o	-	-	+	o	o
		E3B	- -		+	+	+	+	o	+
	DNeasy PowerMax Soil	E4A	-	- -	-	-	NA	NA	o	-
		E4B	- -		o	o	NA	NA	o	+

Description			Logistic		DNA Yields		DNA quality		Community Structure	
Brand	Kit	Protocol	Handling time	Cost	Quantity	Reproductibility	Contamination	Length	Richness (ASV)	Intra-replicate dissimilarity
Macherey-Nagel	Nucleospin Soil	E5A1 (Buffer SL1)	o	+	-	-	-	NA	+	-
		E5A2 (Buffer SL2)	o		-	o	-	NA	+	-
Zymo	ZymoBIOMICS DNA	E6A	o	o	-	-	NA	+	o	-
MP Biomedicals	FastDNA SPIN	E7A	o	o	o	-	NA	+	+	o
Experiment L (12 samples)										
Qiagen	DNeasy PowerLyzer PowerSoil	E1D	- -	o	+	+	-	+	-	+
		E1E	+		+	+	+	-	-	+
		E1F	+		+	+	+	-	- -	+
		E1G	+		+	+	+	-	-	+

Concluding, the fundamental challenge that emerged here is the balance between
recovering sufficient and good quality DNA versus effects on downstream sequencing
output (reads and ASVs) (Table 3). Our results suggested that commercial DNA
extraction kits, and protocol modifications - of lysis and elution - can affect both yield
and the inferred prokaryotic community composition in freshwater samples. These
effects were kit-specific: for example, intensifying the PowerLyzer workflow increased
total DNA, whereas varying FastPrep settings did not change yield but did alter
diversity patterns. Because we evaluated a single freshwater matrix, the magnitude
and direction of these effects may differ in other matrices [62]. Choosing a protocol is
therefore a multi-criteria decision that balances logistics (equipment, hands-on time,
cost) against a clear trade-off between total DNA recovered and downstream
sequencing output (reads and ASVs) (Table 3). In our data, read counts and ASV
richness tended to decrease as total DNA increased. If high yield and reproducibility
are the primary goals and longer handling is acceptable, PowerLyzer with all
modifications (E1D) is a good compromise, while stronger mechanical lysis (e.g E1E)
provides similar yield with less hands-on time. However, both settings reduced ASV
richness and read counts, so if the priority is maximizing usable reads, unmodified
manufacturer protocols are preferable, which also preserve DNA integrity for long-read
sequencing.
Finally, these findings can be complemented by additional tests, to better distinguish
DNA recovery, integrity, and downstream performance: (i) explicitly test for PCR
inhibition using dilution series and internal spike-ins, and consider post-extraction
cleanup; (ii) include absolute quantification (e.g. qPCR or ddPCR) targeting total 16S
rRNA genes and selected taxa to evaluate how lysis conditions affect the detection of
low-abundance groups.
Last but not least, these findings are of primary importance for metagenomic-based
studies of microbial communities since for this approach, high DNA quantities are
usually required to process the sequencing step.

Acknowledgements:
We thank P. Bertolino for laboratory assistance with DNA extraction and amplification
procedures. We acknowledge PRISMARCTYC project (ANR-21-SOIL-0003), the
graduate school IFSEA (ANR-21-EXES- 0011) and the CPER- IDEAL (2022-2027). A.

Szylit was funded by a Ph.D. grant from the 'Région Hauts-de-France' and the
'Université du Littoral Côte d'Opale (ULCO)'.

Conflict of interest statement:
The authors declare that they have no financial or non-financial conflicts of interest in
relation to the content of the article

Data availability statement:
Sequence data have been deposited in the European Nucleotide Archive under project
accession number PRJEB94317.
The complete R pipeline (v1.0.0) and datasets supporting this study are available on
Zenodo: https://doi.org/10.5281/zenodo.18421570

Author contributions
Conceptualization: A.S., L.J., MC.C. Methodology: A.S., MC.C. Investigation: A.S.
Data curation: A.S. Writing – original draft: A.S. Writing – review and editing: A.S., L.J.,
MC.C., L.C., M.B., U.C. Supervision: L.J., M.B., L.C., U.C. Funding acquisition: U.C.,
L.J. All authors have read and agreed to the published version of the manuscript.

References:

1. Hobbie JE, Daley RJ, Jasper S. Use of nuclepore filters for counting bacteria by
fluorescence microscopy. *Appl Environ Microbiol* 1977;**33**:1225–8.
https://doi.org/10.1128/aem.33.5.1225-1228.1977.

2. Staley J, Konopka A. Measurement of In Situ Activities of Nonphotosynthetic
Microorganisms in Aquatic and Terrestrial Habitats. *Annual Review of Microbiology*
1985;**39**:321–46. https://doi.org/10.1146/annurev.micro.39.1.321.

3. Rappé MS, Giovannoni SJ. The Uncultured Microbial Majority. *Annual Review of*
Microbiology 2003;**57**:369–94. https://doi.org/10.1146/annurev.micro.57.030502.090759.

4. Hofer U. The majority is uncultured. *Nat Rev Microbiol* 2018;**16**:716–7.
https://doi.org/10.1038/s41579-018-0097-x.

5. Maxam AM, Gilbert W. A new method for sequencing DNA. *Proceedings of the National*
Academy of Sciences 1977;**74**:560–4. https://doi.org/10.1073/pnas.74.2.560.

6. Sanger F, Nicklen S, Coulson AR. DNA sequencing with chain-terminating inhibitors.
Proceedings of the National Academy of Sciences 1977;**74**:5463–7.
https://doi.org/10.1073/pnas.74.12.5463.

7. Caron DA, Worden AZ, Countway PD *et al.* Protists are microbes too: a perspective. *ISME*
J 2009;**3**:4–12. https://doi.org/10.1038/ismej.2008.101.

8. van Dijk EL, Auger H, Jaszczyszyn Y *et al.* Ten years of next-generation sequencing
technology. *Trends in Genetics* 2014;**30**:418–26. https://doi.org/10.1016/j.tig.2014.07.001.

9. Eren K, Taktakoğlu N, Pirim I. DNA Sequencing Methods: From Past to Present. *Eurasian*
J Med 2022;**54**:S47–56. https://doi.org/10.5152/eurasianjmed.2022.22280.

10. Debroas D, Domaizon I, Humbert J-F *et al.* Overview of freshwater microbial eukaryotes
diversity: a first analysis of publicly available metabarcoding data. *FEMS Microbiol Ecol*
2017;**93**:fix023. https://doi.org/10.1093/femsec/fix023.

11. Lever MA, Torti A, Eickenbusch P *et al.* A modular method for the extraction of DNA and
RNA, and the separation of DNA pools from diverse environmental sample types. *Front*
Microbiol 2015;**6**. https://doi.org/10.3389/fmicb.2015.00476.

12. Ten Hoopen P, Cochrane G. Micro B3 Consortium (2014). Ocean Sampling Day
Handbook. *Version of June 2014 Micro B3 Consortium* 2014.

13. Battelle Memorial Institute. NEON DNA Extraction Standard Operating Procedure v.7.
2022.

14. Chomczynski P, Sacchi N. Single-step method of RNA isolation by acid guanidinium
thiocyanate-phenol-chloroform extraction. *Analytical Biochemistry* 1987;**162**:156–9.
https://doi.org/10.1016/0003-2697(87)90021-2.

15. Zielińska S, Kidawa D, Stempniewicz L *et al.* Environmental DNA as a valuable and unique
source of information about ecological networks in Arctic terrestrial ecosystems. *Environ Rev*
2017;**25**:282–91. https://doi.org/10.1139/er-2016-0060.

16. Vishnivetskaya TA, Layton AC, Lau MCY *et al.* Commercial DNA extraction kits impact
observed microbial community composition in permafrost samples. *FEMS Microbiology*
Ecology 2014;**87**:217–30. https://doi.org/10.1111/1574-6941.12219.

17. Feng S, DeKlotz M, Taş N. Comparison of three DNA extraction methods for recovery of
microbial DNA from Arctic permafrost. *MicroPubl Biol*
2023:10.17912/micropub.biology.000834. https://doi.org/10.17912/micropub.biology.000834.

18. Cabrol L, Malhautier L, Poly F *et al.* Assessing the bias linked to DNA recovery from
biofiltration woodchips for microbial community investigation by fingerprinting. *Appl Microbiol*
Biotechnol 2010;**85**:779–90. https://doi.org/10.1007/s00253-009-2253-8.

19. Cruaud P, Vigneron A, Lucchetti-Miganeh C *et al.* Influence of DNA Extraction Method,
16S rRNA Targeted Hypervariable Regions, and Sample Origin on Microbial Diversity
Detected by 454 Pyrosequencing in Marine Chemosynthetic Ecosystems. *Applied and*
Environmental Microbiology 2014;**80**:4626–39. https://doi.org/10.1128/AEM.00592-14.

20. Abdel-Latif A, Osman G. Comparison of three genomic DNA extraction methods to obtain
high DNA quality from maize. *Plant Methods* 2017;**13**:1. https://doi.org/10.1186/s13007-016-
0152-4.

21. M. Carpi F, Di Pietro F, Vincenzetti S *et al.* Human DNA Extraction Methods: Patents and
Applications. *Recent Patents on DNA & Gene Sequences* 2011;**5**:1–7.
https://doi.org/10.2174/187221511794839264.

22. Elie C, Perret M, Hage H *et al.* Comparison of DNA extraction methods for 16S rRNA gene
sequencing in the analysis of the human gut microbiome. *Sci Rep* 2023;**13**:10279.
https://doi.org/10.1038/s41598-023-33959-6.

23. Gamba C, Hanghøj K, Gaunitz C *et al.* Comparing the performance of three ancient DNA
extraction methods for high-throughput sequencing. *Molecular Ecology Resources*
2016;**16**:459–69. https://doi.org/10.1111/1755-0998.12470.

24. Walden C, Carbonero F, Zhang W. Assessing impacts of DNA extraction methods on next
generation sequencing of water and wastewater samples. *Journal of Microbiological Methods*
2017;**141**:10–6. https://doi.org/10.1016/j.mimet.2017.07.007.

25. Cabrol L, Delleuze M, Szylit A *et al.* Assessing the diversity of plankton-associated
prokaryotes along a size-fraction gradient: A methodological evaluation. *Marine Pollution*
Bulletin 2023;**197**:115688. https://doi.org/10.1016/j.marpolbul.2023.115688.

26. Walters W, Hyde ER, Berg-Lyons D *et al.* Improved Bacterial 16S rRNA Gene (V4 and
V4-5) and Fungal Internal Transcribed Spacer Marker Gene Primers for Microbial Community
Surveys.   *mSystems*   2015;**1**:10.1128/msystems.00009-15.
https://doi.org/10.1128/msystems.00009-15.

27. Callahan BJ, McMurdie PJ, Rosen MJ *et al.* DADA2: High-resolution sample inference
from   Illumina   amplicon   data.   *Nat*   *Methods*   2016;**13**:581–3.
https://doi.org/10.1038/nmeth.3869.

28. Bolyen E, Rideout JR, Dillon MR *et al.* Reproducible, interactive, scalable and extensible
microbiome   data   science   using   QIIME   2.   *Nat*   *Biotechnol*   2019;**37**:852–7.
https://doi.org/10.1038/s41587-019-0209-9.

29. Quast C, Pruesse E, Yilmaz P *et al.* The SILVA ribosomal RNA gene database project:
improved data processing and web-based tools. *Nucleic Acids Research* 2013;**41**:D590–6.
https://doi.org/10.1093/nar/gks1219.

30. Bokulich NA, Subramanian S, Faith JJ *et al.* Quality-filtering vastly improves diversity
estimates   from   Illumina   amplicon   sequencing.   *Nat*   *Methods*   2013;**10**:57–9.
https://doi.org/10.1038/nmeth.2276.

31. R Core Team. R: A language and environment for statistical computing.2013.

32. Fox J, Weisberg S. *An R Companion to Applied Regression*. SAGE Publications, 2018.

33. McMurdie PJ, Holmes S. phyloseq: An R Package for Reproducible Interactive Analysis
and   Graphics   of   Microbiome   Census   Data.   *PLOS*   *ONE*   2013;**8**:e61217.
https://doi.org/10.1371/journal.pone.0061217.

34. Wickham H, François R, Henry L. dplyr: A Grammar of Data Manipulation. 2023.

35. Wickham H, Vaughan D, Girlich M *et al.* tidyr: Tidy Messy Data. 2024.

36. Wickham H. ggplot2: Elegant Graphics for Data Analysis. 2016.

37. Wright E. Using DECIPHER v2.0 to Analyze Big Biological Sequence Data in R. *The R*
Journal 2016.

38. Schliep KP. phangorn: phylogenetic analysis in R. *Bioinformatics* 2011;**27**:592–3.
https://doi.org/10.1093/bioinformatics/btq706.

39. Lozupone C, Lladser ME, Knights D *et al.* UniFrac: an effective distance metric for
microbial   community   comparison.   *ISME   J*   2011;**5**:169–72.
https://doi.org/10.1038/ismej.2010.133.

40. Anderson MJ. A new method for non-parametric multivariate analysis of variance. *Austral*
Ecology 2001;**26**:32–46. https://doi.org/10.1111/j.1442-9993.2001.01070.pp.x.

41. Oksanen J, Simpson GL, Blanchet FG *et al.* vegan: Community Ecology Package. 2022.

42. Lin H, Peddada SD. Analysis of compositions of microbiomes with bias correction. *Nat*
Commun 2020;**11**:3514. https://doi.org/10.1038/s41467-020-17041-7.

43. Duarte CM, Prairie YT. Prevalence of Heterotrophy and Atmospheric CO2 Emissions from
Aquatic Ecosystems. *Ecosystems* 2005;**8**:862–70. https://doi.org/10.1007/s10021-005-0177-
4.

44. Rosentreter JA, Borges AV, Deemer BR *et al.* Half of global methane emissions come
from   highly   variable   aquatic   ecosystem   sources.   *Nat   Geosci*   2021;**14**:225–30.
https://doi.org/10.1038/s41561-021-00715-2.

45. Lynch MDJ, Neufeld JD. Ecology and exploration of the rare biosphere. *Nat Rev Microbiol*
2015;**13**:217–29. https://doi.org/10.1038/nrmicro3400.

46. Chen Q-L, Ding J, Zhu D *et al.* Rare microbial taxa as the major drivers of ecosystem
multifunctionality in long-term fertilized soils. *Soil Biology and Biochemistry* 2020;**141**:107686.
https://doi.org/10.1016/j.soilbio.2019.107686.

47. Abellan-Schneyder I, Matchado MS, Reitmeier S *et al.* Primer, Pipelines, Parameters:
Issues in 16S rRNA Gene Sequencing. *mSphere* 2021;**6**:10.1128/msphere.01202-20.
https://doi.org/10.1128/msphere.01202-20.

48. He J, Zhou T, Shen X *et al.* Primer selection impacts the evaluation of microecological
patterns in environmental microbiomes. *iMeta* 2023;**2**:e135. https://doi.org/10.1002/imt2.135.

49. Li D, Li Z, Yu J *et al.* Characterization of Bacterial Community Structure in a Drinking Water
Distribution System during an Occurrence of Red Water. *Applied and Environmental*
Microbiology 2010;**76**:7171–80. https://doi.org/10.1128/AEM.00832-10.

50. Olalemi AO, Ola HO, Ogundiran BA *et al.* Assessment of bacterial diversity in hand rinse,
stored and drinking water in low-income neighbourhoods of Ibule-Soro, Nigeria. *Water*
Practice and Technology 2024;**19**:4245–53. https://doi.org/10.2166/wpt.2024.249.

51. Chen F, Ye J, Chio C *et al.* A simplified quick microbial genomic DNA extraction via freeze–
thawing cycles. *Mol Biol Rep* 2020;**47**:703–9. https://doi.org/10.1007/s11033-019-05176-w.

52. Xin Y, Xie J, Nan B *et al.* Freeze-Thaw Pretreatment Can Improve Efficiency of Bacterial
DNA   Extraction   From   Meconium.   *Front   Microbiol*   2021;**12**.
https://doi.org/10.3389/fmicb.2021.753688.

53. Miller DN, Bryant JE, Madsen EL *et al.* Evaluation and Optimization of DNA Extraction and
Purification Procedures for Soil and Sediment Samples. *Applied and Environmental*
Microbiology 1999;**65**:4715–24. https://doi.org/10.1128/AEM.65.11.4715-4724.1999.

54. He J, Xu Z, Hughes J. Pre-lysis washing improves DNA extraction from a forest soil. *Soil*
Biology and Biochemistry 2005;**37**:2337–41. https://doi.org/10.1016/j.soilbio.2005.04.016.

55. Islam MdR, Sultana T, Melvin Joe M *et al.* Comparisons of direct extraction methods of
microbial DNA from different paddy soils. *Saudi Journal of Biological Sciences* 2012;**19**:337–
42. https://doi.org/10.1016/j.sjbs.2012.04.001.

56. Desneux J, Pourcher A-M. Comparison of DNA extraction kits and modification of DNA
elution procedure for the quantitation of subdominant bacteria from piggery effluents with real-
time PCR. *MicrobiologyOpen* 2014;**3**:437–45. https://doi.org/10.1002/mbo3.178.

57. Jørgensen T, Haile J, Möller P *et al.* A comparative study of ancient sedimentary DNA,
pollen and macrofossils from permafrost sediments of northern Siberia reveals long-term
vegetational stability. *Molecular Ecology* 2012;**21**:1989–2003. https://doi.org/10.1111/j.1365-
294X.2011.05287.x.

58. Taskin B, Gozen AG, Duran M. Selective Quantification of Viable Escherichia coli Bacteria
in Biosolids by Quantitative PCR with Propidium Monoazide Modification. *Applied and*
Environmental Microbiology 2011;**77**:4329–35. https://doi.org/10.1128/AEM.02895-10.

59. Douglas CA, Ivey KL, Papanicolas LE *et al.* DNA extraction approaches substantially
influence the assessment of the human breast milk microbiome. *Sci Rep* 2020;**10**:123.
https://doi.org/10.1038/s41598-019-55568-y.

60. Pearman JK, Keeley NB, Wood SA *et al.* Comparing sediment DNA extraction methods
for assessing organic enrichment associated with marine aquaculture. *PeerJ* 2020;**8**:e10231.
https://doi.org/10.7717/peerj.10231.

61. Iturbe-Espinoza P, Brandt BW, Braster M *et al.* Effects of DNA preservation solution and
DNA extraction methods on microbial community profiling of soil. *Folia Microbiol* 2021;**66**:597–
606. https://doi.org/10.1007/s12223-021-00866-0.

62. Demkina A, Slonova D, Mamontov V *et al.* Benchmarking DNA isolation methods for
marine metagenomics. *Sci Rep* 2023;**13**:22138. https://doi.org/10.1038/s41598-023-48804-z.

63. Bürgmann H, Pesaro M, Widmer F *et al.* A strategy for optimizing quality and quantity of
DNA extracted from soil. *Journal of Microbiological Methods* 2001;**45**:7–20.
https://doi.org/10.1016/S0167-7012(01)00213-5.

64. Carrigg C, Rice O, Kavanagh S *et al.* DNA extraction method affects microbial community
profiles from soils and sediment. *Appl Microbiol Biotechnol* 2007;**77**:955–64.
https://doi.org/10.1007/s00253-007-1219-y.

65. Salonen A, Nikkilä J, Jalanka-Tuovinen J *et al.* Comparative analysis of fecal DNA
extraction methods with phylogenetic microarray: Effective recovery of bacterial and archaeal
DNA using mechanical cell lysis. *Journal of Microbiological Methods* 2010;**81**:127–34.
https://doi.org/10.1016/j.mimet.2010.02.007.

66. Bouillaut L, McBride SM, Sorg JA. Genetic Manipulation of Clostridium difficile. *Current*
Protocols in Microbiology 2011;**20**:9A.2.1-9A.2.17.
https://doi.org/10.1002/9780471729259.mc09a02s20.

67. Pal D, Bhardwaj A, Kaur N *et al.* Fictibacillus aquaticus sp. nov., isolated from downstream
river water. *International Journal of Systematic and Evolutionary Microbiology* 2018;**68**:160–
4. https://doi.org/10.1099/ijsem.0.002474.

68. Wang M, Nie Y, Wu X-L. Extracellular heme recycling and sharing across species by novel
mycomembrane vesicles of a Gram-positive bacterium. *ISME J* 2021;**15**:605–17.
https://doi.org/10.1038/s41396-020-00800-1.

69. Kang M, Chhetri G, Kim J *et al.* Tumebacillus amylolyticus sp. nov., isolated from garden
soil in Korea. *International Journal of Systematic and Evolutionary Microbiology*
2022;**72**:005376. https://doi.org/10.1099/ijsem.0.005376.

70. Chang S-C, Kao M-R, Saldivar RK *et al.* The Gram-positive bacterium Romboutsia ilealis
harbors a polysaccharide synthase that can produce (1,3;1,4)-β-d-glucans. *Nat Commun*
2023;**14**:4526. https://doi.org/10.1038/s41467-023-40214-z.

71. Lear G, Dickie I, Banks J *et al.* Methods for the extraction, storage, amplification and
sequencing of DNA from environmental samples. *New Zealand Journal of Ecology*
2018;**42**:10-50A.

72. Whitehouse CA, Hottel HE. Comparison of five commercial DNA extraction kits for the
recovery of *Francisella tularensis* DNA from spiked soil samples. *Molecular and Cellular*
Probes 2007;**21**:92–6. https://doi.org/10.1016/j.mcp.2006.08.003.

73. Lemarchand K, Berthiaume F, Maynard C *et al.* Optimization of microbial DNA extraction
and purification from raw wastewater samples for downstream pathogen detection by
microarrays.   *Journal   of   Microbiological   Methods*   2005;**63**:115–26.
https://doi.org/10.1016/j.mimet.2005.02.021.

74. Eland LE, Davenport R, Mota CR. Evaluation of DNA extraction methods for freshwater
eukaryotic   microalgae.   *Water   Research*   2012;**46**:5355–64.
https://doi.org/10.1016/j.watres.2012.07.023.

75. Li P, Yang SF, Lv BB *et al.* Comparison of extraction methods of total microbial DNA from
freshwater. *Genet Mol Res* 2015;**14**:730–8. https://doi.org/10.4238/2015.January.30.16.

76. Deiner K, Walser J-C, Mächler E *et al.* Choice of capture and extraction methods affect
detection of freshwater biodiversity from environmental DNA. *Biological Conservation*
2015;**183**:53–63. https://doi.org/10.1016/j.biocon.2014.11.018.

77. Djurhuus A, Port J, Closek CJ *et al.* Evaluation of Filtration and DNA Extraction Methods
for Environmental DNA Biodiversity Assessments across Multiple Trophic Levels. *Front Mar*
Sci 2017;**4**. https://doi.org/10.3389/fmars.2017.00314.

78. Hinlo R, Gleeson D, Lintermans M *et al.* Methods to maximise recovery of environmental
DNA   from   water   samples.   *PLoS   One*   2017;**12**:e0179251.
https://doi.org/10.1371/journal.pone.0179251.

79. Hermans SM, Buckley HL, Lear G. Optimal extraction methods for the simultaneous
analysis of DNA from diverse organisms and sample types. *Molecular Ecology Resources*
2018;**18**:557–69. https://doi.org/10.1111/1755-0998.12762.

80. Mateus-Barros E, Meneghine AK, Bagatini IL *et al.* Comparison of two DNA extraction
methods widely used in aquatic microbial ecology. *Journal of Microbiological Methods*
2019;**159**:12–7. https://doi.org/10.1016/j.mimet.2019.02.005.

81. Liu M, Xue Y, Yang J. Rare Plankton Subcommunities Are Far More Affected by DNA
Extraction   Kits   Than   Abundant   Plankton.   *Front   Microbiol*   2019;**10**.
https://doi.org/10.3389/fmicb.2019.00454.

82. Muñoz-Colmenero M, Sánchez A, Correa B *et al.* Evaluation of DNA Extraction Methods and Bioinformatic Pipelines for Marine Nano- and Pico-Eukaryotic Plankton Analysis. *Front Mar Sci* 2021;**7**. https://doi.org/10.3389/fmars.2020.584253.

83. Stojan I, Trumbić Ž, Lepen Pleić I *et al.* Evaluation of DNA extraction methods and direct PCR in metabarcoding of mock and marine bacterial communities. *Front Microbiol* 2023;**14**. https://doi.org/10.3389/fmicb.2023.1151907.