1   **Controls of spatio-temporal patterns of soil respiration in a mixed forest**

2   Julian Brzozon*[1], Patricia Schwarzkopf[1], Teja Kattenborn[2], Julian Frey[3], Friederike Lang[1], Helmer Schack-Kirchner[1]

4   [1]Chair for Soil Ecology, University of Freiburg, Bertoldstraße 17, 79098 Freiburg, Germany

5   [2]Chair Sensor-based Geoinformatics, University of Freiburg, Tennenbacher Str. 4, 79106 Freiburg, Germany

6   [3]Chair of Forest Growth and Dendroecology, Tennenbacher Str. 4, 79106 Freiburg, Germany

*Corresponding author: julian.brzozon@soil.uni-freiburg.de

Key-words: Tree distribution, vapour pressure, time-lag effects, hot spot, hot moments

Abstract

Patterns of soil respiration are heterogeneous on temporal and spatial scale. The most important controlling factors
of soil respiration are microclimatic conditions such as soil temperature and soil moisture. However, a strong
pronounced seasonality shifts soil respiration patterns from temperature to moisture-controlled regimes. Rarely
investigated patterns are time-lagged effects prior to soil respiration measurements and influences of trees in mixed
forests on large spatial and temporal scales. We investigated soil respiration over two years on a weekly to fortnightly
measurement rhythm at an approximately 1 ha area in a mixed forest using 35 predefined locations. By tendency, R_s
decreased with increasing distance to the tree and we observed significantly higher respiration in broadleaf patches
(*Fagus sylvatica*) compared to coniferous and mixed patches during the summer season (+27 %, +18 %, respective).
Our data confirmed that soil temperature and soil moisture were important controlling variables in explaining the
temporal patterns of Rs. Yet, our results highlight an additional predictor, which explained a higher proportion in
variability: the current vapour pressure in the atmosphere. In contrast to soil moisture and soil temperature this
predictor was able to track a collapse in respiration due to drought and extreme increases due to rewetting. We
conclude that current water vapour in the atmosphere might be a valuable indicator for CO_2 emission from soils.

Introduction

Soil respiration (R_s), the release of carbon dioxide (CO_2) from soils, represents the second largest terrestrial CO_2 flux and the biggest contributor to total ecosystem respiration (R_e) (Valentini et al. 2000; Davidson et al. 2006; Longdoz et al. 2000; Goulden et al. 1996). Even though temporal patterns showed similar trends, contrasting results of higher R_s compared to R_e were reported at various ecosystem scales (Barba et al. 2018). A certain proportion of this discrepancy may be explained by heterogeneity of R_s in ecosystems, as well as the existence and abundance of hotspots (Barba et al. 2018; Hereș et al. 2021; Scapucci et al. 2025). R_s is composed of heterotrophic respiration (R_{SH}) and autotrophic respiration (R_{SA}) (Kuzyakov 2006), while R_{SA} is seen as the main contributor to total R_s (Raich and Tufekciogul 2000). Thus, the big spatio-temporal variability of R_s may be controlled by the ecosystem structure (Štursová et al. 2016; Lejon et al. 2005; Søe and Buchmann 2005; Köstler et al. 1968).

Across forest ecosystems, only few studies have examined spatio-temporal patterns of R_s at broader scales (Darenova et al. 2024; Hereș et al. 2021; Ruehr and Buchmann 2010; Søe and Buchmann 2005; Ruehr et al. 2010). Among the known controlling variables, microclimatic conditions such as soil temperature (T_s) or soil moisture (Θ_S) are most commonly studied (Wang et al. 2014a; Wang et al. 2014b; Schindlbacher et al. 2012; Darenova et al. 2024; Ruehr et al. 2010; Subke et al. 2003; Epron et al. 1999; Søe and Buchmann 2005; Hereș et al. 2021). A widely accepted relationship between R_s and T_s can be described by site-specifically fitted exponential functions, which can often explain a high proportion of variability in R_s. From these functions, the increase in R_s by changes in T_s of 10 °C can be derived (Q_{10}). However, a simple use of this value neglects seasonality (Janssens and Pilegaard 2003) and important interaction effects with Θ_S (Qi et al. 2002; Lloyd and Taylor 1994). Even though R_s models based on T_s perform well (e.g. Ruehr et al. 2010; Lloyd and Taylor 1994), a rather minor (Ekblad et al. 2005) or indirect (Kuzyakov and Gavrichkova 2010) role of T_s was suggested due to time lag-effects. An increase in solar radiation during spring, peaking during the summer season (in temperate ecosystems), also drives changes in T_s (Kuzyakov and Gavrichkova 2010). Such meteorological conditions prior to R_s measurements represent rarely investigated control variables for R_s, but they might play a crucial role, as they drive ecosystem activity, highlighting the importance of plants as contributors to R_s in ecosystems due to the dependency of R_{SA} on photosynthesis (Davidson et al. 1998; Kuzyakov and Gavrichkova 2010; Tang et al. 2005a; Ekblad and Högberg 2001; Ruehr and Buchmann 2010; Moyano et al. 2008). Ekblad et al. (2005) and Bowling et al. (2002) highlighted the importance of lag-effects of the vapour pressure deficit (VPD) and are, to our knowledge, the only researchers using the VPD to explain patterns of isotopic signatures (^{13}C) of R_s, even though the VPD can be used as a proxy for ecosystem stress due to plant-specific stomatal behaviour as a response to VPD changes (e.g. Hsiao 1973; McAdam and Brodribb 2015; Sanginés de Cárcer et al. 2018). However, shorter-term lag effects of 2-4 hours were also determined for T_s, depending on soil depth (Ruehr et al. 2010).

Despite the fact that microclimatic conditions provide a good explanation for spatio-temporal patterns of R_s, the effect of forest structure is also of crucial importance observed by various studies. In a mixed forest with a high density of various tree species, Vesterdal et al. (2012) observed the highest R_s associated with European beech trees while in a mixed forest containing both pure patches and mixed patches, Khomik et al. (2006) observed the highest R_s associated with deciduous compared to coniferous patches, as well as a higher R_s associated with deciduous trees within the mixed patches. Similar to these results, Longdoz et al. (2000) reported higher R_s associated with European beech compared to Douglas fir patches within one forest. Jochheim et al. (2022) also report differences between deciduous and coniferous patches, but these varied between the measurement years depending on drought conditions. In an early literature review, Raich and Tufekciogul (2000) summarized in general 10 % lower R_s in coniferous forests in close proximity to deciduous forests. These contrasts in R_s associated with species composition may be due to tree specific habitats, as root systems (Köstler et al. 1968), as well as microbial and fungal community compositions (e.g. Hofmann et al. 2023; Prescott and Grayston 2013) and nutrient availability (e.g. Spielvogel et al. 2016; Vesterdal 1999; Vesterdal et al. 2013) depend on the tree species. Additionally, differences in photosynthetic activity between evergreen coniferous trees and the seasonal change in foliation of deciduous trees (Jochheim et al. 2022; Hansen et al. 1997; Curiel Yuste et al. 2004; Longdoz et al. 2000) are important drivers for spatio-temporal patterns of R_s. While tree species effects are common in the literature, only few studies highlight the distance to stemfoot effects on R_s as controlling variable (Jochheim et al. 2022; Søe and Buchmann 2005; Scott-Denton 2003), while Søe and Buchmann (2005) discovered no significant relationship between the distance to the closest tree and R_s rate in a European beech forest. However, Jochheim et al. (2022) observed constantly the highest R_s closest to the trees (1.25 m), compared to more distant positions (2.5 and 3.5 m), with more pronounced species-specific effects during the summertime compared to the wintertime in pure stocks. In a higher elevated subalpine coniferous forest,

Scott-Denton (2003) determined a negative correlation between the distance to the tree and the associated R_s, which
was, however, more likely to be explained by the forest floor thickness. A further positive correlation regarding forest
structural features was determined for the average diameter at breast height (DBH) of trees in proximity of up to 4 m
to measurement position, which explained a considerable proportion in variability (Søe and Buchmann 2005). Due
to these complex dependencies of R_s on spatio-temporal patterns, future trends of total R_s in the scope of climate
change with extreme weather conditions such as heavy rain or extended drought periods (IPCC 2023) are uncertain
because contributors to R_s as microorganism activity (Schindlbacher et al. 2011; Schindlbacher et al. 2012), root
distribution (Brunner et al. 2015; Brunn et al. 2022) as well as spatially variable substrate availability, might alter with
climate change (Brunn et al. 2023; Brunn et al. 2022). Dry periods can alter the pattern or R_s with collapse during
drought and an increase in R_s due to rewetting, which was early discovered as the "birch-effect" (Birch 1958).
However, this change in R_s was not evident in all cases, although there was a collapse and a re-establishment in R_s
but not an extreme increase (Schindlbacher et al. 2011).

The goal of our study is to obtain reliable data for R_s from a mixed forest, to determine it's controlling factors and to
quantify the contribution of predictor variables on R_s. We hypothesize that 1.) temporal patterns of R_s depends on T_s
but an additional proportion in variation can be explained by meteorological parameters. 2.) R_s collapses during
natural occurring drought and increases due to rewetting. 3.) due to species specific phenology R_s associated with
deciduous trees is higher during the summertime, while R_s associated with coniferous trees is higher during the
winter season and we also expect higher R_s closer to tree stems compared to R_s with a larger distance to tree stems,
due to respiratory processes of trees. To test our stated hypothesises, we measured R_s on an area of around
10,000 m^2 at 35 predefined positions over 2 years and fitted it into a statistical model. This approach had also the
aim to identify hot spots and hot moments in R_s. Following, we defined hot spots (Stutz and Lang 2023) as locations
that, on average across the entire area (average of all measurement locations), reflect significantly higher R_s while
cold spots are locations with significantly lower R_s than the seasonal averages. As hot moments, we defined phases
in the temporal development of respiration that deviate from the pattern with unusual reactions (McClain et al.
2003), which we defined as changes in control variables (T_s). Our investigations are part of the collaborative research
center ECOSENSE (Werner et al. 2024)with its joint research area called the ECOSENSE forest (Tesch et al. 2025).

Material & Methods

Site description: The ECOSENSE Forest (Werner et al. 2024) is located in the south-western foothills of the Black
Forest in Germany (48.2684336, 7.8779861, WGS84) with an average elevation of 465-475 m a.s.l.. Dominant tree
species are European beech (*Fagus sylvatica* L.) and Douglas fir (*Pseudotsuga menziesii* Mirbel). Silver fir (*Abies alba*),
European larch (*Larix decidua* Mill.) and Norway spruce (*Picea abies* L.) also occur sporadically, and understorey

Figure 1: Location of the ECOSENSE Forest in Germany (right) and the setup
of R_s measurement plots throughout the forest with location of soil profiles
(left).

vegetation is only sparsely developed. The entire stock is approximate 55-110 years old. The long-year (1981 – 2010)
mean precipitation derived from the HYRAS dataset is 1000 mm yr^{-1}, with a mean annual temperature of 10°C (DWD
2025).

Soil Sampling: In November 2024, we excavated two soil profiles to a depth of 1 m to describe the existing soil types
in the ECOSENSE Forest and described according to (IUSS 2022). We then sampled the soil according to the defined
horizons. Bulk samples were taken from the corresponding depth directly next to undisturbed rings (200 cm^3).
Samples were then stored at 4°C until preparation for analysis.

The organic forest floor consists of a patchy mull type with an averaging thickness of 2-4 cm. Two geological parent
materials are present, resulting in two different soil types. In the southern part of the research area, we identified a
Dystric Stagnic Cambisol (loamic) (IUSS 2022) derived from carbonate-free quaternary loess covering mesozoic
sediments that are strongly weathered with redoximorphic features below a depth of 40 cm. In the northern part of
the research area, we identified a Dystric Skeletic Cambisol (siltic, humic) (IUSS 2022) derived from carbonate-free
quaternary loess, covering mesozoic sandstone with a high skeleton content. Below a depth of 40 cm only thin layers
of fine earth between platy stones can be found. Detailed soil chemical data are presented in table 1. Soil physical
measurements (table 2) were carried out on the 200 cm^3 soil rings.

Soil chemical analysis: Bulk samples were air dried (40°C) for at least 7 days and well mixed in between the days.
After air drying, samples were sieved to 2 mm, with all visible roots, organic material and stones being sorted out. To
determine the carbon to nitrogen ratio (C:N), soil was ground in a ball-mill (MM 400, Retsch GmbH, Germany), dried
at 105°C and analysed using a CHNS elemental analyzer (vario EL cube, Elementar Analysensysteme GmbH,
Germany). For cation exchange capacity (CEC), we followed the methods according to Trüby and Aldinger (1989) and
analysed the extract with an ICP-OES (5800, Agilent, USA). For P$_{available}$, we used citric acid (1%) as extractant and
followed tandard methods (Gutachterausschuss Forstliche Analytik 2022). Analysis for P$_{available}$ was done with an ICP-
OES (5800 ICP-OES, Agilent, USA). The pH-value was measured in H$_2$O following standard methods
(Gutachterausschuss Forstliche Analytik 2022) using an automatic sampler (905 Titrando, Metrohm, Germany).
Dissolved organic carbon (DOC) was extracted using a 1:2 water extract (König and Fortmann 1996) and measured
using a total organic carbon analyzer (TOC-L, Shimadzu GmbH, Germany). Iron oxalate (Fe$_{OX}$) and iron dithionite (Fe$_{DI}$)
were determined using the methods of Tamm (1922) modified by Schwertmann (1964) and measured on an ICP-OES
(5800, Agilent, USA). Chemical soil properties of both profiles are presented in Table 1.

Table 1: Chemical soil properties of the two soil profiles in the ECOSENSE Forest. Values represent average values out
of 3 laboratory analyses for the corresponding soil depth. Organic carbon content (C$_{org}$), carbon to nitrogen ratio (C:N),
effective cation exchange capacity (CEC), base saturation (BS), dissolved organic carbon (DOC), available phosphorous
(P$_{available}$), iron oxides as iron oxalate (Fe$_{OX}$) and iron dithionite (Fe$_{DI}$)

Soil type	depth [cm]	pH$_{H2O}$ [-]	C$_{org}$ [mg g^{-1}]	C:N [-]	CEC [µmol$_c$ g^{-1}]	BS [%$_{molc}$]	DOC [mg g^{-1}]	**P$_{available}$** [mg g^{-1}]	Fe$_{OX}$ [g kg^{-1}]	Fe$_{DI}$ [g kg^{-1}]
Dystric Stagnic Cambisol (loamic)	0-7	4.8	21.6	11.8	74	3	0.4	26.4	4.2	16.2
	7-25	3.8	9.5	9.3	73	12	0.2	14.1	4.4	15.6
	25-38	4.7	3.6	5.1	96	5	0.1	3.9	3.3	16.3
	38-60	3.7	3.1	4.7	104	3	0.1	2.2	2.8	19.1
	60-80	4.9	1.5	2.4	100	3	0.0	0.5	1.4	17.2
	80-100	3.6	0.5	0.9	87	2	0.0	1.2	1.5	13.9
Dystric Skeletic Cambisol (siltic, humic)	0-7	5.0	25.7	14.5	75	5	0.4	43.2	3.6	9.3
	7-20	3.5	13.3	12.0	59	10	0.2	28.5	3.7	10.0
	20-42	4.6	11.0	12.0	56	14	0.2	21.4	3.7	10.3
	42-100	3.7	4.7	6.7	75	24	0.1	8.1	2.6	10.0

Soil physical analysis: Undisturbed ring samples were taken from the cooling storage and were placed in the
laboratory with their lids closeduntill the cylinders reached approximately room temperature. Cylinders were
weighted and the gas diffusivity was subsequently measured using diffusivity chambers (900 cm^3). We used neon (5

ml) as tracer gas, which was well mixed by fans in a chamber below the sample. After injection, we waited 15 minutes
before performing the first gas analysis to ensure the tracer gas had diffused through the sample. Gas from the
chamber was then drawn into a micro gas chromatograph (CP2002, Chrompack, The Netherlands) for six seconds. A
total of six measurements were done for each sample with one flushing process before each measurement. The
apparent diffusion coefficient (D_s) was then calculated by first Fick's law for gas flow. The relative diffusion coefficient
(D_s/D_0) (Schack-Kirchner et al. 2001) was then calculated using the D_0 of Neon (0.297 $m^2 s^{-1}$) at 10°C. The samples
were then saturated through a filter from the bottom side of the ring over the period of one week with a stepwise
increase of water column. A water retention curve was created using the steps of -60 hPa, -300 hPa and -15000 hPa.
After each drainage step, we also measured gas diffusivity. After -300 hPa we determined bulk density and porosity
using a vacuum pycnometer. To determine -15000 hPa, we used a sub-sample with a height of 1 cm from the bottom
part of the undisturbed ring sample. Afterwards the sub-sample and the sample were dried at 105°C to determine
the residual water content. The fraction of the sub-sample at total dry weight was then used to calculate the -15000
hPa on the whole sample. To determine particle size, we took parts of the bulk samples after preparation for chemical
analysis and analysed particles according to (Hartge and Horn 1989).

Table 2: Physical soil properties of the two soil profiles in the ECOSENSE Forest. Values represent mean values out of
3 laboratory analyses for the corresponding depth. D_sD_0 directly after field extraction (FF) and at water potential of -
60 hPa and -300 hPa, porosity (ϕ_{soil}) and bulk density (ρ_{bulk}). Number of samples (n) corresponds to number of
undisturbed rings available for these variables. Stones were estimated in field per horizon. Samples for particle size (n
= 3 for each horizon) were taken from bulk samples. Due to high skeleton content below 42 cm in the Skeletic Cambisol,
no undisturbed rings could be taken, therefore physical parameters were not determinable (n.d.). Standard deviations
were ≤ 0.05 except for marked (*) values.

Soil type (IUSS 2022)	depth [cm]	n	D_s/D_0 FF [-]	D_s/D_0 -60 hPa [-]	D_s/D_0 -300 hPa [-]	ϕ_{soil} [-]	ρ_{bulk} [g cm^{-3}]	stones [%$_{vol}$]	Particle size			Texture class
									Clay [%]	Silt [%]	Sand [%]	
Dystric Stagnic Cambisol (loamic)	0-7	7	0.05	0.05	0.07	0.56	1.15*	0	34.4	58.5	7.0	
	7-25	8	0.05	0.04	0.05	0.53	1.24*	0	32.5	60.4	7.2	
	25-38	6	0.05	0.03	0.05	0.51	1.37*	3	39.2	53.5	7.3	Tu3
	38-60	10	0.03	0.01	0.05	0.45	1.54*	3	41.9	52.4	5.6	
	60-80	9	0.04	0.01	0.01	0.38	1.69*	8	43.5	52.5	4.0	
	80-100	9	0.01	0.01	0.01	0.35	1.79	3	34.8	58.2	7.0	
Dystric Skeletic Cambisol (siltic, humic)	0-7	7	0.06	0.04	0.06	0.57	1.07*	0	26.4	56.5	17.2	
	7-20	7	0.07	0.04	0.07	0.54	1.22*	0	24.9	56.1	17.5	Lu
	20-42	6	0.09	0.05	0.07	0.53	1.24*	17	26.5	54.3	19.2	
	42-100	-	n.d.	n.d.	n.d.	n.d.	n.d.	63	34.8	49.2	16.0	Lt2

Soil respiration measurements: R_s was measured manually over two years on an approximately weekly- to bi-weekly
basis (January 2023-December 2024). The 35 measuring plots as basic spatial investigation units (Figure 1) had an
area of 2.25m^2 where the open-bottom measuring chambers where randomly placed for the measurements without
a permanently installed collar in the soil. For each measurement two open bottom chambers each equipped with a
solid-state CO_2-probe (Vaisala GMP343, Vaisala Oyj, Helsinki, Finland) were placed directly on the forest floor. We
did not use collars to prevent any disturbances to the soil, as collars were shown to potentially influence soil
respiration measurements negatively depending on the insertion depth (Jovani-Sancho et al. 2017; Snell et al. 2014).
During R_s-measurements, we also measured T_s in ca. 13 cm soil depth (GMH 3700 GES401, Seseca GmbH, Germany)
and volumetric soil moisture (Θ_s) (SM150T, Delta-T Devices, UK) in ca. 5 cm depth. The plots for R_s measurements
were established in December 2022 with distances of 10 m between each plot on a north-south transect (21 plots).
Further plots were established (14 plots) east and west of the transect at a distance of approximately 20 m to the
transect (Figure 1). To compensate for potential lateral leakage, we used a correction factor which was determined
in the laboratory for measurements with the foam rubber (CF_F, Equation 1) and for measurements without any
leakage prevention (CF_N, Equation 2). The measurement time was adapted to the expected flux (Fiedler et al. 2022):
at the beginning of the measurement campaign, we started measurements with a length of 390 seconds (20

measurements, gap of 20 seconds). Initial analysis revealed that closing time could be reduced and was set to 190
seconds (20 measurements, gap of 10 seconds). Since the measurement time was shortened, we were able to
measure all plots within one day (since June 2023). Prior to this change, the plots were selected randomly and not
every plot was measured within one day. On three days, measurements were not completed due to weather
conditions or system error (once only 33/35 plots and twice only 26/35 plots). Both individual measurements have
been averaged to a plot mean per measuring day. Due to technical problems in 89 of 2295 (4%) cases R_s was
measured only with one chamber. The order of plots to measure was set randomly for each measuring day to prevent
any bias to diurnal patterns (Ruehr et al. 2010). For R_s flux calculation 50 seconds were cut off at the beginning and
at the end of each single measurement due to saturation effects at the end or slow increase at the beginning. In very
few cases we needed to adapt the time manually, as the device stopped the recording (< 1 %).

$$CF_F = 1.49 + 0.007 * (\frac{T_a + T_s}{2}) \tag{1}$$
$$CF_N = 1.96 + 0.013 * (\frac{T_a + T_s}{2}) \tag{2}$$

With T_a = air temperature in °C and T_s = soil temperature in °C.

Forest stock features: A forest inventory was conducted in the core area of the ECOSENSE forest (circa 3 ha; Tesch
et al. 2025; Werner et al. 2024). All trees with a diameter at breast height (DBH) ≥ 7 cm were recorded in the field,
including species identity, DBH, and spatial position. Tree locations were initially determined using an EMLID RTK
GNSS system (EMLID REACH RS2, Emlid Tech Kft. Budapest, Hungary). Subsequently, terrestrial laser scanning (TLS)
was carried out using a Riegl VZ-400i (RIEGL Laser Measurement Systems GmbH, Horn, Austria) along multiple
transects to capture the stand structure in high detail. The resulting point clouds were used to refine (optimize) the
geolocation of individual tree positions and to derive tree heights for the complete inventory.

Comparable to Søe and Buchmann (2005) we calculated the distance to the nearest neighbouring tree. Additionally,
we buffered a zone around the measurement position (5 m radius) and calculated the average distance to all trees,
the number of trees and the average DBH within the buffered area. We used the distance of 5 m to include all plots
into analysis, as the maximum nearest neighbouring distance was 4.5 m. Since not only *Fagus sylvatica* and
Pseudotsuga menziesii were within the buffer zone we grouped the tree species according to the species
composition, which are also referred to as these groups in the following: deciduous (pure *Fagus sylvatica* composition
[18 plots]), coniferous (either pure *Pseudotsuga menziesii*, [8 plots] pure *Abies alba* [1 plot] or *Pseudotsuga menziesii*
mixed with *Abies alba* [2 plots]) and mixed (always containing *Fagus sylvatica* mixed with different contributions of
Pseudotsuga menziesii, Abies alba, Larix dedidua, Picea abies [6 plots]). The number of trees within the buffer zone
of 5 m varied between 1-10 trees.

Meteorological data: At the beginning of the measurement period no reliable weather data were available from the
ECOSENSE weather monitoring, so we decided to use the HYRAS dataset (DWD 2025) which represents modelled
data of daily precipitation (Rauthe et al. 2013), minimum, maximum and average air temperature (T_a), relative
humidity (H_{rel}) (Frick et al. 2014; Razafimaharo et al. 2020) and global radiation (Brendel et al. 2023) for the specified
cell covering the ECOSENSE Forest (5x5 km cell-size). We calculated the saturated vapour pressure (VP_{sat}) using
Teten's equation with corrections for positive and negative average T_a, which was then converted to actual vapour
pressure (VP_{act}) by multiplication with H_{rel}. The vapour pressure deficit (VPD) was calculated by subtraction of VP_{act}
from VP_{sat}. Wind velocity data were extracted from ERA5-Land datasets (Copernicus Climate Change Service 2019;
Muñoz-Sabater et al. 2021), which also represents a modelled grid covering the ECOSENSE Forest (11 x 7 km cell size).
As a smooth model parameter of seasonality we calculated the negative cosine of the Julian day of the year (cos_{doy}),
(e.g. Sims 1974) with positive, increasing values during the mid of the year (+1) and decreasing, negative values during
the start and end of the year (till -1). We also calculated the negative sine of the Julian day of the year (sin_{doy}), which
then creates the same effect but shifted by three months, and therefore represents the spring and autumn season.
This method takes into account a transition with a development, apart from standard seasonal classification, which
cumulates three months to only one variable and is therefore specifically well suited for model predictions. In order
to obtain more general statements about the seasonal effects in addition to the cos_{doy} we also used the classical
seasonal classification according to the German weather service (DWD) and averaged the values over the seasons
for a more common evaluation. Seasons were classified as Winter: December – February, Spring: March – May,
Summer: June – August, Autumn: September – November.

Statistical analysis: To determine the predictors for **temporal patterns** in R_s we used linear models (lm from stats-
package, R Core Team 2023) applied to the average R_s values from the measuring days of all 35 sampling plots. As
predictors, we used the respective average weather parameters from the day of the measurements of R_s as well as
aggregated mean or summarised values of these predictors 1-7 days prior to the measurement day. To prevent model
overfitting and to identify predictors that provide meaningful information on the development of R_s only predictors
with an R^2 above 0.4 were taken into consideration for the further model optimization. We also tested the model for
normal distribution and heteroscedasticity using analytical plots. If the common problem of heteroscedasticity was
present, we calculated robust standard errors using HC4 (Hayes and Cai 2007). Additionally, we also applied different
methods to determine temporal patterns by checking dependencies of R_s on T_s alone (equation 3) and in combination
with soil moisture Θ_S (equation 4). We also calculated Q_{10} values (equation 5) using non least square models (nls from
stats-package, R Core Team 2023) with fitted parameters.

$$R_s = a * e^{b*T_s} \qquad (3)$$
$$R_s = c * \Theta_s * e^{d*T_s} \qquad (4)$$
$$Q_{10} = e^{10*b} \qquad (5)$$

With R_s = the modelled soil respiration [µmol CO_2 m^{-2} s^{-1}], T_s = soil temperature [°C] at reference depth (13 cm) and
Θ_s [%$_{vol}$]. a, b, c, d were fitted using non least square models.

To visualize and determine **spatial patterns,** we used inverse distance weighting (idw from phylin-package, Tarroso
et al. 2019; Tarroso et al. 2015; Pebesma and Bivand 2023; Pebesma 2018). Dependencies of R_s on forest structural
features were tested using linear models (lm from from stats-package, R Core Team 2023) and pairwise wilcox rank
sum tests (from stats-package, R Core Team 2023) for group comparisons. Answering the question whether
(seasonal) **hot spots** for R_s exist in the ECOSENSE Forest, we calculated a mixed effects model (lmer from lme4, Bates
et al. 2015) with the measurement plot as explanatory variable, but the day of the measurement as a random effect.
From this, we calculated estimated marginal mean values (emmeans from emmeans package, Lenth 2023) and set
those in contrast (contrast from stats-package, R Core Team 2023) to the averaged values of the corresponding
season, to identify plots with statistically significant higher or lower deviations from the averaged seasonal R_s. To
identify **hot moments**, we used linear models (lm from stats-package, R Core Team 2023) and spearman correlation
(cor from stats-package, R Core Team 2023) tests to check for unusual patterns between R_s and T_s. All statistical
analysis was done in RStudio 2023.06.0 based on R4.2.3 (R Core Team 2023). For data organization we used the
tidverse (Wickham et al. 2019), ggplot2 (Wickham 2016) and lubridate packages (Grolemund and Wickham 2011).

Results

Weekly to fortnightly soil respiration measurements

In total, 4507 single R_s measurements were available for data evaluation, which were merged to 2295 measurements
on the plot level on a total of 72 measurement days. Annually averaged R_s in 2024 (3.10 ± 1.86 CO_2 µmol m^{-2} s^{-1}) was
higher than in 2023 (2.74 ± 1.89 CO_2 µmol m^{-2} s^{-1}). The highest daily averaged R_s was reached in 2023 (7.74 ± 2.0 µmol
CO_2 m^{-2} s^{-1}), while the highest daily averaged R_s was lower by almost 1 µmol CO_2 m^{-2} s^{-1} in 2024 (6.83 ± 2.44
µmol CO_2 m^{-2} s^{-1}). On plot level, the highest R_s was reached in 2023 with 27.1 µmol CO_2 m^{-2} s^{-1} which was lower by
11.4 µmol CO_2 m^{-2} s^{-1} in 2024. We identified strong seasonal fluctuations with peak R_s during summertime with only
small differences between the two years, but high differences between the seasons within one year (Table 3).
Unfortunately, the measurement device for Θ_S broke down in January 2024 due to high skeleton content in one plot
and measurements were not possible thereafter. The two years differed in the precipitation amount. In 2023 annual
precipitation was 37 mm below the long year average, in 2024 it was 106 mm above. The VP$_{cur}$ was similar for both
years but showed deviations during the summertime with an averaging higher VP$_{cur}$ during summer 2023, while the
VPD also deviated by averaging 0.19 kPa during both summer seasons (Table 3). In both years the VP$_{cur}$ and VPD
showed seasonal variation as well and followed the development of R_s. Also, the Q_{10} values (Equation 5) determined
using the daily average R_s and T_s revealed seasonal variability with the highest Q_{10} for both winter season in both
years. However, the highest Q_{10} during winter 2023 exceeded the Q_{10} in winter 2024 by a factor of 2.6. In 2023 the
Q_{10} continuously decreased till autumn, while in 2024, the lowest Q_{10} was reached in the summer season. Calculating
the Q_{10} based on values throughout the whole year, the Q_{10} for both years were similar with 3.32 ± 1.22 and 3.11 ±
1.12 in 2023 and 2024, respectively. The correlation matrix separated by season (Table 4) shows the different

responses of R_s in relation to the climate predictors. Interestingly, the daily average R_S correlates differently with the presented meteorological parameters across seasons, showing the lowest correlations and even negative correlations in some cases during the summer for all parameters except for Θ_S in 2023, although positive correlations were observed during the other seasons.

Table 3: Seasonal averaged soil respiration (R_S), with number of measurement days (n) within the respective season, mean soil temperature (T_s) during R_s measurement campaigns and soil moisture (Θ_S). All values are averaged per season for the corresponding years of measurement on the ECOSENSE Forest and given with standard deviation. Q_{10} values were calculated for each season and years separately and given with standard errors from the fitted model. From the HYRAS dataset we summarised precipitation ($\sum$ Prec.) and averaged air temperature (T_a), current vapour pressure (VP_{cur}) and vapour pressure deficit (VPD).

Year	Season	R_S	n	T_s	Q_{10}	Θ_S	$\sum$ Prec.	T_a	VP_{cur}	*VPD*
		μmol CO$_2$ m^{-2} s^{-1}		°C		%vol	mm	°C	kPa	kPa
2023	Winter	**1.07 ± 0.53**	8	4.8 ± 1.4	20.4 ± 2.1	33.0 ± 3.5	174.2	3.9 ± 3.9	0.67 ± 0.18	0.16 ± 0.09
	Spring	**1.98 ± 1.12**	10	8.7 ± 2.8	2.9 ± 1.9	33.2 ± 3.8	234.1	9.6 ± 4.5	0.92 ± 0.27	0.33 ± 0.2
	Summer	**4.83 ± 2.07**	10	15.8 ± 1.7	2.8 ± 2.1	16.3 ± 4.5	279.6	19.4 ± 3.1	1.53 ± 0.33	0.75 ± 0.34
	Autumn	**2.76 ± 0.96**	10	13.4 ± 3.1	2.1 ± 1.4	17.7 ± 9.9	275.1	12.1 ± 5.8	1.16 ± 0.36	0.35 ± 0.24
2024	Winter	**1.14 ± 0.44**	7	5.1 ± 1.3	7.9 ± 2.9	n.d.	181.4	3.7 ± 4.3	0.68 ± 0.19	0.15 ± 0.09
	Spring	**2.30 ± 1.09**	10	8.9 ± 2.1	5.1 ± 1.5	n.d.	374.3	10.5 ± 4.3	0.98 ± 0.23	0.33 ± 0.22
	Summer	**5.36 ± 0.93**	9	16.1 ± 2.1	1.4 ± 1.4	n.d.	282.3	19.0 ± 3.0	1.67 ± 0.28	0.56 ± 0.23
	Autumn	**3.27 ± 1.49**	8	12.6 ± 3.2	3.5 ± 1.2	n.d.	267.9	10.5 ± 4.9	1.13 ± 0.35	0.20 ± 0.13

Table 4: Correlation matrix (Spearman-correlation coefficients) between average R_s and potential meteorological predictors (current daily values compared to daily averaged R_s) and the seven day (7-d.) aggregated values prior to the measurement day separated by season for the two measurement years. Climate predictors are the vapour pressure deficit (VPD), current vapour pressure (VP_{cur}), air temperature (T_{air}), global radiation (Glob-Rad), daily soil temperature (T_s) and soil water content (Θ_S).

Year	Season	VPD		VP_{cur}		T_a		Glob-Rad		Prec		T_s	Θ_S
		daily	7-d.	daily	7-d.	daily	7-d.	daily	7-d.	daily	7-d.	daily	daily
2023	Winter	0.17	-0.36	0.76	0.62	0.69	0.57	-0.07	-0.29	0.30	0.26	0.88	0.05
	Spring	0.2	0.26	0.78	0.77	0.65	0.59	0.53	0.46	0.29	0.05	0.64	0.6
	Summer	**-0.6**	**-0.32**	**0.48**	**0.55**	**0.03**	**0.14**	**-0.39**	**-0.55**	**-0.33**	**0.65**	**0.53**	**0.6**
	Autumn	0.22	0.37	0.79	0.78	0.38	0.67	0.49	0.58	-0.07	-0.04	0.64	-0.3
2024	Winter	0.71	0.64	0.68	0.89	0.68	0.82	0.21	0.39	0	0.57	0.64	-
	Spring	0.42	0.58	0.83	0.96	0.73	0.9	0.2	0.54	0.66	0.08	0.95	-
	Summer	**-0.25**	**-0.22**	**0.17**	**0.33**	**0.22**	**0.16**	**0.07**	**-0.1**	**0**	**0.65**	**0.42**	-
	Autumn	0.14	0.33	0.68	0.81	0.43	0.81	0.38	0.48	-0.07	-0.29	0.48	-

Hot moments of R_s

During the transition between spring and summer in 2023 average R_s collapsed from 10[th] May until 16[th] June by 3 μmol CO$_2$ m^{-2} s^{-1}. This phase was characterised by increasing T_s and T_a but low precipitation and no precipitation (Figure 2) on 29/38 days. After the last higher rain event on 21[st] May (14.5 mm d^{-1}), the weather remained dry and only a low rainfall event on 4[th] June with 1.4 mm d^{-1} (0.14 % of 30-year average) occurred, causing a slight increase in R_s (7[th] June) by 0.72 μmol CO$_2$ m^{-2} s^{-1} but a further decrease in the following week. During this whole phase, Θ_S dropped on average by 24 %$_{vol}$. With an onset of rainfall from 18[th] June till 22[nd] June with a total of 29.5 mm (approx. 3 % of 30-year average) out of which 20 mm d^{-1} fell within just in one day (13[th] highest rainfall event over the two years) a sharp increase in R_s by 6.26 μmol CO$_2$ m^{-2} s^{-1} was measured on 23[rd] June, while average T_s and Θ_S increased only slightly by 1.5 °C and 1.3 %$_{vol}$, respective. The day after the strong rainfall event represented the highest R_s (7.74 ± 2.0 μmol CO$_2$ m^{-2} s^{-1}) in the measurement period. Additionally, VPD increased from 22[nd] May until 12[th] June by 0.94 kPa but decreased directly afterwards till 23[rd] June by 0.33 kPa following the larger precipitation events. During this phase the maximum average VPD was 1.37 kPa. The correlations between R_s and T_s as well as VPD, were negative during this phase (ρ_s = -0.77 and ρ_s = -0.94, respective), but positive between R_s and Θ_S (ρ_s = 0.83). After the stronger rainfall event R_s dropped after the maximum efflux by averaging 4.4 μmol CO$_2$ m^{-2} s^{-1} just six days later. The second highest R_s (7.3 ± 4.1 μmol CO$_2$ m^{-2} s^{-1}) over the two years was reached two months later in August. It is also noticeable, that in total R_s was higher during 2024, but the two highest rates in R_s were reached in 2023 and exceed those from 2024 by 0.5 – 1.2 μmol CO$_2$ m^{-2} s^{-1}. However, we also identified further phases characterized by low

322 precipitation and low Θ_S i.e. during the end of August 2023 which was not as prolonged as the aforementioned phase.
323 In comparison to VPD, we identified around 10 further but shorter time periods in which VPD and R_s were correlated
324 negatively.

Figure 2: Extract of the timeseries from May – June 2023 for mean and median for soil respiration (R_s), mean soil temperature (T_s), mean soil water content (Θ_S), daily summed precipitation ($\sum prec.$), mean daily air temperature (T_a) and mean daily vapour pressure deficit (VPD).

Regression models with meteorological and microclimatic parameters

To explain the temporal variability, we compared three models using a linear and exponential fits. All models based
on the daily average of R_s from all 35 plots and the abiotic predictors (Table 5). All correlation coefficients between
the predictors are listed in the supplemental materials (Figure S1).

Table 5: Explanation of variability (R^2) of averaged daily R_s for climate variables as predictors with p-value determined with linear
models and spearman correlation coefficients (ρ_s) between R_s and climate variables over the two measurement years. Θ_S^* only
for the year 2023. Note: values were round to two decimal digits.

Predictor	R^2	p-value	ρ_s	Aggregated predictors	R^2	p-value	ρ_s
VP_{cur}	0.73	***	0.88	7-day mean	0.78	***	0.91
T_s	0.68	***	0.86	-	-	-	-
T_a	0.57	***	0.79	7-day mean	0.64	***	0.83
$\cos_{doy}$	0.50	***	0.73	-	-	-	-
Global radiation	0.31	***	0.61	7-day sum	0.32	***	0.62
Θ_S^*	0.20	*	-0.48	-	-	-	-
VPD	0.17	***	0.59	7-day mean	0.30	***	0.66
$\sin_{doy}$	0.16	***	0.42	-	-	-	-
Wind-velocity	0.08	***	-0.25	7-day sum	0.22	***	-0.54
H_{rel}	0.01	/	-0.31	7-day mean	0.03	/	-0.25
Precipitation	-0.01	/	0.16	7-day sum	0.02	/	0.19

In the following we present three models:

1. *Meteorological model*: The highest contribution to explain the variation of R_s is represented by the 7-day-
mean VP_{cur} (Table 5). We did not include T_s into our model due to high co-linearity with the VP_{cur} and the 7-
day mean VP_{cur} which created variance inflation factor (VIF) > 3. T_a is already combined in the VP_{cur} and does
not need to be included as a separate predictor. As the next highest predictor, we included the $\cos_{doy}$ which
resulted in VIF-values of < 3 ($R^2 = 0.78$). However, residuals of this model contained high information content
and were systematically explainable in all seasons by positive relationship with VPD (Supplemental material

Figure S3). We therefore needed to introduce a new predictor, which were the residuals the model,
explaining the variation of VPD by using the VP_{cur} (R^2 = 0.75). The residuals of this model represent low, wet
VPD as negative value and high, dry VPD as positive value and represents therefore the humidity of the
atmosphere. We could not include the original VPD due to high co-linearity between VP_{cur} and VPD. Including
the new predictor increased explanation in variability by 6 % (R^2 = 0.84) with VIF-values < 3.

2. *Exponential T_s model*: The exponential model fit using T_s as a predictor (Equation 3) described a similar
proportion in variability (R^2 = 0.68) as the linear fitted model (Table 5). However, comparisons on aggregated
values prior to the measurement day were not possible as T_s was only captured during R_s-measurement.
Exponential T_s and Θ_S model: This model could only be used for the 2023 due to missing Θ_S in 2024. For
comparison, the prescribed models were also fitted only for 2023. The exponential fit of T_s and Θ_S (Equation
4) resulted in a lower explanation in variability (R^2 = 0.72) than the model using the exponential fit of T_s
alone (Equation 3, R^2 = 0.78). A similar explanation in variability was represented by the *meteorological*
model (R^2 = 0.77). If we compare the models only for 2024, the exponential fit of T_s delivered the highest
explanation in variability (R^2 = 0.82). However, if we separate the models by years, the 7-day mean VP_{cur} still
delivers the highest explanation in variance in 2024 (R^2 = 0.89) but lower explanation in variance in 2023
(R^2 = 0.69).

Application of the *meteorological model*

For the further analyses we focus on the best model, the meteorological (Equation 6) model explaining 87% of the
total variability. This model left lower residuals compared to both other models using T_s and Θ_S as predictors
especially during "hot moments" such as dry and hot weeks in 2023 (Supplemental Material Figure S4). As can be
seen in Figure 3 (B) the model follows generally the measured values well, but smoothes the to a certain degree the
data. This is shown in the boxplots of the model residuals (Figure 3, B) which contain still a seasonal pattern with a
systematic overestimation in spring and autumn and an underestimation in summer. These seasonal effects could
obviously not be covered by the seasonal term ($\cos_{doy}$) in the model. Due to heteroscedasticity, we calculated robust
standard errors (Supplemental Material, Table S1).

$$R_s = -0.69 + (3.24 * 7d.VP_{cur}) + (0.7 * \cos_{doy}) + (-2.11 * res_{VPD}) \tag{6}$$

366 With R_s in µmol CO_2 m^{-2} s^{-1}, 7d. VP_{cur} as 7-day mean VP_{cur}, $\cos_{doy}$ as cosine of the day of the year and res_{VPD} as residuals
367 from model explaining the VPD using VP_{cur}.

Figure 3: Temporal model (A) for R_s [µmol CO_2 m^{-2} s^{-1}] for both measurement years in comparison to measured R_s (daily averaged) and residuals of the model (B) separated by year.

Spatial predictors – hot spots of R_s and the influence of forest structure

The comparison of plots at the individual plot level, separated by season, shows the development over the two years
per season (Figure 4, A) and illustrates a spatial pattern of R_s within the ECOSENSE Forest. Analysing the hot spots
and cold spots (significantly higher or lower R_s) separated by season and tree species composition reveals particularly
one plot, which is always considered a hot spot, while all other hot and cold spots occur only in specific seasons
(Figure 4, A). The largest occurrence in hot spots was determined for the summer season within deciduous species
composition. The plot-wise deviation from the averaged R_s (over the whole measurement period) separated by
species composition and season (Figure 4, B) was significantly different between every season except for the
comparison between mixed composition during spring and autumn. In contrast, within a season, deviations from the
average R_s separated by species composition were only significantly different within the summer season. In general,

Figure 4: Spatial variability of soil respiration (R_s) over two measurement years (2023 - 2024) plotted on a part of the ECOSENSE-Forest area using inverse-distance-weighting separated by season (A) and distribution of average deviation from mean R_s separated by season and tree species composition (B).

Figure 5: Soil respiration (R_s) in dependence to distance to trees (A) and in dependence of tree species composition and season (B).

the deviation from the overall average R_s was less pronounced during the autumn season, with the slightest average
deviation for deciduous composition.

Due to the strong seasonal patterns (Table 4, Figure 4), we averaged R_s per plot per season and analysed the
dependence of R_s on stand structural features of the surrounding area. The R_s associated with tree compositions was
significantly different between every season (Figure 5, B), while within the single seasons, only during the
summertime, the R_s associated with deciduous trees were significantly different from the other two species
compositions. We detected a slight negative relation between average distance to trees within the buffer zone and
R_s, as well as with the nearest neighbouring tree, which was more pronounced during the vegetative seasons (spring,
summer, autumn), but not significant (Figure 5, A). A comparison to the average DBH of surrounding trees within the
5 m buffer zone showed a positive relationship between R_s rate and deciduous and mixed composition during the
summertime, while the relationship between R_s rate and DBH was negative for coniferous composition. In
comparison to tree height, we determined a negative relationship of tree height for deciduous and coniferous species
during the vegetative season, while mixed composition showed a positive relationship Explaining the variability in R_s
as average per year with average distance to trees or nearest neighbouring distance was only significant, if
seasonality was included as predictor, while seasonality explained the main part of the variability in R_s. The same
pattern was determined for the DBH and tree height as predictors (data not shown).

Discussion

Temporal patterns in R_s are determined by meteorological conditions prior to measurements

Our results of average R_s are similar to those reported from other European research areas (e.g. Ruehr et al. 2010;
Scapucci et al. 2025; Ruehr and Buchmann 2010; Søe and Buchmann 2005; Hereș et al. 2021; Ekblad et al. 2005;
Jochheim et al. 2022; Subke et al. 2003) with a strongly pronounced seasonality. We hypothesised, that R_s depends
on T_s but also on meteorological conditions prior to the measurement. A comparison of the three models presented,
clearly shows that the *meteorological model* we proposed performed best. The strong, linear relationship between
R_s and the VP_{cur} stands out and has not been described so far. The VP_{cur} reflects the current vapour pressure in the air
and combines T_a and H_{rel} in one variable. The positive relationship indicates the importance of available atmospheric
water for temporal R_s patterns. However, we must also point out a very high correlation between the VP_{cur} and T_s (ρ_s
= 0.92) and a high co-linearity between T_s and the VP_{cur} (R^2 = 84%). This could be a reason why we were able to replace
the usually used T_s by the VP_{cur} in our model. T_s might reflect a better predictor for R_s estimation in other research,
as reference depths were different. We measured T_s in 13 cm depth, while e.g. Moyano et al. (2008) found highest
correlation with T_s in 2 cm depth. Ruehr et al. (2010) suggested T_s measurements within 1-5 cm depth for lag
estimations. However, for depths of 10 cm an exceptionally high Q_{10} was found by Ruehr et al. (2010), which can not
be supported by our findings of Q_{10} values. The values determined in this study were similar to the Q_{10} determined
by Khomik et al. (2006) from 2 cm soil depth. We also calculated seasonally separated Q_{10} values (Khomik et al. 2006)
with the limitation that temperature differences of up to 10°C within one season are unrealistic. The higher Q_{10} values
during the wintertime but lower Q_{10} during the summertime suggest a shift from a temperature-controlled to a
climate/moisture-controlled regime. Also, the change in correlations between R_s and Θ_s and the summarized
precipitation (Table 4) underlines our suggestions, as correlation was higher during the spring and summer season,
while reversed for the other predictors. A similar change in controls in R_s was also suggested by Hereș et al. (2021).

As T_s was only measured in combination with R_s, we cannot make any more precise statements about lag-effects of
T_s on R_s, as e.g. Ruehr et al. (2010). However, we also point out to a strong correlation between T_s and global radiation
and the summarised global radiation (ρ_s = 0.67, ρ_s = 0.72, respective) and linear relationship in T_s (R^2 = 44% and
R^2 = 51%, respective) and therefore highlight the effect of global radiation on T_s development (Kuzyakov and
Gavrichkova 2010). For long-term available predictors in every case, the aggregated values prior to measurements
correlated highest with R_s and we thereby support the assumptions that lag-effects (Kuzyakov and Gavrichkova 2010;
Ekblad et al. 2005; Bowling et al. 2002) are crucial for R_s patterns on temporal scale.

Even though the *meteorological model* performed well, the residuals showed larger range during the vegetation
season. These differences could be driven by the activity of the ecosystem. During the wintertime, the ecosystem
stays dormant due to lower temperatures, short daylight and the absence of foliage of deciduous trees.
Photosynthetic activity is at rest except for coniferous evergreen trees (Hansen et al. 1997; Jochheim et al. 2022) and
due to lower T_s and low nutrient availability, microbial activity as well. Photosynthetic activity with its peak during

the summer drives R_{SA} and therefore the total R_s (Kuzyakov and Gavrichkova 2010; Tang et al. 2005a). Microbial
activity is also higher during the summer season, because enzymatic activity depends on T_S (Carey et al. 2016;
Schindlbacher et al. 2011; Schindlbacher et al. 2012) but also root on exudate availability, which in turn is driven by
photosynthesis (Kuzyakov and Gavrichkova 2010). The inverse or lower correlations with most of the presented
meteorological predictors (Table 4) further underline the importance of the phenology on temporal R_s patterns. Yet
for this research the solar radiation is the only available predictor as a proxy for photosynthetic activity, besides the
manual observation of leaf developmental stage. The latter resulted in a direct increase in R_s due to foliation and
decrease in R_s due to senescence (Figure 3, A). We implemented smooth seasonal shifts with the $\cos_{doy}$ in our model,
with the peak during the summertime. This still lacks the true representation of the phenology. The overestimation
during spring could be due to the fact that trees invest sugars into biomass at first (Kuzyakov and Gavrichkova 2010)
and therefore R_s does not follow the steady increase of the VP_{cur}. Furthermore, R_{SH} might be low, due to low
exudation of easily available substrates. However, our model shows during the summertime in 2023 a broader range
in residuals which can simply be attributed to the dry phase (hot moment), as during this phase, the model is not
capable in representing the high variability in responses to the rainfall event after the dry phase. This suggests a
strong spatial and individual response on plot-level. The widely applied models using T_s as the only predictor or in
combination with Θ_s, especially during the hot moment in R_s, did not deliver the same performance as our presented
model (Supplemental Material Figure S4). For models using T_s as predictor it was already shown, that available Θ_s is
a limiting factor (Ruehr et al. 2010).

Collapse of R_s and increase due to rewetting

In our second hypothesis we stated that R_s will collapse during drought periods and increases after rewetting. We
were able to clearly identify one time period as a hot moment. The usual pattern of positive correlation between T_s
and R_s was reversed due to a decline in Θ_s caused by a precipitation deficit. Rewetting led to an extreme increase in
R_s. This increase was also reported in the literature (Birch 1958; Tang et al. 2005b; Ruehr et al. 2010; Jochheim et al.
2022). However, this effect has not always been confirmed. Schindlbacher et al. (2012) observed a decline due to
drought and a regeneration of R_s due to rewetting, but not a peak. The authors attributed the lack of this effect to
the importance in homogeneous rewetting of the forest floor, which might not have occurred due to preferential
flow paths created by drought induced hydrophobicity (Schindlbacher et al. 2012). Overall, the data obtained confirm
our hypothesis and we suggest, that the strong increase was an effect of reactivation of microbial activity (R_{SH}) (Wang
et al. 2014a; Wang et al. 2014b; Ruehr et al. 2010; Jochheim et al. 2022; Birch 1958). Jochheim et al. (2022)
additionally suggested a potential collapse in R_{SA} associated with *Fagus sylvatica* due to a stronger decline associated
with these species, which cannot be supported by our data, as all plots were affected similarly. However, we cannot
exclude that R_{SA} was not affected by the drought period. The VPD can also be used as an estimate for aboveground
plant stress, as trees open or close their stomata when VPD is low or high, respective (McAdam and Brodribb 2015).
In comparison, the summer and autumn of 2023 showed a higher VPD than 2024, indicating drier aboveground
conditions, especially during the hot moment. The strongly differing correlations of VPD and R_s in the individual
seasons underline the VPD as an important factor for R_s patterns. Sanginés de Cárcer et al. (2018) showed the effects
of species specific tree growth reduction for VPD values above 1.5 kPa, decoupled from soil water limitations. This
value can be used as an approximate threshold at which the stomata of *Fagus sylvatica* (Lendzion and Leuschner
2008) and *Pseudotsuga menziesii* trees (Meinzer 1982) reduce conductance, which affects carbon uptake and
therefore also carbon allocation and respiration by tree roots (Hsiao 1973). There was only one day in the two years,
which represented a VPD > 1.5 kPa. This might be due to the type of calculation of VPD using the daily averaged T_a.
The VPD is therefore smoothed and represents only the average for the corresponding day, neglecting peaks. VPD
values > 1 kPa were more present during 2023 (23 days) compared to 2024 (4 days). If VPD is calculated using the
maximum T_a, values above 1.5 kPa were more common in 2023 (29 days) compared to 2024 (7 days) and the
aforementioned hot moment is particularly affected by the high potential. Similar to daily averaged VPD > 0.75 kPa
(or maximum VPD > 1 kPa) also leads to a negative relationship with R_s which is in contrast to the general captured
pattern of positive relationship. The general comparison between available dates with Θ_s measurements and daily
as well as averaged VPD prior to measurements also reveals a negative correlation (ρ_s = -0.69 and ρ_s = -0.71,
respectively), indicating drier soil conditions with increasing VPD. Interestingly, we detected a higher average R_s in
2024 compared to 2023, but the two highest R_s rates were reached in 2023, which again shows the precipitation

deficit within the ecosystem and reflects ecosystem stress within the soil. Our results highlight the importance of
high-resolution measurements to capture these specific hot moments as an interpolation of R_s during these phases
might lead to false conclusions.

Forest structural features and hot spots in the ECOSENSE Forest

In agreement with our hypotheses, we observed larger R_s for deciduous than for coniferous patches during summer.
However, tree species did not affect soil respiration in winter, which is in contrast to Jochheim et al. (2022) and our
hypothesis. However, a generally higher R_s for broadleaf species compared to coniferous species were also reported
in the literature (Raich and Tufekciogul 2000). Longdoz et al. (2000) also showed differences in R_s associated especially
with *Fagus sylvatica* and *Pseudotsuga menziesii* patches in a slightly different climate. The authors presented various
reasons of which they underlines differences in fresh leaf litter and lower C:N and lignin:N as indicators for
decomposition. A further reason might be the difference in transport processes through phloem, which was
summarized by Kuzyakov and Gavrichkova (2010) to be lower for coniferous species compared to angiosperms, while
the latter includes *Fagus sylvatica*. The differences in tree species also seem to reflect the spatial pattern in the
ECOSENSE Forest, as abundance of *Fagus sylvatica* is higher in the northern part of the area. The pattern was still
homogeneous which is in comprehension to research focusing on spatial patterns of R_s, (Knohl et al. 2008; Hereş et
al. 2021). Especially during the summer season we detected a larger variability compared to the winter season
(Khomik et al. 2006). Regarding the strong heterogeneous and seasonal pattern, surprisingly, the deviation in R_s
during autumn season from the average over the two-year measurement period was lowest, which would suggest
that R_s during autumn could reflect the average R_s best. However, this might also be a specific effect of our research
area.

We also point out to a slightly higher diffusivity of the soil associated with the soil profile in the northern area.
Differences between the soil types in regard to the other specified physical and chemical properties are not too
different, except for a slightly higher carbon content in the topsoil horizons, as well as the CEC and the phosphorous
content. However, soil properties could be very heterogeneous throughout forest soils (Stutz and Lang 2023) and a
clear border within the ECOSENSE Forest between the two existing soil types is yet to be determined. A transition
zone could be located approximately in the middle of the area. Due to rather similar chemical and physical properties
of the described soil profiles, we do not expect larger influences of soil properties on R_s Further biological analysis as
root distribution and microbial biomass, chemical and physical properties of each of the 35 plots within the ECOSENSE
Forest, might underline our ideas that tree species and their associated micro-habitat (Köstler et al. 1968; Hofmann
et al. 2023; Prescott and Grayston 2013; Spielvogel et al. 2016; Vesterdal 1999; Vesterdal et al. 2013) shape R_s
(Brzozon et al., in preparation).

The positive relationship with the average DBH for deciduous composition during the summertime also reflects the
vegetative phase of trees also found by Søe and Buchmann (2005) within a distance of 4 m, while the negative effect
associated with coniferous composition might be biased, as the relationship between the distance to the tree and
the DBH is strongly positive for coniferous composition. The relation was also positive for mixed and deciduous
composition but not as strong as for coniferous composition. This could explain the negative effect of coniferous
composition with increasing DBH, as for the average distance to trees we also showed the negative relationship.
Surprisingly the tree height showed negative relationship with R_s which is one the one hand again biased as trees in
coniferous compositions were averaging higher compared to trees in deciduous composition. However, as trends
within the compositions (deciduous, coniferous) were also negative, we also suggest a negative effect of tree height
on R_s despite the bias between the composed groups. To our knowledge no effects of tree height have been stated
in the literature until now.

Considerations on the sensor systems and spatial analysis

The system we used for measurements was a manual system and errors occurred in some cases (R_s measurements
in Material and Methods). However, errors were only on a small percentage and we do not expect a large difference.
Our analysis showed a small-scale heterogeneity in our forest, and therefore if the three measurement days
highlighted in the section did not include hot-spots or cold spots of the forest area, this would skew the average value

of the day upwards or downwards, respectively. Out of the three days that did not include all plots within one
measurement day from June 2024, one day was during winter (December, 26/35), one in Spring (May 26/35) and
one in autumn (September, 33/35). Since spatial variability was highest during summer, we do not assume largely
distorted averages during winter but we must stress, that mean values from the two days during vegetative season
might be distorted, as some plots that built hot spots were omitted during these days. This in any case reflects a
dilemma, as heterogeneity was found on small scale, therefore a high spatial resolution is needed, but might be
prone to weather conditions or device error. We must also note that four measurement days during the winter
months in 2023 and April 2023 covered less than 20/35 plots. Either way, here we also do not expect large distortions
of the averaged value, as these measurements primarily covered the winter season.

Conclusion:

Our results highlight the strong dependency of R_s on meteorological conditions. However, our presented model
outperformed the usually applied models containing T_s and Θ_S emphasizing the high relevance of VP_{cur}. We showed
that especially during the vegetative season important phenology factors are required to explain R_s. The spatial
pattern could be partly explained by species composition with significantly higher R_s associated with deciduous
composition during the summer season, which was in our case always a pure composition of *Fagus sylvatica* within
5 m around the measurement location. Yet we are not able to fully explain the spatial pattern within our research
site, maybe soil chemical, biological and physical properties will mediate tree species effects. However, with our long-
term weekly- bi-weekly measurements on large spatial resolution we were able to capture a hot moment in R_s and
presented the immediate effect of dry conditions and rewetting on R_s. It is important to consider that hot and cold
spots in our studied area were not constantly present and rather changed throughout seasons, with higher or lower
contributions to overall seasonal R_s. The high seasonality highlights the importance of high-resolution measurements
covering the seasonality of a year, as not only on spatial scale but also on temporal scale measurements at hot spots,
cold spots or during hot moments could falsify conclusions made using extrapolation.

CRediT authorshop contribution statement

Julian Brzozon: Writing – original draft, Investigation, Methodology, Formal analysis. Patricia Schwarzkopf: Writing –
review & editing, Investigation. Friederike Lang: Writing – review & editing, Funding acquisition. Teja Kattenborn:
Writing – review & editing, Investigation, Methodology, Formal analysis. Julian Frey: Writing – review & editing,
Investigation, Methodology, Formal analysis. Helmer Schack-Kirchner: Writing – review & editing, Methodology,
Conceptualization.

Funding

Financial Support: This project was Funded by the Deutsche Forschungsgemeinschaft (DFG) - Project-ID 459819582 -
SFB 1537.

Declaration of competing interest

The authors declare that they have no known competing financial interest or personal relationships that could have
appeared to influence the work reported in this paper.

Acknowledgements

We thank the laboratory team of the chair for soil ecology for supporting the laboratory analysis. Furthermore, we
want to thank Luna Reimer, Chris Huck and Fiona Bauer for their support during the start of the measurement
campaign. We also thank the team members of the Chair of Sensor-based Geoinformatics and the Chair of Forest
Growth and Dendroecology who worked on the forest tree inventory and helped in field, digitizing and validating the
dataset.

Publication bibliography

Barba, Josep; Cueva, Alejandro; Bahn, Michael; Barron-Gafford, Greg A.; Bond-Lamberty, Benjamin; Hanson, Paul J. et al. (2018): Comparing ecosystem and soil respiration: Review and key challenges of tower-based and soil measurements. In *Agricultural and Forest Meteorology* 249, pp. 434–443. DOI: 10.1016/j.agrformet.2017.10.028.

Bates, Douglas; Maechler, Martin; Bolker, Ben; Walker, Steve (2015): Fitting Linear Mixed-Effects Models Using lme4 (67(1)), pp. 1–48. DOI: 10.18637/jss.v067.i01.

Birch, H. F. (1958): The effect of soil drying on humus decomposition and nitrogen availability. In *Plant and Soil* 10 (1), pp. 9–31. DOI: 10.1007/BF01343734.

Bowling, David R.; McDowell, Nate G.; Bond, Barbara J.; Law, Beverly E.; Ehleringer, James R. (2002): 13C content of ecosystem respiration is linked to precipitation and vapor pressure deficit. In *Oecologia* 131 (1), pp. 113–124. DOI: 10.1007/s00442-001-0851-y.

Brendel, Christoph; Rauthe, Monika; Deutschländer, Thomas (2023): The new-high resolution HYRAS global radiation dataset for Germany and its river catchments. Copernicus Meetings (Copernicus Meetings, EMS2023-295). Available online at https://meetingorganizer.copernicus.org/ems2023/ems2023-295.html.

Brunn, Melanie; Hafner, Benjamin D.; Zwetsloot, Marie J.; Weikl, Fabian; Pritsch, Karin; Hikino, Kyohsuke et al. (2022): Carbon allocation to root exudates is maintained in mature temperate tree species under drought. In *New Phytologist* 235 (3), pp. 965–977. DOI: 10.1111/nph.18157.

Brunn, Melanie; Krüger, Jaane; Lang, Friederike (2023): Experimental drought increased the belowground sink strength towards higher topsoil organic carbon stocks in a temperate mature forest. In *Geoderma* 431, p. 116356. DOI: 10.1016/j.geoderma.2023.116356.

Brunner, Ivano; Herzog, Claude; Dawes, Melissa A.; Arend, Matthias; Sperisen, Christoph (2015): How tree roots respond to drought. In *Front. Plant Sci.* 6, p. 547. DOI: 10.3389/fpls.2015.00547.

Carey, Joanna C.; Tang, Jianwu; Templer, Pamela H.; Kroeger, Kevin D.; Crowther, Thomas W.; Burton, Andrew J. et al. (2016): Temperature response of soil respiration largely unaltered with experimental warming. In *Proceedings of the National Academy of Sciences of the United States of America* 113 (48), pp. 13797–13802. DOI: 10.1073/pnas.1605365113.

Copernicus Climate Change Service (2019): ERA5-Land hourly data from 1950 to present.

Curiel Yuste, J.; Janssens, I. A.; Carrara, A.; Ceulemans, R. (2004): Annual Q 10 of soil respiration reflects plant phenological patterns as well as temperature sensitivity. In *Global change biology* 10 (2), pp. 161–169. DOI: 10.1111/j.1529-8817.2003.00727.x.

Darenova, Eva; Adamič, Pia Caroline; Čater, Matjaž (2024): Effect of temperature, water availability, and soil properties on soil CO2 efflux in beech-fir forests along the Carpathian Mts. In *CATENA* 240, p. 107974. DOI: 10.1016/j.catena.2024.107974.

Davidson, E. A.; Richardson, A. D.; Savage, K. E.; Hollinger, D. Y. (2006): A distinct seasonal pattern of the ratio of soil respiration to total ecosystem respiration in a spruce-dominated forest. In *Global change biology* 12 (2), pp. 230–239. DOI: 10.1111/j.1365-2486.2005.01062.x.

Davidson, EriC. A.; Belk, Elizabeth; Boone, Richard D. (1998): Soil water content and temperature as
independent or confounded factors controlling soil respiration in a temperate mixed hardwood forest. In
Global change biology 4 (2), pp. 217–227. DOI: 10.1046/j.1365-2486.1998.00128.x.

DWD (2025): Index of /climate_environment/CDC/grids_germany/daily/hyras_de. Available online at
https://opendata.dwd.de/climate_environment/CDC/grids_germany/daily/hyras_de/, updated on
6/10/2025, checked on 6/10/2025.

Ekblad, A.; Högberg, P. (2001): Natural abundance of 13C in CO2 respired from forest soils reveals speed
of link between tree photosynthesis and root respiration. In *Oecologia* 127 (3), pp. 305–308. DOI:
10.1007/s004420100667.

Ekblad, Alf; Boström, Björn; Holm, Anders; Comstedt, Daniel (2005): Forest soil respiration rate and
delta13C is regulated by recent above ground weather conditions. In *Oecologia* 143 (1), pp. 136–142.
DOI: 10.1007/s00442-004-1776-z.

Epron, Daniel; Farque, Lætitia; Lucot, Éric; Badot, Pierre-Marie (1999): Soil CO2 efflux in a beech forest:
dependence on soil temperature and soil water content. In *Ann. For. Sci.* 56 (3), pp. 221–226. DOI:
10.1051/forest:19990304.

Fiedler, Jan; Fuß, Roland; Glatzel, Stephan; Hagemann, Ulrike; Huth, Vytas; Jordan, Sabine et al. (2022):
BEST PRACTICE GUIDELINE Measurement of carbon dioxide, methane and nitrous oxide fluxes between
soil-vegetation-systems and the atmosphere using non-steady state chambers. With assistance of FID
GEO: Arbeitsgruppe Bodengase, Deutsche Bodenkundliche Gesellschaft.

Frick, Claudia; Steiner, Heiko; Mazurkiewicz, Alex; Riediger, Ulf; Rauthe, Monika; Reich, Thomas; Gratzki,
Annegret (2014): Central European high-resolution gridded daily data sets (HYRAS): Mean temperature
and relative humidity. In *metz* 23 (1), pp. 15–32. DOI: 10.1127/0941-2948/2014/0560.

Goulden, Michael L.; Munger, J. William; Fan, Song-Miao; Daube, Bruce C.; Wofsy, Steven C. (1996):
Exchange of Carbon Dioxide by a Deciduous Forest: Response to Interannual Climate Variability. In
Science 271 (5255), pp. 1576–1578. DOI: 10.1126/science.271.5255.1576.

Grolemund, Garrett; Wickham, Hadley (2011): Dates and Times Made Easy with lubridate. In *J. Stat. Soft.*
40 (3), pp. 1–25. DOI: 10.18637/jss.v040.i03.

Gutachterausschuss Forstliche Analytik (2022): Handbuch Forstliche Analytik: Eine Loseblatt-Sammlung
der Analysemethoden im Forstbereich. Bundesministerium für Verbraucherschutz, Ernährung und
Landwirtschaft, Bonn. Base edition, June 2005, with supplements, 6th edition (February 2022). Available
online at https://blumwald.thuenen.de/fileadmin/blumwald/BZE/HFA_Gesamtdatei_2022.pdf, checked
on 6/17/2025.

Hansen, Jens; Turkington, Roy; Vogg, Gerd; Heim, Richard; Beck, Erwin (1997): Conifer carbohydrate
physiology : Updating classical views. Leiden: Backhuys (Trees - contributions to modern tree
physiology). Available online at https://eref.uni-bayreuth.de/id/eprint/20731.

Hartge, Karl Heinrich; Horn, Rainer (1989): Die physikalische Untersuchung von Böden. Stuttgart,
Germany: Enke (Vol. 2). Available online at https://library.wur.nl/webquery/titel/505652.

Hayes, Andrew F.; Cai, Li (2007): Using heteroskedasticity-consistent standard error estimators in OLS
regression: an introduction and software implementation. In *Behavior research methods* 39 (4), pp. 709–
722. DOI: 10.3758/bf03192961.

Hereș, Ana-Maria; Bragă, Cosmin; Petritan, Any Mary; Petritan, Ion Catalin; Curiel Yuste, Jorge (2021):
Spatial variability of soil respiration (R s) and its controls are subjected to strong seasonality in an even-
aged European beech (Fagus sylvatica L.) stand. In *European J Soil Science* 72 (5), pp. 1988–2005. DOI:
10.1111/ejss.13116.

Hofmann, Benjamin; Dreyling, Lukas; Dal Grande, Francesco; Otte, Jürgen; Schmitt, Imke (2023): Habitat
and tree species identity shape aboveground and belowground fungal communities in central European
forests. In *Front. Microbiol.* 14, p. 1067906. DOI: 10.3389/fmicb.2023.1067906.

Hsiao, Theodore C. (1973): Plant Responses to Water Stress. In *Annu. Rev. Plant. Physiol.* 24 (1), pp. 519–
570. DOI: 10.1146/annurev.pp.24.060173.002511.

IPCC (2023): Climate Change 2022 – Impacts, Adaptation and Vulnerability: Working Group II
Contribution to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change:
Cambridge University Press.

IUSS (2022): World reference base for soil resources 2022 : International soil classification system for
naming soils and creating legends for soil maps / IUSS Working Group WRB. With assistance of IUSS.
4.edition. Vienna, Austria: International Union of Soil Sciences.

Janssens, Ivan A.; Pilegaard, Kim (2003): Large seasonal changes in Q10 of soil respiration in a beech
forest. In *Global change biology* 9 (6), pp. 911–918. DOI: 10.1046/j.1365-2486.2003.00636.x.

Jochheim, Hubert; Wirth, Stephan; Gartiser, Valentin; Paulus, Sinikka; Haas, Christoph; Gerke, Horst H.;
Maier, Martin (2022): Dynamics of Soil CO2 Efflux and Vertical CO2 Production in a European Beech and
a Scots Pine Forest. In *Front. For. Glob. Change* 5, Article 826298, p. 826298. DOI:
10.3389/ffgc.2022.826298.

Jovani-Sancho, A. Jonay; Cummins, Thomas; Byrne, Kenneth A. (2017): Collar insertion depth effects on
soil respiration in afforested peatlands. In *Biol Fertil Soils* 53 (6), pp. 677–689. DOI: 10.1007/s00374-017-
1210-4.

Khomik, Myroslava; Arain, M. Altaf; McCaughey, J. H. (2006): Temporal and spatial variability of soil
respiration in a boreal mixedwood forest. In *Agricultural and Forest Meteorology* 140 (1-4), pp. 244–256.
DOI: 10.1016/j.agrformet.2006.08.006.

Knohl, Alexander; Søe, Astrid R. B.; Kutsch, Werner L.; Göckede, Mathias; Buchmann, Nina (2008):
Representative estimates of soil and ecosystem respiration in an old beech forest. In *Plant and Soil* 302
(1-2), pp. 189–202. DOI: 10.1007/s11104-007-9467-2.

König, Nils; Fortmann, Heike (1996): Berichte des Forschungszentrum Waldökosysteme. Reihe B, Bd. 49.

Köstler, Josef Nikolaus; Brückner, Ernst; Bibelriether, Hans (1968): Die Wurzeln der Waldbäume.
Untersuchungen zur Morphologie der Waldbäume in Mitteleuropa. Germany: Verlag Paul Parey,
Hamburg and Berlin.

Kuzyakov, Y. (2006): Sources of CO2 efflux from soil and review of partitioning methods. In *Soil Biology*
and Biochemistry 38 (3), pp. 425–448. DOI: 10.1016/j.soilbio.2005.08.020.

Kuzyakov, Yakov; Gavrichkova, Olga (2010): REVIEW: Time lag between photosynthesis and carbon
dioxide efflux from soil: a review of mechanisms and controls. In *Global change biology* 16 (12),
pp. 3386–3406. DOI: 10.1111/j.1365-2486.2010.02179.x.

Lejon, David P. H.; Chaussod, Rémi; Ranger, Jacques; Ranjard, Lionel (2005): Microbial community
structure and density under different tree species in an acid forest soil (Morvan, France). In *Microbial*
ecology 50 (4), pp. 614–625. DOI: 10.1007/s00248-005-5130-8.

Lendzion, Jasmin; Leuschner, Christoph (2008): Growth of European beech (Fagus sylvatica L.) saplings is
limited by elevated atmospheric vapour pressure deficits. In *Forest Ecology and Management* 256 (4),
pp. 648–655. DOI: 10.1016/j.foreco.2008.05.008.

Lenth, Russel V. (2023): emmeans: Estimated Marginal Means, aka Least-Squares Means. Version R
package version 1.8.5.

Lloyd, J.; Taylor, J. A. (1994): On the Temperature Dependence of Soil Respiration. In *Functional Ecology*
8 (3), p. 315. DOI: 10.2307/2389824.

Longdoz, B.; Yernaux, M.; Aubinet, M. (2000): Soil CO 2 efflux measurements in a mixed forest: impact of
chamber disturbances, spatial variability and seasonal evolution. In *Global change biology* 6 (8), pp. 907–
917. DOI: 10.1046/j.1365-2486.2000.00369.x.

McAdam, Scott A. M.; Brodribb, Timothy J. (2015): The evolution of mechanisms driving the stomatal
response to vapor pressure deficit. In *Plant Physiol* 167 (3), pp. 833–843. DOI: 10.1104/pp.114.252940.

McClain, Michael E.; Boyer, Elizabeth W.; Dent, C. Lisa; Gergel, Sarah E.; Grimm, Nancy B.; Groffman,
Peter M. et al. (2003): Biogeochemical Hot Spots and Hot Moments at the Interface of Terrestrial and
Aquatic Ecosystems. In *Ecosystems* 6 (4), pp. 301–312. DOI: 10.1007/s10021-003-0161-9.

Meinzer, Frederick C. (1982): The effect of vapor pressure on stomatal control of gas exchange in
Douglas fir (Pseudotsuga menziesii) saplings. In *Oecologia* 54 (2), pp. 236–242. DOI:
10.1007/BF00378398.

Moyano, Fernando E.; Kutsch, Werner L.; Rebmann, Corinna (2008): Soil respiration fluxes in relation to
photosynthetic activity in broad-leaf and needle-leaf forest stands. In *Agricultural and Forest*
Meteorology 148 (1), pp. 135–143. DOI: 10.1016/j.agrformet.2007.09.006.

Muñoz-Sabater, Joaquín; Dutra, Emanuel; Agustí-Panareda, Anna; Albergel, Clément; Arduini, Gabriele;
Balsamo, Gianpaolo et al. (2021): ERA5-Land: a state-of-the-art global reanalysis dataset for land
applications. In *Earth Syst. Sci. Data* 13 (9), pp. 4349–4383. DOI: 10.5194/essd-13-4349-2021.

Pebesma, Edzer (2018): Simple Features for R: Standardized Support for Spatial Vector Data. In *The R*
Journal 10 (1), p. 439. DOI: 10.32614/rj-2018-009.

Pebesma, Edzer J.; Bivand, Roger (2023): Spatial data science. With applications in R. First edition. Boca
Raton: CRC Press Taylor & Francis Group (Chapman & Hall/CRC Press the R series). Available online at
https://www.taylorfrancis.com/books/mono/10.1201/9780429459016/spatial-data-science-edzer-
pebesma-roger-bivand.

Prescott, Cindy E.; Grayston, Sue J. (2013): Tree species influence on microbial communities in litter and
soil: Current knowledge and research needs. In *Forest Ecology and Management* 309, pp. 19–27. DOI:
10.1016/j.foreco.2013.02.034.

Qi, Ye; Xu, Ming; Wu, Jianguo (2002): Temperature sensitivity of soil respiration and its effects on
ecosystem carbon budget: nonlinearity begets surprises. In *Ecological Modelling* 153 (1-2), pp. 131–142.
DOI: 10.1016/S0304-3800(01)00506-3.

R Core Team (2023): R: A Language and Environment for Statistical Computing. Version 4.2.3: R
Foundation for Statistical Computing, Vienna, Austria. Available online at https://www.R-project.org/.

Raich, James W.; Tufekciogul, Aydin (2000): Vegetation and soil respiration: Correlations and controls. In
Biogeochemistry 48 (1), pp. 71–90. DOI: 10.1023/A:1006112000616.

Rauthe, Monika; Steiner, Heiko; Riediger, Ulf; Mazurkiewicz, Alex; Gratzki, Annegret (2013): A Central
European precipitation climatology – Part I: Generation and validation of a high-resolution gridded daily
data set (HYRAS). In *metz* 22 (3), pp. 235–256. DOI: 10.1127/0941-2948/2013/0436.

Razafimaharo, Christène; Krähenmann, Stefan; Höpp, Simona; Rauthe, Monika; Deutschländer, Thomas
(2020): New high-resolution gridded dataset of daily mean, minimum, and maximum temperature and
relative humidity for Central Europe (HYRAS). In *Theor Appl Climatol* 142 (3-4), pp. 1531–1553. DOI:
10.1007/s00704-020-03388-w.

Ruehr, Nadine K.; Buchmann, Nina (2010): Soil respiration fluxes in a temperate mixed forest: seasonality
and temperature sensitivities differ among microbial and root-rhizosphere respiration. In *Tree physiology*
30 (2), pp. 165–176. DOI: 10.1093/treephys/tpp106.

Ruehr, Nadine K.; Knohl, Alexander; Buchmann, Nina (2010): Environmental variables controlling soil
respiration on diurnal, seasonal and annual time-scales in a mixed mountain forest in Switzerland. In
Biogeochemistry 98 (1-3), pp. 153–170. DOI: 10.1007/s10533-009-9383-z.

Sanginés de Cárcer, Paula; Vitasse, Yann; Peñuelas, Josep; Jassey, Vincent E. J.; Buttler, Alexandre;
Signarbieux, Constant (2018): Vapor-pressure deficit and extreme climatic variables limit tree growth. In
Global change biology 24 (3), pp. 1108–1122. DOI: 10.1111/gcb.13973.

Scapucci, Liliana; Krebs, Luana; Burri, Susanne; Hörtnagl, Lukas; Buchmann, Nina (2025): Soil and forest
floor respiration already acclimated to increasing temperatures in a mixed deciduous forest. In *Ecological*
processes 14 (1), p. 71. DOI: 10.1186/s13717-025-00639-4.

Schack-Kirchner, Helmer; Gaertig, Thorsten; v. Wilpert, Klaus; Hildebrand, Ernst E. (2001): A modified
McIntyre and Phillip approach to measure top-soil gas diffusivity in-situ. In *Journal of Plant Nutrition and*
Soil Science 164 (3), pp. 253–258. DOI: 10.1002/1522-2624(200106)164:3<253::AID-JPLN253>3.0.CO;2-G.

Schindlbacher, A.; Rodler, A.; Kuffner, M.; Kitzler, B.; Sessitsch, A.; Zechmeister-Boltenstern, S. (2011):
Experimental warming effects on the microbial community of a temperate mountain forest soil. In *Soil*
Biology and Biochemistry 43 (7), pp. 1417–1425. DOI: 10.1016/j.soilbio.2011.03.005.

Schindlbacher, Andreas; Wunderlich, Steve; Borken, Werner; Kitzler, Barbara; Zechmeister-Boltenstern,
Sophie; Jandl, Robert (2012): Soil respiration under climate change: prolonged summer drought offsets
soil warming effects. In *Global change biology* 18 (7), pp. 2270–2279. DOI: 10.1111/j.1365-
2486.2012.02696.x.

Schwertmann, U. (1964): Differenzierung der Eisenoxide des Bodens durch Extraktion mit
Ammoniumoxalat-Lösung. In *Journal of Plant Nutrition and Soil Science* 105 (3), pp. 194–202. DOI:
10.1002/jpln.3591050303.

Scott-Denton, L. (2003): Spatial and temporal controls of soil respiration rate in a high-elevation,
subalpine forest. In *Soil Biology and Biochemistry* 35 (4), pp. 525–534. DOI: 10.1016/S0038-
0717(03)00007-5.

Sims, Christopher A. (1974): Seasonality in Regression. In *Journal of the American Statistical Association*
69 (347), pp. 618–626. DOI: 10.1080/01621459.1974.10480178.

Snell, Helen S. K.; Robinson, David; Midwood, Andrew J. (2014): Minimising methodological biases to
improve the accuracy of partitioning soil respiration using natural abundance 13C. In *Rapid*
Communications in Mass Spectrometry 28 (21), pp. 2341–2351. DOI: 10.1002/rcm.7017.

Søe, Astrid R. B.; Buchmann, Nina (2005): Spatial and temporal variations in soil respiration in relation to
stand structure and soil parameters in an unmanaged beech forest. In *Tree physiology* 25 (11), pp. 1427–
1436. DOI: 10.1093/treephys/25.11.1427.

Spielvogel, Sandra; Prietzel, Jörg; Kögel-Knabner, Ingrid (2016): Stand scale variability of topsoil organic
matter composition in a high-elevation Norway spruce forest ecosystem. In *Geoderma* 267, pp. 112–122.
DOI: 10.1016/j.geoderma.2015.12.001.

Štursová, Martina; Bárta, Jiří; Šantrůčková, Hana; Baldrian, Petr (2016): Small-scale spatial heterogeneity
of ecosystem properties, microbial community composition and microbial activities in a temperate
mountain forest soil. In *FEMS microbiology ecology* 92 (12). DOI: 10.1093/femsec/fiw185.

Stutz, Kenton P.; Lang, Friederike (2023): Forest ecosystems create pedogenic patchworks through
woody debris, trees, and disturbance. In *Geoderma* 429, p. 116246. DOI:
10.1016/j.geoderma.2022.116246.

Subke, Jens-Arne; Reichstein, Markus; Tenhunen, John D. (2003): Explaining temporal variation in soil
CO2 efflux in a mature spruce forest in Southern Germany. In *Soil Biology and Biochemistry* 35 (11),
pp. 1467–1483. DOI: 10.1016/S0038-0717(03)00241-4.

Tamm, Olof (1922): Om bestämning av de oorganiska komponenterna i markens gelkomplex. Available
online at https://pub.epsilon.slu.se/id/eprint/10087/contents.

Tang, Jianwu; Baldocchi, Dennis D.; Xu, Liukang (2005a): Tree photosynthesis modulates soil respiration
on a diurnal time scale. In *Global change biology* 11 (8), pp. 1298–1304. DOI: 10.1111/j.1365-
2486.2005.00978.x.

Tang, Jianwu; Misson, Laurent; Gershenson, Alexander; Cheng, Weixin; Goldstein, Allen H. (2005b):
Continuous measurements of soil respiration with and without roots in a ponderosa pine plantation in
the Sierra Nevada Mountains. In *Agricultural and Forest Meteorology* 132 (3-4), pp. 212–227. DOI:
10.1016/j.agrformet.2005.07.011.

Tarroso, P.; Velo-Anton, G.; Carvalho, S. B. (2015): PHYLIN: an R package for phylogeographic
interpolation. 15(2), pp. 349–357.

Tarroso, Pedro; Carvalho, Sílvia B.; Velo-Antón, Guillermo (2019): Phylin 2.0: Extending the
phylogeographical interpolation method to include uncertainty and user-defined distance metrics. In
Molecular Ecology Resources 19 (4), pp. 1081–1094. DOI: 10.1111/1755-0998.13010.

Tesch, Jasmin; Kühnhammer, Kathrin; Wagner, Delon; Christen, Andreas; Dormann, Carsten; Frey, Julian
et al. (2025): The ECOSENSE forest: A distributed sensor and data management system for real-time
monitoring of ecosystem processes and stresses.

Trüby, Peter; Aldinger, Eberhard (1989): Eine Methode zur Bestimmung austauschbarer Kationen in
Waldböden. In *J Plant Nutrition & Soil* 152 (3), pp. 301–306. DOI: 10.1002/jpln.19891520307.

Valentini, R.; Matteucci, G.; Dolman, A. J.; Schulze, E. D.; Rebmann, C.; Moors, E. J. et al. (2000):
Respiration as the main determinant of carbon balance in European forests. In *Nature* 404 (6780),
pp. 861–865. DOI: 10.1038/35009084.

Vesterdal, Lars (1999): Influence of soil type on mass loss and nutrient release from decomposing foliage
litter of beech and Norway spruce. In *Can. J. For. Res.* 29 (1), pp. 95–105. DOI: 10.1139/x98-182.

Vesterdal, Lars; Clarke, Nicholas; Sigurdsson, Bjarni D.; Gundersen, Per (2013): Do tree species influence
soil carbon stocks in temperate and boreal forests? In *Forest Ecology and Management* 309, pp. 4–18.
DOI: 10.1016/j.foreco.2013.01.017.

Vesterdal, Lars; Elberling, Bo; Christiansen, Jesper Riis; Callesen, Ingeborg; Schmidt, Inger Kappel (2012):
Soil respiration and rates of soil carbon turnover differ among six common European tree species. In
Forest Ecology and Management 264, pp. 185–196. DOI: 10.1016/j.foreco.2011.10.009.

Wang, Xin; Liu, Lingli; Piao, Shilong; Janssens, Ivan A.; Tang, Jianwu; Liu, Weixing et al. (2014a): Soil
respiration under climate warming: differential response of heterotrophic and autotrophic respiration. In
Global change biology 20 (10), pp. 3229–3237. DOI: 10.1111/gcb.12620.

Wang, Yanfen; Hao, Yanbin; Cui, Xiao Yong; Zhao, Haitao; Xu, Chengyuan; Zhou, Xiaoqi; Xu, Zhihong
(2014b): Responses of soil respiration and its components to drought stress. In *J Soils Sediments* 14 (1),
pp. 99–109. DOI: 10.1007/s11368-013-0799-7.

Werner, Christiane; Wallrabe, Ulrike; Christen, Andreas; Comella, Laura; Dormann, Carsten; Göritz, Anna
et al. (2024): ECOSENSE - Multi-scale quantification and modelling of spatio-temporal dynamics of
ecosystem processes by smart autonomous sensor networks. In *RIO* 10, Article e129357. DOI:
10.3897/rio.10.e129357.

Wickham, Hadley (2016): Data Analysis. In *ggplot2*, pp. 189–201. DOI: 10.1007/978-3-319-24277-4_9.

Wickham, Hadley; Averick, Mara; Bryan, Jennifer; Chang, Winston; McGowan, Lucy; François, Romain et
al. (2019): Welcome to the Tidyverse. In *JOSS* 4 (43), p. 1686. DOI: 10.21105/joss.01686.