Large temperature excursions have modest impacts on community composition in the high

diversity gut microbiome of omnivorous American cockroaches (*Periplaneta americana*)

Kevin C. Riedmuller[1], Josey E. Dyer[2], Elizabeth A. Ottesen[1]

[1]Department of Microbiology, University of Georgia, Athens, Georgia, United States of America

[2]College of Veterinary Medicine, University of Georgia, Athens, Georgia, United States of

America

Abstract

Microbial residents of ectothermic hosts are exposed to variations in temperature that have the potential to impact their physiology and the host-microbe symbiotic relationship. In this experimental warming study, laboratory populations of American cockroaches (*Periplaneta americana*) were kept at a baseline low room temperature of 20-22°C or a high temperature of 30°C for two weeks. We quantified bacterial load and performed high-throughput 16S rRNA gene sequencing to assess the hindgut microbiome's response to a near 10°C shift in environmental temperature. We report modest impacts of temperature on cockroach gut microbiome composition. The high temperature (30°C treatment) induced increases in the relative abundance of Proteobacteria and Euryarchaeota phyla as well as the Lactobacillaceae and Enterococcaceae. We also observed increased interindividual variability. There were no significant differences in the dominant Bacteroidota or Firmicutes phyla and no significant losses or reductions in taxa or bacterial load, respectively. This suggests that the gut community of American cockroaches is largely resilient to prolonged increases in temperature and has implications for the cockroach to withstand the impacts of climate change.

Importance

Insects, as with most animals, often harbor microbial symbionts that play an essential role in host health and nutrition. As insects are ectotherms, these microbial symbionts are subject to the same temperature fluctuations as their hosts, potentially impacting host temperature responses. Here, we demonstrate that the American cockroach (*Periplaneta americana*) gut microbiome exhibits only modest changes following an ~10°C increase in environmental temperature. This contrasts with studies in other insects, whose microbiota were highly

46 responsive to temperature variation. This work illustrates that the microbiota of insects may vary

47 in their sensitivity to long-term temperature shifts, providing a more comprehensive

48 understanding of potential variability in insect responses to climate shifts.

50 Introduction

Insects are some of the most abundant multicellular organisms on earth with an estimated

5.5 million species (1). They are key drivers of ecological and biological processes ranging from

nutrient cycling (2,3) to disease dynamics (4,5) and agricultural productivity (6,7). As

ectotherms, environmental temperature strongly governs their biological timing, physiology,

distribution, and fitness (8). Consequently, changes in environmental temperature can have

outsized and varied impacts on insects' roles in these critical systems. For example, simulated

warming in mesocosms containing *Formica subsericea*, an ant native to eastern North America,

led to decreased decomposition rates and nitrogen availability (9). In contrast, a field study of

Formica Manchu nests on the southeastern Tibetan Plateau in China found that decomposition

rates and nitrogen availability remained stable under simulated warming conditions (10). As

2024 was the warmest year on record and global surface temperatures are rising faster than any

other period over the last 2000 years (11,12), understanding how insects are affected by warming

climate has become increasingly important.

Insects have evolved a wide variety of behavioral and physiological thermoregulatory

responses for survival that modulate metabolism, stress response, and growth and development

(8). Studies frequently report faster growth and/or development rates in response to increased

temperature, which is often accompanied by a decrease in body size (13–15). However, one

aspect of insect adaptation to temperature that is less well-understood is the impact of

temperature on the insect gut microbiome. The gut microbiome plays a critical role in host

nutrition (16–18), development (19–21), and immunity (22–24). As symbionts of ectotherms,

gut microbes (and the services they provide to their hosts) are subject to changes in

environmental temperature alongside their hosts. Studies in three species of stink bugs have

reported reduction or complete loss of gut bacterial symbionts in response to increased

environmental temperature by as little as 2.5°C (25,26). Similarly, a paired observational and

experimental study in desert arboreal ants found that increased environmental temperature

resulted in decreased bacterial loads as well as moderate shifts in microbial community

composition. Notably, these changes in the gut microbiome were due to only a 1°C shift from

baseline temperature in the experimental treatment (27). These data parallel studies in other

ectotherms like corals, sponges, and mussels where increased temperature saw substantial shifts,

reductions, or complete loss of symbiotic microbes (28–30). Overall, studies in a variety of

invertebrates thus far tend to observe marked changes in both high and low diversity gut

microbiomes and subsequent decreases in host fitness as a result of elevated temperatures

ranging from ~1-9°C from baseline temperatures.

Omnivorous cockroaches carry a high-diversity hindgut microbiome (31–33) which is

believed to play an important role in facilitating host digestion of complex polysaccharides and

other refractory dietary components (34,35). The cockroach gut microbiome has been shown to

play a vital role in shaping host development (36,37), immune activity (38,39), and behavior

(40). It also shows evidence of long-term co-evolution with its host (41) and likely plays a key

role in their effectiveness as decomposers and scavengers—functions that contribute

significantly to global nutrient cycling due to their ubiquity across terrestrial ecosystems. For

example, in central Amazonian inundation forests, at least eight different species of cockroaches

were identified, one of which was estimated to consume up to 6% of annual leaf litter (42).

Their effectiveness as decomposers and scavengers also contribute to pest species' ability to

infest virtually all human spaces (43). Pest species of cockroaches pose significant human health

risks as they are a common cause of allergy and asthma (44,45) and vectors for a variety of

pathogens (46,47). Moreover, they contribute to economic losses by driving up medical costs

(48), destroying property and goods (43), and necessitating continual pest control (49). Given

their ecological and economic importance, it is critical that we understand the effects of warming

temperatures on the cockroach and its closely associated gut microbiome.

While little is known about the impact of temperature on the cockroach gut microbiome,

studies of their close relatives the termites (50) show marked temperature impacts. An

experimental study in *Reticulitermes flavipes* comparing the gut microbiota at 15°C, 27°C

(baseline) and 35°C found dramatic shifts in gut microbiome composition. Colder temperatures

were associated with increased relative abundances of Proteobacteria while heat stress was

associated with decreased bacterial richness, loss of protist associated bacteria and increased

relative abundances of Bacteroidota (51). In contrast, a field study in Janesville, Wisconsin

between July and December observed only modest seasonal shifts in gut community composition

of *R. flavipes*. However, the mean environmental air temperature during the collection period

ranged from 22°C to -2°C, which was substantially lower than the warming treatment in the

experimental study (52).

The aim of this work was to investigate the impacts of environmental temperature on the

gut microbiome of the model insect *Periplaneta americana*. It has been observed that their

thermal preference ranges from ~24-33°C (53). However, in many cases experimental insects

are kept at a variety of temperatures within or below the thermal preference range (54–56) or at

an unspecified room temperature (33,57–59). While such experimental differences likely make

little difference to the gut microbiota of endotherms such as mice, it poses a potential

complicating factor in studies of microbial symbioses of ectotherms. As a result, we sought to

examine the impact of temperature on the gut microbiota of omnivorous cockroaches. To do so,

we employed bacterial load quantification and high throughput 16S rRNA gene sequencing to

assess the hindgut microbiome's response to an increase in environmental temperature from a

'low' laboratory room temperature of 20-22°C to 30°C, a 'high' temperature used to speed

reproduction and development (60). This work provides insight into how temperature can

impact the gut microbiota of ectothermic hosts; This knowledge has implications for both

organismal responses to climate shifts and experimental practice for laboratory investigation of

ectotherm symbioses.

Materials and methods

Insects and experimental design

P. americana cockroach stock colonies have been maintained in captivity at room

temperature (20-22°C) for over 15 years. Stock colonies live in 10-gallon glass aquarium tanks

with wood chip bedding (PJ Murphy Forest Products SaniChips) and cardboard tubes for

housing. They were provided dog chow (Purina One) and water (via cellulose sponge) *ad

libitum*.

For temperature treatments, 20-24 mixed-sex, adult cockroaches were selected from stock

colonies, divided into two plastic terrariums, and maintained under stock colony conditions at

room temperature (20-22°C) or 30°C for 14 days. Dog chow with any visible mold growth was

removed daily. The experiment was performed in duplicate. Cohort 1 and 2 treatments were

138 completed in April of 2022; cohort 2 treatments were initiated one week after cohort 1 began.

139 We also include data from a pilot experiment conducted November of 2020 (cohort 3). The

140 room temperature treatment for this cohort was slightly warmer, ranging from 22-25°C.

Sample collection and DNA extraction

Cockroaches were placed on ice in sterile petri dishes until torpid (~5 min). The entire

gut (foregut to rectum) was excised with a scalpel and forceps; any visible fat body and

exoskeleton was removed. The gut tract was then moved to a sterile aluminum dish, placed over

dry ice, and the hindgut was resected. Hindguts were transferred to 500 µL of sterile 1x

phosphate buffered saline and stored at -20°C.

Hindguts were thawed and homogenized with a sterile microcentrifuge pestle. DNA was

extracted from a 200 µL aliquot of sample homogenate using the E.Z.N.A Bacterial DNA Kit

(Omega Bio-tek, Norcross, GA) per manufacturer's protocol with the following modifications:

30 min lysozyme incubation, optional glass bead beating step included, and RNase A step

excluded. Sample DNA was eluted in 50 µL of elution buffer. DNA concentrations and

A260/280 values were measured with a BioTek Synergy HTX Multimode plate reader (Agilent,

Santa Clara, CA). Sample DNA was stored at -20°C and concentrations ranged from 30 to 300

ng/µL (Table S1).

Bacterial load quantification

Total number of bacteria per sample was assessed via qPCR of the 16S rRNA gene as

described in Nadkarni et al. (61). qPCR was performed in triplicate, 10µL reactions using a

QuantStudio 6 flex instrument and QuantStudio Real Time PCR Software (v1.3)

(Applied Biosystems, Waltham, MA). Reactions consisted of 1x TaqMan Multiplex Master Mix

(Applied Biosystems), 0.3 µM forward primer 5′-TCCTACGGGAGGCAGCAGT-3′ (Tm, 59.4

°C), 0.3 µM reverse primer 5′-GGACTACCAGGGTATCTAATCCTGTT-3′ (Tm, 58.1 °C), 0.25

µM hydrolysis probe (6-FAM)-5′-CGTATTACCGCGGCTGCTGGCAC- 3′-(TAMRA) (Tm,

69.9 °C), and 1µL sample DNA (see Table S1 for DNA quantities) under the following

conditions: 95°C for 20 s followed by 40 cycles of 95°C for 1 s and 60°C for 60 s. A standard

curve was generated with a 10-fold dilution series of gBlocks Gene Fragment of the

Frigididesulfovibrio cuneatus 16S gene (GenBank: X99503.1, position 330-798) with a linear

dynamic range of 10-10^8 copies. The slope, y-intercept, and r^2 of the standard curve was -3.529,

41.67, and 0.994, respectively. Amplification efficiency was 92.04%.

qPCR of a *Blattabacterium* MIP/aquaporin family protein coding sequence was

performed to account for the cockroach endosymbiont's contribution to bacterial load.

Triplicate, 10 µL reactions consisted of 1x TaqMan Environmental Master Mix 2.0 (Applied

Biosystems), 0.1 µM forward primer 5′-CAGCTAATGCCCATCCCATAG-3′ (Tm, 55.4 °C), 0.1

µM reverse primer 5′-GGAAATGGAGTGGTGGCTAAT-3′ (Tm, 55.0 °C), 0.1 µM hydrolysis

probe (6-FAM)-5′-TTGTGACCCATCCACCATCTCCAT-3′-(MGB-NFQ) (Tm, 63.7 °C), and

1µL sample DNA (see Table S1 for DNA quantities) under the following conditions: 95°C for 10

min followed by 40 cycles of 95°C for 15 s and 60°C for 60 s. A standard curve was generated

with a 10-fold dilution series of gBlocks Gene Fragment

(CAGCTAATGCCCATCCCATAGTGATTGTGACCCATCCACCATCTCCATGTCCTTTTGTT

TTAGACAATAGAACATTAGCCACCACTCCATTTCC) with a linear dynamic range of 10-10^8

copies. The slope, y-intercept, and r^2 of the standard curve was -3.526, 38.97, and 0.993,

respectively. Amplification efficiency was 92.15%.

C_q values for both assays were auto selected by the QuantStudio Real Time PCR

Software (v1.3). All primers and gBlocks were synthesized by Integrated DNA Technologies

with standard desalting purification. Hydrolysis probes were synthesized by Applied

Biosystems. Primer set specificity was validated via endpoint PCR and agarose gel

electrophoresis.

Hindgut bacterial load L was calculated as follows:

190
$$L = ((s - b)/c) \times t$$

191   where s = 16S copy number, b = *Blattabacterium* MIP/aquaporin family protein copy number,

192   c = DNA quantity in qPCR reaction (ng), and t = total sample DNA (ng). Reported 16S and

193   *Blattabacterium* MIP/aquaporin family protein copy numbers are averages from triplicate qPCR

194   reactions. Bacterial load boxplots were generated with the Tidyverse package (62).

Library preparation and sequencing

The V4 region of the 16S rRNA gene from each hindgut sample was amplified using a

two-step PCR protocol as described in Tinker et al. (63). Initial amplification of the V4 region

was performed with a 10 µL PCR reaction that consisted of 1x Q5 hot start reaction buffer, 0.02

U/µL Q5 hot start high-fidelity DNA polymerase (New England BioLabs, Ipswitch, MA), 0.5

µM 515F primer 5´-GTGCCAGCMGCCGCGGTAA-3´, 0.5 µM 806R primer 5´-

GGACTACHVGGGTWTCTAAT-3´, 200 µM deoxynucleotide triphosphates (dNTPs), and 0.8

ng/µL sample DNA under the following conditions: 98°C for 30 s, followed by 15 cycles of

98°C for 10 s, 52°C for 30 s, and 72°C for 30 s, followed by a final extension step at 72°C for 2

min. The product of reaction 1 was reamplified in a subsequent 30 µL PCR reaction, containing

1x Q5 hot start reaction buffer, 0.02 U/µL Q5 hot start high-fidelity DNA polymerase, 0.5 µM

515F barcoded primer, 0.5 µM 806R barcoded primer, 200 µM dNTPs, and 9 µL of reaction 1

product under the following conditions: 98°C for 30 s, followed by 4 cycles of 98°C for 10 s,

52°C for 30 s, and 72°C for 30 s, followed by 6 cycles of 98°C for 10 s and 72°C for 1 min,

followed by a final extension step at 72°C for 2 min. All primers were synthesized by Integrated

DNA Technologies with standard desalting purification (Coralville, IA).

PCR products were verified by agarose gel electrophoresis and purified with the E.Z.N.A

Cycle Pure Kit (Omega Bio-tek) per manufacturer's protocol and eluted in 30 µL elution buffer.

Purified amplicon concentrations and A260/280 values were measured with a BioTek Synergy

HTX Multimode plate reader. Amplicons were normalized and pooled to a final concentration of

10 ng/µL. Quality of the prepared amplicon library was assessed with the Agilent 2100

Bioanalyzer DNA-HS assay (Agilent) before submission to the Georgia Genomics and

Bioinformatics Core for paired-end 250bp sequencing on an Illumina MiSeq platform (Illumina

Inc., San Diego, CA).

Sequence processing

Demultiplexed, paired-end fastq files were analyzed with the standard DADA2 (64)

pipeline. Forward reads were truncated at 240 bp and reverse reads were truncated at 160 bp.

Samples were pooled during the sample inference step to increase sensitivity to sequence

variants present at low abundances in multiple samples. Taxonomy was assigned to amplicon

sequence variants (ASVs) with the DADA2 package and the Silva reference database (v138.1)

(65); reads corresponding to *Blattabacterium* (cockroach endosymbiont located in the fat body),

chloroplasts, and mitochondria were removed. All downstream analyses of sequencing and

bacterial load data were performed using R Statistical Software (v4.2.1) (66) and Geneious

Prime 2025.0.3 (https://www.geneious.com).

Alpha and beta diversity

All libraries were resampled to a depth of 10,797 reads with the Vegan package (v2.6-4)

(67) for alpha and beta diversity metrics. Alpha diversity was assessed by calculating the

Shannon diversity index, richness (number of observed taxa), and Pielou's evenness (Shannon

diversity index/log(richness)) (67). Alpha diversity was visualized with violin plots via the

Tidyverse packages (v2.0.0) (62). Beta diversity was assessed by calculating weighted and

unweighted (incidence based, equivalent to Sorenson index) (68) Bray-Curtis dissimilarities (67).

Beta diversity was visualized with nonmetric multidimensional scaling ordinations and violin

plots via the Tidyverse and Vegan packages (62,67).

Differential abundance analysis

DESeq2 (v1.38.3) (69) was implemented for ASV level differential abundance analysis to

identify temperature responsive taxa. The ASV raw counts table was passed to the DESeq

command with default parameters and 'design = ~ cohort + temperature', testing for the effect of

temperature while controlling for cohort variation. ASVs with an adjusted p value < 0.05 were

considered significant (Table S2). A phylogenetic tree of differentially abundant ASVs present in

the 12 most abundant families (maximum relative abundance $> 10\%$) was generated in Geneious

Prime. Sequences were aligned with Clustal Omega (v1.2.3) (70) and the phylogenetic tree was

built using FastTree (v2.1.11) (71,72). The *Desulfovibrionaceae* clade was used as an outgroup

for rooting, and tree visualization was performed in R with the Tidyverse and ggtree (v3.6.2) (73)

packages. A heatmap of differentially abundant ASVs present in the 12 most abundant families

was generated with the ComplexHeatmap package (74,75).

Data availability

256   The sequences generated from this experiment were submitted to the NCBI Sequence Read

257   Archive and are available under BioProject PRJNA1380472.

Results

Hindgut bacterial load is stable across temperature treatments.

261   To assess the effect of temperature on hindgut bacterial load, we performed qPCR of the

16S rRNA gene with a universal primer/probe set as described in Nadkarni et al (61). To account

for 16S rRNA copy contribution from the cockroaches' *Blattabacterium* endosymbiont, we also

performed qPCR of a *Blattabacterium* gene in the MIP/aquaporin family. Wilcoxon rank-sum

tests, a non-parametric test comparing the distribution of independent groups, were used for two-

group comparisons. Kruskal-Wallis tests, an extension of the Wilcoxon rank-sum test, were used

for the overall comparison of three or more groups (non-parametric ANOVA analogue).

Significant Kruskal-Wallis tests were followed up with post-hoc Dunn's test with Bonferroni

adjustment to determine which groups' distributions were significantly different (FSA v0.9.6)

(76). A Wilcoxon rank-sum test found no significant difference in hindgut bacterial load across

temperature treatments ($p = 0.26$). A Kruskal-Wallis test found a significant difference in

hindgut bacterial load among cohorts ($p = 0.04$). However, Dunn's test did not indicate

significant differences between cohorts: cohort 1 and 2 ($p = 0.10$), cohort 1 and 3 ($p = 1.00$), and

cohort 2 and 3 ($p = 0.07$) (Fig 1).

Hindgut microbial community composition

A total of 4,526,160 rRNA gene sequences were obtained and 2,580,005 passed filtering

parameters, resulting in an average of 42,295 reads per sample. One sample was removed prior

to downstream analysis due low sequencing depth (Table S3). 2,923 unique ASVs were

identified across all samples.

Overall gut microbiome composition exhibited modest changes in response to

temperature. Wilcoxon rank sum tests found no significant differences in the Shannon diversity,

Pielou's evenness, or richness (total number of ASVs) of samples at the ASV or family level

across temperature treatments (Shannon: ASV $p = 0.86$, family $p = 0.57$; Pielou's: ASV $p = 0.88$,

family $p = 0.56$; richness: ASV $p = 0.64$, family $p = 0.47$) (Figs 2A and S1A). However, beta

diversity measures suggest modest but significant impacts of temperature on gut microbiome

community composition. Dunn's test indicated that between-temperature treatment

dissimilarities were significantly greater than within- room temperature treatment dissimilarities

at the ASV ($p < 0.001$) and family level ($p < 0.001$) (Figs 2B and S1B). While nonmetric

multidimensional scaling (NMDS) of weighted Bray-Curtis dissimilarities did not indicate clear

separation of samples by temperature, permutational multivariate analysis of variance

(PERMANOVA) (67) found a significant effect at the ASV ($p = 0.001$, $r^2 = 0.05$) and family ($p =$

0.001, $r^2 = 0.05$) levels (Figs 2C and S1C). Temperature was also observed to have impacts on

beta dispersion, with the 30°C treatment having higher within-group dissimilarity than the room

temperature treatment (Dunn's test: ASV $p < 0.001$, family $p < 0.001$) (Figs 2B and S1B).

Given the large time gap between experiments, we also examined cohort impacts on

community composition. Notably, cohort 3 samples tended to cluster separately from other

cohorts by weighted and unweighted Bray-Curtis metrics and PERMANOVA found a significant

effect for cohort on community composition at the ASV (weighted: $p = 0.001$, $r^2 = 0.07$;

unweighted: $p = 0.001$, $r^2 = 0.13$) and family (weighted: $p = 0.04$, $r^2 = 0.04$, unweighted: $p =$

0.001, $r^2 = 0.22$) level (Figs 2C and S1C). Dunn's test indicated significantly greater between-

cohort dissimilarity for all comparisons involving cohort 3 compared to the between-cohort

dissimilarity of 1 and 2 by weighted and unweighted metrics at the ASV level ($p < 0.001$) (Fig

S2). This suggests that some observed taxa in cohort 3 differ from the observed taxa in cohorts 1

and 2. This trend aligns temporally with each experiment, where cohort 1 and 2 treatments were

carried out within a week of each other while cohort 3 was carried out over one year earlier. This

indicates minor, detectable drift in taxonomic composition of our stock colonies. To investigate

whether cohort effects were obscuring temperature responses, we examined temperature effects

on gut microbiome composition separately for cohorts 1 and 2, which were combined due to

their more similar community composition, and cohort 3 (Figs 2C and 3). Accounting for cohort

variation, samples separated more clearly by temperature at the ASV level (weighted Bray-Curtis

dissimilarity) than at the family level (Fig 3), although the effect size of temperature calculated

by PERMANOVA was still modest (cohort 1 and 2: $p = 0.003$, $r^2 = 0.07$, cohort 3: $p = 0.001$, $r^2 =$

0.13). This suggests that, while we observe stability of the gut microbiome at the family-level in

response to temperature, there may be some ASV-level differences contributing to observed

stability.

Taxa exhibiting significant responses to temperature differences

All ASVs present in more than 50% of the room temperature samples were also present

All ASVs present in more than 50% of the room temperature samples were also present

in the 30°C samples, and vice versa, suggesting that temperature differences in community

composition were primarily the result of changes in ASV relative abundance. We performed

differential abundance analysis with DESeq2 to further explore ASV level responses to

environmental temperature. Accounting for cohort variation, 231 ASVs were identified as

significantly differentially abundant ($p < 0.05$) across temperature treatments (Table S2). Of

those 231 ASVs, 123 belonged to the most abundant families (maximum relative abundance >

10%) in our samples (Figs 4 and S4). Ruminococcaceae and Lachnospiraceae families had a

comparable number of ASVs enriched in both temperature treatments. This may be indicative of

ASV-level switching among functionally redundant taxa within these families. However, we did

not see this trend for the remaining families. All but one Bacteroidaceae ASV was enriched in

the room temperature treatment and all Lactobacillaceae and Enterococcaceae ASVs were

enriched in the 30°C treatment. We found that most differentially abundant ASVs in the

Tannerellaceae, Dysgonomonadaceae, Rikenellaceae, Christensenellacae, and

Desulfovibrionaceae families were enriched in the 30°C treatment (Figs 4 and S4).

These ASV level differences translated into significant differences in family and phylum

level abundances. At the family level, Wilcoxon rank sum tests found that Bacteroidaceae were

significantly more abundant at room temperature (median difference = 2.07%, $p < 0.001$) while

Lactobacillaceae and Enterococcaceae were significantly more abundant at 30°C (median

difference = 1.94%, $p = 0.01$ and median difference = 0.41%, $p = 0.005$) (Figs 5A and S5).

When summed at the phylum level, Bacteroidota and Firmicute (Bacillota) abundances were not

significantly different across temperature treatments ($p = 0.83$ and $p = 0.89$). In contrast, the

sulfate-reducing Desulfobacterota (Thermodesulfobacteridota) were significantly more abundant

at room temperature (median difference = 3.5%, *p* = 0.04) while the Euryarchaeota (containing

exclusively *Methanibrevibacter* ASVs) were significantly more abundant at 30°C (median

difference = 0.17%, *p* = 0.005). Proteobacteria were significantly more abundant in the 30°C

treatment (median difference = 0.26%, *p* = 0.04) (Figs 5B and S6).

Discussion

The goal of this study was to examine how changes in environmental temperature affect

gut microbial community composition and stability in omnivorous cockroaches. We observed

modest differences between room temperature (20-22°C) and 30°C treatments at the phylum and

family level (Figs 2 and S1). At the phylum level, there were significantly more Proteobacteria

in the 30°C treatment than the room temperature treatment. The observed shift in Proteobacteria

relative abundance was subtle, with a median difference of 0.26% (Fig S1). This response

parallels studies in other insects and invertebrates. In a study of rough wood lice (*Porcellio*

scaber), those maintained at 22°C had higher abundances of Proteobacteria (Enterobacteriaeae)

compared to those maintained at 15°C (77). Experimental warming of the invertebrate nematode

Caenorhabditis elegans also resulted in higher abundances of Proteobacteria (*Agrobacterium*) as

temperature increased (15, 20, and 25°C treatments) (78). However, those studies observed more

pronounced changes in Proteobacteria abundances compared to our findings in the American

cockroach and were concurrent with substantial decreases in other common gut microbiota such

as the Actinobacteria in *P. scaber* and *Sphingobacterium* in *C. elegans*. In the closely related

termite *Reticulitermes flavipes*, Arango et al. saw a considerable increase in the abundance of

Proteobacteria (*Acinetobacter,* Betaproteobacteriales, *Enterobacter*) at their lowest temperature

treatment of 15°C compared to 27 or 35°C as well as subsequent decreases in Bacteroidota

abundances (51). Studies in the fruit fly *Drosophila melanogaster* and cricket *Gryllus veletus*

also report that Proteobacteria are responsive to temperature (79,80). However, the responsive

Proteobacteria in the fruit fly and cricket studies are primarily *Wolbachia*, an intracellular

bacterial symbiont that is not directly associated with the gut microbiome.

In vertebrate studies, increases in Proteobacteria abundance are associated with gut

dysbiosis, pronounced shifts in community composition and diseases like irritable bowel

syndrome and Crohn's disease (81–83). The Proteobacteria are generally composed of

opportunists that can survive a broad range of stressors compared to other phyla. It may be that

changes in their abundance is indirectly determined by other community members response to

temperature. In this scenario, Proteobacteria are well suited to 'take over' under conditions where

more typical symbiotic populations are under environmental stress. We observed similar patterns

with the Enterococcaceae and Lactobacillaceae in the 30°C treatment (Figs 4, 5 and S4), both of

which are also commonly associated with gut dysbiosis (84,85). These taxa appear to bloom in a

subset of our 30°C samples, consistent with our observation of increased beta-dispersion under

high temperature conditions. This aligns with the "Anna Karenina" model where dysbiotic

microbiota are unique and highly variable compared to healthy microbiota (86,87). We note a

similar pattern in *Cephalotes rohweri* ant colonies exposed to high temperatures in field warming

studies (27).

We also observed a significant decrease in Desulfobacterota at higher temperatures. The

Desulfobacterota are highly active in *P. americana*, primarily acting as sulfate reducers (34,88).

As sulfate reducers, they are likely in competition with methanogens for hydrogen (89,90).

Cockroach gut methanogenic taxa include the Euryarchaeota which were significantly more

abundant in the 30°C treatment. This suggests that hydrogen utilization may tip in favor of

methanogens at higher temperatures, or that methanogens themselves fare better at higher

temperatures, increasing hydrogen utilization and decreasing Desulfobacterota abundances.

These observations align with studies in soils and batch reactors, where methanogens

outcompete sulfate reducers as temperatures rise (91–93). Further studies are needed to better

quantify the methanogenic archaea, as universal 16S rRNA primers used in this study are not

optimized for their detection.

The modest shifts in gut microbiome composition that we report here contrasts with the

experimental warming of *R. flavipes* termites, in which a shift from 27°C to 35°C resulted in

large changes in phylum-level composition. Arango et al. reported that Bacteroidetes

(Bacteroidota) almost doubled in relative abundance while Spirochaetes, Euryarchaeota and

Elusimicrobia decreased. However, the authors noted that many of the microbial taxa lost

following warming tended to be associated with ciliated protists that do not fare well at these

higher temperatures (51). These protists are critical for breakdown of lignocellulosic material in

lower termites (94). While similar protists have been reported in the cockroach, they are not

essential for cockroach survival and energy acquisition (95). Microscopic observation of

samples from our stock tanks indicate that a few protists are present but that they tend to be

present at low abundance and low frequency. It may be that the 'high' temperature tested here

was not high enough to negatively impact the protists, or that any reduction in protists had a

negligible impact on overall community composition.

Together, these results suggest that the cockroach gut microbiome is relatively stable

across a wide range of environmental temperatures. While the 30°C temperature used in our

study is within the preferred temperature range of this species, we observed that all cockroaches

maintained at this temperature had a darker cuticle by the end of the experiment. Cuticle

melanization has been directly linked to dessication tolerance in *Blattella germanica* (96),

suggesting that our warming treatment was high enough to elicit a melanization stress response

in *Periplaneta americana*. Additionally, it is common for bacteria to have lower tolerances for

higher-than-preferred than lower-than-preferred temperatures (97). The high temperature tested

shows some signs of increased variability and growth of opportunistic Proteobacteria,

Enterococcaceae, and Lactobacillaceae which may have implications for the ability of these

insects to serve as disease vectors in less climate-controlled settings. Finally, we highlight the

potential for temperature to impact microbial competition between essential hydrogenotrophic

organisms in the gut. This study suggests that the microbiome of ectotherms can be surprisingly

resilient to temperature shifts of up to 10°C and has important implications for the ability of

these insects to tolerate the impacts of climate change.

Acknowledgements

425   This work was supported by the National Institute of General Medical Sciences of the National

426   Institutes of Health under award number R35GM133789 and the National Science Foundation

427   Graduate Research Fellowship Program under Grant No. 2236869. Any opinions, findings, and

428   conclusions or recommendations expressed in this material are those of the author(s) and do not

429   necessarily reflect the views of the National Science Foundation.

Fig 1. Comparison of hindgut bacterial load across temperature treatments and cohort.

Violin plots of hindgut bacterial load across (A) temperature and (B) cohort where bars represent

the median and points represent individual samples. Wilcoxon rank-sum test was used to

compare across temperature treatments. Kruskal-Wallis and post-hoc Dunn's test with

Bonferroni adjustment was used to compare across cohorts. *Blattabacterium* counts subtracted

from total 16S rRNA copy number. Bacterial load normalized by total sample DNA. RT = room

temperature, ns = no significance.

Fig 2. ASV level comparison of hindgut microbial community composition across temperature treatments.

(A) Violin plots of alpha diversity measurements (Shannon diversity, observed richness, Pielou's evenness) where points represent the medians. (B) Violin plots of weighted (left) and unweighted (right) Bray-Curtis dissimilarities within and between temperature treatments where points represent the medians. Wilcoxon rank-sum tests were used to compare alpha diversity measures. Kruskal-Wallis and post-hoc Dunn's test with Bonferroni adjustment was used to compare Bray-Curtis dissimilarities. (C) Nonmetric multidimensional scaling (NMDS) of weighted (top) and unweighted (bottom) Bray-Curtis dissimilarities. NMDS stress was calculated with the metaMDS() function from the Vegan package. PERMANOVA was used to calculate r^2 and p values for temperature and cohort. All libraries were resampled to a depth of 10,797 reads. RT = room temperature, * = $p < 0.05$, ** = $p < 0.01$, *** = $p < 0.001$, ns = no significance.

464
Fig 3. ASV and family level comparison of Bray-Curtis dissimilarities accounting for cohort

differences.

Nonmetric multidimensional scaling (NMDS) of weighted Bray-Curtis dissimilarities at the (A)

ASV and (B) family level. NMDS stress was calculated with the metaMDS() function from the

Vegan package. PERMANOVA was used to calculate r^2 and p values for temperature and cohort.

All libraries were resampled to a depth of 10,797 reads. RT = room temperature, $* = p < 0.05$,

$** = p < 0.01$, $*** = p < 0.001$.

Fig 4. Phylogenetic tree of differentially abundant ASVs across temperature treatments.

FastTree depicting differentially abundant ASVs in the most abundant families (maximum

relative abundance > 10%). Highlights represent family level classifications. Leaf tip labels

represent genus level classifications. ASVs without a leaf tip label were not classifiable at the

genus level. Leaf tip colors depict which treatment ASVs were enriched in. Branch lengths not

shown. RT = room temperature.

Fig 5. Comparison of family and phylum level relative abundances across temperature treatments.

(A) Relative abundance of the 12 most abundant families (maximum relative abundance > 10%). Each bar represents an individual sample. ++ = higher in room temperature treatment, + = higher in 30°C treatment. (B) Violin plots showing the relative abundance of select phyla where bars represent the median and points represent individual samples. All libraries were resampled to a depth of 10,797 reads. Wilcoxon rank-sum tests were used to compare relative abundance across temperature treatments. RT = room temperature, * = $p < 0.05$, ** = $p < 0.01$, ns = no significance.

References

1. Stork NE. How Many Species of Insects and Other Terrestrial Arthropods Are There on Earth? Annu Rev Entomol. 2018;63:31–45.

2. Whiles MR, Callaham MA, Meyer CK, Brock BL, Charlton RE. Emergence of Periodical Cicadas (Magicicada cassini) From a Kansas Riparian Forest: Densities, Biomass and Nitrogen Flux. Am Midl Nat. 2001;145(1):176–87.

3. Metcalfe DB, Asner GP, Martin RE, Silva Espejo JE, Huasco WH, Farfán Amézquita FF, et al. Herbivory makes major contributions to ecosystem carbon and nutrient cycling in tropical forests. Ecol Lett. 2014;17(3):324–32.

4. Emerson PM, Lindsay SW, Walraven GEL, Faal H, Bøgh C, Lowe K, et al. Effect of fly control on trachoma and diarrhoea. Lancet. 1999;353:1401–3.

5. Effler PV, Pang L, Kitsutani P, Vorndam V, Nakata M, Ayers T, et al. Dengue Fever, Hawaii, 2001–2002. Emerg Infect Dis. 2005;11(5):742–9.

6. Rader R, Bartomeus I, Garibaldi LA, Garratt MPD, Howlett BG, Winfree R, et al. Non-bee insects are important contributors to global crop pollination. Proc Natl Acad Sci. 2016;113(1):146–51.

7. Bowling RD, Brewer MJ, Kerns DL, Gordy J, Seiter N, Elliott NE, et al. Sugarcane Aphid (Hemiptera: Aphididae): A New Pest on Sorghum in North America. J Integr Pest Manag. 2016;7(1):12.

8. González-Tokman D, Córdoba-Aguilar A, Dáttilo W, Lira-Noriega A, Sánchez-Guillén RA, Villalobos F. Insect responses to heat: physiological mechanisms, evolution and ecological implications in a warming world. Biol Rev. 2020;95(3):802–21.

9. Toro ID, Ribbons RR, Ellison AM. Ant-mediated ecosystem functions on a warmer planet: effects on soil movement, decomposition and nutrient cycling. J Anim Ecol. 2015;84(5):1233–41.

10. Luo B, Huang M, Wang W, Niu J, Shrestha M, Zeng H, et al. Ant nests increase litter decomposition to mitigate the negative effect of warming in an alpine grassland ecosystem. Proc R Soc B. 290:20230613.

11. Temperatures Rising: NASA Confirms 2024 Warmest Year on Record - NASA [Internet]. [cited 2025]. Available from: https://www.nasa.gov/news-release/temperatures-rising-nasa-confirms-2024-warmest-year-on-record/

12. IPCC, 2023: Climate Change 2023: Synthesis Report. Contribution of Working Groups I, II and III to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change [Core Writing Team, H. Lee and J. Romero (eds.)]. Geneva, Switzerland: IPCC; 2023 p. 35–115.

531 13. Atkinson D. Temperature and Organism Size—A Biological Law for Ectotherms? Adv Ecol
532  Res. 1994;25:1–58.

533 14. Mas-Martí E, Muñoz I, Oliva F, Canhoto C. Effects of increased water temperature on leaf
534  litter quality and detritivore performance: a whole-reach manipulative experiment. Freshw
535  Biol. 2015;60(1):184–97.

536 15. Garrad R, Booth DT, Furlong MJ. The effect of rearing temperature on development, body
537  size, energetics and fecundity of the diamondback moth. Bull Entomol Res.
538  2016;106(2):175–81.

539 16. Cummings JH, Pomare EW, Branch WJ, Naylor CPE, Macfarlane GT. Short chain fatty acids
540  in human large intestine, portal, hepatic and venous blood. Gut. 1987;28(10):1221–7.

541 17. Ridaura VK, Faith JJ, Rey FE, Cheng J, Duncan AE, Kau AL, et al. Gut Microbiota from
542  Twins Discordant for Obesity Modulate Metabolism in Mice. Science. 2013;341:1241214.

543 18. Kenny DJ, Plichta DR, Shungin D, Koppel N, Hall AB, Fu B, et al. Cholesterol Metabolism
544  by Uncultured Human Gut Bacteria Influences Host Cholesterol Level. Cell Host Microbe.
545  2020;28:245–57.

546 19. Reinhardt C, Bergentall M, Greiner TU, Schaffner F, Östergren-Lundén G, Petersen LC, et
547  al. Tissue factor and PAR1 promote microbiota-induced intestinal vascular remodelling.
548  Nature. 2012;483:627–31.

549 20. Gordon HA, Bruckner-Kardoss E. Effect of normal microbial flora on intestinal surface area.
550  Am J Physiol-Leg Content. 1961;201(1):175–8.

551 21. Banasaz M, Norin E, Holma R, Midtvedt T. Increased Enterocyte Production in Gnotobiotic
552  Rats Mono-Associated with *Lactobacillus rhamnosus* GG. Appl Environ Microbiol.
553  2002;68(6):3031–4.

554 22. Bouskra D, Brézillon C, Bérard M, Werts C, Varona R, Boneca IG, et al. Lymphoid tissue
555  genesis induced by commensals through NOD1 regulates intestinal homeostasis. Nature.
556  2008;456:507–10.

557 23. Sanos SL, Bui VL, Mortha A, Oberle K, Heners C, Johner C, et al. RORγt and commensal
558  microflora are required for the differentiation of mucosal interleukin 22–producing NKp46+
559  cells. Nat Immunol. 2009;10(1):83–91.

560 24. Cash HL, Whitham CV, Behrendt CL, Hooper LV. Symbiotic Bacteria Direct Expression of
561  an Intestinal Bactericidal Lectin. Science. 2006;313:1126–30.

562 25. Prado SS, Hung KY, Daugherty MP, Almeida RPP. Indirect Effects of Temperature on Stink
563  Bug Fitness, via Maintenance of Gut-Associated Symbionts. Appl Environ Microbiol.
564  2010;76(4):1261–6.

26. Kikuchi Y, Tada A, Musolin DL, Hari N, Hosokawa T, Fujisaki K, et al. Collapse of Insect
Gut Symbiosis under Simulated Climate Change. mBio. 2016;7(5):e01578-16.

27. McMunn MS, Hudson AI, Zemenick AT, Egerer M, Bennett L, Philpott SM, et al. Thermal
sensitivity and seasonal change in the gut microbiome of a desert ant, *Cephalotes rohweri*.
FEMS Microbiol Ecol. 2022;98(7):1–11.

28. Ramsby BD, Hoogenboom MO, Whalan S, Webster NS. Elevated seawater temperature
disrupts the microbiome of an ecologically important bioeroding sponge. Mol Ecol.
2018;27:2124–37.

29. Jokiel PL, Coles SL. Effects of temperature on the mortality and growth of Hawaiian reef
corals. Mar Biol. 1977;43:201–8.

30. Li YF, Yang N, Liang X, Yoshida A, Osatomi K, Power D, et al. Elevated Seawater
Temperatures Decrease Microbial Diversity in the Gut of *Mytilus coruscus*. Front Physiol.
2018;9:839.

31. Renelies-Hamilton J, Germer K, Sillam-Dussès D, Bodawatta KH, Poulsen M. Disentangling
the Relative Roles of Vertical Transmission, Subsequent Colonizations, and Diet on
Cockroach Microbiome Assembly. mSphere. 2021;6(1):e01023-20.

32. Cazzaniga M, Domínguez-Santos R, Marín-Miret J, Gil R, Latorre A, García-Ferris C.
Exploring Gut Microbial Dynamics and Symbiotic Interaction in *Blattella germanica* Using
Rifampicin. Biology. 2023;12(7):955.

33. Tinker KA, Ottesen EA. Differences in Gut Microbiome Composition Between Sympatric
Wild and Allopatric Laboratory Populations of Omnivorous Cockroaches. Front Microbiol.
2021;12:703785.

34. Dukes HE, Tinker KA, Ottesen EA. Disentangling hindgut metabolism in the American
cockroach through single-cell genomics and metatranscriptomics. Front Microbiol.
2023;14:1156809.

35. Dockman RL, Ottesen EA. Purified fibers in chemically defined synthetic diets destabilize
the gut microbiome of an omnivorous insect model. Front Microbiomes. 2024;3:1477521.

36. Jahnes BC, Poudel K, Staats AM, Sabree ZL. Microbial colonization promotes model
cockroach gut tissue growth and development. J Insect Physiol. 2021;133:104274.

37. Jahnes BC, Herrmann M, Sabree ZL. Conspecific coprophagy stimulates normal
development in a germ-free model invertebrate. PeerJ. 2019;7:e6914.

38. Turner M, Van Hulzen L, Pietri JE. The gut microbiota induces melanin deposits that act as
substrates for *fimA*-mediated aggregation of *Salmonella* Typhimurium and enhance infection
of the German cockroach vector. Microbiol Spectr. 2023;11(5):e02119-23.

39. Turner M, Van Hulzen L, Guse K, Agany D, Pietri JE. The gut microbiota confers resistance
against *Salmonella* Typhimurium in cockroaches by modulating innate immunity. iScience.
2024;27:111293.

40. Wada-Katsumata A, Zurek L, Nalyanya G, Roelofs WL, Zhang A, Schal C. Gut bacteria
mediate aggregation in the German cockroach. Proc Natl Acad Sci. 2015;112(51):15678–83.

41. Tinker KA, Ottesen EA. Phylosymbiosis across Deeply Diverging Lineages of Omnivorous
Cockroaches (Order Blattodea). Appl Environ Microbiol. 2020;86(7):e02513-19.

42. Irmler L, Furch K. Production, energy, and nutrient turnover of the cockroach *Epilampra*
irmleri Rocha e Silva & Aguiar in a Central-Amazonian inundation forest. AMAZONIANA.
1979;6(4):497–520.

43. Wang C, Lee CY, Rust MK. Biology and Management of the German Cockroach. Clayton
South VIC, Australia: CSIRO Publishing; 2021.

44. Arbes SJ, Gergen PJ, Elliott L, Zeldin DC. Prevalences of positive skin test responses to 10
common allergens in the US population: Results from the Third National Health and
Nutrition Examination Survey. J Allergy Clin Immunol. 2005;116(2):377–83.

45. Kang B, Vellody D, Homburger H, Yunginger JW. Cockroach cause of allergic asthma: Its
specificity and immunologic profile. J Allergy Clin Immunol. 1979;63(2):80–6.

46. Memona H, Manzoor F, Anjum AA. Cockroaches (Blattodea: Blattidae): A Reservoir of
Pathogenic Microbes in Human-Dwelling Localities in Lahore. J Med Entomol.
2017;54(2):435–40.

47. Liu J, Yuan Y, Feng L, Lin C, Ye C, Liu J, et al. Intestinal pathogens detected in cockroach
species within different food-related environment in Pudong, China. Sci Rep. 2024;14:1947.

48. Rosenstreich DL, Eggleston P, Kattan M, Baker D, Slavin RG, Gergen P, et al. The Role of
Cockroach Allergy and Exposure to Cockroach Allergen in Causing Morbidity among Inner-
City Children with Asthma. N Engl J Med. 1997;336(19):1356–63.

49. Gulati J. Pest Control in the US. IBIS World Industry Report; 2025 Nov. Report No.: 56171.

50. Inward D, Beccaloni G, Eggleton P. Death of an order: a comprehensive molecular
phylogenetic study confirms that termites are eusocial cockroaches. Biol Lett. 2007;3:331–5.

51. Arango RA, Schoville SD, Currie CR, Carlos-Shanley C. Experimental Warming Reduces
Survival, Cold Tolerance, and Gut Prokaryotic Diversity of the Eastern Subterranean
Termite, *Reticulitermes flavipes* (Kollar). Front Microbiol. 2021;12:632715.

52. Arango RA, Bishell AB, Ohno KM, Shelton TG, Schoville SD, Carlos-Shanley C. Seasonal
shifts in gut microbiota and cold tolerance metrics in a northern population of *Reticulitermes*
flavipes (Blattodea: Rhinotermitidae). Environ Entomol. 2024;53(3):447–56.

53. Gunn DL. The Temperature and Humidity Relations of the Cockroach: III. A Comparison of
Temperature Preference, and Rates of Desiccation and Respiration of *Periplaneta*
Americana, *Blatta Orientalis* and *Blatella Germanica*. J Exp Biol. 1935;12(2):185–90.

54. Ayayee PA, Ondrejech A, Keeney G, Muñoz-Garcia A. The role of gut microbiota in the
regulation of standard metabolic rate in female *Periplaneta americana*. PeerJ. 2018;6:e4717.

55. Lee S, Kim JY, Yi MH, Lee IY, Lee WJ, Moon HS, et al. Comparative Microbiome Analysis
of Three Species of Laboratory-Reared *Periplaneta* Cockroaches. Korean J Parasitol.
2020;58(5):537–42.

56. Vera-Ponce de León A, Jahnes BC, Otero-Bravo A, Sabree ZL. Microbiota Perturbation or
Elimination Can Inhibit Normal Development and Elicit a Starvation-Like Response in an
Omnivorous Model Invertebrate. mSystems. 2021;6:e00802-21.

57. Chen Z, Wen S, Shen J, Wang J, Liu W, Jin X. Composition and diversity of the gut
microbiota across different life stages of American cockroach (*Periplaneta americana*). Bull
Entomol Res. 2023;1–7.

58. Bertino-Grimaldi D, Medeiros MN, Vieira RP, Cardoso AM, Turque AS, Silveira CB, et al.
Bacterial community composition shifts in the gut of *Periplaneta americana* fed on different
lignocellulosic materials. SpringerPlus. 2013;2:609.

59. Zurek L, Keddie BA. Contribution of the colon and colonic bacterial flora to metabolism and
development of the American cockroach *Periplaneta americana* L. J Insect Physiol.
1996;42(8):743–8.

60. Rollo CD, Gunderman MW. Variation among individuals and the effect of temperature on
food consumption and reproduction in the cockroach, *Periplaneta americana* (Orthoptera:
Blattidae). Can Entomol. 1984;116(6):785–93.

61. Nadkarni MA, Martin FE, Jacques NA, Hunter N. Determination of bacterial load by real-
time PCR using a broad-range (universal) probe and primers set. Microbiology.
2002;148:257–66.

62. Wickham H, Averick M, Bryan J, Chang W, McGowan LD, François R, et al. Welcome to
the Tidyverse. J Open Source Softw. 2019;4(43):1686.

63. Tinker KA, Ottesen EA. The Core Gut Microbiome of the American Cockroach, *Periplaneta*
americana, Is Stable and Resilient to Dietary Shifts. Appl Environ Microbiol.
2016;82(22):6603–10.

64. Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP. DADA2: High-
resolution sample inference from Illumina amplicon data. Nat Methods. 2016;13(7):581–3.

65. Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, et al. The SILVA ribosomal
RNA gene database project: improved data processing and web-based tools. Nucleic Acids
Res. 2013;41:D590–6.

66. R Core Team. R: A Language and Environment for Statistical Computing [Internet]. Vienna,
Austria: R Foundation for Statistical Computing; 2022. Available from: https://www.R-
project.org/

67. Oksanen J, Simpson GL, Blanchet FG, Kindt R, Legendre P, Minchin PR, et al. vegan:
Community Ecology Package [Internet]. 2022. Available from: https://cran.r-
project.org/web/packages/vegan/index.html

68. Legendre P, De Cáceres M. Beta diversity as the variance of community data: dissimilarity
coefficients and partitioning. Ecol Lett. 2013;16:951–63.

69. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-
seq data with DESeq2. Genome Biol. 2014;15:550.

70. Sievers F, Wilm A, Dineen D, Gibson TJ, Karplus K, Li W, et al. Fast, scalable generation of
high-quality protein multiple sequence alignments using Clustal Omega. Mol Syst Biol.
2011;7:539.

71. Price MN, Dehal PS, Arkin AP. FastTree: Computing Large Minimum Evolution Trees with
Profiles instead of a Distance Matrix. Mol Biol Evol. 2009;26(7):1641–50.

72. Price MN, Dehal PS, Arkin AP. FastTree 2 – Approximately Maximum-Likelihood Trees for
Large Alignments. PLOS ONE. 2010;5(3):e9490.

73. Yu G, Smith DK, Zhu H, Guan Y, Lam TTY. ggtree: an R package for visualization and
annotation of phylogenetic trees with their covariates and other associated data. Methods
Ecol Evol. 2017;8:28–36.

74. Gu Z. Complex heatmap visualization. iMeta. 2022;1:e43.

75. Gu Z, Eils R, Schlesner M. Complex heatmaps reveal patterns and correlations in
multidimensional genomic data. Bioinformatics. 2016;32(18):2847–9.

76. Ogle DH, Doll JC, Wheeler AP, Dinno A. FSA: Simple Fisheries Stock Assessment Methods
[Internet]. 2025. Available from: https://CRAN.R-project.org/package=FSA

77. Horváthová T, Babik W, Kozłowski J, Bauchinger U. Vanishing benefits - The loss of
actinobacterial symbionts at elevated temperatures. J Therm Biol. 2019;82:222–8.

78. Berg M, Stenuit B, Ho J, Wang A, Parke C, Knight M, et al. Assembly of the *Caenorhabditis
elegans* gut microbiota from diverse soil microbial environments. ISME J. 2016;10:1998–
2009.

79. Ferguson LV, Dhakal P, Lebenzon JE, Heinrichs DE, Bucking C, Sinclair BJ. Seasonal shifts
in the insect gut microbiome are concurrent with changes in cold tolerance and immunity.
Funct Ecol. 2018;32:2357–68.

80. Moghadam NN, Thorshauge PM, Kristensen TN, De Jonge N, Bahrndorff S, Kjeldal H, et al. Strong responses of *Drosophila melanogaster* microbiota to developmental temperature. Fly (Austin). 2018;12(1):1–12.

81. Willing B, Halfvarson J, Dicksved J, Rosenquist M, Järnerot G, Engstrand L, et al. Twin Studies Reveal Specific Imbalances in the Mucosa-associated Microbiota of Patients with Ileal Crohn's Disease. Inflamm Bowel Dis. 2009;15(5):653–60.

82. Willing BP, Dicksved J, Halfvarson J, Andersson AF, Lucio M, Zheng Z, et al. A Pyrosequencing Study in Twins Shows That Gastrointestinal Microbial Profiles Vary With Inflammatory Bowel Disease Phenotypes. Gastroenterology. 2010;139:1844–54.

83. Su Q, Tun HM, Liu Q, Yeoh YK, Mak JWY, Chan FKL, et al. Gut microbiome signatures reflect different subtypes of irritable bowel syndrome. Gut Microbes. 2023;15(1):2157697.

84. Dahal RH, Kim S, Kim YK, Kim ES, Kim J. Insight into gut dysbiosis of patients with inflammatory bowel disease and ischemic colitis. Front Microbiol. 2023;14:1174832.

85. Pietrucci D, Cerroni R, Unida V, Farcomeni A, Pierantozzi M, Mercuri NB, et al. Dysbiosis of gut microbiota in a selected population of Parkinson's patients. Parkinsonism Relat Disord. 2019;65:124–30.

86. Zaneveld JR, McMinds R, Thurber RV. Stress and stability: applying the Anna Karenina principle to animal microbiomes. Nat Microbiol. 2017;2:17121.

87. Ma Z. Testing the Anna Karenina Principle in Human Microbiome-Associated Diseases. iScience. 2020;23:101007.

88. DePoy AN, Wall HE, Tinker KA, Black IM, Ottesen EA. Metabolic flexibility among fiber degraders supports gut microbiome stability across host diets in the American cockroach [Internet]. bioRxiv; 2025 [cited 2026]. p. 2024.10.31.621369. Available from: https://www.biorxiv.org/content/10.1101/2024.10.31.621369v3

89. Gibson GR, Cummings JH, Macfarlane GT. Competition for hydrogen between sulphate-reducing bacteria and methanogenic bacteria from the human large intestine. J Appl Bacteriol. 1988;65:241–7.

90. Lovley DR, Dwyer DF, Klug MJ. Kinetic Analysis of Competition Between Sulfate Reducers and Methanogens for Hydrogen in Sediments. Appl Environ Microbiol. 1982;43(6):1373–9.

91. Bodegom PMV, Stams AJM. Effects of alternative electron acceptors and temperature on methanogenesis in rice paddy soils. Chemosphere. 1999;39(2):167–82.

92. Muller E, Guélard J, Sissmann O, Tafit A, Poirier S. Evidencing the influence of temperature and mineralogy on microbial competition for hydrogen consumption: Implications for underground hydrogen storage (UHS). Int J Hydrog Energy. 2024;82:1101–13.

737   93. Jung H, Kim J, Lee C. Temperature Effects on Methanogenesis and Sulfidogenesis during
738   Anaerobic Digestion of Sulfur-Rich Macroalgal Biomass in Sequencing Batch Reactors.
739   Microorganisms. 2019;7:682.

740   94. Todaka N, Moriya S, Saita K, Hondo T, Kiuchi I, Takasu H, et al. Environmental cDNA
741   analysis of the genes involved in lignocellulose digestion in the symbiotic protist community
742   of *Reticulitermes speratus*. FEMS Microbiol Ecol. 2007;59:592–9.

743   95. Gijzen HJ, Barugahare M. Contribution of anaerobic protozoa and methanogens to hindgut
744   metabolic activities of the American cockroach, *Periplaneta americana*. Appl Environ
745   Microbiol. 1992;58(8):2565–70.

746   96. Bai TT, Pei XJ, Liu TX, Fan YL, Zhang SZ. Melanin synthesis genes *BgTH* and *BgDdc*
747   affect body color and cuticle permeability in *Blattella germanica*. Insect Sci. 2022;29:1552–
748   68.

749   97. Bennett AF, Lenski RE. Evolutionary Adaptation to Temperature II. Thermal Niches of
750   Experimental Lines of *Escherichia Coli*. Evolution. 1993;47(1):1–12.