Transcriptional responses of *Solanum lycopersicum* to three distinct parasites reveal host hubs and networks underlying parasitic successes

Julia Truch[1]*†, Maëlle Jaouannet[1]*, Martine Da Rocha[1], Emmanuelle Kulhanek-Fontanille[1], Cyril Van Ghelder[1], Corinne Rancurel[1], Olivier Migliore[1], Arthur Péré[1], Stéphanie Jaubert[1], Christine Coustau[1], Eric Galiana[1] and Bruno Favery[1]†

* These authors contributed equally to the current study

† Corresponding authors (julia.truch@inrae.fr, bruno.favery@inrae.fr)

Affiliations

[1]Institut Sophia Agrobiotech, INRAE, Université Côte d'Azur, CNRS, 400 routes des Chappes, 06903, Sophia-Antipolis, France.

Abstract

Crops face attacks from a wide range of pathogens and pests that deploy various parasitic strategies. All aggressors must manipulate major plant functions, such as immune signaling, hormone pathways, metabolism, and development, to successfully establish themselves and complete their life cycle. Here, we compared the tomato plant's response to three evolutionary distant pathogens (nematodes, aphids and oomycetes) during compatible interactions using a transcriptomic approach. We identified differentially expressed genes and biological processes, and highlighted potential key host hubs associated with successful parasitism. By integrating recent published datasets, we refined our understanding of the global and tissue-specific mechanisms targeted during compatible interactions and, through co-expression analysis, identified clusters showing shared dysregulation patterns enriched in specific GO terms. Finally, model-to-crop translational analysis using the Arabidopsis interactome network, repositioned tomato candidates within larger interaction networks and emphasized the key positions occupied by some of them. Identifying these pivotal tomato targets is crucial to decipher processes underlying parasitism and, consequently, offers new opportunities to develop sustainable multi-pathogen control strategies.

Keywords

Comparative transcriptomic meta-analyses, tomato, hub, network, parasitism, plant pathogens, nematode, oomycete, aphid

Introduction

Plants are severely challenged by soil-borne and aerial pathogens that threaten crop production worldwide. Cultivated tomato (*Solanum lycopersicum*), the most important vegetable crop by total production and second by harvested area globally (FAOSTAT2022; https://www.fao.org/faostat/en/#home; 186 Mt/year; 5 M of ha of surface harvested), is grown across a wide range of climates and serves as a host for numerous pathogens, including oomycetes, nematodes, and insects. In resistant plants, early detection of the pathogen through

basal immunity (pathogen-associated molecular patterns PAMP-triggered immunity, PTI) or later
through the effector-triggered immunity (ETI) modulates MAPK signaling, transcription factors
(WRKY, ERF, MYB…), epigenetic modifications, hormone imbalance (salicylic acid, jasmonate,
auxins…) leading to productions of reactive oxygen species, pathogenesis-related proteins and
secondary metabolites (Campos et al., 2021). Thus, compatible interactions between host plants
and parasites rely on the avoidance of plant immunity and the reprogramming of the host's
metabolic activity (Jones & Dangl, 2006). Consequently, pathogens hijack plant resources for their
own profit, ultimately leading to reduced plant growth, yield losses, and potentially premature plant
death (Kong & Yang, 2023).
As endoparasites, oomycetes and root-knot nematodes (RKN) have mastered long-term
interactions within the host roots (Blok et al., 2008; Kamoun et al., 2015). On one hand, zoospores
of the oomycete *Phytophthora parasitica* encyst and adhere to the plant surface before penetrating
intercellularly. In the roots, *P. parasitica* hyphae invaginate host cell membranes, form haustoria
and secrete apoplastic molecules, effectors, that promote pathogen development (Attard et al.,
2008). At 15 hours post- infection, *P. parasitica* uses appressoria to penetrate host roots during an
initial biotrophic phase (Attard et al., 2010). On the other hand, the mobile infective second-stage
juveniles of the RKN *Meloidogyne incognita* penetrate host roots, and migrate intercellularly until
reaching the central cylinder, where they become sedentary and induce a feeding structure
composed of several giant multinucleate cells. At 21 days post infection (dpi), giant cells are fully
differentiated and function as mature, metabolically active feeding structures that sustain RKN
development (Abad et al., 2008; Favery et al., 2020).
In contrast, aphids such as *Macrosiphum euphorbiae* are ectoparasites establishing durable yet
transient interactions by feeding on host plants via sap-sucking (Walling, 2008). Aphid colonies
develop on the leaf surface and winged adults ensure the dissemination of new colonies. At 24
hours post-infection in a confined space, aphids have induced their feeding and the host displays
a peak response (Jaouannet et al., 2015). Similar to *M. incognita*, *M. euphorbiae* uses a stylet to
deliver effectors that manipulate host immunity and metabolism, enabling nutrient acquisition and
colony establishment (Favery et al., 2020; F. Wang et al., 2021).
Characterizing both the host targets of effectors and the host functions manipulated is essential to
decipher the mechanisms underlying parasitic success, and represents a crucial step to develop
new strategies to control pathogens. The diversity of life traits is associated with effector multiplicity
which, overall, converges to target directly or indirectly fundamental host components. Interestingly,
evolutionary distant pathogens deploy effectors that interact with the same host targets (Mukhtar
et al., 2011; Carella et al., 2018). Such shared targets are often interconnected key molecules,
hubs, whose perturbation can compromise the robustness of the host regulatory networks. In this
context, monitoring plant transcriptional activity is a relevant approach to identify key factors and
pathways commonly targeted by pathogens. Some independent works individually analyzed the
transcriptomic response of *S. lycopersicum* to the parasites used in our study (i.e. *M. incognita*, *M.*
euphorbiae and *P. parasitica*) (Coppola et al., 2013; Naveed & Ali, 2018; Mejias et al., 2021).
However none of them investigated and integrated all the responses in a cordonnated experimental
procedure and single analysis. Comparative transcriptome meta-analyses focusing on a single
model plant or crop species exposed to multiple pathogens, are gaining momentum. These studies
reanalyze public RNA-seq datasets to generate multi-pathogens plant response profiles (Biniaz et
al., 2022; Y. Wang et al., 2022). Conversely to these studies, which rely on conditions defined by
the original experiments, our study used the same tomato variety, matching the tomato reference
genome for optimal mapping of RNA-seq data, plants grown under the same conditions and
carefully chosen infection time points. To this end, we assessed the transcriptomic response of *S.*
lycopersicum var. 'Heinz 1706' infected or not by the RKN *M. incognita*, the oomycete *P. parasitica*
or the aphid *M. euphorbiae*, using comparative RNA-seq approaches and pathway analyses.
Beyond the identification of genes that are strongly and commonly dysregulated upon infection, we

implemented a co-expression analysis to pinpoint candidate hubs associated with parasitic success
and to position homologs of these candidate gene products within their interactome network
environment.

Methods

Material production

Six four-week-old plants of *S. lycopersicum var. 'Heinz 1706'* per replicate were grown in a growth
chamber under a 16-h photoperiod at 24°C and inoculated with 1000 J2s of *M. incognita* or 500,000
zoospores of *P. parasitica* in the soil, or 50 mixed stages *M. euphorbiae*. Root galls resulting from
M. incognita infection were collected at 21 dpi, *P. parasitica*-infected roots were harvested 15 hpi,
and leaves attacked by *M. euphorbiae* were sampled 24 hpi. For each condition, equivalent roots
or leaves tissues were collected from non-infected plants. Four independent biological replicates
were produced per condition.

RNA extraction

Samples were ground with mortars in liquid nitrogen and transferred to new tubes. Four volumes
of extraction buffer (0.1 M Glycine-NaOH pH 9, 0.1 M NaCl, 0.1% SDS and 1% deoxycholic acid
sodium salt) were added. Samples were incubated 5 min on ice, centrifuged for 5 min and
supernatants were transferred to fresh tubes. One volume of phenol:chloroform:isoamyl alcohol
(PCI; 25:24:1, v/v/v) was added. Samples were vortexed for 15 sec, incubated 2 min at room
temperature and centrifuged at 15,000 × g for 15 min at 4°C. The aqueous phase was collected
and a second extraction with PCI was performed. The aqueous phase was washed twice with one
volume of chloroform before been centrifuged at 12,000 × g for 15 min at 4°C. Sodium acetate was
added to a final concentration of 0.3 M, followed by one volume of cold 100% isopropanol. Samples
were incubated overnight at -20°C and centrifuged at 12,000 × g for 15 min at 4°C. RNA pellets
were washed twice with 1 ml of 70% ethanol, centrifuged at 12,000 × g for 5 min at 4°C, air dried
for 10 min, resuspended in 100 µl of RNase free water and 33µl of 8M Lithium Chloride and
incubated for one hour at -20°C and centrifuged at 12,000 × g for 5 min at 4°C. Pellets were washed
twice with 1ml of cold 70% ethanol, air-dried 10 min and resuspended in 25 µl of RNase free water
at 55°C for 5 min. gDNA contamination was removed using Qiagen RNeasy Plus kit following the
manufacturer's instructions.

Library preparation and sequencing

117 Library preparation and sequencing were performed by Eurofins Genomics Europe Sequencing
118 GmbH. RNA samples were assessed for integrity and quality and strand-specific, directional polyA
119 selected library preparation was performed using NEBNext Ultra II Directional RNA Library Prep
120 Kit for Illumina with TruSeq Adapter sequences. The samples were sequenced on Illumina
121 NovaSeq 6000 platform using 2x150 Sequence mode.

Raw data analysis

Raw data were analyzed using an in-house Nextflow based pipeline. Briefly, quality controls of raw
data were performed using FastQC (fastqc%3A0.11.9--hdfd78af_1) and MultiQC
(multiqc%3A1.10.1--pyhdfd78af_1) before and after data cleaning including trimming and filtering
for reads of at least 100 bp using FastP (fastp%3A0.23.2--h79da9fb_0). Reads were then mapped
with STAR (star%3A2.7.8a--h9ee0642_1) with the options --genomeSAindexNbases 10 for the

indexing and --alignEndsType EndToEnd and --twopassMode Basic for the mapping. For each
interaction, reads were mapped on a composite genome combining *S. lycopersicum* v4 genome
(solgenomics.net) with either the *M. incognita* V3 genome (Blanc-Mathieu et al., 2017), the *M.*
euphorbiae transcriptome (NCBI ref GAOM01000000) or the *P. parasitica* genome
GCA_000247585.2 (https://protists.ensembl.org) in order to remove potential pathogen reads.
Then, all reads mapping to pathogen genome/transcriptome were removed to conserve only the
reads mapping to the *S. lycopersicum* v4 genome. Read counts were obtained with Feature-Counts
(subread:2.0.1--hed695b0_0) with the options -t mRNA and -g ID. Differential analyses were
performed using DESeq2 and EdgeR. The gene annotation was based on the *S. lycopersicum*
annotation ITAG4.1 (solgenomics.net). Principal component analysis (PCA), VENN diagrams and
co-expression analyses in roots and leaves were performed using DiCoExpress (Lambert et al.,
2020). Graphical representations were performed with ggplot2 (ggplot2_3.5.0). Filtering,
normalization and quality control (including PCA analysis) of the dataset were performed using the
Quality_Control option with default parameters (CPM_Cutoff=1 and Normalization_Method =
"TMM"). Datasets were analyzed using GLM_Contrasts model taking into account the interaction.
DiffAnalysis.edgeR was run with default parameters (Alpha_DiffAnalysis = 0.05, NbGenes_Profiles
= 20, NbGenes_Clustering = 50). Venn diagrams were performed using Venn_IntersectUnion
option. Co-expression analyses were performed using Coexpression_coseq option based on
Gaussian mixture models using default parameters on gene candidates commonly dysregulated in
roots or leaves pathogens therefore excluding the candidates present in the respective interactions.
Singularity prebuilt images were retrieved from the Galaxy Project Singularity Depot
(https://depot.galaxyproject.org/singularity/).

Subtilase and TCP gene family analyses

Fifty-five protein sequences were retrieved from TAIR (Araport11), matching with the 56 subtilases
described in A. Figueiredo et al. (2014), with AT3G44262 removed because it is a pseudogene).
Similarly, 103 protein sequences annotated as subtilase were extracted from the *S. lycopersicum*
protein dataset (ITAG4.1). The 158 sequences were aligned using MUSCLE (v3.8.31) with default
settings for highest accuracy. No outgroup has been added as the objective was to position *S.*
lycopersicum sequences according to *A. thaliana* sequences and the known subtilase sub-families.
The phylogenetic tree was reconstructed using the maximum likelihood method implemented in the
PhyML program (v3.1/3.0 aLRT). The WAG substitution model was selected assuming an
estimated proportion of invariant sites (of 0.000) and 4 gamma-distributed rate categories to
account for rate heterogeneity across sites. The gamma shape parameter was estimated directly
from the data (gamma=1.197). Reliability for internal branch was assessed using the aLRT test
(SH-Like). Graphical representation and edition of the phylogenetic tree were performed with iTOL
v7. Similarly, 24 and 32 protein sequences annotated as TCP transcription factors were retrieved
from the *A. thaliana* (Araport11) and the *S. lycopersicum* (ITAG4.1) protein datasets, respectively.
These 56 sequences were aligned with MEGA X using MUSCLE. The evolutionary history was
inferred using the Neighbor-Joining method. The bootstrap consensus tree was inferred from 1000
replicates. The evolutionary distances were computed using the JTT matrix-based method.

Gene ontology (GO) and transcription factor analyses

The GO analyses were performed with GOFuncR (GOfuncR_1.22.2) except the co-expression
analysis part for which the GO analyses were performed using DiCoExpress using Enrichment
option with default parameters (Alpha_Enrichment=0.01) (Lambert et al., 2020). Visualisation was
done with ggplot2 (3.5.0). The list of transcription factors was downloaded from the Plant

Transcription Factor Database (PlantTFDB v5.0) (https://planttfdb.gao-lab.org/index.php?sp=Sly)
(Tian et al., 2019).

Ortholog identification, effector interactor and interactome network analyses

Protein sequences from *S. lycopersicum* (ITAG4.1), *Arabidopsis thaliana* (Araport11), *Amborella*
trichopoda (AMTR1.0), *Brassica napus* (AST_PRJEB5043_v1), *Capsicum annuum*
(ASM51225v2), *Coffea arabica* (GCF_003713225.1_Cara_1.0), *Helianthus annuus* (HanXRQr2.0-
SUNRISE), *Medicago truncatula* (MedtrA17_4.0), *Nicotiana benthamiana* (Niben261), *Oryza sativa*
(IRGSP-1.0), *Prunus persica* (NCBIv2), *Sorghum bicolor* (NCBIv3), *Selaginella moellendorffii*
(v1.0), *Solanum tuberosum* (SolTub_3.0), *Vitis vinifera* (PN40024.v4) and *Zea mays* (Zm-B73-
REFERENCE-NAM-5.0) were functionally annotated using InterProScan (5.51-85.0) and the
ortholog analysis was performed using OrthoFinder (2.5.4). The identification of effectors
interacting with candidate homologs was performed using EffectorK (González-Fuente et al.,
2020). *A. thaliana* based interactome was generated with BIOGRID-ORGANISM (v4.4.227) using
BIOGRID-ORGANISM-Arabidopsis_thaliana_Columbia-4.4.227.psi25   and   analysed   with
Cytoscape (v3.10.1).

Results

Global transcriptomic response of S. lycopersicum to M. incognita, P. parasitica and M.
euphorbiae

We performed comparative RNA-seq analyses on *S. lycopersicum* roots inoculated or not with
either *M. incognita* or *P. parasitica* and leaves inoculated or not with *M. euphorbiae,* to assess the
transcriptional reprogramming occurring in tomato under biotic stresses. To be considered as
differentially expressed genes (DEGs), candidates had to display both, an adjusted p-value < 0.05
in DEseq2 analysis and FDR < 0.05 in EdgeR analysis. Thus, we identified 12,231 DEGs at 21 dpi
with *M. incognita*, 12,018 DEGs at 15 hpi with *P. parasitica* and 11,001 DEGs at 24 hpi with *M.*
euphorbiae (Figure 1a and Figure S1). Each interaction displayed a high number of DEGs with
similar proportions of up- and down-regulated candidates (Figure 1a). A total of 4,461 DEGs were
identified in response to the three pathogen infections (Figure 1b). Among them, 961 were down-
regulated and 645 were up-regulated across all three conditions, whereas 2,855 showed variable
dysregulation patterns depending on the specific interaction studied (Figure 1c). To obtain an
overview of the biological processes commonly targeted by the three pathogens, we performed GO
enrichment analyses on the lists of DEGs identified. We identified 53 biological processes GO
terms shared across the three interactions (Figure 1d). Among them, 45 GO terms were all either
over- or under-represented (Figure S2a), whereas eight showed variability depending on the
specific interaction (Figure S2b). All over-represented GO terms were related to metabolic
processes, while the negative regulation of metabolic processes were among the under-
represented GO terms. These results are consistent with the enhanced metabolic activity of the
host associated with pathogen infection.
Focusing on the DEGs most strongly responsive to all three pathogens, we identified only 11 genes
with a log2 fold change (log2FC) < -2 across all interactions (Figure 1e). This short list includes the
TOLERANT TO CHILLING AND FREEZING 1 (*AtTCF1*) Arabidopsis ortholog, *Solyc05g052950.*
This stress response-associated gene encodes for a regulator of chromosome condensation that
indirectly affects lignin accumulation (Ji et al., 2015). Surprisingly, the list also contains two *A.*
thaliana orthologs of transcription factors that are involved in plant pigment regulation (flavonol and
chlorophyll, respectively): the *Solyc06g005330,* related to the AtMYB48 transcription factor and
Solyc12g009240, a putative ethylene-responsive transcription factor (homolog of ERF017 in *A.*

thaliana). Conversely, 10 DEGs displayed a log2FC >2 in three interactions (Figure 1f). Among
them, we identified genes that are involved in a variety of biological responses. Notably, we
identified tomato orthologs of *A. thaliana* proteins known to regulate systemic acquired resistance
(SAR), such as the NON-INDUCIBLE IMMUNITY1 (NIM1)-INTERACTING 2-like protein
(Solyc04g007140); proteins induced upon biotic stress, such as N-acetyltransferase NATA induced
by insect feeding and bacterial infection (Solyc08g006765) (Adio et al., 2011; Lou et al., 2016) or
abiotic stresses, such as the chromatin remodeling factor OXIDATIVE STRESS 3 (OXS3,
Solyc06g066420) (Blanvillain et al., 2009). Finally, we identified two susceptibility factors: (i) the
osmotin-like protein OSM34 (Solyc08g080670), which is subverted in *A. thaliana* by the turnip
mosaic virus (TuMV) thus decreases plasmodesmata callose deposition to promote viral cell-to-
cell movement and replication (He et al., 2025), and (ii) DOWNY MILDEW RESISTANT6 DMR6,
an enzyme that converts SA into its inactive form and whose loss of function in tomato confers
broad-spectrum disease resistance (Thomazella et al., 2021). Consistently, both *DMR6-1* and
DRM6-2 (*Solyc03g080190* and *Solyc06g073080*) were upregulated in the three interactions tested
(Figure 1f and Figure S2c).

Figure 1 - Transcriptomic response of *S. lycopersicum* to *M. incognita*, *P. parasitica* and *M. euphorbiae*. a. Barplot showing the number of DEGs up- and down-regulated depending on the tested parasites. b. Venn diagram illustrating the overlap between the DEGs identified in the experiments with the different parasites. c. Barplot comparing the change in gene expression of the 4461 candidate genes differentially expressed during the three interactions. D. Venn Diagrams of GO overlap between the three interactions considering all the DEGs. e. DEGs with a log2FC < -2 in the three interactions tested. f. DEGs with a log2FC > 2 in the three interactions tested. Me: *M. euphorbiae*, Mi: *M. incognita* and Pp: *P. parasitica*.

243 Furthermore, we identified two subtilases (or subtilisin-like proteases) among the top 10 most
244 strongly upregulated DEGs in response to the three parasites. These two proteases encoding
245 genes (*Solyc01g087800* and *Solyc01g087810*) were arranged in tandem within a larger subtilase
246 cluster (cluster 1) on the chromosome 1 (Figure 2). The subtilase gene family was particularly well
247 represented in our DEG dataset. Indeed, 65 of the 103 annotated subtilase genes in tomato were
248 differentially expressed in response to at least one pathogen. A deeper focus on the subtilase
249 cluster 1 revealed additional subtilase candidates differentially expressed in each plant-parasites
250 interaction (Figure 2). The phylogenetic tree built with the subtilases of *A. thaliana* and *S.*
251 *lycopersicum* showed that subtilases from cluster 1 are all related to the *A. thaliana* gene *SBT1.9*
252 (*At5g67090*). In addition, the subtilase cluster 8, previously characterized in tomato (Reichardt et
253 al., 2018), also contained numerous pathogen-responsive genes (Figure 2). Interestingly, both
254 clusters included subtilases identified as phytaspases, caspases-like proteases regulating
255 programmed cell death, including *Solyc01g087800,* the most highly expressed candidate. Our
256 results showed that subtilases in Arabidopsis have more gene members from the subfamilies SBT3
257 and SBT4 than in tomato. Conversely, tomato subtilases undergo a massive expansion in the SBT1
258 subfamily that includes subtilases from the clusters 1 and 8 (Figure 2).

Cluster 1

SL4.1ch01: 74 912 271..   ..74 962 050

Figure 2 - Phylogenic tree of subtilases of the cluster 1 and 8 in the chromosome 1
and 8 of *S. lycopersicum*. The ML phylogenetic tree (WAG model, aLRT test)
contains 55 subtilases from A. thaliana and 103 subtilases from *S. lycopersicum*.
SBT1-6 subfamilies were assigned following (A. Figueiredo et al., 2014) and clades

were collapsed to facilitate the visualization (the extended developed phylogenetic
tree is available in Figure S3). In brackets, the number of *A. thaliana*
sequences/number of sequences from *S. lycopersicum*. The differential expression
(Log2 Fold Change) of genes from the cluster 1 and 8 (from chromosome 1 and 8 of
S. lycopersicum) in response to the three pathogens (*M. incognita*, *P. parasitica* and
M. euphorbiae) is presented on the right panel. Asterisks indicate the two highly
overexpressed genes in response to the three pathogens. Physical representations
of the Cluster 1 and 8, including their coordinates on the *S. lycopersicum*
chromosome 1 and 8, respectively, are presented with differentially expressed
subtilase genes in response to at least one pathogen in dark blue and light blue for
the others.

In-depth identification of targeted hubs during plant-pathogen interactions

Following the identification of the most strongly dysregulated genes, we next analyzed DEGs
through the lens of common stress-response pathways. We first examined the salicylic acid (SA)
pathway and, in addition to *DMR6s* and *NIM1-INTERACTING 2-like,* we identified five DEGs
encoding: two MACPF domain-containing proteins (Solyc01g005220 and Solyc02g077780), the
latter being implicated in the negative regulation of the SA-related defense responses (Noutoshi et
al., 2006), and three NIM1-like proteins (Solyc07g040690, Solyc02g069310 and Solyc07g044980)
(Figure 3a). All of these genes were upregulated in the presence of root pathogens, whereas their
expression pattern were more variable in response to the leaf parasite *M. euphorbiae*. We, then,
identified six candidates associated with auxin response, including two members of the small auxin
upregulated RNA (SAUR) family (Solyc01g110907 and Solyc07g042470), known to regulate cell
elongation and the activation of plasma membrane H$^+$-ATPases (Spartz et al., 2016) (Figure 3b).
Finally, we identified 11 DEGs associated to ethylene response in the three interactions, which held
the attention given the pivotal role of ethylene in term of regulation of plant development and stress
resilience (Figure 3c). Among these DEGs, we detected several transcription factor encoding
genes, including *Solyc05g052050,* which encodes Pti4, a protein interacting with the resistance Pto
kinase protein and cis regulatory elements (Zhou et al., 1997) and *Solyc06g053710,* which encodes
the ethylene receptor ETR4 (Figure 3c). Using the Plant Transcription Factor Database (Tian et al.,
2019), we further identified 34 up-regulated (Figure 3d) and 92 down-regulated (Figure 3e) genes
across the three interactions within the "*Solanum lycopersicum* Transcription Factors database",
covering 29 transcription factor families. The representations between TF families differ between
up- and down- regulated candidates. Up-regulated genes in all three interactions showed a top
enrichment within the ERF and WRKY families (Figure 3d), whereas down-regulated candidates
displayed the bHLH and MYB-related families as top enrichments (Figure 3e). We then assessed
whether the putative hubs identified by transcriptomics in *S. lycopersicum* might have *A. thaliana*
homologs encoding characterized effector hubs, *i.e.* host proteins interacting with several pathogen
effectors. We identified orthologs/homologs between *S. lycopersicum* and *A. thaliana* using
OrthoFinder (Emms and Kelly, 2019) and search the EffectorK database compiling *A. thaliana*
proteins interacting with pathogen effectors (González-Fuente et al., 2020). Among the *S.*
lycopersicum orthologs/homologs of *A. thaliana* effector interactors, we identified, among others,
candidate hubs encoding WRKYs (*Solyc03g095770* and *Solyc03g116890*), bHLH
(*Solyc05g009880*) and TCP (*Solyc01g008230*) transcription factors, a protein phosphatase 2C
(PP2CA) (Solyc03g096670), a protein homologous to RESPONSE TO LOW SULFUR 2 (LSU2)
(*Solyc03g096780*) and several paralogs of plasma membrane intrinsic proteins (*Solyc06g011350,*
Solyc09g007760 and *Solyc09g007770*) (Figure 3f). Altogether, these results pointed out a short
panel of putative key tomato hubs that could be directly or indirectly affected by at least these three
pathogens and acting at multiple levels to promote parasitic success.

Figure 3 - Characterization of candidate genes with a common response in the three interactions. a. DEGs associated with salicylic acid pathways in the three interactions tested. b. DEGs associated with auxin response in the three interactions tested. c. DEGs associated with ethylene response in the three interactions tested. d. Distribution of candidate genes identified in Plant Transcription Factor Database and

318   up-regulated in the three interactions tested. e. Distribution of candidate genes
319   identified in Plant Transcription Factor Database and down-regulated in the three
320   interactions tested. f. Selection of candidate hubs in *S. lycopersicum* encoding for
321   orthologs of effector interactors in *A. thaliana*. Me: *M. euphorbiae*, Mi: *M. incognita*
322   and Pp: *P. parasitica*

Root and leaf parasites trigger preferentially different pathways

To dissect the regulatory networks underlying parasitic success, we performed a co-expression
analysis. We took advantage of publicly available RNA-seq generated from *S. lycopersicum* leaves
in presence (24 hpi) or absence of the South American tomato pinworm *Tuta absoluta* (Roumani
et al., 2022) to further explore tissue-dependent co-expression patterns. PCA revealed a clear
separation between leaves and roots along the first component, accounting for more than 45% of
the variance (Figure S4a), consistent with the distinct gene expression patterns observed in roots
and leaves (Attard et al., 2010; Pirona et al., 2023). Therefore, we examined the co-expression
profiles for root and leaf infections separately using DiCoExpress (Lambert et al., 2020), taking into
account the interaction effect between biological factors to identify the genes displaying a common
response to the pathogens tested. Multifactorial analysis of the RNA-seq data from the root
pathogen experiments (*M. incognita* and *P. parasitica*) pointed out 2,119 candidate genes showing
a common response to root pathogens (Figure 4a).
Co-expression analysis of these candidates resulted in five clusters (Figure 4b and Figure S4b).
Clusters 1, 2 and 3 contained genes generally up-regulated in response to both root pathogens (*M.
incognita* and *P. parasitica*), whereas clusters 4 and 5 were characterized by a general pattern of
downregulation in response to root pathogens. We identified over-represented cluster-specific
biological processes using a GO enrichment analysis for each cluster (Figure 4c). Cluster 1 was
characterized by an overrepresentation of processes linked with the regulation of SAR and
ethylene-activated signaling pathway, including the candidate hub NIM1-INTERACTING 2-like
(*Solyc04g007140*) (Figure 1f). In contrast, clusters 2 and 3 showed an enrichment in processes
associated with programmed cell death (PCD) and regulation of jasmonic acid mediated signaling
pathway (including the NIM1-like protein 1 encoded by *Solyc02g069310*) (Figure 3a). Both clusters
were also enriched for GO term "regulation of SA mediated signaling pathway". Processes linked
to trehalose biosynthetic process and response to hormone were overrepresented in clusters 4 and
5, respectively (Figure 4c). Notably, clusters 3 and 4 shared an enrichment in processes linked to
protein phosphorylation despite displaying opposite co-expression patterns. This suggests that
genes belonging to the same pathway may have opposite roles (positive /negative regulators), and
that parasites may need to up-regulate some components while downregulating others to achieve
parasitic success (Figure 4c).

Figure 4 - Co-expression analysis of root response to telluric parasites. (a) Venn
diagram of DEGs in roots in response to *M. incognita*, *P. parasitica* taking into
account the interaction of both factors. (b) Co-expression showing the five clusters
with boxplot expression profiles per sample per conditions. (c) GO enrichment of
biological processes for the five co-expression clusters containing gene candidates
commonly dysregulated in presence of root pathogens.

We then performed a similar analysis with leaf pathogens (*M. euphorbiae* and *T. absoluta*).
Multifactorial analysis identified 938 genes showing a common response to leaf pathogens (Figure
5a). Co-expression analysis of these candidates also resulted in five clusters (Figure 5b and Figure
S4c). As observed with root pathogens, GO enrichments were cluster specific (Figure 5c). Cluster
1 showed an overrepresentation of processes related to vesicle trafficking and carbohydrate
metabolic process, with candidates associated with cell wall remodeling such as the beta-
mannosidase (*Solyc03g119080*) (Pirona et al., 2023). Additional cell wall-associated genes, such
as *Solyc09g098510* (extensin-2-like), were present in cluster 2, which was enriched in plant-type
cell wall organization processes. Cluster 3 displayed enrichments in processes associated with
protein phosphorylation and photosynthesis. Among the genes in cluster 3, the MAP kinase kinase
kinase encoding gene *Solyc02g065110* has been associated with the induction of PCD (Roberts
et al., 2019). Cluster 4 was enriched in ion transport processes with cyclic nucleotide-gated ion
channel candidates, consistent with the changes in ion fluxes associated with plant immune
response to pathogens (Moeder et al., 2011) and in hydrogen peroxide catabolic process. Finally,
cluster 5 was enriched in inositol catabolism with downregulated inositol oxygenases, a gene family
involved in the cell wall biosynthesis and ascorbic acid biosynthesis pathway (Munir et al., 2020).
Although several candidate genes responded to both root and leaf pathogens, the co-expression
analyses revealed a panel of biological processes significantly over-represented in a cluster and
tissue specific manner. These results unveiled clear distinct modules of underlying regulations
associated with pathogen responses.

Figure 5 - (a) Venn diagram of DEG in leaf in response to *M. euphorbiae* and *T. absoluta* taking into account the interaction of both factors. (b) Co-expression clusters with the boxplot expression profiles per sample per conditions in leaves. (c) GO enrichment of biological processes per co-expression cluster gene candidates commonly dysregulated in presence of leaf pathogens.

Putative *S.* lycopersicum hubs targeted by parasites are homologs of *A. thaliana* hubs

Finally, we evaluated the relevance of our candidate hubs within the experimentally tested *A. thaliana* interactome network in a model-to-crop translational work using OrthoFinder-identified orthologs as proxies. We characterized over 900 homologs of the putative hubs identified in this study within the *A. thaliana* interactome network, most of which taking part in complex, highly interconnected networks (in orange in Figure 6a). To reconstruct the networks affected by the three parasites, we then considered only the interactions between the homologs of our candidate hubs. This analysis revealed the central positions of several effector-interacting homologs, such as the PP2CA (Solyc03g096670 ortholog) and the plasma membrane intrinsic protein PIP2A (associated with paralogs Solyc06g011350, Solyc09g007760 and Solyc09g007770), among numerous other candidates (orange nodes in Figure 6b). Furthermore, the analysis pinpointed the pivotal position of NATA1 (AT2G39030, Solyc08g006765 homolog) (in red in Figure 6c). NATA1 interacts with two transcription factors, HB-12 (Solyc01g096320 homolog) and TCP2 (Solyc08g048370 homolog), thereby bridging two important regulatory network subsets. ATHB12 acts as positive transcriptional regulators of *PP2C* genes, and thus as a negative regulator of ABA signaling (Valdés et al., 2012). On the other branch of this network, we identified three TCP transcription factors, TCP2, TCP21 and TCP24, which centralize interactions involving numerous DEGs identified in our study. TCP2 and TCP21 are known to be directly targeted by pathogen's effectors (González-Fuente et al., 2020; Drcelic et al., 2024; Fu et al., 2025). Consistently, in our study, Solyc08g048370 and Solyc01g008230, the tomato homologs of TCP2 and TCP21, were similarly dysregulated in response to the three parasites, making them strong candidate hubs (Figure S5).

Figure 6 - S. lycopersicum orthologs in the context of experimentally tested A. thaliana regulatory network. In orange, the orthologs/homologs of candidate hubs identified in S. lycopersicum (this study). (a) A. thaliana full interactome network highlighting, the orthologs/homologs of candidate hubs identified in S. lycopersicum

417   (this study). (b) A. thaliana interactome subset showing only the interactions between
418   the orthologs/homologs of candidate hubs identified in this study. (c) A. thaliana
419   interactome subset highlighting the key position of NATA1 (AT2G39030 in red)
420   among the network of orthologs/homologs of candidate hubs. In yellow: effector
421   interacting proteins.

Discussion

Compatible plant-pathogen interactions rely on the pathogens ability to manipulate host gene
expression in order to (1) facilitate pathogen entry through interference with the integrity of physical
barriers such as modulations of the cell wall structure or stomatal opening, (2) suppress host
immune responses by altering gene expression regulation and hormonal signaling pathways and
(3) promote pathogen access to host nutrients via the regulation of metabolite synthesis and
nutrient transport (van Schie & Takken, 2014; D. Wang et al., 2025). Such genes can be
evolutionary conserved across species (Koseoglou et al., 2022) and may be targeted, directly or
indirectly, by effectors from trans-kingdom pathogens. Understanding these hub targets in depth
could therefore contribute to the development of broad-spectrum resistance through their targeted
impairment.
Although pathogens generally downregulate defense pathways while activating processes that
favor their own growth, host responses reflect complex regulatory networks with redundancy and
feedback loops, complicating transcriptomic interpretation. Illustrating this conundrum, we found
that *Solyc08g080670* was among the most strongly upregulated genes in response to all three
parasites tested. This gene is the tomato homolog of *OSM34* in *A. thaliana,* a stress-responsive
gene, functioning in the ABA signaling pathway, and potentially playing a role in plant protection
against fungi infections (Mukherjee et al., 2010; Hao et al., 2015; Park & Kim, 2021). A recent study
demonstrated that the TuMV hijacked *AtOSM34* to promote viral infection. Indeed, *AtOSM34*
upregulation reduces plasmodesmata callose deposition to promote viral intercellular movement
and targeted viral replication complex to promote viral replication. Additionally, it suppressed the
ROS-mediated antiviral response (He et al., 2025). Whether multiple pathogens trigger the
upregulation of *Solyc08g080670* to promote infection remains to be determined, but this possibility
highlights its potential importance as a key factor to target. Another set of key candidate genes
identified in this study are *SlDMR6-1* and *SlDMR6-2,* the closest homologs of the salicylic Acid 5-
Hydroxylase DMR6 and DLO, which are well-known susceptibility factors to oomycetes and
bacteria. In Arabidopsis, these enzymes function redundantly, in different tissues, to suppress plant
immunity. They catalyze the formation of 2,5-Dihydroxybenzoic acid2 (5-DHBA) by hydroxylating
SA, thereby inactivating SA (Zeilmaker et al., 2015; van Butselaar et al., 2024). Here, we showed
that both genes are strongly overexpressed in tomato in response to all three pathogens tested.
Interestingly, *SlDMR6-1* is only slightly induced in response to aphids, consistent with the variable
spatial expression mentioned in the literature and with the possibility that *SlDMR6-1* and SlDMR6-
2 respond differently depending on the pathogen. For instance, Thomazella et al. (2021) reported
an upregulation of *SlDMR6-1* whereas SlDMR6-2 expression was not detected upon infection by
Xanthomonas gardneri (Thomazella et al., 2021). Although such genes represent potential targets
for gene inactivation to enhance plant resistance, their role in maintaining SA homeostasis
highlights the trade-off between immunity and growth.
Another important multi-pathogen target identified in our study is the *NIMIN-2* homolog
(Solyc04g007140). NIMIN proteins interact with NPR1, the key positive regulator of SAR acting in
the SA-mediating pathway (Zavaliev et al., 2020). Interestingly, NIMIN proteins act as negative
regulators of NPR1, therefore, the upregulation of *Solyc04g007140* is likely to favor pathogen

infection by compromising SAR, making it a prominent target (Weigel et al., 2005). Pathogen
infections also affected host epigenetic factors. Indeed, *Solyc06g066420* belongs to the *OXS 3*
family, which, like OSM34, functions in ABA signaling and acts as a chromatin-associated factor
involved in tolerance to heavy metals and oxidative stress (Blanvillain et al., 2009). *AtOXS3* has
been associated with the regulation of gene expression via modulation of histone γ-H2A.X
occupancy (Xiao et al., 2021). *Solyc06g066420* was consistently upregulated across all
interactions and is therefore likely to modulate a wide range of gene expression programs.
Conversely, *Solyc05g052950*, *AtTCF1* ortholog, was strongly downregulated. AtTCF1 has been
studied in the context of abiotic stress, particularly cold acclimatation (Ji et al., 2015). TCF1
indirectly affects lignin accumulation though its association with chromatin at specific loci.
Interestingly, loss of TCF1 modified H3K4me2 and H3K27me3 levels, resulting in reduced lignin
content (Ji et al., 2015). Because lignin-based barrier restricts pathogen spread and contributes to
plant resistance (Lee et al., 2019), *Solyc05g052950* is therefore an excellent epigenetic factor to
target, acting upstream lignin accumulation. Together, these findings underscore the importance of
chromatin-level regulation during parasitism. Furthermore,
several candidate hubs identified here are also associated with abiotic stress responses (Pirona et
al., 2023) reflecting conserved physiological responses—such as ABA signaling, lignin deposition,
and stomatal regulation—between biotic and abiotic stress adaptation. This overlap highlights the
potential cost of manipulating such hubs for resistance breeding.
Besides, subtilases are involved in a broad range of biological functions including plant defense
responses or in plant-pathogen interactions (Weng et al., 2014; Serrano et al., 2016; Stegmann et
al., 2017; J. Figueiredo et al., 2018; Zhang et al., 2024). Within this large gene family (around one
hundred in *S. lycopersicum*), we identified two genes that were strongly upregulated in response
to all three pathogens, while 65 tomato subtilase encoding genes were differentially expressed in
response to at least one of them. This result suggests that subtilases are primarily involved in
pathogen-specific plant responses. In *S. lycopersicum*, research has mainly focused on the
subtilase cluster encoding the P69 proteases, which has been linked to pathogenesis (Jordá et al.,
1999; Zhang et al., 2024). For instance, *Pseudomonas syringae* and *Ralstonia solanacearum*
infections have been associated with the induction of *P69B* among others (Jordá et al., 1999; Zhang
et al., 2024). In our study, *P69B* (*Solyc08g079870*) was also up-regulated in response to *M.*
incognita, *P. parasitica* and *M. euphorbiae*. In contrast, little is known about the role of the subtilases
of the cluster 1, highlighting an important avenue for future studies to determine their role in
pathogenesis.
SA-activated responses are effective against biotrophic pathogens, which retrieve nutrients from
living plant cells. In contrast, JA and ET elicit responses that combat necrotrophic microbes, which
kill plant cells and feed on the resulting tissues. While activation of the JA pathway suppresses SA
biosynthesis and vice versa (Zheng et al., 2012), SA signaling also interferes with JA-ET pathways.
For example, SA affects JA-induced transcription by inducing the degradation of transcription
factors that activate JA signaling, as shown for the ERF transcription factor ORA59. Additionally,
SA can induce negative regulators, among them several WRKY transcription factors, that directly
or indirectly inhibit JA-responsive gene expression (Zheng et al., 2012). More than one hundred
transcription factors were commonly dysregulated in response to the three pathogens tested. We
observed a divergence in the TF family enrichments between the up and down regulated
candidates. Similar results were also observed in transcriptomic analysis of *S. lycopersicum* in
response to osmotic stress (Pirona et al., 2023). Interestingly, our results and those of (Pirona et
al., 2023) showed that, in response to both biotic and abiotic stresses, ERF and bHLH families of
transcription factors were among the top enrichment in up and down regulated candidates,
respectively, with several common candidate genes responding to both types of stresses.
GO enrichment analyses revealed common overrepresentation of metabolic processes across
nematode, aphid, and oomycete infections, supporting the view that parasites enhance host

metabolic activity. Beyond a set of commonly dysregulated candidate hubs, roots and leaves diverged markedly in the significantly overrepresented processes identified across the different co-expression clusters. These findings, along with previous studies reporting tissue-specific responses between leaves and roots (Hermanns et al., 2003; Attard et al., 2010), indicate that combining organ-specific with common hubs may be a more effective strategy for engineering broad-spectrum resistance, potentially through rootstock–scion approaches.

Finally, we identified *Solyc08g006765*, the tomato homolog of *AtNATA1*, as a central multi-pathogen hub. Insect feeding and *P. syringae* inoculation both upregulate NATA1, leading to the production of N^δ-acetylornithine via the jasmonate pathway (Adio et al., 2011). While this metabolite affects the reproduction of the aphid, *Myzus persicae*, *P. syringae* growth was impaired in *nata1* mutant. In addition to jasmonic acid, the coronatine effector produced by DC3000 *P. syringae* strain also induces *NATA1* expression (Lou et al., 2016). Via the acetylation of putrescine, a precursor required for the defense-related hydrogen peroxide, NATA1 acts as a susceptibility factor competing for this common substrate impeding plant defense response (Lou et al., 2016). Furthermore, we noted that several NATA1 interactors, including TCP transcription factors, are also direct targets of known effectors (González-Fuente et al., 2020; Drcelic et al., 2024; Fu et al., 2025). We identified tomato homologs of AtTCP2 and AtTCP21 as consistently dysregulated in presence of all three pathogens. Although *SlTCP2* and *SlTCP21* belong to different classes and are probably involved in two different pathways, their expression is clearly affected by multiple pathogens. Additionally, TCP21, also named CHE for CCA1 Hiking Expedition, was recently described to be the missing link between local infection and the SAR making it a compelling hub (Cao et al., 2024). We propose that these transcription factors and especially *TCP21* (*CHE*), should be considered as a key regulatory hub in future studies of tomato pathogen interactions.

Far beyond the trivial production of a repertoire of commonly dysregulated genes, we implemented an in-depth co-expression analysis that pinpoint compelling candidate hubs covering a large range of functions and pathways. By pairing our results obtained in tomato with Arabidopsis interactome network data, we found crucial factors associated with parasitic success that should be of interest for plant breeding programs.

Acknowledgments

The authors are grateful to Marie-Laure Martin and Etienne Delannoy (IPS2, Orsay, France) for their guidance with the co-expression analysis and Romain Larbat (IRHS, Angers, France) for sharing the RNA-seq datasets of tomato response to *T. absoluta* (Roumani et al., 2022). The authors thank the bioinformatics and genomics platform BIG (ISA, Sophia Antipolis, France, plantBIOs, https://doi.org/10.15454/qyey-ar89) for computing and storage resources and the Genotoul bioinformatics platform (Bioinfo Genotoul, Toulouse, France https://doi.org/10.15454/1.5572369328961167E12) for computing resources.

Data, scripts, code, and supplementary information availability

The datasets generated for this study are available under the BioProject PRJNA1375716. The datasets of tomato response to *T. absoluta* infection at 24 hpi (Roumani et al., 2022) were retrieved through the accession number GSE200795. Supplementary information and code are available at https://doi.org/10.57745/GL5MLM, https://forge.inrae.fr/GNet/dicoexpress/-/tree/985351e4e49c854f1bcb894de585c6939374f8bc/ and https://forge.inrae.fr/plantbios_big/nextflow/mapping_count_featurecount_rsem/-/tree/f9f4b7dd59fb5cc295c446ab9fd067a41956071c/.

Conflict of interest disclosure

The authors declare that they have no financial conflicts of interest in relation to the content of the article.

Funding

This work was funded by the Plant Health and Environment (SPE) division from INRAE (2021-2023 TRIP project) and the CSI-2021 IDEX (2021 ICARE project). BF is supported by INRAE and the French Government (National Research Agency, ANR) through the "Investments for the Future" LabEx SIGNALIFE (#ANR-11-LABX-0028-01) and the MASH project (#ANR-21-CE20-0002).

References

Abad P, Gouzy J, Aury JM, Castagnone-Sereno P, Danchin EG, Deleury E, Perfus-Barbeoch L, Anthouard V, Artiguenave F, Blok VC, Caillaud MC, Coutinho PM, Dasilva C, De Luca F, Deau F, Esquibet M, Flutre T, Goldstone JV, Hamamouch N, Hewezi T, Jaillon O, Jubin C, Leonetti P, Magliano M, Maier TR, Markov GV, McVeigh P, Pesole G, Poulain J, Robinson-Rechavi M, Sallet E, Segurens B, Steinbach D, Tytgat T, Ugarte E, van Ghelder C, Veronico P, Baum TJ, Blaxter M, Bleve-Zacheo T, Davis EL, Ewbank JJ, Favery B, Grenier E, Henrissat B, Jones JT, Laudet V, Maule AG, Quesneville H, Rosso MN, Schiex T, Smant G, Weissenbach J, Wincker P (2008) Genome sequence of the metazoan plant-parasitic nematode Meloidogyne incognita. *Nat Biotechnol,* **26**, 909-915. https://doi.org/10.1038/nbt.1482

Adio AM, Casteel CL, De Vos M, Kim JH, Joshi V, Li B, Juéry C, Daron J, Kliebenstein DJ, Jander G (2011) Biosynthesis and defensive function of N δ-acetylornithine, a jasmonate-induced Arabidopsis metabolite. *The Plant Cell,* **23**, 3303-3318. https://doi.org/10.1105/tpc.111.088989

Attard A, Gourgues M, Callemeyn-Torre N, Keller H (2010) The immediate activation of defense responses in Arabidopsis roots is not sufficient to prevent Phytophthora parasitica infection. *New Phytologist,* **187**, 449-460. https://doi.org/10.1111/j.1469-8137.2010.03272.x

Attard A, Gourgues M, Galiana E, Panabieres F, Ponchet M, Keller H (2008) Strategies of attack and defense in plant–oomycete interactions, accentuated for Phytophthora parasitica Dastur (syn. P. Nicotianae Breda de Haan). *Journal of Plant Physiology,* **165**, 83-94. https://doi.org/10.1016/j.jplph.2007.06.011

Biniaz Y, Tahmasebi A, Tahmasebi A, Albrectsen BR, Poczai P, Afsharifar A (2022) Transcriptome meta-analysis identifies candidate hub genes and pathways of pathogen stress responses in Arabidopsis thaliana. *Biology,* **11**, 1155. https://doi.org/10.3390/biology11081155

Blanvillain R, Kim JH, Wu S, Lima A, Ow DW (2009) OXIDATIVE STRESS 3 is a chromatin-associated factor involved in tolerance to heavy metals and oxidative stress. *The Plant Journal,* **57**, 654-665. https://doi.org/10.1111/j.1365-313X.2008.03717.x

Blok VC, Jones JT, Phillips MS, Trudgill DL (2008) Parasitism genes and host range disparities in biotrophic nematodes: the conundrum of polyphagy versus specialisation. *BioEssays,* **30**, 249-259. https://doi.org/10.1002/bies.20717

Campos MD, Félix MdR, Patanita M, Materatski P, Varanda C (2021) High throughput sequencing unravels tomato-pathogen interactions towards a sustainable plant breeding. *Horticulture Research,* **8**, 171. https://doi.org/10.1038/s41438-021-00607-x

Cao L, Karapetyan S, Yoo H, Chen T, Mwimba M, Zhang X, Dong X (2024) H2O2 sulfenylates CHE, linking local infection to the establishment of systemic acquired resistance. *Science,* **385**, 1211-1217. https://doi.org/10.1126/science.adj7249

Carella P, Evangelisti E, Schornack S (2018) Sticking to it: phytopathogen effector molecules may converge on evolutionarily conserved host targets in green plants. *Current opinion in plant biology,* **44**, 175-180. https://doi.org/10.1016/j.pbi.2018.04.019

Coppola V, Coppola M, Rocco M, Digilio MC, D'Ambrosio C, Renzone G, Martinelli R, Scaloni A, Pennacchio F, Rao R (2013) Transcriptomic and proteomic analysis of a compatible tomato-aphid interaction reveals a predominant salicylic acid-dependent plant response. *BMC genomics,* **14**, 515. https://doi.org/10.1186/1471-2164-14-515

Drcelic M, Skiljaica A, Polak B, Bauer N, Seruga Music M (2024) 'Candidatus Phytoplasma solani'Predicted Effector SAP11-like Alters Morphology of Transformed Arabidopsis Plants and Interacts with AtTCP2 and AtTCP4 Plant Transcription Factors. *Pathogens,* **13**, 893. https://doi.org/10.3390/pathogens13100893

Favery B, Dubreuil G, Chen M-S, Giron D, Abad P (2020) Gall-inducing parasites: convergent and conserved strategies of plant manipulation by insects and nematodes. *Annual Review of Phytopathology,* **58**, 1-22. https://doi.org/10.1146/annurev-phyto-010820-012722

Figueiredo A, Monteiro F, Sebastiana M (2014) Subtilisin-like proteases in plant-pathogen recognition and immune priming: a perspective. *Front Plant Sci,* **5**, 739. https://doi.org/10.3389/fpls.2014.00739

Figueiredo J, Sousa Silva M, Figueiredo A (2018) Subtilisin-like proteases in plant defence: the past, the present and beyond. *Mol Plant Pathol,* **19**, 1017-1028. https://doi.org/10.1111/mpp.12567

Fu M, Zhou Y, Zhang X, Yang K, Xu Y, Wang X, Chen Z, Wang Y, Shi Y, Ma L (2025) A pathogen effector HaRxL10 hijacks the circadian clock component CHE to perturb both plant development and immunity. *Nature Communications,* **16**, 1538. https://doi.org/10.1038/s41467-025-56787-w

González-Fuente M, Carrère S, Monachello D, Marsella BG, Cazalé AC, Zischek C, Mitra RM, Rezé N, Cottret L, Mukhtar MS, Lurin C, Noël LD, Peeters N (2020) EffectorK, a comprehensive resource to mine for Ralstonia, Xanthomonas, and other published effector interactors in the Arabidopsis proteome. *Mol Plant Pathol,* **21**, 1257-1270. https://doi.org/10.1111/mpp.12965

Hao J, Wu W, Wang Y, Yang Z, Liu Y, Lv Y, Zhai Y, Yang J, Liang Z, Huang K (2015) Arabidopsis thaliana defense response to the ochratoxin A-producing strain (Aspergillus ochraceus 3.4412). *Plant Cell Reports,* **34**, 705-719. https://doi.org/10.1007/s00299-014-1731-3

He R, Li Y, Bernards MA, Wang A (2025) Turnip mosaic virus selectively subverts a PR-5 thaumatin-like, plasmodesmal protein to promote viral infection. *New Phytologist,* **245**, 299-317. https://doi.org/10.1111/nph.20233

Hermanns M, Slusarenko AJ, Schlaich NL (2003) Organ-specificity in a plant disease is determined independently of R gene signaling. *Molecular Plant-Microbe Interactions,* **16**, 752-759. https://doi.org/10.1094/MPMI.2003.16.9.752

Jaouannet M, Morris JA, Hedley PE, Bos JI (2015) Characterization of Arabidopsis transcriptional
responses to different aphid species reveals genes that contribute to host susceptibility and non-host
resistance. *PLoS pathogens,* **11**, e1004918. https://doi.org/10.1371/journal.ppat.1004918

Ji H, Wang Y, Cloix C, Li K, Jenkins GI, Wang S, Shang Z, Shi Y, Yang S, Li X (2015) The Arabidopsis
RCC1 family protein TCF1 regulates freezing tolerance and cold acclimation through modulating
lignin biosynthesis. *PLoS genetics,* **11**, e1005471. https://doi.org/10.1371/journal.pgen.1005471.
eCollection 2015

Jones JD, Dangl JL (2006) The plant immune system. *Nature,* **444**, 323-329.
https://doi.org/10.1038/nature05286

Jordá L, Coego A, Conejero V, Vera P (1999) A Genomic Cluster Containing Four Differentially
Regulated Subtilisin-like Processing Protease Genes Is in Tomato Plants *. *Journal of Biological
Chemistry,* **274**, 2360-2365. https://doi.org/10.1074/jbc.274.4.2360

Kamoun S, Furzer O, Jones JD, Judelson HS, Ali GS, Dalio RJ, Roy SG, Schena L, Zambounis A,
Panabières F (2015) The Top 10 oomycete pathogens in molecular plant pathology. *Molecular plant
pathology,* **16**, 413-434. https://doi.org/10.1111/mpp.12190

Kong F, Yang L (2023) Pathogen-triggered changes in plant development: Virulence strategies or host
defense mechanism? *Frontiers in Microbiology,* **14**, 1122947.
https://doi.org/10.3389/fmicb.2023.1122947

Koseoglou E, van der Wolf JM, Visser RGF, Bai Y (2022) Susceptibility reversed: modified plant
susceptibility genes for resistance to bacteria. *Trends in Plant Science,* **27**, 69-79.
https://doi.org/https://doi.org/10.1016/j.tplants.2021.07.018

Lambert I, Paysant-Le Roux C, Colella S, Martin-Magniette ML (2020) DiCoExpress: a tool to process
multifactorial RNAseq experiments from quality controls to co-expression analysis through differential
analysis based on contrasts inside GLM models. *Plant Methods,* **16**, 68.
https://doi.org/10.1186/s13007-020-00611-7

Lee MH, Jeon HS, Kim SH, Chung JH, Roppolo D, Lee HJ, Cho HJ, Tobimatsu Y, Ralph J, Park OK
(2019) Lignin-based barrier restricts pathogens to the infection site and confers resistance in plants.
The EMBO journal, **38**, e101948. https://doi.org/10.15252/embj.2019101948

Lou YR, Bor M, Yan J, Preuss AS, Jander G (2016) Arabidopsis NATA1 Acetylates Putrescine and
Decreases Defense-Related Hydrogen Peroxide Accumulation. *Plant Physiol,* **171**, 1443-1455.
https://doi.org/10.1104/pp.16.00446

Mejias J, Bazin J, Truong N-M, Chen Y, Marteu N, Bouteiller N, Sawa S, Crespi MD, Vaucheret H, Abad
P (2021) The root-knot nematode effector MiEFF18 interacts with the plant core spliceosomal protein
SmD1 required for giant cell formation. *The New Phytologist,* **229**, 3408-3423.
https://doi.org/10.1111/nph.17089

Moeder W, Urquhart W, Ung H, Yoshioka K (2011) The Role of Cyclic Nucleotide-Gated Ion Channels
in Plant Immunity. *Molecular Plant,* **4**, 442-452. https://doi.org/https://doi.org/10.1093/mp/ssr018

Mukherjee AK, Carp M-J, Zuchman R, Ziv T, Horwitz BA, Gepstein S (2010) Proteomics of the response
of Arabidopsis thaliana to infection with Alternaria brassicicola. *Journal of proteomics,* **73**, 709-720.
https://doi.org/10.1016/j.jprot.2009.10.005

Mukhtar MS, Carvunis AR, Dreze M, Epple P, Steinbrenner J, Moore J, Tasan M, Galli M, Hao T,
Nishimura MT, Pevzner SJ, Donovan SE, Ghamsari L, Santhanam B, Romero V, Poulin MM,
Gebreab F, Gutierrez BJ, Tam S, Monachello D, Boxem M, Harbort CJ, McDonald N, Gai L, Chen
H, He Y, Vandenhaute J, Roth FP, Hill DE, Ecker JR, Vidal M, Beynon J, Braun P, Dangl JL (2011)
Independently evolved virulence effectors converge onto hubs in a plant immune system network.
Science, **333**, 596-601. https://doi.org/10.1126/science.1203659

Munir S, Mumtaz MA, Ahiakpa JK, Liu G, Chen W, Zhou G, Zheng W, Ye Z, Zhang Y (2020) Genome-
wide analysis of Myo-inositol oxygenase gene family in tomato reveals their involvement in ascorbic
acid accumulation. *BMC genomics,* **21**, 284. https://doi.org/10.1186/s12864-020-6708-8

Naveed ZA, Ali GS (2018) Comparative transcriptome analysis between a resistant and a susceptible
wild tomato accession in response to Phytophthora parasitica. *International Journal of Molecular*
Sciences, **19**, 3735. https://doi.org/10.3390/ijms19123735

Noutoshi Y, Kuromori T, Wada T, Hirayama T, Kamiya A, Imura Y, Yasuda M, Nakashita H, Shirasu K,
Shinozaki K (2006) Loss of Necrotic Spotted Lesions 1 associates with cell death and defense
responses in Arabidopsis thaliana. *Plant Mol Biol,* **62**, 29-42. https://doi.org/10.1007/s11103-006-
9001-6

Park E-J, Kim T-H (2021) Arabidopsis OSMOTIN 34 functions in the ABA signaling pathway and is
regulated by proteolysis. *International Journal of Molecular Sciences,* **22**, 7915.
https://doi.org/10.3390/ijms22157915

Pirona R, Frugis G, Locatelli F, Mattana M, Genga A, Baldoni E (2023) Transcriptomic analysis reveals
the gene regulatory networks involved in leaf and root response to osmotic stress in tomato. *Front*
Plant Sci, **14**, 1155797. https://doi.org/10.3389/fpls.2023.1155797

Reichardt S, Repper D, Tuzhikov AI, Galiullina RA, Planas-Marquès M, Chichkova NV, Vartapetian AB,
Stintzi A, Schaller A (2018) The tomato subtilase family includes several cell death-related
proteinases with caspase specificity. *Scientific Reports,* **8**, 10531. https://doi.org/10.1038/s41598-
018-28769-0

Roberts R, Hind SR, Pedley KF, Diner BA, Szarzanowicz MJ, Luciano-Rosario D, Majhi BB, Popov G,
Sessa G, Oh C-S (2019) Mai1 protein acts between host recognition of pathogen effectors and
mitogen-activated protein kinase signaling. *Molecular Plant-Microbe Interactions,* **32**, 1496-1507.
https://doi.org/10.1094/MPMI-05-19-0121-R

Roumani M, Le Bot J, Boisbrun M, Magot F, Péré A, Robin C, Hilliou F, Larbat R (2022) Transcriptomics
and Metabolomics Analyses Reveal High Induction of the Phenolamide Pathway in Tomato Plants
Attacked by the Leafminer Tuta absoluta. *Metabolites,* **12**. https://doi.org/10.3390/metabo12060484

Serrano I, Buscaill P, Audran C, Pouzet C, Jauneau A, Rivas S (2016) A non canonical subtilase
attenuates the transcriptional activation of defence responses in Arabidopsis thaliana. *eLife,* **5**,
e19755. https://doi.org/10.7554/eLife.19755

Spartz AK, Lor VS, Ren H, Olszewski NE, Miller ND, Wu G, Spalding EP, Gray WM (2016) Constitutive
Expression of Arabidopsis SMALL AUXIN UP RNA19 (SAUR19) in Tomato Confers Auxin-
Independent Hypocotyl Elongation. *Plant physiology,* **173**, 1453-1462.
https://doi.org/10.1104/pp.16.01514

Stegmann M, Monaghan J, Smakowska-Luzan E, Rovenich H, Lehner A, Holton N, Belkhadir Y, Zipfel
C (2017) The receptor kinase FER is a RALF-regulated scaffold controlling plant immune signaling.
Science, **355**, 287-289. https://doi.org/10.1126/science.aal2541

Thomazella DPT, Seong K, Mackelprang R, Dahlbeck D, Geng Y, Gill US, Qi T, Pham J, Giuseppe P,
Lee CY, Ortega A, Cho MJ, Hutton SF, Staskawicz B (2021) Loss of function of a DMR6 ortholog in
tomato confers broad-spectrum disease resistance. *Proc Natl Acad Sci U S A,* **118**.
https://doi.org/10.1073/pnas.2026152118

Tian F, Yang D-C, Meng Y-Q, Jin J, Gao G (2019) PlantRegMap: charting functional regulatory maps in
plants. *Nucleic Acids Research,* **48**, D1104-D1113. https://doi.org/10.1093/nar/gkz1020

Valdés AE, Övernäs E, Johansson H, Rada-Iglesias A, Engström P (2012) The homeodomain-leucine
zipper (HD-Zip) class I transcription factors ATHB7 and ATHB12 modulate abscisic acid signalling
by regulating protein phosphatase 2C and abscisic acid receptor gene activities. *Plant Molecular*
Biology, **80**, 405-418. https://doi.org/10.1007/s11103-012-9956-4

van Butselaar T, Silva S, Lapin D, Bañales I, Tonn S, van Schie C, Van den Ackerveken G (2024) The
role of salicylic acid in the expression of RECEPTOR-LIKE PROTEIN 23 and other immunity-related
genes. *Phytopathology®,* **114**, 1097-1105. https://doi.org/10.1094/PHYTO-10-23-0413-KC

van Schie CC, Takken FL (2014) Susceptibility genes 101: how to be a good host. *Annual Review of*
Phytopathology, **52**, 551-581. https://doi.org/ 10.1146/annurev-phyto-102313-045854

Walling LL (2008) Avoiding effective defenses: strategies employed by phloem-feeding insects. *Plant*
physiology, **146**, 859-866. https://doi.org/10.1104/pp.107.113142

Wang D, Mandal P, Rahman MS, Yang L (2025) Engineering tomato disease resistance by manipulating
susceptibility genes. *Frontiers in Genome Editing,* **7**, 1537148.
https://doi.org/10.3389/fgeed.2025.1537148

Wang F, Park Y-L, Gutensohn M (2021) Epidermis-specific metabolic engineering of sesquiterpene
formation in tomato affects the performance of potato aphid Macrosiphum euphorbiae. *Frontiers in*
plant science, **12**, 793313. https://doi.org/10.3389/fpls.2021.793313

Wang Y, Li T, Sun Z, Huang X, Yu N, Tai H, Yang Q (2022) Comparative transcriptome meta-analysis
reveals a set of genes involved in the responses to multiple pathogens in maize. *Frontiers in plant*
science, **13**, 971371. https://doi.org/10.3389/fpls.2022.971371

Weigel RR, Pfitzner UM, Gatz C (2005) Interaction of NIMIN1 with NPR1 modulates PR gene expression
in Arabidopsis. *The Plant Cell,* **17**, 1279-1291. https://doi.org/10.1105/tpc.104.027441

Weng K, Li Z-Q, Liu R-Q, Wang L, Wang Y-J, Xu Y (2014) Transcriptome of Erysiphe necator-infected
Vitis pseudoreticulata leaves provides insight into grapevine resistance to powdery mildew.
Horticulture Research, **1**. https://doi.org/10.1038/hortres.2014.49

Xiao S, Jiang L, Wang C, Ow DW (2021) Arabidopsis OXS3 family proteins repress ABA signaling
through interactions with AFP1 in the regulation of ABI4 expression. *Journal of Experimental Botany,*
72, 5721-5734. https://doi.org/ 10.1093/jxb/erab237

Zavaliev R, Mohan R, Chen T, Dong X (2020) Formation of NPR1 condensates promotes cell survival
during the plant immune response. *Cell,* **182**, 1093-1108. e1018.
https://doi.org/10.1016/j.cell.2020.07.016

Zeilmaker T, Ludwig NR, Elberse J, Seidl MF, Berke L, Van Doorn A, Schuurink RC, Snel B, Van den
Ackerveken G (2015) DOWNY MILDEW RESISTANT 6 and DMR 6-LIKE OXYGENASE 1 are
partially redundant but distinct suppressors of immunity in Arabidopsis. *The Plant Journal,* **81**, 210-
222. https://doi.org/10.1111/tpj.12719

Zhang W, Planas-Marquès M, Mazier M, Šimkovicová M, Rocafort M, Mantz M, Huesgen PF, Takken
FLW, Stintzi A, Schaller A, Coll NS, Valls M (2024) The tomato P69 subtilase family is involved in
resistance to bacterial wilt. *Plant J,* **118**, 388-404. https://doi.org/10.1111/tpj.16613

Zheng XY, Spivey NW, Zeng W, Liu PP, Fu ZQ, Klessig DF, He SY, Dong X (2012) Coronatine promotes
Pseudomonas syringae virulence in plants by activating a signaling cascade that inhibits salicylic
acid accumulation. *Cell Host Microbe,* **11**, 587-596. https://doi.org/10.1016/j.chom.2012.04.014

Zhou J, Tang X, Martin GB (1997) The Pto kinase conferring resistance to tomato bacterial speck
disease interacts with proteins that bind a cis-element of pathogenesis-related genes. *Embo j,* **16**,
3207-3218. https://doi.org/10.1093/emboj/16.11.3207
768