1 Neutrophils in patients with chronic coronary syndrome exhibit

2 delayed spontaneous apoptosis and resistance to regulatory T cell-

3 induced apoptosis – Brief Report

Maike Schneider[1], Camilla Skoglund[1], Rosanna WS Chung[1], Lena Jonasson[1,2]

[1]Department of Health, Medicine, and Caring Sciences, Linköping University,

Linköping, Sweden, [2]Clinical Department of Cardiology, Region Östergötland,

Linköping, Sweden

Short title: Delayed neutrophil apoptosis in coronary patients

Corresponding author: Lena Jonasson, lena.jonasson@liu.se

Division of Diagnostics and Specialist Medicine, Department of Health, Medicine, and

Caring Sciences, Linköping University, SE-58183 Linköping, Sweden.

Total word count: 4924

Abstract

Background: Persistent inflammation is linked to poor outcomes in patients with chronic coronary syndrome (CCS). The inflammatory state in atherosclerotic disease has been associated with activation of neutrophils as well as with regulatory T cell (T_{reg}) deficiency. The role of T_{regs} in the regulation of neutrophil survival has been postulated recently. Here, we investigated neutrophil apoptosis along with the potential impact of T_{regs} on neutrophil apoptosis in patients with CCS, compared to healthy controls.

Methods: Twenty patients with CCS and 19 healthy controls were included. Neutrophil apoptosis was assessed after 5h culture with or without interleukin(IL-)10, TNF or LPS. Neutrophil phenotype was evaluated through flow cytometry analysis of surface receptors (CD66b and CXCR4) and *ex vivo* release of cytokines and granule proteins. The ability of T_{regs} to induce neutrophil apoptosis was examined in autologous neutrophil-T_{reg} co-cultures.

Results: Spontaneous neutrophil apoptosis was significantly delayed in patients compared to controls (10.3% vs. 19.2%, p=0.025). Also, neutrophils from patients overexpressed CD66b and CXCR4 and were more prone to release proinflammatory mediators. Notably, T_{regs} induced neutrophil apoptosis in healthy subjects, but not in patients, indicating a loss of T_{reg}-mediated regulation in the latter. There was no evidence that IL-10 had any influence on neutrophil apoptosis. However, cell-to-cell contact was found essential for T_{reg}-induced neutrophil apoptosis.

Conclusions: Patients with CCS display delayed neutrophil apoptosis and a pro-inflammatory neutrophil phenotype that is resistant to T_{reg}-mediated apoptosis.

44   Neutrophil dysfunction may contribute to persistent inflammation in patients with

45   CCS, and as such constitute a novel therapeutic target.

47   **Key words:**

48   Coronary artery disease, Chronic coronary syndrome, Inflammation, Apoptosis,

49   Neutrophils, Regulatory T cells

Introduction

The relationship between inflammation and atherosclerosis is well established, as

shown in experimental, epidemiological and clinical studies [1]. Despite optimal

management of classical risk factors like hyperlipidemia and hypertension, systemic

low-grade inflammation promotes recurrent cardiovascular events in patients with

chronic coronary syndrome (CCS), a condition former known as stable coronary

artery disease [2].

Growing evidence implicates neutrophils as pivotal mediators in atherosclerotic

inflammation [3]. Epidemiological studies consistently link elevated neutrophil counts to

incident atherosclerotic cardiovascular disease and mortality [4]. Increased neutrophil

activation is also a well-known feature of the acute coronary syndrome (ACS) [5-7],

while the occurrence of neutrophil activation in patients with CCS has been less

studied. However, elevated neutrophil counts as well as neutrophil granule proteins,

including myeloperoxidase (MPO), matrix metalloprotease (MMP)-8 and MMP-9,

have been reported in patients with CCS, indicating a heightened neutrophil

activation status [8-11]. A few previous studies have also investigated whether

neutrophils from patients with CCS are more prone to activation *ex vivo* compared to

neutrophils from healthy controls, although with inconsistent results [12-14].

Neutrophils have a very short life, around 6 h, in circulating blood, They are

inherently programmed to migrate back to the bone marrow via CXCR4 expression

and die by constitutive apoptosis [15]. Neutrophil apoptosis is considered to be a critical

mechanism for resolving acute inflammation. A delay in the apoptotic process has been

associated with increased neutrophil effector functions [16,17]. Accordingly, delayed

neutrophil apoptosis has been observed in a number of chronic inflammatory

disorders [18-20]. A few previous studies have also described delayed neutrophil

apoptosis in patients with coronary artery disease, though with focus on ACS [21,22].

Whether a delay in neutrophil apoptosis occurs in the stabilized phase after an ACS

has not been clarified.

In patients with CCS, reduced numbers as well as impaired function of regulatory T

cells (T_{regs}) have been a consistent finding [23-25]. T_{regs} are primarily recognized for their

ability to modulate adaptive immunity. However, it has also been demonstrated that

murine as well as human T_{regs} are able to promote neutrophil apoptosis and inhibit

neutrophil activity [26,27]. Interestingly, the role of dysregulated neutrophil-T_{reg}

interaction in chronic inflammatory diseases, such as rheumatoid arthritis and

systemic lupus erythematosus, has been proposed [28,29]. Whether a dysregulated

neutrophil-T_{reg} interaction takes place in patients with CCS has not previously been

explored.

In the present study, we hypothesized that delayed neutrophil apoptosis is a feature

of the persistent inflammatory state in patients with CCS. We examined neutrophil

apoptosis, along with neutrophils´ activation/aging status and their susceptibility to

autologous T_{regs}, in patients with CCS and healthy control subjects.

Methods

The data that support the findings of this study are available from the corresponding

author upon reasonable request. A detailed, expanded method sectionis is available

in the Data Supplement.

Study population

Twenty patients with CCS were recruited; all had angiographically verified coronary

artery disease and a prior history of ACS. Patients were not eligible if they were more

than 75 years of age, suffered from severe heart failure, immunological disorders,

neoplastic disease, received treatment with immunosuppressive or anti-inflammatory

agents (except low-dose aspirin), or had a history of acute or recent (<2 months)

infection, major trauma/surgery or any revascularization procedure. For the control

group (n = 19), volunteers with generally equal age and sex distribution were

randomly selected from the Swedish Population Register. The study was conducted

in accordance with the ethical guidelines of the Declaration of Helsinki, and the

protocol was approved by the Ethical Review Board of Linkoping. Written informed

consent was obtained from all study participants.

Cell isolation and culture

Venous blood was collected in the morning in a fasting state. $CD4^+CD25^+CD127^{-/low}$

T_{regs} were isolated from peripheral blood mononuclear cells (PBMCs) using an

EasySep Human CD4+CD127lowCD25+ Regulatory T Cell Isolation Kit, and CD4+ T

cells were isolated from PBMCs using a CD4+ T Cell Isolation Kit human. Neutrophils

were isolated from whole blood using Polymorphprep.

For immediate staining, neutrophils were incubated with mouse anti-human CXCR4

APC and mouse anti-human CD66b PE antibodies for 15 min (Table S1). They were

analysed within 1h using a Gallios Flow Cytometer.

Pure neutrophils, or co-cultures with neutrophils and T_{regs} or $CD4^+$ T cells were

seeded in 96-well plates at concentrations of 0.5 x10^6 cells/ml for 2.5 h or 5 h, with or

without interleukin (IL)-10, tumor necrosis factor (TNF) or lipopolysaccharide from *E.*

coli (LPS).

Neutrophil apoptosis and activation

Neutrophils were incubated for either 2.5 h or 5 h in different series of experiments.

After 2.5 h, supernatants were stored at -80°C until further analysis. After 5 h, cells

were stained with mouse anti-human CD66b PE, anti-human Annexin V BV42 and

mouse anti-human CD4 BV510 antibodies (if the sample contained T_{regs} or CD4[+]

cells) in Annexin V Binding Buffer, followed by staining with SYTOX Red dead cell

stain. Cells were analysed within 1 h using a Gallios Flow Cytometer to monitor

apoptosis and activation status of neutrophils. Gating strategies are shown in Figure

S1.

<u>Transwell experiments</u>

Isolated neutrophils and T_{regs} were seeded in a 48-well cell culture plate with or

without Millicell Cell Culture Inserts.

<u>Proteins in plasma and supernatants</u>

Tumor necrosis factor (TNF), interferon(IFN)-γ, interleukin(IL)-6, IL-1β and IL-18

were measured in plasma via Olink Target 48, with an inter-assay CV of 7% and an

intra-assay CV of 6%. Matrix metalloproteinase(MMP)-9, MMP-8, myeloperoxidase

(MPO) in plasma were measured using a Luminex Multiplex Assay, with an intra-

assay CV of 3.2%. MMP-9, MMP-8, MPO, TNF, IFN-γ, IL-6, IL-10, IL-1β and IL-18

were also measured in supernatants using a Luminex Multiplex Assay, with an inter-

assay CV of 16.6% and an intra-assay CV of 8.1%. Measuements were performed by

the Affinity Proteomics-Unit of SciLifeLab, Stockholm, Sweden.

<u>Statistics</u>

All statistics were performed using the software IBM SPSS Statistics version

29.0.2.0. For comparisons between groups, a Mann-Whitney U test was performed.

For comparison of paired data from the same individual, a Wilcoxon signed rank test

was performed. For analysis of concentration-dependency, the Greenhouse-Geisser

Test was performed.

Results

<u>Cohort characteristics</u>

All patients with CCS were clinically and metabolically stable with no symptoms of

angina. There were no differences between patients and controls regarding age

(median age 65 *vs* 64 years), sex distribution (females 50 % *vs* 53 %), body mass

index or smoking status. Also, the protein levels in plasma were comparable between

groups, except for higher levels of TNF and IL-10, as well as slightly but not

significantly higher levels of IL-6, in the CCS group. Blood samples were collected

from patients at a median of 11.5 months (interquartile range: 5.8-13 months) after an

ACS. Cohort characteristics are presented in Table S2.

<u>Neutrophil apoptosis is delayed in patients with CCS</u>

The apoptotic rate of neutrophils was assessed after 5 h culture in medium only

(spontaneous apoptosis) or in the presence of either anti-inflammatory (IL-10) or pro-

inflammatory (TNF, LPS) stimuli. As shown in Figure 1A, the spontaneous apoptosis

was significantly reduced in CCS patients compared to healthy controls (10.3% vs.

19.2%, $p = 0.025$). A similar result was seen after 5 h treatment with IL-10 (10.1% vs.

19.8%, $p = 0.020$). Under inflammatory conditions (TNF or LPS), the differences in

neutrophil apoptosis between patients and controls did not reach statistical

significance.

<u>Neutrophils exhibit an aged and pro-inflammatory phenotype in patients with CCS</u>

The surface expression of CD66b was used as a marker of neutrophil activation [30].

After 5 h in medium only or in the presence of IL-10, TNF, or LPS the expression of

CD66b was significantly higher in patients than in controls (Figure 1B).

The surface expression of CD66b and CXCR4, the latter representing neutrophil

aging [15], was also assessed immediately after cell isolation. Neutrophils from CCS

patients exhibited increased surface expression of CD66b as well as increased

expression of CXCR4, compared to neutrophils from healthy subjects (Figure 1C-D).

To further investigate the neutrophil activation status, we measured the release of

neutrophil granule proteins (MMP-9, MMP-8, MPO) and cytokines (TNF, IFN-γ, IL-6,

IL-18, IL-1β) after a shorter period of incubation (2.5 h) with or without IL-10, TNF or

LPS. There were no differences between patients and controls when neutrophils

were cultured in medium only. However, upon LPS stimulation, the neutrophils from

patients secreted higher levels of MMP-9, MMP-8, TNF, IFN-γ, IL-6 and IL-1β

compared to controls. Also, the neutrophils from patients released higher levels of

MMP-9 upon TNF treatment (Figure 1E).

<u>T_{regs} induce neutrophil apoptosis in healthy controls, but not in patients with CCS</u>

We then proceeded to examine the effect of autologous T_{regs} on neutrophil apoptosis

by co-culturing the cells at varying ratios (neutrophils:T_{regs} 1:0, 10:1, 4:1 and 1:1) for

5 h with or without IL-10, TNF or LPS. In the controls, T_{regs} induced neutrophil

apoptosis in a concentration-dependent manner under all conditions, whereas no

such effect was observed in the patients (Figure 2A-B). When T_{reg}-induced neutrophil

apoptosis was compared between patients and controls, the controls exhibited a

significantly higher apoptotic rate at 5 h (with or without IL-10) at all ratios (Figures

S2A-B). Autologous CD4$^+$ T cells did not induce neutrophil apoptosis, neither in

healthy controls nor in CCS patients (Figure S2A-D for between-groups analysis,

Figure S3 for within-group analysis).

T_{reg} survival and IL-10 secretion in neutrophil-T_{reg} co-cultures do not differ between

patients with CCS and healthy controls

To determine whether differences in T_{reg} survival contributed to the discrepancy in

T_{reg}-induced neutrophil apoptosis between groups, we assessed the T_{reg} viability

under various conditions. Neither T_{reg} nor CD4$^+$ T cell survival differed between

patients and controls under any conditions (Figure S4A-B).

To gain an insight into the functionality of T_{regs}, IL-10 was measured in co-culture

supernatants of patients and controls. In controls, the concentrations of IL-10

appeared to increase at ratios 4:1 and 1:1 (Figure S4C-D). However, the absolute

levels were low (0.2 - 1.1 pg/ml), and no significant differences were observed

between patients and controls.

T_{reg}-induced neutrophil apoptosis is not dependent on IL-10, but rather on cell-to-cell

contact

Next, we investigated two possible mechanisms of T_{reg}-induced neutrophil apoptosis.

To determine whether IL-10 was a potential mediator of T_{reg}-induced neutrophil

apoptosis, neutrophils from healthy volunteers were treated with increasing

concentrations of IL-10 (1 pg/ml, 1 ng/ml, 100 ng/ml) for 5 h, mimicking

concentrations in co-culture supernatants (mentioned above) as well as

concentrations used in previous *in vitro* studies [31,32]. None of the IL-10 concentrations

showed any effect on the apoptotic rate (Figure 2D). To further assess whether T_{regs}

used cell-to-cell contact to regulate neutrophil apoptosis, transwell experiments were

performed to physically separate T_{regs} and neutrophils via a membrane (mesh) at

ratio 4:1. As shown in Figure 2E, the separation completely abrogated T_{reg}-induced

neutrophil apoptosis.

Discussion

The main finding of the present study is that spontaneous neutrophil apoptosis, a key

process in the resolution of inflammation, was significantly delayed in patients with

CCS compared to healthy controls. Concomitantly, the neutrophils in patients

exhibited an aged and proinflammatory phenotype as well as a resistance to T_{reg}-

induced neutrophil apoptosis.

Delayed neutrophil apoptosis has been described in several chronic inflammatory

diseases [18-20]. Only a few earlier studies have investigated neutrophil death in

coronary artery disease, with emphasis on ACS, a condition characterized by acute

inflammation. In 2004, Garlich *et al.* [21] demonstrated that neutrophil death, including

both apoptosis and necrosis, was delayed in patients with ACS (n=30), compared to

patients with stable angina (n=15) and controls (n=15) after 24 h culture. Some years

later, Biasucci *et al.* [22] measured spontaneous neutrophil apoptosis in patients with

ACS (n=30), patients with stable angina (n=13) and controls (n=34). They found

delayed neutrophil apoptosis in ACS patients after 4 h culture (10.1%), whereas the

apoptotic rate in stable angina patients (28.5%) did not differ from that in controls

(41.8%). In the present study, we detected a delay in neutrophil apoptosis already

after 5 h in patients with CCS compared to controls. Comparisons with earlier study

groups should however be drawn with caution, e.g. the apoptotic rates reported by

Biasucci *et al.* [22] after 4 h culture were remarkably higher than what we found, which

might indicate methodological differences. Also, we included patients who had

survived ACS, while Biasucci *et al.* [22] included stable angina patients without prior

ACS. Despite being asymptomatic and without overt heart failure, it cannot be

excluded that our patient cohort had a more severe, complicated coronary artery

disease, hence a more proinflammatory neutrophil phenotype.

It is well established that patients with ACS exhibit a more active neutrophil

phenotype, either within culprit coronary lesions or circulating in peripheral blood [3].

Whether the active neutrophil phenotype is maintained in those stabilized after a prior

ACS has been less clear. Results from *ex vivo* functional assays are sparse; some

have reported that circulating neutrophils in patients with CCS are primed, i.e. more

prone to stimulation *ex vivo*, as compared to neutrophils in healthy controls [12,13],

while others have not been able to detect any differences [14,33]. In the present study,

delayed spontaneous neutrophil apoptosis in patients with CCS was clearly

accompanied by increased expression of CD66b, a sensitive marker of neutrophil

activation in human samples [30]. Also, the finding that patients´neutrophils were more

prone to secrete cytokines and granule proteins upon stimulation *ex vivo* supported

their state of heightened activity. Furthermore, the increased expression of CXCR4

indicate an accumulation of aged neutrophils in the circulation of patients with CCS.

Aged neutrophils have been shown to exhibit a distinct, highly reactive phenotype,

both in experimental models and in chronic inflammatory disease [34,35].

Statins are known to exert pleiotropic anti-inflammatory effects, one of which is the

suppression of neutrophil activation [36,37]. Furthermore, two single-center randomized

trials have shown that statins promote neutrophil apoptosis in acute inflammatory

conditions [38,39]. Despite extensive treatment with statins, CCS patients in our study

exhibited an increase in proinflammatory neutrophils with delayed apoptosis. These

findings indicate that statin-mediated immunomodulatory effects on innate immunity,

while robust *in vitro*, may be less effective or relevant in chronic inflammation.

T_{regs} are well known for their role in controling the adaptive immune response [29].

Emerging evidence indicates that T_{regs} also have a regulatory role in the innate

immune response [29]. Of note, a deficit in circulating T_{regs} is a characteristic finding in

patients with both acute and chronic manifestations of coronary artery disease [23-25],

as well as in patients with other chronic inflammatory diseases [40-42]. Earlier

experimental studies, mainly performed on murine cells, have shown that LPS-

activated T_{regs} are able to inhibit neutrophil function and promote neutrophil apoptosis

[26,27]. In the present study, T_{regs} promoted neutrophil apoptosis both with and without

LPS in healthy controls, while this phenomenon did not occur in patients with CCS.

The reason for resistance to T_{reg}-induced neutrophil apoptosis may be multifactorial,

depending on anti-apoptotic features of pro-inflammatory neutrophils [43] but also on

functional defects of T_{regs}. As previously described by our group, the capacity of T_{regs}

to regulate proliferation and secretion of IFN-γ and IL-10 in autologous T responder

cells is impaired in patients with CCS compared to healthy controls [25]. Several

studies have also shown that a chronic inflammatory environment may lead to

impaired T_{reg} function depending on molecular mechanisms, such as perturbed

Foxp3 expression and epigenetic modifications [44].

IL-10 has been described as a mediator of anti-inflammatory effects exerted by T_{regs}

[45,46]. In the present study, we used a wide range of concentrations of IL-10 in both

neutrophil cultures and neutrophil-T_{reg} co-cultures, but were not able to demonstrate

any effect on neutrophil apoptosis after 5 h. This is in agreement with earlier studies

reporting that IL-10 does not affect spontaneous neutrophil apoptosis, neither after

short (6 h) nor long (20 h) incubation [47,48]. However, these earlier studies have also

reported that, under *in vitro* conditions mimicking severe sepsis, IL-10 is capable of

counteracting LPS-induced delay of neutrophil apoptosis [31,47,48]. Moreover, LPS-

stimulated T_{regs} have been shown to induce IL-10 production in apoptotic neutrophils

after 5 h co-culture, while LPS alone or unstimulated T_{regs} do not [49]. In the present

study, we detected very low concentrations of IL-10 in supernatants after 5 h co-

culture, both with and without LPS, speaking against a significant role of IL-10 in this

experimental model for T_{reg}-induced neutrophil apoptosis. On the other hand, our

findings clearly show that T_{reg}-induced neutrophil apoptosis was dependent on cell-

to-cell contact. The results are in agreement with previous studies reporting that

human T_{regs} exert suppressive effects through cell-to-cell contact [49,50].

Several limitations of our study should be acknowledged. The neutrophil-T_{reg} co-

cultures were performed at cell ratios that do not reflect physiological conditions in

peripheral blood where the neutrophil:T_{regs} ratio is normally 100:1. However, the *in*

vitro approach using ratios from 1:1 to 10:1 is in line with previous mechanistic

studies [26,27,51]. Another major limitation is that we did not measure the phenotype of

isolated T_{regs}, which limits the understanding of why neutrophil-T_{reg} interaction was

impaired in patients compared to controls. Lastly, the sample size was small and our

findings should be considered preliminary until validated in a larger cohort.

Nevertheless, a strength of the study is the *ex vivo* investigation of peripheral

neutrophils in human disease, addressing a critical gap in translational cardiovascular

research. While numerous epidemiological studies have shown that high neutrophil

counts is a causal risk factor for cardiovascular disease [4], mechanistic insights have

been mainly derived from animal models [3].

To conclude, delayed spontaneous neutrophil apoptosis, along with a more active

neutrophil phenotype showing resistance to T_{reg}-induced apoptosis, indicate that

neutrophils may be a major contributor to the non-resolved inflammatory state in

patients with CCS. Of note, neutrophil dysfunction was observed in clinically stable

patients who received optimal medical therapy according to current guidelines.

Although the findings need to be confirmed in larger samples, and also explored in

further mechanistic studies, they allow us to speculate about neutrophil apoptosis as

a future target of therapy in patients with CCS.

Acknowledgements

We acknowledge the Core Facility at the Faculty of Medicine and Health Sciences,

Linköping University for providing assistance in flow cytometry and the Affinity

Proteomics Unit at SciLifeLab in Stockholm for supporting with protein quantification.

Sources of funding

This study was financially supported by Heart–Lung Foundation, Sweden

(20210436), and Swedish Research Council (2021–02262).

Disclosures

None.

References

1.   Libby P, Hansson GK. From Focal Lipid Storage to Systemic Inflammation: JACC

Review Topic of the Week. *Journal of the American College of Cardiology*.

2019;74:1594–1607. doi: https://doi.org/10.1016/j.jacc.2019.07.061

2.   Libby P, Soehnlein O. Inflammation in atherosclerosis: Lessons and therapeutic

implications. *Immunity*. 2025;58:2383–2401. doi:

https://doi.org/10.1016/j.immuni.2025.09.012

3.   Döring Y, Drechsler M, Soehnlein O, Weber C. Neutrophils in Atherosclerosis.

Arteriosclerosis, Thrombosis, and Vascular Biology. 2015;35:288–295. doi:

doi:10.1161/ATVBAHA.114.303564

4.   Luo J, Thomassen JQ, Nordestgaard BG, Tybjærg-Hansen A, Frikke-Schmidt R.

Neutrophil counts and cardiovascular disease. *European Heart Journal*.

2023;44:4953–4964. doi: 10.1093/eurheartj/ehad649

5.   Bonaventura A, Montecucco F, Dallegri F, Carbone F, Lüscher TF, Camici GG,

Liberale L. Novel findings in neutrophil biology and their impact on cardiovascular

disease. *Cardiovasc Res*. 2019;115:1266–1285. doi: 10.1093/cvr/cvz084

6.   Talmor N, Pillinger MH, Xia Y, Leonard A, Curovic F, Shah B. Neutrophil Activation

and Adhesiveness in Coronary Artery Disease: Results From the

COLCHICINE‐PCI Biomarker Substudy. *Journal of the American Heart

Association*. 2024;13:e036701. doi: doi:10.1161/JAHA.124.036701

7.   Maréchal P, Tridetti J, Nguyen ML, Wéra O, Jiang Z, Gustin M, Donneau AF, Oury C,

Lancellotti P. Neutrophil Phenotypes in Coronary Artery Disease. *J Clin Med*. 2020;9.

doi: 10.3390/jcm9051602

8.   Momiyama Y, Ohmori R, Tanaka N, Kato R, Taniguchi H, Adachi T, Nakamura H,

Ohsuzu F. High plasma levels of matrix metalloproteinase-8 in patients with unstable

angina. *Atherosclerosis*. 2010;209:206–210. doi:

10.1016/j.atherosclerosis.2009.07.037

9.   Caselli C, Di Giorgi N, Ragusa R, Lorenzoni V, Smit J, El Mahdiui M, Buechel RR,

Teresinska A, Pizzi MN, Roque A, et al. Association of MMP9 with adverse features of

plaque progression and residual inflammatory risk in patients with chronic coronary

syndrome (CCS). *Vascul Pharmacol*. 2022;146:107098. doi:

10.1016/j.vph.2022.107098

10. Nicholls SJ, Hazen SL. Myeloperoxidase and Cardiovascular Disease.

Arteriosclerosis, Thrombosis, and Vascular Biology. 2005;25:1102–1111. doi:

doi:10.1161/01.ATV.0000163262.83456.6d

11. Avanzas P, Arroyo-Espliguero R, Cosín-Sales J, Quiles J, Zouridakis E, Kaski JC.

Multiple complex stenoses, high neutrophil count and C-reactive protein levels in

patients with chronic stable angina. *Atherosclerosis*. 2004;175:151–157. doi:

10.1016/j.atherosclerosis.2004.03.013

12. Paulsson J, Dadfar E, Held C, Jacobson SH, Lundahl J. Activation of peripheral and

in vivo transmigrated neutrophils in patients with stable coronary artery disease.

Atherosclerosis. 2007;192:328–334. doi: 10.1016/j.atherosclerosis.2006.08.003

13. Jönsson S, Lundberg A, Kälvegren H, Bergström I, Szymanowski A, Jonasson L.

Increased Levels of Leukocyte-Derived MMP-9 in Patients with Stable Angina

Pectoris. *PLOS ONE*. 2011;6:e19340. doi: 10.1371/journal.pone.0019340

14. Särndahl E, Bergström I, Brodin VP, Nijm J, Lundqvist Setterud H, Jonasson L.

Neutrophil activation status in stable coronary artery disease. *PLoS One*.

2007;2:e1056. doi: 10.1371/journal.pone.0001056

15. Martin C, Burdon PC, Bridger G, Gutierrez-Ramos JC, Williams TJ, Rankin SM.

Chemokines acting via CXCR2 and CXCR4 control the release of neutrophils from

the bone marrow and their return following senescence. *Immunity*. 2003;19:583–593.

doi: 10.1016/s1074-7613(03)00263-2

16. Brostjan C, Oehler R. The role of neutrophil death in chronic inflammation and cancer.

Cell Death Discovery. 2020;6:26. doi: 10.1038/s41420-020-0255-6

17. Ocaña MG, Asensi V, Montes AH, Meana A, Celada A, Valle-Garay E. Autoregulation

mechanism of human neutrophil apoptosis during bacterial infection. *Mol Immunol*.

2008;45:2087–2096. doi: 10.1016/j.molimm.2007.10.013

18.   Midgley A, McLaren Z, Moots RJ, Edwards SW, Beresford MW. The role of neutrophil

404   apoptosis in juvenile-onset systemic lupus erythematosus. *Arthritis Rheum*.

405   2009;60:2390–2401. doi: 10.1002/art.24634

406   19.   Weinmann P, Moura RA, Caetano-Lopes JR, Pereira PA, Canhão H, Queiroz MV,

407   Fonseca JE. Delayed neutrophil apoptosis in very early rheumatoid arthritis patients

408   is abrogated by methotrexate therapy. *Clin Exp Rheumatol*. 2007;25:885–887.

409   20.   Brannigan AE, O'Connell PR, Hurley H, O'Neill A, Brady HR, Fitzpatrick JM, Watson

410   RW. Neutrophil apoptosis is delayed in patients with inflammatory bowel disease.

411   *Shock*. 2000;13:361–366. doi: 10.1097/00024382-200005000-00003

412   21.   Garlichs CD, Eskafi S, Cicha I, Schmeisser A, Walzog B, Raaz D, Stumpf C, Yilmaz

413   A, Bremer J, Ludwig J, et al. Delay of neutrophil apoptosis in acute coronary

414   syndromes. *J Leukoc Biol*. 2004;75:828–835. doi: 10.1189/jlb.0703358

415   22.   Biasucci LM, Liuzzo G, Giubilato S, Della Bona R, Leo M, Pinnelli M, Severino A,

416   Gabriele M, Brugaletta S, Piro M, et al. Delayed neutrophil apoptosis in patients with

417   unstable angina: relation to C-reactive protein and recurrence of instability. *Eur Heart*

418   *J*. 2009;30:2220–2225. doi: 10.1093/eurheartj/ehp248

419   23.   Mor A, Luboshits G, Planer D, Keren G, George J. Altered status of CD4(+)CD25(+)

420   regulatory T cells in patients with acute coronary syndromes. *Eur Heart J*.

421   2006;27:2530–2537. doi: 10.1093/eurheartj/ehl222

422   24.   George J, Schwartzenberg S, Medvedovsky D, Jonas M, Charach G, Afek A,

423   Shamiss A. Regulatory T cells and IL-10 levels are reduced in patients with

424   vulnerable coronary plaques. *Atherosclerosis*. 2012;222:519–523. doi:

425   10.1016/j.atherosclerosis.2012.03.016

426   25.   Hasib L, Lundberg AK, Zachrisson H, Ernerudh J, Jonasson L. Functional and

427   homeostatic defects of regulatory T cells in patients with coronary artery disease. *J*

428   *Intern Med*. 2016;279:63–77. doi: 10.1111/joim.12398

26.   Lewkowicz P, Lewkowicz N, Sasiak A, Tchórzewski H. Lipopolysaccharide-activated

CD4+CD25+ T regulatory cells inhibit neutrophil function and promote their apoptosis

and death. *J Immunol*. 2006;177:7155–7163. doi: 10.4049/jimmunol.177.10.7155

27.   Okeke EB, Mou Z, Onyilagha N, Jia P, Gounni AS, Uzonna JE. Deficiency of

Phosphatidylinositol 3-Kinase δ Signaling Leads to Diminished Numbers of

Regulatory T Cells and Increased Neutrophil Activity Resulting in Mortality Due to

Endotoxic Shock. *J Immunol*. 2017;199:1086–1095. doi: 10.4049/jimmunol.1600954

28.   Okeke EB, Uzonna JE. The Pivotal Role of Regulatory T Cells in the Regulation of

Innate Immune Cells. *Front Immunol*. 2019;10:680. doi: 10.3389/fimmu.2019.00680

29.   Ou Q, Power R, Griffin MD. Revisiting regulatory T cells as modulators of innate

immune response and inflammatory diseases. *Front Immunol*. 2023;14:1287465. doi:

10.3389/fimmu.2023.1287465

30.   Ledderose C, Hashiguchi N, Valsami EA, Rusu C, Junger WG. Optimized flow

cytometry assays to monitor neutrophil activation in human and mouse whole blood

samples. *J Immunol Methods*. 2023;512:113403. doi: 10.1016/j.jim.2022.113403

31.   Cox G. IL-10 enhances resolution of pulmonary inflammation in vivo by promoting

apoptosis of neutrophils. *Am J Physiol*. 1996;271:L566–571. doi:

10.1152/ajplung.1996.271.4.L566

32.   Laichalk LL, Danforth JM, Standiford TJ. Interleukin-10 inhibits neutrophil phagocytic

and bactericidal activity. *FEMS Immunol Med Microbiol*. 1996;15:181–187. doi:

10.1111/j.1574-695X.1996.tb00084.x

33.   Mazzone A, De Servi S, Ricevuti G, Mazzucchelli I, Fossati G, Pasotti D, Bramucci E,

Angoli L, Marsico F, Specchia G, et al. Increased expression of neutrophil and

monocyte adhesion molecules in unstable coronary artery disease. *Circulation*.

1993;88:358–363. doi: 10.1161/01.cir.88.2.358

34.   Uhl B, Vadlau Y, Zuchtriegel G, Nekolla K, Sharaf K, Gaertner F, Massberg S,

Krombach F, Reichel CA. Aged neutrophils contribute to the first line of defense in the

acute inflammatory response. *Blood*. 2016;128:2327–2337. doi: 10.1182/blood-2016-

05-718999

35. Chen J, Bai Y, Xue K, Li Z, Zhu Z, Li Q, Yu C, Li B, Shen S, Qiao P, et al. CREB1-

driven CXCR4hi neutrophils promote skin inflammation in mouse models and human

patients. *Nature Communications*. 2023;14:5894. doi: 10.1038/s41467-023-41484-3

36. Maher BM, Dhonnchu TN, Burke JP, Soo A, Wood AE, Watson RW. Statins alter

neutrophil migration by modulating cellular Rho activity--a potential mechanism for

statins-mediated pleotropic effects? *J Leukoc Biol*. 2009;85:186–193. doi:

10.1189/jlb.0608382

37. Antoniellis Silveira AA, Dominical VM, Morelli Vital D, Alves Ferreira W, Trindade

Maranhão Costa F, Werneck CC, Ferreira Costa F, Conran N. Attenuation of TNF-

induced neutrophil adhesion by simvastatin is associated with the inhibition of Rho-

GTPase activity, p50 activity and morphological changes. *International*

Immunopharmacology. 2018;58:160–165. doi:

https://doi.org/10.1016/j.intimp.2018.03.025

38. Chello M, Anselmi A, Spadaccio C, Patti G, Goffredo C, Di Sciascio G, Covino E.

Simvastatin increases neutrophil apoptosis and reduces inflammatory reaction after

coronary surgery. *Ann Thorac Surg*. 2007;83:1374–1380. doi:

10.1016/j.athoracsur.2006.10.065

39. Mandal P, Chalmers JD, Graham C, Harley C, Sidhu MK, Doherty C, Govan JW,

Sethi T, Davidson DJ, Rossi AG, et al. Atorvastatin as a stable treatment in

bronchiectasis: a randomised controlled trial. *Lancet Respir Med*. 2014;2:455–463.

doi: 10.1016/s2213-2600(14)70050-5

40. Li W, Deng C, Yang H, Wang G. The Regulatory T Cell in Active Systemic Lupus

Erythematosus Patients: A Systemic Review and Meta-Analysis. *Front Immunol*.

2019;10:159. doi: 10.3389/fimmu.2019.00159

41.   Jiang Q, Yang G, Liu Q, Wang S, Cui D. Function and Role of Regulatory T Cells in

483   Rheumatoid Arthritis. *Front Immunol*. 2021;12:626193. doi:

484   10.3389/fimmu.2021.626193

485   42.   Jalalvand M, Enayati S, Akhtari M, Madreseh E, Jamshidi A, Farhadi E, Mahmoudi M,

486   Amirzargar A. Blood regulatory T cells in inflammatory bowel disease, a systematic

487   review, and meta-analysis. *Int Immunopharmacol*. 2023;117:109824. doi:

488   10.1016/j.intimp.2023.109824

489   43.   Noseykina EM, Schepetkin IA, Atochin DN. Molecular Mechanisms for Regulation of

490   Neutrophil Apoptosis under Normal and Pathological Conditions. *J Evol Biochem*

491   *Physiol*. 2021;57:429–450. doi: 10.1134/s0022093021030017

492   44.   Dominguez-Villar M, Hafler DA. Regulatory T cells in autoimmune disease. *Nature*

493   *Immunology*. 2018;19:665–673. doi: 10.1038/s41590-018-0120-4

494   45.   Chaudhry A, Samstein RM, Treuting P, Liang Y, Pils MC, Heinrich JM, Jack RS,

495   Wunderlich FT, Brüning JC, Müller W, et al. Interleukin-10 signaling in regulatory T

496   cells is required for suppression of Th17 cell-mediated inflammation. *Immunity*.

497   2011;34:566–578. doi: 10.1016/j.immuni.2011.03.018

498   46.   Laidlaw BJ, Cui W, Amezquita RA, Gray SM, Guan T, Lu Y, Kobayashi Y, Flavell RA,

499   Kleinstein SH, Craft J, et al. Production of IL-10 by CD4+ regulatory T cells during the

500   resolution of infection promotes the maturation of memory CD8+ T cells. *Nature*

501   *Immunology*. 2015;16:871–879. doi: 10.1038/ni.3224

502   47.   Keel M, Ungethüm U, Steckholzer U, Niederer E, Hartung T, Trentz O, Ertel W.

503   Interleukin-10 counterregulates proinflammatory cytokine-induced inhibition of

504   neutrophil apoptosis during severe sepsis. *Blood*. 1997;90:3356–3363.

505   48.   Ward C, Murray J, Clugston A, Dransfield I, Haslett C, Rossi AG. Interleukin-10

506   inhibits lipopolysaccharide-induced survival and extracellular signal-regulated kinase

507   activation in human neutrophils. *Eur J Immunol*. 2005;35:2728–2737. doi:

508   10.1002/eji.200425561

49.   Lewkowicz N, Mycko MP, Przygodzka P, Ćwiklińska H, Cichalewska M, Matysiak M,

Selmaj K, Lewkowicz P. Induction of human IL-10-producing neutrophils by LPS-

stimulated Treg cells and IL-10. *Mucosal Immunol*. 2016;9:364–378. doi:

10.1038/mi.2015.66

50.   Thornton AM, Shevach EM. CD4+CD25+ immunoregulatory T cells suppress

polyclonal T cell activation in vitro by inhibiting interleukin 2 production. *J Exp Med*.

1998;188:287–296. doi: 10.1084/jem.188.2.287

51.   Lewkowicz N, Klink M, Mycko MP, Lewkowicz P. Neutrophil--CD4+CD25+ T

regulatory cell interactions: a possible new mechanism of infectious tolerance.

Immunobiology. 2013;218:455–464. doi: 10.1016/j.imbio.2012.05.029

Figure 1: Neutrophil apoptosis, cell surface marker expression and neutrophil activation in patients with chronic coronary syndrome and healthy controls.

Isolated neutrophils from 20 patients with chronic coronary syndrome (CCS) and 19 healthy controls (HC) were cultured in medium only or medium with interleukin (IL)-10 (20 ng/ml), tumor necrosis factor (TNF, 1 ng/ml) or lipopolysaccharide (LPS, 100 ng/ml). After 5 h, flow cytometry was used to assess neutrophil apoptosis; data are presented as the percentage of apoptotic (Annexin$^+$/SYTOX$^-$) neutrophils (**A**). Flow cytometry was also used to assess the expression of CD66b; data are presented as mean fluorescence intensity (MFI) values (**B**).

Furthermore, the neutrophil expression of CD66b and CXCR4 was assessed by flow cytometry immediately after cell isolation in the same study cohort. Data are presented as MFI values of CD66b (**C**) and MFI values of CXCR4 (**D**).

Additionally, protein concentrations were measured in the same study cohort after 2.5 h of incubation in medium with or without IL-10 (20 ng/ml), TNF (1 ng/ml) or LPS (100 ng/ml) (**E**). Concentrations of matrix metalloprotease (MMP)-9, MMP-8 and myeloperoxidase (MPO), TNF, interferon (IFN)γ, IL-6, IL-18 and IL-1β are presented as pg/ml. Significances between CCS and HC in normal medium and IL-10 were all $p > 0.05$, except for MMP-9 measurement in IL-10 treated sample.

Significances are indicated between CCS and HC and all values are given as median ± 95% confidence interval.

Figure 2: The effect of neutrophil-T_{reg} co-cultures on neutrophil apoptosis in patients with chronic coronary syndrome and healthy controls.

Neutrophils and T_{regs} were isolated from 20 patients with chronic coronary syndrome (CCS) and 19 healthy controls (HC). The cells were co-cultured at different ratios (neutrophils: T_{regs} 1:0, 10:1, 4:1 and 1:1) in medium only or medium with interleukin (IL)-10 (20 ng/ml), tumor necrosis factor (TNF, 1 ng/ml) or lipopolysaccharide (LPS, 100 ng/ml). After 5 h of co-culture, neutrophil apoptosis (Annexin$^+$/SYTOX$^-$) was determined by flow cytometry. The percentages of apoptotic neutrophils are presented as fold change compared to pure neutrophil culture in patients with CCS (**A**) and HC (**B**). Significances are compared to 1:0 culture within CCS and HC groups, respectively, *p*-values are adjusted for multiple comparison via Dunn; CCS (**A**): *p=0.015 in medium only ratio 4:1, *p=0.036 in TNF ratio 1:1; HC (**B**): *p=0.014 in medium only ratio 4:1, *p=0.030 in medium only ratio 1:1, *p=0.025 in IL-10 ratio 4:1, **p=0.002 in TNF ratio 1:1, **p=0.009 in LPS ratio 4:1, **p=0.006 in LPS ratio 1:1; and significance for trend is given as *p*-value of a Greenhouse-Geisser test. All values are given as median ± 95% confidence interval. Representative examples for the gating of apoptotic neutrophils as Annexin V$^+$/SYTOX$^-$ cells are shown in **C**, the full gating strategy can be reviewed in Figure S1.

Additionally, isolated neutrophils from healthy volunteers (n=5, < 50 years of age) were treated with different doses of IL-10 (1 pg/ml, 1ng/ml and 100ng/ml). Flow cytometry was used to assess neutrophil apoptosis after 5 h; data are presented as the percentage of apoptotic (Annexin$^+$/SYTOX$^-$) neutrophils (**D**). In another series of experiments, using cells from the same volunteers described in (**D**), neutrophils alone or neutrophil-T_{reg} co-cultures at ratio 4:1 were seeded +/- mesh (as graphically indicated) in medium only for 5 h, followed by flow cytometry to assess neutrophil apoptosis (**E**). *p*-values were adjusted for multiple comparison via Tukey. All values are given as mean ± SEM.