**Quantifying the oxygen preferences of bacterial communities using a metagenome-based**

**approach**

Clifton P. Bueno de Mesquita[1,2#], Elías Stallard-Olivera[1,2], Noah Fierer[1,2#]

[1] Cooperative Institute for Research in Environmental Sciences, University of Colorado,

Boulder, CO 80309-0216, USA

[2] Department of Ecology and Evolutionary Biology, University of Colorado, Boulder, CO

80309-0334, USA

[#] Corresponding authors: cliff.buenodemesquita@colorado.edu, noah.fierer@colorado.edu

Keywords: oxygen, oxygen tolerance, aerobes, anaerobes, bioindicators

Running title: Quantifying bacterial oxygen preferences

**Abstract**

Oxygen is a primary driver of the distribution and activity of microbial life. Since oxygen

levels are often difficult to measure *in situ*, one potential solution is to use bacteria as

bioindicators of oxygen levels. As bacteria range from obligate aerobes to obligate anaerobes,

quantification of bacterial community oxygen preferences could be used to infer variation in

environmental oxygen levels and bacterial metabolic strategies. After using ensemble machine

learning to select the 20 most important genes that predict oxygen tolerances in individual

bacteria, we established a relationship between the abundance ratio of aerobic: anaerobic

indicator genes and the proportional abundance of aerobic bacteria using simulated metagenomes

with varying ratios of known aerobic and anaerobic bacteria. We developed a tool, OxyMetaG,

that takes metagenomic reads as input, extracts bacterial reads, maps reads to the 20 genes, and

predicts the proportion of aerobic versus anaerobic bacteria in any given sample. We tested

OxyMetaG on a suite of metagenomes with measured or inferred oxygen levels across a variety

of environmental and host-associated samples. To demonstrate the utility of our approach, we

applied OxyMetaG to 540 surface soils, showing that surface soils are typically dominated by

aerobes, but wetter sites with finer textures have relatively more anaerobes. Lastly, we applied

OxyMetaG to 73 human gut samples, showing that in the first three years of life, human guts

progress from having up to 61% aerobes to being completely dominated by anaerobes. We

expect OxyMetaG to have broad utility for characterizing both modern and ancient

environments.

**Importance**

Oxygen is one of the most important environmental variables affecting microbial activity and

composition but is often difficult to measure *in situ*. We developed a tool, OxyMetaG, that

leverages differences in bacterial gene content across known aerobic and anaerobic taxa to

predict the proportion of aerobes and anaerobes in a given sample directly from shotgun

metagenomic reads. OxyMetaG works on samples with low sequencing depth and avoids

computationally expensive genome assembly, which often captures only a fraction of the

microbial community in a given environment. With OxyMetaG, bacteria can be used as

bioindicators of oxygen availability over broader time scales than just a single measurement and

provide crucial environmental context in cases where oxygen has not or cannot be measured.

OxyMetaG is publicly available and can be used to answer a wide variety of ecological questions

in both environmental and host-associated systems.

**Introduction**

Throughout geologic history, atmospheric oxygen levels have been variable. For the first ~1.5 billion years that life likely existed on Earth, atmospheric oxygen was limited and anaerobic bacteria predominated (1). Only following the Great Oxidation Event (2) did the atmosphere become oxygenated, with levels over the past ~2.4 billion years ranging from 15% to 35% before reaching the current concentration of 21% (3). Even across modern environments, oxygen concentrations can be highly variable (e.g., waterlogged hypoxic wetland sediments to oxic surface ocean water). This spatial and temporal variation in oxygen concentrations is often a primary determinant of bacterial distributions and their metabolic activities. Bacteria can differ widely with respect to their oxygen tolerances and preferences with the oxygen levels in any given environment dictating what taxa can thrive in that environment and their potential metabolic strategies. We sought to take advantage of these patterns to develop a predictive tool that can work directly on metagenomic reads (avoiding computationally expensive assembly, binning, and annotation) to predict the percent relative abundances of aerobic and anaerobic bacteria, which in turn can be used to help infer the relative amount of oxygen in any given environmental or host-associated sample.

Bacterial oxygen preferences range from taxa that are obligate anaerobes that are unable to grow in the presence of oxygen and instead generate energy via fermentation or anaerobic respiration, to obligate aerobes that require atmospheric levels of oxygen to sustain growth. Between these end members exist facultative aerobes/anaerobes that can tolerate oxygen or a lack of oxygen and switch their metabolisms accordingly, and microaerophiles, which are defined as organisms that grow at low (below atmospheric, usually 2-10%) oxygen levels (4). In short, bacterial oxygen preferences span a spectrum and, for most cultivated bacteria, we know

from experimental assays where they fall along this spectrum. Even for the majority of bacteria

that are resistant to cultivation and have not been studied *in vitro*, oxygen preferences can often

be inferred from genome-based models, including models that rely on specific sets of genes (5–

14) or other genomic attributes such as amino acid frequencies (15, 16). While such genome-

based models enable inferences of oxygen preferences for individual taxa, inferring community-

level oxygen preferences would require having high-quality genomes for the majority of taxa in a

given community, a challenging task in many systems with high levels of bacterial diversity (17).

We thus sought to develop a metagenome-based tool that can work directly on metagenomic

reads to predict community-level oxygen preferences, namely the percent of aerobic versus

anaerobic bacteria in any given sample. Not only would such a metagenome-based model make

it possible to quantify oxygen preferences as a community aggregated trait (18), we could also

use inferred community-level oxygen preferences as a means to quantify oxygen availability in

individual samples, assuming that the oxygen preferences of bacterial communities should, over

time, reflect oxygen availability. In this way, microbes would be used as bioindicators of oxygen

availability, as has been done previously for other important but hard-to-measure variables such

as phosphorus availability (19), heavy metal contamination, and fecal contamination (20).

Despite its importance, oxygen is notoriously hard to measure in environmental and host-

associated samples due to a combination of methodological constraints and the fact that oxygen

concentrations are often highly variable across space and time. Take, for example, soils, where

oxygen concentrations can vary at the millimeter scale, for example between oxic pore spaces

and neighboring anoxic microsites in aggregates (21). Soil oxygen concentrations are also

constantly in flux as oxygen is consumed by microbial respiration and supplied by diffusion from

the atmosphere. Yet understanding soil oxygen is important for understanding microbial ecology,

soil health, and soil biogeochemical process rates (22, 23). As another example, measuring

oxygen concentrations in vertebrate guts is notoriously challenging as sensors must be placed

inside the individual (24). Yet understanding both human and non-human animal gut oxygen

concentrations is crucial for understanding digestion, metabolism, and disease (25–27). Thus, it

would be beneficial to infer relative oxygen levels in soils, vertebrate guts, and other

environmental or host-associated samples directly from the ever-expanding metagenomic

datasets now available.

Here, we describe the development and validation of the new tool OxyMetaG (Figure 1).

Our goal was to identify a narrow set of genes with high predictive power at the genome level

and then use a read-mapping and abundance-ratio methodology to rapidly profile the raw reads

of metagenomes, thereby avoiding the limitations of having to assemble high-quality genomes

from metagenomes. After comprehensive validation using simulated metagenomes, we tested

OxyMetaG on a wide variety of environmental and host-associated sample types that capture

broad gradients in oxygen availability, including a subset of samples with measured oxygen

concentrations. We then present two example case studies of how OxyMetaG can be used to

answer ecological questions in one environmental system (surface soils) and one host-associated

system (human guts). For surface soils, demonstrate how OxyMetaG can be used to quantify

continental-scale variation in the predicted relative abundances of aerobic bacteria and the

climate or edaphic variables associated with this variation. For human guts, we used

metagenomic data from selected individuals sampled over time to quantify how the predicted

relative abundance of aerobic bacteria changes during the first three years of life. Together these

examples demonstrate how OxyMetaG can be used to identify community-level shifts in

bacterial oxygen preferences directly from metagenomic data and infer differences in oxygen

availability within and across environments.

**Results**

We identified a set of 20 genes that were highly predictive of oxygen tolerance, including

13 genes associated with aerobes and 7 genes associated with anaerobes (Table 1, Figure S1).

The classification model of oxygen tolerance using just those 20 genes had high precision, recall,

and F1-score (Table 2), and this was true when tested on all taxa and, the four main phyla

(Pseudomonadota, Actinomycetota, Bacillota, Bacteroidota) combined, on each of the main four

phyla individually, and on all other rare phyla combined (Table S1). Accuracy per phylum was >

85% for all 31 phyla, and > 93% for 29 of 31 phyla (Table S2).

In simulated metagenomes with known amounts of aerobic and anaerobic bacterial

genomes, the ratio of the summed RPK abundance of the 13 aerobic indicator genes to the 7

anaerobic indicator genes was well correlated with the percent of aerobic bacterial taxa in

simulated metagenomes, regardless of sequencing depth (Figure 2). The relationship was non-

linear and a generalized additive model was fit to the data ($F = 1605$, $R^2 = 0.99$, $p < 0.001$) that

was then used to make predictions on real metagenomes with unknown proportions of aerobic

and anaerobic bacteria.

The predicted percentages of aerobic bacteria closely matched the expected differences in

oxygen availability across different habitat types. Samples from cattle rumen, human gut, and

marine sediments had the greatest proportion of anaerobes, while surface ocean, grassland soil,

and forest soil had significantly more aerobes (Tukey posthoc, $p < 0.05$, Figure 3). Irrigated

temperate cropland soils and tropical forest soils had the greatest relative abundance of

anaerobes in any of the upland soil habitats, potentially reflecting greater soil moisture content

and consequently lower oxygen concentrations in those soils. Furthermore, in the three datasets

where metagenomic data were available with corresponding measurements of oxygen

concentrations (Baltic Sea sediment, Black Sea water, Lake Tanganyika water), the predicted

relative abundances of aerobic bacteria increased with measured oxygen concentrations (Figure

4). Results were similar when comparing the default cutoffs (percent identity $\geq 60$, e-value $<$

0.001, and bitscore $\geq 50$) and more stringent cutoffs (percent identity $\geq 60$, e-value $< 1e\text{-}10$,

bitscore $\geq 60$, and amino acid alignment length $\geq 40$) (Figure S2).

Next, we tested OxyMetaG across surface soils that are generally considered to be oxic

but can have variable oxygen concentrations depending on moisture and texture, with some

surface soils harboring anaerobes in microsites (21). Across 331 Australian soils, ranging from

arid sites in western Australia to humid sites in Tasmania and northeastern Australia, the

predicted relative abundance of aerobic bacteria ranged from 60% to 100% (mean = 95%, SD =

9%, median = 100%). The majority of these soils were dominated by aerobic bacteria, with 204

samples predicted to have 100% aerobes and 127 samples predicted to have some anaerobes.

Across 209 soils collected from 26 sites across the U.S., ranging from tropical to subarctic

climates, the predicted relative abundance of aerobic bacteria ranged from 40% to 100% (mean =

95%, SD = 10%, median = 100%). Similar to Australian soils, most U.S. soils were dominated

by aerobic bacteria, with 145 samples predicted to have 100% aerobes and 64 samples predicted

to have some anaerobes. Ecosystem type significantly affected the proportion of aerobic bacteria

in Australia (ZIBR, $p < 0.001$) but not in the U.S. (ZIBR, $p = 0.22$) (Figure 5). Climate class did

not significantly affect the proportion of aerobes in Australia (ZIBR, $p = 0.13$) or the U.S.

(ZIBR, $p = 0.96$) (Figure S3). As for continuous soil and climate variables, gravimetric water

content was the only variable included in our models that was significantly associated with the

predicted proportion of aerobes in both datasets (Table S3, Figure S4). Sand content, organic

carbon content, and mean annual temperature were also significantly associated with the

predicted proportion of aerobes in the Australian dataset (Table S3).

Lastly, we tested OxyMetaG with metagenomes from the human gut, which is generally

considered to be anoxic, but can be oxic in the first few months of life (28). Across the 73

metagenomes tested, the predicted relative abundance of aerobic bacteria ranged from 0 to 61%.

The proportion of aerobic bacteria declined significantly with time after birth (LMER, $\chi^2 = 18.9$,

$p < 0.001$), but there was considerable variation across individuals (Figure 6), with the model

including random slopes and intercepts for subject IDs significantly outperforming the model

with only random intercepts (ANOVA, $\chi^2 = 7.3$, $p = 0.03$). All individuals eventually had 0%

aerobic bacteria by the later time points, but the timepoint at which this occurred varied, with one

individual still having aerobes past day 600 (Figure 6).

**Discussion**

We developed a metagenome-based tool, OxyMetaG, that uses a gene abundance ratio to

infer the percent relative abundance of aerobic bacteria in a wide variety of environmental

samples. OxyMetaG outputs the predicted percent relative abundance of aerobic bacteria (out of

total bacteria) in a sample, which can be used as a bioindicator of general oxygen availability.

We identified 20 genes whose presence or absence could accurately predict whether a given

bacterial taxon was aerobic or anaerobic across the majority of bacterial phyla (Figure S1, Table

S1). These 20 genes are typically involved in various pathways of aerobic or anaerobic metabolic

processes or in free radical scavenging, with a subset of these genes (9 of 20) having also been

identified as predictors of bacterial oxygen tolerance in previous studies (6, 12–14). The gene

abundance ratio was strongly related to the relative abundance of known aerobic bacteria in

simulated metagenomes (Figure 2), demonstrating the predictive power of this metric. We first

tested OxyMetaG on a diverse set of 203 terrestrial, aquatic, marine, and host-associated

metagenomes. We then used OxyMetaG on 540 surface soils across two continents to investigate

how the percent of aerobic bacteria varies across surface soil bacterial communities, and on 73

infant to toddler-age human gut samples to investigate patterns in the percent aerobic bacteria in

the gut over time.

OxyMetaG accurately captured expected differences in community-level bacterial

oxygen tolerances, predicting a dominance of anaerobes in known hypoxic to anoxic

environments such as cattle rumens, human guts, anaerobic digesters, and wetland sediments,

with aerobes found to dominate in environments that we typically consider to be oxic, such as

surface soils and surface marine and freshwater samples (Figure 3). These results were similar to

those obtained using other tools that rely on classifying individual genomes from those sample

types (15, 16). One advantage of OxyMetaG compared to genome-based approaches is that it

works on raw reads and does not require genome assembly, which can be a major limitation in

many samples where obtaining metagenome-assembled genomes (MAGs) can be difficult, or

when the MAGs that are obtained only represent a fraction of the diversity (17). Furthermore, in

metagenomes from the Black Sea, the Baltic Sea, and Lake Tanganyika, the inferred changes in

the relative abundances of aerobes within environments were well correlated with known oxygen

gradients (29) (Figure 4). Together, these validation tests demonstrated how the OxyMetaG

output (predicted percent aerobes) can be used for broad comparisons among sample types as

well as more specific comparisons within sample types collected across gradients.

As a test case for demonstrating the utility of OxyMetaG for quantifying community-

level patterns in bacterial oxygen tolerances across environmental gradients, we analyzed 540

metagenomes representing a broad range of soil types, habitat types, and climate conditions

across Australia and the US. Most samples across both datasets were overwhelmingly dominated

by aerobes, with OxyMetaG suggesting 100% relative abundance of aerobes in a majority (74%

of samples). While these samples may in fact harbor a small proportion of anaerobic taxa, the

high ratio ($> 30$) of aerobic indicator genes to anaerobic indicator genes in these samples

suggests that any anaerobes may be relatively rare and temporally transient. These results

suggest that, despite the presence of anaerobic microsites, surface soils generally select for, and

are dominated by, aerobic bacteria. In line with predictions (21), soil moisture was the top

variable associated with the percent aerobic bacteria in both datasets (Table S3), with wetter soils

having slower diffusion of atmospheric oxygen (Figure S4). Still, the variation explained by soil

moisture was low (Table S3), likely due to moisture being a single time-of-sampling

measurement and the complexity of variables that influence soil oxygen availability.

As a second test case for OxyMetaG, we analyzed 73 human gut metagenomes, focusing

on individuals that were sampled repeatedly over their first 3 years of life. Prior work using 16S

rRNA marker gene sequencing combined with trait databases (matched by genus and species

name or with hidden state prediction algorithms) showed that the abundance-weighted mean

oxygen tolerance decreased markedly from months 3-15 (28). Results from our metagenome-

based approach generally supports the idea of elevated oxygen availability in the gut during

infancy (30), with a trend towards anaerobic dominance by year three that is likely driven by

increases in *Bifidobacterium* and members of Bacillota such as Clostridiaceae (31, 32).

Importantly, our data show a greater degree of variation than the prior work, likely due to the

fact that OxyMetaG considers all bacterial reads and the gene-abundance-based method does not

rely on trait databases biased towards well-characterized taxa.

One limitation of OxyMetaG is that we do not know the temporal scale that is integrated

by our metagenome-based metrics. Standard metagenomic sequencing approaches do not

discriminate between live and dead cells (33), nor do they capture variation in taxon-specific

growth rates (34). Thus, OxyMetaG inferences of oxygen availability likely integrate over longer

time periods that reflect both current conditions at the time of sampling and past conditions. This

is beneficial in that OxyMetaG predictions are more representative of the sample history and

general sample conditions, not just the single sampling moment, and the predictions are likely

more robust to rapid variations in oxygen levels. The downside is that OxyMetaG would not be

useful for studying changes in oxygen availability over short time scales. Although we currently

do not know the exact time period that is integrated, as this will depend on community turnover

rates, future work could investigate the temporal changes in aerobe versus anaerobe ratios in

communities sampled over time where oxygen levels are experimentally manipulated. Another

limitation of OxyMetaG is its omission of facultative aerobes/anaerobes and microaerophiles

during training and gene identification; a useful next step would also be to incorporate genomic

features from these taxa into OxyMetaG to more fully capture the spectrum of bacterial oxygen

preferences.

We have presented results from OxyMetaG for a total of 816 modern metagenomic

samples and expect OxyMetaG to be useful for inferring oxygen levels and answering ecological

questions in more studies on modern metagenomes. Additionally, we also suggest that

OxyMetaG could be useful in ancient metagenomic studies, where past oxygen levels are

unknown and can only be inferred using geochemical proxies. OxyMetaG could be used in

concert with other geochemical paleo-oxybarometers (35) for more robust inference about

sample history. In a first pilot study on ancient metagenomes, OxyMetaG predictions were in

line with isotopic data and helped identify two clusters of communities (aerobic and anaerobic)

in subglacial lake samples from Antarctica (De Sanctis et al. 2025). OxyMetaG could contribute

to the toolkit of paleo-oxygen proxies in that it is universal (as long as the sample has bacterial

reads) and does not need to be calibrated on each individual sample or sample type. However,

OxyMetaG still requires further tuning and optimization to handle issues specific to ancient

DNA analyses, such as DNA damage and shorter fragment lengths; such improvements will be

released in future updates to OxyMetaG. We hope OxyMetaG is ultimately useful and

contributes to a better understanding of both microbial ecology and environmental conditions in

general.

**Materials and Methods**

Our workflow consisted of first modeling oxygen tolerance at the genome level to

identify key genes that are highly predictive of oxygen tolerance ("indicator genes"), then using

simulated metagenomes with known percentages of aerobic bacteria to establish a relationship

between the percentage of aerobic bacteria and the ratio of aerobic indicator genes to anaerobic

indicator genes, and then testing with actual metagenomes both for validation and application

(Figure 1).

We first downloaded 5520 bacterial genomes from 31 different phyla that had oxygen

tolerance information available in BacDive in June 2025 (36) (Table S4, Figure S5). Since our

goal was to identify the most important genes associated with clearly aerobic and anaerobic

bacteria, we only used taxa classified as "obligate aerobe", "aerobe", "anaerobe" or "obligate

anaerobe", and omitted taxa classified as "facultative aerobe", "facultative anaerobe", or

"microaerophile". We acquired additional information about the genomes, including taxonomic

information, by matching accessions to GTDB r226 (37). We used hmmer v3.4 (38) to annotate

the genomes with the Pfam v37.3 database (39). We then used an ensemble machine learning

approach to select the top 20 genes most predictive of oxygen tolerance. This approach uses

CatBoost, XGboost, and RandomForests, an 80:20 ratio of training to testing data, and SHAP

scores to assess variable importance (40). After building the best model using all genes, the

model was rerun with just the top 20 genes with the highest SHAP scores. The model with only

20 genes performed as well as the model with all genes in terms of precision, recall, and F1-

score (weighted averages all 0.96 for both models), identifying 13 genes associated with aerobes

and 7 with anaerobes (Table 1, Figure S1). The prevalence of the 13 aerobic indicator genes

ranged from 54% to 99% in aerobic taxa and from 2% to 47% in anaerobic taxa, while the

prevalence of the 7 anaerobic indicator genes ranged from 51% to 89% in anaerobic taxa and 1%

to 17% in aerobic taxa (Table 1). Furthermore, the accuracy of the model was high even when

tested on individual phyla (F1 > 0.90). Tests were conducted on each of the four phyla with the

most genomes (Pseudomonadota, Actinomycetota, Bacillota, Bacteroidota), as well as on all of

the remaining 27 phyla combined. Furthermore, accuracy of the main model (trained using all

phyla, tested using all phyla) was assessed for each phylum to quantify model performance per

phylum. The scripts to implement our approach are publicly available on Zenodo

(https://doi.org/10.5281/zenodo.18331946).

We then used ART (41) to simulate metagenomes (Illumina, 150 bp) using 500 randomly

selected genomes from the original set of 5520 genomes. To test for any effect of sequencing

depth, simulations were done using fold coverages of 0.5 and 2.5, which generated 5-8 million

reads and 26-41 million reads, respectively, depending on the randomly selected subset of

genomes used in the simulation. We used a gradient of 0, 20, 40, 60, 80, and 100% of the

genomes as aerobes. We used three replicates per aerobe percentage, with the specified number

of aerobic and anaerobic genomes randomly selected from the available pool of 5520 genomes.

Reads were trimmed and quality filtered with Trimmomatic v0.39 to remove Illumina adapters

and retain reads with $Q \geq 20$ and length $\geq 100$ bp (42). We then downloaded all of the reviewed

protein sequences for each Pfam as a fasta file from the Pfam website, concatenated them into a

single fasta file, and generated a DIAMOND database with DIAMOND v2.1.13 (43). We then

used DIAMOND BLASTX to calculate the abundance of the 20 genes in the metagenomes. We

retained hits with percent identity $\geq 60$, e-value $< 0.001$, and bitscore $\geq 50$ (44). We then

normalized by mean Pfam length to calculate reads per kilobase (RPK) for each Pfam. We

summed the RPK of the 13 aerobic genes and the 7 anaerobic genes and then calculated the

aerobic: anaerobic gene RPK ratio. Because the relationship was nonlinear, we fit a generalized

additive model (GAM) of the relationship between the percent aerobe genomes and the ratio

(Figure 2) using the mgcv R package (45). The model relationship was highly similar across the

two sequencing depths tested (Figure 2).

We then validated the model on 203 modern metagenomes that we expected to capture

broad differences in oxygen concentrations, including a subset of samples with paired

measurements of oxygen concentrations. The 203 metagenomes spanned 16 distinct

environments, including cattle rumen (46), human gut (47), surface ocean water (top 5 m) (48),

Black Sea water (surface and deep) (49), Baltic sea sediment (0-2 cm below sea floor) (50),

anaerobic sludge digester (publicly available but unpublished), wetland sediment (51),

freshwater lake water (surface and $> 150$ m deep) (29), surface soil (0-10 cm) from arid

grasslands in Australia (52), surface soil (0-10 cm) from tropical forests in Panama (53), other

soils ranging from 0 to 70 cm depth from temperate forests, grasslands, and meadows (54), and

surface bulk and rhizosphere soil from maize crops (55). Approximately 15 samples were

selected from each study (Table S5). For further validation, we conducted additional analyses on

a subset of samples with measured gradients in oxygen concentrations. We compared 15

metagenomes from the Black Sea that spanned a depth profile from 50 m to 2000 m, with

measured oxygen and hydrogen sulfide concentrations (49), 15 metagenomes from Baltic Sea

sediments that spanned dissolved oxygen levels from 0.13 to 12.45 mg $L^{-1}$ (50), and 15

freshwater lake metagenomes ranging from 0 m to 1200 m depth with oxygen saturation levels

ranging from 1.3 to 91.4% (29). We downloaded raw metagenomic data from the NCBI SRA

and trimmed and filtered the reads with Trimmomatic as above. Because not all of the 20

predictor Pfams are exclusive to bacteria, we extracted only bacterial reads from the

metagenomes, as hits to archaeal or eukaryotic reads could introduce errors in our model. We

used Kraken2 v2.1.3 (56) to classify all of the quality-filtered metagenomic reads and then

KrakenTools (57) to extract only the bacterial reads from each metagenome. The number of

bacterial reads per metagenome averaged 10.9 million, with only four samples below 256000

reads. Then, as above, we used DIAMOND BLASTX to calculate the abundances of each Pfam

and then calculated the RPK ratio of aerobic indicator genes to anaerobic indicator genes to

predict the relative abundance of aerobic bacteria in the sample. Predictions over 100% were set

to 100%, predictions below 0% were set to 0%, and any ratios above 30 (the maximum value in

simulations) were set to 100%.

OxyMetaG is implemented as a Python package with three main functions, one for each

of the steps outlined above. The 'extract' function is a wrapper for Kraken2 and KrakenTools,

which together classify each read and then extract reads classified as bacterial. The 'profile'

function takes the bacterial reads as input, runs DIAMOND BLASTX on the database of the 20

Pfams, and records the information about hits for each sample. Lastly, the 'predict' function

takes the DIAMOND output files, filters them to the top significant hit per read, calculates the

summed aerobic: anaerobic RPK ratio, and predicts the proportion of aerobic bacteria using the

GAM established with the simulated metagenomes. While we used cutoffs of percent identity $\geq$

60, e-value < 0.001, and bitscore $\geq$ 50, users can also specify their own percent identity, e-value,

and bitscore cutoffs based on their specific data characteristics, such as read length, and desired

balance in sensitivity and specificity (58). This mode also enables users to test for consistency of

results across multiple tested cutoff values.

To test the effect of sequencing depth on the predictions, we chose one oxic sample, one

anoxic sample, and one mixed sample that each had high sequencing depth (>32 million reads

per sample). We randomly subsampled each of the three metagenomes to 16 different sequencing

depths (ranging from 0.001 to 32.768 million reads) running OxyMetaG separately at each

sequencing depth. The predicted relative abundance of aerobic bacteria stabilized at only 256000

reads, demonstrating the utility of this method for analyzing metagenomes with relatively low

sequencing depths that would preclude assembling genomes (Figure S6). This was also the point

at which there were anaerobic genes detected in the oxic sample, enabling the ratio to be

calculated and the prediction to be made (Figure S6).

After validation, we applied OxyMetaG on soil and human gut datasets to answer

ecological questions in those systems. First, we used 331 surface soil (0-10 cm) metagenomes

collected using standardized methodology from across Australia (52). These metagenomes have

been previously used in other research (59) and were selected because they are from natural

areas, had at least 10 million reads per metagenome, and had defined upland vegetation types.

Quality-filtered shotgun metagenomes (150 bp) and metadata (Table S6) were downloaded from

the Bioplatforms Australia data portal (https://data.bioplatforms.com/bpa/otu/metagenome).

Mean annual precipitation and mean annual temperature were determined for each sample using

the WorldClim2 database (60), and aridity index was determined using the Global Aridity

Database version 3 map at 30 arc-second resolution (61). The number of bacterial reads in these

Australian metagenomes ranged from 3 to 19 million (mean = 7.9 million). For comparison and

to include more humid sites, we used an additional, independent dataset of 209 surface soil (0-30

cm) shotgun metagenomes (Table S7) collected by the National Ecological Observatory Network

(NEON) using standardized methodology from 26 sites across the United States. We used

samples collected in 2023 and sequenced at the U.S. Department of Energy Joint Genome

Institute (JGI). Because OxyMetaG is assembly-free and does not require high sequencing depth

(Figure S6), we subsampled all metagenomes down to 5 million reads before running OxyMetaG

to reduce computational costs. The number of bacterial reads ranged from ~1.1 to 2.2 million

reads (mean = 1.7 million). Climate, soil, and other general metadata associated with each

metagenome were downloaded using a combination of the phyloNEON R package (62) and

neonUtilities R package (63). For statistical analyses, we converted the percent aerobe values,

which had a highly skewed distribution towards 100% values, to a proportion anaerobe value, for

use as a response in zero-inflated beta regression (ZIBR), where the response must be bounded

between 0 and 1 (64). Relationships between the predicted proportion of anaerobes and

ecosystem type, climate class, and continuous climate (mean annual precipitation and

temperature, aridity index) and soil variables (gravimetric water content, pH, total carbon, total

nitrogen, carbon: nitrogen ratio, nitrate, phosphorus, percent sand) were tested with ZIBR

implemented in the gamlss R package (65). Models including each predictor variable were

compared to a null model with the likelihood ratio test.

Second, we used 73 human gut metagenomes from Russia and Finland collected with

standardized methodology for the DIABIMMUNE cohort study (66). From all of the available

metagenomes in the study, we selected metagenomes from 9 individuals that were sampled

across at least 7 time points, had the first sample before the 93rd day of life (minimum days after

birth was 43), and had a sampling time range of at least 634 days (maximum day after birth was

1111) (Table S8). These metagenomes had an average of 1.5 million bacterial reads. We ran

linear mixed effect regression (LMER) models in the lme4 R package (67) between predicted

relative abundance of aerobic bacteria and days after birth. We compared two models with the

'anova' function in R: one that only allowed random intercepts for each subject ID, and one that

allowed random intercepts and slopes for each subject ID. This enabled us to assess the effect of

time (days since birth) and subject ID on the predicted relative abundance of aerobic bacteria. All

downstream statistical and graphical analyses were performed in R 4.5.1 (68).

**Acknowledgements**

We thank the NovoNordisk Foundation and Wellcome Trust for funding. We thank Hugh Cross

for assistance in acquiring the NEON metagenomes and metadata. We thank Antonio Fernandez-

Guerra for feedback on OxyMetaG.

**Data availability**

The 5520 genomes used in the genome-level models are publicly available on NCBI, with

accessions and metadata listed in Table S4. The 203 shotgun metagenomes used for the cross-

habitat validation are publicly available on NCBI, with accessions and metadata listed in Table

S5. The 331 BASE metagenomes from Australia are publicly available on the Bioplatforms

Australia data portal (https://data.bioplatforms.com/bpa/otu/metagenome); sample IDs and

metadata are provided in Table S6. The 209 NEON metagenomes from the United States in 2023

are publicly available on the JGI data portal (https://data.jgi.doe.gov/) under JGI_ID 509938,

with "2023" contained in the sample name; sample IDs and metadata are provided in Table S7.

The 73 human gut metagenomes are a subset of the publicly available DIABIMMUNE

metagenomes available on NCBI BioProject PRJNA290380; exact accessions and metadata are

detailed in Table S8.

**Code availability**

OxyMetaG is available from GitHub (https://github.com/cliffbueno/oxymetag) and PyPI. Data

analysis scripts for this manuscript can be found on Zenodo

(https://doi.org/10.5281/zenodo.18331946).

**References**

1.    Nutman AP, Bennett VC, Friend CRL, Van Kranendonk MJ, Chivas AR. 2016. Rapid

emergence of life shown by discovery of 3,700-million-year-old microbial structures. Nature

537:535–538.

2.    Poulton SW, Bekker A, Cumming VM, Zerkle AL, Canfield DE, Johnston DT. 2021. A 200-

million-year delay in permanent atmospheric oxygenation. Nature 592:232–236.

3.    Canfield DE. 2014. Oxygen: A Four Billion Year History. Princeton University Press,

Princeton.

4.    Krieg NR, Hoffman PS. 1986. Microaerophily and Oxygen Toxicity. Annual Review of

Microbiology 40:107–130.

5.  Davín AA, Woodcroft BJ, Soo RM, Morel B, Murali R, Schrempf D, Clark JW, Álvarez-

Carretero S, Boussau B, Moody ERR, Szánthó LL, Richy E, Pisani D, Hemp J, Fischer

WW, Donoghue PCJ, Spang A, Hugenholtz P, Williams TA, Szöllősi GJ. 2025. A

geological timescale for bacterial evolution and oxygen adaptation. Science 388:eadp1853.

6.  Koblitz J, Reimer LC, Pukall R, Overmann J. 2025. Predicting bacterial phenotypic traits

through improved machine learning using high-quality, curated datasets. Commun Biol

8:897.

7.  Edirisinghe JN, Goyal S, Brace A, Colasanti R, Gu T, Sadhkin B, Zhang Q, Kamimura R,

Henry CS. 2023. Machine Learning-Driven Phenotype Predictions based on Genome

Annotations. bioRxiv https://doi.org/10.1101/2023.08.11.552879.

8.  Weber Zendrera A, Sokolovska N, Soula HA. 2021. Functional prediction of environmental

variables using metabolic networks. Sci Rep 11:12192.

9.  Jabłońska J, Tawfik DS. 2019. The number and type of oxygen-utilizing enzymes indicates

aerobic vs. anaerobic phenotype. Free Radical Biology and Medicine 140:84–92.

10.  Weimann A, Mooren K, Frank J, Pope PB, Bremges A, McHardy AC. 2016. From

Genomes to Phenotypes: Traitar, the Microbial Trait Analyzer. mSystems

1:10.1128/msystems.00101-16.

11.  Jensen DB, Ussery DW. 2013. Bayesian prediction of microbial oxygen requirement.

F1000Res 2:184.

12.  Lingner T, Mühlhausen S, Gabaldón T, Notredame C, Meinicke P. 2010. Predicting

phenotypic traits of prokaryotes from protein domain frequencies. BMC Bioinformatics

11:481.

13. Hoffert M, Gorman E, Lladser ME, Fierer N. 2025. A periodic table of bacteria?: Mapping

bacterial diversity in trait space. bioRxiv https://doi.org/10.1101/2025.07.11.664459.

14. Wan S, Liu H, Zhu G, Geng Y, Li W, Chen L, Zhang Y, Han G. 2025. Decoding oxygen

preference: Machine learning discovers functional genes in Bacteria. Genomics

117:111095.

15. Barnum TP, Crits-Christoph A, Molla M, Carini P, Lee HH, Ostrov N. 2024. Predicting

microbial growth conditions from amino acid composition. bioRxiv

https://doi.org/10.1101/2024.03.22.586313.

16. Flamholz AI, Goldford JE, Richter PA, Larsson EM, Jinich A, Fischer WW, Newman DK.

2024. Annotation-free prediction of microbial dioxygen utilization. mSystems 9:e00763-24.

17. Sczyrba A, Hofmann P, Belmann P, Koslicki D, Janssen S, Dröge J, Gregor I, Majda S,

Fiedler J, Dahms E, Bremges A, Fritz A, Garrido-Oter R, Jørgensen TS, Shapiro N, Blood

PD, Gurevich A, Bai Y, Turaev D, DeMaere MZ, Chikhi R, Nagarajan N, Quince C, Meyer

F, Balvočiūtė M, Hansen LH, Sørensen SJ, Chia BKH, Denis B, Froula JL, Wang Z, Egan

R, Don Kang D, Cook JJ, Deltel C, Beckstette M, Lemaitre C, Peterlongo P, Rizk G,

Lavenier D, Wu Y-W, Singer SW, Jain C, Strous M, Klingenberg H, Meinicke P, Barton

MD, Lingner T, Lin H-H, Liao Y-C, Silva GGZ, Cuevas DA, Edwards RA, Saha S, Piro VC,

Renard BY, Pop M, Klenk H-P, Göker M, Kyrpides NC, Woyke T, Vorholt JA, Schulze-

Lefert P, Rubin EM, Darling AE, Rattei T, McHardy AC. 2017. Critical Assessment of

Metagenome Interpretation—a benchmark of metagenomics software. Nat Methods

14:1063–1071.

18. Fierer N, Barberán A, Laughlin DC. 2014. Seeing the forest for the genes: using

metagenomics to infer the aggregated traits of microbial communities. Front Microbiol 5.

19. Fierer N, Wood SA, Bueno de Mesquita CP. 2021. How microbes can, and cannot, be
used to assess soil health. Soil Biology and Biochemistry 153:108111.

20. Choix FJ, Palacios OA, Nevarez-Moorillón GV. 2023. Traditional and new proposals for
environmental microbial indicators—a review. Environ Monit Assess 195:1521.

21. Lacroix EM, Aeppli M, Boye K, Brodie E, Fendorf S, Keiluweit M, Naughton HR, Noël V,
Sihi D. 2023. Consider the Anoxic Microsite: Acknowledging and Appreciating
Spatiotemporal Redox Heterogeneity in Soils and Sediments. ACS Earth Space Chem
7:1592–1609.

22. Usharani K, Roopashree K, Naik D. 2019. Role of soil physical, chemical and biological
properties for soil health improvement and sustainable agriculture. Journal of
Pharmacognosy and Phytochemistry 8:1256–1267.

23. Brevik EC. 2010. Soil Health And Productivity, p. 106–137. *In* Soils, Plant Growth and Crop
Production. Eolss Publishers, United Kingdom.

24. Gray ME, Marland JRK, Dunare C, Blair EO, Meehan J, Tsiamis A, Kunkler IH, Murray AF,
Argyle D, Dyson A, Singer M, Potter MA. 2019. In vivo validation of a miniaturized
electrochemical oxygen sensor for measuring intestinal oxygen tension. American Journal
of Physiology-Gastrointestinal and Liver Physiology 317:G242–G252.

25. Rigottier-Gois L. 2013. Dysbiosis in inflammatory bowel diseases: the oxygen hypothesis.
ISME J 7:1256–1261.

26. Colgan SP, Campbell EL. 2017. Oxygen metabolism and innate immune responses in the
gut. Journal of Applied Physiology 123:1321–1327.

27.  Rivera-Chávez F, Lopez CA, Bäumler AJ. 2017. Oxygen as a driver of gut dysbiosis. Free
Radical Biology and Medicine 105:93–101.

28.  Guittar J, Shade A, Litchman E. 2019. Trait-based community assembly and succession of
the infant gut microbiome. Nat Commun 10:512.

29.  Tran PQ, Bachand SC, McIntyre PB, Kraemer BM, Vadeboncoeur Y, Kimirei IA,
Tamatamah R, McMahon KD, Anantharaman K. 2021. Depth-discrete metagenomics
reveals the roles of microbes in biogeochemical cycling in the tropical freshwater Lake
Tanganyika. ISME J 15:1971–1986.

30.  Dotinga BM, Mintzer JP, Moore JE, Hulscher JBF, Bos AF, Kooi EMW. 2020. Maturation of
Intestinal Oxygenation: A Review of Mechanisms and Clinical Implications for Preterm
Neonates. Front Pediatr 8.

31.  de Muinck EJ, Trosvik P. 2018. Individuality and convergence of the infant gut microbiota
during the first year of life. Nat Commun 9:2233.

32.  Sanidad KZ, Zeng MY. 2020. Neonatal gut microbiome and immunity. Curr Opin Microbiol
56:30–37.

33.  Carini P, Marsden PJ, Leff JW, Morgan EE, Strickland MS, Fierer N. 2016. Relic DNA is
abundant in soil and obscures estimates of soil microbial diversity. 3. Nat Microbiol 2:1–6.

34.  Foley MM, Stone BWG, Caro TA, Sokol NW, Koch BJ, Blazewicz SJ, Dijkstra P, Hayer M,
Hofmockel K, Finley BK, Mack M, Marks J, Mau RL, Monsaint-Queeney V, Morrissey E,
Propster J, Purcell A, Schwartz E, Pett-Ridge J, Fierer N, Hungate BA. 2024. Growth rate
as a link between microbial diversity and soil biogeochemistry. Nat Ecol Evol 8:2018–2026.

35.    Kendall B. 2021. Recent Advances in Geochemical Paleo-Oxybarometers. Annual Review

of Earth and Planetary Sciences 49:399–433.

36.    Schober I, Koblitz J, Sardà Carbasse J, Ebeling C, Schmidt ML, Podstawka A, Gupta R,

Ilangovan V, Chamanara J, Overmann J, Reimer LC. 2025. BacDive in 2025: the core

database for prokaryotic strain data. Nucleic Acids Research 53:D748–D756.

37.    Parks DH, Chuvochina M, Rinke C, Mussig AJ, Chaumeil P-A, Hugenholtz P. 2022. GTDB:

an ongoing census of bacterial and archaeal diversity through a phylogenetically

consistent, rank normalized and complete genome-based taxonomy. Nucleic Acids

Research 50:D785–D794.

38.    Eddy SR. 2011. Accelerated Profile HMM Searches. PLOS Computational Biology

7:e1002195.

39.    Mistry J, Chuguransky S, Williams L, Qureshi M, Salazar GA, Sonnhammer ELL, Tosatto

SCE, Paladin L, Raj S, Richardson LJ, Finn RD, Bateman A. 2021. Pfam: The protein

families database in 2021. Nucleic Acids Research 49:D412–D419.

40.    Walsh C, Stallard-Olivera E, Fierer N. 2023. Nine (not so simple) steps: a practical guide to

using machine learning in microbial ecology. mBio 0:e02050-23.

41.    Huang W, Li L, Myers JR, Marth GT. 2012. ART: a next-generation sequencing read

simulator. Bioinformatics 28:593–594.

42.    Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina

sequence data. Bioinformatics 30:2114–2120.

43.    Buchfink B, Xie C, Huson DH. 2015. Fast and sensitive protein alignment using DIAMOND.

Nat Methods 12:59–60.

44.    Ramoneda J, Fan K, Lucas JM, Chu H, Bissett A, Strickland MS, Fierer N. 2024.

544            Ecological relevance of flagellar motility in soil bacterial communities. The ISME Journal

545            18:wrae067.

546    45.    Wood SN. 2017. Generalized Additive Models: An Introduction with R2nd edition.

547            Chapman and Hall/CRC.

548    46.    Li F, Hitch TCA, Chen Y, Creevey CJ, Guan LL. 2019. Comparative metagenomic and

549            metatranscriptomic analyses reveal the breed effect on the rumen microbiome and its

550            associations with feed efficiency in beef cattle. Microbiome 7:6.

551    47.    Candeliere F, Sola L, Raimondi S, Rossi M, Amaretti A. 2024. Good and bad dispositions

552            between archaea and bacteria in the human gut: New insights from metagenomic survey

553            and co-occurrence analysis. Synthetic and Systems Biotechnology 9:88–98.

554    48.    Sunagawa S, Acinas SG, Bork P, Bowler C, Eveillard D, Gorsky G, Guidi L, Iudicone D,

555            Karsenti E, Lombard F, Ogata H, Pesant S, Sullivan MB, Wincker P, de Vargas C. 2020.

556            Tara Oceans: towards global ocean ecosystems biology. Nat Rev Microbiol 18:428–445.

557    49.    Villanueva L, Bastiaan von Meijenfeldt FA, Westbye AB, Yadav S, Hopmans EC, Dutilh

558            BE, Sinninghe Damsté JS. 2021. Bridging the membrane lipid divide: bacteria of the FCB

559            group superphylum have the potential to synthesize archaeal ether lipids. The ISME

560            Journal 15:168–182.

561    50.    Broman E, Izabel-Shen D, Rodríguez-Gijón A, Bonaglia S, Garcia SL, Nascimento FJA.

562            2022. Microbial functional genes are driven by gradients in sediment stoichiometry,

563            oxygen, and salinity across the Baltic benthic ecosystem. Microbiome 10:126.

51.    Borton MA, Oliverio AM, Narrowe AB, Villa JA, Rinke C, Hoyt DW, Liu P, McGivern BB,

Bechtold EK, Ellenbogen JB, Daly RA, Smith GJ, Angle JC, Flynn RM, Freiburger AP,

Louie KB, Stemple B, Northen TR, Henry C, Miller CS, Morin TH, Bohrer G, Wrighton KC.

2025. Mapping the soil microbiome functions shaping wetland methane emissions. bioRxiv

https://doi.org/10.1101/2024.02.06.579222.

52.    Bissett A, Fitzgerald A, Meintjes T, Mele PM, Reith F, Dennis PG, Breed MF, Brown B,

Brown MV, Brugger J, Byrne M, Caddy-Retalic S, Carmody B, Coates DJ, Correa C,

Ferrari BC, Gupta VVSR, Hamonts K, Haslem A, Hugenholtz P, Karan M, Koval J, Lowe

AJ, Macdonald S, McGrath L, Martin D, Morgan M, North KI, Paungfoo-Lonhienne C,

Pendall E, Phillips L, Pirzl R, Powell JR, Ragan MA, Schmidt S, Seymour N, Snape I,

Stephen JR, Stevens M, Tinning M, Williams K, Yeoh YK, Zammit CM, Young A. 2016.

Introducing BASE: the Biomes of Australian Soil Environments soil microbial diversity

database. GigaSci 5:21.

53.    Oliverio AM, Bissett A, McGuire K, Saltonstall K, Turner BL, Fierer N. 2020. The Role of

Phosphorus Limitation in Shaping Soil Bacterial Communities and Their Metabolic

Capabilities. mBio 11:e01718-20.

54.    Brewer TE, Aronson EL, Arogyaswamy K, Billings SA, Botthoff JK, Campbell AN, Dove

NC, Fairbanks D, Gallery RE, Hart SC, Kaye J, King G, Logan G, Lohse KA, Maltz MR,

Mayorga E, O'Neill C, Owens SM, Packman A, Pett-Ridge J, Plante AF, Richter DD, Silver

WL, Yang WH, Fierer N. 2019. Ecological and Genomic Attributes of Novel Bacterial Taxa

That Thrive in Subsurface Soil Horizons. mBio 10:10.1128/mbio.01318-19.

55.    Wu X, Bei S, Zhou X, Luo Y, He Z, Song C, Yuan H, Pivato B, Liesack W, Peng J. 2023.

Metagenomic insights into genetic factors driving bacterial niche differentiation between

bulk and rhizosphere soils. Science of The Total Environment 891:164221.

56.    Wood DE, Lu J, Langmead B. 2019. Improved metagenomic analysis with Kraken 2.

Genome Biology 20:257.

57.    Lu J, Rincon N, Wood DE, Breitwieser FP, Pockrandt C, Langmead B, Salzberg SL,

Steinegger M. 2022. Metagenome analysis using the Kraken software suite. Nat Protoc

17:2815–2839.

58.    Treiber ML, Taft DH, Korf I, Mills DA, Lemay DG. 2020. Pre- and post-sequencing

recommendations for functional annotation of human fecal metagenomes. BMC

Bioinformatics 21:74.

59.    Bueno de Mesquita CP, Olm MR, Bissett A, Fierer N. 2025. High strain-level diversity of

Bradyrhizobium across Australian soils. ISME J wraf222.

60.    Fick SE, Hijmans RJ. 2017. WorldClim 2: new 1-km spatial resolution climate surfaces for

global land areas. International Journal of Climatology 37:4302–4315.

61.    Zomer RJ, Xu J, Trabucco A. 2022. Version 3 of the Global Aridity Index and Potential

Evapotranspiration Database. Sci Data 9:409.

62.    Cross H. 2024. phyloNEON: Utilities for working with genetic NEON   data. R package

version 0.1.1. https://github.com/NEONScience/phyloNEON.

63.    Lunch C, Laney C, Mietkiewicz N, Sokol E, Cawley K, NEON. 2025. neonUtilities: Utilities

for   Working with NEON Data. R package version 3.0.1. https://CRAN.R-

project.org/package=neonUtilities.

64.    Tang B, Frye HA, Gelfand AE, Silander JA. 2023. Zero-Inflated Beta Distribution

Regression Modeling. JABES 28:117–137.

65.   Rigby RA, Stasinopoulos DM. 2005. Generalized additive models for location, scale and

shape. Journal of the Royal Statistical Society: Series C (Applied Statistics) 54:507–554.

66.   Zhao L, Yang X, Liang Y, Zhang Z, Ding Y, Wang Y, Chen B, Wu J, Jin C, Zhao G, Li Z,

Zhang L. 2024. Temporal development and potential interactions between the gut

microbiome and resistome in early childhood. Microbiology Spectrum 12:e03177-23.

67.   Bates D, Mächler M, Bolker BM, Walker SC. 2015. Fitting linear mixed-effects models

using lme4. Journal of Statistical Software 67:1–48.

68.   R Core Team. 2025. R: A Language and Environment for Statistical   Computing.

https://www.R-project.org/.

**Tables and Figures**

Table 1. The 20 Pfams selected for predicting the oxygen tolerance of individual isolates in
BacDive, as well as the Pfam name, whether it is indicative of aerobic or anaerobic metabolism,
the percent prevalence in 4227 aerobic bacteria, the percent prevalence in 1293 anaerobic
bacteria, and the aggregated SHAP feature importance score.

| Pfam | Name | Oxygen | Aerobe prev. | Anaerobe prev. | FI |
|------|------|--------|--------------|----------------|-----|
| PF00510 | Cytochrome c oxidase subunit III | aerobic | 94.58 | 12.22 | 0.155 |
| PF00296 | Luciferase-like monooxygenase | aerobic | 89.21 | 13.92 | 0.060 |
| PF16870 | 2-oxoglutarate dehydrogenase C-terminal | aerobic | 96.59 | 17.63 | 0.038 |
| PF00116 | Cytochrome C oxidase subunit II, periplasmic domain | aerobic | 94.42 | 14.23 | 0.038 |
| PF01521 | Iron-sulphur cluster biosynthesis | aerobic | 98.77 | 31.25 | 0.028 |
| PF05425 | Copper resistance protein D | aerobic | 77.88 | 9.59 | 0.023 |
| PF17773 | UPF0176 acylphosphatase like domain | aerobic | 66.57 | 6.19 | 0.022 |
| PF00115 | Cytochrome C and Quinol oxidase polypeptide I | aerobic | 98.11 | 23.28 | 0.020 |
| PF00916 | Sulfate permease family | aerobic | 90.51 | 47.10 | 0.019 |
| PF08530 | X-Pro dipeptidyl-peptidase C-terminal non-catalytic domain | aerobic | 54.67 | 16.71 | 0.019 |
| PF01152 | Bacterial-like globin | aerobic | 74.50 | 6.11 | 0.018 |
| PF05721 | Phytanoyl-CoA dioxygenase (PhyH) | aerobic | 60.70 | 2.24 | 0.017 |
| PF00042 | Globin | aerobic | 53.58 | 5.49 | 0.016 |
| PF01871 | AMMECR1 | anaerobic | 3.38 | 50.66 | 0.074 |
| PF10371 | EKR | anaerobic | 6.60 | 65.82 | 0.058 |
| PF17910 | FeoB cytosolic helical domain | anaerobic | 10.34 | 74.25 | 0.043 |
| PF02579 | Dinitrogenase iron-molybdenum cofactor | anaerobic | 7.55 | 64.89 | 0.041 |
| PF13597 | Anaerobic ribonuceoside-triphosphate reductase | anaerobic | 16.87 | 88.86 | 0.032 |
| PF03063 | Prismane/CO dehydrogenase family | anaerobic | 6.53 | 68.99 | 0.028 |
| PF02906 | Iron only hydrogenase large subunit, C-terminal domain | anaerobic | 1.14 | 58.62 | 0.018 |

625 Table 2. Ensemble model statistics on the 20% holdout test dataset.

| | Precision | Recall | F1-score | Support |
|---|---|---|---|---|
| Aerobe | 0.97 | 0.98 | 0.97 | 845 |
| Anaerobe | 0.92 | 0.90 | 0.91 | 259 |
| | | | | |
| accuracy | | | 0.96 | 1104 |
| macro avg | 0.94 | 0.94 | 0.94 | 1104 |
| weighted avg | 0.96 | 0.96 | 0.96 | 1104 |

Figure 1. Overview of the development and use of OxyMetaG. 20 predictive Pfams were first

selected using a genome-level ensemble machine learning model (Random Forest, CatBoost,

XGboost) that utilized oxygen tolerance information from BacDive. Then a relationship between

the ratio of selected aerobic indicator genes (n = 13) and anaerobic indicator genes (n = 7) was

built based on simulated metagenomes with varying levels of known aerobes and anaerobes.

Lastly, the model was tested on 203 metagenomes from various habitats with known or inferred

differences in oxygen levels, as well as on 331 surface soil metagenomes from Australia and 209

surface soil metagenomes from the United States. The steps of extracting bacterial reads, running

DIAMOND BLASTX, and predicting the percent aerobe relative abundance are implemented in

the standalone Python package OxyMetaG, with the functions oxymetag extract, oxymetag

profile, and oxymetag predict.

Figure 2. Relationship between the percent of aerobe genomes in simulated metagenomes, and

the ratio of the RPK of aerobic indicator genes to anaerobic indicator genes, across two ranges of

sequencing depths. The model used for prediction in OxyMetaG is the one trained with data from

26 to 41 million reads.

Figure 3. Predicted relative abundance of aerobic bacteria across 16 habitat types. The x-axis is

sorted from left to right according to mean predicted percent abundance of aerobes. Different

letters represent significant differences (Tukey posthoc, $p < 0.05$).

Figure 4. The predicted percent aerobe relative abundance significantly increased a) with

increasing ratio of oxygen to hydrogen sulfide across the Black Sea water depth gradient (GAM,

p < 0.05), b) with increasing oxygen levels across sediments from the Baltic Sea (linear

regression, p < 0.05), and c) in surface waters compared to deeper hypoxic waters in Lake

Tanganyika (t-test, p < 0.05). Shown are a loess function in panel a, a linear model in panel b,

and different letters representing a significant difference according to a t-test in panel c.

Figure 5. Predicted relative abundance of aerobic bacteria across habitats in a) Australia and b)

the United States. The x-axis within each panel is sorted from left to right by mean predicted

percent aerobes. Insets show histograms; note the difference in axes scales between panels.

Grassland, forest, and shrubland habitats were shared between the two datasets, while only

Australia had dune samples and only the U.S. had wetland samples. Habitat significantly affected

the predicted relative abundance of aerobic bacteria in Australia (ZIBR, p = 0.0002) but not in

the U.S. (ZIBR, p = 0.22).

Figure 6. Predicted relative abundance of aerobic bacteria over time across 9 individuals from

the DIABIMMUNE cohort study (66). The predicted relative abundance of aerobic bacteria

significantly decreased over time (LMER, $\chi^2 = 18.9$, p < 0.001).