Title: Efficient transgene-free multiplexed genome editing via viral delivery of an

engineered TnpB.

Authors: Trevor Weiss[1], Maris Kamalu[1], Honglue Shi[2, 3, 4], Gabriel Wirnowski[1], Alice

Ingelsson[1], Jasmine Amerasekera[1], Kamakshi Vohra[2, 4, ‡], Marena I. Trinidad[2, 5], Zheng

Li[1], Emily Freitas[1], Noah Steinmetz[1], Charlie Ambrose[1], Kerry Chen[1], Jennifer A.

Doudna[2, 3, 4, 6, 7, 8, 9, 10, 11]*, Steven E. Jacobsen[1, 12]*

* Correspondence: doudna@berkeley.edu and jacobsen@ucla.edu

Affiliations:

[1]Department of Molecular, Cell and Developmental Biology, University of California at

Los Angeles, Los Angeles, CA, USA, 90095

[2]Innovative Genomics Institute, University of California, Berkeley, CA, USA, 94720

[3]Howard Hughes Medical Institute, University of California, Berkeley, CA, USA, 94720

[4]California Institute for Quantitative Biosciences (QB3), University of California,

Berkeley, Berkeley, CA, USA, 94720

[5]University of California, Berkeley-University of California, San Francisco Graduate

Program in Bioengineering, University of California, Berkeley, Berkeley, CA, USA,

94720

[6]Department of Molecular and Cell Biology, University of California, Berkeley, CA, USA,

94720

[7]Li Ka Shing Center for Genomic Engineering, University of California, Berkeley,

Berkeley, CA, USA, 94720

[8]Department of Chemistry, University of California, Berkeley, Berkeley, CA, USA, 94720

[9]Molecular Biophysics and Integrated Bioimaging Division, Lawrence Berkeley National

Laboratory, Berkeley, CA, USA, 94720

[10]Gladstone Institute of Data Science and Biotechnology, San Francisco, CA, USA,

94158

[11]Gladstone-UCSF Institute of Genomic Immunology, San Francisco, CA, USA, 94158

[12]Howard Hughes Medical Institute, University of California at Los Angeles, Los

Angeles, CA, USA, 90095

‡Present Address: Biochemistry, Quantitative Biology, Biophysics and Structural

Biology, Yale University, New Haven, CT, USA, 06511

Abstract:

Virus-induced genome editing (VIGE) using compact RNA-guided endonucleases is a

transformational new approach in plant biotechnology, enabling tissue-culture-

independent and transgene-free genome editing (Hu et al. 2025; Weiss et al. 2025; Liu

et al. 2025). We recently established a VIGE approach for heritable editing at single loci

in *Arabidopsis* by delivering ISYmu1 TnpB (Ymu1) and its guide RNA (gRNA) via

Tobacco Rattle Virus (TRV) (Weiss et al. 2025). Here, we greatly improved this

approach by devising a multiple gRNA expression system and by utilizing an

engineered high-activity Ymu1 variant (Ymu1-WFR) (Zhou et al. 2026) to develop an

efficient multiplexed genome editing platform.

Main Text:

To evaluate TRV-mediated multiplexing capabilities, we co-delivered RNA1 with two

RNA2 vectors targeting either *AtPDS3* gRNA12 or *AtCHLI1* gRNA4 to *Arabidopsis*

(Figure S1A). Amplicon sequencing (amp-seq) revealed editing almost exclusively at

one target site or the other (Figure S1B), suggesting viral superinfection exclusion

(Perdoncini Carvalho et al. 2022). We therefore sought to develop a system in which

both gRNAs could be expressed on a single RNA2 vector.

Through small RNA sequencing (RNA-seq) and protoplast editing assays using various

ωRNA lengths, we identified 127 nucleotides as the optimal ωRNA for genome editing

(Figures S2A-D). Next, we tested multiplexed arrays featuring tRNA, HDV, HDV-HH, or

a repeat as gRNA processing elements. Amp-seq analysis showed that while all

designs enabled editing, HDV-based designs performed best at simultaneously editing

both sites in protoplasts (Figure S2E-G). Furthermore, polymerase chain reaction (PCR)

using primers spanning both sites suggested the occurrence of large deletions between

the two targets (Figure S2H).

Next, we tested the HDV multiplexed array design for TRV-mediated editing in planta

using vectors targeting *AtCHLI1* (gRNA4) and *AtPDS3* (gRNA12), incorporating a

tRNA$^{\text{Ile}}$ mobility sequence at the 3' end of the cargo (Figure S3A). gRNA4 targets the

gene body of *AtCHLI1* (biallelic edits create yellow tissue sectors) and gRNA12 targets

the promoter region upstream of the *AtPDS3* transcription start site (no visible

phenotype). After delivering TRV vectors, we did not observe any phenotypic evidence

of editing. Suspecting inefficient mobility or processing of the RNA2 cargo, we tested

three additional constructs containing a tRNA$^{\text{Ile}}$ downstream of each HDV ribozyme

(Figure 1A). After TRV delivery, yellow sectors appeared on leaves for all three vectors,

indicating biallelic edits at *AtCHLI1* (Figure 1B). Yellow sectored plants infected with

pTW2278 and pTW2279 displayed average editing efficiencies of 25.6% for *AtCHLI1*

and 30.2% for *AtPDS3* (Figure 1C), and those infected with pTW2498 showed 31.3% at

AtCHL1 gRNA4 and 5.2% at the *AtCHL1* gRNA6 (Figure 1D).

To characterize the inheritance of edited alleles, we selected two plants: pTW2278_17

and pTW2278_54. We observed yellow progeny at frequencies of 4.7% and 1.7%,

respectively (Figure 1E, S3B). Among the yellow seedlings, 76.9% (pTW2278_17) and

66.7% (pTW2278_54) of them harbored biallelic edits at both loci (Figure 1E, S3C).

These data demonstrate that TRV effectively delivers Ymu1 and multiple gRNAs for

heritable multiplexed editing, and that biallelic editing at one locus is highly predictive of

biallelic editing at the second target site.

We recently developed an engineered Ymu1 variant (Ymu1-WFR) with improved editing

capabilities (Zhou et al. 2026). To evaluate Ymu1-WFR efficiency using TRV, we

targeted three published *AtCHLI1* target sites: gRNA4, gRNA6, and gRNA9 (Weiss et

al. 2025). Infected plants showed a strong yellow phenotype for all three gRNAs (Figure

1F). Amp-seq revealed average editing efficiencies of 18.6%, 14.7%, and 9.8%,

respectively, much higher (up to 9.8-fold) than WT Ymu1 (Weiss et al. 2025) (Figure

1G).

To assess the impact of the WFR variant on multiplexed editing efficiency, we replaced

the wild type (WT) Ymu1 sequence in pTW2278, pTW2279, and pTW2498 with Ymu1-

WFR (Figure S3D). Following TRV delivery, we observed much more pronounced

phenotypic evidence of editing than with WT Ymu1 (Figure 1H compared with Figure

1B). Amp-seq of yellow sectored plants confirmed enhanced editing: pTW2657 and

pTW2658 averaged 48.9% for *AtCHLI1* (gRNA4) and 41.3% for *AtPDS3* (gRNA12)

(Figure 1I), while pTW2655 averaged 45.3% and 28.1% at the two *AtCHLI1* sites

(Figure 1J). Consistent with the protoplast result (Figure S2H), PCR analysis using

primers spanning the *AtCHLI1* gRNA4 and gRNA6 sites revealed large deletions

(Figures S3E, S3F).

By optimizing the gRNA array design, and incorporating the highly active engineered

Ymu1-WFR variant, we developed an efficient multiplexed editing platform that

bypasses the need for transgenesis. Given the broad host range of TRV, we anticipate

this approach will be adaptable to commercially relevant crop species. Additionally, the

ability to generate large deletions should expand this system's utility for regulatory

element engineering. Finally, this multiplexed system may enable the study of

embryonic lethal genes by utilizing *AtCHLI1* as a visual marker; the yellow somatic

sectors should facilitate the identification of tissue harboring biallelic knockout of a gene

of interest.

Acknowledgements:

Supported by NSF Plant Genome Research Program grant (2334027) to S.E.J. and

J.A.D. S.E.J. and J.A.D. are HHMI Investigators. H.S. was supported by the Jane Coffin

Childs Memorial Fund for Medical Research and a K99 award from the NIH

(K99GM160778).

Author Contributions:

T.W. and S.E.J. designed research. T.W., H.S., J.A.D. and S.E.J. interpreted data. T.W.

and S.E.J. wrote the paper. T.W., M.K., H.S., G.W., A.I., J.A., K.V., M.T., Z.L., E.F.,

N.S., C.A., and K.C. performed experiments.

Data Availability:

Amp-seq data will be accessible at NCBI Sequence Read Archive upon publication.

RNA-seq data is available at GEO accession: GSE316183.

Competing interests:

T.W., M.K, H.S., J.A.D. and S.E.J. have filed patents related to this work. S.E.J. is a
cofounder and consultant for Inari Agriculture and a consultant for Terrana Biosciences,
Invaio Sciences, Sail Biomedicines and Zymo Research. J.A.D. is a cofounder of
Azalea Therapeutics, Caribou Biosciences, Editas Medicine, Evercrisp, Scribe
Therapeutics and Mammoth Biosciences, a scientific advisory board member at
Evercrisp, Caribou Biosciences, Scribe Therapeutics, Mammoth Biosciences, The
Column Group and Inari, an advisor for Aditum Bio, the Chief Science Advisor to Sixth
Street, a Director at Johnson & Johnson, Altos and Tempus, and has a research project
sponsored by Apple Tree Partners.

References:

Hu, Manman, Lingran Zhang, Luis Herrera-Estrella, and Degao Liu. 2025.
"Heritable, Tissue Culture-Independent and Transgene-Free Genome Editing in
Plants via Viral Delivery of CRISPR/AsCas12f." *Plant Biotechnology Journal*, no.
pbi.70315 (September). https://doi.org/10.1111/pbi.70315.

Liu, Ye, Trevor Weiss, Jinhee Lee, et al. 2025. "Virus Induced Transgene- and
Tissue-Culture Free Heritable Genome Editing in Tomato." In *bioRxiv*. November 9.
https://doi.org/10.1101/2025.11.08.687402.

Perdoncini Carvalho, Camila, Ruifan Ren, Junping Han, and Feng Qu. 2022.
"Natural Selection, Intracellular Bottlenecks of Virus Populations, and Viral
Superinfection Exclusion." *Annual Review of Virology* 9 (1): 121–137.

Weiss, Trevor, Maris Kamalu, Honglue Shi, et al. 2025. "Viral Delivery of an RNA-
Guided Genome Editor for Transgene-Free Germline Editing in Arabidopsis."
Nature Plants 11 (5): 967–976.

Zhou, Zehan, Iren Saffarian-Deemyad, Honglue Shi, et al. 2026. "Stepwise DNA

Unwinding Gates TnpB Genome-Editing Activity." In *bioRxiv*. BioRxiv, January 9. https://doi.org/10.64898/2026.01.09.698545.

Figure 1: Heritable and transgene-free multiplexed genome editing in *Arabidopsis*

via viral delivery of Ymu1. (A) Schematic representation of the cargo being expressed

by the PEBV promoter in TRV RNA2. *AtCHLI1* gRNA4 (yellow), *AtPDS3 gRNA12*

(black), and *AtCHLI1* gRNA6 (green) were used in the multiplex TRV experiments. The

plasmid ID is listed to the left of each construct. **(B)** Yellow sector phenotype observed

from plants infected with the vectors from panel A. The plant ID (construct_plant) is

listed below each picture. **(C and D)** Editing efficiency (y-axis) of plants displaying the

yellow sector phenotype (x-axis) infected with pTW2278, pTW2279 and pTW2498. **(E)**

Heritability data from plants infected with pTW2278_17 and pTW2278_54. **(F)** Yellow

sector phenotype observed from plants infected with Ymu1-WFR targeting *AtCHLI1*

(gRNA4, gRNA6, and gRNA9). The site being targeted is listed below each picture. **(G)**

Editing efficiency (y-axis) of plants infected with Ymu1-WFR targeting *AtCHLI1* (x-axis).

Each dot represents an individual plant. The yellow dashed line on each bar indicates

the average editing efficiency previously reported (Weiss et al. 2025). **(H)** Yellow sector

phenotype observed from plants infected with Ymu1-WFR targeting *AtCHLI1* gRNA4

and *AtPDS3* gRNA12 (pTW2657 and pTW2658) or *AtCHLI1* gRNA4 and *AtCHLI1*

gRNA6 (pTW2655). The plant ID (construct_plant) is listed below each picture. **(I and J)**

Editing efficiency (y-axis) of plants displaying the yellow sector phenotype (x-axis)

infected with pTW2657, pTW2658 and pTW2655.

    Supplementary Materials:

    Supplementary Figure 1: Co-delivery of RNA1 and two RNA2 vectors does not

    enable multiplexed genome editing. (A) Schematic representation of TRV. The cargo

    and plasmid ID are listed below RNA2. **(B)** Editing efficiency (y-axis) of plants ($n = 84$)

    co-infected with RNA1 and two RNA2 vectors targeting *AtCHLI1* gRNA4 and *AtPDS3*

    gRNA12 (x-axis). Each gray line represents a single plant, with the black dot at each

    end of the gray line corresponding to the editing efficiency of each target site.

Supplementary Figure 2: Development of Ymu1 multiplexed gRNA arrays for plant

gene editing. (A) Schematic of plasmid used in *E. coli* RNA-seq experiment. The

tetracycline promoter (pTet, arrow) was used to express the Ymu1-ωRNA (black),

gRNA (yellow), and HDV ribozyme (red). **(B)** RNA-seq in *E. coli* using the plasmid in

panel A. Read coverage (y-axis) is displayed for each Ymu1 single transcript feature (x-

axis). **(C)** Schematic of plasmids expressing the Ymu1-ωRNA in a single transcript

expression format or split expression format targeting *AtPDS3* gRNA2. In the split

expression plasmids the Ymu1 coding sequence was expressed by the *AtUBQ10*

promoter (UBQ10p) and the ωRNA was expressed using the U6 promoter (U6p). **(D)**

Editing efficiency (y-axis) using single expression and split expression constructs with

varying lengths of ωRNA (x-axis). Each dot represents an individual transfection with

bars indicating the standard error of the mean (SEM). **(E)** Schematic of *AtPDS3* gene

with the arrows indicating the orientation of gRNA2 and gRNA9, located roughly 2.4kb

apart. **(F)** Schematic of vectors used in the *Arabidopsis* protoplast multiplexed editing

experiment. **(G)** Editing efficiency (y-axis) of the multiplex arrays (x-axis) tested in

Arabidopsis protoplast targeting *AtPDS3* (gRNA2 and gRNA9). The subscript below

each gRNA indicates its placement in the multiplex array as depicted in panel F. **(H)**

PCR gel electrophoresis image using primers spanning *AtPDS3* gRNA2 and *AtPDS3*

gRNA9 target sites. Deletion between the two target sites corresponds to a ~850bp

band.

Supplementary Figure 3: TRV-delivery of Ymu1 for multiplexed genome editing.

(A) Schematic of multiplex vectors targeting *AtCHLI1* gRNA4 and *AtPDS3* gRNA12,

containing a single 3' tRNA[Ileu] . **(B)** Representative image of yellow and green progeny

seedlings from plant pTW2278_17. All yellow seedlings harbored biallelic edits at

AtCHLI1 gRNA4 target site. **(C)** Sanger sequencing trace file screenshots from a yellow

seedling in panel B, harboring homozygous edited alleles at *AtCHLI1* gRNA4 (4bp

deletion) and *AtPDS3* gRNA12 (11bp deletion). **(D)** Schematic representation of the

Ymu1-WFR multiplexed gRNA cargo being expressed by the PEBV promoter in TRV

RNA2. **(E)** Schematic of *AtCHLI1* gene with the arrows indicating the orientation of

gRNA4 and gRNA6, located approximately 275bp apart. **(F)** PCR gel electrophoresis

image using primers spanning *AtCHLI1* gRNA4 and *AtCHLI1* gRNA6 target sites.

Deletion between the two target sites corresponds to a ~420bp band. The plant ID

(construct_plant) is listed above each well. The TnpB (Ymu1 or Ymu1-WFR) and

gRNA(s) is listed above the plant ID.

Plasmid	Plasmid description
pTW1169	TRV RNA1
pTW2154	TRV2 expressing wild type Ymu1 protein and *AtCHLI1* gRNA4
pTW2082	TRV2 expressing wild type Ymu1 protein and *AtPDS3* gRNA12
pHS516	Wild type Ymu1 single transcript expression plasmid for RNA-seq in E.coli
pTW2064	Wild type Ymu1 with no ωRNA
pTW2071	Wild type Ymu1 with 100 nt ωRNA targeting *AtPDS3* gRNA2
pTW2072	Wild type Ymu1 with 110 nt ωRNA targeting *AtPDS3* gRNA2
pTW2073	Wild type Ymu1 with 120 nt ωRNA targeting *AtPDS3* gRNA2
pTW2074	Wild type Ymu1 with 127 nt ωRNA targeting *AtPDS3* gRNA2
pTW2075	Wild type Ymu1 with 130 nt ωRNA targeting *AtPDS3* gRNA2
pTW2076	Wild type Ymu1 with 150 nt ωRNA targeting *AtPDS3* gRNA2
pTW2077	Wild type Ymu1 with 170 nt ωRNA targeting *AtPDS3* gRNA2
pTW2078	Wild type Ymu1 with 190 nt ωRNA targeting *AtPDS3* gRNA2
pTW2079	Wild type Ymu1 with 210 nt ωRNA targeting *AtPDS3* gRNA2
pTW2080	Wild type Ymu1 with 230 nt ωRNA targeting *AtPDS3* gRNA2
pTW2081	Wild type Ymu1 with 250 nt ωRNA targeting *AtPDS3* gRNA2
pTW2109	Repeat design multiplex vector expressing *AtPDS3* gRNA2 first in the array and *AtPDS3* gRNA9 second
pTW2110	Repeat design multiplex vector expressing *AtPDS3* gRNA9 first in the array and *AtPDS3* gRNA2 second
pTW2137	tRNA design multiplex vector expressing *AtPDS3* gRNA2 first in the array and *AtPDS3* gRNA9 second
pTW2140	tRNA design multiplex vector expressing *AtPDS3* gRNA9 first in the array and *AtPDS3* gRNA2 second

pTW2148	HDV design multiplex vector expressing *AtPDS3* gRNA2 first in the array and *AtPDS3* gRNA9 second
pTW2138	HDV design multiplex vector expressing *AtPDS3* gRNA9 first in the array and *AtPDS3* gRNA2 second
pTW2136	HDV-HH design multiplex vector expressing *AtPDS3* gRNA2 first in the array and *AtPDS3* gRNA9 second
pTW2139	HDV-HH design multiplex vector expressing *AtPDS3* gRNA9 first in the array and *AtPDS3* gRNA2 second
pMK061	Wild type Ymu1 and *AtPDS3* gRNA2
pMK067	Wild type Ymu1 and *AtPDS3* gRNA9
pTW2229	TRV2 multiplex vector expressing wild type Ymu1, *AtCHLI1* gRNA4 and *AtPDS3* gRNA12
pTW2230	TRV2 multiplex vector expressing wild type Ymu1, *AtPDS3* gRNA12 and *AtCHLI1* gRNA4
pTW2658	TRV2 multiplex vector expressing Ymu1-WFR, *AtCHLI1* gRNA4 and *AtPDS3* gRNA12
pTW2657	TRV2 multiplex vector expressing Ymu1-WFR, *AtPDS3* gRNA12 and *AtCHLI1* gRNA4
pTW2655	TRV2 multiplex vector expressing Ymu1-WFR, *AtCHLI1* gRNA6 and *AtCHLI1* gRNA4
pTW2278	TRV2 multiplex vector expressing wild type Ymu1, *AtCHLI1* gRNA4 and *AtPDS3* gRNA12
pTW2279	TRV2 multiplex vector expressing wild type Ymu1, *AtPDS3* gRNA12 and *AtCHLI1* gRNA4
pTW2498	TRV2 multiplex vector expressing wild type Ymu1, *AtCHLI1* gRNA6 and *AtCHLI1* gRNA4
pTW2652	TRV2 expressing Ymu1-WFR protein and *AtCHLI1* gRNA4

pTW2653	TRV2 expressing Ymu1-WFR protein and *AtCHLI1* gRNA6
pTW2660	TRV2 expressing Ymu1-WFR protein and *AtCHLI1* gRNA9
pMK435	Cloning vector for PaqCI Golden Gate assembly of Ymu1 guides into TRV2

351 **Supplementary Table 1:** Plasmids and their descriptions used in this study.

Gene ID	Target site(s)	Target site sequence	Forward primer sequence	Reverse primer sequence	Purpose
AT4G14210	*AtPDS3* gRNA2	aaggcaaattcgccgc	ACACTCTTT CCCTACAC GACGCTCT TCCGATCTg aagcagttgtga gttaagttggaga	GTGACTGGA GTTCAGACG TGTGCTCTT CCGATCTttgt cttaagcgcttgag aagtgg	NGS amplicon sequencing
AT4G18480	*AtCHLI1* gRNA4	CTGTTACCTGA GATTA	ACACTCTTT CCCTACAC GACGCTCT TCCGATCT GTGGTGTT ATGATTATG GGAGATAG AG	GTGACTGGA GTTCAGACG TGTGCTCTT CCGATCTTC GCAATAACA GGAACTTGC TC	NGS amplicon sequencing
AT4G18480	*AtCHLI1* gRNA6	GAAGTTAATCT CTTGG	ACACTCTTT CCCTACAC GACGCTCT TCCGATCTA AGCCTTTGA GCCTGGTT TG	GTGACTGGA GTTCAGACG TGTGCTCTT CCGATCTCG GGTGAGAAA TCGAAATCC C	NGS amplicon sequencing
AT4G18480	*AtCHLI1* gRNA9	CGGTTTGGTAT GCATG	ACACTCTTT CCCTACAC GACGCTCT TCCGATCT GCGAGGTT TATCTTGAT	GTGACTGGA GTTCAGACG TGTGCTCTT CCGATCTTC ACGGAAATC CTTTGGGTT	NGS amplicon sequencing

			CGGTTC	ACTA	
AT4G14210	*AtPDS3* gRNA12	gcgttggagcatataa	ACACTCTTT CCCTACAC GACGCTCT TCCGATCTg aaccgacccga gaagagatttg	GTGACTGGA GTTCAGACG TGTGCTCTT CCGATCTtgg aatacacacatttg tacaacca	NGS amplicon sequencing
AT4G14210	*AtPDS3* gRNA9	agctttgaaccggttt	ACACTCTTT CCCTACAC GACGCTCT TCCGATCTg cgctaaactttat aaaccctgatg	GTGACTGGA GTTCAGACG TGTGCTCTT CCGATCTtatg attgccagtgaga taaagattc	NGS amplicon sequencing
AT4G14210	*AtPDS3* gRNA12	see above	tgttacacaactt attatgattgggct	ggaagtagccga taacaaaatgga g	Sanger sequencing
AT4G18480	*AtCHLI1* gRNA4	see above	ACACTCTTT CCCTACAC GACGCTCT TCCGATCTg aagttaggaaca tgattgttgttggt	GTGACTGGA GTTCAGACG TGTGCTCTT CCGATCTTC ACGGAAATC CTTTGGGTT ACTA	Sanger sequencing
AT4G14210	*AtPDS3* gRNA2 and *AtPDS3* gRNA9	see above	ACACTCTTT CCCTACAC GACGCTCT TCCGATCTa ctccattttgttatc ggctacttcc	GTGACTGGA GTTCAGACG TGTGCTCTT CCGATCTgtg cttccattagtgagt aagaaact	PCR spanning both target sites

			ACACTCTTT CCCTACAC GACGCTCT TCCGATCTC ATTTGCAGC TATAGTAG GGCA	GTGACTGGA GTTCAGACG TGTGCTCTT CCGATCTTC ACGGAAATC CTTTGGGTT ACTA	
AT4G18480	*AtCHLI1* gRNA4 and *AtCHLI1* gRNA6	see above			PCR spanning both target sites

377 **Supplementary Table 2:** Target sites and primers used in this study.

Methods:

<u>Plasmids used in this study</u>

To generate the Ymu1 plasmids with varying ωRNA lengths, NEB HiFi assembly was performed using two PCR amplicons and pTW2064 (the no ωRNA plasmid backbone). One amplicon contained the U6 promoter and the other amplicon contained the ωRNA. First, pTW2064 underwent a restriction enzyme digestion with SpeI (R3133) and Quick CIP (M0525) to linearize the plasmid. Next, both amplicons and the digested and purified pTW2064 plasmid were used in an NEBuilder HiFi DNA Assembly (E2621) reaction.

Multiplex vectors used in the protoplast experiment expressing wild type Ymu1, *AtPDS3* gRNA2 and *AtPDS3* gRNA9 were created using a golden gate reaction. PCR amplicons or gene fragments (Twist Bioscience) containing the first gRNA, processing sequence, 127-nt ωRNA and the second gRNA were inserted into the pMK025 vector (Weiss et al. 2025) using a PaqCI (R0745) golden gate reaction.

Multiplex TRV2 vectors were created using a golden gate reaction assembly. First, PCR was performed to generate two amplicons; one amplicon contained the first gRNA in the array, an HDV ribozyme and tRNA$^{\text{Ile}}$, and the second amplicon contained the 127-nt ωRNA and second gRNA in the array. Amplicons contained overhangs with PaqCI (R0745) sites to facilitate assembly. Both amplicons and pMK435 (Weiss et al. 2025) were then used in a PaqCI golden gate reaction.

TRV2 vectors containing the Ymu1-WFR variant were created using NEBuilder HiFi DNA Assembly (E2621). The TRV2 vectors were constructed by digesting the wild type Ymu1 TRV2 vectors with HpaI (R0105) and Eco53KI (R0116) restriction enzymes. A gene fragment (Twist Bioscience) containing the Ymu1-WFR sequence was then cloned into the digested plasmid using NEBuilder HiFi DNA Assembly (E2621).

430   Following assembly, all plasmids were transformed into NEB 10-beta (C3019) *E.coli*
431   and validated by whole-plasmid sequencing. Plasmid names and descriptions can be
432   found in Table S1.
433
<u>Plant gene editing experiments</u>
Arabidopsis mesophyll protoplasts were isolated according to previously described
methods (Yoo et al. 2007). For protoplast transfection, 20 µg of plasmid was placed at
the bottom of each tube and brought to a total volume of 20 µl with water. Then, 200 µl
of protoplasts were added to the plasmid solution, followed by 220 µl of fresh, sterile
polyethylene glycol (PEG)-CaCl$_2$ solution. The samples were mixed by gentle tapping
and incubated at room temperature for 10 minutes. Transfection was terminated by
adding 880 µl of W5 solution and inverting the tubes two to three times. Protoplasts
were subsequently pelleted by centrifugation at 100 RCF for 3 minutes, resuspended in
1 ml of WI solution, and plated in 6-well plates precoated with 5% calf serum. The cells
were incubated at 26°C for 48 hours. During this incubation period, a 37°C heat-shock
treatment was applied for 2 hours, beginning 16 hours post-transfection. At 48 hours
post-transfection, the protoplasts were collected for genomic DNA extraction.
447
TRV delivery was performed as previously described (Weiss et al. 2025; Nagalakshmi
et al. 2022). TRV1 and TRV2 vectors were introduced into *Agrobacterium tumefaciens*
strain GV3101, and cultures were grown in 200 ml of lysogeny broth (LB) with
antibiotics for 18 hours at 28°C. The cultures were centrifuged at 3,500 × g for 20
minutes; the supernatant was discarded, and the pellets were resuspended in 200 ml of
sterile water. This suspension was centrifuged again at 2,109 × g for 10 minutes. After
discarding the supernatant, the pellet was resuspended in sterile agro-infiltration buffer
(10 mM MgCl$_2$, 10 mM MES, and 250 µM acetosyringone) to a final optical density
OD$_{600}$ of 1.5 and incubated at 23°C for 3 hours with slow shaking. Following incubation,
the *Agrobacterium* cultures harboring TRV1 and TRV2 were mixed in a 1:1 ratio (co-
delivery experiment used a ratio of two parts TRV1, one part pTW2154, and one part
pTW2082). 15 ml of this mixture was delivered to 10-day-old seedlings via agroflood co-

culture. After 4 days, the seedlings were transplanted to soil and grown at room

temperature for the remainder of their life cycle.

Approximately 12 weeks after viral delivery, seeds were harvested from TRV-infected

plants. Seeds were sown on ½ MS plates supplemented with 3% sucrose and stratified

at 4°C in the dark for 5 days. Following stratification, the seeds were moved to a growth

room under a 16-h light/8-h dark cycle at 23°C for 10–12 days. A subset of plants was

then sampled for genotyping by collecting a single piece of leaf tissue. DNA was

extracted using the Invitrogen Platinum Direct PCR Universal Master Mix (A44647500)

according to the manufacturer's instructions. Finally, the extracted DNA was analyzed

via PCR and Sanger sequencing using the primers listed in Table S2.

<u>Amplicon sequencing and analysis</u>

Genomic DNA was extracted from protoplast samples using the Qiagen DNeasy Plant

Mini Kit (Qiagen, 69106). For plants subjected to agroflood, three leaf tissue samples

distal to the TRV delivery site were pooled per plant about 3-4 weeks after viral delivery.

The ratio of green to yellow tissue samples collected were selected in a manner that

most accurately reflected the phenotype of the entire plant. Once collected, all tissue

samples were frozen overnight at −80°C and ground. DNA was subsequently extracted

using the Invitrogen Platinum Direct PCR Universal Master Mix (A44647500) according

to the manufacturer's instructions. The DNA was then used for next-generation

amplicon sequencing.

Amplicon sequencing was performed using single-end sequencing on the Illumina

NovaSeqX platform. Libraries were constructed using a two-step PCR approach. First,

target regions were amplified for 25 cycles using primers flanking the TnpB target site

(Table S2), followed by purification with 1.0× AMPure XP beads (Beckman Coulter,

A63881). Then, samples underwent 12 additional cycles of amplification to attach

Illumina indexing primers, followed by a 0.7× AMPure XP cleanup. Final libraries were

normalized and pooled for sequencing. Amplicon sequencing analysis was performed

using the CrispRvariants R package (v.1.14.0) as previously described (Weiss et al.

2025).

<u>Bacterial RNA expression and sequencing</u>

For heterologous RNA expression, *E. coli* transformed with pHS516 (Figure S2A, Table

S1) and grown in LB medium to an OD_{600} of 0.6 and induced with 10 mM arabinose at

16°C overnight. Following overnight growth, total RNA from cells was extracted using a

hot formamide method, in which pelleted *E. coli* were resuspended in an 18 mM EDTA

and 95% formamide solution and lysed by incubating at 65ºC for 5 minutes. After

centrifugation to pellet cell debris, the supernatant containing total RNA was purified

using the RNA Clean & Concentrator-5 Kit (Zymo Research R1013) following

manufacturer protocols. Approximately 200 ng of purified total RNA was subjected to

rRNA depletion using the NEBNext rRNA Depletion Kit for Bacteria (New England

Biolabs E7850L) following manufacturer protocols. rRNA-depleted samples were again

purified using the RNA Clean & Concentrator-5 Kit and subjected to end repair and

small-RNA-seq library prep as previously described (Zhou et al. 2026). Sequencing was

performed using an Illumina NextSeq 1000/2000 P2 v3 kit. Following sequencing,

paired-end reads were trimmed and merged using fastp (v0.23.2) (Chen et al. 2018).

Merged reads of 100–150 nucleotides were selected with fastq-filter (v0.3.0;

https://github.com/LUMC/fastq-filter) and aligned to the Ymu1 locus using BWA

(v0.7.17) (Li 2013). Alignments were sorted with SAMtools (v1.17) (Li et al. 2009) and

converted to BigWig format using deepTools2 (v3.5.1) (Ramírez et al. 2014), then

visualized in the Integrative Genomics Viewer (IGV) (Robinson et al. 2011).

Supplementary Materials References:

Chen, Shifu, Yanqing Zhou, Yaru Chen, and Jia Gu. 2018. "Fastp: An Ultra-Fast All-in-
One FASTQ Preprocessor." *Bioinformatics (Oxford, England)* 34 (17): i884–i890.

Li, Heng. 2013. "Aligning Sequence Reads, Clone Sequences and Assembly Contigs
with BWA-MEM." In *arXiv [q-bio.GN]*. March 16. arXiv.
http://arxiv.org/abs/1303.3997.

Li, Heng, Bob Handsaker, Alec Wysoker, et al. 2009. "The Sequence Alignment/Map
Format and SAMtools." *Bioinformatics (Oxford, England)* 25 (16): 2078–2079.

Nagalakshmi, Ugrappa, Nathan Meier, Jau-Yi Liu, Daniel F. Voytas, and Savithramma
P. Dinesh-Kumar. 2022. "High-Efficiency Multiplex Biallelic Heritable Editing in
Arabidopsis Using an RNA Virus." *Plant Physiology* 189 (3): 1241–1245.

Ramírez, Fidel, Friederike Dündar, Sarah Diehl, Björn A. Grüning, and Thomas Manke.
2014. "deepTools: A Flexible Platform for Exploring Deep-Sequencing Data."
Nucleic Acids Research 42 (Web Server issue): W187–91.

Robinson, James T., Helga Thorvaldsdóttir, Wendy Winckler, et al. 2011. "Integrative
Genomics Viewer." *Nature Biotechnology* 29 (1): 24–26.

Weiss, Trevor, Maris Kamalu, Honglue Shi, et al. 2025. "Viral Delivery of an RNA-
Guided Genome Editor for Transgene-Free Germline Editing in Arabidopsis."
Nature Plants 11 (5): 967–976.

Yoo, Sang-Dong, Young-Hee Cho, and Jen Sheen. 2007. "Arabidopsis Mesophyll
Protoplasts: A Versatile Cell System for Transient Gene Expression Analysis."
Nature Protocols 2 (7): 1565–1572.

Zhou, Zehan, Iren Saffarian-Deemyad, Honglue Shi, et al. 2026. "Stepwise DNA
Unwinding Gates TnpB Genome-Editing Activity." In *bioRxiv*. BioRxiv, January 9.
https://doi.org/10.64898/2026.01.09.698545.