Oculomotor dance learning task: Implications for audio-visual cued spatial learning

Michael Petrovski, Susu Beheiry, Udichi U. Das, Simran Rooprai, Ashkan Karimi, Jenny R. Simon, Rachel J. Bar, Joseph F. X. DeSouza

Abstract

Learning dance of a motor sequence learning involves the coordination of both oculomotor and manual motor systems through the practiced repetition of a fixed sequence of actions, resulting in automatized execution of movement through habit learning. This study aims to address whether a sequence-based learning paradigm centered on the visual-motor system can feasibly be measured while listening to music (Bar and DeSouza 2016). It aims to develop a new visual-motor-based learning paradigm with music, potentially promoting neuroplasticity and creating new interventional tools, building upon prior research that shows behavioural and putative neural changes following dance-based neurorehabilitation in people with Parkinson's disease (Bearss et al. 2024). Eye movements of 10 participants (8 female, 2 male) were tracked using the Eyelink 1000 Plus system during a 68-second eye-dance sequence. The experiment consisted of a learning phase, where participants observed the sequence five times with 30-second breaks, and a performance phase, where they performed the sequence five times from memory on a grey screen without visual cues. Music was incorporated into both phases to aid memorization of the 4 spatial locations. After each performance, the participant was shown a visual reinforcer and asked for their thoughts on how well they executed the dance. A visual reinforcer flashes one of three different colours: red, yellow, or green. Each colour corresponds to how many steps in the dance a participant performed correctly, with key points being: under one third, between one to two thirds, and over two thirds of total steps correct. Participants were

scored based on timing of the steps as well for exact (1.00), good (0.66), slightly off (0.33) or missed (0) steps. Data was analyzed using R4.3.1, MATLAB, and Experiment Builder: Data Viewer software. Results showed a significant improvement in performance accuracy between the first session (g1; M = 40%, SD = 7.2%) and the last session (g5; M = 69.7%, SD = 22.8%). A repeated-measures ANOVA revealed a significant main effect of session on performance accuracy, $F(4, 36) = 6.99$, $p < 0.001$, $\eta^2_G = 0.26$, indicating that accuracy significantly improved over sessions. Post-hoc Bonferroni comparisons showed that accuracy in later sessions was significantly higher than earlier sessions, suggesting a defined learning curve and consolidation of performance pattern across repeated practice. Similarly, there was significant improvement in timing accuracy between the first session g1; M = 0.29, SD = 0.06) and the fifth session (g5; M = 0.46, SD = 0.12). A repeated-measures ANOVA revealed a significant main effect of session on timing precision, $F(4, 36) = 11.67$, $p < 0.001$, $\eta^2_G = 0.25$, indicating significant improvements in temporal control and coordination over sessions. Post-hoc Bonferroni comparisons showed that timing precision significantly improved between early and late sessions (e.g, g1-g4, $p < 0.01$; g1-g5, $p < 0.001$), suggesting a defined learning curve and increase in precision across repeated practice. These findings suggest that visual-motor-based interventions have the potential to enhance motor and non-motor symptoms like depression and anxiety for neurodegenerative diseases such as Parkinson's Disorder (PD). The results provide a foundation for developing targeted therapies that integrate learning paradigms to improve functional outcomes, warranting further exploration of their long-term efficacy.

Introduction

Motor sequence learning involves the coordination of both oculomotor and manual motor systems through the practiced repetition of a fixed sequence of actions, resulting in automatized

execution of movement (Rubino et al., 2025). While extensive research regarding motor sequence learning has been explored involving neural bases of behaviour (Jäger et al., 2022, Tremblay et al., 2021), the oculomotor context remains an emerging field of motor sequence learning. Typical design of motor sequence learning tasks generally assess reaction times using limb movement in response to stimuli (Jeyarajan et al., 2024; Maceira-Elvira et al., 2022; Rubino et al., 2025; Sailer et al., 2005; Vieluf et al., 2015). Although a smaller subset of motor sequence learning literature, various studies found that eye movements predict motor sequence learning similar to that of physical movement (Coiner et al., 2019; Medimorec et al., 2021; Rubino et al., 2024), and have activational patterns closely resembling physical motor response (Rubino et al., 2024; Rubino et al., 2025; Sailer et al., 2005). Other studies have found auditory stimuli to enhance motor sequence learning (Leow et al., 2025), aiding response to visual stimuli. The use of auditory cues synchronously within sequence learning has been associated with increased development of explicit knowledge of learned sequences (Han et al., 2024). In addition, cues of this nature are found to aid in movement sequences that occur over longer periods of time (Bar & DeSouza, 2016; Leow et al., 2025). Understanding the components behind sequence learning that reinforce behaviours and show greater results in combination rather than individually creates a significant demand for future task design to seamlessly incorporate both facets.

Defining the underlying mechanisms of motor sequence learning, specifically the neurological communications between areas discussed prior, is increasingly critical. Neuroplasticity is the brain's ability to reorganize neural pathways in response to degeneration or loss of function within such pathways. To prompt neuroplastic change is to engage brain function of specified regions of interest with a task and to understand the compensations or new

connections made. Various methods of quantifying neuroplastic changes are documented (Barnstaple et al., 2021; Ferreira et al., 2025; Herzberg et al., 2024; Hänggi et al., 2010; Hehl et al., 2025; Tremblay et al., 2021), however the developments brought about by physical intervention or tasks related to movement will be pertinent to the following body of work. Systematic reviews have highlighted the importance that physical exercise or engagement can have in improving executive function and wellbeing in old age (Cheng et al., 2024; Du et al., 2025; Quan et al., 2025). Even in tasks involving minimal physical effort, greater activation of regions involved such as frontal eye fields (FEF) (Gonzalez et al., 2016), superior colliculus (Soetedjo et al., 2009), cerebellar oculomotor vermis (Kurkin et al., 2014), and SMA-related motor networks have been shown (Bonzano et al., 2015; Coiner et al., 2019; Rubino et al., 2024; Rubino et al., 2025). In addition, it has been noted that eye tracking proves to be a promising tool for assessment and clinical applications as they can potentially be applied towards measurement of neuroplastic changes in neurodegenerative diseases (Culicetto et al., 2025; Diotaiuti et al., 2025). Providing analysis of specific areas involved in neuroplastic change brought about through motor engagement, whether ocular or physical, remains pivotal.

As a form of physical movement combined with audio influence, and common amongst all cultures and ages, dance has great quantifiable effect on motor sequence learning and neuroplasticity (Gao et al., 2025; Jeyarajan et al., 2024; Johansson et al., 2022; Tsai et al., 2024). Dance actively engages an individual physically, cognitively, and sensorially, and it has been widely recognized as a practical intervention to promote neuroplasticity and execute motor sequence learning (Barnstaple et al., 2021; Cheng et al., 2024; Dhami et al., 2015; Du et al., 2025; Simon et al., 2024; Volpe et al., 2025). While a vast amount of literature involving

measures of neuroplasticity within the last decade suggests physical exercise or dance can lead to improvements or mitigation of impairments, the need for both simplified and accessible tasks is paramount. The physical demands of exercise and other coordinated movement such as dance continue to pose the challenge of being inaccessible to various groups. It has been widely urged within the surrounding literature to create simplified versions of both dance intervention (Dhami et al., 2015; Du et al., 2025), and physical exercise intervention (Johansson et al., 2022). For old-aged, neurodegenerative disordered, or disabled groups, severity of cognitive or physical decline is a key issue (Gorges et al., 2016; Hill-Briggs et al., 2007), and it is acknowledged that interventions should come to address individuals beyond the mild-moderate scale.

The aim with the present eye tracking task is to develop a feasible, non-exertive task that minimizes the physical response needed from a participant whilst still activating similar regions of the brain to that of coordinated physical movement and auditory cues such as dance. Within the current study, the usage of eye movement as an alternative to physical movement in motor sequence learning was investigated. In combination with auditory cues of a musical piece, the additional goal of diversifying current literature on multisensory sequence learning was sought. We hypothesized that a within-subjects design with 5 sessions of eye dance practice over 5 weeks of time would result in improvements regarding sequence-specific learning, namely performance accuracy and timing precision.

Methods

Participants:

Ten healthy adults were recruited for the study all of which from the population of York University students and provided written consent. Exclusion criteria for the study included visual or auditory impairment. Total n = 10 (female = 8, male = 2), Ages: 20-25, M = 22.

Procedure:

Participants were taught a complex eye movement sequence to a 68 second musical piece by watching a choreographed eye-dance pattern. The video and music of the dance were played five times with thirty second break intervals featuring a gray interface with no audio in between. The set of prior events is referred to as the learning phase of the test. Following the learning phase, participants are tasked to perform the same sequence of eye movements using only the musical piece and target markers on a gray interface. Executions of the dance sequence were recorded five times with a thirty second break with a gray interface similar to the learning phase, and are referred to as the performance phase. At the end of each 68 second performance recording, 5 out of the 30 seconds within the break consisted of a colour flash of the interface corresponding to accuracy of the steps correctly made. Three colours were shown: red, yellow, and green; corresponding to total score being <33% correct, 33%< x <66% correct, and >66% respectively. The visual reinforcer was shown to cue accuracy of the pattern recorded to each participant. Participants were also asked to rate their performance after each trial on a 1-100% scale. After learning and performance phases the session was completed and participants were instructed to complete their next session within 7-10 days. All eye-movements were directionally cued along the X and Y axes recorded within an eye-tracking paradigm on an Eyelink 1000 plus (EYELINK I CL v5.04 Sep 25 2014 CAMERA: Eyelink GL Version 1.2 Sensor=AJ7).

Results

Data plotting and analysis were conducted using softwares R 4.3.1 and Matlab(version). Results of accuracy statistics were found using a repeated-measures ANOVA. Eye gaze data was sampled every 2ms, all blinks and positions off-screen were omitted from data processing. Target thresholds were located within 200 pixel squares around exact targets (Fig. 3). Movements were only marked if 200 samples remained within target threshold, then compared with the dance sequence to be matched. A repeated-measures ANOVA revealed a significant main effect of sessions on Performance Accuracy, $F(4, 36) = 6.99$, $p < 0.001$, an effect size of $\eta^2_G = 0.26$, and Mauchly's test confirmed that sphericity was met ($p = 0.32$). An additional repeated-measures ANOVA revealed a significant main effect of sessions on Timing Precision, $F(4,36) = 11.67$ $p < 0.001$, an effect size of $\eta^2_G = 0.25$. Mauchly's test indicated sphericity was violated ($p = 0.022$); the Greenhouse-Geisser correction confirmed the effect remained significant ($p = 0.0086$).

Post-hoc pairwise comparisons using Bonferroni correction showed that Session 5 performance accuracy was significantly higher than Session 1 ($p = 0.008$). No other session pairs differed significantly, though the difference in first and last sessions suggests a gradual increase in performance each session leading to significant improvement.

Post-hoc pairwise comparisons using Bonferroni correction showed that timing accuracy of movements improved significantly between early and later sessions, with significant differences observed between sessions 1 and 4 ($p = 0.044$), sessions 1 and 5 ($p = 0.0038$), sessions 2 and 4 ($p = 0.0286$), sessions 2 and 5 ($p = 0.0055$) and sessions 3 and 5 ($p = 0.0055$). No significant

differences were found between later sessions (g4 & g5, p = 1.00), suggesting that timing

accuracy plateaued prior to the end of training.

Fig 1. Participant Performance Accuracy Across Sessions. Accuracy was recorded based on the proportion of movements correctly executed to the dance sequence (X/46). A line of best fit and shaded 95% confidence interval is shown as well.

Fig 2. Participant Timing Precision Across Sessions. Participant timing was measured based on proximity to the movements synced to video runtime.. A line of best fit and shaded 95% confidence interval is shown as well.

Fig 3. XY Value positions of Training Performance. Exact sequence positions shown in pixels of expected (A) X values and (B) Y values with video/audio timestamp.

Fig 4. Participant Trial Position and Timing of Actual Performance. Exact sequence positions shown in pixels of individual participant performance in session 5. (A) X values and (B) Y values compared from training position to actual performance with video/audio timestamp.

Discussion

  This study investigated the potential of motor sequence learning explicitly using oculomotor function and auditory cues. This novel visual-motor sequence learning task demonstrated significant improvements in both performance accuracy and timing precision, suggesting procedural learning and cognitive-motor coordination can be enhanced through gaze-based repetition. The statistical analyses show a well defined learning curve, proving the efficacy of the task in consolidation of learning following practice. Further analysis of the timing precision can be used to suggest audio cues played a significant role in learning of the sequence. Participants also tended to lag behind audio cue timestamps, in line with findings from Han et al. (2024) and Leow et al. (2025), that suggested auditory stimuli influence motor sequence learning preceding oculomotor movement. Furthermore, it is suggested that audio cues played congruently with visual stimuli encourages stronger development of explicit knowledge within reaction tasks compared to solely visual tasks (Lagarrigue et al., 2021; Silva et al., 2017). The distinct roles of audio and visual stimuli towards influencing eye movement provides a strong foundation for future motor sequence learning studies to consist of neurodegenerative disease or tetraplegic populations, therefore expanding the field on severe motor deficit groups. By using this focused oculomotor approach, the task can potentially aid in designing non-invasive, low-burden rehabilitation tools for enhancing functional outcomes. This gives the opportunity, should interventional and rehabilitational use prove effective, that the task be simplified for frequent/daily use as well.

  Various structures of the brain associated with motor-sequence learning and furthermore, the associated oculomotor components of learning, are expected to be pertinent to the results within the current body of work. To elaborate, while motor sequence learning is multifaceted due

to various regions of the brain involved, the basal ganglia has been cited throughout prior literature as a core region associated with learning and rewards systems (Di Filippo et al., 2009; Joti et al., 2007). Another region of interest is the ventral tegmental area (VTA), also associated with reward systems (Laviolette & Van Der Kooy, 2001; Morales & Margolis, 2017). These defined areas that encompass the reward system driving learning mechanisms, can be fully traced using fMRI to account for increased functional connectivity by measuring cortical white matter (Hänggi et al., 2010; Tremblay et al., 2021). Furthermore, such increases are noted post-training after motor sequence learning tasks (Jäger et al., 2022), similar results would likely be evident should the present study be replicated with a fMRI component. To further elucidate the mechanisms behind motor sequence learning, neurotransmitter function and measurement must also be accounted for. Once more, neurotransmitter function of motor sequence learning is varied, but namely dopamine (Ferreira et al., 2025), and GABA (Ferreira et al., 2025, Hehl et al., 2025), are promoted by visual tasks, (Tsai et al., 2024), and physical activity (Gao et al., 2025; Petzinger et al., 2013). As elaborated prior, the oculomotor system of the brain has potential to mimic results of physical activity and prompt neurotransmitter signalling. Areas of the oculomotor circuit to include in further analysis would be the supplementary motor area (DeSouza et al 2003; Chan, Kucyi & DeSouza, 2015; Jäger et al., 2022; Simon et al., 2024), the frontal eye field (DeSouza et al 2003; Chan, Kucyi & DeSouza, 2015; Rubino et al., 2024), and the supplementary eye field (DeSouza et al 2003), as they are crucial in motor sequence learning (Gonzalez et al., 2016). This novel eye tracking task in future studies, should therefore incorporate a fMRI scanning element to explore this oculomotor task in having potential results comparable to that of other motor sequence learning tasks.

Throughout our dance for Parkinson's work many eye movements are made that are incorporated to gaze to the hand and/or body parts (Bearss et al., 2017). It is made clear that similar results from oculomotor influence can benefit those within neurodegenerative disease and/or those physically disabled. Various forms of exercise or dance have been identified previously within the body of work as promoting neuroplastic effects (Dhami et al., 2015). While limited motor movement of the body would likely pose differing results to typical eye-limb motor sequence tasks, the results found in a solely oculomotor task could expand knowledge of an area that lacks specific motor interventions. Parkinson's disease is an ideal group the novel oculomotor task can be targeted towards, as they have deficits that are directly engaged by this learning paradigm (Hamedani et al., 2020). The use of musical cues and reinforcement feedback may promote dopaminergic engagement related to reward circuitry very relevant for PD populations as well (Tsai et al., 2024; D'Allesandro & DeSouza 2025). While not tested within the present study, the exploration of dopaminergic reinforcement involving the score indicator would be of great interest for future studies. Furthermore, dopaminergic changes can modulate anxiety (Baik, 2020; Dong et al., 2020) which future applications of the task can aim to analyze in addition to reinforcement paradigms. Prior studies have discovered physical dance to be cognitively engaging and to promote neuroplastic changes or mitigate future degradation of neural pathways (Barnstaple et al., 2021; Bar & DeSouza, 2016; Cheng et al., 2024; Simon et al., 2024; Quan et al., 2025), thus further supporting the demand for simplified movement related tasks for severe disease cases. Longitudinal studies have urged, (Wong et al., 2018), as understanding mitigative effects of interventions are best imparted when repeated throughout illness progression. Through the implementation of reinforcement paradigms within the task, and

engaging areas in which clear deficits are known, there could be potential in promoting greater response or mitigation in neurodegenerative groups such as those with PD.

Within the present study there were several limitations observed. The small sample size of ten participants, while significant, could be increased to get a broader scope of the learning curve and how findings may differ. A participant expressed an issue with the repetitive nature of the task causing dryness in the eyes, which is to be made note of for future studies involved with eye movement tracking. To understand the full extent of our findings would require fMRI analysis of regions of interest involved in motor sequence learning and furthermore, other pathways exemplifying neuroplastic changes. Additionally, EEG is another measurement tool that could be considered in motor learning applications (Titone et al., 2022). Extra sessions or an extended follow-up session period up to 1 month would be beneficial in a future eye task study to ensure motor sequence learning consolidation and forgetting. A lack of implicit learning is noted, as the explicit learning of the repeated sequence was the central focus of our study. To properly measure implicit learning it is advised to follow learning of the eye dance sequence with a novel sequence. The measurement of saccades is necessary to include for implicit learning (Jiang et al., 2014; Medimorec et al., 2021; Rubino et al., 2025), to assess whether saccadic movements after a novel or random sequence is greater after post sequence training.

These preliminary findings within the current body of work suggest that this novel eye dance task is feasible and sufficient in participant learning of a specific sequence, therefore it can be further investigated in several ways. The simplicity and low physical strain on participants are key components of the eye tracking dance task design, and should prove to be cost-effective in training and execution. A psychomotor vigilance task used in the study of passive and active exercise effects on mental fatigue from Jeyarajan et al. (2024), is a task that can be modified

under the context of the novel eye tracking task's results. Its design uses randomized targets appearing on a screen and gauges reaction time measurement by clicking targets, which can be replaced with eye movements and blinks to supplement lack of limb use for future studies. Through the application of our eye tracking test, a future study involving this task could be to account for those with disability or limited limb movement could prove beneficial in promoting neuroplastic effects. Additionally, this would allow for a comparison of other learning paradigms and tasks within the relevant literature to reinforce similarities in sequence learning and elaborate differences regarding the novel task. In doing so first with healthy participants, a clear definition of how such connections develop and adapt over time can be used to compare with a population of those with neurodegenerative diseases.

Author Contributions

Petrovski, M. - developed task environment, collected participant data, developed code, performed analysis, and first draft of manuscript.

Beheiry, S. - collected participant data, edited manuscript.

Das, U. U. - created study design and initial task environment.

Muhammad, M. - created study design and initial task environment.

Simon, R. - conceptualization, funding acquisition

Rooprai, S.- conceptualization, initial pilot testing and study design

Karimi, A. - conceptualization, initial pilot testing and study design, funding acquisition

Bar, R. - conceptualization, funding acquisition

DeSouza, J. F. X - conceptualization, initial pilot testing, contributed testing and analysis tools, edited manuscript, and finalized manuscript, funding acquisition.

Ethics Statement

This study was approved by the Office of Research Ethics Committee (ORE) of York University Certificate #2025-032.

Conflict of Interest

The authors declare that the research was conducted in the absence of any commercial or financial influence that could be a potential conflict of interest.

Acknowledgements

The authors would like to thank all the participants that committed their time over the course of several weeks to this study, whose dedication made this possible. We also thank all members of the joeLAB. This research was supported by the Anxiety Research Fund powered by Beneva awarded to York University Faculty of Health.

References

Baik, J.-H. (2020). Stress and the dopaminergic reward system. *Experimental & Molecular Medicine*, *52*(12), 1879–1890. https://doi.org/10.1038/s12276-020-00532-4

Bar, R. J., & DeSouza, J. F. X. (2016). Tracking plasticity: Effects of long-term rehearsal in expert dancers encoding music to Movement. *PLOS ONE*, *11*(1). https://doi.org/10.1371/journal.pone.0147731

Barnstaple, R., Protzak, J., DeSouza, J. F. X., & Gramann, K. (2021). Mobile Brain/Body Imaging in dance: A dynamic transdisciplinary field for applied research. *European Journal of Neuroscience*, *54*(12), 8355–8363. https://doi.org/10.1111/ejn.14866

Bearss, K. A., Barnstaple, R. E., Bar, R. J., & DeSouza, J. F. X. (2024). Impact of Weekly Community-Based Dance Training Over 8 Months on Depression and Blood Oxygen Level–Dependent Signals in the Subcallosal Cingulate Gyrus for People With Parkinson Disease: Observational Study. *JMIRx Med*, *5*(1), e44426. https://doi.org/10.2196/44426

Bearss, K. A., Barnstaple, R. E., Bar, R. J., & DeSouza, J. F. X. (2024). Impact of Weekly Community-Based Dance Training Over 8 Months on Depression and Blood Oxygen Level–Dependent Signals in the Subcallosal Cingulate Gyrus for People With Parkinson Disease: Observational Study. *JMIRx Med*, *5*(1), e44426. https://doi.org/10.2196/44426

Bearss, K. A., McDonald, K. C., Bar, R. J., & DeSouza, J. F. X. (2017). Improvements in balance and gait speed after a 12 week dance intervention for Parkinson's disease. *Advances in Integrative Medicine*, *4*(1), 10–13. https://doi.org/10.1016/j.aimed.2017.02.002

Bearss, K. A., McDonald, K. C., Bar, R. J., & DeSouza, J. F. X. (2017). Improvements in balance

    and gait speed after a 12 week dance intervention for Parkinson's disease. *Advances in*

    Integrative Medicine, *4*(1), 10–13. https://doi.org/10.1016/j.aimed.2017.02.002

Bonzano, L., Palmaro, E., Teodorescu, R., Fleysher, L., Inglese, M., & Bove, M. (2015).

    Functional connectivity in the resting-state motor networks influences the kinematic

    processes during motor sequence learning. *European Journal of Neuroscience*, *41*(2),

    243–253. https://doi.org/10.1111/ejn.12755

Chan J.L., Kucyi A. and DeSouza J.F.X. (2015) Oculomotor System. In: Arthur W. Toga, editor.

    Brain Mapping: An Encyclopedic Reference, vol. 2, pp. 483- 488. Academic Press:

    Elsevier. https://doi.org/10.1016/B978-0-12-397025-1.00031-2

Cheng, W.-H., Quan, Y., & Thompson, W. F. (2024). The effect of dance on mental health and

    quality of life of people with Parkinson's disease: A systematic review and three-level

    meta-analysis. *Archives of Gerontology and Geriatrics*, *120*, 105326.

    https://doi.org/10.1016/j.archger.2024.105326

Coiner, B., Pan, H., Bennett, M. L., Bodien, Y. G., Iyer, S., O'Neil-Pirozzi, T. M., Leung, L.,

    Giacino, J. T., & Stern, E. (2019). Functional neuroanatomy of the Human Eye

    Movement Network: A review and Atlas. *Brain Structure and Function*, *224*(8),

    2603–2617. https://doi.org/10.1007/s00429-019-01932-7

Culicetto, L., Cardile, D., Marafioti, G., Lo Buono, V., Ferraioli, F., Massimino, S., Di Lorenzo,

    G., Sorbera, C., Brigandì, A., Vicario, C. M., Quartarone, A., & Marino, S. (2025).

    Recent advances (2022–2024) in eye-tracking for parkinson's disease: A promising tool

for diagnosing and monitoring symptoms. *Frontiers in Aging Neuroscience*, 17.

https://doi.org/10.3389/fnagi.2025.1534073

DeSouza, J. F. X., Menon, R. S., & Everling, S. (2003). Preparatory Set Associated With

Pro-Saccades and Anti-Saccades in Humans Investigated With Event-Related fMRI.

Journal of Neurophysiology. https://doi.org/10.1152/jn.00562.2002

Dhami, P., Moreno, S., & DeSouza, J. F. X. (2015). New framework for rehabilitation – fusion of

cognitive and physical rehabilitation: The hope for dancing. *Frontiers in Psychology*, 5.

https://doi.org/10.3389/fpsyg.2014.01478

Di Filippo, M., Picconi, B., Tantucci, M., Ghiglieri, V., Bagetta, V., Sgobio, C., Tozzi, A.,

Parnetti, L., & Calabresi, P. (2009). Short-term and long-term plasticity at corticostriatal

synapses: Implications for learning and memory. *Behavioural Brain Research*, *199*(1),

108–118. https://doi.org/10.1016/j.bbr.2008.09.025

Diotaiuti, P., Marotta, G., Di Siena, F., Vitiello, S., Di Prinzio, F., Rodio, A., Di Libero, T.,

Falese, L., & Mancone, S. (2025). Eye tracking in parkinson's disease: A review of

oculomotor markers and clinical applications. *Brain Sciences*, *15*(4), 362.

https://doi.org/10.3390/brainsci15040362

Dong, M.-X., Chen, G.-H., & Hu, L. (2020). Dopaminergic System Alteration in Anxiety and

Compulsive Disorders: A Systematic Review of Neuroimaging Studies. *Frontiers in*

Neuroscience, *14*. https://doi.org/10.3389/fnins.2020.608520

Du, M., Hancox, J. E., Hooper, O., Sandford, R., & Huang, C. (2025). Dancing towards

wellbeing: A scoping review of dance interventions for therapeutic purposes in

educational settings. *International Review of Sport and Exercise Psychology*, *0*(0), 1–37. https://doi.org/10.1080/1750984X.2025.2471759

D'Alessandro, E. M., Karimi, A., Bearss, K. A., Bar, R. J., & DeSouza, J. F. X. (2025). Reward network modulation while learning movements to music during weekly multisensory training program for people with Parkinson's disease: Mood effects over 7-yrs (p. 2025.07.19.25331836). *medRxiv*. https://doi.org/10.1101/2025.07.19.25331836

Ferreira, M., Matias, F., Abrunhosa, A., Martins, R., & Castelo-Branco, M. (2025). Evidence for neuroplasticity in the human brain in health and disease: A systematic review focusing on molecular imaging using PET. *Neural Plasticity*, *2025*(1). https://doi.org/10.1155/np/9423232

Gao, B., Wang, L., Gong, J., Zhu, Z., Liu, Q., Yuan, H., & Wang, H. (2025). The interplay between physical exercise and autophagy signaling in brain health, neurodegenerative diseases and aging. *Frontiers in Aging Neuroscience*, *17*. https://doi.org/10.3389/fnagi.2025.1579208

Gonzalez, C. C., Billington, J., & Burke, M. R. (2016). The involvement of the fronto-parietal brain network in oculomotor sequence learning using fMRI. *Neuropsychologia*, *87*, 1–11. https://doi.org/10.1016/j.neuropsychologia.2016.04.021

Gorges, M., Müller, H.-P., Lulé, D., Pinkhardt, E. H., Ludolph, A. C., Kassubek, J., & LANDSCAPE Consortium. (2016). The association between alterations of eye movement control and cerebral intrinsic functional connectivity in Parkinson's disease. *Brain Imaging and Behavior*, *10*(1), 79–91. https://doi.org/10.1007/s11682-015-9367-7

Hamedani, A. G., Abraham, D. S., Maguire, M. G., & Willis, A. W. (2020). Visual Impairment Is

　　More Common in Parkinson's Disease and Is a Risk Factor for Poor Health Outcomes.

　　Movement Disorders, *35*(9), 1542–1549. https://doi.org/10.1002/mds.28182

Han, Z., Sanchez, D., Levitan, C. A., & Sherman, A. (2024). Stimulus-locked auditory

　　information facilitates real-time visuo-motor sequence learning. *Psychonomic Bulletin &*

　　Review, *31*(2), 828–838. https://doi.org/10.3758/s13423-023-02378-z

Hänggi, J., Koeneke, S., Bezzola, L., & Jäncke, L. (2010). Structural neuroplasticity in the

　　sensorimotor network of professional female ballet dancers. *Human Brain Mapping*,

　　31(8), 1196–1206. https://doi.org/10.1002/hbm.20928

Hehl, M., Malderen, S. V., Blashchuk, S., Sunaert, S., Edden, R. A. E., Swinnen, S. P., &

　　Cuypers, K. (2025). The Reciprocal Relationship Between Short‑ and Long‑Term Motor

　　Learning and Neurometabolites. *Human Brain Mapping*, *46*(4), n/a-n/a.

　　https://doi.org/10.1002/hbm.70170

Herzberg, M. P., Nielsen, A. N., Luby, J., & Sylvester, C. M. (2024). Measuring neuroplasticity

　　in human development: The potential to inform the type and timing of mental health

　　interventions. *Neuropsychopharmacology*, *50*(1), 124–136.

　　https://doi.org/10.1038/s41386-024-01947-7

Hill-briggs, F., Dial, J., Morere, D., & Joyce, A. (2007). Neuropsychological assessment of

　　persons with Physical Disability, visual impairment or blindness, and hearing impairment

　　or deafness. *Archives of Clinical Neuropsychology*, *22*(3), 389–404.

　　https://doi.org/10.1016/j.acn.2007.01.013

Jäger, A.-T. P., Huntenburg, J. M., Tremblay, S. A., Schneider, U., Grahl, S., Huck, J., Tardif, C. L., Villringer, A., Gauthier, C. J., Bazin, P.-L., & Steele, C. J. (2022). Motor sequences; separating the sequence from the motor. A longitudinal rsfMRI study. *Brain Structure and Function*, *227*(3), 793–807. https://doi.org/10.1007/s00429-021-02412-7

Jeyarajan, G., Buwadi, L., Ayaz, A., Nagamatsu, L. S., Haile, D., Zou, L., & Heath, M. (2024). Passive and active exercise do not mitigate mental fatigue during a sustained vigilance task. *Experimental Brain Research*, *243*(1), 19. https://doi.org/10.1007/s00221-024-06950-4

Jiang, Y. V., Won, B.-Y., & Swallow, K. M. (2014). First saccadic eye movement reveals persistent attentional guidance by implicit learning. *Journal of Experimental Psychology. Human Perception and Performance*, *40*(3), 1161–1173. https://doi.org/10.1037/a0035961

Johansson, M. E., Cameron, I. G. M., Van der Kolk, N. M., de Vries, N. M., Klimars, E., Toni, I., Bloem, B. R., & Helmich, R. C. (2022). Aerobic Exercise Alters Brain Function and Structure in Parkinson's Disease: A Randomized Controlled Trial. *Annals of Neurology*, *91*(2), 203–216. https://doi.org/10.1002/ana.26291

Joti, P., Kulashekhar, S., Behari, M., & Murthy, A. (2007). Impaired inhibitory oculomotor control in patients with Parkinson's disease. *Experimental Brain Research*, *177*(4), 447–457. https://doi.org/10.1007/s00221-006-0687-0

Kurkin, S., Akao, T., Fukushima, J., Shichinohe, N., Kaneko, C. R. S., Belton, T., & Fukushima, K. (2014). No-go neurons in the cerebellar oculomotor vermis and caudal fastigial nuclei: Planning tracking eye movements. *Experimental Brain Research*, *232*(1), 191–211. https://doi.org/10.1007/s00221-013-3731-x

Lagarrigue, Y., Cappe, C., & Tallet, J. (2021). Regular rhythmic and audio-visual stimulations enhance procedural learning of a perceptual-motor sequence in healthy adults: A pilot study. *PLOS ONE*, *16*(11), e0259081. https://doi.org/10.1371/journal.pone.0259081

Laviolette, S. R., & Van Der Kooy, D. (2001). GABAA receptors in the ventral tegmental area control bidirectional reward signalling between dopaminergic and non-dopaminergic neural motivational systems. *The European Journal of Neuroscience*, *13*(5), 1009–1015. https://doi.org/10.1046/j.1460-9568.2001.01458.x

Leow, L., Nguyen, A., Corti, E., & Marinovic, W. (2025). Informative auditory cues enhance motor sequence learning. *European Journal of Neuroscience*, *61*(10), 1–10. https://doi.org/10.1111/ejn.70140

Maceira-Elvira, P., Timmermann, J. E., Popa, T., Schmid, A.-C., Krakauer, J. W., Morishita, T., Wessel, M. J., & Hummel, F. C. (2022). Dissecting motor skill acquisition: Spatial coordinates take precedence. *Science Advances*, *8*(29), eabo3505. https://doi.org/10.1126/sciadv.abo3505

Medimorec, S., Milin, P., & Divjak, D. (2021). Inhibition of eye movements disrupts spatial sequence learning. *Experimental Psychology*, *68*(4), 221–228. https://doi.org/10.1027/1618-3169/a000528

Morales, M., & Margolis, E. B. (2017). Ventral tegmental area: cellular heterogeneity, connectivity and behaviour. *Nature Reviews. Neuroscience*, *18*(2), 73–85. https://doi.org/10.1038/nrn.2016.165

Petzinger, G. M., Fisher, B. E., McEwen, S., Beeler, J. A., Walsh, J. P., & Jakowec, M. W. (2013). Exercise-enhanced neuroplasticity targeting motor and cognitive circuitry in Parkinson's disease. *The Lancet Neurology*, *12*(7), 716–726. https://doi.org/10.1016/S1474-4422(13)70123-6

Quan, Y., Lo, C. Y., Olsen, K. N., & Thompson, W. F. (2025). The effectiveness of aerobic exercise and dance interventions on cognitive function in adults with mild cognitive impairment: An overview of meta-analyses. *International Review of Sport and Exercise Psychology*, *18*(2), 1026–1047. https://doi.org/10.1080/1750984X.2024.2332989

Rubino, C., Andrushko, J. W., Rinat, S., Harrison, A. T., & Boyd, L. A. (2024). Oculomotor functional connectivity associated with Motor Sequence Learning. *Cerebral Cortex*, *34*(11). https://doi.org/10.1093/cercor/bhae434

Rubino, C., Harrison, A. T., & Boyd, L. A. (2025). Oculomotor learning is evident during implicit motor sequence learning. *Scientific Reports*, *15*(1), 9498. https://doi.org/10.1038/s41598-025-93498-0

Sailer, U., Flanagan, J. R., & Johansson, R. S. (2005). Eye–Hand Coordination during Learning of a Novel Visuomotor Task. *Journal of Neuroscience*, *25*(39), 8833–8842. https://doi.org/10.1523/JNEUROSCI.2658-05.2005

Silva, A. E., Barakat, B. K., Jimenez, L. O., & Shams, L. (2017). *Multisensory Congruency Enhances Explicit Awareness in a Sequence Learning Task*. https://doi.org/10.1163/22134808-00002587

Simon, J. R., Bek, J., Ghanai, K., Bearss, K. A., Barnstaple, R. E., Bar, R. J., & DeSouza, J. F. (2024). Neural effects of multisensory dance training in parkinson's disease: Evidence

from a longitudinal neuroimaging single case study. *Frontiers in Aging Neuroscience*, *16*.

https://doi.org/10.3389/fnagi.2024.1398871

Soetedjo, R., Fuchs, A. F., & Kojima, Y. (2009). Subthreshold Activation of the Superior

Colliculus Drives Saccade Motor Learning. *Journal of Neuroscience*, *29*(48),

15213–15222. https://doi.org/10.1523/JNEUROSCI.4296-09.2009

Titone, S., Samogin, J., Peigneux, P., Swinnen, S., Mantini, D., & Albouy, G. (2022).

Connectivity in Large-Scale Resting-State Brain Networks Is Related to Motor Learning:

A High-Density EEG Study. *Brain Sciences*, *12*(5), 530.

https://doi.org/10.3390/brainsci12050530

Tremblay, S. A., Jäger, A.-T., Huck, J., Giacosa, C., Beram, S., Schneider, U., Grahl, S.,

Villringer, A., Tardif, C. L., Bazin, P.-L., Steele, C. J., & Gauthier, C. J. (2021). White

matter microstructural changes in short-term learning of a continuous visuomotor

sequence. *Brain Structure and Function*, *226*(6), 1677–1698.

https://doi.org/10.1007/s00429-021-02267-y

Tsai, C.-L., Pan, C.-Y., Wang, T.-C., Tseng, Y.-T., Ukropec, J., Ukropcová, B., & Lin, T.-K.

(2024). Effects of acute aerobic exercise with different intensities on cerebral

dopamine/norepinephine/serotonin metabolites and executive-related oculomotor control

in individuals with Parkinson's disease. *Mental Health and Physical Activity*, *26*, 100582.

https://doi.org/10.1016/j.mhpa.2024.100582

Vieluf, S., Massing, M., Blandin, Y., Leinen, P., & Panzer, S. (2015). The role of Eye Movements

in Motor Sequence Learning. *Human Movement Science*, *40*, 220–236.

https://doi.org/10.1016/j.humov.2015.01.004

Volpe, D., Baldassarre, M. G., Bakdounes, L., Campo, M. C., Ferrazzoli, D., & Ortelli, P. (2025).

"Dance Well"—A Multisensory Artistic Dance Intervention for People with Parkinson's

Disease: A Pilot Study. *Brain Sciences*, *15*(4), 357.

https://doi.org/10.3390/brainsci15040357

Wong, O. W., Chan, A. Y., Wong, A., Lau, C. K., Yeung, J. H., Mok, V. C., Lam, L. C., & Chan,

S. (2018). Eye movement parameters and cognitive functions in Parkinson's disease

patients without dementia. *Parkinsonism & Related Disorders*, *52*, 43–48.

https://doi.org/10.1016/j.parkreldis.2018.03.013