Title

Ontogeny of vocalizations in Adlie penguin (*Pygoscelis adeliae*) chicks.

Authors

Michele L. Adams[1,3], Danielle T. Fradet[1,3], Megan A. Cimino[2], Easton R. White[1], Laura N.

Kloepper[1,3]

Affiliations

Department of Biological Sciences, University of New Hampshire, Durham, NH.[1]

Department of Ocean Sciences, University of California, Santa Cruz, CA.[2]

Center for Acoustics Education and Research, University of New Hampshire, Durham, NH.[3]

Title

Ontogeny of vocalizations in Adélie penguin (*Pygoscelis adeliae*) chicks.

Abstract

Passive acoustic monitoring (PAM) is an efficient method to monitor dense aggregations

of vocal animals but requires understanding the acoustic ecology of the species under

examination. Avian vocal development is largely understood from songbirds, limiting its

application to non-passerine taxa with different social and environmental pressures. As an

example, colonial seabirds such as the Adélie penguin (*Pygoscelis adeliae*) inhabit acoustically

crowded environments and rely on vocal cues, in addition to spatial information, for parent-

offspring recognition. While adult penguin vocal communication is well studied, chick vocal

development remains poorly characterized. Using the deep learning-based system DeepSqueak,

we aimed to characterize the vocal development of wild *P. adeliae* chicks in the West Antarctic

Peninsula. We found that acoustic features of chick calls changed systematically with age, with

calls becoming longer and more frequency modulated over time. Characterizing chick vocal

development from hatch to fledge provides important information to study phenological

communication patterns in vocal-dependent seabirds and supports the application of PAM to

assess climate-driven impacts on indicator species.

Keywords

Introduction

Vocal learning pertains to the capacity of specific mammal and bird species to learn and adjust their vocalizations through imitation and social interaction (Brenowitz & Beecher, 2005; Elie & Theunissen, 2020). Animal communication often relies on various contextual cues to recognize kinship, especially in large groups (Beecher, 1982). Among bird species, oscine songbirds (Passeriformes) are renowned for their vocal learning abilities and have been the primary focus of research on avian communication, contributing significantly to our understanding of neuroanatomical motor control of learned songs (Vicario et al., 2002) and revealing a critical developmental threshold for vocal learning in most oscine songbirds (Tramontin & Brenowitz, 1999). Vocal non-learners, such as seabirds, present a challenge in understanding avian communication because it is believed they rely on innate vocalizations rather than learning their vocalizations through mimicry and practice (Jones & Ryan, 2010). Vocal non-learning has been studied less extensively than vocal learning in birds, and it remains unclear whether vocal non-learners exhibit thresholds for adult vocalizations.

Research on the development of vocalizations in colonial breeding seabirds has primarily focused on how these vocalizations are used in social and environmental contexts (Aubin et al, 2000; Aubin & Jouventin, 2002; Beecher, 1982; Clark et al, 2006; Favaro et al., 2015; Favaro et al., 2016; Jouventin & Aubin, 2002; Kidawa et al, 2023; Lynch & Lynch, 2017; Marks et al, 2010; Robisson et al, 1993; Sladen, 1958; Spurr, 1975; Terranova et al, 2023). Such research has shown how vocalizations are essential for mating, territory defense, and offspring recognition (Jouventin & Aubin, 2002). One study suggests that changes in pitch in the vocalizations of Little Blue Penguins (*Eudyptula minor*) were variable between individuals and could act as an identity marker (Nakagawa et al., 2001). However, limited research exists on the vocal

development of penguin chicks before they fledge, making it challenging to understand how

vocalizations emerge and evolve in the early life stages of colonial seabirds. Here, we aim to test

if specific acoustic features in non-passerine chicks change predictably with age as part of

developmental growth.

Penguins are considered vocal non-learners, demonstrating limited vocal learning in

specific social contexts. For example, breeding adult penguins perform vocal adjustments in

pitch, duration, syllables, and rhythm to communicate identity (Robsion et al., 1993; Jouventin &

Aubin, 2002). *P. adeliae* perform a loud mutual display call during the breeding season that

likely evolved for the purpose of mutual recognition between individuals within loud, congested

breeding colonies (Aubin & Jouventin, 2002). Chicks perform a begging peep from the day they

hatch, and they have been known to perform immature versions of the loud mutual display call

(Davis, 1982; Spurr, 1975). Acoustic modulation in these calls facilitates mate and parent-chick

recognition and ensures parental investment is given to the genetic offspring within the colony

(Speirs et al., 1991). Mutual vocal recognition is particularly crucial for species with high

parental investment and a risk of offspring misidentification (Kidawa et al., 2023). However, the

stage at which chicks begin to exhibit vocal modulation patterns or if they participate in vocal

learning during brooding remains unclear. Addressing these gaps could improve our

understanding of seabird communication and its ecological and evolutionary implications and

provide a measure to track potential phenological changes due to climate change in polar

regions.

Chick Age Classes

P. adeliae chicks are semi-altricial and require significant parental care. The initial phase,

known as the "guard stage," spans from 0 to 20 days of age (Sladen, 1958). During the first 10 days, chicks are unable to thermoregulate and are immobile, making them highly susceptible to predation (Spurr, 1975). In this stage, both parents alternate between guarding the nest and foraging at sea, with one adult always present to tend to the chicks (Spurr, 1975). On average, the guard stage lasts until the chicks are 20 days old, at which point they become homeothermic, allowing both parents to shift their focus primarily to foraging. Following the guard stage, chicks enter the "crèche stage," which occurs from approximately 21 to 41 days of age, and is marked by rapid growth (Taylor & Roberts, 1962). During this period, chicks become more mobile and begin to form crèches, aggregations that provide warmth and protection from predators while the parents are at sea (Davis, 1982; Spurr, 1975; Taylor & Roberts, 1962). Selective pressures for individual recognition may be driven by this behavior, enabling parents to identify and retrieve their offspring from the crèche (Kidawa et al., 2023). In this study, the final developmental stage before the chicks fledge is referred to as the "post-crèche" stage. During this stage, chicks exhibit increased independence as they prepare to leave the colony and live at sea. Large groups of chicks gather on beaches to fledge at approximately 54 days post-hatch (Cimino et al, 2014; Salihoglu et al, 2001).

This study explores the vocal development of *P. adeliae* chicks using Passive Acoustic Monitoring (PAM) combined with machine learning. Vocalizations were recorded across different chick developmental stages to examine how these calls change over time. PAM is a minimally invasive, cost-effective data collection method that involves the use of unsupervised, autonomous acoustic sensors to continuously record sounds in challenging or sensitive environments. This method is particularly valuable for long-term monitoring of wildlife distribution and behavior, especially in colonial, highly vocal species where breeding grounds

are costly and invasive to access by traditional monitoring methods. However, PAM generates

large volumes of data, making manual annotation time-consuming and inefficient. To address the

challenge of data volume, machine learning was introduced for efficient processing. Machine

learning uses computational models to learn representations of natural sight, vision, and hearing

for detection and classification (LeCun et al., 2015). Machine learning, particularly deep

learning, offers powerful tools to automate the detection and classification of vocalizations. By

leveraging computational models that can learn complex patterns in sound, machine learning

significantly reduces the time and effort required to analyze acoustic data, making PAM more

efficient and scalable for studying wildlife behavior over extended periods.

Materials and Methods

Five Adélie penguin colonies of various sizes were continuously recorded during the

austral summer from November 2022 – February 2023 near Palmer Station in the West Antarctic

Peninsula using Wildlife Acoustics Song Meter Minis (SMMs, Wildlife Acoustics, Maynard,

MA, USA). Two colonies were located on Humble Island, and three were located on Torgerson

Island. Penguin nests remained in the same location throughout this period, and nests were

monitored daily by visual inspection of 30 nests per island.

Acoustic Recorder Deployment and Programming

Five SMMs were deployed on recording rigs adjacent to *P. adeliae* colonies. Each rig

consisted of a PVC structure with a recorder zip-tied at a height of 1 m to keep it out of reach of

the penguins and positioned within 1 m of the colony edge (Fig. 1). At Humble Island, two rigs

were placed near the colonies due to the size of the colony, while at Torgersen Island, a single rig

was positioned near the approximate center of the three colonies, yet still at the edge of the

colonies. Rocks were placed at the base of each rig to provide additional stability. The SMMs

recorded five-minute audio clips at the start of every hour, sampled at 24 kHz with 16-bit depth

and a 6 dB gain. Memory cards and batteries were replaced monthly. Recordings were collected

during the *P. adeliae* breeding season, from 27 November 2022 to 27 February 2023, with

deployment occurring after the peak egg-laying date to minimize the risk of nest abandonment.

In total, 13,932 five-minute recordings were obtained, all wild *P. adeliae* vocalizations. The

original audio files were used for all subsequent analyses.

132

Figure 1: Photograph of recording rig deployed near the edge of an Adélie penguin colony with

the acoustic sensor at the top.

Data Processing

This study employs DeepSqueak, a deep learning-based system, to investigate the vocal

development of *P. adeliae* chicks. DeepSqueak is a call recognition software designed for non-

expert users to detect, measure, and classify mice and rat vocalizations (Coffey et al., 2019).

Because DeepSqueak's architecture uses spectrogram images, it is important to include different

call types into training categories. Prior to training and testing of DeepSqueak, call types were

differentiated through manual visual inspection of spectrograms, with particular attention given

to identifying chick calls and separating them from adult vocalizations. Calls were broken into

three categories based on spectrographic characteristics: begging peep (BEG), multi-syllable beg

(MSB), and loud mutual display (LMD). Early spectral BEG calls were identified as very short,

high-frequency vocalizations that appeared as small, discrete "carrot-shaped" elements above the

frequency range of adult calls. MSB calls were characterized by multi-syllable begging

vocalizations that lacked consistent contour structure and showed substantial variability across

syllables. In contrast, LMD calls were longer in duration and exhibited a more refined and

consistent contour, with repeated syllables forming a stable spectrotemporal pattern. These

visually distinct features guided the assignment of calls to training categories, and representative

spectrograms of each call type are shown in Fig. 2.

The training dataset consisted of 81 audio files containing *P. adeliae* chick vocalizations.

These included 43 BEG recordings, all from the Cornell Macaulay Library; 28 MSB recordings,

all obtained from the Macaulay Library; and 10 LMD recordings, of which 2 were sourced from

the Macaulay Library and 8 were sourced from our own recordings and excluded from the test

data set. Macaulay Library recordings spanned a wide temporal range, with collection dates of 3-

5 February 1988, 9 January 1989, 11 January 1989, 23 January 2010, 9 January 2014, 26 January

2022, 8 February 2024, and 5 January 2025. The test data set consisted of 1,848 five-minute

segments of our total original recordings, collected between 13 December 2022 and 27 February

2023 and filtered to exclude files where a) chicks were not visually confirmed to be present on

the islands and b) recordings where wind noise dominated the soundscape.

After we built our training dataset, we ran our detector on the test data set. Detections

were manually reviewed and edited to select the lowest harmonic band for each true positive

detection and false positives were labeled within DeepSqueak. For each detection DeepSqueak

measured multiple acoustic parameters; the parameters relevant to and reported in our study are

listed and described in Table 1. The audio data were divided into three sampling groups

distinguished by age class consistent with *P. adeliae* life stage and verified by visual

observations: guard stage (1), crèche stage (2), and post-crèche stage (3). The guard stage (weeks

1-3) included all recordings between 13 December 2022 and 02 January 2023. The crèche stage

included all recordings between 03 January 2023 and 16 February 2023 (weeks 4-9). The post-

crèche stage (week 10) group included all recordings between 17 February 2023 and 27 February

2023, but due to low numbers of chick calls and poor signal-to-noise ratio, we did not include

these files in our analysis.

Figure 2. Spectrographic examples of the progression from single syllable begs to multi-syllable

P. adeliae chick vocalizations highlight the structural changes over times; (a) beg (BEG), (b)

multi-syllable beg (MSB) and (c) immature loud mutual display (LMD).

Table 1. DeepSqueak acoustic parameters and definitions used in this study.

Parameter	Definition
Duration (s)	Total call time, End Time - Begin Time
High Frequency (kHz)	Highest frequency of the contour
Low Frequency (kHz)	Lowest frequency of the contour

Delta Frequency (kHz)	Change in frequency, High Freq - Low Freq; renamed to Bandwidth in our analysis
Sinuosity	Length of the path between the first and last points on the contour, divided by the Euclidean distance between the first and last points.

Statistical Analyses

To investigate temporal change in chick calls we first applied multinomial regression modeling to quantify developmental shifts in call-type probability across eight weeks. Second, we used a series of generalized additive models (GAMs) to examine how each acoustic feature (bandwidth, duration, high frequency, low frequency, peak frequency, and sinuosity) varied for week, call-type (BEG, MSB, and LMD calls) and the interaction between week and call-type.

To characterize population-level changes in acoustic features of *P. adeliae* chick calls across time, we analyzed all verified detected calls pooled across call types. For each acoustic parameter (bandwidth, duration, high frequency, low frequency, peak frequency, and sinuosity), we fit a generalized additive model (GAM) using week as a continuous predictor. GAMs were implemented using thin-plate regression splines with restricted maximum likelihood estimation to flexibly capture non-linear temporal trajectories while avoiding overfitting. To identify periods of abrupt change in the direction or rate of each trajectory, we applied segmented (piecewise linear) regression to the GAM-predicted values, using week as the segmentation variable. This two-stage approach allowed breakpoint detection to be driven by smooth population-level trends rather than by noise in individual observations. For each acoustic feature, the segmented regression yielded an estimated breakpoint (in weeks) and slopes before and after the breakpoint.

Results

Call Type Variation Across Developmental Week

DeepSqueak detected 2579 calls, all of which were manually verified resulting in 1636

true detections (1395 BEG; 111 MSB; 127 LMD). Over the eight weeks of sampling, the

composition of call types shifted noticeably throughout the phenological stages (Table 2). BEG

calls dominated early in the sampling period (guard stage), comprising nearly all vocalizations in

weeks 1 and 2 (>99%), while LMD and MSB calls were initially rare, with proportions below

1%. By week 3 (the transition from guard to crèche stage), the proportion of BEG calls decreased

to 58%, while MSB calls increased substantially to 31%, and LMD calls rose to 11%, indicating

the emergence of more complex or varied vocalizations. Interestingly, the proportion of BEG

calls rebounded in Week 4 to 91%, with LMD and MSB call types remaining low. Later weeks

demonstrated further heterogeneity. In week 5, MSB calls (58%) surpassed LMD (42%), while

BEG was absent. Weeks 6 and 7 showed BEG calls regaining dominance (71–87%), with LMD

and MSB present in smaller proportions. By week 8, BEG calls comprised 70%, and LMD 30%,

with MSB absent.

Overall, BEG calls were consistently the most frequent, but the data indicate a clear

temporal trend toward increasing contribution of LMD and MSB calls during the middle of the

observation period, followed by a slight return to BEG dominance in the final weeks. These

results were reflected in the predicted probabilities of each call from multinomial regression (Fig.

3).

Table 2. Proportions of call types from 24 December 2022 to 16 February 2023 shown as the total number of verified detections per call type per week, with the proportion of each call type relative to the total number of calls.

Week	BEG (n / prop)	LMD (n / prop)	MSB (n / prop)	Total
1	612 / 0.990	2 / 0.003	4 / 0.006	618
2	290 / 0.990	3 / 0.010	-	293
3	147 / 0.583	28 / 0.111	77 / 0.306	252
4	48 / 0.906	5 / 0.094	-	53
5	-	11 / 0.423	15 / 0.577	26
6	137 / 0.706	41 / 0.211	16 / 0.083	194
7	120 / 0.870	17 / 0.123	1 / 0.007	138
8	41 / 0.695	18 / 0.305	-	59

Figure 3. Predicted probabilities of chick call types over the brooding period. Predicted probabilities for each call type (BEG, LMD, MSB) are shown across weeks 1–8, illustrating the changing likelihood of each call type with age. Early development is dominated by BEG calls;

mid-development shows greater variability among call types, and later weeks return to higher

BEG probability.

Temporal Trends in Acoustic Features

Changes in acoustic features over time for each call type are shown in Fig. 4. Bandwidth

increased over time in BEG (0.268–0.361 kHz) and MSB calls (0.242–0.561 kHz), whereas

LMD calls peaked mid-season (0.498 kHz at week 4) before declining to 0.337 kHz by week 8.

Duration showed strong temporal increases in BEG (0.153–0.302 s) and MSB calls (0.220–0.552

s), while LMD calls exhibited higher overall durations early in development (0.802 s at week 1)

followed by a gradual decline toward week 8 (0.597 s). Consistent with high model fit (Table 3),

duration showed the strongest temporal structure across call types.

High-frequency components were relatively stable in BEG calls (3.88–4.33 kHz),

increased substantially in MSB calls (2.79–5.31 kHz), and peaked mid-season in LMD calls

(4.92 kHz at week 4) before declining. Low-frequency components decreased slightly in BEG

calls (3.56–2.90 kHz), increased gradually in LMD calls (2.29–2.76 kHz), and showed a mid-

season peak in MSB calls. Although these frequency-related features exhibited weaker overall

model fit, significant temporal effects were detected for selected call types (Table 3).

Peak frequency remained largely stable in BEG calls, while MSB and LMD calls exhibited mid-

season peaks. Sinuosity increased gradually over time in BEG calls (1.52–1.82) and MSB calls

(2.03–2.30), whereas LMD calls maintained consistently higher sinuosity values with little

temporal change. Consistent with model-fit results, significant nonlinear temporal change in

sinuosity was detected only for BEG calls, possibly indicating seasonal modulation of call

structure rather than differences in overall complexity.

Figure 4. Developmental trajectories of acoustic features in chick calls. GAM-derived fitted

values (± SE) are shown across weeks 1–8 on the x-axis for three call types (BEG, LMD, MSB)

for each acoustic feature: bandwidth, duration, high frequency, low frequency, peak frequency,

and sinuosity.

Table 3. Model fit statistics for generalized additive models (GAMs) describing temporal

variation in acoustic features across chick development. Temporal patterns indicate whether

nonlinear change across weeks was detected consistently across call types or restricted to

specific call types.

Acoustic feature	n	Adjusted R²	Deviance explained (%)	Temporal Pattern Detected
Bandwidth	1633	0.135	14.1	All call types
Duration	1633	0.705	70.7	All call types
High frequency	1633	0.046	5.23	All call types

Low frequency	1633	0.117	12.1	BEG only
Peak frequency	1633	0.062	6.77	BEG, MSB
Sinuosity	1633	0.384	38.6	BEG only

All models used a Gaussian error distribution with identity link. Temporal variation was modelled using penalized smooths of week fitted separately for each call type (s(week, by = call_type), k = 4). "Temporal pattern detected" summarizes whether the smooth term of week was significant (p < 0.05) for all call types or only a subset, reflecting call-type–specific temporal structure rather than overall differences among call types.

262 *Breakpoint Analysis (Segmented Regression)*

263   Recognizing that the transition between call types may be gradual, especially the

transition between BEG and MSB, we pooled all call categories together and used a breakpoint

analysis to identify major shifts in the vocal behavior during chick development. Segmented

regression identified significant shifts in the rate of change for each acoustic feature (Fig. 5). The

greatest change in frequency parameters (high frequency, low frequency, bandwidth) occurred

during weeks 3 and 4 (crèche stage), whereas the greatest change in duration and sinuosity

occurred around week 5 (also in crèche stage).

Figure 5. Breakpoint analysis of chick call features. Data represents all call types pooled

together. Vertical lines indicate weeks where segmented regression identified a significant shift in

the rate of change for each acoustic feature, highlighting potential critical periods of vocal

development. Shaded regions illustrate the slope before and after each breakpoint, with

corresponding values provided in Table 4.

Table 4. Breakpoints were estimated using segmented linear regressions fitted to GAM-predicted values averaged across call types. Slopes represent the rate of change per week before and after the estimated breakpoint.

Feature	BP Week	Slope before BP	Slope after BP	Directional change
Bandwidth	3.57	0.016	0.01	Increase → slower increase
Duration	5.56	0.047	−0.005	Increase → plateau / slight decline
High freq.	3.27	−0.154	0.048	Decrease → increase
Low freq.	3.38	−0.320	0.022	Decrease → increase
Peak freq.	3.61	−0.203	0.001	Decrease → stabilization
Sinuosity	4.69	0.106	0.064	Increase → slower increase

Discussion

As *P. adeliae* chicks develop, their vocalizations also change in both temporal pattern and acoustic features. During the guard stage, vocalizations are overwhelmingly BEG calls, characterized by short (< 0.3 seconds) durations, higher (4.5 kHz) frequencies, and low sinuosity. As the chicks transition to the crèche stage, MSB and LMD calls begin to emerge, characterized by longer durations, lower frequencies, and more sinuosity compared to the BEG calls (Fig. 2). MSB predicted probabilities peak around week 4 (start of crèche) with an associated decline in BEG call probabilities, whereas LMD predicted probabilities steadily increase over development (Fig. 3). Within BEG and MSB call types, we see further structural changes as chicks age. Across our observed development period, BEG and MSB calls became more broadband, longer, and more tonally complex (sinuosity), while LMD calls peaked in bandwidth around week 4 and demonstrated consistent sinuosity across weeks (Fig. 4). Together, these patterns suggest progressive modification of call structure across developmental stages, with some call types

showing more pronounced temporal changes than others. These changes may be related to

colony dynamics due to developmental changes, behavioral transitions, and/or shifts in colony

activity that favor specific call types. For example, during the guard stage, when the chicks are

primarily near their parents, BEG calls would be expected, as these calls are used to stimulate the

parent to feed the chick (Spurr, 1975). In contrast, during the crèche stage, chicks are unguarded

by parents for several hours a day and left in clusters with other chicks. Upon reunion with

parents, both "immature versions of the adult LMD" and complete LMD are made prior to

begging calls, which play a crucial role in parent-offspring recognition (Spurr, 1975), with this

recognition emerging between 8 and 17-days post-hatch (Thompson and Emlen, 1968). Our

results are consistent with the description from Spurr (1975), especially if our classified MSB is

an immature version of the LMD. The similarities in spectral features (Fig. 2) and the peak in

MSB at the start of crèche (Fig. 3), suggest the MSB should be considered a precursor to the

LMD. The MSB/LMD reunification calls occur with a behavioral display clearly distinguishable

from that of begging (Spurr 1975).

Breakpoint analysis further reinforces the suggestion that changes in the chick call

features are associated with developmental stage. All spectral features (bandwidth, high

frequency, peak frequency, low frequency), demonstrate a significant shift in the rate of change

between weeks 3 and 4, which is at the start of the crèche period. This is consistent with the

emergence (Fig. 3) and subsequent stabilization of frequency for the MSB and LMD calls (Fig.

4), suggesting early refinement of critical call spectral features, which may be important for

parental recognition. Although the spectral features of calls remain relatively stable after week 4,

the duration and sinuosity of calls continue to increase over development, reflecting later

enhancement of complex calls possibly used for social signaling or competition, which may be

significant during the crèche stage.

In "open-ended" vocal learners such as songbirds (passerines), high vocal sinuosity in

early developmental stages have represented exploratory or "babbling" behavior as a way for

juveniles to practice sounds before vocal motor control is refined (Arato et al, 2021; Elie &

Theunissen et al, 2020; Aronov et al., 2008). As they mature and experience vocal plasticity, the

variability in sinuosity decreases and aligns with increased control and stability of sound (Arato

et al, 2021). This process is characteristic of species that undergo a distinct learning phase via

practice, feedback mechanisms, and maturation of vocal organs (Elie & Theunissen et al, 2020).

However, in "closed-ended-learners" (i.e., species with limited vocal modification potential) that

rely on innate vocal patterns, vocal development may not rely on such high variability (Arato et

al, 2021). While sinuosity cannot capture the combined effect of phenological stage and call type

together, it may provide valuable ontological information when the context of each factor is

interpreted separately. In this context, an increase in sinuosity with age could suggest an increase

in control over vocalizations, suggesting vocal refinement in older chicks. Conversely, very low

sinuosity in younger chicks may reflect less control or simple vocal structures. In contrast to the

vocal development of passerine birds, these data may suggest that high sinuosity corresponds

with more complex or modulated calls in the penguin chicks, as opposed to indicating refined,

less variable calls in passerine species. In other words, the complexing of learned avian *song*

may decrease to signify mature vocalizations, whereas the complexing of avian *calls* in non-

passerines may increase to signify mature vocalizations.

For our analysis, we grouped calls based on call types from distinct spectrographic

features, but there is the possibility that these call types are instead the same call in different

forms. When we accounted for this by grouping all call types together, we still see shifts in vocal

parameter across weeks (Fig. 5), indicating chicks undergo gradual vocal development as they

age. This could suggest developmental fine-tuning within a fixed call type, supporting a form of

vocal maturity without the acquisition of new calls. If the call functions primarily for a limited

purpose, like signaling hunger or distress, they will not require complex vocal repertoires at early

developmental ages. However, we know that *P. adeliae* adults modulate calls in courtship rituals

and identification (Aubin & Jouventin et al, 2002; Clark et al, 2006) and descriptive studies of

chicks indicate multiple call types associated with different behaviors (Spurr, 1975). We

therefore propose that even in early developmental stages, chicks use different call types, with

the emergence of modulated calls (MSB and LMD) occurring at the start of the crèche.

Beyond our study period, there is likely further refinement of call modulation as chicks

transition their LMD calls to their adult form. In adult *P. adeliae,* LMDs are individually

distinctive and play a key role in identifying mates or communicate to conspecifics. Males

discriminate more than females to identify individuals and are more territorial with higher

fidelity to offspring than females (Speirs et al., 1991). Given the role the LMD plays in

individual recognition, learning likely plays a role, supporting the hypothesis that simpler vocal

learning mechanisms may exist outside the current understanding of vocal learning in birds (ten

Cate, 2021; Baciadonna et al., 2022) and underscoring the need for more diverse models than

those used for the complex vocal development seen in passerine birds (Vernes et al., 2021).

Given the role that physical displays play with the associated vocalizations (Spurr, 1975), further

analysis could incorporate behavioral elements to investigate whether specific social contexts

lead to different call patterns, potentially indicating some degree of plasticity. For instance, does

the presence or absence of parents or other adults near the nest affect the call rate or acoustic

features of chick vocalizations? Additionally, what role does sinuosity play in adult courtship

vocalizations, and how might sinuosity and call duration relate to chick survival or reproductive

success in subsequent years?

Sources of Error and DeepSqueak Utility

Our analysis also relies on general knowledge of breeding phenology and timelines

supported by literature (Cimino et al., 2023; Spurr, 1975; Taylor & Robertson, 1962) and visual

observations, but it is important to note that these timelines are approximate, and visual

observations can be imperfect, especially considering not every nest is monitored. Additionally,

we categorized our calls based on unique spectrographic differences, but some call types may be

inherently more variable or transitional, such as the MSB call, making them more prone to

detection errors. This could introduce biases in observed patterns, both in DeepSqueak's

detection networks and under variable environmental conditions, such as wind. For example,

wind may artificially broaden measured bandwidth, potentially confounding true biological

changes with recording artifacts, or affect penguin vocal behavior (Zhao et al., 2022). The 2022-

2023 Antarctic summer was an extremely windy period (Fradet et al., 2025), so continued

recording of chick vocalizations with larger sample sizes can further test the patterns we

observed with our data.

The performance of DeepSqueak in detecting chick calls depends largely on user

experience and on the supplemental programs used alongside it. Even so, its object-detection

framework makes it an appealing option for non-experts, especially compared to more

specialized, expert-driven tools that lack this feature. For users seeking both call detection and

extraction of acoustic variables, DeepSqueak provides a straightforward and cost-effective

solution. Nevertheless, our detection networks encountered some challenges. A single network

could not reliably distinguish adults, chicks, and other species, nor could it separate detections by

age class when trained on sampling dates rather than observed spectrographic differences among

chick calls. Our networks focused on specific call types, identified through contour

characteristics in spectrograms, and performed more consistently for detecting *P. adeliae* chick

calls. The unsupervised classifier also struggled to group chick calls, likely due to minimal

variation among a large number of similar data points.

Conclusion

These findings demonstrate the development of vocal features during early *P. adeliae*

development. In the first few weeks, chicks produce primarily begging calls, characterized by

short-duration, low sinuosity, and high frequencies. Around the start of the crèche period, chicks

begin to produce multi-syllable beg calls, which may be a precursor to the loud mutual display

calls. Over development, acoustic features of calls continue to change, suggesting vocal

refinement and a shift to producing signals longer in duration and with more frequency

modulation compared to early development. These results raise the possibility of a vocal learning

mechanism in penguins, but additional data spanning other breeding seasons and penguin species

is needed to continue this investigation.

Acknowledgements

This project is funded by the National Science Foundation EAGER Award 2226886 and

National Science Foundation OPP 2026045. The authors would like to thank Darren Roberts and

408 Megan Roberts for collecting acoustic data and conducting colony surveys, and Kevin R. Coffey

409 for guidance with DeepSqueak. Work was approved by the Antarctic Conservation Act permit

410 #ACA 2021-002.

References

Arato, J., & Fitch, W. T. (2021). Phylogenetic signal in the vocalizations of vocal learning and vocal non-learning birds. Philosophical Transactions - Royal Society. Biological Sciences, 376(1836), 20200241–20200241. https://doi.org/10.1098/rstb.2020.0241

Aronov, D., Andalman, A. S., & Fee, M. S. (2008). Specialized Forebrain Circuit for Vocal Babbling in the Juvenile Songbird. *Science (American Association for the Advancement of Science)*, 320(5876), 630–634. https://doi.org/10.1126/science.1155140

Aubin, T., Jouventin, P., Hildebrand, C. (2000). Penguins use the two–voice system to recognize each other. Proceedings - Royal Society. Biological Sciences/Proceedings - Royal Society. *Biological Sciences*, 267(1448), 1081–1087. https://doi.org/10.1098/rspb.2000.1112

Aubin, T. & Jouventin, P. (2002). How to vocally identify kin in the crowd: The penguin model. *Advances in the Study of Behavior*, 31, 243-277. https://doi.org/10.1016/S0065-3454(02)80010-9

Baciadonna, L., Solvi, C., del Vecchio, F., Pilenga, C., Baracchi, D., Bandoli, F., Isaja, V., Gamba, M., and Favaro, L. (2022). Vocal accomodation in penguins (*Spheniscus demersus*) as a result of social environment. *Proceedings of the Royal Society of London B*, 289, 20220626.

Beecher, MD. (1982). Signature Systems and Kin Recognition. *American Zoologist*, 22:3, 477-490. https://doi.org/10.1093/icb/22.3.477

Brenowitz, EA., Beecher, MD. (2005). Song learning in birds: diversity and plasticity, opportunities and challeng*e*s. *Trends in Neurosciences* (Regular Ed.), 28(3), 127–132. https://doi.org/10.1016/j.tins.2005.01.004

Cimino, MA., Conroy, JA., Connors, E., Bowman, J., Corso, A., Ducklow, H., Fraser, W., Friedlaender, A., Kim, HH., Larsen, GD., Moffat, C., Nichols, R., Pallin, L., Patterson-Fraser, D., Roberts, D., Roberts, M., Steinberg, DK., Thibodeau, P., Trinh, R., … Stammerjohn, S. (2023). Long-term patterns in ecosystem phenology near Palmer Station, Antarctica, from the perspective of the Adélie penguin. *Ecosphere*, 14(2). https://doi.org/10.1002/ecs2.4417

Cimino, M. A., Fraser, W. R., Patterson-Fraser, D. L., Saba, V. S., Oliver, M. J. (2014). Large-scale climate and local weather drive interannual variability in Adélie penguin chick fledging mass. Marine Ecology. Progress Series, 513, 253–268. https://doi.org/10.3354/meps10928

Clark, J. A., Boersma, P. D., Olmsted, D. M. (2006). Name that tune: call discrimination and individual recognition in Magellanic penguins. *Animal Behaviour*, 72(5), 1141–1148. https://doi.org/10.1016/j.anbehav.2006.04.002

Coffey, K. R., Marx, R. E., Neumaier, J. F. (2019). DeepSqueak: a deep learning-based system for detection and analysis of ultrasonic vocalizations. *Neuropsychopharmacology* (New York, N.Y.), 44(5), 859–868. https://doi.org/10.1038/s41386-018-0303-6

Davis, S. L. (1982) Creching behaviour of Adelie penguin chicks (Pygoscelis adeliae). *New Zealand Journal of Zoology*, 9:2, 279-285. https://doi.org/10.1080/03014223.1982.10423856

Elie, J. E., & Theunissen, F. E. (2020). The Neuroethology of Vocal Communication in Songbirds: Production and Perception of a Call Repertoire. *The Neuroethology of Birdsong* (Vol. 71, pp. 175–209). Springer International Publishing AG. https://doi.org/10.1007/978-3-030-34683-6_7

Faiß, M., Riede, T., Goller, F. (2022). Tonality over a broad frequency range is linked to vocal learning in birds. Biological Sciences/Proceedings - Royal Society. *Biological Sciences*. https://doi.org/10.1098/rspb.2022.0792

Favaro, L., Gamba, M., Alfieri, C., Pessani, D., and McElligot, A. G. (2015). Vocal individuality cues in the African penguin (*Spheniscus demersus*): a source-filter theory approach. *Scientific Reports*, 5, 17255.

Favaro, L., Gili, C., Da Rugna, C., Gnone, G., Fissore, C., Sanchez, D., McElligot, A. G., Gamba, M., and Pessani, D. (2016). Vocal individuality and species divergence in the contact calls of banded penguins. *Behavioural Processes*, 128, 83-88.

Fradet, D.T., Cimino, M. A., White, E. R., and Kloepper, L. N. (2025). Designing a BirdNET classifier for high wind detection in passive acoustic recordings to support wildlife monitoring. *The Journal of the Acoustical Society of America*, 157, 4502-4512.

Jones, I. L., Ryan, P. G. (2010). Seabird vocalizations and their use in species recognition. *Seabird Islands: Ecology, Invasion, and Restoration* (pp. 78-88).

Jouventin, P., Aubin, T. (2002). Acoustic systems are adapted to breeding ecologies: individual recognition in nesting penguins. *Animal Behavior*, 64, 747-757. https://doi.org/10.1006/anbe.2002.4002

Kidawa, D., Wojczulanis-Jakubas, K., Jakubas, D., Palme, R., Barcikowski, M. (2023). Mine or

my neighbours' offspring: an experimental study on parental discrimination of offspring in

a colonial seabird, the little auk Alle alle. *Scientific Reports*, 13, 15088.

https://doi.org/10.1038/s41598-023-41925-5

LeCun, Y., Bengio, Y., Hinton, G. (2015). Deep learning. *Nature* (London), 521(7553), 436–444.

https://doi.org/10.1038/nature14539

Lynch, M.A., Lynch, HJ. (2017). Variation in the ecstatic display call of the Gentoo Penguin

(Pygoscelis papua) across regional geographic scale*s. The Auk Ornithological Advances*,

134, 894–902.

Marks, E., Rodrigo, A., Brunton, D. (2010). Ecstatic display calls of the Adélie penguin honestly

predict male condition and breeding success. *Behaviour*, 147(2), 165–184.

https://doi.org/10.1163/000579509X12512863752751

Nakagawa, S., Waas, J. R., & Miyazaki, M. (2001). Heart Rate Changes Reveal That Little Blue

Penguin Chicks (*Eudyptula minor*) Can Use Vocal Signatures to Discriminate Familiar

from Unfamiliar Chicks. *Behavioral Ecology and Sociobiology*, 50(2), 180–188.

https://doi.org/10.1007/s002650100355

Robission, P., Aubin, T., Bremond, J.C. (1993). Individuality in the Voice of the Emperor Penguin

Aptenodytes forsteri: Adaptation to a Noisy Environment. *Ethology*, 94:4, 279-290.

https://doi.org/10.1111/j.1439-0310.1993.tb00445.x

Salihoglu, B., Fraser, W. R., HOFMANN, E. E. (2001). Factors affecting fledging weight of Adélie

penguin (Pygoscelis adeliae) chicks: a modeling study. *Polar Biology*, 24(5), 328–337.

https://doi.org/10.1007/s003000000215

Sladen, W. J. 1958. The Pygoscelid Penguins: I. Methods of Study. II. The Adélie Penguin. *Falkland Islands Dependencies Survey Scientific Reports* 17, 1-97.

Speirs, E. A. H., Davis, L. S. (1991). Discrimination by adélie penguins, Pygoscelis adeliae, between the loud mutual calls of mates, neighbours and strangers. *Animal Behaviour*, 41(6), 937–944. https://doi.org/10.1016/S0003-3472(05)80631-1

Spurr, E. B. (1975). Behavior of the Adélie Penguin Chick. *The Condor*, 77(3), 272–280. https://doi.org/10.2307/1366222

Taylor, J. R. E. (1985). Ontogeny of thermoregulation and energy metabolism in pygoscelid penguin chicks. *Journal of Comparative Physiology B*, 155, 615-627.

Taylor, R. H. & Roberts, H. S. (1962). Growth of Adélie penguin (*Pygoscelis adeliae* Hombron and Jacquinot) chicks. *New Zealand Journal of Science*, 5, 191-197.

ten Cate, C. (2021). Re-evaluating vocal production learning in non-oscine birds. Philosophical Transactions of the Royal Society of London. Series B. *Biological Sciences*, 376(1836), 20200249–20200249. https://doi.org/10.1098/rstb.2020.0249

Terranova, F., Baciadonna, L., Maccarone, C. et al. (2023). Penguins perceive variations of source- and filter-related vocal parameters of species-specific vocalisations. *Animal Cognition*, 26, 1613–1622.

Thompson, D. H. and Emlen, J. T. (1968). Parent-chick individual recognition in the Adélie penguin. *Antarctic Journal of the United States*, 3, 132.

Tramontin, A. D., Brenowitz, E. A. (1999). A field study of seasonal neuronal incorporation into the song control system of a songbird that lacks adult song learning. *Journal of*

Neurobiology, 40(3), 316–326. https://doi.org/10.1002/(SICI)1097-
4695(19990905)40:3<316::AID-NEU4>3.0.CO;2-S

Vernes, C., S., Kriengwatana, Pralle, B., Beeck, C., V., Fischer, Julia, Tyack, L., P., Cate, t., Carel,
Janik & M., V. (2021). *The multi-dimensional nature of vocal learning*. Philosophical
Transactions of the Royal Society B: Biological Sciences 376.
https://royalsocietypublishing.org/doi/full/10.1098/rstb.2020.0236

Vicario, D., Raksin, J., Naqvi, N., Thande, N., Simpson, H. (2002). The relationship between
perception and production in songbird vocal imitation: what learned calls can teach us.
Journal of Comparative Physiology, 188(11–12), 897–908.
https://doi.org/10.1007/s00359-002-0354-2

Zhao, K., Chen, G., Liu, Y., Møller, A. P., and Zhang, Y. (2022). Population size assessment of
Adélie penguin (*Pygoscelis adeliae)* chicks based on vocal activity rate index. *Global
Ecology and Conservation*, 38, e02263.

529