Task goals reorganize visual codes in human ventral temporal cortex via medial frontal modulation

Robert Kim[1], Tomas G. Aquino[2], Adam N. Mamelak[3], Chrystal M. Reed[1], Nuttida Rungratsameetaweemana[2] *, and Ueli Rutishauser[1,3,4,5] *

[1] Department of Neurology, Cedars-Sinai Medical Center, Los Angeles, CA, USA

[2] Department of Biomedical Engineering, Columbia University, New York, NY, USA

[3] Department of Neurosurgery, Cedars-Sinai Medical Center, Los Angeles, CA, USA

[4] Center for Neural Science and Medicine, Department of Biomedical Sciences, Cedars-Sinai Medical Center, Los Angeles, CA, USA

[5] Division of Biology and Biological Engineering, California Institute of Technology, Pasadena, CA, USA.

*Joint senior authors

Abstract

How higher cortical regions dynamically modulate sensory representations to support cognitive flexibility remains a fundamental open question. Using rare simultaneous single-neuron recordings across multiple human brain regions, we show that representations in ventral temporal cortex (VTC), not traditionally associated with context-dependent coding, are dynamically reconfigured by the timing of task instructions. Identical visual stimuli evoked distinct responses in VTC depending on when task instructions were provided, with categorical representations sharpened when goals were specified in advance. In contrast, dorsal anterior cingulate cortex (dACC) and hippocampus carried strong signals related to instruction timing, consistent with known regional specialization in top-down control. Coupling between dACC and VTC increased with uncertainty and for high cognitive demand, and its strength predicted correct performance. These findings uncover a dynamic feedback mechanism through which medial frontal cortex reorganizes sensory population codes in VTC, linking flexible cortical coordination to successful goal-directed computation.

Introduction

Goal-directed behavior requires the brain to dynamically adjust how sensory inputs are represented and interpreted depending on task demands. Human ventral temporal cortex (VTC) has traditionally been viewed as faithfully encoding properties of objects and faces present in the external visual input [1–3]. More recently, it has become clear that these representations are not fixed but can instead be modulated by behavior and task contexts [4, 5]. In this new framework, identical visual stimuli can evoke distinct neural responses in VTC depending on the current task, expectation, and previous experience. Whether such context-dependent modulation occurs in human sensory areas remains unclear. In particular, it is not yet known whether the human VTC, a region canonically associated with stable visual category coding [1, 6–9], can flexibly reorganize its activity as a function of task context, and if so, how quickly this can occur.

Historically, prior studies in both non-human primates and rodents have characterized the ventral visual stream as a primarily feedforward system specialized for encoding stimulus features. In non-human primates, neurons in inferotemporal (IT) cortex, the homologue of the human VTC, exhibit selectivity for visual categories such as faces and objects, forming hierarchical representations that are remarkably invariant to low-level stimulus changes [10–15]. Similarly, neurons in rodent lateral visual areas encode low-level visual features such as orientation, contrast, and shape, with limited sensitivity to behavioral state or task demands [16, 17].

More recent studies in animals have challenged this purely sensory-encoding view, demonstrating that sensory representations can flexibly change depending on behavioral context, attentional state, or task goals. For example, in rodents, visual cortical responses are modulated by learned associations, expectations, and reward contingencies even when the physical stimuli remain identical [18–23]. In contrast, evidence for such context-dependent modulation in non-human primate visual cortex remains limited. Even less is known in humans, although recent fMRI work has provided evidence that visual cortical representations can be flexibly reorganized by task context [5]. However, there is presently no direct neuronal evidence for such modulation and the circuit mechanisms by which higher-order areas exert top-down influence on human sensory cortex remain unknown.

We developed a visual decision-making paradigm that is designed to disentangle sensory encoding from modulation driven by the timing of task instructions. By combining this paradigm with

rare simultaneous single-neuron and local field potential (LFP) recordings from multiple human brain regions spanning the cortical hierarchy, ranging from the VTC to higher-order control areas, we examined how instruction timing shapes sensory codes at both single-neuron and inter-areal levels. We show that the temporal availability of task instructions modulates sensory codes in human VTC, and further demonstrate that coupling from the dorsal anterior cingulate cortex (dACC) to VTC was critical for successful performance under increased uncertainty and cognitive demand, suggesting that dACC serves as a key source of dynamic top-down modulation of VTC. Together, these findings reveal a previously uncharacterized mechanism at the human single-neuron and LFP level through which higher-order control regions dynamically recalibrate sensory representations to optimize goal-directed perception and behavior.

Results

Single-neuron recordings across multiple brain areas during a flexible visual decision-making task. We recorded activity from single neurons (773 neurons) in six brain areas, including ventral temporal cortex (VTC; $n = 164$), dorsal anterior cingulate cortex (dACC; $n = 74$), ventromedial prefrontal cortex (vmPFC; $n = 70$), pre-supplementary motor area (preSMA; $n = 185$), hippocampus (HIPP; $n = 137$), and amygdala (AMY; $n = 143$), from 11 patients with refractory epilepsy who underwent depth electrode implantation for clinical monitoring (Fig.1b; see Extended Data Table 1).

Participants performed a flexible visual decision-making task designed to dissociate sensory input from task context (Fig. 1a). On each trial, two visual stimuli were presented sequentially, and participants were instructed either before ("early" instruction) or after ("late" instruction) the first stimulus how to choose between the two (a manipulation we refer to as "instruction timing"). Instructions specified either a "pro" condition, in which participants followed the specified rule directly, or an "anti" condition, in which they applied the opposite rule (see Extended Data Table 2 for examples instructions). The anti condition imposed greater cognitive demands due to its double-negative structure. Crucially, the same images were presented in both early and late instruction conditions, enabling us to investigate how VTC and other regions represent identical visual inputs under different task contexts.

Behaviorally, participants performed the task with high accuracy (mean $\pm$ s.d.; $75.28 \pm 12.45\%$).

Fig. 1 | Task design, electrode locations, and behavior. a, Two example trials from the flexible visual decision-making task. On each trial, two stimuli were presented sequentially. An instruction was given either before the first stimulus (early instruction; top) or after the first stimulus (late instruction; bottom), specifying whether participants should follow the rule directly (pro) or apply the opposite rule (anti) when making their choice. **b**, Electrode coverage. Colored dots represent microwire bundle electrode locations in Montreal Neurological Institute (MNI) space. **c–f**, Behavioral performance. Task accuracy (**c** and **e**) and reaction times (**d** and **f**) are shown for early vs. late instruction timing conditions (**c** and **d**) and for pro vs. anti tasks (**e** and **f**). Participants showed significantly higher accuracy and faster reaction times in the pro compared to the anti condition, while no significant differences were observed between early and late instruction timing conditions. $*P < 0.01$ and $***P < 0.001$ by two-sided Wilcoxon rank-sum test.

Accuracy was significantly higher for pro compared to anti trials, consistent with greater cognitive

demands in the anti condition (Fig.1c; mean $\pm$ s.d., $85.70 \pm 11.14\%$ vs. $64.87 \pm 17.04\%$; $P < 0.001$

by two-sided Wilcoxon signed-rank test). Reaction times were also longer for anti compared to pro

trials (mean $\pm$ s.d.; 1.41 ± 0.16s vs. 1.17 ± 0.23s; $P < 0.005$ by two-sided Wilcoxon signed-rank

test). In contrast, no significant differences in accuracy or reaction times were observed between

early and late instruction trials (accuracy: $76.14 \pm 12.34\%$ for early vs. $74.43 \pm 14.30\%$ for late;

reaction time: 1.29 ± 0.16s vs. 1.29 ± 0.20s; $P > 0.5$, two-sided Wilcoxon signed-rank test). To

further examine behavioral effects, we compared early versus late instruction trials within pro and

anti conditions separately. Although task performance did not differ significantly between early

and late trials within either the pro or anti conditions, early pro trials showed significantly faster

reaction times than late pro trials ($P < 0.05$ by one-sided Wilcoxon signed-rank test; Extended

Data Fig. 1). No significant difference in reaction time was observed between early and late trials
for the anti condition.

VTC neurons are modulated by both stimulus category and instruction timing. We
first asked whether single-neuron activity in human VTC reflects not only the visual category of
the stimulus but also the timing of task instruction, which determines when the goal is known
relative to stimulus onset. We first examined the firing rates of single-neurons following onset of
stimulus 1 (stim 1). We observed robust category selectivity in many VTC neurons (examples
shown in Fig. 2a,b), consistent with the established role of VTC in forming sensory representations
of visual categories [24].

Performing a two-way 4×2 ANOVA with interactions with factors of visual category (there
were four possible categories) and instruction timing during the stim 1 window revealed a large
proportion of VTC neurons that were category-selective (69 of 164 neurons, 42.07%; Fig. 2c,d). The
preferred category was determined as the category that elicited the highest mean firing rate during
the stim 1 window for the early instruction trials only. In contrast, only 8 neurons (4.9%) exhibited a
significant main effect of instruction timing in the absence of a significant category effect, indicating
that contextual modulation alone was uncommon. Importantly, we also identified a subgroup of
neurons that exhibited significant main effects of both stimulus category and instruction timing
(e.g., joint selectivity), with category responses that varied depending on whether the instruction
was presented early or late (Fig. 2c,d; 17 of 164 neurons, 10.37%). Within this jointly selective
subpopulation, some neurons showed enhanced responses to their preferred category during early
instruction trials, whereas others showed suppressed responses under the same condition (Fig. 2e).
These results demonstrate that while category selectivity dominated in VTC, a subset of neurons
flexibly integrated both sensory and contextual information, suggesting that VTC may be subject
to context-dependent top-down modulation.

Several of the category-selective neurons were also active during the instruction period when
their preferred category was mentioned, for early (7 of 57 category-selective VTC neurons) and
late instruction (15 of 57) trials, suggesting that verbal task cues can activate or sustain category-
specific representations in VTC (see Extended Data Fig. 2). In addition, some neurons exhibited
pronounced instruction-dependent modulation of category responses: activity was either enhanced

Fig. 2 | VTC neurons encode stimulus category and instruction timing. a, Example VTC neuron showing category-selective responses (to car images in this case) during both the stimulus and instruction windows. **b**, Two additional sample VTC neurons illustrating instruction-related modulation. Neuron 2 showed enhanced responses to preferred categories during early instruction trials, while Neuron 3 showed diminished responses under the same condition. Raster plots (left) and peristimulus time histograms (right) are aligned to stimulus 1 onset. Each row in the raster plot corresponds to a single image, shown under both early and late instruction conditions. **c**, Two-way ANOVA (category × instruction timing) during the stim 1 window revealed three classes of VTC neurons: category-selective (cyan), instruction-timing-selective (magenta), jointly selective (orange), and non-selective (black). **d**, Proportion of neurons in each category, showing that most were category-selective, with a smaller but notable jointly selective population. The dashed line indicates the 5% chance level. **e, f**, Average firing rates during the stim 1 window for jointly selective neurons. These neurons could be divided into two groups: one subgroup showed enhanced responses to their preferred category during early instruction trials (**e**), while the other subgroup showed suppressed responses under the same condition (**f**). Boxplot central lines, median; bottom and top edges, lower and upper quartiles; whiskers, 1.5*interquartile range; outliers not plotted. *$P < 0.01$ and ***$P < 0.001$ by two-sided Wilcoxon signed-rank test.

(Fig. 2b, left) or suppressed (Fig. 2b, right) when the instruction preceded stimulus 1, indicating

flexible gain control dependent on task context.

Population-level decoding of instruction timing and category in VTC. Given the modu-

lation related to instruction timing we observed at the single-neuron level in VTC, we next asked

whether we could reliably decode the presence of instruction during the stimulus windows from

VTC population activity. Using a linear SVM decoder during the stim 1 window, we found that

VTC population activity (pseudopopulation of all recorded neurons across participants) reliably

differentiated early from late instruction trials, with decoding accuracy significantly above chance

(Fig. 3a; mean $\pm$ s.d., $64.78 \pm 6.68\%$ vs. $50.66 \pm 7.12\%$ for shuffled data). We next visualized

the structure of VTC population activity by projecting it into a low-dimensional latent space using

Contrastive Embedding for Behavioral and Neural Analysis (CEBRA; [25]). CEBRA is a non-linear

dimensionality reduction framework designed for uncovering latent population dynamics from neu-

ral data. In this analysis, we used an unsupervised implementation (i.e., without providing any

task or behavioral labels) to visualize the intrinsic structure of VTC activity across the two instruc-

tion timing conditions (early and late; see Methods). Even without explicit task information, this

approach revealed distinct neural trajectories for early and late instruction conditions during the

stim 1 window (Fig. 3b). Computing the distance between the two trajectories revealed that early

and late instruction trials were reliably separated during the stim 1 period compared to shuffled

controls (Fig. 3c).

How did instruction timing information modify representations of categories in VTC? When the

instruction was provided before stim 1 (i.e., early instruction trials), category representations were

sharpened, as indicated by larger pairwise distances between category trajectories in the CEBRA

latent space (Fig. 3d) and greater overall category separation (Fig. 3e). This increase in represen-

tational distance suggests that advance knowledge of task goals allows VTC population activity to

more effectively differentiate between stimulus categories. In contrast, when the instruction was

given after the first stimulus (i.e., late instruction trials), trajectories corresponding to different

categories were more overlapping (Fig. 3d, blue), reflecting a less differentiated sensory represen-

tation during the initial encoding phase. Consistent with these findings, stimulus category could

be decoded with significantly higher accuracy in early compared to late instruction trials (Fig. 3f;

Fig. 3 | VTC population activity encodes instruction timing and sharpens category representations. a, Linear SVM decoding during the stim 1 window using a pseudopopulation of all recorded VTC neurons revealed that instruction timing (early vs. late) could be decoded significantly above chance. **b**, Low-dimensional neural trajectories obtained with CEBRA for early (blue) and late (red) instruction trials during the stim 1 window. **c**, Distance between early and late trajectories shown in b, compared with shuffled controls. **d**, Category trajectories in the CEBRA latent space for early versus late instruction trials during the stim 1 window (at 0.70 s after stim 1 onset). **e**, Average pairwise category distances during the stim 1 window. **f**, Linear SVM decoding accuracy for category identity during the stim 1 window for early and late instruction trials. **g-j**, Similar analyses as in **a**–**f**, performed during the stim 2 window. Ellipsoids in the CEBRA trajectories (**b**, **h**) show trial variability, estimated from the covariance of neural activity at selected time points and plotted as orthogonal projections in latent space. Boxplot central lines, median; bottom and top edges, lower and upper quartiles; whiskers, 1.5*interquartile range; outliers not plotted. **$P < 0.005$ and ***$P < 0.001$ by two-sided Wilcoxon rank-sum test (**a**, **e**, **g**, **k**) or Kruskal-Wallis test with post hoc Dunn's multiple comparisons test (**f**, **l**). Thick horizontal black bars in **c** and **i** indicate time periods of significant differences identified using a cluster-based bootstrap procedure ($P < 0.05$, cluster-level corrected; see Methods). Circles and triangles mark the onset and offset of the stim 1 analysis window, respectively.

mean $\pm$ s.d., 64.62 $\pm$ 8.67% for early, 52.12 $\pm$ 10.52% for late, and 26.75 $\pm$ 10.88% for the shuffled

null condition), confirming that contextual information enhances the fidelity of category-specific

coding in VTC.

We next performed the same analyses during the stim 2 window. During this time window,

instructions have always been shown but in some instances just before stim 2 was shown (late) vs.

already before the stim 1 window. By this stage both the early and late conditions thus contained

equivalent amounts of sensory and contextual information. Despite this, at the population level,

VTC still differed between the early and late conditions (Fig. 3a), albite with less accuracy than

during the stim 1 window (Fig. 3a; mean $\pm$ s.d., 53.81 $\pm$ 7.35% vs. 50.50 $\pm$ 7.69% for shuffled data).

Neural trajectories in the CEBRA latent space also remained well separated between early and late

instruction conditions (Fig. 3h,i), indicating that contextual timing differences in VTC population

activity persisted throughout the trial. Pairwise category distances were large for both early and late

instruction trials at this stage (Fig. 3j,k), suggesting that category information became robustly

represented in VTC regardless of instruction timing. However, decoding again revealed higher

category decodability when the instruction immediately preceded the stimulus, corresponding to

the stim 2 window for late instruction trials (Fig. 3l; mean $\pm$ s.d., 59.06 $\pm$ 11.80% for early, 69.75 $\pm$

10.72% for late, and 26.12 $\pm$ 11.53% for the shuffled null condition). Together, these results indicate

that VTC population activity flexibly integrates sensory information with instruction timing across

time, with category representations sharpened when task context immediately preceded stimulus

presentation.

Signals related to instruction timing in hippocampus and dACC. Having shown that VTC

encodes not only stimulus category but also exhibits modulation by instruction timing for the same

stimuli, we next investigated potential sources of these top-down signals, defined here as contextual

influences from higher-order brain regions that shape sensory responses. Recent animal studies

demonstrate that top-down feedback signals from frontal cortex can dynamically modulate visual

cortical processing. For example, Ährlund-Richter et al. showed that mouse prefrontal subregions,

especially anterior cingulate cortex, exert behavioral-state-dependent feedback onto primary visual

cortex [15]. Motivated by these insights, we next asked whether higher-order human brain areas,

including hippocampus and dorsal anterior cingulate cortex (dACC), encode instruction timing

signals that could provide top-down inputs necessary to flexibly shape VTC sensory representations.

We first quantified single-neuron selectivity in hippocampus and dACC using a two-way $4 \times$
2 ANOVA with factors of category and instruction timing, as in VTC (Fig. 4a). Both regions
contained more neurons selective for instruction timing than VTC, with dACC showing the highest
proportion (HIPP, 20 of 137 neurons, 14.60%; dACC, 17 of 74 neurons, 22.97%). Decoding similarly
followed this hierarchy, with accuracy for distinguishing early versus late instruction trials highest
in dACC (Fig. 4b). We also extended the two-way ANOVA analysis to the amygdala (AMY),
pre-supplementary motor area (preSMA), and ventromedial prefrontal cortex (vmPFC) (Extended
Data Fig. 3). The amygdala showed a pattern similar to the hippocampus, with a moderate
proportion of neurons selective for instruction timing. In contrast, both preSMA and vmPFC
exhibited patterns resembling dACC, with a larger fraction of neurons showing instruction-related
modulation, consistent with recent human single-neuron recordings demonstrating that preSMA
neurons encode abstract decision variables and integrated choice signals during flexible behavior
[26]. Consistent with this pattern, decoding of instruction timing revealed higher accuracy in
medial frontal regions (preSMA, dACC, and vmPFC) compared to VTC during the stim 1 window
(Extended Data Fig. 4 left). Although this hierarchical separation was less pronounced during the
stim 2 window, medial frontal areas continued to show robust instruction-related signals relative to
VTC (Extended Data Fig. 4 right). These findings confirm that instruction timing is represented
across a distributed network of higher-order regions, with the strongest contextual modulation
emerging in the medial frontal cortical areas dACC, preSMA, and vmPFC.

Visualization of population activity using CEBRA revealed that trajectories in both hippocam-
pus and dACC were well separated between early and late instruction conditions, demonstrating
robust representation of instruction timing at the population level (Fig. 4c,d).

We then examined how stimulus category information was represented across these regions.
Decoding revealed that category decoding ability decreased systematically along the hierarchy,
with VTC showing the highest decoding and HIPP and dACC showing lower decoding accuracy;
however, decoding in HIPP remained significantly above chance (Fig. 4e,f; Matched for number
of neurons across areas). Extended Data Fig. 5 further illustrates this pattern by showing cate-
gory decoding for early and late instruction trials across all regions. Consistent with the hierarchy

Fig. 4 | Signals related to instruction timing in hippocampus (HIPP) and dorsal anterior cingulate cortex (dACC). a, Proportion of neurons selective for stimulus category (cyan), instruction timing (magenta), or both (orange), based on two-way ANOVA (category × instruction timing). **b**, Linear SVM decoding accuracy for distinguishing early versus late instruction trials in HIPP and dACC during the stim 1 window (left) and stim 2 window (right). **c**, Low-dimensional population trajectories obtained with CEBRA for early (blue) and late (red) instruction trials in HIPP and dACC during the stim 1 window (top) and stim 2 window (bottom). Ellipsoids indicate trial variability estimated from the covariance of trajectories at selected time points. **d**, Linear SVM decoding accuracy for stimulus category across regions, showing highest decoding in VTC and reduced decoding in HIPP and dACC. *$P < 0.05$ and ***$P < 0.001$ by Kruskal-Wallis test with post hoc Dunn's multiple comparisons test. Circles and triangles mark the onset and offset of the stim 1 analysis window, respectively.

observed in the main analysis, VTC exhibited the highest category decodability during both stimu-

lus windows, whereas medial frontal and medial temporal regions showed markedly lower decoding

accuracy, reinforcing the distinction between lower-level sensory regions that primarily encode stim-

ulus features and higher-order areas that emphasize task context. VTC again showed sharpened

category representations when the instruction immediately preceded the stimulus window of inter-

est. The results here differ slightly from Fig. 3f,l because the number of neurons was matched across

regions to the smallest sample size, which was 74 in dACC. This pattern is consistent with VTC's

established role in bottom-up sensory coding, whereas higher-order regions primarily emphasized

instruction-related information.

Together, these analyses suggest that higher-order regions such as dACC may provide top-down

signals that modulate VTC, enabling sensory representations in VTC to be flexibly shaped by task

context.

dACC-VTC modulation reflects uncertainty, cognitive demand, and behavioral per-

formance. We next employed spike-field coherence (SFC) analyses to examine potential sources

of top-down modulation to VTC. Previous studies in non-human primates demonstrated that feed-

back projections from prefrontal and cingulate areas modulate sensory processing in visual cortices

through rhythmic synchronization across broad frequency ranges, including the theta (4-7 Hz) and

beta (12-32 Hz) bands [15, 27, 28]. Motivated by these findings, we tested whether VTC spikes

phase-locked to ongoing oscillations in the LFP in the other simultaneously recorded areas, includ-

ing dACC (Fig. 5a; see Methods). Focusing on the stim 1 window and comparing early versus late

instruction trials revealed that VTC spikes were more strongly phase-locked to LFPs from dACC

across a broad frequency range in early relative to late trials, especially in the beta band (12-32 Hz;

Fig. 5b). Investigating whether LFP activity from other regions was coordinated with VTC spiking

revealed that the amygdala also exhibited strong late-instruction modulation, primarily within the

theta range (Extended Data Fig. 6). Examining the reverse direction, i.e. whether spikes in other

areas were phase-locked to VTC LFP, showed robust early-instruction modulation from VTC to

dACC, potentially indicating that early contextual information sharpens dACC-VTC interactions

(Extended Data Fig. 6).

These findings suggest that dACC exerts modulation of VTC during periods of uncertainty,

specifically in late instruction trials within the stim 1 window during which instructions have

not yet been provided by the time the stimulus appears. We next asked whether this top-down

modulation was behaviorally relevant by examining whether the strength of dACC-VTC SFC was

associated with task performance. We compared dACC-VTC SFC between correct and incorrect

trials, separately for early (Fig. 5c) and late (Fig. 5d) instruction conditions. We ensured that the

Fig. 5 | dACC–VTC spike–field coherence (SFC) reflects uncertainty, cognitive demand, and behavioral performance. a, Example recording configuration showing simultaneous dACC local field potential (LFP) signal and VTC spiking activity. **b**, SFC aligned to stimulus 1 for early vs. late instruction trials. Left, schematic of trial types. Middle, SFC spectra showing increased beta-band (12–32 Hz) coherence for early compared to late instruction trials. Right, paired comparisons of beta-band SFC across neurons ($n = 47$). **c**, SFC during stimulus 1 for correct vs. incorrect trials. Left, schematic. Middle, spectra showing comparable coherence between correct and incorrect trials. Right, paired comparisons in the beta band ($n = 40$). **d**, SFC during stimulus 1 for correct vs. incorrect trials, showing significantly increased beta-band coherence on correct compared to incorrect trials ($n = 42$). **e**, SFC for pro vs. anti trials. Left, schematic. Middle, spectra showing elevated theta-band (4–7 Hz) coherence in anti compared to pro trials. Right, paired comparisons of average theta-band SFC ($n = 44$). **f**, Average theta-band SFC for correct vs. incorrect trials separated by task condition, shown for pro (left) and anti (right) trials. Thick horizontal black bars indicate time periods of significant differences identified using a cluster-based bootstrap procedure ($P < 0.05$, cluster-level corrected. Boxplot central lines, median; bottom and top edges, lower and upper quartiles; whiskers, 1.5*interquartile range; outliers not plotted. *$P < 0.05$, **$P < 0.01$, and ***$P < 0.001$ by two-sided Wilcoxon signed-rank test.

number of trials and spikes was matched between correct and incorrect sets within each instruction

timing condition (see Methods). During early instruction trials, dACC-VTC SFC did not differ

significantly between correct and incorrect responses (Fig. 5c). In contrast, during late instruction

trials, dACC-VTC coupling in the beta band was significantly stronger on correct compared to

incorrect trials (Fig. 5d). Repeating this analysis in the reverse direction (i.e., SFC between VTC
LFP and dACC spikes) revealed no significant increases for correct trials (Extended Data Fig. 7),
indicating that performance-related modulation was specific to top-down dACC $\rightarrow$ VTC inter-
actions rather than bottom-up signaling. These results indicate that when task uncertainty was
high (because instructions hadn't been provided yet), stronger top-down modulation from dACC
to VTC was associated with successful performance.

We next asked whether strong dACC-VTC coupling was also observed under conditions of high
cognitive demand. To test this, we compared SFC between pro and anti trials during the stim 1
window, noting that participants performed significantly worse and slower on anti trials than on
pro trials (Fig. 1c), thereby indicating that anti trials were more cognitively demanding. SFC was
significantly elevated in the theta-band (4-7 Hz) for anti trials compared to pro trials (Fig. 5e). Fur-
thermore, when trials were separated by accuracy, theta-band coherence was significantly stronger
for correct compared to incorrect responses in anti trials, whereas no such difference was observed
for pro trials (Fig. 5f). These effects were confirmed using paired Wilcoxon signed-rank tests per-
formed separately within each frequency band, with false discovery rate (FDR) correction applied
across four frequency bands (theta 4-7 Hz; alpha 8-12 Hz; beta 12-32 Hz; gamma 32-80 Hz). After
FDR correction, accuracy-related differences were significant only in the theta band for anti trials
($P = 0.0368$), with no significant effects observed in the alpha, beta, or gamma bands (all $P > 0.7$),
nor in any frequency band for pro trials (all $P > 0.45$).

These findings suggest that dACC-VTC coupling increases not only during uncertainty but also
during high cognitive demand, and that stronger coupling supports successful performance in these
conditions. Such dynamic modulation of VTC by dACC may reflect a flexible top-down mechanism
through which medial frontal cortices synchronize sensory representations to optimize goal-directed
performance.

Discussion

274 In this study, we investigated how the timing of task instructions dynamically shapes sensory
275 representations in the human brain using simultaneous single-neuron recordings across multiple
276 cortical and medial temporal regions. We show that identical visual stimuli evoke distinct patterns
277 of activity in VTC depending on when task instructions are provided, demonstrating that sensory

representations in human VTC are flexibly reorganized by instruction timing even when sensory input is held constant. While the present study focuses on how task instruction timing modulates sensory coding in VTC, the question of where and how instruction content itself is represented is addressed in a companion study using the same task framework. To our knowledge, the current study provides the first human single-neuron and population-level evidence that the temporal availability of task goals can dynamically reshape representations in a high-level visual area traditionally viewed as primarily sensory. At the same time, medial frontal regions, particularly dACC, carried robust signals related to instruction timing and exhibited frequency-specific, performance-predictive interactions with VTC, revealing a dynamic top-down mechanism through which sensory population codes were modulated to support goal-directed computation. Importantly, this is the first evidence in humans that top-down interactions between medial frontal cortex and a high-level visual area are selectively strengthened under instructional uncertainty and increased cognitive demand in a manner that predicts behavioral performance.

Converging evidence from studies in animals suggests that top-down feedback from frontal and cingulate cortices modulates sensory processing through frequency-specific oscillatory coupling. Beta-band synchronization has been linked to the maintenance of task sets and the transmission of predictive signals from higher-order areas to sensory cortices [27, 29, 30], whereas theta-band synchronization supports adaptive updating and cognitive control under uncertainty [31]. The increase in dACC-VTC beta coherence we observed during late-instruction trials likely reflects top-down stabilization of sensory codes when goal information must be retroactively integrated with already-encoded stimuli. Critically, stronger beta coupling predicted successful performance specifically when uncertainty was highest (in late instruction trials), suggesting that this mechanism is behaviorally consequential. These findings are aligned with previously described roles of beta-band coupling in mediating cognitive control [15, 32–34]. Conversely, elevated theta-band coupling during anti trials may index dynamic control processes required to apply inverted task rules. Similarly, theta enhancement also predicted correct performance specifically under high cognitive demand, supporting its role in coordinating flexible rule application. These frequency-dependent interactions are consistent with hierarchical predictive coding frameworks in which beta oscillations convey top-down predictions while theta oscillations coordinate cognitive control and error monitoring [35–

38]. The dissociation between beta-mediated modulation during uncertainty and theta-mediated
modulation during cognitive demand suggests that dACC employs frequency-multiplexed channels
to flexibly coordinate with VTC. This architecture may enable the same circuit to simultaneously
maintain stable sensory representations while allowing for rapid, context-dependent reconfiguration
when task demands shift.

At the microcircuit level, feedback from medial frontal regions could engage specific classes
of inhibitory interneurons in VTC to implement the context-dependent modulation we observed.
In line with this possibility, prior studies in mice have demonstrated that top-down projections
from higher-order cortical areas specifically engage vasoactive intestinal peptide (VIP)-expressing
inhibitory neurons in visual cortex, which in turn inhibit somatostatin (SST)-expressing interneu-
rons, resulting in disinhibition of pyramidal neurons [20, 21, 28]. These studies further show
that such disinhibitory circuits are critical for enabling context-dependent modulation of visual
responses, including task-related enhancement of behaviorally relevant features and suppression
of irrelevant information. In our data, the sharpening of category representations in VTC when
instructions preceded stimuli (Fig. 3) might reflect precisely this type of disinhibitory mechanism:
dACC feedback may selectively release task-relevant VTC neurons from local inhibition, amplifying
their responses while leaving the underlying categorical tuning intact. In addition, feedback could
recruit parvalbumin (PV)-expressing interneurons to sharpen sensory representations through lat-
eral inhibition, as has been observed during attention-demanding tasks in rodents [39, 40]. Testing
these microcircuit hypotheses in humans remains challenging, but future studies combining lami-
nar recordings with computational modeling could help disambiguate the specific inhibitory motifs
underlying the top-down modulation we observed.

The robust dACC-VTC coupling we observed raises important questions about the anatomical
pathways supporting this long-range communication. In non-human primates, anatomical tracing
studies have revealed that anterior cingulate cortex sends direct feedback projections to inferotem-
poral cortex, terminating predominantly in superficial layers where they can modulate the activity
of both excitatory and inhibitory neurons [41, 42]. These feedback projections follow a hierarchical
gradient, with higher-order frontal regions preferentially targeting superficial layers of lower-order
sensory areas, a pattern consistent with predictive coding models in which top-down predictions

modulate sensory processing [43]. In humans, diffusion tensor imaging and tract-tracing studies have similarly identified white matter pathways connecting medial frontal cortex to ventral temporal regions, including connections through the inferior fronto-occipital fasciculus and the cingulum bundle [44, 45]. However, direct monosynaptic connections between dACC and VTC in humans may be sparse, raising the possibility that dACC modulation could also operate through polysynaptic routes involving intermediate relay stations such as the thalamus, posterior cingulate cortex, or lateral prefrontal regions [46–48]. The pulvinar nucleus of the thalamus represents a particularly intriguing candidate, as it receives inputs from prefrontal cortex and projects extensively to temporal visual areas, providing a potential substrate for rapid routing of contextual signals to sensory cortex [37, 46]. The pulvinar's role in coordinating cortical communication through cortico-thalamo-cortical loops could enable dACC to dynamically gate sensory processing in VTC depending on behavioral demands [49, 50]. The specific contribution of monosynaptic versus polysynaptic pathways to the observed spike-field coupling remains an important question for future investigation, potentially addressable through combined recordings with electrical microstimulation or optogenetic perturbations in animal models.

Our findings have important implications for understanding cognitive flexibility and its breakdown in clinical populations. The dynamic reconfiguration of VTC representations we observed suggests that even canonical sensory areas are continuously shaped by top-down goal states rather than operating as passive feature detectors. Critically, the strength of dACC-VTC coupling predicted performance under conditions of high uncertainty and cognitive demand, suggesting that individual differences in this mechanism may underlie variability in cognitive control abilities. Dysfunction in such top-down modulation could contribute to cognitive inflexibility observed in neuropsychiatric disorders including schizophrenia, where patients show impaired context-dependent modulation of sensory processing [51], and autism spectrum disorder, where atypical sensory responses may reflect reduced top-down predictive signaling [52, 53]. Furthermore, neurodegenerative conditions such as Alzheimer's disease, which preferentially affect frontal and temporal cortical regions, as well as age-related decline in prefrontal function, could weaken dACC-VTC coupling and contribute to reduced cognitive flexibility in older adults. Our findings thus provide a mechanistic framework for understanding how goal-directed modulation of sensory cortex supports adaptive behavior, and how

365 disruptions to this mechanism may contribute to cognitive and perceptual deficits across clinical

366 populations.

References

1.   Haxby, J. V. et al. Distributed and overlapping representations of faces and objects in ventral temporal cortex. *Science* **293,** 2425–2430 (Sept. 2001).

2.   Kanwisher, N. Functional specificity in the human brain: a window into the functional architecture of the mind. *Proceedings of the national academy of sciences* **107,** 11163–11170 (June 2010).

3.   Konkle, T. & Caramazza, A. Tripartite organization of the ventral stream by animacy and object size. *Journal of neuroscience* **33,** 10235–10242 (June 2013).

4.   Çukur, T., Nishimoto, S., Huth, A. G. & Gallant, J. L. Attention during natural vision warps semantic representation across the human brain. *Nature neuroscience* **16,** 763–770 (June 2013).

5.   Henderson, M. M., Serences, J. T. & Rungratsameetaweemana, N. Dynamic categorization rules alter representations in human visual cortex. *Nature communications* (2025).

6.   Bugatus, L., Weiner, K. S. & Grill-Spector, K. Task alters category representations in prefrontal but not high-level visual cortex. *Neuroimage* **155,** 437–449 (July 2017).

7.   Margalit, E. et al. Ultra-high-resolution fMRI of Human Ventral Temporal Cortex Reveals Differential Representation of Categories and Domains. *Journal of neuroscience* **40,** 3008–3024 (Apr. 2020).

8.   Fan, X., Wang, F., Shao, H., Zhang, P. & He, S. The bottom-up and top-down processing of faces in the human occipitotemporal cortex. *Elife* **9,** e48764 (Jan. 2020).

9.   Masharipov, R., Knyazeva, I., Korotkov, A., Cherednichenko, D. & Kireev, M. Comparison of whole-brain task-modulated functional connectivity methods for fMRI task connectomics. *Communications biology* **7,** 1402 (Oct. 2024).

10.   Tanaka, K. Inferotemporal cortex and object vision. *Annual review of neuroscience* **19,** 109–139 (1996).

11.   Li, N. & DiCarlo, J. J. Unsupervised natural experience rapidly alters invariant object representation in visual cortex. *Science* **321,** 1502–1507 (2008).

12.   Freiwald, W. A. & Tsao, D. Y. Functional compartmentalization and viewpoint generalization within the macaque face-processing system. *Science* **330,** 845–851 (Nov. 2010).

13. DiCarlo, J. J., Zoccolan, D. & Rust, N. C. How does the brain solve visual object recognition?
Neuron **73,** 415–434 (2012).

14. Bao, P., She, L., McGill, M. & Tsao, D. Y. A map of object space in primate inferotemporal
cortex. en. *Nature* **583,** 103–108 (2020).

15. Ährlund Richter, S. et al. Distinct roles of prefrontal subregion feedback to the primary visual
cortex across behavioral states. *Neuron* (Nov. 2025).

16. Marshel, J. H., Garrett, M. E., Nauhaus, I. & Callaway, E. M. Functional Specialization of
Seven Mouse Visual Cortical Areas. *Neuron* **72,** 1040–1054 (2011).

17. Niell, C. M. & Stryker, M. P. Modulation of Visual Responses by Behavioral State in Mouse
Visual Cortex. *Neuron* **65,** 472–479 (2010).

18. Fiser, A. et al. Experience-dependent spatial expectations in mouse visual cortex. *Nature*
neuroscience **19,** 1658–1664 (Dec. 2016).

19. Liu, D. et al. Orbitofrontal control of visual cortex gain promotes visual associative learning.
Nature communications **11,** 2784 (June 2020).

20. Kirchberger, L. et al. The essential role of recurrent processing for figure-ground perception
in mice. en. *Science advances* **7,** eabe1833 (2021).

21. Keller, A. J. et al. A Disinhibitory Circuit for Contextual Modulation in Primary Visual
Cortex. *Neuron* **108,** 1181–1193.e8 (Dec. 2020).

22. Hajnal, M. A. et al. Continuous multiplexed population representations of task context in the
mouse primary visual cortex. *Nature communications* **14,** 6687 (Oct. 2023).

23. Faulkner, A. D. et al. Context flexibly modulates cue representations in visual cortex. *Nature*
communications **16,** 5516 (2025).

24. Wadia, V. S. et al. A shared code for perceiving and imagining objects in human ventral
temporal cortex. *Biorxiv.* Preprint (2024).

25. Schneider, S., Lee, J. H. & Mathis, M. W. Learnable latent embeddings for joint behavioural
and neural analysis. *Nature* **617,** 360–368 (2023).

26. Aquino, T. G., Cockburn, J., Mamelak, A. N., Rutishauser, U. & O'Doherty, J. P. Neurons in
human pre-supplementary motor area encode key computations for value-based choice. *Nature*
human behaviour **7,** 970–985 (June 2023).

27. Veniero, D. et al. Top-down control of visual cortex by the frontal eye fields through oscillatory
realignment. *Nature communications* **12,** 1757 (Mar. 2021).

28. Bastos, G. et al. Top-down input modulates visual context processing through an interneuron-
specific circuit. *Cell reports* **42,** 113133 (2023).

29. Bastos, A. et al. Visual Areas Exert Feedforward and Feedback Influences through Distinct
Frequency Channels. *Neuron* **85,** 390–401 (2015).

30. Van Kerkoerle, T. et al. Alpha and gamma oscillations characterize feedback and feedforward
processing in monkey visual cortex. *Proceedings of the national academy of sciences* **111,**
14332–14341 (2014).

31. Cavanagh, J. F. & Frank, M. J. Frontal theta as a mechanism for cognitive control. *Trends in*
cognitive sciences **18,** 414–421 (Aug. 2014).

32. Buschman, T. J. & Miller, E. K. Top-down versus bottom-up control of attention in the
prefrontal and posterior parietal cortices. *Science* **315,** 1860–1862 (Mar. 2007).

33. Engel, A. K. & Fries, P. Beta-band oscillations–signalling the status quo? *Current opinion in*
neurobiology **20,** 156–165 (Apr. 2010).

34. Cohen, M. X. A neural microcircuit for cognitive conflict detection and signaling. *Trends in*
neurosciences **37,** 480–490 (Sept. 2014).

35. Arnal, L. H. & Giraud, A.-L. Cortical oscillations and sensory predictions. *Trends in cognitive*
sciences **16,** 390–398 (July 2012).

36. Bastos, A. M. et al. Canonical microcircuits for predictive coding. *Neuron* **76,** 695–711 (Nov.
2012).

37. Fiebelkorn, I. C. & Kastner, S. A Rhythmic Theory of Attention. *Trends in cognitive sciences*
23, 87–101 (Feb. 2019).

38. Han, H.-B., Brincat, S. L., Buschman, T. J. & Miller, E. K. Working memory readout varies
with frontal theta rhythms. *Neuron* (2025).

39. Zhang, S. et al. Selective attention. Long-range and local circuits for top-down modulation of
visual cortex processing. *Science* **345,** 660–665 (Aug. 2014).

40. Kim, H., Ährlund Richter, S., Wang, X., Deisseroth, K. & Carlén, M. Prefrontal Parvalbumin
Neurons in Control of Attention. *Cell* **164,** 208–218 (Jan. 2016).

41. Barbas, H. & Pandya, D. N. Architecture and intrinsic connections of the prefrontal cortex in
the rhesus monkey. *The journal of comparative neurology* **286,** 353–375 (Aug. 1989).

42. Saleem, K. S., Miller, B. & Price, J. L. Subdivisions and connectional networks of the lateral
prefrontal cortex in the macaque monkey. *Journal of comparative neurology* **522,** 1641–1690
(May 2014).

43. Markov, N. T. et al. Anatomy of hierarchy: feedforward and feedback pathways in macaque
visual cortex. *The journal of comparative neurology* **522,** 225–259 (Jan. 2014).

44. Schmahmann, J. D. et al. Association fibre pathways of the brain: parallel observations from
diffusion spectrum imaging and autoradiography. *Brain: a journal of neurology* **130,** 630–653
(Mar. 2007).

45. Catani, M. & Thiebaut de Schotten, M. A diffusion tensor imaging tractography atlas for
virtual in vivo dissections. *Cortex* **44,** 1105–1132 (Sept. 2008).

46. Saalmann, Y. B., Pinsk, M. A., Wang, L., Li, X. & Kastner, S. The pulvinar regulates informa-
tion transmission between cortical areas based on attention demands. *Science* **337,** 753–756
(Aug. 2012).

47. Fiebelkorn, I. C., Pinsk, M. A. & Kastner, S. The mediodorsal pulvinar coordinates the
macaque fronto-parietal network during rhythmic spatial attention. *Nature communications*
10, 215 (Jan. 2019).

48. Mukherjee, A., Lam, N. H., Wimmer, R. D. & Halassa, M. M. Thalamic circuits for indepen-
dent control of prefrontal signal and noise. *Nature* **600,** 100–104 (Dec. 2021).

49. Sherman, S. M. Thalamus plays a central role in ongoing cortical functioning. *Nature neuro-
science* **19,** 533–541 (Apr. 2016).

50. Halassa, M. M. & Kastner, S. Thalamic functions in distributed cognitive control. *Nature
neuroscience* **20,** 1669–1679 (Dec. 2017).

51.   Silverstein, S. M. & Keane, B. P. Perceptual organization impairment in schizophrenia and
associated brain mechanisms: review of research from 2005 to 2010. *Schizophrenia bulletin* **37,**
690–699 (July 2011).

52.   Pellicano, E. & Burr, D. When the world becomes 'too real': a Bayesian explanation of autistic
perception. *Trends in cognitive sciences* **16,** 504–510 (Oct. 2012).

53.   Van de Cruys, S. et al. Precise minds in uncertain worlds: predictive coding in autism. *Psy-*
chological review **121,** 649–675 (Oct. 2014).

54.   Minxha, J., Mamelak, A. N. & Rutishauser, U. in *Extracellular Recording Approaches* (ed
Sillitoe, R. V.) Series Title: Neuromethods, 267–293 (Springer New York, New York, NY,
2018).

55.   Rutishauser, U., Schuman, E. M. & Mamelak, A. N. Online detection and sorting of extracel-
lularly recorded action potentials in human medial temporal lobe recordings, in vivo. *Journal*
of neuroscience methods **154,** 204–224 (June 2006).

56.   Mormann, F. et al. Latency and selectivity of single neurons indicate hierarchical processing
in the human medial temporal lobe. *Journal of neuroscience* **28,** 8865–8872 (2008).

57.   Minxha, J., Adolphs, R., Fusi, S., Mamelak, A. N. & Rutishauser, U. Flexible recruitment
of memory-based choice representations by the human medial frontal cortex. *Science* **368,**
eaba3313 (June 2020).

58.   Fu, Z. et al. The geometry of domain-general performance monitoring in the human medial
frontal cortex. *Science* **376,** eabm9922 (May 2022).

59.   Courellis, H. S. et al. Abstract representations emerge in human hippocampal neurons during
inference. *Nature* **632,** 841–849 (Aug. 2024).

60.   Rutishauser, U., Ross, I. B., Mamelak, A. N. & Schuman, E. M. Human memory strength
is predicted by theta-frequency phase-locking of single neurons. *Nature* **464,** 903–907 (Apr.
2010).

Acknowledgements

We are grateful to the staff of the epilepsy monitoring unit at Cedars-Sinai Medical Center, including the neuromonitoring staff, nursing staff, and epileptologists (Drs. Lisa Bateman and Jeffrey Chung), for their dedicated patient care and clinical support. We thank members of the Rutishauser laboratory, including Jonathan Daume, Hristos Courellis, Varun Wadia, Natasha Kurilenko, Jake Gavenas, Sophia Cheng, and Daniel Deng, for their helpful discussions and feedback on the manuscript. We give special thanks to Varun Wadia and Hristos Courellis for their invaluable assistance with data collection for several patients. We also thank Jorge Aldana for assistance with computing resources. Above all, we are deeply grateful to the patients and their families for their willingness to participate in this research. This work was funded by the American Academy of Neurology (AAN) Resident Research Scholarship (to R.K.), the Simons Foundation's SCGB program (to U.R.), and the BRAIN initiative through U01NS117839 (to U.R.). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Author contributions

N.R. and T.G.A. conceived the study. N.R., U.R. and T.G.A. designed the behavioral task. R.K. and T.G.A. collected the data. R.K. and T.G.A. performed data processing. R.K. performed formal analysis. R.K., N.R., and U.R. wrote the initial draft of the manuscript. T.G.A., C.M.R. and A.N.M. reviewed and edited the manuscript. N.R. and U.R. supervised the study. C.M.R. provided patient care and facilitated experiments. A.N.M. performed all surgical procedures. U.R. acquired funding.

Declaration of interests

The authors declare no competing interests.

Methods

Participants. Eleven adult patients (6 female; age range: 25 - 55 years; see Extended Data Table 1) with drug-resistant epilepsy participated in this study. All patients were undergoing stereotactic depth electrode implantation for seizure localization and potential surgical treatment at Cedars-Sinai Medical Center. Electrode placement was determined solely on clinical grounds by the clinical care team. Each patient provided written informed consent to participate in research protocols approved by the Institutional Review Board of Cedars-Sinai Medical Center.

Task design. Participants performed a flexible visual decision-making task designed to dissociate sensory encoding from task-related contextual modulation. Each participant completed one recording session consisting of four blocks of 48 trials each. The blocks alternated between early and late instruction conditions.

In the early instruction condition, each trial began with a 1-s fixation period followed by a 2.5-s instruction period. Two color images, drawn from one of four visual categories (cars, animals, fruits, and human faces), were then presented sequentially for 1 s each. Based on the instruction, participants indicated whether the first or second image satisfied the rule by pressing a corresponding button. The mapping between the first/second choice and left/right response buttons was randomized across trials to prevent motor preparation biases. In the late instruction condition, each trial also began with a 1-s fixation period but was followed by two sequential images before the 2.5-s instruction period, after which participants made their choice.

There were two types of instructions: pro and anti. In pro trials, participants followed the rule as stated (e.g., "Choose the most colorful car"), whereas in anti trials they applied the opposite rule (e.g., "Do not choose the most colorful car"). Because of their double-negative structure, anti trials imposed greater cognitive demands than pro trials.

All conditions were balanced across blocks such that each image appeared twice, as the first and second stimulus, and was presented equally often under early and late instruction conditions.

Electrophysiology and spike sorting. Intracranial recordings were obtained at Cedars-Sinai Medical Center from bilateral hybrid depth electrodes containing eight microwires each (AdTech Medical). Electrode implantation sites were determined exclusively by clinical needs for seizure

localization and included the ventral temporal cortex (VTC), dorsal anterior cingulate cortex (dACC), pre-supplementary motor area (preSMA), ventromedial prefrontal cortex (vmPFC), hippocampus (HIPP), and amygdala (AMY). See Extended Data Table 1. Broadband extracellular signals (0.1–9,000 Hz) were continuously recorded from each microwire at a sampling rate of 32 kHz using the ATLAS system (Neuralynx Inc.). Recordings were locally referenced within each electrode bundle to a designated low-impedance microwire.

Electrode localization was performed by co-registering postoperative computed tomography (CT) scans with preoperative magnetic resonance imaging (MRI) using standard alignment procedures implemented in FreeSurfer (v5.3.0 and v7.4.1) and Advanced Normalization Tools (ANTs v2.1) [54]. Individual pre-operative MRIs were aligned to the MNI152-space CIT168 probabilistic atlas for visualization of electrode locations. All electorde localization was performed in the subjects native space. Electrode contacts were confirmed to be within gray matter through visual inspection in the native space.

Spike detection and sorting were performed offline using the semi-automated template-matching algorithm OSort (v4.1 [55]). The raw broadband signal was band-pass filtered between 300 and 3,000 Hz, and candidate spikes were detected using an adaptive threshold. Putative single units were isolated based on waveform shape, refractory period violations, and cluster separation. Units that did not meet these isolation criteria were excluded. Only patients with at least one well-isolated single neuron in one of the regions of interest were included in the analyses.

LFP preprocessing. All local field potential (LFP) analyses were performed on broadband signals (0.1 - 9,000 Hz, sampled at 32 kHz) recorded from the microwires of the hybrid depth electrodes. To prevent contamination of low-frequency activity by action-potential (spiking) artifacts, we removed spike waveforms by linearly interpolating the raw signal from -1 to $+2$ ms around each detected spike time on that microwire. This interpolation was applied across the continuous broadband recording prior to downsampling. Because the same spike can occasionally appear on multiple nearby microwires, interpolation was performed across all wires within the same bundle whenever a spike was detected on any one of them.

After spike removal, signals underwent band-stop filtering to remove 60 Hz line noise and its harmonics up to 180 Hz using a zero-phase Butterworth filter (2 Hz bandwidth around each

harmonic). The data were then low-pass filtered at 500 Hz (8th-order Butterworth, zero-phase) and downsampled to 1 kHz using MATLAB (MathWorks, R2023a). All filtering and downsampling steps were implemented using zero-phase (forward-reverse) filtering to avoid phase distortions in the LFP signal.

Following these preprocessing steps, each trial and channel were individually reviewed by visual inspection. Trials containing interictal epileptiform discharges, large-amplitude movement arti-facts, amplifier saturation, or abnormally low signal variance were excluded from further analysis. Channels showing persistent interictal discharges or excessive noise throughout the session were removed. On average, 60.02 ± 69.27 trials (31.3% of the data) were excluded per microwire during this artifact rejection process. Only artifact-free trials and channels were included in subsequent analyses.

Single-neuron selectivity analysis. Neuronal selectivity for stimulus category, instruction tim-ing, and their interaction was quantified using a two-way analysis of variance (ANOVA) performed independently for each neuron. Analyses were conducted separately for responses to the first and second stimuli (stim 1 and stim 2). For VTC neurons, firing rates were calculated within a 0.1 - 1.0 s window following stimulus onset (t=0 is stimulus onset). For non-VTC regions (dorsal ante-rior cingulate cortex, pre-supplementary motor area, ventromedial prefrontal cortex, hippocampus, and amygdala), the analysis window was shifted to 0.2 - 1.0 s after stimulus onset to account for the longer onset latencies in higher-order areas [56].

For each neuron, trial-by-trial firing rates were entered into a two-way 2×4 ANOVA with instruction timing (early vs. late) and stimulus category (cars, animals, fruits, faces) as fixed factors, including the interaction term. Neurons were classified as category-selective or instruction-selective based on significant main effects ($P < 0.05$, uncorrected). Neurons showing a significant main effect of category were labeled category-selective, and those with a significant main effect of instruction timing were labeled instruction-selective. Neurons showing significant main effects for both factors were grouped as jointly selective.

The proportion of neurons in each class was computed for each brain region. For population-level summaries (Fig. 2d and Fig. 4a), neurons were counted once per region, and chance-level proportions (5%) were estimated via shuffling of category and instruction labels across trials.

Population decoding analysis. Population decoding was performed using a pseudo-population approach, in which single-trial firing rates from all isolated neurons recorded within a given brain region were aggregated across participants, as has been done in prior studies [57–59]. This approach allowed us to estimate population-level discriminability despite the limited number of simultaneously recorded neurons per participant. For each neuron, firing rates were averaged within the stimulus analysis window (0.1-1.0 s after stimulus onset for VTC; 0.2-1.0 s for non-VTC regions).

Decoding analyses were implemented in MATLAB (MathWorks, R2023a) using a linear support vector machine (SVM) classifier (fitcecoc function with default parameters). For category decoding, data were balanced across the four stimulus categories (cars, animals, fruits, faces), with four trials per category used for training and four for testing (16 trials total per split). This process was repeated for 100 random iterations, and decoding accuracy was averaged across iterations.

For instruction-order decoding, analyses were conducted separately within each stimulus category to ensure that sensory information was held constant. Within each category, four trials from each instruction condition (early vs. late) were used for training and four for testing (eight trials total per split). Trial assignments were randomized across 100 iterations, and the mean decoding accuracy across iterations was used as the final measure.

All decoding analyses were performed on datasets in which the number of trials was explicitly matched across conditions to avoid bias from unequal trial counts. Decoding performance was evaluated on held-out test data in each iteration, and reported accuracies represent mean cross-validated decoding performance across iterations for each brain region.

For comparing SVM decoding accuracy across brain areas (Fig. 4b,d), the number of neurons used for decoding was fixed to the number of isolated neurons from the region with the smallest sample size (dACC, 74 neurons). For regions containing more than 74 neurons, 74 neurons were randomly subsampled at each iteration to ensure matched population sizes across regions.

State-space analysis using CEBRA. To visualize the low-dimensional structure of population activity, we used Consistent Embeddings of high-dimensional Recordings using Auxiliary variables (CEBRA; [25]). Unlike linear dimensionality reduction methods such as principal component analysis (PCA), CEBRA learns nonlinear latent embeddings that preserve the temporal continuity and intrinsic geometry of neural population dynamics.

We used an unsupervised version of CEBRA, fitting the model exclusively to neural activity

without auxiliary behavioral or task labels. The model was trained on concatenated single-trial

firing rates from all neurons within each region, using the same stimulus-aligned analysis windows

defined above. Prior to training, we performed a grid search to identify optimal model parameters

(output dimensions, time offsets, hidden units, and temperature) that minimized the InfoNCE

contrastive loss. The final model was trained using the "offset10-model" architecture with a batch

size of 1024, learning rate of 3×10^{-4}, temperature of 1.1, and cosine distance metric. The model

was conditioned on time with a time offset of 5, and trained for up to 3000 iterations with 64

hidden units and an output dimensionality of 6. All training was performed using the official

CEBRA Python package (v0.3.0) with GPU acceleration enabled when available.

Each trained model produced low-dimensional embeddings for every trial, which were used to

visualize neural trajectories for early and late instruction conditions. To quantify the separation

between trajectories, we computed the Euclidean distance between the mean latent trajectories of

the two conditions at each time point. These distances provided a measure of representational

divergence in the latent space, with shuffled-label controls used to confirm that observed trajectory

differences were above chance. To ensure that our results were not dependent on a single model

initialization, we trained 10 independent CEBRA models using identical hyperparameters but dif-

ferent random seeds. For each brain region, low-dimensional trajectories and Euclidean distance

time courses of early and late instruction conditions were computed separately within each trained

model and then averaged across models to obtain the final trajectory estimates and distance curves.

Spike-field coherence. To quantify the strength of phase coupling between neuronal spiking

activity and LFPs, we computed spike–field coherence (SFC) [60] during the stim 1 window (0.1–

1.0 s after stimulus onset for VTC; 0.2–1.0 s for non-VTC regions). Analyses were restricted to

neuron–LFP channel pairs recorded within the same hemisphere.

Instantaneous phase information from LFPs was extracted using complex Morlet wavelets span-

ning 2 - 150 Hz in 40 logarithmically spaced frequency steps. The number of cycles for each wavelet

increased logarithmically from 3 to 10 across the frequency range to ensure adequate spectral and

temporal resolution. For each neuron-LFP pair, we computed the phase of the LFP at each spike

time and quantified SFC as the mean vector length (MVL) of the spike-phase distribution across

all spikes and trials.

To control for differences in trial counts and spike rate across conditions, the number of trials
and the spike-time distribution were matched between the two conditions being compared (e.g.,
early vs. late instruction, correct vs. incorrect, pro vs. anti). For each matched dataset, SFC was
computed using the procedure described above. This process was repeated 100 iterations, and the
mean SFC across iterations was used as the condition-specific estimate.

To assess statistical significance, we generated a null distribution of SFC values by shuffling
spike times within each trial while preserving the overall firing rate profile. For each iteration,
spike times during the stim 1 window were shuffled, and SFC was recomputed using the same
procedure. This shuffling process was repeated 100 times, and the resulting surrogate distribution
was used to compute a z-scored SFC for the oringinal data.

Statistics. Throughout the study, statistical analyses were performed using non-parametric meth-
ods unless otherwise stated. Pairwise comparisons between independent samples used the Wilcoxon
rank-sum test, and paired comparisons used the Wilcoxon signed-rank test. All tests were two-
sided unless noted, and $P < 0.05$ was considered significant. When applicable, trial counts and
spike numbers were matched across conditions to avoid biases arising from unequal sample sizes.
Additional task- or analysis-specific statistical procedures (e.g., ANOVA for single-neuron selectiv-
ity, Kruskal–Wallis tests with Dunn's correction for comparing decoding accuracy across regions,
and cluster-based bootstrap testing for time-resolved effects) are described below and in the corre-
sponding sections.

Cluster-based bootstrap testing. To identify time periods showing significant differences between
two conditions, we used a nonparametric cluster-based bootstrap procedure. For each comparison,
we first computed the observed time-pointwise difference between conditions. We then generated
a null distribution by randomly permuting condition labels across trials (or across neurons for
pseudo-population analyses) and recomputing the time-pointwise difference on each iteration (1,000
bootstrap iterations). Significant time points were identified by thresholding the observed difference
against the 95th percentile of the null distribution (one-sided). Contiguous suprathreshold time
points were grouped into clusters, and a cluster-level statistic was computed as the sum of the
absolute differences within each cluster. The same procedure was applied to each bootstrap iteration

to obtain a null distribution of cluster-level statistics. Clusters in the observed data whose cluster

statistic exceeded the 95th percentile of the bootstrap null distribution were considered significant

($P < 0.05$, cluster-level corrected). All significant clusters are indicated in figures by thick horizontal

black bars.

Code Availability. The code for the analyses performed in this work will be made publicly avail-

able upon acceptance.

Data Availability. Data used in this study will be made publicly available upon acceptance.

Extended Data Tables

Session	Gender	Age	Seizure onset zone	VTC	HIPP	AMY	preSMA	dACC	vmPFC
P81cs	F	28	Left mesial temporal	25 (7, 18)	9 (1, 8)	38 (18, 20)	33 (16, 17)	9 (9, 0)	22 (6, 16)
P82cs	M	42	Bilateral mesial temporal	0	15 (2, 13)	18 (14, 4)	5 (5, 0)	0	1 (1, 0)
P86cs	F	39	Not localized	37 (23, 14)	1 (0, 1)	19 (3, 16)	25 (20, 5)	18 (7, 11)	16 (5, 11)
P90cs	M	32	Left mesial temporal	1 (0, 1)	17 (14, 3)	6 (5, 1)	3 (3, 0)	13 (5, 8)	11 (11, 0)
P92cs	F	30	Left mesial occipital	17 (8, 9)	17 (4, 13)	12 (6, 6)	22 (8, 14)	3 (3, 0)	13 (0, 13)
P97cs	M	32	Right mesial temporal	3 (0, 3)	12 (7, 5)	10 (5, 5)	4 (0, 4)	1 (1, 0)	7 (1, 6)
P98cs	F	30	Bilateral mesial temporal	21 (15, 6)	22 (22, 0)	4 (4, 0)	8 (4, 4)	0	18 (16, 2)
P99cs	F	39	Left lateral temporal	2 (0, 2)	13 (2, 11)	1 (1, 0)	17 (13, 4)	6 (6, 0)	7 (4, 3)
P100cs	M	25	Right mesial temporal	18 (3, 15)	6 (0, 6)	12 (4, 8)	18 (15, 3)	0	12 (10, 2)
P102cs	F	55	Right mesial temporal	32 (16, 16)	15 (10, 5)	12 (7, 5)	20 (4, 16)	15 (4, 11)	11 (11, 0)
P103cs	M	51	Bilateral mesial temporal	8 (6, 2)	10 (2, 8)	11 (7, 4)	30 (18, 12)	0	13 (9, 4)

Extended Data Table 1 | Summary of patient demographics and single-neuron recordings across brain regions. Each entry indicates the total number of well-isolated neurons obtained from spike sorting. Values in parentheses correspond to left and right hemisphere counts, respectively.

Pro condition	Anti condition
Choose the older car	Do not select the older car
Select the image with fewer objects	Do not select the image with fewer objects
Choose the more colorful animal	Avoid the more colorful animal
Select the less modern vehicle	Do not select the less modern vehicle
Choose the image with more animals	Do not select the image with more animals
Select the vehicle with more colors	Do not select the vehicle with more colors
Choose the younger face	Avoid the younger face
Select the creature with fewer colors	Do not select the creature with fewer colors
Choose the image with more creatures	Do not select the image with more creatures
Select the more modern vehicle	Do not select the more modern vehicle
Choose the image with fewer items	Do not select the image with fewer items
Select the more colorful car	Avoid the more colorful car

Extended Data Table 2 | Example task instructions for Pro and Anti conditions.

Stim 1					
Region 1	Region 2	Mean diff	CI low	CI high	*P*-value
VTC	HIPP	-183.61	-112.28	-40.94	0.0001
VTC	AMY	-34.14	37.20	108.53	0.8693
VTC	preSMA	-228.67	-157.33	-85.99	< 0.0001
VTC	dACC	-353.24	-281.91	-210.57	< 0.0001
VTC	vmPFC	-142.86	-71.53	-0.19	0.0488
HIPP	AMY	78.13	149.48	220.89	< 0.0001
HIPP	preSMA	-116.39	-45.06	26.28	0.6310
HIPP	dACC	-240.97	-169.63	-98.29	< 0.0001
HIPP	vmPFC	-29.58	40.75	112.09	0.7724
AMY	preSMA	-194.37	-124.55	-63.28	< 0.0001
AMY	dACC	-265.86	-194.56	-123.17	< 0.0001
AMY	vmPFC	-80.95	-8.72	37.28	0.0001
preSMA	dACC	-189.46	-120.18	-37.38	0.0001
preSMA	vmPFC	14.47	85.85	157.14	0.0064
dACC	vmPFC	139.04	210.38	281.72	< 0.0001
Stim 2					
Region 1	Region 2	Mean diff	CI low	CI high	*P*-value
VTC	HIPP	-353.90	-282.41	-210.91	< 0.0001
VTC	AMY	-448.01	-376.56	-305.03	< 0.0001
VTC	preSMA	-198.15	-126.66	-55.16	< 0.0001
VTC	dACC	-425.43	-353.94	-282.45	< 0.0001
VTC	vmPFC	-222.03	-150.54	-79.04	< 0.0001
HIPP	AMY	165.61	94.12	22.62	0.0017
HIPP	preSMA	-84.26	155.75	227.24	< 0.0001
HIPP	dACC	-143.03	-71.54	-0.04	0.0497
HIPP	vmPFC	160.37	31.87	103.36	< 0.0001
AMY	preSMA	178.37	249.87	321.36	0.9936
AMY	dACC	-48.91	22.76	94.07	< 0.0001
AMY	vmPFC	154.49	225.98	297.47	< 0.0001
preSMA	dACC	-298.78	-227.29	-155.79	< 0.0001
preSMA	vmPFC	-95.38	-23.89	47.61	0.9974
dACC	vmPFC	131.91	203.40	274.89	< 0.0001

Extended Data Table 3 | Pairwise post-hoc comparisons of instruction-order decoding accuracy across regions (related to Extended Data Fig. 4). Differences across the six regions were assessed using a Kruskal-Wallis nonparametric ANOVA, followed by Dunn's multiple comparisons test. The table reports all pairwise contrasts for both the stim 1 and stim 2 windows, including group indices, mean rank differences, confidence intervals, and adjusted *P*-values.

Region 1	Region 2	CI low	Mean diff	CI high	P-value
VTC Early	VTC Late	-41.7522	121.3750	284.5022	0.5610
VTC Early	HIPP Early	243.6328	406.7600	569.8872	< 0.0001
VTC Early	HIPP Late	145.7528	308.8800	472.0072	< 0.0001
VTC Early	AMY Early	216.2428	379.3700	542.4972	< 0.0001
VTC Early	AMY Late	444.2728	607.4000	770.5272	< 0.0001
VTC Early	preSMA Early	342.4028	505.5300	668.6572	< 0.0001
VTC Early	preSMA Late	542.4128	705.5400	868.6672	< 0.0001
VTC Early	dACC Early	527.2078	690.3350	853.4622	< 0.0001
VTC Early	dACC Late	472.2078	635.3350	798.4622	< 0.0001
VTC Early	vmPFC Early	447.9878	611.1150	774.2422	< 0.0001
VTC Early	vmPFC Late	226.8328	389.9600	553.0872	< 0.0001
VTC Late	HIPP Early	122.2578	285.3850	448.5122	< 0.0001
VTC Late	HIPP Late	24.3778	187.5050	350.6322	0.0074
VTC Late	AMY Early	94.8678	257.9950	421.1222	< 0.0001
VTC Late	AMY Late	322.8978	486.0250	649.1522	< 0.0001
VTC Late	preSMA Early	221.0278	384.1550	547.2822	< 0.0001
VTC Late	preSMA Late	421.0378	584.1650	747.2922	< 0.0001
VTC Late	dACC Early	405.8328	568.9600	732.0872	< 0.0001
VTC Late	dACC Late	350.8328	513.9600	677.0872	< 0.0001
VTC Late	vmPFC Early	326.6128	489.7400	652.8672	< 0.0001
VTC Late	vmPFC Late	105.4578	268.5850	431.7122	< 0.0001
HIPP Early	HIPP Late	-261.0072	-97.8800	65.2472	0.9477
HIPP Early	AMY Early	-190.5172	-27.3900	135.7372	1.0000
HIPP Early	AMY Late	37.5128	200.6400	363.7672	0.0024
HIPP Early	preSMA Early	-64.3572	98.7700	261.8972	0.9405
HIPP Early	preSMA Late	135.6528	298.7800	461.9072	< 0.0001
HIPP Early	dACC Early	120.4478	283.5750	446.7022	< 0.0001
HIPP Early	dACC Late	65.4478	228.5750	391.7022	0.0002
HIPP Early	vmPFC Early	41.2278	204.3550	367.4822	0.0017
HIPP Early	vmPFC Late	-179.9272	-16.8000	146.3272	1.0000
HIPP Late	AMY Early	-92.6372	70.4900	233.6172	1.0000
HIPP Late	AMY Late	135.3928	298.5200	461.6472	< 0.0001
HIPP Late	preSMA Early	33.5228	196.6500	359.7772	0.0034
HIPP Late	preSMA Late	233.5328	396.6600	559.7872	< 0.0001
HIPP Late	dACC Early	218.3278	381.4550	544.5822	< 0.0001
HIPP Late	dACC Late	163.3278	326.4550	489.5822	< 0.0001
HIPP Late	vmPFC Early	139.1078	302.2350	465.3622	< 0.0001
HIPP Late	vmPFC Late	-82.0472	81.0800	244.2072	0.9986
AMY Early	AMY Late	64.9028	228.0300	391.1572	0.0002
AMY Early	preSMA Early	-36.9672	126.1600	289.2872	0.4618

Region 1	Region 2	CI low	Mean diff	CI high	*P*-value
AMY Early	preSMA Late	163.0428	326.1700	489.2972	< 0.0001
AMY Early	dACC Early	147.8378	310.9650	474.0922	< 0.0001
AMY Early	dACC Late	92.8378	255.9650	419.0922	< 0.0001
AMY Early	vmPFC Early	68.6178	231.7450	394.8722	0.0001
AMY Early	vmPFC Late	-152.5372	10.5900	173.7172	1.0000
AMY Late	preSMA Early	-264.9972	-101.8700	61.2572	0.9100
AMY Late	preSMA Late	-64.9872	98.1400	261.2672	0.9457
AMY Late	dACC Early	-80.1922	82.9350	246.0622	0.9976
AMY Late	dACC Late	-135.1922	27.9350	191.0622	1.0000
AMY Late	vmPFC Early	-159.4122	3.7150	166.8422	1.0000
AMY Late	vmPFC Late	-380.5672	-217.4400	-54.3128	0.0005
preSMA Early	preSMA Late	36.8828	200.0100	363.1372	0.0025
preSMA Early	dACC Early	21.6778	184.8050	347.9322	0.0092
preSMA Early	dACC Late	-33.3222	129.8050	292.9322	0.3910
preSMA Early	vmPFC Early	-57.5422	105.5850	268.7122	0.8624
preSMA Early	vmPFC Late	-278.6972	-115.5700	47.5572	0.6831
preSMA Late	dACC Early	-178.3322	-15.2050	147.9222	1.0000
preSMA Late	dACC Late	-233.3322	-70.2050	92.9222	1.0000
preSMA Late	vmPFC Early	-257.5522	-94.4250	68.7022	0.9700
preSMA Late	vmPFC Late	-478.7072	-315.5800	-152.4528	< 0.0001
dACC Early	dACC Late	-218.1272	-55.0000	108.1272	1.0000
dACC Early	vmPFC Early	-242.3472	-79.2200	83.9072	0.9992
dACC Early	vmPFC Late	-463.5022	-300.3750	-137.2478	< 0.0001
dACC Late	vmPFC Early	-187.3472	-24.2200	138.9072	1.0000
dACC Late	vmPFC Late	-408.5022	-245.3750	-82.2478	< 0.0001
vmPFC Early	vmPFC Late	-384.2822	-221.1550	-58.0278	0.0003

Extended Data Table 4 | Pairwise Dunn-test comparisons across early/late conditions in all regions during stim 1 window (related to Extended Data Fig. 5a). Each row lists a pairwise contrast between regional conditions (e.g., VTC Early vs. HIPP Late), including the confidence-interval lower bound, mean rank difference, confidence-interval upper bound, and Dunn corrected *P*-value.

Region 1	Region 2	CI low	Mean diff	CI high	P-value
VTC Early	VTC Late	-263.8578	-100.6350	62.5878	0.9238
VTC Early	HIPP Early	282.3722	445.5950	608.8178	< 0.0001
VTC Early	HIPP Late	275.4422	438.6650	601.8878	< 0.0001
VTC Early	AMY Early	280.2522	443.4750	606.6978	< 0.0001
VTC Early	AMY Late	-20.4678	142.7550	305.9778	0.1954
VTC Early	preSMA Early	159.9972	323.2200	486.4428	< 0.0001
VTC Early	preSMA Late	446.9772	610.2000	773.4228	< 0.0001
VTC Early	dACC Early	399.0022	562.2250	725.4478	< 0.0001
VTC Early	dACC Late	182.1872	345.4100	508.6328	< 0.0001
VTC Early	vmPFC Early	368.1972	531.4200	694.6428	< 0.0001
VTC Early	vmPFC Late	214.2872	377.5100	540.7328	< 0.0001
VTC Late	HIPP Early	383.0072	546.2300	709.4528	< 0.0001
VTC Late	HIPP Late	376.0772	539.3000	702.5228	< 0.0001
VTC Late	AMY Early	380.8872	544.1100	707.3328	< 0.0001
VTC Late	AMY Late	80.1672	243.3900	406.6128	< 0.0001
VTC Late	preSMA Early	260.6322	423.8550	587.0778	< 0.0001
VTC Late	preSMA Late	547.6122	710.8350	874.0578	< 0.0001
VTC Late	dACC Early	499.6372	662.8600	826.0828	< 0.0001
VTC Late	dACC Late	282.8222	446.0450	609.2678	< 0.0001
VTC Late	vmPFC Early	468.8322	632.0550	795.2778	< 0.0001
VTC Late	vmPFC Late	314.9222	478.1450	641.3678	< 0.0001
HIPP Early	HIPP Late	-170.1528	-6.9300	156.2928	1.0000
HIPP Early	AMY Early	-165.3428	-2.1200	161.1028	1.0000
HIPP Early	AMY Late	-466.0628	-302.8400	-139.6172	< 0.0001
HIPP Early	preSMA Early	-285.5978	-122.3750	40.8478	0.5414
HIPP Early	preSMA Late	1.3822	164.6050	327.8278	0.0452
HIPP Early	dACC Early	-46.5928	116.6300	279.8528	0.6626
HIPP Early	dACC Late	-263.4078	-100.1850	63.0378	0.9282
HIPP Early	vmPFC Early	-77.3978	85.8250	249.0478	0.9950
HIPP Early	vmPFC Late	-231.3078	-68.0850	95.1378	1.0000
HIPP Late	AMY Early	-158.4128	4.8100	168.0328	1.0000
HIPP Late	AMY Late	-459.1328	-295.9100	-132.6872	< 0.0001
HIPP Late	preSMA Early	-278.6678	-115.4450	47.7778	0.6871
HIPP Late	preSMA Late	8.3122	171.5350	334.7578	0.0269
HIPP Late	dACC Early	-39.6628	123.5600	286.7828	0.5166
HIPP Late	dACC Late	-256.4778	-93.2550	69.9678	0.9758
HIPP Late	vmPFC Early	-70.4678	92.7550	255.9778	0.9779
HIPP Late	vmPFC Late	-224.3778	-61.1550	102.0678	1.0000
AMY Early	AMY Late	-463.9428	-300.7200	-137.4972	< 0.0001
AMY Early	preSMA Early	-283.4778	-120.2550	42.9678	0.5862

Region 1	Region 2	CI low	Mean diff	CI high	*P*-value
AMY Early	preSMA Late	3.5022	166.7250	329.9478	0.0386
AMY Early	dACC Early	-44.4728	118.7500	281.9728	0.6181
AMY Early	dACC Late	-261.2878	-98.0650	65.1578	0.9467
AMY Early	vmPFC Early	-75.2778	87.9450	251.1678	0.9918
AMY Early	vmPFC Late	-229.1878	-65.9650	97.2578	1.0000
AMY Late	preSMA Early	17.2422	180.4650	343.6878	0.0133
AMY Late	preSMA Late	304.2222	467.4450	630.6678	< 0.0001
AMY Late	dACC Early	256.2472	419.4700	582.6928	< 0.0001
AMY Late	dACC Late	39.4322	202.6550	365.8778	0.0020
AMY Late	vmPFC Early	225.4422	388.6650	551.8878	< 0.0001
AMY Late	vmPFC Late	71.5322	234.7550	397.9778	0.0001
preSMA Early	preSMA Late	123.7572	286.9800	450.2028	< 0.0001
preSMA Early	dACC Early	75.7822	239.0050	402.2278	0.0001
preSMA Early	dACC Late	-141.0328	22.1900	185.4128	1.0000
preSMA Early	vmPFC Early	44.9772	208.2000	371.4228	0.0012
preSMA Early	vmPFC Late	-108.9328	54.2900	217.5128	1.0000
preSMA Late	dACC Early	-211.1978	-47.9750	115.2478	1.0000
preSMA Late	dACC Late	-428.0128	-264.7900	-101.5672	< 0.0001
preSMA Late	vmPFC Early	-242.0028	-78.7800	84.4428	0.9993
preSMA Late	vmPFC Late	-395.9128	-232.6900	-69.4672	0.0001
dACC Early	dACC Late	-380.0378	-216.8150	-53.5922	0.0005
dACC Early	vmPFC Early	-194.0278	-30.8050	132.4178	1.0000
dACC Early	vmPFC Late	-347.9378	-184.7150	-21.4922	0.0094
dACC Late	vmPFC Early	22.7872	186.0100	349.2328	0.0084
dACC Late	vmPFC Late	-131.1228	32.1000	195.3228	1.0000
vmPFC Early	vmPFC Late	-317.1328	-153.9100	9.3128	0.0961

Extended Data Table 5 | Pairwise Dunn-test comparisons across early/late conditions in all regions during stim 2 window (related to Extended Data Fig. 5b). Each row lists a pairwise contrast between regional conditions (e.g., VTC Early vs. HIPP Late), including the confidence-interval lower bound, mean rank difference, confidence-interval upper bound, and Dunn corrected *P*-value.

Extended Data Figures

Extended Data Fig. 1 | Behavioral performance for early versus late instruction trials within pro and anti conditions. a, Pro task. Task accuracy (left) and reaction time (right) shown for early and late instruction trials. Each gray line represents an individual participant, and colored points denote group means (blue: early; red: late). Accuracy did not differ between early and late pro trials. However, reaction times were significantly faster for early compared to late pro trials ($P < 0.05$, one-sided Wilcoxon signed-rank test). **b**, Anti task. Task accuracy (left) and reaction time (right) for early versus late instruction trials. Neither accuracy nor reaction time differed significantly between early and late anti trials.

Extended Data Fig. 2 | Subset of category-selective VTC neurons show instruction-dependent firing-rate modulation. a, average firing rates (mean ± s.e.m.) across the 7 VTC category-selective neurons (from 3 participants) that were identified, using a bootstrap null distribution of preferred-versus-non-preferred firing-rate differences, as exhibiting significantly elevated activity in the instruction window during early-instruction trials. **b**, average firing rates across the 13 VTC category-selective neurons (from 5 participants) that met the same bootstrap-null criterion for the late-instruction condition. Between the two instruction-timing conditions, 5 neurons contributed to both groups. Thick horizontal black bars indicate time periods of significant preferred-versus-non-preferred differences based on the cluster-based bootstrap procedure ($P < 0.05$, cluster-level corrected). Vertical dashed lines mark instruction onset and offset, respectively.

Extended Data Fig. 3 | Category-, instruction-, and jointly selective neurons in additional higher-order regions. Proportion of neurons exhibiting selectivity for stimulus category (Cat; magenta), instruction timing (Instr; teal), or both factors (Cat+Instr; orange) based on a two-way ANOVA (category × instruction timing) in (**a**) amygdala (AMY), (**b**) ventromedial prefrontal cortex (vmPFC), and (**c**) pre-supplementary motor area (preSMA). Dashed line indicates the 5% chance level.

Extended Data Fig. 4 | Instruction-order decoding accuracy across brain regions. Violin plots show linear SVM decoding accuracy for distinguishing early versus late instruction trials during the stim 1 (**a**) and stim 2 (**b**) windows across six regions: ventral temporal cortex (VTC), hippocampus (HIPP), amygdala (AMY), pre-supplementary motor area (preSMA), dorsal anterior cingulate cortex (dACC), and ventromedial prefrontal cortex (vmPFC). White dots denote median decoding accuracy, and the dashed horizontal line indicates chance level (50%). Kruskal–Wallis tests were performed across regions, followed by pairwise post-hoc Dunn's multiple comparisons tests to identify significant differences (results reported in Extended Data Table 3.

Extended Data Fig. 5 | Category decoding accuracy across brain regions for stim 1 and stim 2 windows. Violin plots show linear SVM decoding accuracy for decoding stimulus category during the stim 1 (**a**) and stim 2 (**b**) windows across six regions: ventral temporal cortex (VTC), hippocampus (HIPP), amygdala (AMY), pre-supplementary motor area (preSMA), dorsal anterior cingulate cortex (dACC), and ventromedial prefrontal cortex (vmPFC). Early and late instruction trials are plotted separately for each region. White dots denote median decoding accuracy, and the dashed horizontal line marks chance level (25%). Kruskal–Wallis tests were performed across regions, followed by pairwise post-hoc Dunn's multiple comparisons tests to identify significant differences (results reported in Extended Data Table 4–5).

Extended Data Fig. 6 | Spike–field coherence (SFC) differences between early and late instruction trials across VTC–network pairs. a-b, Stim 1 window. Population-averaged SFC (z-scored) is shown for early (dark blue) and late (dark red) instruction trials for all VTC-involving pairs, plotted separately for VTC as the LFP region (**a**) and VTC as the spike region (**b**). **c-d**, Stim 2 window. Same analyses as in **a** and **b**, computed during the stim 2 window. Shaded regions denote mean ± s.e.m. across neurons. Thick horizontal black bars indicate time periods of significant preferred-versus-non-preferred differences based on the cluster-based bootstrap procedure ($P < 0.05$, cluster-level corrected).

Extended Data Fig. 7 | Spike–field coherence (SFC) differences between correct and incorrect trials for VTC LFP and dACC spikes. SFC (z-scored) is shown for dACC spikes phase-locked to VTC LFP signals during early (**a**) and late (**b**) instruction trials. Orange and blue traces denote correct and incorrect trials, respectively; shaded regions indicate mean ± s.e.m. Thick horizontal black bars indicate time periods of significant preferred-versus-non-preferred differences based on the cluster-based bootstrap procedure ($P < 0.05$, cluster-level corrected).