A chromosome-scale, haplotype-resolved genome assembly of the barley stripe rust

pathogen *Puccinia striiformis* f. sp. *hordei*

Rita Tam[1], Shideh Mojerlou[2], Mareike Möller[1], John P. Rathjen[1], Benjamin Schwessinger[1],

Julian Rodriguez-Algaba[2*]

*= corresponding author e-mail: julianr.algaba@agro.au.dk

[1] Research School of Biology, Australian National University, Canberra, Australia.

[2] Department of Agroecology, Faculty of Sciences and Technology, Aarhus University, Slagelse, Denmark.

Abstract

Puccinia striiformis f. sp. *hordei* (*Psh*) causes barley stripe rust, an economically important

disease affecting barley production across multiple temperate regions. Unlike its well-studied

wheat-infecting relative *P. striiformis* f. sp. *tritici*, genomic resources for *Psh* remain limited,

with no fully documented chromosome-scale, haplotype-resolved reference genomes publicly

available. Here we present a high-quality, haplotype-resolved genome assembly of the

dikaryotic *Psh* isolate NP85002, generated using Oxford Nanopore long-read sequencing

combined with Hi-C chromatin conformation capture. The assembly comprises 18

chromosomes per haplotype (151.3 Mb total) including 33 telomere-to-telomere

chromosomes, telomeric repeats at 69/72 chromosome ends, and six internal gaps. The

assembly shows high consensus accuracy (QV >72) and strong haplotype separation (97.54%

within-haplotype Hi-C contacts). Genome completeness reached 90.1% complete BUSCOs,

and a combined lift-over plus *ab initio* annotation achieved 94.4% complete BUSCOs. Repeat

annotation indicates 45% repetitive content. We additionally provide a 102,058 bp

mitochondrial genome assembly with 41 annotated genes. This genome resource provides a

chromosome-scale framework for comparative and population genomic analyses of barley

stripe rust and related cereal rust pathogens.

1. Background & Summary

Puccinia striiformis f. sp. *hordei* (*Psh*), the causal agent of barley stripe rust, is an economically

important fungal pathogen affecting barley production across multiple temperate regions,

where epidemics result in substantial yield and grain quality losses[1,2]. The disease poses

particular challenges in South Asia, East Africa, and parts of Central and North America, where

Psh causes high disease pressure and resistance breakdown is recurrent[2-4]. As a key crop for

food, feed, and malting industries[5], barley requires durable rust control, making *Psh* a priority

target for resistance breeding and integrated crop protection efforts[2,6].

Psh belongs to the *P. striiformis* species complex, a group of obligate biotrophic rust fungi

comprising host-adapted lineages specialized to different cereal species[4,7]. These fungi display

a complex macrocyclic, heteroecious life cycle that requires two botanically distinct hosts,

with haploid (n), dikaryotic (n+n), and diploid (2n) nuclear phases occurring across the sexual

and asexual stages[8,9]. The rapid evolution of cereal rusts through sexual recombination, clonal
propagation, and somatic hybridization (nuclear exchange without meiosis) coupled with
their long-distance dispersal capacity, enables frequent breakdown of host resistance and
poses persistent threats to global cereal production[10-12]. The epidemic phase on cereals is
dominated by dikaryotic urediniospores that contain two haploid nuclei, which can differ
markedly in sequence composition and gene content[13,14]. This divergence between the two
haplotypes often leads to fragmented representations in genome assembly approaches.
Comparative studies between wheat stripe rust (*P. striiformis* f. sp. *tritici*, *Pst*) and *Psh* indicate
that host adaptation within the *P. striiformis* complex involves gene gain and loss, structural
variation, and changes in effector repertoires[15,16]. Many underlying genomic features of
barley-adapted lineages remain poorly characterized, particularly regarding chromosome-
level organization, telomere structure, and detailed haplotype-specific content[15]. Over the
last decade, several high-quality genome assemblies have been generated for *Pst* including
fully haplotype-resolved, chromosome-scale references based on long-read sequencing[17-19].
These resources are transforming our understanding of inter-haplotype diversity, structural
variation and candidate avirulence genes in wheat stripe rust, providing a robust framework
for population genomics and functional studies[20]. In contrast, fully documented
chromosome-scale, haplotype-resolved genome assemblies for *Psh* remain absent.
Currently, only a single PacBio contig-level assembly representing a collapsed haplotype and
lacking chromosome-scale scaffolding is available in public repositories[16]. Although recent
efforts to generate additional *Psh* assemblies have begun, fully documented chromosome-
scale, haplotype-resolved references are not yet publicly accessible[21]. High-quality,
haplotype-resolved reference genomes for *Psh* are therefore essential for reference-based
population resequencing and association studies targeting barley stripe rust resistance, and
for enabling chromosome-level comparative and evolutionary analyses across cereal rust
pathogens.
Here, we present a haplotype-resolved, chromosome-scale, and near telomere-to-telomere
(T2T) genome assembly of a dikaryotic *Psh* isolate generated using Oxford Nanopore
Technologies (ONT) long-read sequencing combined with chromatin conformation capture
(Hi-C). We additionally provide a circular mitochondrial genome assembly. The assembly is
complemented by repeat annotation, combining lift-over and *ab initio* gene prediction, and
extensive quality-control analyses, providing a valuable genomic framework for future studies
of *Psh* population biology, host adaptation, and effector diversity within the *P. striiformis*
species complex.
75
2. Methods
Isolate selection and propagation
Puccinia striiformis f. sp. *hordei* isolate NP85002 was selected from historical collections at the
Global Rust Reference Center (GRRC), Aarhus University (Denmark) and was originally

collected from infected barley in Nepal in 1985. This isolate provides a valuable temporal and
geographic reference point for comparative genomics, preceding recent barley breeding
efforts and documented pathogen population shifts. Fresh urediniospores were propagated
on the susceptible barley cultivar Afzal under controlled greenhouse conditions (17±2 °C, 16
h photoperiod, 70-80% relative humidity). Seedlings at the two-leaf stage were spray-
inoculated with urediniospore suspensions in Novec™ 7100 fluid (3M™), incubated in a dew
chamber at 10 °C with >95% relative humidity for 24 h in darkness, and subsequently
transferred to spore-proof greenhouse cabins[22]. Fresh urediniospores were harvested 15-20
days post-inoculation and stored at -80°C.
89

DNA extraction and long-read genome sequencing
High-molecular-weight genomic DNA was extracted from 300 mg fresh urediniospores
following the protocol described in[23]. Long-read sequencing was performed using Oxford
Nanopore Technologies (ONT) on a PromethION platform. An ONT library was prepared using
the Ligation Sequencing Kit v14 (SQK-LSK114) and sequenced on a single R10.4.1 flow cell with
Q20+ chemistry. Raw signals were basecalled with Dorado (v0.7.2) using the super-accuracy
model dna_r10.4.1_e8.2_400bps_sup@v5.0.0 to generate simplex reads. Base called reads
were evaluated with NanoPlot v1.44.1 and seqkit v2.10.0, yielding 985,067 reads and a total
of 9.51 Gb, with a mean read length of ~9.7 kb, and read-length N50 of ~19.2 kb. These data
correspond to ~60× coverage of the estimated 160 Mb diploid genome and were used for
downstream *de novo* assembly.
101

Hi-C sequencing
Fresh urediniospores (~100 mg) were cross-linked in 1% formaldehyde at room temperature
for 20 min with periodic mixing and quenched with 125 mM glycine, then washed twice in 1×
PBS and pelleted, following the Phase Genomics Proximo Hi-C (Fungal) sample preparation
guidelines. Cross-linked pellets were frozen and shipped on dry ice to Phase Genomics
(Seattle, WA, USA). Hi-C libraries were prepared using the Proximo Hi-C (Fungal) Kit (KT6040,
Protocol v4.0) with restriction enzymes DpnII, HinfI, MseI, and DdeI and sequenced on an
Illumina NovaSeq 6000 (2 × 150 bp), generating 77.6 million read pairs (~23.3 Gb),
corresponding to ~145× physical coverage for chromatin contact mapping.
111

Genome size estimation
Prior to genome assembly, basic genomic parameters were estimated from k-mer frequency
analysis of high-quality ONT reads. K-mer counting was performed using Jellyfish v2.2.10 with
k = 21, and the resulting k-mer frequency histogram was analyzed with GenomeScope v2.0
[24,25]. The analysis yielded an estimated haploid genome size of ~75.7 Mb (diploid ~151 Mb), a
heterozygosity rate of ~0.8%, and ~72% unique sequence content (Figure 1).
118

Figure 1. GenomeScope k-mer spectrum analysis of the barley stripe rust pathogen *Puccinia striiformis* f. sp. *hordei* isolate NP85002. The 21-mer frequency distribution and fitted model were used to estimate genome size, heterozygosity, sequencing error rate, and the proportions of unique and duplicated sequences.

Haplotype resolved genome assembly

A haplotype-resolved genome assembly was generated from ONT and Hi-C data using hifiasm v0.25.0-r726 configured for ONT reads and Hi-C-guided phasing[26]. Simplex ONT reads were supplied with the --ont option and Hi-C read pairs with --h1/--h2. The assembler was run with the --dual-scaf to self-scaffold contigs using the homologous region on the alternative haplotype for improved contiguity, and --telo-m CCCTAA was applied to preserve telomeric repeats. This produced two haplotype-specific scaffold/contig sets (hereafter *Psh*_hap1 and *Psh*_hap2.

To remove non-nuclear and low-confidence sequences, we combined sequence-similarity and coverage filtering. All contigs were queried against the NCBI nt database using BLASTN v2.15.0+ with parameters -perc_identity 75 -evalue 1e-5 to check for contaminants and mitochondrial DNA (mtDNA). Contigs with mtDNA hits (pident ≥ 80%) were discarded. ONT reads were then mapped back to the assembly with minimap2 v2.28-r1209 (-ax map-ont --secondary=no)[27] and mean per-contig coverage was calculated using bamtocov v2.7.0[28]. Contigs shorter than 20 kb or with mean coverage <10× were considered low confidence and removed.

Hi-C contact maps were used to confirm nuclear haplotype phasing and to scaffold contigs into chromosomes. Hi-C reads were mapped to the cleaned assembly using Juicer v1.6[29] and the 3D-DNA pipeline v180922[30]. Draft scaffolds were manually curated in the visualization

software Juicebox Assembly Tools v2.20.00[31]. The majority of scaffolds and contigs produced
by hifiasm were already at full chromosome scale where no curation was needed. The final
assembly comprises 18 scaffolds per haplotype (36 in total), totaling 151.3 Mb with scaffold
N50 up to 4.6 Mb, and GC content ~44.3% (Table 1). Chromosome naming was assigned based
on homology to the Pst104E T2T genome assembly[19]. Telomere completeness was assessed
by searching for TTAGGG/CCCTAA motifs within the first and last 50 bp of each scaffold using
FindTelomeres.py   (https://github.com/JanaSperschneider/FindTelomeres).   Of   the   72
expected chromosome ends (18 chromosomes × 2 haplotypes × 2 ends), 69 contained
telomeric repeats; the 5' ends of chr2_hap1, chr15_hap1 and chr15_hap2 lacked detectable
telomeric tracts. Six assembly gaps remained in chr4_hap1/2, chr6_hap1, chr8_hap2 and
chr13_hap1/2 (Table 1), most of which were associated with repetitive regions, such as the
LTR retrotransposon-dense region near the *PR* locus on chr6, and ribosomal DNA repeats on
chr13[19].

168
169 **Table 1.** Assembly statistics for the *Puccinia striiformis* f. sp. *hordei* NP85002 genome assembly. Metrics are
170 reported for the combined assembly (*Psh*_hap1/2) and for each haplotype separately (*Psh*_hap1 and *Psh*_hap2).
171

Assembly statistics	*Psh*_hap1/2	*Psh*_hap1	*Psh*_hap2
Total size (bp)	151,290,331	75,244,898	76,045,433
Number of T2T chromosomes	33	17	16
Scaffold N50 (bp)	4,580,210	4,519,295	4,580,210
Longest scaffold (bp)	5,598,305	5,377,700	5,598,305
GC content (%)	44.33	44.32	44.34
Telomeres detected/expected	69/72	35/36	34/36
Number of internal gaps	6	3	3

172
Gene prediction and annotation
Transposable elements (TEs) were predicted and annotated for each haplotype using the
REPETv4 snakemake pipeline with default parameters[32]. This process identified 44.93%
repetitive content in *Psh*_hap1 and 45.04% in *Psh*_hap2, dominated by Class I
retrotransposons (15.78%) and Class II DNA transposons (14.97%). Identified repeats were
soft-masked using bedtools maskfasta -soft[33]. Gene annotation was performed using two
complementary approaches. First, a reference annotation lift-over strategy was employed.
Dikaryotic evidence-based gene annotations of two high-quality *Pst* assemblies, Pst104E and
AZ2, were projected onto each *Psh* haplotype using Liftoff v1.6.3[34] chromosome by
chromosome with the -chrom option[18,19]. Transferred models with invalid ORFs were filtered
using agat_sp_filter_feature_by_attribute_value.pl --attribute valid_ORFs --value 0 --test "="
(https://github.com/NBISweden/AGAT/tree/master). Second, *ab initio* prediction was run on
each haplotype with funannotate predict to recover additional gene models. The Liftoff-
derived and *ab initio* gene sets were merged into a single GFF3 file using
agat_sp_merge_annotations.pl. Redundant CDS were deduplicated using custom Python

scripts (see Code Availability). Only the isoform with the longest CDS was retained per gene, and genes encoding proteins shorter than 50 amino acids were excluded. For functional annotation, putative secreted proteins were first predicted by identifying signal peptides with SignalP v3.0 (neural network; -t euk -m nn) using the default D-score cutoff (0.433)[35,36]. To reduce false positives, mature (signal peptide-cleaved) protein sequences were screened for transmembrane domains using Phobius v1.01[37] and TMHMM v2.0[37]. Only proteins with transmembrane predictions supported by both tools were excluded from the secretome set to ensure a conservative approach. Secondary metabolite biosynthetic gene clusters were identified with antiSMASH v8.0.4[38]. Interproscan v5.64-96.0.0[39] was then used to assign protein domains and functional signatures, and all results were parsed with funannotate annotate to produce the final functional annotation. The final annotation comprised 17,238 protein-coding genes on *Psh*_hap1 and 17,691 on *Psh*_hap2, along with 571 and 593 tRNA genes, respectively. The mean coding sequence length was 1,076 bp with an average of 4.3 exons per gene, with a total of 4,969 single-exon protein-coding genes. These features are consistent with previously published *P. striiformis* gene structure statistics[18]. Centromere locations were visually estimated from the Hi-C contact maps based on the characteristic "bowtie" patterns indicative of inter-chromosomal centromere-to-centromere interaction enrichment. Where no clear enrichment was observed, centromere positions were estimated by aligning corresponding centromere sequences from Pst104E isolate. Gene density was calculated in non-overlapping 10 kb windows and visualized alongside TE density and gap positions, and telomere locations using KaryplotR[40] to provide an overview of genome organization (Figure 2).

Figure 2. Karyotype representation of the haplotype-resolved *Puccinia striiformis* f. sp. *hordei* genome assembly. Each chromosome pair (1-18) displays haplotype 1 (upper tracks) and haplotype 2 (lower tracks). Gene density is displayed as a heatmap (orange gradient) and transposable element (TE) density is shown as a filled area plot (blue), calculated in 10 kb non-overlapping windows. Centromeric regions are indicated by semi-transparent grey shading boxes across both gene density and TE tracks. Assembly gaps are marked by red circular symbols positioned on the gene density tracks. Black triangular arrowheads indicate telomeric sequences at chromosome ends, with missing telomeres absent on chromosomes 2B, 15A, and 15B. Chromosomes are ordered numerically (Chr1-Chr18) and scaled proportionally to their assembled lengths.

Mitogenome assembly and annotation

The *Psh* mitogenome was assembled separately from the nuclear genome. Mitochondrial reads were first baited by mapping ONT reads against a concatenated FASTA file containing the *Psh* chromosome sequences and a mtDNA contig sourced from AZ2 *Pst* isolate[18]. Reads mapping to the mtDNA contig were extracted, length-filtered (20-120 kb), and assembled with hifiasm (--ont --primary), generating a circular mitogenome assembly[26]. Protein-coding and non-coding RNA genes were annotated with MFannot, an intron- and mini-exon-aware mitogenome annotator, using the genetic code table 4 (Mold, protozoan, and Coelenterate Mitochondrial; Mycoplasma/Spiroplasma)[41]. Protein sequences identified by MFannot were confirmed via BLASTP search against the ClusteredNR database. The final mitogenome assembly (total size 102,058 bp) was re-oriented to begin at the start of the *cox1* gene. A total of 41 mitochondrial genes were detected, including 14 protein-coding genes, 24 tRNA genes, two ribosomal RNA genes (*rns* and *rnl*) and a ribonuclease P RNA (*rnpB*) gene. The circular mitogenome assembly and annotations were visualized using the Proksee web-tool (Figure 3)[42].

236
Figure 3. Circular representation of the mitochondrial genome assembly of *Puccinia striiformis* f. sp. *hordei*
NP85002 isolate (size 102,058 bp), re-oriented to begin at the start of the *cox1* gene. The mitogenome contains
41 annotated genes, including 14 protein-coding genes, 24 tRNA genes, two ribosomal RNA genes (*rns and rnl*),
and one RNase P RNA gene (*rnpB*). Protein-coding genes are shown in red, tRNA genes in brown, and other non-
coding RNA genes (*rns, rnl, rnpB*) in green. The innermost track represents GC content variation across the
mitochondrial genome.

3. Data Records

ONT long-reads and Hi-C reads generated in this study have been deposited in the NCBI
Sequence Read Archive (SRA) under BioProject PRJNA1406799 and BioSample
SAMN54803514. The haplotype-resolved nuclear genome assemblies and the mtDNA
genome assembly have been submitted to NCBI and will be released upon processing under
accessions PRJNA1406819 (haplotype 1), PRJNA1406818 (haplotype 2), and x (mtDNA). The
assemblies (haplotype 1, haplotype 2, and mtDNA) and associated resources, including gene
and transposable element annotations, are available via Zenodo (DOI:
10.5281/zenodo.18323365).

4. Technical Validation

We evaluated the *Psh* assembly using k-mer-based consensus accuracy, read-mapping support, Hi-C contact structure and gene-space completeness. These complementary metrics provide an integrated assessment of assembly quality and haplotype resolution of the dikaryotic *Psh* genome.

K-mer-based consensus accuracy

Reference-free consensus accuracy and completeness were estimated with Merqury v1.3 using a k-mer database (k = 21) built from ONT reads[43]. The combined assembly achieved a QV of 72.01 (6.30×10⁻⁸ error rate), with similarly high-quality values for *Psh*_hap1 and *Psh*_hap2 (QV 71.41 and 72.69, respectively). K-mer completeness exceeded 99.999% for both haplotypes with error rates of approximately 1 in 100 million bases. Spectra-cn plots showed the expected dikaryotic pattern, with most k-mers present once per haplotype and only a small fraction of higher-copy k-mers, consistent with limited haplotype collapse (Figure 4).

Figure 4. Spectra-cn plot of 21-mer multiplicity for the haplotype-resolved *Puccinia striiformis* f. sp. *hordei* assembly. The distribution shows the expected dikaryotic pattern, with most k-mers assigned to the 1× and 2× copy-number classes and only a small fraction at higher copy number (>2×), consistent with limited haplotype collapse.

Read mapping and coverage

ONT reads were mapped back to the final assembly with minimap2 v2.28-r1209[27] (-ax map-ont), followed by sorting, indexing, and coverage estimation with samtools v1.16[44]. Approximately 90% of ONT reads mapped to the assembled nuclear genome, with sequencing coverage ranging from ~98% to 100% across both haplotypes. Hi-C paired-end reads were mapped with BWA-MEM v0.7.18-r1243 to the final assembly[45]. Approximately 95% of Hi-C read pairs properly mapped, with sequencing coverage of >99%, providing robust support for chromosome-scale scaffolding and contact-map-based curation.

Hi-C scaffolding and haplotype phasing

A Hi-C contact matrix for the full dikaryotic genome was generated using the Juicer/3D-DNA pipeline and visualized in Juicebox Assembly Tools (Figure 5). The contact heatmap clearly supported the spatial separation of homologous chromosomes into two nuclear haplotypes, *Psh*_hap1 and *Psh*_hap2. Within each haplotype, the map showed strong along-diagonal contact signals for all 18 chromosomes with no apparent off-diagonal blocks indicative of major mis-joins or misplaced contigs. Centromere positions were inferred from strong inter-chromosomal contact signals and were used to confirm chromosome orientations from the shorter p-arm to the q-arm.

Figure 5. Hi-C contact map of the haplotype-resolved *Puccinia striiformis* f. sp. *hordei* assembly visualized in Juicebox. Strong within-haplotype diagonal signal across all 18 chromosomes and minimal off-diagonal structure support correct scaffolding and separation of haplotypes A and B.

Nuclear haplotype phasing correctness

To quantitatively evaluate haplotype separation and phasing accuracy, HiC-Pro v3.1.0 was used to generate a contact matrix from MAPQ≥20 Hi-C alignments[46]. *Cis-* and *trans-* chromosome contacts were quantified from raw matrix (bin size 20 kb) and visualized on a

circos plot using https://github.com/RunpengLuo/HiC-Analysis[19]. The analysis demonstrated accurate phase separation with 97.54% of Hi-C contacts occurring within-haplotype contained in one nucleus, and only 2.46% representing cross-haplotype links (Table 2). The low cross-haplotype contact rate confirms successful separation of the two nuclear genomes and high-quality haplotype resolution (Figure 6).

Table 2. Hi-C contact counts used to assess haplotype phasing accuracy in the haplotype-resolved *Puccinia striiformis* f. sp. *hordei* assembly. Contacts were quantified from MAPQ≥20 Hi-C alignments and summarized as within-haplotype links (cis- and trans-chromosome contacts within *Psh*_hap1 or *Psh*_hap2) and cross-haplotype links (contacts between haplotypes). Values are shown as counts with percentages of total contacts.

Hi-C contact map types	Contacts (%)
Within-haplotype links	432,454 (97.54%)
Cis-chromosome, *Psh*_hap1	176,512 (39.81%)
Cis-chromosome, *Psh*_hap2	221,582 (49.98%)
Trans-chromosome, *Psh*_hap1	15,158 (3.42%)
Trans-chromosome, *Psh*_hap2	19,202 (4.33%)
Cross-haplotype links (total)	10,928 (2.46%)

Figure 6. Circos plot visualizing within- and cross-haplotype Hi-C contact links, confirming that *Psh* haplotypes were accurately phased into two separate nuclei (red = haplotype 1, blue = haplotype 2). Each chromosome sequence was divided into 20 kb windows, and Hi-C contacts (MAPQ≥20) were counted per window. Cross-haplotype links between haplotype 1 and haplotype 2 are shown in green. A region on chr17_hap2 displayed abundant cross-haplotype contacts with chr2_hap1, chr5_hap1, and chr7_hap1; these contacts overlapped the estimated centromeric regions and likely reflect mapping noise associated with repetitive centromeric sequences.

Gene-space completeness

Assembly completeness was assessed using BUSCO v6.0.0 with the basidiomycota_odb12 lineage dataset[47]. In genome mode, haplotype 1 and haplotype 2 each showed high completeness (90.9% and 91.0% complete BUSCOs, respectively), with single-copy BUSCOs comprising 89.0% and 86.9% of each haplotype, respectively. When both haplotypes were analyzed together, overall completeness was 90.1%, with most BUSCOs classified as duplicated (85.7%), consistent with the expected pattern for a haplotype-resolved assembly where orthologous genes are present in both haplotypes.

For gene annotation, lift-over projections from Pst104E (87.8% complete BUSCOs; 28,436 genes) and AZ2 (92.4%; 26,336 genes), together with *ab initio* gene predictions (86.1%; 19,084 genes), identified a higher number of BUSCOs when combined (94.4%; 34,929 genes). This supports the effectiveness of integrating lift-over and *ab initio* annotation approaches to resolve gene models despite the high repeat load and dikaryotic genome structure (Table 3).

Table 3. BUSCO completeness assessment of protein sequences annotated in the *Psh* genome assembly. BUSCO v6.0.0 (basidiomycota_odb12) was run on the combined assembly (*Psh*_hap1/2) and each haplotype (*Psh*_hap1 and *Psh*_hap2). Values report the percentage of complete BUSCOs (split into single-copy and duplicated), as well as fragmented and missing BUSCOs.

Haplotype	Complete (%)	Single-copy (%)	Duplicated (%)	Fragmented (%)	Missing (%)
*Psh*_hap1/2	94.4	2.2	92.2	1.4	4.2
*Psh*_hap1	93.6	89.0	4.5	1.5	5.0
*Psh*_hap2	93.5	86.9	6.6	1.5	4.9

5. Code availability

Analyses were performed using the software and workflows described in the Methods section. Where relevant, command-line options and key parameters are reported; otherwise, tools were run using standard or recommended settings. Custom scripts used for data processing and figure generation are available on GitHub at: (https://github.com/ritatam/PshNP85002-genome-assembly).

6. References

1.   Marshall, D. & Sutton, R.L. Epidemiology of stripe rust, virulence of Puccinia striiformis f.sp. hordei, and yield loss in barley. *Plant Disease* **79**, 732–737 (1995).
2.   Line, R.F. Stripe rust of wheat and barley in North America: a retrospective historical review. *Annu Rev Phytopathol* **40**, 75-118 (2002).

3.   Visioni, A. *et al.* Genome Wide Association Mapping of Seedling and Adult Plant
Resistance to Barley Stripe Rust (Puccinia striiformis f. sp. hordei) in India. *Frontiers in*
Plant Science **Volume 9 - 2018**(2018).

4.   Wellings, C.R. Global status of stripe rust: a review of historical and current threats.
Euphytica **179**, 129-141 (2011).

5.   Baik, B.-K. & Ullrich, S.E. Barley for food: Characteristics, improvement, and renewed
interest. *Journal of Cereal Science* **48**, 233-242 (2008).

6.   Chen, X. & Line, R.F. Identification of genes for resistance to Puccinia striiformis f. sp.
hordei in 18 barley genotypes. *Euphytica* **129**, 127-146 (2003).

7.   Anikster, Y. The *formae speciales*. in *The Cereal Rusts*, Vol. I (eds. Bushnell, W.R. &
Roelfs, A.P.) 124-137 (Academic Press, Orlando, FL., 1984).

8.   Zhao, J., Wang, M., Chen, X. & Kang, Z. Role of alternate hosts in epidemiology and
pathogen variation of cereal rusts. *Annu. Rev. Phytopathol.* **54**, 207-28 (2016).

9.   Mojerlou, S., Schwessinger, B. & Rodriguez-Algaba, J. Sexual recombination in cereal
rust fungi: Knowns and unknowns of pathogen evolution and adaptation. *PLOS*
Pathogens **21**, e1012908 (2025).

10. Hovmoller, M.S., Sorensen, C.K., Walter, S. & Justesen, A.F. Diversity of Puccinia
striiformis on cereals and grasses. *Annu Rev Phytopathol* **49**, 197-217 (2011).

11. Brown, J.K. & Hovmøller, M.S. Aerial dispersal of pathogens on the global and
continental scales and its impact on plant disease. *Science* **297**, 537-41 (2002).

12. Hovmøller, M.S. *et al.* Long-term surveillance suggests multiple hybridization events
by nuclear reassortment and accelerated intercontinental spread of wheat yellow
rust. *bioRxiv*, 2025.05.22.655633 (2025).

13. Schwessinger, B. *et al.* Distinct Life Histories Impact Dikaryotic Genome Evolution in
the Rust Fungus Puccinia striiformis Causing Stripe Rust in Wheat. *Genome Biol Evol*
12, 597-617 (2020).

14. Schwessinger, B. *et al.* A Near-Complete Haplotype-Phased Genome of the Dikaryotic
Wheat Stripe Rust Fungus Puccinia striiformis f. sp. tritici Reveals High Interhaplotype
Diversity. *mBio* **9**, e02275-17 (2018).

15. Xia, C. *et al.* Genomic insights into host adaptation between the wheat stripe rust
pathogen (Puccinia striiformis f. sp. tritici) and the barley stripe rust pathogen
(Puccinia striiformis f. sp. hordei). *BMC Genomics* **19**, 664 (2018).

16. Xia, C. *et al.* Genome Sequence Resources for the Wheat Stripe Rust Pathogen
(Puccinia striiformis f. sp. tritici) and the Barley Stripe Rust Pathogen (Puccinia
striiformis f. sp. hordei). *Molecular Plant-Microbe Interactions®* **31**, 1117-1120
(2018).

17. Schwessinger, B. *et al.* A Chromosome Scale Assembly of an Australian Puccinia
striiformis f. sp. tritici Isolate of the PstS1 Lineage. *Molecular Plant-Microbe*
Interactions® **35**, 293-296 (2022).

18. Wang, J. *et al.* A fully haplotype-resolved and nearly gap-free genome assembly of
wheat stripe rust fungus. *Scientific Data* **11**(2024).

19. Tam, R. *et al.* Long-read genomics reveal extensive nuclear-specific evolution and
allele-specific expression in a dikaryotic fungus. (Cold Spring Harbor Laboratory,
2024).

20. Pereira, E.C. *et al.* Defence Recognition of a Stripe Rust Fungal Effector Is Uncoupled
from Disease Outcomes in Wheat. *Molecular Plant-Microbe Interactions®* (2026).

21.   Holden, S. *et al.* Uncovering the history of recombination and population structure in
394   western Canadian stripe rust populations through mating type alleles. *BMC Biology*
395   **21**, 233 (2023).

396   22.   Hovmøller, M.S., Rodriguez-Algaba, J., Thach, T. & Sørensen, C.K. Race Typing of
397   *Puccinia striiformis* on Wheat. in *Wheat Rust Diseases, Methods and Protocols* (ed.
398   Periyannan, S.) 29-40 (Springer Protocols, Methods in Molecular Biology, N.Y., U.S.A.,
399   2017).

400   23.   Schwessinger, B. & Rathjen, J.P. Extraction of High Molecular Weight DNA from
401   Fungal Rust Spores for Long Read Sequencing. in *Wheat Rust Diseases: Methods and*
402   *Protocols* (ed. Periyannan, S.) 49-57 (Springer New York, New York, NY, 2017).

403   24.   Cabanettes, F. & Klopp, C. D-GENIES: dot plot large genomes in an interactive,
404   efficient and simple way. *PeerJ* **6**, e4958 (2018).

405   25.   Vurture, G.W. *et al.* GenomeScope: fast reference-free genome profiling from short
406   reads. *Bioinformatics* **33**, 2202-2204 (2017).

407   26.   Cheng, H., Concepcion, G.T., Feng, X., Zhang, H. & Li, H. Haplotype-resolved de novo
408   assembly using phased assembly graphs with hifiasm. *Nature Methods* **18**, 170-175
409   (2021).

410   27.   Li, H. Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* **34**,
411   3094-3100 (2018).

412   28.   Birolo, G. & Telatin, A. BamToCov: an efficient toolkit for sequence coverage
413   calculations. *Bioinformatics* **38**, 2617-2618 (2022).

414   29.   Durand, N.C. *et al.* Juicer Provides a One-Click System for Analyzing Loop-Resolution
415   Hi-C Experiments. *Cell Syst* **3**, 95-8 (2016).

416   30.   Dudchenko, O. *et al.* De novo assembly of the Aedes aegypti genome using Hi-C
417   yields chromosome-length scaffolds. *Science* **356**, 92-95 (2017).

418   31.   Durand, N.C. *et al.* Juicebox Provides a Visualization System for Hi-C Contact Maps
419   with Unlimited Zoom. *Cell Systems* **3**, 99-101 (2016).

420   32.   Bardet, E., Wan, M., Confais, J. & Quesneville, H. REPET v4.0: A Comprehensive Tool
421   for Transposable Element Analysis. in *JOBIM 2025* (Bordeaux, France, 2025).

422   33.   Quinlan, A.R. & Hall, I.M. BEDTools: a flexible suite of utilities for comparing genomic
423   features. *Bioinformatics* **26**, 841-842 (2010).

424   34.   Shumate, A. & Salzberg, S.L. Liftoff: accurate mapping of gene annotations.
425   *Bioinformatics* **37**, 1639-1643 (2021).

426   35.   Dyrløv Bendtsen, J., Nielsen, H., von Heijne, G. & Brunak, S. Improved Prediction of
427   Signal Peptides: SignalP 3.0. *Journal of Molecular Biology* **340**, 783-795 (2004).

428   36.   Sperschneider, J., Williams, A.H., Hane, J.K., Singh, K.B. & Taylor, J.M. Evaluation of
429   Secretion Prediction Highlights Differing Approaches Needed for Oomycete and
430   Fungal Effectors. *Frontiers in Plant Science* **Volume 6 - 2015**(2015).

431   37.   Käll, L., Krogh, A. & Sonnhammer, E.L.L. A Combined Transmembrane Topology and
432   Signal Peptide Prediction Method. *Journal of Molecular Biology* **338**, 1027-1036
433   (2004).

434   38.   Blin, K. *et al.* antiSMASH 7.0: new and improved predictions for detection, regulation,
435   chemical structures and visualisation. *Nucleic Acids Research* **51**, W46-W50 (2023).

436   39.   Jones, P. *et al.* InterProScan 5: genome-scale protein function classification.
437   *Bioinformatics* **30**, 1236-1240 (2014).

438   40.   Gel, B. & Serra, E. karyoploteR: an R/Bioconductor package to plot customizable
439   genomes displaying arbitrary data. *Bioinformatics* **33**, 3088-3090 (2017).

41. Lang, B.F. *et al.* Mitochondrial genome annotation with MFannot: a critical analysis of gene identification and gene model prediction. *Frontiers in Plant Science* **Volume 14 - 2023**(2023).

42. Grant, J.R. *et al.* Proksee: in-depth characterization and visualization of bacterial genomes. *Nucleic Acids Research* **51**, W484-W492 (2023).

43. Rhie, A., Walenz, B.P., Koren, S. & Phillippy, A.M. Merqury: reference-free quality, completeness, and phasing assessment for genome assemblies. *Genome Biology* **21**, 245 (2020).

44. Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078-2079 (2009).

45. Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows–Wheeler transform. *Bioinformatics* **25**, 1754-1760 (2009).

46. Servant, N. *et al.* HiC-Pro: an optimized and flexible pipeline for Hi-C data processing. *Genome Biology* **16**, 259 (2015).

47. Tegenfeldt, F. *et al.* OrthoDB and BUSCO update: annotation of orthologs with wider sampling of genomes. *Nucleic Acids Research* **53**, D516-D522 (2025).

7. Acknowledgements

This work was supported by Villum Fonden, Denmark (https://veluxfoundations.dk/en) under grant number 50161 awarded to J.R-A. R.T. was supported by a Grains Research and Development Corporation Graduate Research Scholarship. Computational resources were supported by the Australian Government through the National Computational Infrastructure (NCI) under the ANU Merit Allocation Scheme. J.R-A. and R.T. thank Ellen Jørgensen and Jakob Sørensen (GRRC, Aarhus University) for assistance with isolate handling and plant multiplication, respectively.

8. Author contributions

R.T., B.S., and J.R-A. designed and conceptualized the study. J.R-A. acquired funding and supervised the research. M.M. conducted ONT long-read sequencing. S.M. extracted high-molecular-weight DNA and prepared fungal spores for Hi-C library preparation. R.T. and J.R-A. performed bioinformatic analyses including genome assembly, annotation, quality control, and data visualization. J.P.R. and B.S. contributed to interpretation of results and provided critical feedback on the manuscript. R.T. and J.R-A. wrote the original draft manuscript. All authors contributed to manuscript revision, reviewed the final version, and approved it for publication.

9. Competing Interests

The authors declare no competing interests.