Basal ppGpp regulation by SpoT coordinates metabolic homeostasis and acid resistance

Yueyue Liu[1#], Muriel Leandra Schicketanz[1#^], Xichuan Zhai[2], Ling Deng[2], Kenn Gerdes[3], Yong

Everett Zhang[1*]

[1] Department of Biology, University of Copenhagen, DK-2200 Copenhagen, Denmark

[2] Section of Food Microbiology, Gut Health, and Fermentation, Department of Food Science,

University of Copenhagen, Rolighedsvej 26, 1958 Frederiksberg C, Denmark

[3] Independent Researcher, Copenhagen, Denmark

[#] Y.Y.L. and M.L.S. contributed equally to this work.

[^] Current address: Novonesis A/S, Kgs. Lyngby, Denmark

*Correspondence: Yong Everett Zhang zhangy830328@gmail.com

Running title: Basal ppGpp links metabolism and acid resistance

Abstract

Bacteria survive fluctuating and hostile environments by coordinating core metabolism with stress-protective responses. How these processes are integrated at the molecular level remains incompletely understood. Here, we uncover a previously unrecognized role for the stringent response enzyme SpoT in coordinating amino acid homeostasis and acid resistance through maintenance of a basal pool of the alarmone ppGpp. Focusing on a conserved histidine residue in SpoT (H414), we show that its disruption causes severe growth defects in minimal medium and catastrophic acid sensitivity in *Escherichia coli*, independently of the canonical RelA-mediated stringent response. Transcriptomic and genetic analyses reveal that a SpoT H414A substitution leads to misallocation of metabolic flux, characterized by excessive arginine biosynthesis and concomitant depletion of intracellular glutamate, the essential substrate for the glutamate-dependent Gad acid resistance system. Intragenic suppressor mutations within *spoT* restore growth and acid resistance by re-establishing a sub-basal yet physiologically sufficient ppGpp pool, demonstrating that very low levels of ppGpp are required for metabolic balance and stress preparedness. Importantly, the requirement for SpoT H414 in supporting growth and acid survival is conserved in pathogenic *Salmonella* and *Shigella*. Together, our findings establish SpoT-derived basal ppGpp as a distinct regulatory regime that integrates metabolism and acid resistance, providing a framework for understanding how bacteria maintain physiological homeostasis and survive acidic environments.

KEYWORDS: basal ppGpp, SpoT, stringent response, metabolic homeostasis, acid resistance, Gad system

Introduction

Bacteria inhabit environments that fluctuate dramatically in nutrient availability and physicochemical stress. To survive these challenges, cells must coordinate metabolic activity with stress-protective responses in a manner that preserves essential resources while maintaining viability. In enteric bacteria, exposure to acidic conditions, such as those encountered during passage through the mammalian stomach, represents a particularly severe and ecologically relevant challenge. Survival under these conditions is a key determinant of fitness and virulence[1], yet the mechanisms that integrate metabolic state with acid resistance remain incompletely understood.

Escherichia coli exemplifies bacterial acid adaptation, surviving exposure to pH values as low as 2.5 through multiple acid resistance (AR) systems[2-4] Among these, the glutamate-dependent Gad system provides the most robust protection under extreme acid stress by coupling proton consumption to amino acid metabolism. Cytoplasmic glutamate decarboxylases (GadA and GadB) convert glutamate to γ-aminobutyrate (GABA), consuming intracellular protons, while the GadC antiporter exchanges GABA for extracellular glutamate to sustain the cycle[5,6]. This system is tightly regulated[3,6] and energetically costly, requiring precise coordination between metabolic capacity, substrate availability, and stress signalling. How cells ensure that sufficient glutamate is available to support acid resistance under nutrient-limited conditions remains a central unresolved question.

Amino acid metabolism is globally regulated by the stringent response, a conserved bacterial signaling network mediated by the alarmones guanosine tetra- and pentaphosphate (ppGpp)[7-10]. In *E. coli*, the ribosome-associated enzyme RelA synthesizes large bursts of ppGpp in response to amino acid starvation[11-15], triggering extensive transcriptional[16,17] and metabolic[18-21] reprogramming. This acute stringent response represses stable RNA synthesis[16,17], remodels nucleotide metabolism[18-21], and promotes amino acid biosynthesis[22-24], thereby restoring metabolic

balance during nutrient stress[25]. While the role of RelA-driven ppGpp accumulation has been

extensively characterized, the function of its paralog SpoT remains less well defined[7,26].

SpoT is unique among stringent response enzymes in possessing both ppGpp synthetase and

hydrolase activities[27,28] . In contrast to RelA, SpoT produces lower levels of ppGpp and responds to

a broader range of environmental cues, including fatty acid limitation[29,30], iron starvation[31], and

other non–amino acid stresses[32-36]. Genetic and physiological studies have implicated SpoT in

virulence, persistence, and stress adaptation across diverse bacterial species[7,33,37], yet its specific

regulatory role and downstream physiological functions remain poorly understood. In particular,

whether SpoT-derived ppGpp represents merely a weaker version of the canonical stringent

response or instead constitutes a distinct regulatory regime has remained an open question.

Here, we address this gap by dissecting the function of SpoT in the absence of RelA. We identify a

conserved histidine residue (H414) in the C-terminal regulatory region of SpoT that is essential for

maintaining a basal level of ppGpp. Mutagenesis of this residue to alanine uncouples amino acid

metabolism from acid resistance, leading to glutamate starvation, transcriptional collapse of the Gad

system, and extreme acid sensitivity. Through intragenic suppressor analysis, enzymatic

perturbations, and physiological rescue experiments, we demonstrate that a sub-basal yet non-zero

pool of ppGpp is both necessary and sufficient to coordinate metabolic homeostasis and acid

defense. Finally, we show that this SpoT-dependent regulatory logic is conserved in pathogenic

enterobacteria, underscoring its broader relevance to bacterial survival and virulence.

Together, our findings reveal SpoT-derived basal ppGpp as a central integrator of metabolism and

stress resistance, expanding the conceptual framework of the stringent response beyond RelA-

mediated acute starvation signaling to include a foundational role in bacterial physiological

homeostasis.

Results

Mutagenesis of the conserved H414 residue of SpoT leads to defective growth in minimal medium

To investigate how SpoT integrates environmental cues independently of RelA, we focused on the C-terminal regulatory region of SpoT, which is highly conserved among RelA/SpoT homologs but poorly understood mechanistically. Sequence alignment revealed a conserved histidine residue (H414 in *E. coli* SpoT) within the TGS domain, corresponding to H432 in RelA (**Extended Data Fig. 1a, 1b**), a critical residue in sensing amino acid starvation[12-15].

To probe the functional role of this residue, we generated a SpoT H414A substitution in a *ΔrelA* background. While the Δ*relA* *spoT^wt* (hereto after *spoT^wt*) strain grew normally, the H414A mutant (*spoT^H414A*) unexpectedly failed to grow on glucose minimal medium (M9Glc), both on solid and in liquid culture (**Fig. 1a, 1b**). This severe growth defect was not anticipated and suggested a critical role for SpoT H414 in supporting growth under nutrient-limited conditions. The unexpected growth failure of *spoT^H414A* cells prompted us to investigate the underlying molecular basis of this phenotype.

Transcriptomic analysis of *spoT^H414A* cells reveals collapse of stress and acid resistance programs

To gain insight into the molecular defects underlying the growth failure of *spoT^H414A* cells, we performed transcriptomic analysis of the *spoT^H414A* and *spoT^wt* strains. Because *spoT^H414A* cells failed to proliferate in M9Glc, cultures were incubated under identical conditions until the *spoT^wt* strain entered early exponential phase, at which point RNA was harvested from both strains (**Extended Data Fig. 1c**).

Unexpectedly, RNA sequencing revealed extensive transcriptional dysregulation in $spoT^{H414A}$ cells, with strong repression of stress adaptation and acid resistance genes. Among the most prominently downregulated loci were the glutamate-dependent Gad acid resistance system (*gadE*, *gadX*, *gadW*, *gadA*, *gadB*, *gadC*), as well as additional components of the acid fitness island, including the periplasmic chaperones *hdeAB* (**Fig. 2a, 2b**). Genes associated with osmotic and general stress responses were similarly reduced; in contrast, genes linked to phage shock and SOS responses were strongly induced (**Extended Data Fig. 1d-f**), suggesting widespread physiological stress. The unexpected transcriptional collapse of acid resistance pathways prompted us to directly examine the acid survival phenotypes of $spoT^{H414A}$ cells.

SpoT, but not RelA, is required for survival under extreme acid stress

Guided by the pronounced transcriptional repression of acid resistance genes in $spoT^{H414A}$ cells, we next examined survival under lethal acid stress. Upon exposure to pH2.5, $spoT^{H414A}$ cells exhibited extreme acid sensitivity, with viability declining rapidly and no detectable survivors after 60 minutes (**Fig. 2c**). In contrast, both the parental wild-type strain and the $spoT^{wt}$ strain survived acid challenge efficiently, demonstrating that RelA is dispensable for acid resistance under these conditions.

Notably, despite the severe acid sensitivity of $spoT^{H414A}$ cells, viability under neutral, well-buffered conditions (M9Glc) was preserved (**Extended Data Fig. 1g**), indicating that SpoT H414 is specifically required for stress preparedness rather than basal viability. Together, these results establish SpoT as the primary stringent response enzyme supporting acid survival in *E. coli*, independently of RelA.

Transcriptomic signatures indicate arginine metabolic overflow and glutamate depletion

To define the metabolic basis of the growth defect observed in $spoT^{H414A}$ cells, we examined

transcriptomic changes in central metabolic pathways. While expression of many amino acid

biosynthetic genes was modestly reduced, one pathway stood out: arginine biosynthesis was

strongly and coordinately upregulated (**Fig. 3a, 3b**). All the nine enzymes of the arginine

biosynthetic pathway exhibited significantly increased expression, together with *carAB*, which

supplies carbamoylphosphate to this pathway. This transcriptional pattern indicates a pronounced

redirection of metabolic flux toward arginine production.

In contrast, genes directing carbamoylphosphate toward pyrimidine nucleotide biosynthesis were

downregulated, suggesting selective diversion of this shared intermediate rather than a generalized

increase in anabolic metabolism. Because arginine biosynthesis consumes glutamate as a precursor,

the coordinated induction of the *arg* operons predicts substantial depletion of intracellular glutamate

pool.

In parallel with arginine pathway induction, genes involved in *de novo* purine nucleotide

biosynthesis were broadly upregulated (**Extended Data Fig. 2**). Many of these reactions consume

glutamine as a nitrogen donor, which is regenerated from glutamate, thereby imposing an additional

indirect drain on glutamate availability. Thus, simultaneous activation of arginine and purine

biosynthetic programs is predicted to exacerbate glutamate depletion, compounding the metabolic

imbalance caused by SpoT H414A mutation.

Together, these transcriptional signatures suggest that SpoT H414A leads to misallocation of

metabolic resources, with excessive flux into arginine biosynthesis and increased purine nucleotide

production collectively depleting glutamate. Given the central role of glutamate in both cellular

metabolism and acid resistance, this metabolic imbalance provides a unifying explanation for the

growth and stress survival defects observed in $spoT^{H414A}$ cells and prompted us to directly test the

role of glutamate availability in these phenotypes.

Glutamate starvation underlies the growth defect of $spoT^{H414A}$ cells

Prompted by the transcriptomic evidence for glutamate depletion, we tested whether exogenous

supplementation could restore growth. Addition of glutamate to M9Glc markedly improved growth

of $spoT^{H414A}$ cells, substantially shortening the lag phase and increasing final biomass (**Fig. 3c**). In

contrast, supplementation with arginine or uracil failed to restore growth, and combining these

metabolites with glutamate provided no additional benefit.

Amond all 20 amino acids, supplementation with glutamine, asparagine, or proline partially

restored growth, consistent with their metabolic proximity to glutamate, but none matched the effect

of glutamate itself (**Extended Data Fig. 3a**). Increasing the concentration of glutamate up to

tenfold failed to further improve growth (**Extended Data Fig. 3b**), and combining glutamate with

glutamine, asparagine, or proline did not enhance regrowth beyond that achieved with glutamate

alone (**Extended Data Fig. 3c**). In contrast, supplementation with all 20 amino acids fully restored

the growth defect of $spoT^{H414A}$ cells to a level exceeding that of the $spoT^{wt}$ strain (**Extended Data**

Fig. 3d), indicating that glutamate starvation represents the dominant but not exclusive metabolic

limitation imposed by SpoT H414A dysfunction. Importantly, glutamate supplementation had only

minimal effects on the growth of a ppGpp⁰ ($\Delta relA$ $\Delta spoT$) strain (**Extended Data Fig. 3e**),

indicating that the rescue of $spoT^{H414A}$ growth depends on residual SpoT activity rather than simple

nutritional supplementation.

To directly test whether arginine biosynthesis drives glutamate depletion, we deleted $argA$,

encoding the first committed enzyme of the arginine biosynthetic pathway. The resulting $spoT^{H414A}$

ΔargA strain failed to grow in M9Glc (**Fig. 3d**). Strikingly, arginine, but not glutamate, restored

growth in this background, demonstrating that blocking arginine synthesis prevents glutamate

depletion and alleviates the growth defect.

Conversely, artificial induction of arginine overflow phenocopied the *spoT^H414A* growth defect.

Overexpression of a feedback-resistant ArgA variant (ArgA^H15Y)[38] delayed growth in M9Glc, and

this delay was significantly alleviated by glutamate supplementation (**Fig. 3e**). Together, these

results demonstrate that SpoT H414 prevents arginine-driven glutamate starvation, thereby

maintaining amino acid homeostasis required for growth in minimal medium.

Basal SpoT activity supports glutamate-dependent acid resistance

Because glutamate is the essential substrate for the Gad acid resistance system, we next asked

whether glutamate depletion contributes to the acid sensitivity of *spoT^H414A* cells. Supplementation

of the acid challenge medium with glutamate fully rescued survival during short-term acid exposure

and substantially improved long-term viability (**Fig. 4a**). In contrast, supplementation with arginine

or lysine provided only modest protection (**Extended Data Fig. 4a**), consistent with the dominant

role of the glutamate-dependent Gad system under extreme acid stress.

To determine whether impaired regulation of the Gad system accounted for the acid resistance

defect, we ectopically expressed *gadE*, the master transcriptional activator of Gad genes. Even low-

level expression of *gadE* significantly restored acid resistance in *spoT^H414A* cells, and combined

supplementation with glutamate further enhanced survival (**Extended Data Fig 4b**). These

interventions did not alter acid resistance in *spoT^wt* cells, indicating that the defect is specific to

SpoT dysfunction. Notably, despite its ability to restore acid survival, ectopic expression of *gadE*

failed to improve the growth of *spoT^H414A* cells in M9Glc (**Extended Data Fig. 4c**). This

dissociation indicates that impaired acid resistance is not the primary cause of the growth defect,

which instead arises from glutamate starvation and broader metabolic misallocation.

We further examined acid tolerance across a range of pH values. In addition to catastrophic

sensitivity at pH 2.5, $spoT^{H414A}$ cells exhibited reduced survival under mildly acidic conditions (pH

4.0–6.0) (**Fig. 4b, Extended Data Fig 4d**), consistent with the broader impairment of various acid

stress responses (**Extended Data Fig 4e**). Notably, these defects occurred despite preserved

viability under neutral conditions, indicating that SpoT-mediated basal regulation specifically

primes cells for acid stress rather than broadly supporting growth.

Intragenic suppressor mutations within *spoT* restore growth

The severe growth and acid resistance defects of $spoT^{H414A}$ cells suggested that H414 of SpoT plays

a central regulatory role. To gain mechanistic insight into how SpoT activity is compromised by the

H414A substitution, we isolated spontaneous suppressor mutants capable of restoring growth of

$spoT^{H414A}$ cells on M9Glc. After prolonged incubation (24–72 h), colonies of varying sizes emerged

on M9Glc agar (**Fig. 5a**).

Analysis of 50 independently isolated suppressors revealed that the strongest suppressors

predominantly carried secondary mutations within the *spoT* gene itself. Among large-colony

suppressors, we identified recurrent substitutions at residues G334 (G334S; supA), E364 (E364K;

supB), and S463 (S463R; sup41) as well as a small in-frame deletion within the synthetase domain

(sup17) (**Fig. 5b**). Whole-genome sequencing of smaller suppressors lacking *spoT* mutations

identified a single mutation in *rpoB* (K1262Q), which is likely similar as the previously described

stringent-response suppressors[39], and these were excluded from further analysis.

Reconstruction of the identified *spoT* suppressor mutations in a clean background fully restored

growth of $spoT^{H414A}$ in M9Glc (**Extended Data Fig 5**), whereas deletion of *spoT* in the suppressor

strains abolished growth (**Fig. 5c**). These results demonstrate that intragenic mutations within SpoT

are both necessary and sufficient to suppress the $spoT^{H414A}$ growth defect.

Suppressor mutations restore acid resistance in proportion to growth recovery

We next asked whether suppression of the growth defect was accompanied by restoration of acid

resistance. All four *spoT* suppressor strains exhibited improved survival under lethal acid challenge

(pH 2.5) compared to the parental $spoT^{H414A}$ strain, although the degree of rescue varied (**Fig. 5d**).

The suppressor (sup41) displayed the most robust acid resistance, followed by supB, supA and

sup17.

Supplementation of the acid challenge medium with glutamate further enhanced survival across all

suppressor strains, with particularly pronounced effects in the weaker suppressors supB and sup17

(**Fig. 5d**). These differences suggest that the suppressor mutations restore SpoT function through

distinct mechanisms that differentially affect glutamate homeostasis and stress preparedness.

Thus, intragenic suppression of SpoT H414A simultaneously restores growth and acid resistance,

reinforcing the link between SpoT activity, metabolic balance, and stress survival.

A sub-basal but non-zero level of ppGpp is sufficient for growth in minimal medium

232 Given SpoT's role as a ppGpp synthase and hydrolase, we hypothesized that the growth defect of

233 $spoT^{H414A}$ cells reflects insufficient ppGpp production and that suppressor mutations restore a

234 physiologically meaningful basal ppGpp pool. To test this, we measured intracellular ppGpp levels

235 using ^{32}P-orthophosphate labeling and thin-layer chromatography[40].

As expected, $spoT^{H414A}$ cells produced barely detectable levels of ppGpp, whereas $spoT^{wt}$ cells accumulated a clear basal ppGpp signal (**Fig. 6a**). Surprisingly, suppressor strains did not exhibit a marked increase in detectable ppGpp compared to $spoT^{H414A}$ cells, suggesting that either very small increases in ppGpp are sufficient for physiological rescue or that ppGpp turnover is highly dynamic in these strains.

To directly test whether ppGpp synthesis is required for suppression, we inactivated the ppGpp synthetase activity of SpoT (E319Q substitution) in the suppressor backgrounds. These mutations dramatically delayed or abolished growth recovery in M9Glc (**Fig. 6b**), demonstrating that ppGpp synthesis is essential for suppression. Conversely, inactivation of SpoT's hydrolase activity (H72A D73A substitution) led to clear accumulation of ppGpp (**Fig. 6c**) and accelerated growth resumption (**Fig. 6d**), indicating that suppressor strains produce ppGpp that is rapidly degraded. Notably, the hydrolase-inactive (SYN) variants of supA and sup17 could not be obtained, likely due to the toxicity of high ppGpp level.

Together, these data demonstrate that a sub-basal but non-zero ppGpp pool, below the detection limit of conventional assays, is sufficient to support growth in minimal medium and that SpoT activity is finely tuned to maintain this pool.

Consistent with this model, systematic truncation of SpoT C-terminal regulatory domains revealed that distinct suppressor mutations alter the contribution of individual domains to growth control (**Extended Data Fig. 6**), without abolishing the requirement for basal ppGpp synthesis. These findings indicate that SpoT H414 functions within a broader intramolecular regulatory network that fine-tunes basal ppGpp production rather than acting as an isolated molecular switch.

Basal ppGpp negatively regulates SpoT abundance

To investigate how SpoT activity is regulated, we examined SpoT protein levels using an antiserum

raised against the C-terminal regulatory domains of SpoT. Immunoblot analysis revealed that SpoT

protein levels were substantially elevated in $spoT^{H414A}$ cells compared to $spoT^{wt}$ and suppressor

strains (**Fig. 6e**).

Because $spoT^{H414A}$ cells produce sub-basal ppGpp levels, whereas suppressor strains restore

physiological function with marginally increased ppGpp, these results suggest that basal ppGpp

negatively regulates SpoT abundance through a feedback mechanism. Such feedback would

stabilize intracellular ppGpp levels by adjusting SpoT protein concentration in response to alarmone

availability. This regulatory logic provides a mechanistic explanation for how SpoT maintains a

narrow basal ppGpp range sufficient for metabolic homeostasis without triggering a full stringent

response.

Suppressor mutations differentially restore global transcriptional programs

To understand how suppressor mutations restore growth and acid resistance, we performed RNA

sequencing on the four suppressor strains. Both parallel coordinates and multidimensional scaling

analysis revealed that strong suppressors (supA and sup41) clustered closely with $spoT^{wt}$ cells,

whereas weaker suppressors (supB and sup17) occupied intermediate positions between $spoT^{H414A}$

and $spoT^{wt}$ (**Extended Data Fig. 7a,7b**).

Consistent with their physiological phenotypes, strong suppressors (supA and sup41) largely

restored expression of acid resistance genes and repressed phage shock and SOS response pathways

(**Extended Data Fig. 7c,7d**). In contrast, weaker suppressors (supB and sup17) exhibited partial

correction of these transcriptional defects (**Extended Data Fig. 8**). Across all suppressors, a core

set of genes involved in amino acid metabolism and stress adaptation was consistently restored

(**Extended Data Fig. 7c**), whereas genes aberrantly induced in $spoT^{H414A}$ cells were broadly

repressed (**Extended Data Fig. 7d**).

These results indicate that suppressor mutations restore SpoT-dependent basal regulation in a

graded manner, with transcriptional recovery correlating with physiological rescue.

SpoT H414 function is conserved in enteric pathogens

Finally, we asked whether the regulatory role of SpoT H414 is conserved in pathogenic

enterobacteria. We introduced the corresponding H414A substitution into SpoT homologs from

Salmonella enterica serovar Typhimurium and *Shigella flexneri* and expressed these proteins in an

E. coli ppGpp⁰ background.

Wild-type SpoT homologs from both species fully restored growth in M9Glc and acid resistance

when expressed in an *E. coli* ppGpp⁰ background, whereas the corresponding H414A variants failed

to support either phenotype (**Fig. 7a–d**). Notably, glutamate supplementation rescued both growth

and acid resistance defects in the *Shigella* SpoT H414A strain but not in the *Salmonella* SpoT

H414A variant. This differential rescue correlates with the degree of sequence divergence in

regulatory regions of SpoT. *S. flexneri* SpoT differs from *E. coli* SpoT at only a single conserved

position (A614G) within the CC domain, whereas *S. Typhimurium* SpoT shares 98.15% overall

identity but contains multiple amino acid substitutions distributed across the C-terminal domains,

including a cluster of nine substitutions within linker-2 between TGS and CC domains (**Extended**

Data Fig. 9). These differences likely modulate intramolecular regulatory interactions within SpoT,

thereby influencing the ability of the H414A variants to respond to glutamate supplementation.

Altogether, these results demonstrate that SpoT H414 plays a conserved role in coordinating growth

and acid resistance across enteric pathogens, underscoring its physiological and evolutionary

significance.

Discussion

Basal ppGpp defines a distinct regulatory regime coordinating metabolism and stress

resistance

The stringent response has classically been viewed as an acute adaptation to amino acid starvation,

driven by RelA-dependent bursts of ppGpp that globally reprogram transcription and metabolism

(**Fig. 7e**, top). Our findings reveal a complementary and previously underappreciated role for SpoT-

derived basal ppGpp as a distinct regulatory regime that operates independently of acute starvation

signalling (**Fig. 7e**, middle). By dissecting the function of a conserved SpoT residue (H414), we

show that maintenance of a sub-basal yet non-zero ppGpp pool is essential for metabolic

homeostasis, growth in minimal medium, and survival under extreme acid stress.

Importantly, the defects observed in *spoT^{H414A}* cells are not attributable to an inability to mount a

canonical stringent response. Rather, they reflect a failure to maintain basal regulatory control over

metabolic flux allocation and stress preparedness (**Fig. 7e**, bottom). This distinction expands the

conceptual framework of ppGpp signalling, positioning SpoT not merely as a backup or modulator

of RelA activity, but as a primary regulator of foundational physiological states.

SpoT-mediated control of glutamate homeostasis integrates growth and acid resistance

A central finding of this study is that SpoT H414 is required to prevent misallocation of metabolic

resources that leads to glutamate depletion. Transcriptomic analysis revealed strong and

coordinated induction of the arginine biosynthesis pathway, which directly consumes glutamate, alongside upregulation of *de novo* purine nucleotide biosynthesis, which indirectly drains glutamate through increased glutamine demand. Genetic and physiological experiments establish arginine biosynthesis as the dominant driver of glutamate depletion, while purine biosynthesis likely exacerbates this imbalance.

Glutamate occupies a unique position in bacterial physiology, serving as a key metabolic hub and as the essential substrate for the glutamate-dependent Gad acid resistance system. Consequently, glutamate depletion provides a unifying explanation for both the growth defect and catastrophic acid sensitivity observed in *spoT^{H414A}* cells. The ability of glutamate supplementation or genetic blockade of arginine biosynthesis to restore growth, together with GadE-dependent rescue of acid resistance, underscores the centrality of glutamate homeostasis in SpoT-mediated regulation.

Basal ppGpp tunes stress preparedness rather than viability

Our data demonstrate that SpoT-derived basal ppGpp does not simply support cell viability, but instead primes cells for environmental stress. Despite severe repression of acid resistance genes, *spoT^{H414A}* cells remain viable under neutral, well-buffered conditions, indicating that SpoT H414 is specifically required for stress preparedness rather than basal survival. This distinction explains why growth defects and acid resistance defects, while mechanistically linked through glutamate homeostasis, manifest under different environmental conditions.

Notably, RelA is dispensable for acid resistance under the conditions tested, highlighting a division of labor between stringent response enzymes: RelA mediates acute starvation responses, whereas SpoT maintains basal regulatory states necessary for survival in fluctuating environments.

SpoT intrinsically tunes a narrow basal ppGpp window

Intragenic suppressor analysis reveals that SpoT encodes an intrinsic regulatory module capable of

fine-tuning basal ppGpp levels. Suppressor mutations distributed across both the catalytic and

regulatory regions of SpoT restore growth and acid resistance without producing large, readily

detectable increases in ppGpp, indicating that very low, rapidly turned-over ppGpp pools are

physiologically sufficient for function.

Perturbation of SpoT's enzymatic balance further supports this model. Inactivation of ppGpp

synthesis abolishes suppression, whereas inactivation of hydrolase activity elevates ppGpp levels

and accelerates growth recovery. Together, these results demonstrate that SpoT actively maintains a

narrow basal ppGpp range and that deviations below this threshold have profound physiological

consequences.

Consistent with this quantitative tuning model, systematic truncation of SpoT C-terminal regulatory

domains revealed that distinct suppressor mutations differentially alter the contribution of

individual domains to growth control, without abolishing the requirement for basal ppGpp

synthesis. These findings indicate that SpoT H414 functions within a broader intramolecular

regulatory network that modulates ppGpp production in a graded manner rather than acting as a

binary on–off switch.

Finally, the observed negative correlation between basal ppGpp levels and SpoT protein abundance

suggests the existence of a feedback mechanism that stabilizes this regulatory regime. Such

feedback would allow cells to dynamically adjust SpoT abundance and activity to maintain

metabolic balance without triggering a full stringent response.

Conservation of SpoT H414 underscores physiological relevance

The functional conservation of SpoT H414 in *Salmonella* and *Shigella* underscores the broader

significance of basal ppGpp regulation in enteric bacteria. While species-specific differences in

SpoT regulatory domains modulate the extent to which glutamate supplementation can compensate

for H414A dysfunction, the core requirement for this residue in supporting growth and acid

resistance is conserved. This conservation highlights basal ppGpp as a fundamental determinant of

bacterial fitness in acidic environments, such as those encountered during host colonization.

Implications and outlook

Together, our findings establish SpoT as a central coordinator of metabolic homeostasis and stress

resistance through maintenance of a basal ppGpp pool. This regulatory regime operates

independently of the acute, RelA-mediated stringent response and integrates amino acid metabolism

with acid resistance, thereby enabling bacterial survival under hostile environmental conditions.

Several important questions remain. Our domain truncation and suppressor analyses indicate that

SpoT H414 functions within a distributed intramolecular regulatory network that quantitatively

tunes basal ppGpp production, rather than acting as a simple on–off switch; however, the precise

nature of the signal sensed by H414, whether metabolic, physicochemical, or intramolecular,

remains unresolved. In addition, defining the absolute intracellular ppGpp concentrations required

for basal regulation will require new methodologies capable of resolving low-abundance alarmone

pools with high temporal resolution, ideally through single-cell reporters coupled to dynamic

growth measurements. More broadly, elucidating how basal ppGpp differentially modulates

specific metabolic and stress-response pathways will be essential for fully understanding this

regulatory logic.

Given the central role of acid resistance in enteric pathogenesis, SpoT-mediated basal ppGpp

regulation represents a potential vulnerability that could be exploited to compromise bacterial

survival. More generally, our work highlights the physiological importance of low-level signaling

states and suggests that basal regulatory regimes may play underappreciated roles in coordinating

metabolism and stress adaptation across diverse microbial systems.

Figures and legends

a

b

Figure 1. A conserved SpoT H414A mutation reveals defective growth on M9Glc medium. (a) Growth of wild type *E. coli* MG1655 (*relA^{wt} spoT^{wt}*), and Δ*relA spoT^{H414A}*, Δ*relA spoT^{wt}* strains on LB and M9Glc solid media. **(b)** Growth dynamics of the same strains in liquid LB and M9Glc media.

Figure 2. Transcriptomic analysis reveals collapse of acid resistance programs. (a) Heatmap of genes encoded in the *E. coli* acid fitness island in *spoT^{H414A}* relative to *spoT^{wt}*. **(b)** Genetic organization of the *E. coli* acid fitness island. (**c**) Acid resistance assay of wild type MG1655 (*relA^{wt} spoT^{wt}*)*, ΔrelA spoT^{H414A}* and *ΔrelA spoT^{wt}*. Survival of indicated strains following challenge at pH 2.5, expressed as the percentage of colony-forming units (CFUs) remaining relative to starting CFUs for each strain. Data are means ± s.d. from biological replicates. u.d., undetectable. Statistical significance was assessed using unpaired two-tailed Student's *t*-tests. **** denote *P* < 0.0001.

Figure 3. Arginine metabolic overflow drives glutamate depletion in *spoT^H414A*. (a) Heatmap of genes involved in amino acid biosynthesis, comparing *spoT^H414A* and *spoT^wt* strains. The arginine biosynthesis genes are highlighted. **(b)** Schematic of the arginine biosynthesis pathway in *E. coli*.

Genes upregulated in *spoT^{H414A}* cells are indicated by red upward arrows, accompanied by average

log$_2$ fold-change values relative to *spoT^{wt}*. **(c)** Normalized lag phase duration of *spoT^{H414A}* cells

grown in M9Glc medium with supplementation of arginine (5.2 mM, R), uracil (0.2 mM, U),

glutamate (0.6 mM, E), or the indicated combinations, expressed relative to the *spoT^{wt}* strain. **(d)**

Normalized lag phase duration of indicated strains grown in M9Glc medium with supplementation

of arginine (R), glutamate (E), or both, relative to *spoT^{wt}*. Red dots indicate failure to resume

growth within the 72-h observation window. **(e)** Normalized lag phase duration of *spoT^{wt}* strains

harboring pCA24N-*argA^{wt}* or the feedback-resistant pCA24N- *argA^{H15Y}* allele, grown with or

without IPTG induction and/or glutamate supplementation (0.6 mM), relative to the uninduced

argA^{wt} control. Data are means ± s.d. from biological replicates. Statistical significance was

assessed using unpaired two-tailed Student's *t*-tests. ****, *** denote $P < 0.0001$ and 0.001,

respectively.

Figure 4. SpoT H414 is required for survival under extreme acid stress.

(a) Acid resistance assay of $\Delta relA\ spoT^{H414A}$ and $\Delta relA\ spoT^{wt}$ as in Figure 2c. Where indicated

(+E), glutamate (0.6 mM) was included in the acid challenge medium. u.d., undetectable. **(b)**

Survival of $spoT^{H414A}$ and $spoT^{wt}$ strains following 30 min challenge in buffers adjusted to the

indicated pH values, normalized to starting CFUs. Data are means ± s.d. from biological replicates.

Statistical significance was assessed using unpaired two-tailed Student's *t*-tests. ****, ***, **, *

denote $P < 0.0001$, 0.001, 0.01, and 0.05, respectively.

Figure 5. Intragenic SpoT suppressor mutations restore growth and acid resistance. (a)

Schematic illustrating the isolation of spontaneous suppressor mutations that enable growth of

$spoT^{H414A}$ cells on M9Glc medium. **(b)** Domain architecture of the SpoT protein, indicating the

locations of suppressor mutations. Amino-acid positions are shown below. L1/L2, linker-1/linker-2;

HYD, hydrolase domain; SYN, synthetase domain; TGS, ThrRS–GTPase–SpoT domain; CC,

conserved cysteine domain; ACT, aspartate kinase–chorismate mutase–TyrA domain. **(c)** Growth

dynamics of four suppressor strains (supA, supB, sup17, sup41) compared with $spoT^{H414A}$, $spoT^{wt}$,

ppGpp⁰ ($\Delta relA\ \Delta spoT$), and wild-type MG1655 ($relA^{wt}\ spoT^{wt}$). Also shown are suppressor strains

in which $spoT$ was deleted ($spoT207::cat$). **(d)** Survival of indicated strains following challenge at

pH 2.5, expressed as normalized CFUs. Where indicated (+E), glutamate (0.6 mM) was included.

Data are means ± s.d. Statistical significance was assessed using unpaired two-tailed Student's *t*-

tests. ***, **, * denote $P < 0.001$, 0.01, and 0.05, respectively.

Figure 6. A sub-basal ppGpp pool is sufficient for physiological function. (a) Thin-layer

chromatography (TLC) analysis of [32]p-labeled nucleotides from indicated strains grown in low-

phosphate MOPS minimal medium supplemented with 0.2% glucose. NC and PC denote *ΔrelA*

ΔspoT and valine-treated wild-type MG1655 controls, respectively. Red circles mark ppGpp; boxed

regions indicate nonspecific spots also present in *ΔrelA ΔspoT* samples. **(b)** Effect of inactivating

SpoT ppGpp synthetase activity (E319Q; HYD) on growth recovery in M9Glc. Lag phase duration

is normalized to *spoT^wt*. **(c)** TLC analysis of [32]p-labeled nucleotides from strains carrying SpoT

hydrolase-inactive mutations (H72A D73A; SYN). **(d)** Growth recovery of hydrolase-inactive

 strains (SYN), quantified as normalized lag phase duration relative to $spoT^{wt}$. Note that the

hydrolase-inactive (SYN) variants of supA and sup17 could not be obtained, thereby not tested. **(e)**

Immunoblot analysis of SpoT protein levels using an antiserum raised against the SpoT C-terminal

(TGS–CC–ACT; anti-TCA) region. Cells were grown in M9Glc to early exponential phase

($OD_{600nm} \approx 0.2$) prior to sampling. $spoT^{H414A}$ samples were collected after prolonged incubation, as

in Extended Data Fig. 1c. $\Delta relA\ \Delta spoT$ cells grown in LB served as a negative control.

Figure 7. Conservation of SpoT H414 function and model for basal ppGpp regulation. (a,b)

Growth of *E. coli* ppGpp⁰ strains complemented with SpoT homologs from *Salmonella enterica*

serovar Typhimurium (**a**) or *Shigella flexneri* (**b**). E, glutamate (0.6 mM); EV, empty vector. Data

represent at least three biological replicates. **(c,d)** Survival of complemented ppGpp⁰ strains

(*Salmonella* (**c**) or *Shigella* (**d**)) following challenge at pH 2.5, normalized to starting CFUs. Where

indicated (+E), glutamate (0.6 mM) was included. **(e)** Under acute amino-acid starvation (top),

RelA is rapidly activated by ribosome-associated signals to generate a high-amplitude ppGpp surge,

globally reprogramming transcription and metabolism to suppress nucleotide pathways and promote

amino-acid availability. Under minimal medium growth (middle), SpoT maintains a low but

physiologically essential basal ppGpp pool that is distinct from and independent of RelA. This basal

ppGpp state coordinates metabolic homeostasis by restraining excessive arginine and purine

biosynthesis, preserving intracellular glutamate levels, and priming multiple acid resistance

programs, including the glutamate-dependent Gad system (AR). In $spoT^{H414A}$ cells (bottom), a sub-

basal level of ppGpp remains below the threshold required for effective metabolic and stress

control. Therefore, arginine and purine biosynthetic pathways become overactivated, intracellular

glutamate is depleted, and acid resistance programs are inadequately induced, resulting in growth

failure in minimal medium and extreme sensitivity to acid stress.

Extended Data Fig. 1 (a) Sequence alignment of 12 TGS domains of RelA, SpoT and their homologous proteins from α, β, γ and δ proteobacteria, Firmicute and Actinomycete. The highly conserved position corresponding to His 432 of RelA and 414 of SpoT of *E. coli* is indicated with an arrow. **(b)** Superpositioning of the TGS domains of RelA and SpoT from *E. coli* (rmsd over 91 of 98 atom pairs is 0.762 Å) generated by AF3. **(c)** Representative growth curves of Δ*relA* *spoT*^{H414A}, Δ*relA* *spoT*^{wt} and the four suppressor strains in M9Glc media for collecting RNA sequencing samples. Cells were collected when their OD$_{600nm}$ approach ca. 0.2, except for *spoT*^{H414A} cells, which could not grow in this medium, thus these cells were collected after incubated for 18 hrs in the same medium. Coloured arrows indicate the sampling points for each strain. **(d)** Heatmap

of genes with greater than 16-fold changes between Δ*relA spoT^{H414A}* and Δ*relA spoT^{wt}* cells growing

in M9Glc medium. **(e)** Gene ontology groups of genes downregulated in the *spoT^{H414A}* strain.

Genes downregulated at least 16-fold were extracted using DEGUST (|Log2(FC)| > 4; FDR < 0.05)

and analysed by AmiGO and Panther. The 58 genes downregulated more than 16-fold (Log2(FC) <

-4) are listed in **Table S3**. **(f)** Gene ontology groups of genes upregulated at least 16-fold in the

spoT^{H414A} strain. The 28 genes upregulated more than 16-fold (Log2(FC) > 4) are listed in **Table**

S3. Note the differences in the Fold Enrichment scales in **(e)** and **(f)**. Gene names in bold in (**e, f**)

are strongly up or downregulated. **(g)** Normalized percentage of surviving viable cells of the *spoT^{wt}*

and *spoT^{H414A}* strains growing in M9Glc medium. Two replicates were performed and presented.

Extended Data Fig. 2 (a) Purine nucleotide de novo biosynthesis pathway in *E. coli*. The

upregulated genes are indicated by a red upward arrow alongside average log2(FC) values of

$spoT^{H414A}$ versus $spoT^{wt}$. R5P, ribose-5-phosphate; PRPP, 5-phosphoribosyl 1-pyrophosphate; PRA,

5-phosphoribosylamine; GAR, 5'-phosphoribosylglycinamide; FGAR, 5'-phosphoribosyl-N-

formylglycineamide; FGAM, 5'-phosphoribosylformylglycinamidine; AIR, 5-aminoimidazole

ribonucleotide; C-AIR, 5-carboxyamino-1-(5-phospho-D-ribosyl)imidazole; SAICAR, 1-(5-

phosphoribosyl)-4-(N-succino-carboxamide)-5-aminoimidazole; AICAR, 5-amino-1-(5-phospho-D-

ribosyl)imidazole-4-carboxamide; FAICAR, 5-formamido-1-(5-phospho-D-ribosyl)imidazole-4-

carboxamide; IMP, inosine 5'- monophosphate. **(b)** Heatmap of genes involved in purine and

pyrimidine nucleotide de novo biosynthesis. The red box indicates purine de novo biosynthesis

genes.

Extended Data Fig. 3 (a) Normalized lag phase length of the *spoT^H414A* strain without or with

supplemented each of the 20 amino acids (in single letters; A, alanine 0.8mM; C, cysteine 0.1mM;

D, aspartic acid 0.4mM; E, glutamate 0.6mM; F, phenylalinine 0.4mM; G, glycine 0.8mM; H,

histidine 0.2mM; I, isoleucine 0.4mM; K, lysine 0.4mM; L, leucine 0.8mM; M, methionine 0.2mM;

N, asparigine 0.4mM; P, proline 0.4mM; Q, glutamine 0.6mM; R, arginine 5.2mM; S, serine 10mM;

T, threonine 0.4mM; V, valine 0.6mM; W, tryptophan 0.1mM; Y, tyrosine 0.2mM), relative to the

spoT^wt strain in M9Glc medium. **(b)** Normalized lag phase length of the *spoT^H414A* strain without or

with supplemented 1x, 2x, 4x, 10x glutamate (E, 1x = 0.6 mM), relative to the *spoT^wt* strain in M9Glc

medium. **(c)** Normalized lag phase length of the *spoT^H414A* strain without or with supplemented

glutamate (0.6 mM) together with glutamine (Q, 0.6 mM), proline (P, 0.4 mM), asparagine (N, 0.4

mM), relative to the *spoT^wt* strain in M9Glc medium. **(d)** Representative growth curves of Δ*relA*

spoT^H414A, Δ*relA spoT^wt* strains without or with all 20 amino acids (+20AA) (see method for details)

in M9Glc medium. Two biological replicates each with two technical replicates were performed and

presented. **(e)** Normalized lag phase length of the Δ*relA*Δ*spoT207::cat* (ppGpp^0) strain without or

with supplemented arginine (5.2 mM, R), uracil (0.2 mM, U), glutamate (0.6 mM, E) or their varied

combinations, relative to the *spoT^wt* strain in M9Glc medium. Red dots indicate fails of regrowth

within the tested time range of 72 hrs.

Extended Data Fig. 4 (a) Normalized percentage of surviving cells of various strains after

challenged with pH=2.5. The data were normalized to the starting CFUs of each respective tested

strain. +K, +R, lysine (0.6 mM), arginine (0.6 mM) included in the pH=2.5 medium, respectively.

(b) Normalized percentage of surviving cells of $spoT^{wt}$ or $spoT^{H414A}$ strains containing various

plasmid vectors (pCA24N [41], pNTR-SD [42]) after challenged with pH=2.5. The data were

normalized to the starting CFUs of each respective tested strain. EV, empty vector pCA24N. +E,

glutamate (0.6 mM) included in the pH=2.5 medium. **(c)** Representative growth curves of $spoT^{H414A}$

and $spoT^{wt}$ strains containing either the empty vector (EV) or a *gadE* gene on pCA24N or pNTR-SD

vectors in M9Glc medium. Two biological replicates each with two technical replicates were

performed and presented. **(d)** Normalized percentage of surviving cells of $spoT^{wt}$ and $spoT^{H414A}$

strains after challenged with media of various pH values. The data were normalized to the starting

CFUs of each respective tested strain. **(e)** Heatmap of genes involved in acid resistance. Colored

bars indicate genes belonging to the same acid resistance systems: orange, *nuo* genes encoding

NADH dehydrogenase I; yellow, *cyo* genes encoding cytochrome *bo* oxidase; green, lysine

dependent acid resistance systems; cyan, ornithine dependent acid resistance system; black, alkaline

stress; dark blue, arginine dependent acid resistance system.

Extended Data Fig. 5 Growth dynamics of re-constructed four suppressors (cyan)

(*supA/supB/sup17/sup41*) in comparison to *spoT^{H414A}*, *spoT^{wt}*, and the four random suppressors

(blue).

Extended Data Fig. 6 (a) Schematic show of sequential domain truncations of the *spoT* alleles in

various strains. **(b-g)** Normalized lag phase length (in percentage) relative to the respective *spoT*-

full-length (FL) parental strains during growth in M9Glc. Red dots indicate that the respective

strains did not grow up in M9Glc within the tested time frame of 72 hrs. Red crosses indicate the

strains could not be made. **(h)** Sum-up of the effect of domain deletions on cell regrowth thus

potential regulation of SpoT activity from panels **b-g**. Blue and red blocks indicate positive and

negative effects respectively. Red crosses indicate strains impossible to obtain. Grey blocks indicate

no effect can be deduced due to insignificant/indistinguishable growth curves. **(i-n)** Normalized lag

phase length (in percentage) relative to the respective *spoT*-full-length (FL) parental strains during

growth in M9Glc and supplemented with glutamate (0.6 mM). **(o)** Sum-up of the effect of domain

deletions on cell regrowth in the presence of supplemented glutamate (0.6 mM) thus potential

regulation of SpoT activity from panels **i-n**. Color coding is same as panel **h**.

Extended Data Fig. 7 (a) Parallel Coordinates plot of the $spoT^{wt}$, $spoT^{H414A}$ and the four suppressor

strains with |Log2(FC)| > 4 and FDR < 0.05. (**b**) MDS plot of the six strains showing individual

samples. Cell growth and sampling are described in Materials and Methods. Panels **a** and **b** were

both generated by DEGUST. (**c**) (Top) Venn diagram of genes downregulated in the $spoT^{H414A}$

strain and upregulated in either of the four suppressor strains. (middle) Genes downregulated in the

$spoT^{H414A}$ strain and upregulated in all four suppressor strains. (bottom) Gene Ontology analysis of

the 124 genes downregulated in the $spoT^{H414A}$ strain and upregulated in all four suppressor strains.

All genes are listed in **Table S4. (d)** (Top) Venn diagram of genes upregulated in the $spoT^{H414A}$

strain and downregulated in either of the four suppressor strains**. (middle)** Genes upregulated in the

$spoT^{H414A}$ strain and downregulated in all four suppressor strains**. (bottom)** Gene Ontology analysis

of the 779 genes upregulated in the $spoT^{H414A}$ strain and downregulated in all four suppressor

strains. All genes are listed in **Table S4.** Genes were extracted from DEGUST using settings of

|Log2(FC)| > 1 and FDR < 0.05. Gene names in bold in GO term are strongly up or downregulated.

Extended Data Fig. 8 (a) Heatmap of $spoT^{H414A}$ versus $spoT^{wt}$ (a setting of |Log2(FC)| > 3.96 and

FDR < 0.05 returned 100 of a total of 4,575 genes analyzed in DEGUST). Highlighted genes

represent phage shock proteins (*psp*, blue), SOS response genes (magenta), acid stress response

genes (deep purple), and stationary-phase-induced genes (gold). Genes marked with an asterisk (*)

are known ppGpp-inducible genes. (**b, e, h, k**) Heatmaps of $spoT^{H414A}$ versus the suppressor strains.

FDR settings were identical to those used in (**a**) while the |Log2(FC)| values were set to yield

approximately 100 genes to be returned. DEGUST returned 102 genes for *supA* (|Log2(FC)| > 3.9),

97 for *sup41* (|Log2(FC)| > 3.4), 94 for *supB* (|Log2(FC)| > 2.6) and 91 for *sup17* (|Log2(FC)|>2.3).

The actual genes returned by DEGUST are listed in **Table S5**. (**c, d, f, g, i, j, l, m**) Venn diagram

analysis of genes dysregulated in $spoT^{H414A}$ and the four suppressor strains. Panel **a** yielded the

genes up- and down- regulated in $spoT^{H414A}$ relative to the $spoT^{wt}$ strain and these genes were used

throughout the Venn diagram analysis.

a

Hydrolase Domain (45 - 144)

```
SpoT_E_coli   1 MYLFESLNQLIQTYLPEDQIKRLRQAYLVARDAHEGQTRSSGEPYITHPVAVACILAEMKLDYETLMAALLHDVIEDTPATYQDMEQLFGKSVAELVEGVS
SpoT_S_flex   1 MYLFESLNQLIQTYLPEDQIKRLRQAYLVARDAHEGQTRSSGEPYITHPVAVACILAEMKLDYETLMAALLHDVIEDTPATYQDMEQLFGKSVAELVEGVS
```

```
SpoT_E_coli 102 KLDKLKFRDKKEAQAENFRKMIMAMVQDIRVILIKLADRTHNMRTLGSLRPDKRRRIARETLEIYSPLAHRLGIHHIKTELEELGFEALYPNRYRVIKEVV
SpoT_S_flex 102 KLDKLKFRDKKEAQAENFRKMIMAMVQDIRVILIKLADRTHNMRTLGSLRPDKRRRIARETLEIYSPLAHRLGIHHIKTELEELGFEALYPNRYRVIKEVV
```

Synthase Domain 152 - 325)

```
SpoT_E_coli 203 KAARGNRKEMIQKILSEIEGRLQEAGIPCRVSGREKHLYSIYCKMVLKEQRFHSIMDIYAFRVIVNDSDTCYRVLGQMHSLYKPRPGRVKDYIAIPKANGY
SpoT_S_flex 203 KAARGNRKEMIQKILSEIEGRLQEAGIPCRVSGREKHLYSIYCKMVLKEQRFHSIMDIYAFRVIVNDSDTCYRVLGQMHSLYKPRPGRVKDYIAIPKANGY
```

Linker 1 (347 - 387)

```
SpoT_E_coli 304 QSLHTSMIGPHGVPVEVQIRTEDMDQMAEMGVAAHWAYKEHGETSTTAQIRAQRWMQSLLELQQSAGSSFEFIESVKSDLFPDEIYVFTPEGRIVELPAGA
SpoT_S_flex 304 QSLHTSMIGPHGVPVEVQIRTEDMDQMAEMGVAAHWAYKEHGETSTTAQIRAQRWMQSLLELQQSAGSSFEFIESVKSDLFPDEIYVFTPEGRIVELPAGA
```

TGS Domain (388 - 482)   Linker 2 (483 - 555)

```
SpoT_E_coli 405 TPVDFAYAVHTDIGHACVGARVDRQPYPLSQPLTSGQTVEIITAPGARPNAAWLNFVVSSKARAKIRQLLKNLKRDDSVSLGRRLLNHALGGSRKLNEIPQ
SpoT_S_flex 405 TPVDFAYAVHTDIGHACVGARVDRQPYPLSQPLTSGQTVEIITAPGARPNAAWLNFVVSSKARAKIRQLLKNLKRDDSVSLGRRLLNHALGGSRKLNEIPQ
```

Coiled-coil Domain (556 - 627)

```
SpoT_E_coli 506 ENIQRELDRMKLATLDDLLAEIGLGNAMSVVVAKNLQHGDASIPPATQSHGHLPIKGADGVLITFAKCCRPIPGDPIIAHVSPGKGLVIHHESCRNIRGYQ
SpoT_S_flex 506 ENIQRELDRMKLATLDDLLAEIGLGNAMSVVVAKNLQHGDASIPPATQSHGHLPIKGADGVLITFAKCCRPIPGDPIIAHVSPGKGLVIHHESCRNIRGYQ
```

ACT Domain (628 - 702)

```
SpoT_E_coli 607 KEPEKFMAVEWDKETAQEFITEIKVEMFNHQGALANLTAAINTTTSNIQSLNTEEKDGRVYSAFIRLTARDRVHLANIMRKIRVMPDVIKVTRNRN 702
SpoT_S_flex 607 KEPEKFMGVEWDKETAQEFITEIKVEMFNHQGALANLTAAINTTTSNIQSLNTEEKDGRVYSAFIRLTARDRVHLANIMRKIRVMPDVIKVTRNRN 702
```

b

Hydrolase Domain (45 - 144)

```
SpoT_E_coli   1 MYLFESLNQLIQTYLPEDQIKRLRQAYLVARDAHEGQTRSSGEPYITHPVAVACILAEMKLDYETLMAALLHDVIEDTPATYQDMEQLFGKSVAELVEGVS
SpoT_S_ent.   1 MYLFESLNQLIQTYLPEDQIKRLRQAYLVARDAHEGQTRSSGEPYITHPVAVACILAEMKLDYETLMAALLHDVIEDTPATYQDMEQLFGKSVAELVEGVS
```

```
SpoT_E_coli 102 KLDKLKFRDKKEAQAENFRKMIMAMVQDIRVILIKLADRTHNMRTLGSLRPDKRRRIARETLEIYSPLAHRLGIHHIKTELEELGFEALYPNRYRVIKEVV
SpoT_S_ent. 102 KLDKLKFRDKKEAQAENFRKMIMAMVQDIRVILIKLADRTHNMRTLGSLRPDKRRRIARETLEIYSPLAHRLGIHHIKTELEELGFEALYPNRYRVIKEVV
```

Synthase Domain 152 - 325)

```
SpoT_E_coli 203 KAARGNRKEMIQKILSEIEGRLQEAGIPCRVSGREKHLYSIYCKMVLKEQRFHSIMDIYAFRVIVNDSDTCYRVLGQMHSLYKPRPGRVKDYIAIPKANGY
SpoT_S_ent. 203 KAARGNRKEMIQKILSEIEGRLQEAGIPCRVSGREKHLYSIYCKMVLKEQRFHSIMDIYAFRVIVHDSDTCYRVLGQMHSLYKPRPGRVKDYIAIPKANGY
```

Linker 1 (347 - 387)

```
SpoT_E_coli 304 QSLHTSMIGPHGVPVEVQIRTEDMDQMAEMGVAAHWAYKEHGETSTTAQIRAQRWMQSLLELQQSAGSSFEFIESVKSDLFPDEIYVFTPEGRIVELPAGA
SpoT_S_ent. 304 QSLHTSMIGPHGVPVEVQIRTEDMDQMAEMGVAAHWAYKEHGETSTTAQIRAQRWMQSLLELQQSAGSSFEFIESVKSDLFPDEIYVFTPEGRIVELPAGA
```

TGS Domain (388 - 482)   Linker 2 (483 - 555)

```
SpoT_E_coli 405 TPVDFAYAVHTDIGHACVGARVDRQPYPLSQPLTSGQTVEIITAPGARPNAAWLNFVVSSKARAKIRQLLKNLKRDDSVSLGRRLLNHALGGSRKLNEIPQ
SpoT_S_ent. 405 TPVDFAYAVHTDIGHACVGARVDRQPYPLSQPLSSGQTVEIITAPGARPNAAWLNFVVSSKARAKIRQLLKNLKRDDSVSLGRRLLNHALGGSRKLAEIPQ
```

Coiled-coil Domain (556 - 627)

```
SpoT_E_coli 506 ENIQRELDRMKLATLDDLLAEIGLGNAMSVVVAKNLQHGDASIPPATQS-HGHLPIKGADGVLITFAKCCRPIPGDPIIAHVSPGKGLVIHHESCRNIRGY
SpoT_S_ent. 506 ENIQRELDRMKLATLDDLLAEIGLGNAMSVVVAKNLQQGEAVVPTVAQSNHGHLPIKGADGVLITFAKCCRPIPGDPIIAHVSPGKGLVIHHESCRNIRGY
```

ACT Domain (628 - 702)

```
SpoT_E_coli 606 QKEPEKFMAVEWDKETAQEFITEIKVEMFNHQGALANLTAAINTTTSNIQSLNTEEKDGRVYSAFIRLTARDRVHLANIMRKIRVMPDVIKVTRNRN 703
SpoT_S_ent. 607 QKEPEKFMAVEWDKETEQEFITEIKVEMFNHQGALANLTAAINTTTSNIQSLNTEEKDGRVYSTFIRLTARDRVHLANIMRKIRVMPDVIKVTRNRN 703
```

Extended Data Fig. 9 Pairwise sequence comparison between *E. coli* SpoT_{ec} and SpoT from *S. typhimurium* SL1334 and *Shigella flexneri*. *Shigella flexneri* SpoT_{sf} is nearly identical as *E. coli* SpoT_{ec} except for a conserved mutation of Alanine 614 to Glycine on the CC domain. *S. typhimurium* SL1334 SpoT_{st} (702 aa) shares 98.15% identity with *E. coli* SpoT_{ec} (701 aa), with several amino acid variations on the HYD (N267H), TGS (T437S), L2 (nine mutations, N501A,

595   H542Q, D544E, S546V, I547V, P549T, A550V, T551A, an insertion of N between 553-554), CC

596   (A621E), and ACT (A668T) domains.

ACKNOWLEDGEMENTS

We thank Paulina Grucella for helping with Western blot. The work is supported by a Novo Nordisk Foundation Project Grant (NNF19OC0058331), a Danmarks Frie Forskningsfond grant (2032-00030B) to Y.E.Z, the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement (Nº 801199) to M.L.S, and the Chinese Scholarship Council program (Nº 202206870009) to Y.Y.L.

AUTHOR CONTRIBUTIONS

Y.E.Z designed this study; Y.Y.L, M.L.S and Y.E.Z acquired fundings; Y.Y.L, M.L.S, Y.E.Z acquired experimental data; X.C.Z, L.D, K.G, Y.E.Z performed the bioinformatics analyses; Y.E.Z, K.G, wrote the draft manuscript; and all authors analysed the data and edited the manuscript.

DECLARATIONS OF INTERESTS

The authors declare that they have no conflicts of interest.

EXPERIMENTAL MODEL AND SUBJECT DETAILS

E. coli DH5α (Stratagene) was used for cloning and *E. coli* K-12 MG1655 for experiments requiring an *E. coli* wt strain background (see **Table S1** for further details). For cloning, nutrient broth (Oxoid), with agar and antibiotics (chloramphenicol (25 µg/ml), ampicillin (100 µg/ml), and/or kanamycin (25 µg/ml)) when appropriate, was used. The different MG1655 strains were cultivated in Lysogeny Broth (LB) as well as M9Glc media (1x M9 salt, pH 7.4 1 mM MgSO4, 0.1 mM CaCl$_2$, Thiamine (0.001 mg/1 ml), 0.2 % (g/ml) glucose) at 37 °C with agitation (160 rpm).

METHOD DETAILS

Isolation of suppressor mutations rescuing the growth of *spoT^{H414A}* in M9Glc

The *spoT^{H414A}* (YZ527) strain was grown overnight in LB media and then washed twice with PBS

buffer. Approx. 10^7 cells were plated on one M9Glc plate and incubated at 30°C. Colonies of different

sizes were selected between 24 – 72 hours of incubation, and the restored growth was confirmed in

solid and liquid M9Glc medium.

Growth assay in liquid minimal media

To measure the growth curves of various strains in M9Glc media, the OD_{600nm} was measured of

overnight cultures grown in LB of each strain, washed twice with 1x Phosphate Buffered Saline (PBS)

and normalised in PBS and inoculated into fresh M9Glc media with a starting $OD_{600nm} = 0.05$ unless

otherwise stated. The growth was measured every 15 min for 72 hrs in a plate reader (BioTek)

presetted at 30°C with double orbital agitation at 548 cpm.

Measurements of (p)ppGpp by autoradiography

(p)ppGpp is a highly labile molecule not amenable to HPLC or mass spectrometry analysis. Therefore,

thin layer chromatography (TLC) was used to quantify both pppGpp and ppGpp. The excess

phosphates in M9Glc affect efficient labelling of nucleotides by $H_3{}^{32}PO_4$, therefore MOPS-M9Glc

hybrid media was prepared wherein the PBS buffer in M9Glc was replaced by the MOPS buffer of

the MOPS media[43] (i.e., containing MOPS-tricine 40 mM pH 7.2, 1 mM MgSO4, 0.1 mM CaCl2,

Thiamine (0.001 mg/1 ml), 0.2 % (g/ml) glucose, NH4Cl 0.001 g/ml, 8.5 mM NaCl, 0.2 mM KHPO4).

The strains were grown for ca. 18 hrs in MOPS-M9Glc media at 37 °C with agitation. Afterwards,

the strains were adjusted to the OD_{600nm} of 0.05 in MOPS-M9Glc and grown at 37 °C with agitation

until OD_{600nm} of 0.2-0.3 was reached. Then, 100 μl cells were transferred to a 2-ml Eppendorf tube

together with 5 μl of 0.005 mCi $H_3{}^{32}PO_4$ (100 μCi/ml, PerkinElmer). After 1h at 37 °C with 600 rpm

agitation, 90 µl were transferred to a new tube with 20 µl 2N formic acid, and the sample was stored

at -20°C until analysed by TLC as described in[18].

Sample preparation for RNA-sequencing

The strains were grown overnight in LB media before washed twice in PBS and inoculated into

M9Glc media (OD_{600nm} = 0.05). The strains were grown in M9Glc media until the early exponential

phase (approximately OD_{600nm} = 0.2) before the cells were collected, and snap frozen with liquid

nitrogen and stored at -80 °C. As the strain $spoT^{H414A}$ is unable to grow in M9Glc media, the starting

OD_{600nm} was set to 0.2, and the cells were collected when the other strains are ready to harvest, i.e.

ca. 18 hrs.

Total RNA was extracted using the RNeasy Mini Kit (Qiagen; 74104). Ribosomal RNA (rRNA) was

depleted from purified total RNA with Ribo-Zero Plus rRNA Depletion kit (Illumina; 20040892).

First and second strand complementary DNA (cDNA) were synthesized by cDNA Synthesis kit

(Illumina; 20040895). RNA Index Anchors (Illumina; 20040899) was used for ligation of pre-index

anchors to the ends of the double-stranded cDNA fragments to enable dual indexing. Barcoded cDNA

libraries were constructed from anchor-ligated DNA fragments and sequenced on Illumina NovaSeq

6000 Sequencing System from Genewiz (Azenta Life Sciences), yielding a median depth of

39,619,482 paired-end 150 bp reads per sample.

Strain construction

All strains and oligonucleotides used in this study are listed in **Tables S1** and **S2**, respectively. All

chromosomal *in situ* mutations of *spoT* alleles were made by using the I-SceI-based λ -Red

recombineering method[44].

The *spoT* E319Q and H72A D73A mutant strains

Briefly, the pWRG99 plasmid was transformed into the target strains (YZ38, YZ359, YZ543, YZ544,

YZ545, YZ546) creating the target strains YZ313, YZ345, YZ826, YZ827, YZ828, and YZ829. The

SceI site of the pWRG100 vector was PCR amplified using the primers pYZ611 and pYZ612 for the

E319Q mutation and the primers pYZ617 and pYZ618 for the H72A D73A mutation. These PCR

DNAs were then electroporated into the above strains (YZ38, YZ359, YZ543, YZ544, YZ545,

YZ546). Then, the oligos pYZ613 and pYZ616 were used to introduce the E319Q and H72A D73A

mutations into the specific *spoT* alleles *in situ*, generating the E319Q mutant strains (YZ872, YZ873,

YZ874, YZ875, YZ876) and H72A D73A mutant strains (YZ900, YZ902, YZ904, YZ905).

The C-terminal serial truncated mutant strains of *spoT*

pYZ796 and pYZ797 were used to amplify the SceI from pWRG100 and introduced into YZ313,

YZ345, YZ826, YZ827, YZ828, and YZ829. The serial truncations were performed using the

repairing oligos pYZ798, pYZ799, pYZ901, pYZ801, pYZ902, deleting the according domains

(ACT, ACT-CC, ACT-CC-Linker2, ACT-CC-Linker2-TGS, ACT-CC-Linker2-TGS-Linker1,

respectively).

Reconstruction of the *spoT^{H414A}* suppressor mutations

Primers pYZ619 and pYZ620 were used to amplify the SceI DNA from pWRG100 vector before it

was introduced into the strain *spoT^{wt}* (YZ313). Then, the primers pYZ328 and pYZ70 were used to

amplify the corresponding *spoT* alleles using the chromosomal DNA of the strains supA (YZ543),

supB (YZ544), sup17 (YZ545), and sup41 (YZ546) respectively as the templates. These DNAs were

used to introduce the specific *spoT* suppressing mutations.

P1 phage transduction

The *spoT* genes of the suppressor strains YZ543, YZ544, YZ545, and YZ546 were inactivated via

P1 phage transduction. The donor strain for making the P1 phage lysate was the *spoT207::cat* strain

(YZ47)[17], which contains a chloramphenicol resistance cassette together with *spoT207* deletion allele

for selection. The phage lysate was used to infect strains YZ543, YZ544, YZ545, and YZ546 and the

chloramphenicol resistant transductants were selected and confirmed via diagnostic PCR.

Plasmids constructions

To construct *E. coli (Ec), Salmonella typhimurium(st) or Shigella flexneri(sf) spoT* wild type (wt) and

H414A mutant plasmids, standard molecular cloning procedures were used. For *E. coli*, the *spoT*

coding sequence was PCR-amplified from MG1655 *spoT^{wt}* (YZ38) or *spoT^{H414A}* (YZ527) using

primers pYZ1201 and pYZ1202. The pBAD33 vector was isolated from *E. coli* DH5α (YZ593). Both

the amplified *spoT* fragments and pBAD33 were digested with *KpnI* and *HindIII* and ligated using

T4 DNA ligase to generate pBAD33- *spoT^{wt}* and pBAD33- *spoT^{H414A}*. For *S. Typhimurium*, the

H414A mutation was introduced into pGEN-MCS-*spoT_{St}* (YZ692)[45] by site-directed mutagenesis

using primers pYZ1312 and pYZ1313, yielding pGEN-MCS-spoT$_{st}$ (H414A). For *S. flexneri*, an

A614G substitution was introduced by overlap-extension PCR using pBAD33-*spoT_{Ec}*(wt) or

pBAD33- *spoT_{Ec}*(H414A) as templates. Upstream and downstream fragments were amplified with

primer pairs pYZ1201/pYZ1346 and pYZ1202/pYZ1345, respectively, fused by overlap PCR using

primers pYZ1201 and pYZ1202, and cloned into pBAD33 following digestion with *KpnI* and *HindIII*

to generate pBAD33-*spoT*$_{Sf}$(WT) and pBAD33-*spoT*$_{Sf}$(H414A). To construct the plasmid pCA24N-

argA(H15Y), quick-change mutagenesis was carried out using pCA24N-argA(wt) as template and

the primers pYZ1168 and pYZ1169. All plasmids were transformed into *E. coli* DH5α (see **Table**

S1), purified using a commercial miniprep kit (NEB-T1110L), and confirmed by sequencing with

primers pYZ404, pYZ405, and pYZ71 (for pBAD33 constructs), pYZ1305, pYZ1306, and pYZ1228

(for pGEN-MCS constructs) or pYZ34 and pYZ35 (for pCA24N constructs; Eurofins Genomics).

Acid resistance assay

Overnight (16 hrs) cultures were grown in LB medium at 37°C with shaking (160 rpm). Cells were

harvested by centrifugation, washed twice with PBS, and normalized to an optical density at 600 nm

(OD_{600nm}) of 0.4 in minimal medium supplemented with 0.2% glucose (M9Glc, pH 7.4). Cultures

were incubated at 37°C with shaking (160 rpm) for 30 min to allow adaptation to M9Glc. After

adaptation, 1 mL cells were pelleted and resuspended in the same volume of M9Glc medium pre-

adjusted to various pH values (from 2.5 to 8.0), with or without supplemented 0.6 mM glutamate (or

other amino acids). Acid challenge was then performed and assessed at 37°C with shaking (160 rpm).

Cell viability was determined at time zero (i.e., before shifting to M9Glc with a low pH), and at 0.5,

1, 2 and 4 h after acid challenges. For viable-cell counts, 40 μL cell culture at each time point was

collected and 10-fold serially diluted (10^0–10^{-7}) in LB medium, and 5 μL of each dilution was spotted

on LB agar plates, allowing colony formation and enumeration and calculation of CFUs the next day.

Western blot

Western blot analysis of protein levels of various mutant SpoT was performed as reported[46]. Briefly,

cells of various strains were collected the same manner as preparation for RNA sequencing (see above,

and **Fig. S1C**). These cells were normalised to have an equal OD_{600nm} per ml and mixed with 2x SDS-

PAGE loading buffer. Samples were heated for 15 min at 95°C and 20 ul of each were separated in

10% SDS-PAGE gels. The proteins were transferred to a PVDF membrane (Amersham) and the SpoT

proteins detected by first incubation with anti-SpoT antiserum (Covalab) and then the secondary HRP

conjugated rabbit IgG antibody (SIGMA). The SpoT bands were visualized by using the Pierce ECL

chemiluminescence substrate (Thermo) and the signals were detected by an Imagequant LAS4100.

Bioinformatic analysis

<u>Preparing RNA-seq data.</u> RNA-seq data (fastq files) were mapped by the BWA-MEM tool of

Galaxy [47] and then analysed by the featureCounts tool in Galaxy https://usegalaxy.eu/). The tabular

format of the featureCounts files were exported from Galaxy and assembled into one single tabular

file from which sub-files containing collections of genes of interest, e.g. amino acid biosynthesis,

purine de novo synthesis, were extracted.

<u>Analysis of RNA-seq data.</u> To generate heatmaps and fold-of-changes, the tabular files containing

the RNA-seq counts were analysed by DEGUST (https://degust.erc.monash.edu/). To generate

Venn diagrams, gene lists were analysed by the jVenn application at

(https://jvenn.toulouse.inrae.fr/app/index.html). Gene Ontology terms were generated by AmiGO

(https://amigo.geneontology.org/amigo) and Panther

(https://pantherdb.org/webservices/go/overrep.jsp). The json-files generated by PANTHER were

analysed by ChatGPT that generated editable SVG files.

<u>Other Bioinformatic Procedures</u>

Protein sequence alignment was accomplished using JalView[48], protein tertiary structure modelling

and visualization by AlphaFold 3 and ChimeraX, respectively[49,50].

QUANTIFICATION AND STATISTICAL ANALYSIS

Signals of (p)ppGpp from autoradiography were quantified using ImageQuant (GE Healthcare). At

least three biological replicates were performed and one representative shown in the **Figure**s.

For other physiological growth experiments, at least two biological replicates each with three

technical replicates were done (see **Figure** legends for details). The length (hours) of lag phases was

defined as the period from the growth start to the time point where the slope of growth curves first

reaches the exponential-phase growth rate. These lag phage length values were then normalized to

the parental strains or conditions, and plotted (see each of the legends for details). The student t-test

was used for statistic analysis wherever applicable.

DATA AVAILABILITY

The RNA-seq raw data produced in this study are available through the NCBI Sequence Read

Archive under BioProject accession number PRJNA1097790.

773   References

1.   Fang, F.C., Frawley, E.R., Tapscott, T., and Vázquez-Torres, A. (2016). Bacterial Stress
Responses during Host Infection. Cell Host Microbe *20*, 133-143.
10.1016/j.chom.2016.07.009.
2.   Lin, J., Smith, M.P., Chapin, K.C., Baik, H.S., Bennett, G.N., and Foster, J.W. (1996).
Mechanisms of acid resistance in enterohemorrhagic Escherichia coli. Appl Environ
Microbiol *62*, 3094-3100. 10.1128/aem.62.9.3094-3100.1996.
3.   Kanjee, U., and Houry, W.A. (2013). Mechanisms of acid resistance in Escherichia coli. Annu
Rev Microbiol *67*, 65-81. 10.1146/annurev-micro-092412-155708.
4.   Lund, P., Tramonti, A., and De Biase, D. (2014). Coping with low pH: molecular strategies
in neutralophilic bacteria. FEMS Microbiol Rev *38*, 1091-1125. 10.1111/1574-6976.12076.
5.   Capitani, G., De Biase, D., Aurizi, C., Gut, H., Bossa, F., and Grütter, M.G. (2003). Crystal
structure and functional analysis of Escherichia coli glutamate decarboxylase. Embo j *22*,
4027-4037. 10.1093/emboj/cdg403.
6.   Castanie-Cornet, M.P., Penfound, T.A., Smith, D., Elliott, J.F., and Foster, J.W. (1999).
Control of acid resistance in Escherichia coli. J Bacteriol *181*, 3525-3535.
10.1128/jb.181.11.3525-3535.1999.
7.   Irving, S.E., Choudhury, N.R., and Corrigan, R.M. (2021). The stringent response and
physiological roles of (pp)pGpp in bacteria. Nat Rev Microbiol *19*, 256-271. 10.1038/s41579-
020-00470-y.
8.   Anderson, B.W., Fung, D.K., and Wang, J.D. (2021). Regulatory Themes and Variations by
the Stress-Signaling Nucleotide Alarmones (p)ppGpp in Bacteria. Annu Rev Genet *55*, 115-
133. 10.1146/annurev-genet-021821-025827.
9.   Bange, G., Brodersen, D.E., Liuzzi, A., and Steinchen, W. (2021). Two P or Not Two P:
Understanding Regulation by the Bacterial Second Messengers (p)ppGpp. Annu Rev
Microbiol *75*, 383-406. 10.1146/annurev-micro-042621-122343.
10.   Hauryliuk, V., Atkinson, G.C., Murakami, K.S., Tenson, T., and Gerdes, K. (2015). Recent
functional insights into the role of (p)ppGpp in bacterial physiology. Nat Rev Microbiol *13*,
298-309. 10.1038/nrmicro3448.
11.   Block, R., and Haseltine, A.W. (1975). Purification and properties of stringent factor. J Biol
Chem *250*, 1212-1217.
12.   Brown, A., Fernández, I.S., Gordiyenko, Y., and Ramakrishnan, V. (2016). Ribosome-
dependent activation of stringent control. Nature *534*, 277-280. 10.1038/nature17675.
13.   Loveland, A.B., Bah, E., Madireddy, R., Zhang, Y., Brilot, A.F., Grigorieff, N., and
Korostelev, A.A. (2016). Ribosome•RelA structures reveal the mechanism of stringent
response activation. Elife *5*. 10.7554/eLife.17029.
14.   Arenz, S., Abdelshahid, M., Sohmen, D., Payoe, R., Starosta, A.L., Berninghausen, O.,
Hauryliuk, V., Beckmann, R., and Wilson, D.N. (2016). The stringent factor RelA adopts an
open conformation on the ribosome to stimulate ppGpp synthesis. Nucleic Acids Res *44*,
6471-6481. 10.1093/nar/gkw470.
15.   Winther, K.S., Roghanian, M., and Gerdes, K. (2018). Activation of the Stringent Response
by Loading of RelA-tRNA Complexes at the Ribosomal A-Site. Mol Cell *70*, 95-105.e104.
10.1016/j.molcel.2018.02.033.

16. Ross, W., Vrentas, C.E., Sanchez-Vazquez, P., Gaal, T., and Gourse, R.L. (2013). The magic spot: a ppGpp binding site on E. coli RNA polymerase responsible for regulation of transcription initiation. Mol Cell *50*, 420-429. 10.1016/j.molcel.2013.03.021.

17. Ross, W., Sanchez-Vazquez, P., Chen, A.Y., Lee, J.H., Burgos, H.L., and Gourse, R.L. (2016). ppGpp Binding to a Site at the RNAP-DksA Interface Accounts for Its Dramatic Effects on Transcription Initiation during the Stringent Response. Mol Cell *62*, 811-823. 10.1016/j.molcel.2016.04.029.

18. Zhang, Y.E., Bærentsen, R.L., Fuhrer, T., Sauer, U., Gerdes, K., and Brodersen, D.E. (2019). (p)ppGpp Regulates a Bacterial Nucleosidase by an Allosteric Two-Domain Switch. Mol Cell *74*, 1239-1249.e1234. 10.1016/j.molcel.2019.03.035.

19. Zhang, Y., Zborníková, E., Rejman, D., and Gerdes, K. (2018). Novel (p)ppGpp Binding and Metabolizing Proteins of Escherichia coli. mBio *9*. 10.1128/mBio.02188-17.

20. Wang, B., Grant, R.A., and Laub, M.T. (2020). ppGpp Coordinates Nucleotide and Amino-Acid Synthesis in E. coli During Starvation. Mol Cell *80*, 29-42.e10. 10.1016/j.molcel.2020.08.005.

21. Wang, B., Dai, P., Ding, D., Del Rosario, A., Grant, R.A., Pentelute, B.L., and Laub, M.T. (2019). Affinity-based capture and identification of protein effectors of the growth regulator ppGpp. Nat Chem Biol *15*, 141-150. 10.1038/s41589-018-0183-4.

22. Sanchez-Vazquez, P., Dewey, C.N., Kitten, N., Ross, W., and Gourse, R.L. (2019). Genome-wide effects on Escherichia coli transcription from ppGpp binding to its two sites on RNA polymerase. Proc Natl Acad Sci U S A *116*, 8310-8319. 10.1073/pnas.1819682116.

23. Traxler, M.F., Summers, S.M., Nguyen, H.T., Zacharia, V.M., Hightower, G.A., Smith, J.T., and Conway, T. (2008). The global, ppGpp-mediated stringent response to amino acid starvation in Escherichia coli. Mol Microbiol *68*, 1128-1148. 10.1111/j.1365-2958.2008.06229.x.

24. Durfee, T., Hansen, A.M., Zhi, H., Blattner, F.R., and Jin, D.J. (2008). Transcription profiling of the stringent response in Escherichia coli. J Bacteriol *190*, 1084-1096. 10.1128/jb.01092-07.

25. Grucela, P.K., Fuhrer, T., Sauer, U., Chao, Y., and Zhang, Y.E. (2023). Ribose 5-phosphate: the key metabolite bridging the metabolisms of nucleotides and amino acids during stringent response in Escherichia coli? Microb Cell *10*, 141-144. 10.15698/mic2023.07.799.

26. Haseltine, W.A., and Block, R. (1973). Synthesis of guanosine tetra- and pentaphosphate requires the presence of a codon-specific, uncharged transfer ribonucleic acid in the acceptor site of ribosomes. Proc Natl Acad Sci U S A *70*, 1564-1568. 10.1073/pnas.70.5.1564.

27. Xiao, H., Kalman, M., Ikehara, K., Zemel, S., Glaser, G., and Cashel, M. (1991). Residual guanosine 3',5'-bispyrophosphate synthetic activity of relA null mutants can be eliminated by spoT null mutations. J Biol Chem *266*, 5980-5990.

28. Fernández-Coll, L., and Cashel, M. (2020). Possible Roles for Basal Levels of (p)ppGpp: Growth Efficiency Vs. Surviving Stress. Front Microbiol *11*, 592718. 10.3389/fmicb.2020.592718.

29. Seyfzadeh, M., Keener, J., and Nomura, M. (1993). spoT-dependent accumulation of guanosine tetraphosphate in response to fatty acid starvation in Escherichia coli. Proc Natl Acad Sci U S A *90*, 11004-11008. 10.1073/pnas.90.23.11004.

30. Battesti, A., and Bouveret, E. (2006). Acyl carrier protein/SpoT interaction, the switch linking SpoT-dependent stress response to fatty acid metabolism. Mol Microbiol *62*, 1048-1063. 10.1111/j.1365-2958.2006.05442.x.

31.   Vinella, D., Albrecht, C., Cashel, M., and D'Ari, R. (2005). Iron limitation induces SpoT-
dependent accumulation of ppGpp in Escherichia coli. Mol Microbiol *56*, 958-970.
10.1111/j.1365-2958.2005.04601.x.
32.   Spira, B., Silberstein, N., and Yagil, E. (1995). Guanosine 3',5'-bispyrophosphate (ppGpp)
synthesis in cells of Escherichia coli starved for Pi. J Bacteriol *177*, 4053-4058.
10.1128/jb.177.14.4053-4058.1995.
33.   Dalebroux, Z.D., Svensson, S.L., Gaynor, E.C., and Swanson, M.S. (2010). ppGpp conjures
bacterial virulence. Microbiol Mol Biol Rev *74*, 171-199. 10.1128/mmbr.00046-09.
34.   Levin-Reisman, I., Ronin, I., Gefen, O., Braniss, I., Shoresh, N., and Balaban, N.Q. (2017).
Antibiotic tolerance facilitates the evolution of resistance. Science *355*, 826-830.
10.1126/science.aaj2191.
35.   Windels, E.M., Michiels, J.E., Fauvart, M., Wenseleers, T., Van den Bergh, B., and Michiels,
J. (2019). Bacterial persistence promotes the evolution of antibiotic resistance by increasing
survival and mutation rates. Isme j *13*, 1239-1251. 10.1038/s41396-019-0344-9.
36.   Lewis, K. (2007). Persister cells, dormancy and infectious disease. Nat Rev Microbiol *5*, 48-
56. 10.1038/nrmicro1557.
37.   Harms, A., Maisonneuve, E., and Gerdes, K. (2016). Mechanisms of bacterial persistence
during stress and antibiotic exposure. Science *354*. 10.1126/science.aaf4268.
38.   Rajagopal, B.S., DePonte, J., 3rd, Tuchman, M., and Malamy, M.H. (1998). Use of inducible
feedback-resistant N-acetylglutamate synthetase (argA) genes for enhanced arginine
biosynthesis by genetically engineered Escherichia coli K-12 strains. Appl Environ Microbiol
64, 1805-1811. 10.1128/aem.64.5.1805-1811.1998.
39.   Zhou, Y.N., and Jin, D.J. (1998). The rpoB mutants destabilizing initiation complexes at
stringently controlled promoters behave like "stringent" RNA polymerases in Escherichia coli.
Proc Natl Acad Sci U S A *95*, 2908-2913. 10.1073/pnas.95.6.2908.
40.   Zhang, Y.E., Baerentsen, R.L., Fuhrer, T., Sauer, U., Gerdes, K., and Brodersen, D.E. (2019).
(p)ppGpp Regulates a Bacterial Nucleosidase by an Allosteric Two-Domain Switch. Mol Cell
74, 1239-1249 e1234. 10.1016/j.molcel.2019.03.035.
41.   Kitagawa, M., Ara, T., Arifuzzaman, M., Ioka-Nakamichi, T., Inamoto, E., Toyonaga, H., and
Mori, H. (2005). Complete set of ORF clones of Escherichia coli ASKA library (a complete
set of E. coli K-12 ORF archive): unique resources for biological research. DNA Res *12*, 291-
299. 10.1093/dnares/dsi012.
42.   Saka, K., Tadenuma, M., Nakade, S., Tanaka, N., Sugawara, H., Nishikawa, K., Ichiyoshi, N.,
Kitagawa, M., Mori, H., Ogasawara, N., and Nishimura, A. (2005). A complete set of
Escherichia coli open reading frames in mobile plasmids facilitating genetic studies. DNA
Res *12*, 63-68. 10.1093/dnares/12.1.63.
43.   Neidhardt, F.C., Bloch, P.L., and Smith, D.F. (1974). Culture medium for enterobacteria. J
Bacteriol *119*, 736-747. 10.1128/jb.119.3.736-747.1974.
44.   Kanjee, U., Gutsche, I., Alexopoulos, E., Zhao, B., El Bakkouri, M., Thibault, G., Liu, K.,
Ramachandran, S., Snider, J., Pai, E.F., and Houry, W.A. (2011). Linkage between the
bacterial acid stress and stringent responses: the structure of the inducible lysine
decarboxylase. Embo j *30*, 931-944. 10.1038/emboj.2011.5.
45.   Chau, N.Y.E., Perez-Morales, D., Elhenawy, W., Bustamante, V.H., Zhang, Y.E., and
Coombes, B.K. (2021). (p)ppGpp-Dependent Regulation of the Nucleotide Hydrolase PpnN
Confers Complement Resistance in Salmonella enterica Serovar Typhimurium. Infect Immun
89. 10.1128/IAI.00639-20.

46.   Towbin, H., Staehelin, T., and Gordon, J. (1979). Electrophoretic transfer of proteins from
910   polyacrylamide gels to nitrocellulose sheets: procedure and some applications. Proc Natl
911   Acad Sci U S A *76*, 4350-4354. 10.1073/pnas.76.9.4350.
912   47.   The Galaxy platform for accessible, reproducible, and collaborative data analyses: 2024
913   update. (2024). Nucleic Acids Res *52*, W83-w94. 10.1093/nar/gkae410.
914   48.   Procter, J.B., Carstairs, G.M., Soares, B., Mourão, K., Ofoegbu, T.C., Barton, D., Lui, L.,
915   Menard, A., Sherstnev, N., Roldan-Martinez, D., et al. (2021). Alignment of Biological
916   Sequences with Jalview. Methods Mol Biol *2231*, 203-224. 10.1007/978-1-0716-1036-7_13.
917   49.   Senior, A.W., Evans, R., Jumper, J., Kirkpatrick, J., Sifre, L., Green, T., Qin, C., Žídek, A.,
918   Nelson, A.W.R., Bridgland, A., et al. (2020). Improved protein structure prediction using
919   potentials from deep learning. Nature *577*, 706-710. 10.1038/s41586-019-1923-7.
920   50.   Pettersen, E.F., Goddard, T.D., Huang, C.C., Meng, E.C., Couch, G.S., Croll, T.I., Morris,
921   J.H., and Ferrin, T.E. (2021). UCSF ChimeraX: Structure visualization for researchers,
922   educators, and developers. Protein Sci *30*, 70-82. 10.1002/pro.3943.
923

Legends to Supplementary Tables

Table S1. Bacterial strains used in this study

Table S2. Oligonucleotides used in this study

Table S3. Differential gene expression analysis of RNA-seq data from *spoT^wt* and *spoT^H414A*

strains . In total, the transcription levels of 4575 genes of MG1655 were analysed. At a setting of

$|\text{Log2(FC)}| > 1$ and FDR < 0.05, 865 and 949 genes were down and up-regulated, respectively. The

change-of-folds $|\text{Log2(FC)}|$ of each gene are also shown.

Table S4. Gene ontology analysis of the four suppressor strains.

Sheet #1 (related to Extended Data Fig. 7C): Genes downregulated in the *spoT^H414A* strain relative

to the *spoT^wt* strain (Column A). Genes upregulated in either of the four suppressor strains *supA*,

sup41, *supB* and *sup17* relative to the *spoT^H414A* strain (Columns B to E). The list of genes in

Columns B to E were used to generate a list of genes that were upregulated in at least one

suppressor strain (Column F) and a list of genes that were upregulated in all 4 suppressor strains

(Column G).

Sheet #2 (related to Extended Data Fig. 7D): Genes upregulated in the *spoT^H414A* strain relative

to the *spoT^wt* strain (Column A). Genes downregulated in either of the four suppressor strains

relative to the *spoT^H414A* strain. The list of genes in Columns B to E were used to generate a list of

genes that were upregulated in at least one suppressor strain (Column F) and a list of genes that

were downregulated in all 4 suppressor strains (Column G). Gene lists in **Table S4** were generated

by DEGUST at a setting of $|\text{Log2(FC)}| > 1$ and FRD < 0.05.

Table S5. Comparison of *spoT^H414A* with suppressor strains (related to Extended Data Fig. 8).

949   The Table yields the folds-of-changes in the suppressor strains relative to *spoT^{H414A}*. The Log2(FC)

950   setting in DEGUST was chosen to yield around 100 genes up and downregulated in each heatmap

951   Panels a, b, e, h and k of **Extended Data Fig. 8**.