Title

Mechanically competitive regulation of cell volume in cytoplasm-sharing cells connected by

intercellular bridges

Short title

Mechanics of cell volume in cytoplasm-sharing germ cells

Authors

Hiroshi Koyama[1,2,*], Kanako Ikami[3,4], Lei Lei[5,6], and Toshihiko Fujimori[1,2]

[1]Division of Embryology, National Institute for Basic Biology, 5-1 Higashiyama, Myodaiji, Okazaki,

Aichi 444-8787, Japan

[2]SOKENDAI (The Graduate University for Advanced Studies), 5-1 Higashiyama, Myodaiji, Okazaki,

Aichi 444-8787, Japan

[3]Howard Hughes Medical Institute, University of California Davis, Davis, California, CA 95616

[4]Department of Microbiology & Molecular Genetics, University of California Davis, Davis, CA 95616

[5]Department of Obstetrics, Gynecology and Women's Health, University of Missouri School of

Medicine, Columbia, MO 65211

[6]Division of Biological Sciences, College of Arts and Sciences, University of Missouri, Columbia,

MO 65211

*Correspondence: Hiroshi Koyama

Division of Embryology, National Institute for Basic Biology, 5-1, Higashiyama, Myodaiji, Okazaki,

Aichi 444-8787, Japan

Phone: +81 564 59 5862,

hkoyama@nibb.ac.jp.

Abstract

In multicellular organisms, various cellular structures exhibit cytoplasmic sharing, where cells remain interconnected. While essential for development and function in contexts such as germ cell formation and insect early embryos, the physical basis of cell volume regulation in these systems remains poorly defined. Germline cysts are formed by interconnected sister cells via intercellular bridges. In mice, germline cysts form during gametogenesis in fetal ovaries and testes. In mouse fetal female cysts, cells with numerous bridges preferentially differentiate into oocytes by selectively increasing their volume, a process that may be mediated through cytoplasmic flow. This volume bias may be influenced by hydrostatic pressure within the cytoplasm. Here, we theoretically investigate how the mechanical properties of cells affect cytoplasmic pressure and volume distribution within interconnected cells. Our soap-bubble model revealed that cells with more bridges exhibit increased volume when they have large cell-cell contact areas, as observed in fetal cysts. We found that incorporating cell cycle (including cell growth and cell division) significantly enhances the likelihood of volume bias in favor of cells with more bridges. These theoretical findings suggest that intrinsic mechanical properties, coupled with cell cycle, establish robust cyst development in fetal female germline cysts. Our findings also provide insights into the volume dynamics observed in adult male germline cysts, which are characterized by smaller cell-cell contact areas.

Impact statement

A theoretical model demonstrates how mechanical properties and cell cycle dynamics regulate volume distribution in germline cysts, providing a physical basis for oocyte differentiation.

Key words

mouse germline cyst, Laplace pressure, cell-cell contact, cytoplasmic flow, mathematical modeling

Introduction

In multicellular organisms, cytoplasm in each cell is typically separated by cell membranes. On the other hand, there are some cell types whose cytoplasm is shared among multiple cells. Notable

examples include the early syncytial embryos in insects like *Drosophila* during cellularization where
the nuclei are partially separated by cell membrane (Lv et al., 2021); the formation of oocytes from
the gonad in nematodes where each germ cell is connected to a central cytoplasmic core called the
rachis through large intercellular bridges (Kimura and Motegi, 2025; Priti et al., 2018); cytoplasmic
connectivity between blastomeres in the *Xenopus* embryo (Landesman et al., 2000); the
plasmodesmata in plants which are microscopic cytoplasmic channels that traverse the cell walls of
plant cells (Bayer and Benitez-Alfonso, 2024); and the germline cysts found across a wide range of
species from *Drosophila* to mammals where each germ cell is connected via interconnected bridges
(Chaigne and Brunet, 2022). In these contexts, precise regulation of individual cell volume within the
shared cytoplasm is important for development, differentiation, and physiological functions.

Our present study focuses on mouse fetal germline cysts, where sister cells derived from a
common progenitor remain interconnected through intercellular bridges formed by incomplete
cytokinesis (Fig. 1A-B) (Chaigne and Brunet, 2022). A key aspect of germline cyst development,
particularly in females, is the highly regulated process of oocyte differentiation, which often involves
a dramatic increase in the volume of a few cells. During this process in mice, both cyst structure and
fragmentation of germline cysts play critical roles (Ikami et al., 2023). The mouse female germline
cysts are not simple linear chains, but rather complex branched structures (Fig. 1A), which are then
partially pruned during fragmentation. Future oocytes tend to be connected by more intercellular
bridges and become physically larger due to cytoplasmic transfer from neighboring cells, even before
cyst fragmentation. Therefore, there is a competitive regulation of cell volume. Active and directional
cytoplasmic transport via microtubules with motor proteins was found in *Drosophila* female cysts
(Clark et al., 2007; Gutzeit, 1986; Lu et al., 2022). Although such transport was investigated in mice
(Lei and Spradling, 2016), it has not been fully characterized.

Recently, mechanical forces have been reported to play critical roles in various
morphogenetic events and differentiation (Heisenberg and Bellaïche, 2013; Koyama et al., 2023;
Kurotaki et al., 2007; Sagner et al., 2012; Savin et al., 2011). Cell volume and shape are also regulated
by mechanical forces (Cuvelier et al., 2023; Devany et al., 2023; Farhadifar et al., 2007; Koyama et
al., 2012; Miyoshi et al., 2006; Okuda et al., 2015; Runser et al., 2024; Xie et al., 2024). Cytoplasm is
flowed by its hydrostatic pressure which is, at least, partially provided from cell surface tensions
(Fujita and Onami, 2012; Sedzinski et al., 2011). While active and directional cytoplasmic transport
may contribute to oocyte growth, it is plausible that such transport would increase hydrostatic pressure
in the larger cells, potentially creating a counteracting force that could offset or reverse cytoplasmic
flow. Therefore, a precise control of cytoplasmic pressure may be involved in robust cyst development

and the stable establishment of cell fate. In the case of *C. elegans* germ cells connected to the rachis

through intercellular bridges as mentioned above, cell surface tensions generate differences in inner

pressures, leading to biased cell volume (Chartier et al., 2021). In contrast to the *C. elegans* germ cells,

mouse female germ cells exhibit a distinct structure: they lack a rachis, and crucially, the number of

intercellular bridges varies among individual cells within the cyst. This complexity necessitates

studying how mechanical factors regulate volume under conditions where connectivity is

heterogeneous.

Here, we theoretically investigated the contribution of purely mechanical properties of cells to

cytoplasmic pressure regulation and subsequent volume distribution within mouse female germline

cysts. We developed a simplified model where each cell is analogous to a soap bubble, allowing us to

explore the relationship between cell surface tension, surface curvature, and internal cytoplasmic

pressure, as described by the Laplace pressure law. Our model also assumes factors, including the

number of bridges, cell-cell contact area, cell cycle, and cyst fragmentation as described later. By

analyzing this model, we elucidated possible contributions of these cell-intrinsic properties to the

biased cell volume distribution in the female germline cyst.

1. Theory

1-1. Definition of mechanical properties of cells

We developed a theoretical model to investigate the mechanical properties governing cell volume

regulation within germline cysts. Figure 1A illustrates the structure of the female fetal germline cysts

in mice. During this stage, cells exhibit robust adhesion via E-cadherin proteins at cell-cell contacts,

where intercellular bridges enable cytoplasmic sharing (Fig. 1A) (Ikami et al., 2023).

To model the mechanical behavior of these cells, we drew an analogy to a soap bubble, considering

cell surface tensions and internal hydrostatic pressures. This simplified model, often employed in cell

mechanics due to its experimental basis and inherent simplicity, relates internal hydrostatic pressure

(P), surface tension (λ), and surface curvature (H) via the Laplace pressure equation ($P = 2\lambda H$) (Fig.

1B) (Sedzinski et al., 2011; Wang et al., 2021). Subscripts for each parameter denote specific cell IDs

(e.g., P_A for A-cell, P_B for B-cells). A critical assumption of typical soap bubble-based models is that

surface tension (λ) remains constant regardless of cell size, because the surface tension may be

determined by the contractility of the cortical actomyosins rather than by elasticity (Sedzinski et al.,

2011). Consequently, as a cell's volume increases, its surface curvature (H) simultaneously decreases.

Under the constant λ assumption, this leads to a decrease in the pressure (P). It is known that in such

soap bubble-based systems (and in real soap bubbles), even a slight difference in volume between

connected cells is temporally amplified (Sedzinski et al., 2011), leading to the complete loss of volume

in one of the cells due to cytoplasmic flow driven by pressure differences (e.g., P_A vs. P_B). This

knowledge implies an instability if not properly regulated.

1-2. Assumption of constant cell-cell contact number throughout the cell cycle

On the basis of the above model, we aimed to evaluate the effect of the number of cell-cell contacts

with intercellular bridges on cell volume. In the real tissues, the number of cell-cell contacts for each

cell should be temporally changeable due to cell division and cyst fragmentation. However, to purely

extract the effect of the number of cell-cell contacts, we constructed a model where the number of cell-

cell contacts was set to be constant throughout the simulations even when cell division occurred as

follows. In Fig. 2A, B-cell has two cell-cell contacts with A_1- and A_2-cells, while A_1- and A_2-cells have

one cell-cell contact. We assumed that the A_2-cell was equivalent to the A_1-cell (i.e., the same

parameter values are shared). In vivo fetal germ cells undergo multiple rounds of cell division, with

each division theoretically reducing the cell volume by 50%. However, the overall cell volume does

not decrease during multiple rounds of the cell cycle (Ikami et al., 2023), indicating that concurrent

cell growth (i.e., nutrient uptake and synthesis (Devany et al., 2023; Phillips et al., 2008; Wu et al.,

2022)) occurs during interphase. Therefore, during a cell cycle, cell volumes increase due to cell

growth, while the cytoplasm flows through the bridges due to pressure differences (Fig. 2A, 1st panel).

Then, these cells synchronously experience cell division in a manner similar to *in vivo* germline cysts.

We assumed a fixed cell division plane so that the connection of the original three cells (B-, A_1-, and

A_2-cells) is maintained (Fig. 2A, 2nd and 3rd panels). Furthermore, although the proliferated cells are

also connected to their sister cells in *in vivo* cysts, we forcibly deleted these cells in our model so that

the number of cell-cell contacts in B-, A_1-, and A_2-cells remains constant (Fig. 2A, 3rd and 4th panels).

Under these settings, combined with the equivalence between A_1- and A_2-cells, the system is virtually

composed of just two cells with a constant number of cell-cell contacts (Fig. 2A, magenta dashed line

boxes). Note that we will also show a full model incorporating all proliferated cells later.

1-3. Assumption of cell-cell contact regions

Next, we modeled the cell-cell contact region (Fig. 1A). Due to the cell-cell contacts via E-cadherin,

the cells show spherical dome shapes (Fig. 2B). In the absence of E-cadherin junctions, the

interconnected bridges are formed, but the cell-cell contacts are not. The radius of the spherical dome

is denoted as R. The ratio of the volume in the spherical dome to that in the sphere whose radius is the

same as R denotes $(1 - \Omega)$ (Fig. 2B). When the number of cell-cell contacts (N) is larger than 1, the

ratio of the volume becomes $(1 - \Omega N)$. According to these relationships, if the cell volume (V) is given,

R can be calculated by considering Ω and N (Fig. 2 and S1). Although Ω would be complicatedly

changed in the *in vivo* tissues, we simply assumed that Ω is constant throughout a simulation to extract

fundamental behaviors in the presence of the cell-cell contacts.

1-4. Calculation of pressure and cytoplasmic flow rate

After the calculation of R, we can calculate P using the Laplace pressure equation: $P = 2\lambda H = 2\lambda/R$

(Fig. 2 and S1). The difference in P between interconnected cells generates intercellular cytoplasmic

flow. The volumetric flow rate through the bridges may be described by the Hagen–Poiseuille equation

(Phillips et al., 2008). This equation states that the volumetric flow rate in a cylindrical tube is

proportional to both the pressure difference and the fourth power of the tube radius, and inversely

proportional to both the tube length and viscosity (μ). In our model, we subsumed these geometric

factors (i.e., radius, length, and other structural properties) into a single parameter G_h ("geometry of

hole"), leading to the equation of the flow rate as shown in Fig. S1. Both G_h and μ are constant

throughout a simulation.

1-5. Overall procedure of simulation

Our simulation procedure is as follows (Fig. S1). As the initial condition, the surface tensions are given

as constant ($\lambda_A = \lambda_B = $ const.), and the initial cell volumes (V_A and V_B) in A₁- and B-cells are given.

After simulations start, the radii of the cells (R_A and R_B) are calculated according to Ω and N; the

pressures (P_A and P_B) are calculated by considering the Laplace pressure equation; the intercellular

hydraulic cytoplasmic flow rates are calculated; the rates of cell volume changes are determined by

the summation of the intercellular cytoplasmic flow rates and cell growth rate which will be defined

later; the volume changes are numerically calculated by using the Euler method. Note that the above

step involving the Laplace pressure equation was assumed to be a quasi-static process, ensuring that

the Laplace law is satisfied at each time step throughout the temporal evolution. After the time period

of one cell cycle, the cells are divided, and the proliferated cells are deleted to maintain the virtual 2-

cell situation.

In our simulations, the initial R_A was set to be 1.0 as the characteristic length scale, and the

time period of one cell cycle was set to be 1.0. Other default parameter values are $\lambda = 1.0$, $\mu = 0.2$, and

$G_h = 0.2$, unless noted otherwise.

2. Results

2-1. Virtual 2-cell system without cell growth and division

2-1-1. The virtual 2-cell system is unstable

We first confirmed the instability of the system composed of interconnected cells, as illustrated in Fig.

1B. We then performed simulations in the absence of cell division. Figure 3A presents simulation

results when the overlapping region (Ω) between the cell-cell contacts was set to zero (i.e., $\Omega = 0\%$).

The A-cell always had one bridge ($N_A=1$), while the B-cell had either one ($N_B=1$) or two bridges (N_B

$=2$) (Fig. 3A, upper or lower panel, respectively); if $N_A=1$ and $N_B=1$, these two cells are equivalent.

When the initial volume ratio of B-cell to A-cell ($V_B(t=0)/V_A(t=0)$) was exactly 1.0, the cell volumes

remained constant over time, even if $N_B > N_A$. However, when the B-cell had a slightly larger volume

at the initial condition, such as $V_B(0)/V_A(0) = 1.01$, the B-cell volume was continuously increased,

whereas the A-cell volume was decreased, eventually resulting in the complete loss of A-cell volume.

This demonstrates that $V_B(0)/V_A(0) = 1.0$ represents an unstable equilibrium state under these

conditions, as we had expected. In other words, *in vivo* germline cysts should have mechanisms for

stabilization, which prevent the complete loss of germ cells in the cyst due to a small degree of

difference in cell volume. Note that we did not include nuclei in our considerations, which cannot pass

through the intercellular bridges.

2-1-2. Overlapping region changes the condition of unstable equilibrium state

Next, we examined the influence of the overlapping region ($\Omega \neq 0\%$) on the instability (Fig. 3B),

recapitulating E-cadherin junctions between cyst germ cells. When $N_B =1$ (Fig. 3B, upper panels), the

simulation results were largely similar to the $\Omega = 0\%$ case (Fig. 3A); an initial ratio of 1.0 led to

constant volumes, while a slight deviation caused volume divergence. However, when $N_B=2$ (Fig. 3B,

lower panels), the initial condition $V_B(0)/V_A(0) = 1.0$ no longer represented an equilibrium state;

instead, it resulted in an increase in B-cell volume and a decrease in A-cell volume. Interestingly, a

new unstable equilibrium state emerged at approximately $V_B(0)/V_A(0) = \sim 0.89$ (Fig. 3B). If the initial

ratio was slightly larger than 0.89 (e.g., $V_B(0)/V_A(0) = 0.90$), B-cell volume was continuously increased.

Conversely, if $N_B = 1$ and the initial ratio was ~ 0.89, the A-cell volume was continuously increased. If

$N_B = 2$ and the initial ratio was less than 0.89, A-cell volume was continuously increased as later shown

in Fig. 3C.

We plotted these unstable equilibrium states for various values of N_B and $V_B(0)/V_A(0)$ (Fig.

3C). The left panel in Fig. 3C shows how to read a phase diagram for a certain value of Ω: the unstable

equilibrium state is shown as a line that represents the critical threshold determining which cell, A or

B, would continuously increase in volume. Conditions above the dotted line led to an increase in B-

cell volume ($V_A(t=T) < V_B(t=T)$), while conditions below the dotted line resulted in an increase in A-

cell volume ($V_A(T) > V_B(T)$), where T is the final simulation time. In the middle panel in Fig. 3C, the

threshold lines for different values of Ω are shown; the threshold lines can be analytically solved by

considering the condition satisfying $P_A = P_B$ (Fig. 3C, right panel). For larger Ω values with $N_B \geq 2$,

the unstable equilibrium states shifted to smaller values of $V_B(0)/V_A(0)$. These results indicate that the

B-cell can more easily increase its volume in the presence of numerous bridges and larger overlapping

regions (Fig. 3D), highlighting the significant role of cell-cell contact regions in modulating volume

dynamics. This can be a candidate mechanism for explaining the biased increase in cell volume in

fetal female germline cysts (Fig. 1A), although this model has not yet considered the cell cycle. In

addition, the values of the surface tension (λ), the geometric effect of the bridges (G_h), and the

coefficient of viscosity (μ) did not affect the fate of cell volume but did affect the rate of the volume

changes (Fig. S2).

2-2. Virtual 2-cell system with cell growth and division

2-2-1. Cell growth and division stabilize the system

We next incorporated the cell cycle, including both cell division and cell growth, into our virtual 2-

cell model. Typical somatic cells undergo cell growth (driven by nutrient uptake and synthesis) during

interphase (Devany et al., 2023; Wu et al., 2022), and subsequently, their volume decreases by 50%

due to cell division, as mentioned in the Introduction. We assumed two scenarios for cell growth rates

(Fig. S3; constant rate or cell cycle-dependent rate). We will show one of the two scenarios below,

because the two scenarios resulted in qualitatively similar simulation outcomes (Fig. S4). Figure 4A

shows simulation outcomes under conditions similar to Figure 3B, but with the inclusion of cell growth

and division. Note that, as previously mentioned, the cell cycle is synchronized in a cyst. Although the

number of cell cycles is limited to about five *in vivo*, we performed simulations over numerous rounds

to assess the stability and robustness of the volume distribution. In the absence of the cell-cell contact

regions (Fig. 4A; $\Omega = 0\%$), each cell showed an oscillatory behavior of its volume due to cell growth

and division. Importantly, although the initial volume ratio was set to be biased ($V_B(0)/V_A(0) = 0.89 \neq$

1.0), the volume in B-cell became equal to that in A-cell, which was contrast to Figure 3 where a slight

difference in the initial volume was continuously enhanced (e.g., Fig. 3B, $\Omega = 0\%$, $V_B(0)/V_A(0) = 1.01$).

Therefore, cell growth and division stabilized the systems, but did not lead to biased cell volume in

the absence of cell-cell contact regions.

2-2-2. Cell growth and division enhance the likelihood of volume increase in cells with numerous

bridges

Next, we evaluated the effect of the overlapping regions of cell-cell contacts in the presence of cell

growth and division. In Figure 4A, with $\Omega = 10\%$ and $N_B=2$ ($N_A=1$), an initial volume ratio of V_B

$(0)/V_A(0) = \sim0.89$ led to the B-cell volume becoming larger than the A-cell volume. Therefore, the

overlapping regions caused the biased volume increase in B-cell ($\Omega = 10$ vs. 0%). With $\Omega = 10\%$, even

if $V_B(0)/V_A(0)$ was quite small (0.46), the B-cell became larger than the A-cell. A new threshold value

of $V_B(0)/V_A(0)$ for volume increase/decrease emerged at approximately 0.32, which was much smaller

than the threshold value in Fig. 3B (0.89). If the initial ratio was smaller than this threshold (e.g., 0.30),

B-cell volume was completely lost. Under the condition of larger Ω (20%) with larger N_B (3), the

threshold was further decreased ($\sim$0.16). These results theoretically indicate that, in proliferating germ

cells, the cells harboring the numerous bridges with larger overlapping regions have a higher

likelihood of volume increase.

2-2-3. Systematic analysis of the likelihood of volume increase in cells with numerous bridges

Then, we systematically analyzed these threshold values of $V_B(0)/V_A(0)$ for volume increase/decrease.

We generated a phase diagram (Fig. 4B) in a manner similar to that in Fig. 3C. As explained in the left

panel in Fig. 4B, the threshold line was calculated; conditions below the line resulted in an increase in

A-cell volume ($V_A(T) > V_B(T)$), while conditions above the line led to an increase in the B-cell volume

($V_A(t=T) < V_B(t=T)$) or to equal volumes between B- and A-cells. We calculated the threshold line

through numerous trials of simulations (i.e., numerically solved but not analytically). The right panel

in Fig. 4B is the phase diagram for different values of Ω. For [N_B = 2 AND Ω = 0] or N_B = 1 (even

with Ω > 0), conditions above the lines resulted in the equal volumes between A- and B-cells (V_A

(T)=V_B(T)). Otherwise, conditions above the lines led to an increase in B-cell volume ($V_A(T) < V_B(T)$).

Conditions below the lines resulted in an increase in A-cell volume ($V_A(T) > V_B(T)$). The asterisk-

marked point on the green dashed line in Figure 4B corresponds to the condition in Figure 4A ($V_B(0)$/

$V_A(0)$ = ~0.32 with Ω = 0.1), and the cross-marked point adjacent to the red dashed line corresponds

to $V_B(0)$/ $V_A(0)$ = ~0.16 with Ω = 0.2. In comparison with Fig. 3C, the threshold lines were largely

shifted downward in the phase diagram, indicating that the likelihood of volume increase in the B-cell

with numerous bridges was enhanced in the presence of cell growth and division. In other words, even

if the initial volume in the B-cell is much smaller by chance, the B-cell can selectively increase its

volume through the cell cycle. This mechanism may contribute to robustly selecting cells whose

volumes should be increased.

2-2-4. Calculation of cell volume at converged state

Finally, we ran the simulations until they reached converged states, by which we obtained the final

cell volume ratio ($V_B(T)/V_A(T)$). We focused on the conditions where the B-cell volume increased (i.e.,

above the threshold lines in Figure 4B). For a certain value of Ω and N_B, the final volume ratios

remained constant regardless of the initial volume ratios ($V_B(0)/V_A(0)$), as long as they were above the

respective threshold lines. For instance, when Ω = 0.2 (20%), the final volume ratio was 1.0 for N_B =

1 or approximately 2.0 for N_B = 2 (Fig. 4C, red broken line, N_B = 1 or = 2). Notably, for N_B = 3, the

ratio became infinite (indicated by ∞), because of the complete loss of the A-cell volume, as also

suggested by the panel of Figure 4A, Ω = 0.2 (20%). As Ω was set to be larger (Fig. 4C; Ω = 0.2 vs.

0.1 (green broken line), 0.05 (light blue broken line), or 0 (purple line)), the final volume ratios became

larger. Similarly, as N_B was set to be larger, the final volume ratios became larger. These results indicate

that both the bridge number (N_B) and its overlapping degree (Ω) contribute to biased cell volume in

specific cells (Fig. 4D).

2-2-5. Summary of virtual 2-cell system simulations

As illustrated in Fig. 4D, both the number of bridges and the overlapping region are critical for biased

cell volume expansion. This interplay provides a potential mechanism for the asymmetric volume

distribution observed in female fetal germline cysts. In contrast, incorporating the cell cycle leads to

an equalized volume distribution in systems with fewer bridges and no overlapping regions—a

configuration that may reflect the characteristics of adult male germline cysts, as further detailed in

the Discussion.

2-3. Full model considering both cyst formation and cyst fragmentation

2-3-1. Definition of Full Model

To investigate cell volume regulation in a more complex and biologically realistic setting, we extended

our previous model (the virtual 2-cell model) to incorporates stochastic processes observed in mouse

germline development. *In vivo* germline cysts exhibit several characteristic features: the cell-cell

connection pattern such as branching varies among the cysts, the cell number in a cyst varies, and the

intercellular bridge number can be temporally increased due to cell division or decreased due to bridge

severing. In addition to both the Laplace pressure and cell cycle in the virtual 2-cell model, our new

model assumed both the cyst formation and cyst fragmentation (bridge severing), by which we

evaluated the effect of these features on cell volume.

As depicted in Figure 5A, which illustrates cyst formation, the cell division plane was

stochastically set (Fig. S5A), leading to various cell-cell connection patterns; i.e., the proliferated cells

were not deleted in this full model. Subsequently, the intercellular bridges are stochastically severed,

yielding multiple cysts with different cell numbers (Fig. 5A, bottom panel). The resulting cysts then

proceed through the next cell cycle, undergoing processes such as cell growth and cytoplasmic flow

as defined in our virtual 2-cell model. For the bridge severing process, we defined two models (Fig.

S5B): model-#1, biased severing according to bridge number, and model-#2, random severing.

2-3-2. Setting of probability of severing

First, we searched for the probability of bridge severing, which can recapitulate the process of cyst

fragmentation *in vivo*. In *in vivo* female germline cysts (Ikami et al., 2023), after starting from cysts

composed of 1 cell, the populations of the cysts composed of 4 and 8 cells were enriched after 2 or 3

cell cycles, and then, the number of cells in a cyst diverges from 1 cell to several tens of cells after an

addition few cell cycles. Basically, there is a decline in the population of cysts with an increasing

number of constituent cells. In our simulations, we set various values of the severing probability for

the model-#1 and -#2 (Fig. S6A), and found an optimal value for each model, which can approximate

the overall patterns of the populations observed *in vivo* (Fig. S6B). Figure 5B shows the populations

under the optimal value in the model-#1 (based on biased severing). Cysts composed of 4 and 8 cells

were enriched at the 8-cell stage (3 cell cycles). At the 32-cell stage (5 cell cycles), cysts composed of

4 and 8 cells were still enriched, while cysts with several tens of cells showed smaller populations.

The model-#2 (based on random severing) was also able to yield the similar population patterns (Fig.

S6A-B). The random severing during cyst formation is sufficient to recapitulate the *in vivo*

phenomenon during cyst formation. Following cyst formation, as we previously reported, cell division

stops and cyst fragmentation continues, with biased severing occurring throughout this process (Ikami

et al., 2023).

We next checked the bridge number in a cell. In *in vivo* female germline cysts (Ikami et al.,

2023), the populations of cells with 1 and 2 bridges were enriched, and the populations of cells with 0

or >=3 bridges exhibited lower. Under the condition of the severing probability in Fig. 5B, our

simulations enriched the populations of cells with 1 and 2 bridges (Fig. 5C and S6C). Together with

Fig. 5B, our model successfully approximated the structures of the cysts during cyst formation from

the viewpoint of both the bridge numbers in a cell and the cell number in a cyst. Note that, as we will

demonstrate in the following section, the probability of bridge severing itself does not significantly

affect cell volume during cyst formation, suggesting that the precise population distribution shown

here is not central to our core findings on volume regulation.

2-3-3. Effect of overlapping region and bridge number on cell volume in the full model

We next used the full model to investigate whether both the overlapping regions of cell-cell contacts

and the bridge number influence individual cell volumes within the cysts. Figure 6A-i shows the

average cell volumes for the cells harboring different numbers of bridges. In the absence of the

overlapping region (left upper panel; $\Omega = 0.0$), the cell volumes were not significantly different among

the cells with different numbers of bridges (N_B). On the other hand, in the presence of the overlapping

regions (right upper panel; $\Omega = 10.0\%$), the cell volumes became larger as the bridge numbers were

increased. These trends were also observed under the condition of the bridge severing probability =

0.0 ($P_0 = 0.0$, lower two panels; i.e., no severing occurs). Similar results were also obtained in the

model-#2 (Fig. S7). In Fig. 6A-ii, we quantified the effect of Ω on cell volume. Both Ω and N_B

positively contributed to the increase in cell volume. Fig. 6A-iii shows the effect of P_0, where P_0 did

not significantly affect the cell volume. In conclusion, the positive effects of both the overlapping

region and the bridge number were detected in both the full model (Fig. 6) and the virtual 2-cell

(Fig. 4). In addition, the bridge severing probability had no effect on cell volume.

In the above simulations, the parameter values are identical to those in the virtual 2-cell

model, except for the cell growth rate, which was adjusted to almost maintain the total volume in a

cyst across the multiple cell cycles (as described in Fig. S3) and $\mu = 0.15$.

2-3-4. Branching is effective to cell volume bias

Then, to assess the impact of branching, we modified the full model so that no branches were formed;

strictly linear cyst structures were formed solely (Fig. 6B-i). In such linear cysts, the maximum bridge

number in a cell is inherently limited to two, as indicated by our simulations (Fig. 6B-i, right panel).

Figure 6B-ii shows the cell volumes in these unbranched cysts. In the presence of the overlapping

region ($\Omega = 10.0\%$), even in the cells with the maximum possible bridge number ($N_B = 2$), the cell

volumes were just slightly increased (approximately ~ 4.03) compared to the cells with 1 bridge. This

contrasts with the results from the full model with branching (e.g., Figure 6A-ii, where the cells with

5 bridges reached a volume of approximately ~ 4.6 when $\Omega = 10.0\%$). This comparison strongly

suggests that the ability to form the branched structures, which allows the cells to acquire a larger

number of intercellular bridges, is crucial for establishing pronounced volume biases.

3. Impact of cell-cell contacts without bridges on cell volume

In the *in vivo* germline cysts, the cells aggregate regardless of the presence of intercellular bridges,

forming tightly packed clusters. (Ikami et al., 2023). At the cell-cell contacts without the bridges,

cytoplasm is not directly shared between the two cells. Here, we investigated the effect of such cell-

cell contacts on cell volume regulation. Figure 7A illustrates the modified virtual 2-cell model used

for this analysis. Similar to the previous simulations in Fig. 4, A-cell and B-cell were initially assumed

to have 1 and 2 cell-cell contacts with bridges, respectively (Fig. 7A, "Default"). We then introduced

additional cell-cell contacts to the B-cell, either with or without the bridges (Fig. 7A, "Contact number

(N_{new}) = 0, 1, 2…"). For example, the B-cells in the rightmost panels show scenarios where B-cells

have a total of 4 cell-cell contacts: either all 4 contacts harbor a bridge (upper panel), or 2 contacts

harbor a bridge and the remaining 2 contacts do not (lower panel). This setup allowed us to directly

compare the impact of bridge-containing versus bridgeless contacts.

Figure 7B presents the threshold values of the initial volume ratio ($V_B(0)/V_A(0)$) that

determine the direction of volume change, similar to Figure 4B, but now incorporating cell-cell

contacts without bridges. With $\Omega > 0$, conditions above the lines resulted in B-cell volume increasing

(i.e., the final volume ratio will be $V_A(T) < V_B(T)$), while conditions below the lines led to A-cell

volume increasing ($V_A(T) > V_B(T)$). With $\Omega = 0$, conditions above the lines resulted in $V_A(T) = V_B(T)$,

whereas below the lines, A-cell volume increased. In comparison between $\Omega = 0$ and $=5.0\%$, the

presence of the cell-cell contacts, even without the bridges, still influenced the initial conditions that

determine the eventual winner of the volume competition (Fig. 7B, yellow vs. red broken line),

although the specific thresholds differ from the cases with only the bridge-containing contacts.

Figure 7C shows the final converged volume ratios ($V_B(T)/V_A(T)$) under the conditions

where the B-cell volume increased, similar to Figure 4C. We observed that the final volume ratios

increased as the number of the cell-cell contacts without bridges increased, although the rate of this

increase was lower compared to the situations where the additional contacts contained bridges. This

finding raises the possibility that the aggregation of the cells in germline cysts contributes to a biased

volume increase via cell-cell contacts.

Discussion

In the present study, we theoretically investigated the mechanism of the competitive regulation of cell

volume in mouse female germline cysts. On the basis of cellular mechanics including the cell surface

tension and hydrostatic pressure, we constructed mathematical models where the cells shared their

cytoplasm through cell-cell contacts with intercellular bridges. In our models, the cells with a larger

number of bridges selectively increased their cell volumes, when the cells are tightly contacted with

each other (i.e., larger value of overlapping region Ω). Furthermore, cell cycle (i.e., cell growth and

cell division) contributes to biased volume increase in the cells with more bridges. This bridge number-

dependent mechanism accompanying tight cell-cell contacts can be a candidate to explain the biased

cell volume observed in the female germline cysts, leading to oocyte differentiation. Our present study

theoretically proposes mechanical bases for competitive volume regulation and for robust maintenance

of volume in the female germline cysts.

In addition to the above critical factors (i.e., overlapping region, bridge number, and cell

cycle (cell growth and cell division)), we identified several factors contributing to the biased cell

volume: branching, cell surface tension, cross-section area of the bridge (i.e., diameter of the bridge),

and viscosity of cytoplasm. The branching process through cell division leads to an increase in the

bridge number, and therefore, is critical for yielding the biased cell volume distributions (Fig. 6). The

cell surface tension provides the hydrostatic pressure, which can induce intercellular cytoplasmic flow.

The rate of the intercellular cytoplasmic flow is increased according to the geometry of the bridge (G_h)

including the radius or inversely proportional to the viscosity of cytoplasm (μ) (Fig. S2). By

considering these factors, our present study provides experimentally testable hypotheses.

Female gametogenesis proceeds through an initial phase of simultaneous cyst formation and

fragmentation, followed by a phase where only fragmentation continues (Fig. 8A). In our present

model, we focus on the cyst formation stage (embryonic days 10–14, E10-14), when cell division and

cyst fragmentation take place. Our previous work, which focused on the stage following cyst formation

stage (E14–P0 (postnatal day 0)), provided evidence for biased bridge severing and variability in cell

volume among cells within a cyst. Specifically, cells with a greater number of bridges preferentially

exhibited larger volumes. Here, we suggest that biased cell volumes begin to emerge during cyst

formation, consistent with our earlier observation that differences in cell volume are first detected at

E14.5 (Ikami et al., 2023). The small differences in cell volume observed at E14.5 are likely amplified

after cyst formation, leading to larger volume disparities among cells.

In the *Drosophila* germline cysts, microtubule-mediated directional transport of cytoplasm

is observed, resulting in a biased cell volume (Clark et al., 2007; Gutzeit, 1986; Lu et al., 2022). Our

present mechanical model is not incompatible with directional transport-based mechanisms. Rather,

our model can support the directional transport-based mechanisms to stably and robustly achieve the

biased cell size distribution. In humans, the fetal germline cysts are likely to have a branched structure

(Gondos, 1987), implying that the proposed hydrostatic pressure-dependent mechanism may also be

applicable to these systems.

Both hydrostatic pressures and surface tensions in cells/tissues are involved not only in

phenomena of cell biology such as cytokinesis, blebbing, etc. (Sedzinski et al., 2011), but also in

phenomena of tissue morphogenesis such as the blastocyst formation in mice, the behaviors of

spheroids (Chan et al., 2019; Dumortier et al., 2019; Mombach et al., 2005). Regulation of these

mechanical parameters can stabilize or unstabilize the systems, leading to their robust behaviors and

morphological changes, respectively. Our theoretical framework provides mechanical insights into

various phenomena observed during germ cell development. We have focused primarily on the

dividing female germline cysts (Fig. 1A and 8A). At a subsequent later stage, these cells stop dividing

(Fig. 8A) and an extremely biased cell volume increase occurs to form oocytes (Fig. 8B). In consistent

with this stage, the results shown in Fig. 3—simulated in the absence of cell cycle—also demonstrated
an extreme volume increase in cells with numerous bridges. In contrast, adult male germline cysts
undergo mitosis before entering meiosis, and their cell volumes are robustly maintained despite the
presence of the intercellular bridges, leading to the proper male germ cell development (Fig. 8C). This
scenario is consistent with our simulation in Fig. 4A, which showed that the cell volumes were not
biased when cell cycle was assumed but without tight cell-cell contacts. We do not rule out other
possibilities, such as a blockage of cytoplasmic flow caused by the small diameter of the bridges, etc.
Taken together, our framework demonstrates that the combination of two key mechanical
parameters—cell-cell contacts and cell cycle (cell growth and division)—allows the systems to exhibit
these three distinct biological states.

Finally, we raise the limitations of our model. Although we assumed that the cells exhibit
spherical shapes including spherical domes, the real cell shapes are not necessarily so spherical. The
cells within adult male germline cysts adhere to the surrounding tissues, leading to distorted cell shapes.
Therefore, the regulation of the hydrostatic pressures becomes more complicated than in our
theoretical model. A similar argument applies to other syncytial cells observed in the *C. elegans* gonad.
In the context of the *C. elegans* gonad, Chartier et al. incorporated its complicated geometry into their
model and demonstrated that mechanical instability, derived from the surface tension and hydrostatic
pressure, is critical for regulating biased cell volume (Chartier et al., 2021). Therefore, in order to
strictly reproduce the *in vivo* situations, more complicated modeling will be required, while our present
model for cytoplasm-sharing cells can be the basic framework.

Source

The virtual 2-cell model was implemented on the excel software. The full model was written in C
language. These are uploaded on figshare as follows (10.6084/m9.figshare.31148572).

Conflicts of interest

The authors declare no competing financial interests.

Funding

This work was supported by following grants: Japan Society for the Promotion of Science (JSPS)

Grant-in-Aid for Scientific Research (C) for H.K. (24K09478), and for Transformative Research Areas (A) for HK (24H01990, 24H01414).

Authorship contributions

Hiroshi Koyama: Writing – review & editing, Writing – original draft, Visualization, Validation, Project administration, Methodology, Investigation, Funding acquisition, Formal analysis, Data curation, Conceptualization. Kanako Ikami: Writing – review & editing, Investigation, Experiment, Conceptualization. Lei Lei: Writing − review & editing, Conceptualization. Toshihiko Fujimori: Writing – review & editing, Conceptualization.

References

Bayer EM, Benitez-Alfonso Y. 2024. Plasmodesmata: Channels Under Pressure. Annual Review of Plant Biology 75:291–317. DOI: https://doi.org/10.1146/annurev-arplant-070623

Chaigne A, Brunet T. 2022. Incomplete abscission and cytoplasmic bridges in the evolution of eukaryotic multicellularity. *Current Biology*. DOI: https://doi.org/10.1016/j.cub.2022.03.021, PMID: 35472432

Chan CJ, Costanzo M, Ruiz-Herrero T, Mönke G, Petrie RJ, Bergert M, Diz-Muñoz A, Mahadevan L, Hiiragi T. 2019. Hydraulic control of mammalian embryo size and cell fate. Nature 571:112–116. DOI: https://doi.org/10.1038/s41586-019-1309-x, PMID: 31189957

Chartier NT, Mukherjee A, Pfanzelter J, Fürthauer S, Larson BT, Fritsch AW, Amini R, Kreysing M, Jülicher F, Grill SW. 2021. A hydraulic instability drives the cell death decision in the nematode germline. Nature Physics 17:920–925. DOI: https://doi.org/10.1038/s41567-021-01235-x

Cuvelier M, Vangheel J, Thiels W, Ramon H, Jelier R, Smeets B. 2023. Stability of asymmetric cell division: A deformable cell model of cytokinesis applied to C. elegans. Biophysical Journal 122:1858–1867. DOI: https://doi.org/10.1016/j.bpj.2023.04.017, PMID: 37085996

Devany J, Falk MJ, Holt LJ, Murugan A, Gardel ML. 2023. Epithelial tissue confinement inhibits cell growth and leads to volume-reducing divisions. Developmental Cell 58:1462–1476. DOI:

https://doi.org/10.1016/j.devcel.2023.05.018, PMID: 37339629

Dumortier JG, Verge-Serandour M Le, Tortorelli AF, Mielke A, Plater L de, Turlier H, Maître J-L.
2019. Hydraulic fracturing and active coarsening position the lumen of the mouse blastocyst.
Science 365:465–468.

Farhadifar R, Röper JC, Aigouy B, Eaton S, Jülicher F. 2007. The Influence of Cell Mechanics, Cell-
Cell Interactions, and Proliferation on Epithelial Packing. Current Biology 17:2095–2104.
DOI: https://doi.org/10.1016/j.cub.2007.11.049, PMID: 18082406

Fujita M, Onami S. 2012. Cell-to-cell heterogeneity in cortical tension specifies curvature of contact
surfaces in caenorhabditis elegans embryos. PLoS ONE 7:e30224. DOI:
https://doi.org/10.1371/journal.pone.0030224, PMID: 22253922

Heisenberg CP, Bellaïche Y. 2013. Forces in tissue morphogenesis and patterning. Cell 153:948–62.
DOI: https://doi.org/10.1016/j.cell.2013.05.008, PMID: 23706734

Ikami K, Shoffner-Beck S, Weh MT, Schnell S, Yoshida S, Miranda EAD, Ko S, Lei L. 2023.
Branched germline cysts and female-specific cyst fragmentation facilitate oocyte determination
in mice. Proceedings of the National Academy of Sciences of the United States of America
120:e2219683120. DOI: https://doi.org/10.1073/pnas.2219683120, PMID: 37155904

Kimura K, Motegi F. 2025. Mechanics of oogenesis: Lessons from C. elegans germline cysts.
Seminars in Cell and Developmental Biology 175:103644. DOI:
https://doi.org/10.1016/j.semcdb.2025.103644

Koyama H, Okumura H, Ito AM, Nakamura K, Otani T, Kato K, Fujimori T. 2023. Effective
mechanical potential of cell–cell interaction explains three-dimensional morphologies during
early embryogenesis. PLoS Computational Biology 19:e1011306. DOI:
https://doi.org/10.1371/journal.pcbi.1011306

Koyama H, Umeda T, Nakamura K, Higuchi T, Kimura A. 2012. A high-resolution shape fitting and
simulation demonstrated equatorial cell surface softening during cytokinesis and its promotive
role in cytokinesis. PLoS ONE 7:e31607. DOI: https://doi.org/10.1371/journal.pone.0031607,
PMID: 22359606

Kurotaki Y, Hatta K, Nakao K, Nabeshima Y-I, Fujimori T. 2007. Blastocyst Axis Is Specified
Independently of Early Cell Lineage But Aligns with the ZP Shape. Science (New York, N.Y.)
316:719–23. DOI: https://doi.org/10.1126/science.1138591, PMID: 17446354

Landesman Y, Goodenough DA, Paul DL. 2000. Gap Junctional Communication in the Early
Xenopus Embryo. The Journal of Cell Biology 150:929–936.

Lv Z, de-Carvalho J, Telley IA, Großhans J. 2021. Cytoskeletal mechanics and dynamics in the
Drosophila syncytial embryo. Journal of Cell Science 134:jcs246496. DOI:
https://doi.org/10.1242/jcs.246496, PMID: 33597155

Miyoshi H, Satoh SK, Yamada E, Hamaguchi Y. 2006. Temporal change in local forces and total
force all over the surface of the sea urchin egg during cytokinesis. Cell Motility and the
Cytoskeleton 63:208–221. DOI: https://doi.org/10.1002/cm.20118, PMID: 16470543

Mombach JCM, Robert D, Graner F, Gillet G, Thomas GL, Idiart M, Rieu J. 2005. Rounding of
aggregates of biological cells : Experiments and simulations. Physica A 352:525–534. DOI:
https://doi.org/10.1016/j.physa.2005.02.008

Okuda S, Inoue Y, Adachi T. 2015. Three-dimensional vertex model for simulating multicellular
morphogenesis. Biophysics and Physicobiology 12:13–20. DOI:
https://doi.org/10.2142/biophysico.12.0

Phillips Rob, Kondev Jane, Theriot Julie. 2008. Physical biology of the cell, 1st ed. Garland Science.

Priti A, Ong HT, Toyama Y, Padmanabhan A, Dasgupta S, Krajnc M, Zaidel-Bar R. 2018. Syncytial
germline architecture is actively maintained by contraction of an internal actomyosin corset.
Nature Communications 9. DOI: https://doi.org/10.1038/s41467-018-07149-2, PMID:
30410005

Runser S, Vetter R, Iber D. 2024. SimuCell3D: three-dimensional simulation of tissue mechanics
with cell polarization. Nature Computational Science 4:299–309. DOI:
https://doi.org/10.1038/s43588-024-00620-9

Sagner A, Merkel M, Aigouy B, Gaebel J, Brankatschk M, Jülicher F, Eaton S. 2012. Establishment
of global patterns of planar polarity during growth of the drosophila wing epithelium. Current
Biology 22:1296–1301. DOI: https://doi.org/10.1016/j.cub.2012.04.066, PMID: 22727699

Savin T, Kurpios N a, Shyer AE, Florescu P, Liang H, Mahadevan L, Tabin CJ. 2011. On the growth
and form of the gut. Nature 476:57–62. DOI: https://doi.org/10.1038/nature10277, PMID:
21814276

Sedzinski J, Biro M, Oswald A, Tinevez JY, Salbreux G, Paluch E. 2011. Polar actomyosin

contractility destabilizes the position of the cytokinetic furrow. Nature 476:462–468. DOI:

https://doi.org/10.1038/nature10286, PMID: 21822289

Wang X, Li L, Shao Y, Wei J, Song R, Zheng S, Li Y, Song F. 2021. Effects of the Laplace pressure

on the cells during cytokinesis. iScience 24:102945. DOI:

https://doi.org/10.1016/j.isci.2021.102945

Wu Y, Pegoraro AF, Weitz DA, Janmey P, Sun SX. 2022. The correlation between cell and nucleus

size is explained by an eukaryotic cell growth model. PLoS Computational Biology 18. DOI:

https://doi.org/10.1371/journal.pcbi.1009400, PMID: 35180215

Xie J, Huck WTS, Bao M. 2024. Unveiling the Intricate Connection: Cell Volume as a Key

Regulator of Mechanotransduction. Annual Review of Biophysics 53:299–317. DOI:

https://doi.org/10.1146/annurev-biophys-030822

Figure 1

Figure 1: Structures of fetal female germline cysts and mechanical properties of cells. A. Structure of
a fetal female cyst. Cells adhere via E-cadherin to generate cell-cell contacts (purple rectangle), where
intercellular bridges (green circle) enable cytoplasmic sharing. The microscopic image is reproduced
from (Ikami et al., 2023); a single clone is labeled by YFP. YFP, cytoplasm; TEX14, intercellular
bridges. B. Soap bubble model. A cell is modeled as a soap bubble or liquid droplet with hydrostatic
pressure (P), surface tension (λ), and curvature (H) related by the Laplace pressure equation ($P=2\lambda H$).
Subscripts denote cell IDs (e.g., P_A for A-cell). Cytoplasmic flow occurs due to pressure differences
between connected cells (e.g., P_A vs. P_B).

Figure 2

Figure 2: Virtual 2-cell model. A. Modeling of a virtual 2-cell system. The number of cell-cell contacts
with intercellular bridges (magenta box) remains constant. One cell cycle includes cell growth,
intercellular cytoplasmic flow, cell division, and neglect of proliferated cells. B. Calculation of inner
pressure. During cell growth and cytoplasmic flow, inner pressures were calculated based on the
Laplace pressure, considering spherical dome shapes due to cell-cell contact regions. Illustrates
spherical dome and spherical cap shapes.

Figure 3

Figure 3: Simulation in virtual 2-cell model without cell division and cell growth. A. Simulation results

without overlapping region ($\Omega = 0\%$). Bridge number in A-cell (N_A) = 1; B-cell (N_B) = 1 or 2. Volume

changes for initial ratio $V_B(0)/V_A(0) = 1.0$ (unstable equilibrium) and 1.01 are shown. B. Simulation

results with overlapping region ($\Omega \neq 0\%$). Bridge number in A-cell (N_A) = 1; B-cell (N_B) = 1 or 2.

Volume changes for initial ratios of 1.0 and ~0.89 are shown. C. Conditions defining unstable

equilibrium states for various Ω values. Lines represent thresholds of $V_B(0)/V_A(0)$ for each Ω value:

above the line, B-cell volume increases ($V_A(T)<V_B(T)$); below the line, A-cell volume increases (V_A

$(T)>V_B(T)$). D. Summary of the results in the absence of cell cycle. In the presence of the overlapping

region, the B-cell is likely to increase its volume, especially when the B-cell has more bridges. In the

absence of the overlapping region, the probability of whether the A-cell or B-cell increases its volume

is even, and one of the cells completely loses its volume.

Figure 4

Figure 4: Simulation in virtual 2-cell model with both cell division and cell growth. A. Simulation

results with $\Omega = 0$, 10, or 20% and N_B=2 or 3 (N_A=1). Volume changes for initial ratio $V_B(0)/V_A(0) =$

~0.89 and 0.30 are shown. In the first panel ($\Omega = 0$), the simulation result is shown for a long

simulation time ($t = 0$~50). In the bottom panel ($\Omega = 0.1$), the behaviors under the different initial

conditions ($V_B(0)/V_A(0)$) are shown. B. Conditions defining threshold values of $V_B(0)/V_A(0)$ for

various Ω values. For $\Omega > 0$, $N_B \geq 2$: above line, B-cell volume increases; below line, A-cell volume

increases. For $\Omega = 0$ or N_B=1: above line, $V_A(T) = V_B(T)$; below line, A-cell volume increases. Asterisk

and cross-marked points correspond to conditions in A. C. Final volume ratios ($V_B(T)/V_A(T)$) at

convergence for various Ω values under conditions where B-cell volume increased. Shows specific

ratios for N_B=1,2,3 when $\Omega = 0.2$. D. Summary of the results in the presence of cell cycle. In the

presence of the overlapping region, the B-cell is likely to increase its volume, especially when the B-

cell has more bridges. In the absence of the overlapping region, both the A- and B-cells can maintain

their volume, and the volume of these cells is equalized if the bridge number in the B-cell is less.

Figure 5

650 Figure 5: Definition of the full model. A. Schematic of the full model. Connection patterns are
651 randomly determined after cell division, and intercellular bridges are stochastically severed, leading
652 to varied cyst structures before the next cell cycle. B. Histograms of cell numbers within cysts at 2-
653 cell (1 cycle), 8-cell (3 cycles), and 32-cell (5 cycles) stages. C. Histogram of bridge numbers per cell
654 for each stage.

Figure 6: Factors affecting cell volume in the full model. A-i. Cell volumes for cells with different bridge numbers under varying Ω and P_0 conditions. A-ii. Effect of Ω on the volume of cells with the different number of the bridges, showing specific volume increase for cells with more bridges at larger Ω. A-iii. Effect of P_0 on cell volume, indicating no significant changes with varying P_0. B-i. Linear cyst model where no branches are formed, limiting bridge number per cell to less than 3. The histogram was obtained at the 64-cell stage under $\Omega = 10\%$ and $P_0 = 0.25$. B-ii. Cell volumes in unbranched cysts. Compares volumes of cells with 1 versus 2 bridges, showing limited volume increase even with the maximum bridge count in linear structures.

Figure 7

Figure 7: Effect of cell-cell contact without bridges on cell volume. A. Virtual 2-cell model assuming

contacts with or without bridges. This panel illustrates scenarios where B-cell has additional contacts,

some with bridges and some without. See text for details. B. Threshold values of $V_B(0)/V_A(0)$ for

volume change, similar to Figure 4B, considering contacts with/without bridges. For $\Omega > 0$ (i.e., 5%;

light blue and red lines): above line, B-cell volume increases; below line, A-cell volume increases. For

$\Omega = 0$ (purple and yellow lines): above line, $V_A(T) = V_B(T)$; below line, A-cell volume increases. C.

Final converged volume ratios ($V_B(T)/V_A(T)$) similar to Figure 4C, considering contacts with/without

bridges. The legend for the lines is shown in B. Cells harboring contacts without bridges increase the

final volume ratio (yellow and red lines), but at a slower rate than those harboring contacts with bridges

(purple and light blue lines).

Figure 8

Figure 8: Other type of germline cysts

A. Schematic representation of developmental stages in female germ cells. See text for details.

B. Structure of a fetal female cyst at later stage. The cells do not undergo cell cycle, and a smaller
population of the cells shows extreme increase in volume, leading to oocyte formation. The
microscopic image was acquired according to the procedures described previously (Ikami et al., 2023).
YFP, cytoplasm; TEX14, intercellular bridges.

C. Structure of an adult male cyst before meiosis. Cells exhibit weaker adhesion, but intercellular
bridges are present. The cells undergo cell cycle. The microscopic image is reproduced from (Ikami
et al., 2023). YFP, cytoplasm; E-cad, E-cadherin.

695