Microtubule occupancy at kinetochores links checkpoint

silencing with mitotic memory

Joana Soares-de-Oliveira[1,2,3#], Naoyuki Okada[1,2,#], Tobias Kletter[1,2,#],

Elias S. Weiler[1,2,3], António J. Pereira[1,2], Helder Maiato[1,2,4,*]

[1] Chromosome Instability & Dynamics Group, i3S - Instituto de Investigação e Inovação em

Saúde, Universidade do Porto, Rua Alfredo Allen 208, 4200-135 Porto, Portugal

[2] Instituto de Biologia Molecular e Celular, Universidade do Porto, Rua Alfredo Allen 208,

4200-135 Porto, Portugal

[3] ICBAS-Instituto de Ciências Biomédicas Abel Salazar, University of Porto, Rua Jorge de

Ferreira 228, 4050-313 Porto

[4] Cell Division Group, Department of Biomedicine, Faculdade de Medicina, Universidade do

Porto, Alameda Prof. Hernâni Monteiro, 4200-319 Porto, Portugal

[#] Equal contribution

* Correspondence to: maiato@i3s.up.pt

Abstract

The spindle assembly checkpoint (SAC) promotes faithful chromosome segregation by delaying mitosis until all kinetochores attach to spindle microtubules. Paradoxically, a p53-dependent memory mechanism - the "mitotic stopwatch" - blocks daughter cell proliferation after unusually prolonged mitoses. Understanding how the SAC coordinates with the mitotic stopwatch is critical to disentangle this conundrum. Here, we found that microtubule occupancy at kinetochores is a cornerstone linking SAC silencing with mitotic memory. By combining live-cell with super-resolution microscopy, FRAP, laser microsurgery and molecular perturbations in Indian muntjac fibroblasts, we show that SAC silencing at kinetochores is gradual, non-uniform and confined to highly-localized microtubule attachments. Augmin promotes timely SAC silencing with maximal microtubule occupancy at kinetochores, whereas MPS1/CDK1 inhibition bypasses this condition. Partial microtubule occupancy delays SAC silencing, increases segregation errors and blocks daughter cell proliferation. Thus, timely SAC silencing with high microtubule occupancy avoids "bad memories" of mitosis to allow daughter cell proliferation.

Introduction

Faithful chromosome segregation during mitosis is essential for tissue homeostasis. Errors in chromosome segregation may lead to aneuploidy compromising cell viability, or drive chromosomal instability, a hallmark of human cancers implicated in tumor evolution, metastasis, therapy resistance, and reduced patient survival [1]. To promote faithful chromosome segregation during mitosis, the spindle assembly checkpoint (SAC) delays anaphase until all kinetochores attach to mitotic spindle microtubules [2]. At the root of the SAC signaling cascade, a MAD1/MAD2 complex recruited to unattached kinetochores initiates a catalytic process that activates free cytosolic MAD2 to bind CDC20, BUBR1 and BUB3, producing a diffusible Mitotic Checkpoint Complex (MCC). The MCC inhibits the Anaphase Promoting Complex/Cyclosome (APC/C), an E3-ubiquitin ligase that otherwise targets Cyclin B1 and Securin for degradation by the proteasome (reviewed in [3-5]). This releases the cysteine-protease Separase to cleave the Cohesin molecules that hold sister-chromatids together, thereby triggering their initial separation in anaphase, while preventing SAC reactivation during anaphase [6,7].

Upon kinetochore attachment to microtubules, SAC strength gradually decreases as result of the removal of MAD1/MAD2, eventually halting the catalytic cascade underlying MCC production [8,9]. The removal of MAD1/MAD2 from kinetochores during SAC silencing is mediated by PP1 and PP2A phosphatases, which are recruited by the outer kinetochore component KNL1 to oppose the action of key mitotic kinases, such as CDK1 and MPS1[10-15]. As microtubules establish end-on attachments, MPS1 is outcompeted from NDC80 complexes at kinetochores[16,17], and the MAD1/MAD2 complex is removed by dynein-mediated stripping as the fibrous corona disassembles[18-22].

Pioneering correlative light and electron microscopy work in PtK1 cells showed that their kinetochores bind more microtubules in anaphase than in metaphase, indicating that SAC silencing during normal mitosis can occur at ~85% microtubule occupancy at kinetochores [23]. Classic micromanipulation experiments in grasshopper spermatocytes further indicated that weakly attached kinetochores with less than 40% microtubule occupancy show greatly

diminished MAD1/MAD2 recruitment when compared with fully unattached kinetochores, delaying but not preventing anaphase onset [24]. Indeed, in this system, high microtubule occupancy was found to be necessary for the reliable and complete removal of MAD1/MAD2 from kinetochores [24]. In contrast, two recent studies in human HeLa cells overexpressing individual or combinations of Hec1 phosphomimetic mutants to create intermediate average microtubule attachment states at kinetochores estimated that 20–50% of the normal metaphase microtubule occupancy is sufficient to trigger complete and uniform removal of MAD1/MAD2 from kinetochores [25,26]. However, this switch-like model remains difficult to reconcile with the finding that the rate of MAD1/MAD2 loss from kinetochores increases with microtubule occupancy, with SAC silencing under conditions of low microtubule occupancy resulting in extensive mitotic delays [24-26]. A third study in human RPE1 cells used a microtubule depolymerizing drug to reduce microtubule occupancy at kinetochores [27]. Based on direct microtubule counting by electron microscopy, this study showed that ~70% microtubule occupancy at metaphase kinetochores is sufficient to silence the SAC, with less than 5 min delay relative to controls, but at the price of increasing the incidence of lagging chromosomes during anaphase [27]. Thus, although cells can silence the SAC with low microtubule occupancy, there may be selective pressure for high microtubule occupancy at kinetochores before initiating anaphase, ensuring proper chromosome segregation and avoiding significant mitotic delays [27,28]. This hypothesis gains weight in light of the recently uncovered "mitotic stopwatch" complex formed by 53BP1 and the deubiquitinase USP28 that stabilizes p53 in response to even moderate delays in mitosis, leading to subsequent cell cycle arrest or death in G1, independently of DNA damage [29-36]. However, how SAC silencing is coordinated with the mitotic stopwatch to control daughter cell proliferation remains unknown.

Here we set to investigate whether and how the SAC responds to highly-localized microtubule attachments at kinetochores, while testing whether the extent of microtubule occupancy underlying SAC silencing impacts daughter cell proliferation. Overall, our data support a gradual, non-uniform (i.e. confined to highly-localized microtubule attachments), SAC-silencing mechanism that is

assisted by Augmin to promote high microtubule occupancy at kinetochores for
timely and faithful chromosome segregation. Importantly, if cells ultimately
silence the SAC after experiencing significant mitotic delays due to low
microtubule occupancy at kinetochores, the resulting daughter cells are
bookmarked to prevent their proliferation and the potential propagation of
segregation errors.

Results

MAD1 is gradually removed from Indian muntjac kinetochores in a microtubule-dependent manner

According to the switch-like SAC-silencing model, kinetochores respond as a single unit to silence the SAC after just a few microtubule attachments are established [25,26]. However, because human kinetochores cannot be resolved by conventional light microscopy, this postulate has so far been impossible to test experimentally. To overcome this limitation, we took advantage of the unique cytological features of female Indian muntjac fibroblasts, which carry only six chromosomes with naturally "super-resolved" kinetochores that arose by tandem and centric fusions of smaller ancestral chromosomes [37-43]. As a first step towards investigating SAC silencing in this system, we determined the time between MAD1 disappearance from kinetochores and anaphase onset by tracking individual kinetochores in live, hTERT-immortalized, female Indian muntjac fibroblasts stably expressing mScarlet-CENP-A (a constitutive kinetochore marker) and Venus-MAD1 [8]. We found that MAD1 signal at kinetochores became essentially undetectable ~20 min before anaphase onset (Fig. S1A, B), in agreement with previous reports in other vertebrate cell lines [44,45] and consistent with the temporal framework of anaphase onset upon laser-mediated ablation of the last unattached kinetochore [2].

To determine the kinetics of SAC silencing in response to microtubule attachments at individual kinetochores in live Indian muntjac fibroblasts, we combined stable expression of fluorescent Venus-MAD1 with carefully titrated SiR-tubulin [41] under slight cell compression to improve visualization [42]. As a control, cytoplasmic Venus-MAD1 signal was simultaneously measured, and photobleaching was found to be negligible throughout the course of the experiment (Fig. S1C). MAD1 decayed gradually at individual kinetochores as chromosomes bi-oriented and aligned at the equator during prometaphase (Figure 1A, C), with a rate of decay that was well fit to a single exponential (Figure 1E and Fig.S2), in agreement with previous reports in Ptk2 cells [46]. Noteworthy, pole-proximal chromosomes aligned last and did so tangentially along the

curvature of the spindle edge. Moreover, their leading kinetochores were enriched with MAD1 from the onset of congression, with the signal gradually decreasing as chromosomes approached the equator (Figure 1A, A', A''). Importantly, MAD1 typically decayed first on trailing kinetochores, likely reflecting differences in the timing or efficiency in the formation of stable end-on microtubule attachments compared with leading kinetochores. (Figure 1A, A', A''). These results are fully consistent with congression models where pole-proximal, mono-oriented chromosomes use plus-end-directed motors on the leading kinetochore to initiate lateral gliding along pre-existing spindle microtubules towards the equator [46-49], while at odds with models where stabilization of end-on attachments and bi-orientation take place before pole-proximal chromosomes initiate congression [50].

To test whether gradual MAD1 decay at kinetochores during mitosis depends on microtubule occupancy, we monitored MAD1 levels in the presence of 1 µM nocodazole, which depolymerized all spindle microtubules. As expected, under these conditions, MAD1 localized on every kinetochore for over 20 min, without any measurable decay (Figure 1B, C and Fig. S2). These data indicate that gradual SAC silencing during mitosis depends on the establishment of microtubule attachments at kinetochores.

MAD1 is rapidly removed from Indian muntjac kinetochores upon MPS1 or CDK1 inactivation, regardless of microtubule occupancy

MPS1 and CDK1 kinases are critical to maintain SAC signaling by counteracting the activities of PP1 and PP2A phosphatases required for mitotic exit [10-15]. Therefore, we next investigated the kinetics of MAD1 decay at kinetochores after acute MPS1 or CDK1 inactivation with MPS1-IN-1 [51] or RO 3306 [52], respectively. We found that MAD1 decayed more abruptly from kinetochores after MPS1 or CDK1 inhibition, regardless of microtubule occupancy, reaching 50% of the initial levels 3.8x or 1.5x faster than during normal mitosis, respectively (Figure 1B, C, E and Fig. S2). These data indicate that inactivation of MPS1 or CDK1 can bypass the requirement of a given microtubule occupancy at kinetochores to silence the SAC.

MAD1 removal depends on kinetochore size

A prediction from switch-like models that favor independent microtubule binding at kinetochores and where a given percentage of occupancy is key to uniformly silence the SAC [25,26] is that small or large kinetochores would simultaneously reach the same percentage of occupancy (and, consequently, have similar MAD1 half reduction times), despite binding a different total number of microtubules. However, despite ample evidence of kinetochore size differences in humans [53-61], this prediction is currently impossible to test experimentally in human cells for the same reasons evoked before. In contrast, the large kinetochore from chromosome 3+X in female Indian muntjac cells can be unequivocally distinguished from those on the smaller chromosomes 1 and 2, allowing us to track MAD1 decay as a function of kinetochore size [38,40,41]. Surprisingly, we found that MAD1 signal on the large kinetochores of chromosome 3+X requires ~38% more time to become undetectable, when compared with smaller kinetochores on chromosomes 1 and 2 (Figure 1D, E, and Fig. S2). These findings suggest that switch-like models do not fully account for how the SAC is normally silenced.

MAD1 is immobile within kinetochores

In order to obtain a deeper understanding of the mechanisms underlying SAC silencing we investigated MAD1 dynamics within kinetochores. Two possible scenarios could be envisioned: 1) if MAD1 is mobile within kinetochores, few microtubule attachments might be sufficient to "extinguish" all MAD1, for example by Dynein-mediated stripping along microtubules [18,19] (Figure 2A); 2) if MAD1 is immobile within kinetochores, Dynein-mediated stripping along microtubules would be confined to the sites or domains where microtubules are attached (Figure 2A). To distinguish between these two scenarios, we photobleached Venus-MAD1 on a fraction (up to 50%) of the large kinetochore from chromosome 3+X in the absence of microtubules. Clear predictions about MAD1 mobility within kinetochores could be made, depending on the FRAP profile. If MAD1 is mobile within kinetochores, the photobleached fraction of the kinetochore is expected to recover due to migration of fluorescent Venus-MAD1 molecules from the unbleached fraction, resulting in the convergence of signals

to an intermediate and uniform level along the entire kinetochore (Figure 2A'). In

contrast, if MAD1 is immobile within kinetochores, the photobleached fraction of

the kinetochore would be expected to only recover fluorescent Venus-MAD1

molecules that are exchanging with the cytoplasmic pool, while the unbleached

fraction would remain stable over time (Figure 2A'). We were successful in

partially photobleaching Venus-MAD1 in a fraction of the kinetochore, with

minimal/unintended photobleaching of Venus-MAD1 on the remaining fraction

(Figure 2B-B''). Importantly, after photobleaching of the intended fraction of the

kinetochore, Venus-MAD1 fluorescence recovered up to ~50%, with a half

recovery time of 2-6 seconds, while remaining stable (or increasing slightly; see

ahead) over time in the unbleached fraction (Figure 2B-B'', D, E). As a control, we

photobleached Venus-MAD1 on entire kinetochores, which resulted in identical

recovery parameters when compared to partial photobleaching, and in line with

previous measurements in other mammalian systems [44,45] (Figure 2C-C'', D, E).

Because fluorescence recovery of the mobile fraction of Venus-MAD1 after

photobleaching entire kinetochores is exclusively due to exchange with the

cytoplasmic pool [44,45], we concluded that this pool also accounts for the observed

fluorescence recovery of Venus-MAD1 on partially photobleached kinetochores.

Taken together, these data indicate that MAD1 is essentially immobile within

kinetochores, suggesting that it can only be removed from spatially confined

regions upon the establishment of highly-localized microtubule attachments or by

a microtubule-independent SAC-silencing mechanism, such as the one involving

CDK1 or MPS1 inactivation.

MAD1 removal along individual kinetochores is non-uniform and confined

to highly-localized microtubule attachments

The conclusion from our FRAP experiments implies that, in a normal mitosis,

there may be low microtubule occupancy states in which MAD1 removal is limited

to regions where microtubule attachments are established. To test this prediction,

we monitored Venus-MAD1 decay along the entire kinetochore length as

microtubules attach during mitosis. Near-simultaneous tracking of individual

kinetochores, MAD1 and microtubules in living cells revealed a non-uniform

decay of MAD1 along kinetochores (Figure 3A, B). To further dissect the
relationship between non-uniform MAD1 decay and microtubule occupancy
along kinetochores, we used super-resolution CH-STED microscopy [62] and 3D
image rendering to inspect (fixed) late prometaphase cells with only partial MAD1
signal along kinetochores (Figure 4A, B and Fig. S3A, B). We found clear
examples where fractions of the kinetochore had end-on attached microtubules
with no detectable MAD1, while other fractions were devoid of microtubules and
clearly enriched for MAD1 (Figure 4A, B and Fig. S3A, B). A line scanning profile
along the entire length of several such individual kinetochores confirmed a
negative correlation between MAD1 signal and microtubule density in 15/21
(71.4%) of the cases (Figure 4C, D and Fig. S3A', B').

To improve the resolution of our analyses we divided the longitudinal axis of
partially attached kinetochores into multiple subsections and quantified MAD1
and microtubule intensity at each subsection (Figure 4E). We found a weak, yet
highly significant, negative correlation between MAD1 and microtubule signal
intensity along kinetochores (Figure 4F and Fig. S3C). Of note, the resulting
L-shape curve indicates that while high microtubule signal is a good predictor of
the absence of MAD1 at kinetochores, the opposite is not the case, suggesting
that there are regions at kinetochores without microtubules that have no
detectable MAD1, consistent with models evoking cooperative microtubule
binding at kinetochores [63,64]. Overall, these data show that SAC silencing at
kinetochores is non-uniform and confined to highly-localized microtubule
attachments.

Local perturbation of kinetochore function results in a local SAC response
to microtubule attachments/detachments
To directly test how the SAC responds to localized microtubule
attachments/detachments, we used laser microsurgery to partially ablate
individual metaphase kinetochores (and, consequently, microtubules that were
attached to the ablated region) upon SAC silencing in live Indian muntjac
fibroblasts (Figure 5A). These cells stably expressed GFP-CENTRIN-1 to mark
spindle pole positions, 2xGFP-CENP-A to define the kinetochore target, and

mScarlet-MAD1 to monitor SAC status before and after surgery (Figure 5B). We
found that, within few minutes after partial kinetochore ablation, the unperturbed
sister kinetochore typically bent towards the attached spindle pole without any
detectable MAD1 (Figure 5A-C). However, ~5 min after bending, MAD1 was
often recruited to the tip of the bent unperturbed sister kinetochore, presumably
due to highly-localized microtubule detachments (Figure 5A-D). In other cases,
MAD1 was recruited to a more central region of the bent unperturbed sister
kinetochore (Fig. S4A, A', B). Strikingly, in all cases MAD1 accumulation was
highly localized and distinct from the uniform accumulation of MAD1 on fully
unattached kinetochores resulting from nocodazole treatment (Figure 5B, D, E).
Therefore, local kinetochore perturbations result in a highly-localized SAC
response.

To assist us in the interpretation of the laser microsurgery data and determine
whether the SAC was responding to microtubule detachments in the bent
unperturbed sister kinetochore, we performed correlative super-resolution
CH-STED microscopy after fixation of cells that showed MAD1 recruitment upon
partial sister kinetochore laser ablation by live imaging (Figure 5F, F' and Fig.
S4C, C'). We found that MAD1 accumulation, either at the tip or at a more central
region of the unperturbed sister kinetochore, negatively correlated with
microtubule density (Figure 5F', F" and Fig. S4C', C"). We reasoned that MAD1
accumulation near the tip of the unperturbed kinetochore may result from highly
localized microtubule detachments due to lack of tension in the partially ablated
sister. In other cases, localized forces applied to both ends of the unperturbed
kinetochore may generate enough tension along its length to allow a few
microtubules to remain stably attached near the tip, resulting in more central
accumulation of MAD1. These results demonstrate that the SAC responds locally
to highly localized microtubule attachment/detachment events along
kinetochores.

Low microtubule occupancy at kinetochores after Augmin depletion
delays MAD1 removal and SAC silencing

Next, we sought to investigate the cellular consequences of low microtubule

occupancy at kinetochores. To do so we monitored SAC silencing after

RNAi-mediated depletion of HAUS6, a subunit of the Augmin complex required

for K-fiber maturation [43,65]. We found that MAD1 decay from kinetochores was

slowed-down in HAUS6-depleted cells, as revealed by ~50% increase in the half

reduction time of MAD1 signal at kinetochores relative to mock-depleted controls

(Figure 6A-C, Figure S2). Moreover, even when HAUS6-depleted cells were

treated with the proteasome inhibitor MG132 for 1h to provide more time for

K-fiber maturation, many more (66% vs. 12%) cells had at least one

MAD1-positive kinetochore and overall higher MAD1 levels at kinetochores when

compared with control metaphase cells (but lower than prometaphase levels)

(Figure 6D, E). Closer inspection of mock- and HAUS6-depleted fixed cells with

CH-STED super-resolution microscopy revealed that, while kinetochores in

control mock-depleted metaphase cells treated with MG132 were abundantly

occupied with microtubules and without detectable MAD1 signal, in

HAUS6-depleted cells treated with MG132 MAD1 signal could still be detected

along the kinetochores, despite interspersed end-on microtubule attachments

(Figure 6F). However, unlike in control cells with clear partially attached

kinetochores that showed a negative correlation between MAD1 and microtubule

signal (r = -0.29, p<1e-10; Figure S5E), in HAUS6-depleted cells this correlation

was lost (r = -0.06, p=0.14; Figure S5D, E and Figure S6C), suggesting that the

resulting perturbations of microtubule occupancy were beyond our resolution

capacity (Fig. S5A-C and Fig. S6A-B'). Overall, these data are in line with

previous findings in human cells [25,26] indicating that low microtubule occupancy at

kinetochores delays MAD1 removal from kinetochores and consequently SAC

silencing.

324　Low microtubule occupancy delays SAC silencing, increases segregation

325　errors, and blocks daughter cell proliferation

To further characterize the cellular consequences of low microtubule occupancy

at kinetochores, we tracked HAUS6-depleted cells over several days by live-cell

microscopy and determined their fates (Figure 7A). Mitoses in HAUS6-depleted

cells were significantly delayed and error-prone, proportional to the extent of
HAUS6 depletion over time (Figure 7A' and B). When inspecting the outcomes
resulting from mitotic delays in HAUS6-depleted cells more closely, we found
three regimens. The majority of cells with mitotic durations below 200 minutes
divided without segregation errors or spindle defects (Figure 7C). However, after
a delay of approx. 200 minutes, virtually all cells divided with mitotic defects
(Figure 7C), pointing towards persistently immature K-fibers at anaphase onset
[27]. Strikingly, most cells that were stuck in mitosis for longer than 400 minutes
eventually died in mitosis (Figure 7C).

To determine the fate of daughter cells from mothers that *did* divide despite
immature K-fibers, we defined a cohort of cells that underwent mitosis within 12
and 36 hours of HAUS6 depletion and followed their offspring for 60 hours
(Figure 7A). Strikingly, 74% of daughter cells arising from mitoses with durations
longer than approx. 90 minutes stopped proliferating, whereas only 32% of
daughter cells derived from shorter mitosis arrested (Figure 7D and S7F), in line
with previous findings in human cells, suggesting active surveillance by the
"mitotic stopwatch" [29,31,32,34]. To confirm that such a "memory" of mitotic duration
is conserved in Indian muntjac cells, we tracked daughter cell fates after
releasing cells from up to 6 hours of mitotic arrest mediated by the Eg5-inhibitor
monastrol (Fig. S7A). In agreement with our observations in HAUS6-depleted
cells, mother cells that experienced mitotic times longer than approx. 90 minutes
gave rise to daughters that showed increased cell cycle lengths or stopped
proliferating (>90 min mitosis - 61% of daughters arrested; <90 min mitosis - 30%
of daughters arrested; Fig. S7B-F).

Together, these observations suggest the presence of a conserved mitotic
stopwatch mechanism surveilling mitotic durations. Indeed, we were able to
detect a significant accumulation of USP28 bound to 53BP1 – two components of
the mitotic stopwatch complex – in Indian muntjac cells that experienced a
prolonged mitosis (note that available anti-p53 monoclonal antibodies do not
recognize the Indian muntjac protein under denatured conditions; Fig. S7G, H).
However, since prolonged mitotic durations under low microtubule occupancy
(i.e. HAUS6 depletion) or monastrol perturbation were tightly associated with

mitotic errors (Figures 7B, C and Fig. S7C, D) that may independently block

daughter cell proliferation [66-69], we sought to disentangle the two events. To this

end, the low chromosome number of Indian muntjac fibroblasts allowed us to

unequivocally exclude cells dividing with errors from our cell fate analysis (Figure

7E and Fig. S7E, E', F). Remarkably, mother cells with increasing mitotic

durations - but dividing error-free - were still increasingly more likely to give rise to

daughter cells that arrest in G1 (Figure 7E and Fig. S7E, E', F), suggesting active

"mitotic stopwatch" surveillance of mitotic duration.

Discussion

While there is a large consensus that SAC silencing can occur with low microtubule occupancy at kinetochores [23-27,46], whether and how SAC signaling along vertebrate kinetochores responds to highly localized microtubule attachments has remained a central open question. Current models suggest that mammalian kinetochores respond as a single unit that globally halts SAC signaling upon partial microtubule attachments [25,26,46]. As so, low microtubule occupancy would cause kinetochores to respond as a whole and predicts a uniform loss of MAD1 along the kinetochores. Moreover, these switch-like models imply the existence of a sensitive microtubule "counting" mechanism that determines SAC silencing when a given occupancy at each individual kinetochore is reached. However, in virtue of the sub-diffraction nature of most vertebrate (humans included) kinetochores, this prediction had never been directly tested and the essence of a putative microtubule "counting" mechanism remains elusive. In this regard, Indian muntjac kinetochores offer a unique setting to test these ideas. In stark contrast with the predictions from switch-like SAC-silencing models, we found that the removal of the SAC protein MAD1 from Indian muntjac kinetochores is gradual, non-uniform, correlates with local microtubule density and kinetochore size, and depends on Augmin-mediated maturation of K-fibers.

In line with previous work [46], MAD1 signal decay from Indian muntjac kinetochores was well fit by a single exponential, indicating that SAC silencing in this system faithfully replicates established kinetics in other vertebrate systems. Switch-like models that favor independent microtubule binding at kinetochores and where a given percentage of occupancy is key to uniformly silence the SAC [25,26], predict that the fraction of unattached sites is expected to decay exponentially, with a timescale that does not depend on kinetochore size. Interestingly, however, we found that MAD1 decay on large kinetochores was 38% slower compared to small kinetochores. While the underlying reasons for these differences remain unclear, this finding suggests that switch-like models do not fully account for how the SAC is normally silenced. Indeed, both live-cell imaging and super-resolution CH-STED analysis in Indian muntjac cells revealed

a non-uniform distribution of MAD1 on partially attached kinetochores that

negatively correlated with microtubule occupancy. Moreover, laser microsurgery

experiments demonstrated that once silenced, the SAC remains sensitive to

highly-localized microtubule detachments from kinetochores, even when more

than 50% occupancy persists. Lastly, experimental perturbation of microtubule

occupancy at kinetochores by interfering with K-fiber maturation significantly

delayed MAD1 removal and SAC silencing. Therefore, although low microtubule

occupancy at kinetochores may eventually silence the SAC, there is an

unavoidable impact on the respective mitotic duration. This has important

implications in light of the recently-uncovered "mitotic stopwatch" mechanism,

where even mild mitotic delays elicit a subsequent

USP28-53BP1-p53-dependent cell cycle arrest in G1 [29,31,32,35,70], and the

observation that SAC silencing with low microtubule occupancy increases the

incidence of lagging chromosomes during anaphase, potentially leading to

aneuploidy [27]. As so, the mechanism underlying SAC silencing at individual

kinetochores is likely to have evolved to maximize microtubule occupancy within

a strict temporal window to avoid "bad memories" of mitosis, and we show here

that the Augmin complex, through its role in K-fiber maturation [43,65], is

instrumental in this process (Figure 8).

But how do partially attached kinetochores that are locally sensitive to

microtubule attachments ultimately remove all MAD1/MAD2? In line with a model

in which the number of unattached kinetochores ultimately determines the rate of

MCC production that proportially inhibits the APC/C [8,9], partially-attached

kinetochores may produce less MCC than fully unattached kinetochores [24].

Consequently, the APC/C would become sufficiently active and allow for slower,

yet steady, Cyclin B1 degradation (and CDK1 inactivation), ultimately silencing

the SAC after an extensive mitotic delay [24]. Whether kinetochores eventually

reach normal microtubule occupancy by the time the SAC is silenced after a

delay remains unclear (Figure 8), but it should be noted that even cells that are

unable to silence the SAC after extremely prolonged mitotic delays due to the

complete absence of microtubules degrade Cyclin B1 and eventually exit mitosis

due to residual APC/C activity [71,72]. Our finding that CDK1 inhibition triggers SAC

silencing irrespective of microtubule occupancy further supports this
interpretation (Figure 8).

One may argue that the compound architecture of Indian muntjac
kinetochores does not reflect what is normally found in other vertebrate species,
including humans. However, we have no evidence that kinetochore architecture
in Indian muntjac reflects a significant deviation at the level of the core
microtubule-binding moduli, or other functional differences, including MAD1
decay during SAC silencing, when compared with other vertebrates [41,43,46,73].
Noteworthy, recent works have revealed that many human, mouse and chicken
centromeres/kinetochores also have a compound, often bipartite, architecture
[59,61,74,75]. In particular, the same human kinetochore could be found with one
sub-domain end-on attached, and the other laterally attached to microtubules via
the fibrous corona[74]. This is consistent with our finding of non-uniform, highly
localized, MAD1 removal in response to local microtubule occupancy in Indian
muntjac kinetochores. Importantly, while in these cases MAD1 and local
microtubule density often negatively correlated along two apparently insulated
adjacent domains on the same large kinetochore of chromosome 3+X (simply
because these extreme cases are easier to detect), there were other cases
(particularly evident after Augmin perturbation) where this negative correlation
was observed along three or more sub-domains, without following a strict spatial
pattern (also evident in the distinct responses after laser microsurgery
experiments). Moreover, our finding that MAD1 is immobile within unattached
kinetochores indicates that it can only be removed upon the establishment of
highly-localized microtubule attachments (or eventually by a
microtubule-independent SAC-silencing mechanism, such as MPS1 or CDK1
inactivation). Overall, our data favors a model where anaphase onset is
determined by the gradual silencing of the SAC in response to highly localized
microtubule attachments along the entire kinetochore length, rather than a
uniform, switch-like mechanism. This model reconciles gradual SAC silencing
with increasing microtubule occupancy at kinetochores, without the need for a
dedicated microtubule "counting" system that triggers SAC inactivation. Most
relevant, our finding that microtubule occupancy at kinetochores is at the root

between timely SAC silencing and mitotic memory, identifies a critical
rate-limiting step with important implications for cell proliferation in mammals.

Materials and Methods

<u>Cell lines and culture conditions</u>

Indian muntjac cell lines were grown in Minimum Essential Media (MEM) (Corning), supplemented with 10% FBS (GIBCO, Life Technologies). Indian muntjac hTERT-immortalized fibroblasts were a gift from Jerry W. Shay [76]. IM fibroblasts stably expressing Venus-MAD1 were generated by lipofection of pVenus-MAD1 (kind gift from Jakob Nilsson [77]). RPE-1 hTERT-immortalized cells were cultured in DMEM (Corning™, 15323531) supplemented with 10% FBS (GIBCO, Life Technologies). All cells were kept at 37ºC in humidified conditions with 5% CO_2. Transfection was performed by incubating the cells with 1:400 Lipofectamine2000 (Invitrogen) in Opti-MEM (GIBCO, Life Technologies) for 6h, and stable and uniformly expressing cells were selected by FACS sorting (as detailed in [42]. For the production of Venus-MAD1 and mScarlet-CENP-A co-expressing cell line, Venus-MAD1 cells were transduced with pLVx-mScarlet-CENP-A lentiviral plasmid (generated in [43]). IM cells co-expressing 2x-GFP-CENP-A, GFP-CENTRIN-1 and mScarlet-MAD1 were produced by lentiviral transduction of pRRL-2x-EGFP-CENP-A (generated in [43]), pLVX-EGFP-C1-CENTRIN-1 (kind gift from Manuel Thery; Addgene plasmid # 73331) and pInducer-mScarlet-MAD1 (generated in this paper). Cells expressing GFP-H2B were produced by lentiviral transduction of LV-GFP (kind gift from Elaine Fuchs; Addgene plasmid # 25999). Lentiviral transduction was performed as detailed in [42]. Cells were incubated with lentivirus particles in the presence of 1:2000 Polybrene (Sigma-Aldrich; TR-1003) for 24h before supplied with fresh complete media. Stable lines with sufficient fluorescence intensity were selected by FACS sorting.

<u>Plasmid design</u>

pInducer-mScarlet-MAD1 was produced by cloning mScarlet (from pRRL-mScarlet-CENP-A) and MAD1 (from pVenus-MAD1) into the pInducer-20 backbone under a Tet-inducible promotor. To construct the empty backbone KASH-mCherry was removed by SalI + XhoI restriction digestion of pinducer 20

DN-KASH (kind gift from Daniel Conway; (Addgene plasmid # 125554)).

Restriction sites were amplified from the plasmid by PCR (fwd:

5'-TTAAAGGAACCAATTCAGGCTAGCACGCGTATATCTAGACCCAGCTTTCT-

3'; rev: 5'-TAAGCGTAGTCTGGGACG-3') and added to the emptied vector

through Gibson assembly. mScarlet and MAD1 were cloned into the NheI

linearized pInducer-20 backbone through Gibson assembly.

<u>siRNA Experiments</u>

Protein knockdown was performed as detailed in [42]. Briefly, IM fibroblasts seeded

at 70% confluency were starved with MEM supplemented with 5% FBS for 30

minutes before transfection. siRNA transfection was performed by adding

Lipofectamine RNAi Max (1:400; Invitrogen) and 50 nM of siRNA in serum

free-medium (Opti-MEM, Gibco) for 6h before replacing the solution with fresh

complete media. Cells were analyzed 72 hours after depletion. In long-term live

cell imaging experiments (see below), siRNAs were not replaced with fresh

medium and imaging started shortly after treatment. For the depletion of HAUS6,

the following sequence was used:

5'-GGUUGGUCCUAAGUUUAUU[dT][dT]-3'[39]. Cells mock transfected

(lipofectamine only) or transfected with luciferase-targeting siRNA were used as

controls.

<u>Drug Treatments</u>

For acute inhibition of Mps1 and CDK1 kinases, 4 µM MPS1-IN-1 (kind gift from

N.Gray [51]) and 10 µM RO 3306 (ChemCruz; CAS 872573-93-8), respectively,

were added shortly after Nuclear Envelop Breakdown (NEB). To arrest cells in

G2, these were treated with 10 µM RO 3306 for 14h. Microtubule

depolymerization was triggered using 1 or 3.3 µM of Nocodazole (Sigma-Aldrich;

CAS 31430-18-9) for 30 mins before the analysis. To increase mitotic duration

cells were arrested in prometaphase with 100 µL of Monastrol (Tocris Bioscience,

Cat. No.1305). Metaphase arrest was obtained using 5 µM MG132

(Sigma-Aldrich; CAS 133407-82-6). Fixed cell analysis using MG132 was

performed in the first 1.5 h after drug addition to avoid cohesion fatigue.
SiR-tubulin (Spirochrome; SC002) [78] was used to visualize microtubules, at 50
nM concentration incubated for 1 h prior to live-cell imaging. To induce p53
enrichment cells were treated with 5 µM etoposide (Selleck Chemicals, S1225) or
3.3 µM Nocodazole for 16h. The same volume of DMSO (Sigma-Aldrich, D4540)
was used as a control.

<u>Immunofluorescence</u>
Indian muntjac fibroblasts were seeded on fibronectin coated coverslips (#1.5
thickness) 24h before the experiment. Depending on the microscopy routine,
different fixation protocols were used: for widefield imaging cells were incubated
with 4% paraformaldehyde (Electron Microscopy Sciences) in PBS for 10 min at
room temperature; for STED microscopy PFA 4% supplemented with 0.1%-0.2 %
Glutaraldehyde (Electron Microscopy Sciences) in Cytoskeleten Buffer (CB -
274 mM NaCl, 10 mM KCl, 2.2 mM Na_2HPO_4, 0.8 mM KH_2PO_4, 4 mM EGTA,
4 mM $MgCl_2$, 10 mM Pipes, 10 mM glucose, pH 6.1) for 10 min at room
temperature was used; for same cell correlative live confocal and
super-resolution STED microscopy pre-warmed 2x concentrated PFA +
Glutaraldehyde diluted in imaging media to a final concentration of 4% and 0.2%,
respectively, was added to the imaging chamber. After a 5 min incubation at the
microscope (37ºC), the fixation solution was replaced with PFA 4%
supplemented with 0.2 % Glutaraldehyde in CB for an additional 5 min at room
temperature. Autofluorescence was quenched by a 0.1% sodium borohydride
solution (Sigma-Aldrich) for 7min. Cells were permeabilized with CB-0.5%Triton
for 20-30 min and blocked with CB-0.05% Tween 20 with 10% FBS (blocking
buffer) for 1 hour at RT. Samples were incubated with primary antibodies diluted
in blocking buffer over-night at 4ºC. The following primary antibodies were used:
human anti-centromere antiserum (ACA, Fitzgerald; 90C-CS1058) 1:2000/1:200
(Widefield/ STED); anti-tyrosinated tubulin (Bio-Rad; MCA77G) 1:2000/1:100
(Widefield/ STED); anti α-tubulin (clone B-5-1-2, Sigma-Aldrich; T5168) 1:100
(correlative live-cell and fixed STED microscopy). Cells were washed with
PBS-0.05% Tween before incubation with the corresponding secondary

antibodies for 1h at RT – Alexa Fluor 568 and 647 (Thermo Fisher Scientific) 1:1000; or abberior STAR 580 (Abberior Instruments; ST580) and abberior STAR-Red (Abberior Instruments; STRED) 1:200 for STED microscopy. DNA was labeled with a quick incubation in the presence of 1 µg/mL 4'6'-Diamidino-2-phenylindole (DAPI) in PBS-0.05% Tween. The samples were washed in PBS and mounted on glass slides with a mounting solution (20 mM Tris pH8, 0.5 N-propyl gallate, 90% glycerol).

<u>Monitoring kinetochore MAD1 levels in live recordings</u>

Indian muntjac fibroblasts stably expressing Venus-MAD1 or co-expressing Venus-MAD1 and mScarlet-CENP-A were plated on fibronectin coated glass bottom 35 mm FluoroDish (World Precision Instruments; FD35-100), 36-48h before imaging. Before imaging, normal culture media was replaced with Leibovitz's L15 medium (GIBCO, Life Technologies) supplemented with 10% FBS and the cell-permeable microtubule dye, SiR-tubulin. Live-cell imaging was performed on a temperature-controlled Nikon-Ti microscope equipped at the camera port with a Yokogawa CSU-X1 spinning-disc head with Borealis and an iXon+ DU-897 EM-CCD (Andor). Cells were imaged with an oil-immersion 60x 1.4 NA Plan-Apo DIC CFI objective (Nikon, lambda series), yielding a 176 nm/pixel sampling, or a 100x 1.4 NA Plan-Apo DIC CFI objective (Nikon, VC series), yielding a 106 nm/pixel. Images were acquired using 488 nm, 561nm and 647 nm (Coherent) laser lines and acquisition was controlled by NIS Elements AR software. To ensure that kinetochore could be tracked overtime with a reduced number of optical sections, cells were imaged under modest cell confinement. An image stack was acquired: every 6 seconds, 9 planes separated by 0.5 µm (Figures 1, 6 and Fig. S1C); 1 min, 11 planes separated by 0.5 µm, Fig. S1A, B); 10 seconds, 9 planes separated by 0.5 µm (Figure 3). Fluorescence intensity of kinetochore Venus-MAD1 was manually tracked throughout mitosis using the image analysis software Fiji 2.16.0 [79]. The recorded image stacks were sum projected and a region was drawn encircling the kinetochore and three different cytoplasmic background sites at each timepoint. MAD1 kinetochore fluorescence intensity (S_{KT}) was calculated by subtracting the averaged

cytoplasmatic signal (S_{BG}), represented in the following equation: $S_{KT, \text{corrected}}$ =

S_{KT} − $S_{BG, \text{averaged}} \times A_{KT}/A_{BG}$ S: raw integrated density; A: Area. The calculated

signal intensity was normalized to the followings: the clear last peak of the signal

decay curve in untreated/control, mock treated and siHAUS6 conditions; the first

frame after adding the inhibitor in MPS1i and CDK1i treatments; the average of

the first 10 timepoints in nocodazole treated cells. A single-phase exponential

decay function was fitted globally across curves polled from all kinetochores in

each experimental condition using the equation: $Y = (Y0 - \text{Plateau}) \times (\exp^{\wedge}(-Kx)) +$

Plateau, in GraphPad Prism 9 (Boston, Massachusetts USA). The half decay time

($t_{1/2}$) for each condition was derived from the fit according to $t_{1/2} = \ln(2)/K$. To test

for differences in the rate of MAD1 decay (K, derived from the global fit) in

untreated vs treated cells or large vs small kinetochores between conditions an

extra sum-of-squares F-test was used. Quantification of the signal intensity ratio

between kinetochore and cytoplasm (Fig. S1B), was performed manually using

Fiji 2.16.0. Three regions were drawn encircling the kinetochore (KT), the

cytoplasm (cyto) and an area outside of the cell (background signal, BG) at define

time points (anaphase onset, 10, 20, 30 and 40 minutes before anaphase). The

kinetochore and cytoplasmic fluorescence intensities were calculated by

subtracting background signal, represented in the following equations: $S_{KT, \text{corrected}}$

$= S_{KT}/A_{KT} − S_{BG}/A_{BG}$; $S_{\text{cyto, corrected}} = S_{\text{cyto}}/A_{\text{cyto}} − S_{BG}/A_{BG}$; S: signal; A: Area. The

ratio of the corrected kinetochore over cytoplasmatic signals was then calculated.

To observe how Mad1 signal spatial distribution within the kinetochore changed

throughout time kymographs were produced from live recordings of IM cells

(Figure 3). Image stacks were sum projected using Fiji 2.16.0 and kymographs

were generated with a custom-written routine in MATLAB (Mathworks, Natick,

MA) that compensates for spindle rotation and translations previously described

in [64].

<u>Cell confinement</u>

For cell confinement, we adopt a cell confiner as previously described [39, 62] using

a custom-designed polydimethylsiloxane (PDMS, RTV615, GE) layout to fit a

35mm diameter fluorodish. A suction cup was custom-made with a 10:1 mixture

(w/w PDMS crosslinker A/ B) and baked on an 80 ºC hot plate for 30mins and left

to dry over-night before unmolding. A PDMS confinement slide was then

prepared with a 10:1 mixture (w/w PDMS crosslinker A/ B) molded into

micropillars with a height of 8 μm (using a SU-8 wafer mold) polymerized onto a

10 mm round coverslip and baked for 95 °C for 15 min. The PDMS confinement

slide was attached to the PDMS suction cup and connected to a vacuum

generator (AF1-dual, Elveflow).

<u>Quantification of MAD1 intra-kinetochore mobility via fluorescence recovery after</u>

<u>photobleaching </u>(FRAP)

For FRAP experiments, IM cells stably expressing Venus-MAD1 were plated on

fibronectin coated 35mm FluoroDish (World Precision Instruments, FD35-100)

36-48 h before imaging. To elicit uniform and maximal kineotchore-MAD1

occupancy, cells were treated with nocodazole for 30 minutes before the

experiment. To ensure that the bleached kinetochore could be tracked overtime

with a reduced number of optical sections, an agar overlay was placed on top of

the cells decreasing the cell volume, as described in [63]. Briefly, a 170 μm thick

layer of 2% low-melting-point agarose in L15 medium supplemented with 10%

FBS was prepared in advance and incubated for 30 minutes with pre-warmed

L15 + 10% FBS containing nocodazole. Immediately before imaging, a small

piece of the agar layer was gently placed over the cells that were maintained in a

minimal amount of medium to prevent cell dehydration while ensuring

compression. FRAP experiments were conducted on a Leica Scanning Confocal

STELLARIS 8 FALCON (Leica Microsystems, Germany), equipped with an

incubation chamber and a power HyD S detector. Cells were imaged with a

water-immersion 86x 1.20 NA HC PL APO STED white objective (Leica

Microsystems, Germany), plus a 4.5x zoom, yielding an 86 nm/pixel sampling.

The 488-nm emission wavelength of a white Light Laser (WLL) was used to

bleach MAD1 signal from either part of or an entire kinetochore with 2 pulses of

0.3 seconds each. An image stack of 3 z-planes at a 0.5 μm interval was acquired

with the following routine: 3 pre-bleach frames, every 1.3 seconds; kinetochore

bleaching; post-bleach 5 frames, every 1.3 seconds for 5 frames, 10 frames,

every 2 seconds and then every 3 seconds for up to 55 seconds. The selected

representative images show sum-intensity projections of the image stacks.

Venus-MAD1 fluorescence at the kinetochore was manually tracked on the sum

projected image stacks using Fiji 2.16.0and a custom-designed script in Python

3.13. Four rectangular ROIS were defined corresponding to: (i) the bleached

region of the experiment kinetochore; (ii) the non-bleached region of the

experiment kinetochore; (iii) the unperturbed kinetochore from the pair (sister KT)

and (iv) a kinetochore from an unperturbed pair (control KT). Additionally, two

square ROIs with 0.86 μm sides were drawn over the cytoplasm to measure

background fluorescence (BG). A custom-designed script was used to draw the

kinetochore ROIs with a constant width of 0.6 μm, and length defined by the

distance of the kinetochore extremities in the pre-bleach frame (manually marked

based on MAD1 signal using Fiji 2.16.0). In the case of the partially ablated

kinetochore the length of the non-bleached region was defined at the first frame

post-bleaching, and the length of the bleached region was defined as the

difference between the total kinetochore length and the length of the

non-bleached region. The long axis of the rectangle was defined based on the

vector described by the points marking the extremities of the non-bleached signal

and was adjusted at every time point to track the kinetochore motion. MAD1

kinetochore fluorescence intensity ($S_{KT, corrected}$) was corrected for fluctuations in

the cytoplasmatic fluorescent signal as described in the following equation: $S_{KT,}$

$_{corrected}$ = S_{KT} / (a x time + b); S_{KT}: kinetochore MAD1 raw integrated density; a, b:

constants provided by a linear fit of the cytoplasmatic MAD1 raw integrated

density. The corrected kinetochore signal was then normalized to the average of

the 3 pre-bleach values ($S_{KT, normalized}$) and fit to a single exponential association

function: Y = Y0 + (Plateau − Y0) x (1-(exp^(-Kx))); Y0 value was constrained to

be the $S_{KT, corrected}$ at the first frame post-bleaching. Half recovery time ($t_{1/2}$) was

derived from the fit according to $t_{1/2}$ = ln(2)/K. Fluorescence recovery percentage

(Recovery) was calculated according to Recovery = (Plateau − Y0) / (1 − Y0).

Fitting and downstream statistical analysis was performed using GraphPad Prism

9 (Boston, Massachusetts USA). Cells where bleaching efficiency was < 50%;

focal plane changes were > 1 μm; or where the normalized MAD1 fluorescence

could not be fitted to a single exponential (if the 95 % confidence interval for any

of the parameters could not be determined) were excluded from the analysis.

Line intensity profiles showing Venus-MAD1 signal along the kinetochore were

measured from a ROI covering the entire kinetochore length and a width of 0.250

μm at selected times.

<u>Quantification of kinetochore MAD1 signal distribution with Stimulated Emission</u>

<u>Depletion (STED) Microscopy</u>

Indian muntjac fibroblasts stably expressing Venus-MAD1 were seeded on glass

coverslips and processed as described in the "Immunofluorescence" section. 3D

image stacks were acquired using an Abberior 'Expert Line' gated-STED

microscope, equipped with a Nikon Lambda Plan-Apo 1.4NA 60x objective lens.

CH-STED mode was implemented as described before [62]. Images were acquired

using excitation wavelengths at 488 nm, 561 nm, and 640 nm and a single

doughnut-shape depletion beam at 775 nm, a 0.8 Airy unit pinhole and the STED

channel had a time-gate threshold of 500 ps. Pixel size was set to 30 nm in xy

and 200 nm in z and Image stacks covering near full spindle volume (~8 μm). The

selected representative images show sum-intensity projections of a fraction of

the Z-slices (5-10). In images that show merged channels, histograms were

cropped to allow visualization of all structures. Venus-MAD1 and tubulin signal

distribution was analyzed along individual kinetochores in sum projected 3D

stacks (5-10 slices) using Fiji 2.16.0. A line of 0.6 μm width was drawn along the

longitudinal axis of the kinetochore (defined by the ACA signal). Intensity line

profiles were extracted along the kinetochore for both MAD1 and tubulin

channels and background subtracted (average cytoplasmatic intensity value).

Corrected signal intensity was normalized to the maximum intensity value within

each kinetochore. To visualize and measure correlations between MAD1 and

tubulin signals, we calculated the intensity z-scores, standardized internally (for

each kinetochore). Each position along the kinetochore is represented as a point

in a 2D space where the z-scores are the x-y axes and an ensemble of

kinetochores was either overlaid in the same scatter plot or displayed and

calculated for isolated kinetochores. The product zMAD1.ztubulin at each data

point is the local contribution to an overall correlation. Averaging over all data

points yields the covariance normalized to the standard deviation product, known

as the Pearson correlation coefficient. The Matlab function *corrcoef* was used,

which calculates Pearson's correlations and p-values with a two-tailed Student

t-test for zero correlation. 3D reconstructions of the image stacks were obtained

with Imaris 9.3.1 (Oxford Instruments).

<u>Laser microsurgery and correlative live-cell confocal and fixed-cell CH-STED</u>

<u>microscopy</u>

Indian muntjac cells expressing 2x-GFP-CENP-A and GFP-CENTRIN-1 and

mScarlet-MAD1 were seeded on fibronectin coated 25 mm, no. 1.5, round

coverslips (CAT 10593054, fisher scientific) 48h prior to the experiment. To

induce mScarlet-MAD1 expression, cells were incubated with 1 ug/mL of

doxycycline for 24h. Before imaging the coverslips were mounted on a 35 mm

Attofluor™ Cell Chamber (CAT A7816, Invitrogen) and the media replaced with in

Leibovitz's L15 medium (GIBCO, Life Technologies) supplemented with 10%

FBS. Cells were imaged using Nikon-Ti spinning-disk confocal microscope with a

100x 1.4 NA oil objective as detailed in the "Monitoring kinetochore MAD1 levels

in live recordings" section. Cells in metaphase were identified by having MAD1

negative kinetochores aligned at the equator and increased inter- kinetochore

and pole-pole distance (assessed by GFP-CENTRIN-1). After recording a

pre-surgery image stack, one of the large kinetochores (from the IM chromosome

3+X) was partially ablated with a 532 nm laser, controlled by a custom routine. A

detailed description of the microsurgery setup can be found in [80]. 2-5 consecutive

pulses with a 0.35 µm step and a 12 Hz repetition rate) were applied. The pulse

width was 10 ns and the pulse energy was 3.9–4.4 µJ. After surgery an image

stack (5 planes, 0.75 µm z-step) was recorded every minute until MAD1 signal

was detected at the kinetochore, for up to 30 min; after MAD1 detection the

acquisition routine was changed: 7 planes, 0.5 µm z-step every 20 seconds for

5 min. As a control for uniform MAD1 signal distribution, non-ablated cells were

imaged upon acute nocodazole treatment (5 to 60 min incubation). For

correlative live-cell confocal and fixed CH-STED microscopy analysis, once

MAD1 signal was detected at the kinetochore a fixation solution was added to the

imaging chamber. The location of the cell of interest was marked with a

laser-engraved pattern on the glass coverslip. The sample was processed for

staining as described in the "Immunofluorescence" section. After locating the

surgery cell with a 10x objective, a super-resolved image stack of the spindle was

acquired with CH-STED as described in the "Quantification of kinetochore MAD1

signal distribution with Stimulated Emission Depletion (STED) Microscopy"

section (pixel size 35 nm in xy and 150 nm in z). Representative images show a

sum (live-cell data) or max-intensity (fixed-cell data) projection of all z-slices that

contain the kinetochore of interest. Histograms were adjusted per frame to allow

better visualization of the representative phenotypes. To measure MAD1

recruitment after laser microsurgery, kinetochores were manually tracked using

the GFP-tagged CENP-A as a reference in Fiji 2.16.0. kinetochore deformation

was defined as the first time point when an increase in the angle between the line

connecting the ends of the surgery kinetochore and the line connecting the ends

of the unperturbed kinetochore from the pair (sister kinetochore) was observed.

MAD1 recruitment was defined as the first time point when mScarlet-MAD1

signal was visible at the sister kinetochore. MAD1 signal distribution along the

sister kinetochore was measured at the time point when signal intensity appeared

highest, based on visual assessment. On the cells where correlative live-cell

spinning-disk confocal and CH-STED microscopy was performed, tubulin signal

distribution was measured in the fixed sample. Measurements were performed

on sum projected stacks containing the kinetochore pair of interest. A segmented

line of 0.8 μm width (for MAD1, live imaging) or 0.9 μm width (for tubulin, fixed cell

imaging) was drawn to span the full length of the kinetochore, as defined by the

CENP-A signal, starting from the end that retained its counterpart (or randomly in

nocodazole treated cells). Intensity line profiles were extracted along the

kinetochore for all channels and background subtracted (cytoplasmatic signal

near the kinetochore). The corrected intensity profile was normalized to the
maximum within each kinetochore, as exemplified for CENP-A: $S_{\text{CENP-A, norm}}$ =
$(S_{\text{CENP-A}} - S_{\text{BG CENP-A}})$ / max $(S_{\text{CENP-A}} - S_{\text{BG CENP-A}})$, S: mean pixel intensity. To
correct for intensity variations caused by kinetochore deformation or movement,
intensity profiles are shown as a ratio of MAD1/tubulin over CENP-A. Kinetochore
length was normalized to a relative scale ranging from 0 (start of the line profile)
to 1 (end of the line profile). The average MAD1/CENP-A ratio intensity profile for
each condition (Nocodazole vs Surgery), was calculated using the 'Multiple
average curves' function with linear interpolation on OriginPro 2022 (OriginLab
Corporation, Northampton, MA, USA). The coefficient of variation of the
MAD1/CENP-A ratio was calculated as the standard deviation of the
MAD1/CENP-A intensity values along each kinetochore, divided by their mean.
Cells that failed to recruit MAD1 within the mean recruitment time + 1 s.d. (~12
minutes) or exhibited only very faint MAD1 signal were excluded from the MAD1
distribution analysis. Statistical analysis and plots were done using GraphPad
Prism 9 (Boston, Massachusetts USA).
798

<u>Quantification of kinetochore MAD1 levels in prometaphase and metaphase</u>
<u>HAUS6 depleted cells</u>
Control and HAUS6 depleted cells expressing Venus-Mad1 were treated with MG
132 for 1h and then fixed and stained for as detailed in the "Immunofluorescence
for wide-field" section. Samples were imaged on AxioImager Z1 equipped with a
CCD camera (ORCA-R2, Hamamatsu) operated by Zen software (Carl Zeiss,
Inc.). 3D image stacks covering the entire cell volume (0.23 µm z-step) were
acquired using a 100x Plan-Apochromatic oil differential interference contrast
objective lens, 1.46 NA (Carl Zeiss Microimaging Inc.) and excitation
wavelengths at 488 nm, 561 nm, and 640 nm. The number of cells with MAD1
positive kinetochores was visually assessed for each condition. Kinetochore
Venus-MAD1 signal intensity was measured on a single focal plane using Fiji
2.16.0. Cytoplasmatic Venus-MAD1 signal was averaged from 3 regions outside
the spindle area. The ration between kinetochore and cytoplasm signal was
calculated according to the equation: S_{KT} / $S_{\text{cytoplasm;}}$ S: mean pixel intensity.

Statistical analysis and plots were done using GraphPad Prism 9 (Boston,

Massachusetts USA).

<u>Cell tracking and daughter fate analysis in live recordings</u>

Indian muntjac fibroblasts stably expressing GFP-H2B were plated on fibronectin

coated µ-Slide 4 Well Glass Bottom (Ibidi GmbH, 80427) with Ham's F-10 media

supplemented with 20% FBS 24h before imaging. Cells were recorded using the

Zeiss LSM 980 Airyscan confocal microscope built on an Axio Observer 7

inverted stand with a motorized piezo stage and active focus stabilization. A

controlled atmosphere chamber (CO2 and temperature) was used. Objective

lens was a dry 0.3NA 10x Plan Neofluar. Detection was performed with two

GaAsP detectors. Image acquisition was carried out with ZEN 3.11 software. 3D

image stacks (pixel size 830 nm in xy, 4 planes separated by 6.7 µm, pinhole Airy

unit of 1.23, scan speed of 7, no averaging) were recorded every 7 min using both

transmitted light and an excitation wavelength at 488 nm. In experiments where

mitotic delay was induced through monastrol treatment, cells were imaged in the

presence of the drug for 6h before washing out the drug. The fate of cell families

was tracked for 72h. In experiments where HAUS6 was depleted, cells were

treated with control or HAUS6 siRNAs shortly before imaging. Cell families were

then tracked for 96h. Cells were tracked manually using TrackMate [81]. Single-cell

lineages and cell cycle durations (frame of anaphase mother cell to frame

anaphase daughter cell) were analysed by modifying previously published [82]

Jupyter notebooks (7.0.8). We defined mitotic duration as the frame of mitotic

onset (condensing chromatin) until the first frame of anaphase (clearly separated

daughter chromatin). For each division event, we determined mitotic defects and

errors, and the fate of the daughters. For downstream analysis, filtering and

plotting, all data were integrated using Jupyter notebooks (Python 3.12) and the

pandas (2.2.2) [83] and Altair (5.4.1) [84] libraries. Fiji/ImageJ [79] macros and

analytical code for the daughter fate analysis is available at

https://github.com/TobiasKletter/MitoticStop.

<u>Co-immunoprecipitation</u>

Indian muntjac fibroblasts stably expressing GFP-H2B were grown on with Ham's

F-10 media supplemented with 20% FBS. To obtain a mitotic arrested population

cells were treated for 16h with 3.3 µM Nocodazole. 10 million mitotic cells were

collected by shake-off. The remaining interphase cells (NOC interphase) and

cells left untreated (asynchronous) were collected by cell scrapping. Cells were

pelleted at 300 g for 5 minutes, washed with cold PBS and cell pellets were snap

frozen in liquid nitrogen. Co-immunoprecipitation assay was optimized from the

protocol described in [29]. Cell were resuspended in lysis buffer (20 mM Tris/HCl,

pH 7.5; 50 mM NaCl; 0.5% Triton X-100; 5 mM EGTA; 1 mM dithiothreitol; 2 mM

$MgCl_2$; Pierce Protease and Phosphatase inhibitor mini tablets, EDTA-free

[Thermo Fisher Scientific; A32961]) and incubated for 45 minutes at 4ºC with

agitation. Cell lysates were centrifuged at 15 000 rpm for 5 minutes at 4ºC to

collect the supernatant. Protein concentration was quantified and a lysate

solution containing 1.5 mg of protein in 1 mL of lysis buffer was prepared for each

condition. Lysates were incubated with 3 ug of anti-53BP1 antibody (Novus,

NB100-304) or control rabbit IgG for 2h at 4ºC with agitation and subsequently

with 20 µL of pre-washed (4x with lysis buffer) Pierce™ Protein A Magnetic Beads

(Thermo Fisher Scientific, 88845) for 1h at 4ºC with agitation. The non-binding

supernatant faction was collected and the beads were washed 4x with lysis

buffer. The bound fraction was eluted from the beads by resuspending in 2x

sample buffer (50 mM Tris-HCl, pH 6.8, 2% SDS, 10% glycerol, 1%

β-mercaptoethanol, 12.5 mM EDTA, and 0.02% bromophenol blue) and

denatured at 95 °C for 5 min. Samples were analyzed through western blot.

<u>Western blotting</u>

Protein samples of 13-20 ug in sample buffer were loaded on a 4–15%

Mini-PROTEAN® TGX™ Precast Protein Gel (Bio-Rad, 4561084), mounted on a

Mini-PROTEAN vertical electrophoresis apparatus (Bio-Rad). NZYColour Protein

Marker II (NZYTech) was used as a ruler. Blotting was performed with a

Transfer-Blot Turbo transfer system (Bio-Rad) onto 0.2 µm Nitrocellulose

membranes. Membranes were blocked in blocking buffer (5% powder milk in

PBS Tween 0.1%) for 1h. Anti-53BP1 rabbit 1:1000 (Novus, NB100-304),

anti-USP28 rabbit (Abcam, ab126604), anti-p53 DO-1 mouse (Santa Cruz

Biotechnology, SC-126) and anti-GAPDH mouse 1:10 000 (Proteintech,

60004-1-Ig) primary antibodies were diluted in blocking buffer and incubated 1 h

(4 °C) with agitation. Anti-rabbit HRP (Jackson ImmunoResearch, 111-035-003)

and anti-mouse light chain specific HRP (Jackson ImmunoResearch,

115-035-174) secondary antibodies were diluted 1:5000 in blocking buffer and

incubated 1 h (4 °C) with agitation. Membranes were developed with Clarity

Western ECL Blotting Substrate (Bio-Rad, 1705061) and detected in the

Chemidoc XRS system and Image Lab software (Bio-Rad).

Acknowledgements

We would like to thank all colleagues that generously provided reagents used in this study, and Paula Sampaio and Maria Azevedo from i3S Advanced Light Microscopy Facility for technical support. We are indebted to Franz Meitinger for critical discussions and technical advice on the mitotic stopwatch. We also thank CID Lab members for discussions and constructive feedback throughout the course of this project, with special thanks to Daniela Andrade and Rui Martins for their strong contribution during mitotic shake-offs. NO, JS-O and EW were respectively supported by fellowships SFRH/BPD/118126/2016, SFRH/BD/07730/2021 and 2022.13008.BD from Fundação para a Ciência e a Tecnologia of Portugal. This project was funded by The European Union through the European Research Council grant KAREVO (project 101140624) to HM.

Author contributions

Conceptualization: HM

Methodology: NO, JS-O, TK, AJP, EW

Investigation: NO, JS-O, TK

Visualization: NO, JS-O, TK, AJP

Funding acquisition: HM

Project administration: HM

Supervision: HM, AJP

Writing – original draft: NO, HM

Writing – review & editing: NO, JS-O, TK, AJP, EW, HM

Competing interests

Authors declare that they have no competing interests.

 Figure legends

Figure 1. SAC protein MAD1 is gradually removed from the kinetochore in a microtubule and mitotic kinases dependent manner.

(A) Representative live-cell recording images of Indian muntjac fibroblasts expressing Venus-MAD1 (orange) and with SiR-tubulin labeled microtubules (grey). Scale bar: 5μm.

Insets highlight an early congressing kinetochore pair (dashed square) and dashed circles show polar kinetochores/chromosomes. Scale bar: 5μm.

(A') Insets focusing on 1 half-spindle to highlight the congression of polar chromosomes (dashed circles) from A. Scale bar: 5μm.

(A'') Kymographs depicting Venus-MAD1 signal of the 2 polar chromosomes shown in A'. The closest pole position is shown by the dashed line. Leading (black asterisk) and trailing kinetochore (white asterisk) and indicated for each pair. Scale bar: 5μm.

(B) Representative live-cell recording images of Indian muntjac fibroblasts acutely treated with Nocodazole, MPS1 and CDK1 inhibitors. Venus-MAD1 (orange) and SiR-tubulin labeled microtubules (grey). Insets highlight a kinetochore pair. Scale bar: 5μm.

(C) Normalized Venus-MAD1 fluorescence decay IM cells untreated (control) or treated with Nocodazole, MPS1 and CDK1 inhibitors. Data represented as mean ± 95% confidence interval (CI; Control n = 53 kinetochores, 7 cells, green; Nocodazole n = 31 kinetochores, 8 cells, grey; MPS1i n = 63 kinetochores, 12 cells, yellow; CDK1i n = 18 kinetochores, 10 cells, blue; data pooled from at least five independent experiments per condition).

(D) Normalized Venus-MAD1 fluorescence decay in large or small kinetochores from untreated cells. Data are represented as mean ± 95% CI. (Large KTs n = 19 kinetochores, 7 cells, 5 independent experiments, dark green; small KTs n = 34 kinetochores, 5 cells, 4 independent experiments, light green).

(E) Summary table of the parameters extracted from a single exponential fitting of Venus-MAD1 fluorescence decay across the condition shown in C and D. P-values were calculated using extra sum-of-squares F-test comparing the rate

of MAD1 decay (K, derived from global fit) in untreated vs treated cells or large

vs small kinetochores.

Figure 2. MAD1 is immobile within kinetochores

(A) Schematic representation of Dynein-mediated MAD1/MAD2 complex

removal from kinetochores.

(A') Predicted curves of MAD1 fluorescence recovery after partial kinetochore

bleaching assuming MAD1 to be mobile (left) or immobile (right) within the

kinetochore. Lines represent the fluorescence levels from the region of the

kinetochore that was bleached (light blue) or left unperturbed (dark blue).

(B), (C) Representative images of Indian muntjac cells pre- and post-bleach of

part (B) or an entire kinetochore (C). Insets: kinetochore pair that was perturbed.

Dashed line boxes indicate the ablated kinetochore region (partial bleach KT:

non-bleached region - dark blue; bleached region – blue; total bleach KT -

yellow).

(B'), (C') Venus-Mad1 signal intensity profile along the kinetochore at pre-bleach

and at selected times post-bleach for kinetochores shown in B and C,

respectively.

(B''), (C'') Normalized Venus-Mad1 fluorescence recovery curves for cells shown

in B and C, respectively (Dark blue line non-bleached region; light blue bleached

region; yellow line totally bleached kinetochore; black and dark grey lines

unperturbed kinetochores from the bleach or a control kinetochore pair,

respectively; light grey cytoplasm; from fit curves are shown by the

semi-translucid line).

(D) Kinetochore Venus-MAD1 fluorescence recovery after partial or total

kinetochore bleach (Partial KT bleach n = 7; total KT bleach n = 8 from at least

two independent experiments). Dots show individual kinetochores. P = 0.2319,

Mann-Whitney test (two-tailed).

(E) Half recovery times of kinetochore Venus-MAD1 fluorescence after partial

or total kinetochore bleach. Dots show individual kinetochores. P = 0.5358,

Mann-Whitney test (two-tailed).

Figure 3. MAD1 removal along an individual kinetochore is not uniform

(A) Live-cell recording images of Indian muntjac fibroblasts stably co-expressing

Venus-MAD1 (orange) and mScarlet-CENP-A (blue) with SiR-tubulin labeled

microtubules (grey). Inset: kinetochore pair that shows non-uniform loss of

MAD1 signal. Dashed arrow indicates the kinetochore depicted in B. Scale bar:

5µm.

(B) Kymographs depicting Venus-MAD1 and mScarlet-CENP-A signal

distribution along the longitudinal axis of the kinetochore shown in A. Each

channel is shown on a greyscale and as a heatmap. Scale bar: 1µm

Figure 4. MAD1 removal along individual kinetochores correlates with local

microtubule density

(A) Representative CH-STED image of an Indian muntjac fibroblasts with

partially attached kinetochores. Microtubules (grey; STED), centromeres (ACA;

blue; STED) and Venus-MAD1 (orange; confocal). Insets: partially attached

kinetochore. Scale bar: 2µm.

(B) Three-dimensional reconstruction of the cell shown in A (different

perspective). Inset: partially attached kinetochore. The kinetochore pair of

interest is traced by a dashed line. Microtubules (grey), centromeres (ACA; blue)

and Venus-MAD1 (orange). Scale bars, Left: 3µm; Right: 1µm.

(C) Max normalized Venus-Mad1 (orange) and tubulin (microtubules; grey)

signal intensity profile along the longitudinal axis of the kinetochore highlighted

in A (arrowhead).

(D) Pearson correlation coefficient between MAD1 and microtubules signal

intensity along the kinetochore. Bars show individual partially attached

kinetochores (n=21 kinetochores, 7 independent experiments).

E) Representative distribution of kinetochore sub-regions showing positive,

negative and no correlation between MAD1 (y-axis) and microtubules (x-axis)

intensity z-scores. Dots represent distinct positions along the "kinetochore"

shown below the graphs. NE and SW quadrants contribute positively, while NW

and SE quadrants contribute negatively to the Pearson correlation coefficient (r).

(F) Correlation map of MAD1 and microtubule intensity z-scores for the

kinetochores shown in D. The underlying scatterplot contains data from multiple
kinetochores, with each dot representing a sub-kinetochore region. For
visualization, the scatterplot was converted into a regional density
map. P-value was calculated with Student t-test for zero correlation (two-sided).
1018

Figure 5. Local perturbation of kinetochore structure results in a localized
SAC response that inversely correlates with microtubule occupancy

(A) Schematic summary of the kinetochore response to partial laser
microsurgery ablation.
(B) Representative live-recording images of Indian muntjac cells expressing
2x-GFP-CENP-A and GFP-CENTRIN-1 (both in blue) and mScarlet-MAD1
(orange/ grey scale) upon partial kinetochore surgery or acute Nocodazole
treatment. Insets: kinetochore pair that was perturbed. Arrows point to the
ablated kinetochore region, pre- and post-surgery (solid and hollow arrow,
respectively). Arrowheads show MAD1 recruitment. Scale bar: 2µm.
(B') Max normalized line intensity profile plot of MAD1/CENP-A signal ratio along
the longitudinal axis of the kinetochores highlighted in A (surgery cell in orange;
nocodazole cell in grey).
(C) Dot plot showing the time from surgery to observable deformation (bending;
n= 12 kinetochores) and MAD1 signal re-appearance (n = 14 kinetochores) at
the sister kinetochore in cells where SAC recruitment was observed. Each dot
represents an individual kinetochore from a different cell, recorded during 13
independent experiments. P <0.0015, Wilcoxon matched-pairs signed rank test.
(D) Max normalized MAD1/CENP-A ratio intensity profile along the longitudinal
axis of kinetochores after partial surgery of their sisters (orange) or upon acute
Nocodazole treatment (NOC; grey). Thin lines show individual kinetochores and
thick lines show average curves (surgery n = 11 kinetochores/cells, 10
independent experiments; Nocodazole n = 19 kinetochores/cells, 2 independent
experiments). The graph depicts a cohort of cells that recruit MAD1 to the
proximal tip of the kinetochore, relative to the surgery region (n = 11/16 MAD1
recruiting kinetochores).
(E) Coefficient of variation of MAD1/CENP-A signal ratio along kinetochores

after partial surgery of their sisters. Dots represent individual kinetochores pooled from all cells where MAD1 recruitment was observed (surgery n = 16 cells, 14 independent experiments; Nocodazole n = 19 cells, 2 independent experiments). P <0.0001, Mann-Whitney test.

(F), (F') Representative images from correlative live cell spinning disk (F) and fixed CH-STED microscopy (F'). Indian muntjac cells expressing 2x-GFP-CENP-A and GFP-CENTRIN-1 (both in blue) and mScarlet-MAD1 (orange/grey scale) were fixed after partial kinetochore surgery and stained to visualize microtubules (grey). Insets: kinetochore pair that was perturbed. Arrows point to the ablated kinetochore region, pre- and post-surgery (solid and hollow arrow, respectively). Arrowheads show MAD1 recruitment. Scale bar: 2µm.

(F'') Max normalized line intensity profile plot of MAD1/CENP-A (live cell data just before fixation, orange) and microtubules/CENP-A (fixed cell data, grey) signal ratio along the longitudinal axis of the kinetochore highlighted in F and F'.

Figure 6. Partial microtubule occupancy at kinetochores after Augmin depletion delays MAD1 removal and SAC silencing

(A) Representative live-cell recording images of Indian muntjac fibroblasts mock or HAUS6 siRNA treated. Venus-MAD1 (orange) and SiR-tubulin labeled microtubules (grey). Scale bar: 5µm. Insets highlight a kinetochore pair. Asterisks indicate the two kinetochores from the pair. Scale bar: 5µm.

(B) Normalized Venus-MAD1 fluorescence decay from A. Data from at least five independent experiments are represented as mean ± 95% confidence interval (Mock n = 48 kinetochores, 10 cells, grey; siHAUS6 n = 47 kinetochores, 10 cells, blue).

(C) Summary table of the parameters extracted from a single exponential fitting of Venus-MAD1 fluorescence decay across the conditions shown in B. P-values were calculated using extra sum-of-squares *F*-test comparing the rate of MAD1 decay (K, derived from global fit) in mock vs siHAUS6 treated cells.

(D) Percentage of cells with at least one MAD1 positive kinetochore in mock (grey) or siHAUS6 (blue) treated cells after 1h of MG132 arrest. Bars show

mean, error bars show s.d. from three independent experiments (n≥293 cells per

condition). P <0.0001, prometaphase; P=0.0008, metaphase; unpaired t test

(two-sided).

(E) Dot plot showing the ratio between kinetochore and cytoplasmatic

Venus-MAD1 signal intensity in mock (grey) or siHAUS6 (blue) treated cells

fixed after 1h of MG132 arrest. Dots show individual kinetochores, bars show

mean, error bars show s.d. from three independent experiments (n≥309

kinetochores, ≥ 17 cells per condition). P<0.0001, Mann-Whitney test

(two-sided).

(F) Representative CH-STED image of an Indian muntjac control or HAUS6

depleted cells treated with MG132 for 1h. Microtubules (grey; STED),

centromeres (ACA; blue; STED) and Venus-MAD1 (orange; confocal). Inset: full

(control) or partially (siHAUS6) attached kinetochore pair. Scale bar: 2µm.

1091

Figure 7. Delayed SAC silencing due to partial microtubule occupancy at

kinetochores activates the mitotic stopwatch to stop cell proliferation

(A) Schematic summary of the experimental setup and downstream analysis.

(A') Mitotic duration of Indian muntjac cells expressing H2B-GFP with decreasing

HAUS6 levels (time in the presence of control or HAUS6 siRNA; grey and blue,

respectively). Data pooled from three independent experiments (siHAUS6 n =

370, siCtrl n = 335). Circles denote mean in each time bin (bin size = 12h),

error bars show the s.e.m.

(B) Bar graph showing the percentage of Indian muntjac cells that exited mitosis

with a particular mitotic defect/ error at decreasing HAUS6 levels. Data pooled

from three independent experiments (n = 469 cells). Bars show mean in each

time bin (bin size = 6h or 24h), error bars show the s.e.m.

(C) Plot of the mitotic cell fate as a function of mitotic duration in Indian muntjac

cells treated with siHAUS6 for 0 to 96h. Bars show individual cells (n = 469 cells

pooled from three independent experiments).

(D) Plot of daughter cells fate (Gen 2) as a function of mother cell mitotic

duration (Gen 1). Only mother cells from the selected cohort (12-36h of siHAUS6

treatment) are shown. Bars show individual daughter cells (n = 272 cells pooled

from three independent experiments).

(E) Bar graph showing the percentage of mother cells from the selected cohort

that has 0, 1 or 2 daughter cells dividing at least once, as a function of mother

cells time in mitosis. Only mother cells from the selected cohort (12-36h of

siHAUS6 treatment) and that completed mitosis with no defects are shown. Data

are binned (< 30 min n = 1 cell, 30-60 min n = 25 cells, 60-90 min n = 33 cells,

90-120 min n = 27 cell, > 120 min n = 10 cells pooled from three independent

experiments).

(G) Analysis of 53BP1 immunoprecipitates from Indian muntjac fibroblast

populations treated with nocodazole (3.3 µM) for 16 h. Cells were separated into

interphase and mitotic cells via mechanical shake-off. As negative control, rabbit

IgG were used. GAPDH as loading control.

(H) Immunoblot on whole-cell lysates from human RPE1 hTERT cells and Indian

muntjac fibroblasts (GFP-H2B), that were treated for 16h either with DMSO,

etoposide (5 µM), or nocodazole (3.3 µM, mitotic shake-off) and probed for p53

enrichment. GAPDH as loading control.

1126

Figure 8 - Microtubule occupancy at kinetochores integrates with the

mitotic stopwatch to ensure faithful daughter cell proliferation

(Top) Augmin-mediated K-fiber maturation promotes maximal microtubule

occupancy at kinetochores through gradual and highly localized MAD1/MAD2

removal, resulting in timely SAC silencing. In this way, mitotic timing is optimal,

fidelity is maintained, and daughter cells continue proliferating.

(Bottom) When K-fiber maturation is impaired, for example through disruption of

Augmin-mediated microtubule amplification, microtubule occupancy at

kinetochores remains partial and SAC silencing is delayed. Eventually, the cell

divides either after achieving full occupancy or while kinetochores are still only

partially occupied. The resulting mitotic delay is registered by the mitotic

stopwatch, which bookmarks daughter cells to halt proliferation.

1139

 Supplemental Figure legends

 Figure S1. MAD1 is undetectable at kinetochores 20 minutes before

 anaphase

 (A) Representative live-cell recording images of Indian muntjac fibroblasts

 expressing Venus-MAD1 (orange, upper panels or grey, lower panels) and

 mScarlet-CENP-A with SiR-tubulin labeled microtubules (grey, upper panels).

 Scale bar: 5µm.

 (B) Dot plot showing the ratio between kinetochore and cytoplasmatic

 Venus-MAD1 signal intensity at selected time points. Dots show individual

 kinetochores, lines show mean ± s.d. (-40 min: 87 kinetochores, $p < 0.0001$; -30

 min: 123 kinetochores, $p < 0.0001$; -20 min: 137 kinetochores, $p > 0.9999$; -10

 min: 126 kinetochores, $p > 0.9444$; Anaphase onset: 126 kinetochores; 15 cells

 from 7 independent experiments)., P value calculated with Kruskal-Wallis test

 with Dunn's comparisons of mean fluorescence in each time point with

 anaphase onset.

 (C) Normalized signal intensity of cytoplasmatic Venus-MAD1 throughout live

 cell recording. Lines show selected cytoplasmatic regions (n= 27 spots, 9 cells, 6

 independent experiments).

 Figure S2. MAD1 tracking on individual kinetochores throughout mitosis

 Single kinetochore tracks of Venus-MAD1 fluorescence decay shown in Figure 1

 and 6. Thin lines represent individual kinetochores (Control n = 53 kinetochores,

 7 cells, green; Control Large KTs n = 19 kinetochores, 7 cells, dark green;

 Control small KTs n = 34 kinetochores, 5 cells, light green; Nocodazole n = 31

 kinetochores, 8 cells, grey; MPS1i n = 63 kinetochores, 12 cells, yellow; CDK1i n

 = 18 kinetochores, 10 cells, blue; Mock n = 48 kinetochores, 11 cells, light grey;

 siHAUS6 n = 47 kinetochores, 10 cells, purple blue; data pooled from at least

 four independent experiments per condition). Thick lines show the single-phase

 exponential decay fit curve for all pooled data (except for Nocodazole: average

 cure curve).

Figure S3. Additional examples of partially attached kinetochores

(A), (A) Representative CH-STED image of Indian muntjac fibroblasts with partially attached kinetochores. Microtubules (grey; STED), centromeres (ACA; blue; STED) and Venus-MAD1 (orange; confocal). Insets: partially attached kinetochores. Scale bar: 2μm.

(A'), (B') Max normalized Venus-Mad1 (orange) and tubulin (microtubules; grey) signal intensity profile along the longitudinal axis of the kinetochores highlighted in A and B, respectively (arrow head).

(C) Correlation map of MAD1 and microtubule intensity z-scores for the all partially attached kinetochores analyzed in untreated cells. Each scatterplot contains data from a single kinetochore, with each dot representing a sub-kinetochore region. P-value was calculated with Student t-test for zero correlation (two-sided).

Figure S4. Additional examples of localized MAD1 recruitment upon kinetochore laser ablation

(A) Representative live-recording images of Indian muntjac cells expressing 2x-GFP-CENP-A and GFP-CENTRIN-1 (both in blue) and mScarlet-MAD1 (orange/ grey scale) upon partial kinetochore surgery. Insets: kinetochore pair that was perturbed. Arrows point to the ablated kinetochore region, pre- and post-surgery (solid and hollow arrow, respectively). Arrowhead shows MAD1 recruitment. Scale bar: 2μm.

(A') Max normalized line intensity profile plot of MAD1/CENP-A signal ratio along the longitudinal axis of the kinetochore highlighted in A.

(B) Max normalized MAD1/CENP-A ratio intensity profile along the longitudinal axis of kinetochores after partial surgery of their sisters (orange) or upon acute Nocodazole treatment (NOC; grey). Thin lines show individual kinetochores and thick lines show average curves (surgery n = 5 kinetochores/cells, 5 independent experiments; Nocodazole n = 19 kinetochores/cells, 2 independent experiments). The graph depicts a cohort of cells that recruit MAD1 to a distal position relative to the surgery region (n = 5/16 MAD1 recruiting kinetochores). Data from Nocodazole treated cells is the same as shown in Figure 5.

(C), (C') Representative images from correlative live cell spinning disk (C) and

fixed CH-STED microscopy (C') in a cell that recruited MAD1 to distal position

relative to the surgery region. Indian muntjac cells expressing 2x-GFP-CENP-A

and GFP-CENTRIN-1 (both in blue) and mScarlet-MAD1 (orange/grey scale)

were fixed after partial kinetochore surgery and stained to visualize microtubules

(grey). Insets: kinetochore pair that was perturbed. Arrows point to the ablated

kinetochore region, pre- and post-surgery (solid and hollow arrow, respectively).

Arrowheads show MAD1 recruitment. Scale bar: 2µm.

(C'') Max normalized line intensity profile plot of MAD1/CENP-A (live cell data

just before fixation, orange) and microtubules/CENP-A (fixed cell data, grey)

signal ratio along the longitudinal axis of the kinetochore highlighted in C and C'.

1215

Figure S5. Uniform perturbation of microtubule occupancy upon Augmin

depletion

(A) Representative CH-STED image of an Indian muntjac fibroblasts with

partially attached kinetochores upon siHAUS6 treatment. Microtubules (grey;

STED), centromeres (ACA; blue; STED) and Venus-MAD1 (orange; confocal).

Insets: partially attached kinetochore. Scale bar: 2µm.

(B) Three-dimensional reconstruction of the cell shown in A. Inset: partially

attached kinetochore. The kinetochore pair of interest is traced by a dashed line.

Microtubules (grey), centromeres (ACA; blue) and Venus-MAD1 (orange). Scale

bars, Left: 3µm; Right: 1µm.

(C) Max normalized Venus-Mad1 (orange) and tubulin (microtubules; grey)

signal intensity profile along the longitudinal axis of the kinetochore highlighted

in A (arrow head).

(D) Pearson correlation coefficient between MAD1 and microtubules signal

intensity along the kinetochore. Bars show individual partially attached

kinetochores in siHAUS6 treated cells (n=13 kinetochores, 9 cells, 2

independent experiments).

(E) Correlation map of MAD1 and microtubule intensity z-scores for kinetochores

in control (data from Figure 3) and in siHAUS6 treated cells (shown in D). The

underlying scatterplot contains data from multiple kinetochores, with each dot

representing a sub-kinetochore region. For visualization, the scatterplot was converted into a regional density map. P-value was calculated with Student t-test for zero correlation (two-sided).

Figure S6. Additional examples of partially attached kinetochores in Augmin depleted cells

(A), (A) Representative CH-STED image of Indian muntjac fibroblasts siHAUS6 treated with partially attached kinetochores. Microtubules (grey; STED), centromeres (ACA; blue; STED) and Venus-MAD1 (orange; confocal). Insets: partially attached kinetochores. Scale bar: 2µm.

(A'), (B') Max normalized Venus-Mad1 (orange) and tubulin (microtubules; grey) signal intensity profile along the longitudinal axis of the kinetochores highlighted in A and B, respectively (arrowhead).

(C) Correlation map of MAD1 and microtubule intensity z-scores for the all partially attached kinetochores analyzed in siHAUS6 cells. Each scatterplot contains data from a single kinetochore, with each dot representing a sub-kinetochore region. P-value was calculated with Student t-test for zero correlation (two-sided).

Figure S7. Extended mitosis activates the mitotic stopwatch to stop cell proliferation in Indian muntjac cells

(A) Schematic summary of the experimental setup and downstream analysis.

(B) Daughter cell cycle duration as a function of mother mitotic duration in Indian muntjac cells expressing H2B-GFP upon monastrol-induced arrest and release. Bars show individual daughter cells (n = 305 cells pooled from two independent experiments). Note that only daughter cells that entered mitosis during imaging were included in the graph (see methods section).

(C) Mitotic duration of Indian muntjac cells that enter mitosis within a given recording time bin. Data pooled from two independent experiments (n = 489). Circles denote mean in each time bin (bin size□=□6h), error bars show the s.e.m.

(D) Bar graph showing the percentage of Indian muntjac cells that exited mitosis

with no defects or a particular mitotic defect/ error after distinct mitotic durations

(bin size = 30 min). Only mother cells from the selected cohort (0-18h of

imaging) are shown. Data pooled from two independent experiments (n = 142

cells). Bars show mean in each time bin (bin size = 6h or 24h), error bars show

the s.e.m.

(E) Daughter cells fate (Gen 2) as a function of mother cell mitotic duration (Gen

1). Only mother cells from the selected cohort (0-18h of imaging) and that

completed mitosis with no defects are shown are shown. Bars show individual

cells (n = 123 cells pooled from two independent experiments).

(E') Percentages of cell families dividing only once (Gen 1 mothers only) versus

a second (Gen 2 daughters) or a third time (Gen 3 daughters), after binning the

cell families by mitotic duration of the Gen 1 mother. Circles show the two

independent experiments, bars show the mean and errors show the s.e.m.

(F) Summary of the percentage of daughter cells that arrest as a function of

mother cell mitotic duration and cause of mitotic delay.

References

1. Bakhoum, S.F., and Cantley, L.C. (2018). The Multifaceted Role of Chromosomal Instability in Cancer and Its Microenvironment. Cell *174*, 1347-1360. 10.1016/j.cell.2018.08.027.
2. Rieder, C.L., Cole, R.W., Khodjakov, A., and Sluder, G. (1995). The checkpoint delaying anaphase in response to chromosome monoorientation is mediated by an inhibitory signal produced by unattached kinetochores. The Journal of cell biology *130*, 941-948.
3. Musacchio, A., and Salmon, E.D. (2007). The spindle-assembly checkpoint in space and time. Nature reviews *8*, 379-393. nrm2163 [pii] 10.1038/nrm2163.
4. Lara-Gonzalez, P., Pines, J., and Desai, A. (2021). Spindle assembly checkpoint activation and silencing at kinetochores. Semin Cell Dev Biol *117*, 86-98. 10.1016/j.semcdb.2021.06.009.
5. McAinsh, A.D., and Kops, G. (2023). Principles and dynamics of spindle assembly checkpoint signalling. Nature reviews *24*, 543-559. 10.1038/s41580-023-00593-z.
6. Vazquez-Novelle, M.D., Sansregret, L., Dick, A.E., Smith, C.A., McAinsh, A.D., Gerlich, D.W., and Petronczki, M. (2014). Cdk1 inactivation terminates mitotic checkpoint surveillance and stabilizes kinetochore attachments in anaphase. Curr Biol *24*, 638-645. 10.1016/j.cub.2014.01.034.
7. Rattani, A., Vinod, P.K., Godwin, J., Tachibana-Konwalski, K., Wolna, M., Malumbres, M., Novak, B., and Nasmyth, K. (2014). Dependency of the spindle assembly checkpoint on Cdk1 renders the anaphase transition irreversible. Curr Biol *24*, 630-637. 10.1016/j.cub.2014.01.033.
8. Collin, P., Nashchekina, O., Walker, R., and Pines, J. (2013). The spindle assembly checkpoint works like a rheostat rather than a toggle switch. Nature cell biology *15*, 1378-1385. 10.1038/ncb2855 ncb2855 [pii].
9. Dick, A.E., and Gerlich, D.W. (2013). Kinetic framework of spindle assembly checkpoint signalling. Nature cell biology *15*, 1370-1377. 10.1038/ncb2842 ncb2842 [pii].
10. Rosenberg, J.S., Cross, F.R., and Funabiki, H. (2011). KNL1/Spc105 recruits PP1 to silence the spindle assembly checkpoint. Curr Biol *21*, 942-947. 10.1016/j.cub.2011.04.011.
11. Moura, M., Osswald, M., Leca, N., Barbosa, J., Pereira, A.J., Maiato, H., Sunkel, C.E., and Conde, C. (2017). Protein Phosphatase 1 inactivates Mps1 to ensure efficient Spindle Assembly Checkpoint silencing. Elife *6*, e25366. 10.7554/eLife.25366.
12. Grallert, A., Boke, E., Hagting, A., Hodgson, B., Connolly, Y., Griffiths, J.R., Smith, D.L., Pines, J., and Hagan, I.M. (2015). A PP1-PP2A phosphatase relay controls mitotic progression. Nature *517*, 94-98. 10.1038/nature14019.
13. Hayward, D., Alfonso-Pérez, T., Cundell, M.J., Hopkins, M., Holder, J.,

Bancroft, J., Hutter, L.H., Novak, B., Barr, F.A., and Gruneberg, U. (2019).
CDK1-CCNB1 creates a spindle checkpoint-permissive state by enabling
MPS1 kinetochore localization. The Journal of cell biology *218*,
1182-1199. 10.1083/jcb.201808014.

14.   Alfonso-Pérez, T., Hayward, D., Holder, J., Gruneberg, U., and Barr, F.A.
(2019). MAD1-dependent recruitment of CDK1-CCNB1 to kinetochores
promotes spindle checkpoint signaling. The Journal of cell biology *218*,
1108-1117. 10.1083/jcb.201808015.

15.   Allan, L.A., Camacho Reis, M., Ciossani, G., Huis In 't Veld, P.J.,
Wohlgemuth, S., Kops, G.J., Musacchio, A., and Saurin, A.T. (2020).
Cyclin B1 scaffolds MAD1 at the kinetochore corona to activate the
mitotic checkpoint. The EMBO journal *39*, e103180.
10.15252/embj.2019103180.

16.   Hiruma, Y., Sacristan, C., Pachis, S.T., Adamopoulos, A., Kuijt, T., Ubbink,
M., von Castelmur, E., Perrakis, A., and Kops, G.J. (2015). CELL
DIVISION CYCLE. Competition between MPS1 and microtubules at
kinetochores regulates spindle checkpoint signaling. Science (New York,
N.Y *348*, 1264-1267. 10.1126/science.aaa4055.

17.   Ji, Z., Gao, H., and Yu, H. (2015). CELL DIVISION CYCLE. Kinetochore
attachment sensed by competitive Mps1 and microtubule binding to
Ndc80C. Science (New York, N.Y *348*, 1260-1264.
10.1126/science.aaa4029.

18.   Howell, B.J., McEwen, B.F., Canman, J.C., Hoffman, D.B., Farrar, E.M.,
Rieder, C.L., and Salmon, E.D. (2001). Cytoplasmic dynein/dynactin
drives kinetochore protein transport to the spindle poles and has a role in
mitotic spindle checkpoint inactivation. The Journal of cell biology *155*,
1159-1172.

19.   Wojcik, E., Basto, R., Serr, M., Scaerou, F., Karess, R., and Hays, T.
(2001). Kinetochore dynein: its dynamics and role in the transport of the
Rough deal checkpoint protein. Nature cell biology *3*, 1001-1007.

20.   Gassmann, R., Holland, A.J., Varma, D., Wan, X., Civril, F., Cleveland,
D.W., Oegema, K., Salmon, E.D., and Desai, A. (2010). Removal of
Spindly from microtubule-attached kinetochores controls spindle
checkpoint silencing in human cells. Genes & development *24*, 957-971.
10.1101/gad.1886810.

21.   Ide, A.H., DeLuca, K.F., Wiggan, O., Markus, S.M., and DeLuca, J.G.
(2023). The role of kinetochore dynein in checkpoint silencing is restricted
to disassembly of the corona. Molecular biology of the cell *34*, ar76.
10.1091/mbc.E23-04-0130.

22.   Barisic, M., Sohm, B., Mikolcevic, P., Wandke, C., Rauch, V., Ringer, T.,
Hess, M., Bonn, G., and Geley, S. (2010). Spindly/CCDC99 is required for
efficient chromosome congression and mitotic checkpoint regulation.
Molecular biology of the cell *21*, 1968-1981. 10.1091/mbc.e09-04-0356.

23.   McEwen, B.F., Heagle, A.B., Cassels, G.O., Buttle, K.F., and Rieder, C.L.
(1997). Kinetochore fiber maturation in PtK1 cells and its implications for
the mechanisms of chromosome congression and anaphase onset. J.
Cell Biol. *137*, 1567-1580. 10.1083/jcb.137.7.1567.

24. Nicklas, R.B., Waters, J.C., Salmon, E.D., and Ward, S.C. (2001).
Checkpoint signals in grasshopper meiosis are sensitive to microtubule
attachment, but tension is still essential. Journal of cell science *114*,
4173-4183. 10.1242/jcs.114.23.4173.
25. Kuhn, J., and Dumont, S. (2019). Mammalian kinetochores count
attached microtubules in a sensitive and switch-like manner. The Journal
of cell biology. 10.1083/jcb.201902105.
26. Etemad, B., Vertesy, A., Kuijt, T.E.F, Sacristan, C., van Oudenaarden, A.,
and Kops, G. (2019). Spindle checkpoint silencing at kinetochores with
submaximal microtubule occupancy. Journal of cell science *132*.
10.1242/jcs.231589.
27. Dudka, D., Noatynska, A., Smith, C.A., Liaudet, N., McAinsh, A.D., and
Meraldi, P. (2018). Complete microtubule-kinetochore occupancy favours
the segregation of merotelic attachments. Nat Commun *9*, 2042.
10.1038/s41467-018-04427-x.
28. Pereira, A.J., and Maiato, H. (2012). Maturation of the
kinetochore-microtubule interface and the meaning of metaphase.
Chromosome Res *20*, 563-577. 10.1007/s10577-012-9298-8.
29. Meitinger, F., Belal, H., Davis, R.L., Martinez, M.B., Shiau, A.K., Oegema,
K., and Desai, A. (2024). Control of cell proliferation by memories of
mitosis. Science (New York, N.Y *383*, 1441-1448.
10.1126/science.add9528.
30. Sparr, C., and Meitinger, F. (2025). Prolonged mitosis: A key indicator for
detecting stressed and damaged cells. Current opinion in cell biology *92*,
102449. 10.1016/j.ceb.2024.102449.
31. Meitinger, F., Anzola, J.V., Kaulich, M., Richardson, A., Stender, J.D.,
Benner, C., Glass, C.K., Dowdy, S.F., Desai, A., Shiau, A.K., and
Oegema, K. (2016). 53BP1 and USP28 mediate p53 activation and G1
arrest after centrosome loss or extended mitotic duration. The Journal of
cell biology *214*, 155-166. 10.1083/jcb.201604081.
32. Lambrus, B.G., Daggubati, V., Uetake, Y., Scott, P.M., Clutario, K.M.,
Sluder, G., and Holland, A.J. (2016). A USP28-53BP1-p53-p21 signaling
axis arrests growth after centrosome loss or prolonged mitosis. The
Journal of cell biology *214*, 143-153. 10.1083/jcb.201604054.
33. Fulcher, L.J., Batley, C., Sobajima, T., and Barr, F.A. (2025). Time as a
danger signal promoting G1 arrest after mitosis. Trends in cell biology.
10.1016/j.tcb.2025.06.001.
34. Fong, C.S., Mazo, G., Das, T., Goodman, J., Kim, M., O'Rourke, B.P.,
Izquierdo, D., and Tsou, M.F. (2016). 53BP1 and USP28 mediate
p53-dependent cell cycle arrest in response to centrosome loss and
prolonged mitosis. Elife *5*. 10.7554/eLife.16270.
35. Uetake, Y., and Sluder, G. (2010). Prolonged prometaphase blocks
daughter cell proliferation despite normal completion of mitosis. Curr Biol
20, 1666-1671. 10.1016/j.cub.2010.08.018.
36. Uetake, Y., and Sluder, G. (2018). Activation of the apoptotic pathway
during prolonged prometaphase blocks daughter cell proliferation.
Molecular biology of the cell *29*, 2632-2643. 10.1091/mbc.E18-01-0026.

37. Brinkley, B.R., Valdivia, M.M., Tousson, A., and Brenner, S.L. (1984).
Compound kinetochores of the Indian muntjac. Evolution by linear fusion
of unit kinetochores. Chromosoma *91*, 1-11.
38. Chi, J.X., Huang, L., Nie, W., Wang, J., Su, B., and Yang, F. (2005).
Defining the orientation of the tandem fusions that occurred during the
evolution of Indian muntjac chromosomes by BAC mapping.
Chromosoma *114*, 167-172. 10.1007/s00412-005-0004-x.
39. Wurster, D.H., and Benirschke, K. (1970). Indian muntjac, Muntiacus
muntjak: a deer with a low diploid chromosome number. Science *168*,
1364-1366. 10.1126/science.168.3937.1364.
40. Comings, D.E., and Okada, T.A. (1971). Fine structure of kinetochore in
Indian muntjac. Exp Cell Res *67*, 97-110.
41. Drpic, D., Almeida, A.C., Aguiar, P., Renda, F., Damas, J., Lewin, H.A.,
Larkin, D.M., Khodjakov, A., and Maiato, H. (2018). Chromosome
Segregation Is Biased by Kinetochore Size. Curr Biol *28*, 1344-1356
e1345. 10.1016/j.cub.2018.03.023.
42. Almeida, A.C., Drpic, D., Okada, N., Bravo, J., Madureira, M., and Maiato,
H. (2020). Functional Dissection of Mitosis Using Immortalized
Fibroblasts from the Indian Muntjac, a Placental Mammal with Only Three
Chromosomes. Methods Mol. Biol. *2101*, 247-266.
10.1007/978-1-0716-0219-5_16.
43. Almeida, A.C., Soares-de-Oliveira, J., Drpic, D., Cheeseman, L.P.,
Damas, J., Lewin, H.A., Larkin, D.M., Aguiar, P., Pereira, A.J., and Maiato,
H. (2022). Augmin-dependent microtubule self-organization drives
kinetochore fiber maturation in mammals. Cell Rep *39*, 110610.
10.1016/j.celrep.2022.110610.
44. Howell, B.J., Moree, B., Farrar, E.M., Stewart, S., Fang, G., and Salmon,
E.D. (2004). Spindle checkpoint protein dynamics at kinetochores in living
cells. Curr Biol *14*, 953-964. 10.1016/j.cub.2004.05.053
S0960982204003860 [pii].
45. Shah, J.V., Botvinick, E., Bonday, Z., Furnari, F., Berns, M., and
Cleveland, D.W. (2004). Dynamics of centromere and kinetochore
proteins; implications for checkpoint signaling and silencing. Curr Biol *14*,
942-952. 10.1016/j.cub.2004.05.046.
46. Kuhn, J., and Dumont, S. (2017). Spindle assembly checkpoint
satisfaction occurs via end-on but not lateral attachments under tension.
The Journal of cell biology *216*, 1533-1542. 10.1083/jcb.201611104.
47. Kapoor, T.M., Lampson, M.A., Hergert, P., Cameron, L., Cimini, D.,
Salmon, E.D., McEwen, B.F., and Khodjakov, A. (2006). Chromosomes
can congress to the metaphase plate before biorientation. Science (New
York, N.Y *311*, 388-391. 311/5759/388 [pii]
10.1126/science.1122142.
48. Barisic, M., Aguiar, P., Geley, S., and Maiato, H. (2014). Kinetochore
motors drive congression of peripheral polar chromosomes by
overcoming random arm-ejection forces. Nature cell biology *16*.
10.1038/ncb3060.
49. Itoh, G., Ikeda, M., Iemura, K., Amin, M.A., Kuriyama, S., Tanaka, M.,

Mizuno, N., Osakada, H., Haraguchi, T., and Tanaka, K. (2018). Lateral
attachment of kinetochores to microtubules is enriched in prometaphase
rosette and facilitates chromosome alignment and bi-orientation
establishment. Scientific Reports *8*, 3888. 10.1038/s41598-018-22164-5.
50.   Vukusic, K., and Tolic, I.M. (2025). CENP-E initiates chromosome
congression by opposing Aurora kinases to promote end-on attachments.
Nat Commun *16*, 8537. 10.1038/s41467-025-64148-w.
51.   Kwiatkowski, N., Jelluma, N., Filippakopoulos, P., Soundararajan, M.,
Manak, M.S., Kwon, M., Choi, H.G., Sim, T., Deveraux, Q.L., Rottmann,
S., et al. (2010). Small-molecule kinase inhibitors provide insight into
Mps1 cell cycle function. Nat Chem Biol *6*, 359-368.
10.1038/nchembio.345
nchembio.345 [pii].
52.   Vassilev, L.T., Tovar, C., Chen, S., Knezevic, D., Zhao, X., Sun, H.,
Heimbrook, D.C., and Chen, L. (2006). Selective small-molecule inhibitor
reveals critical mitotic functions of human CDK1. Proceedings of the
National Academy of Sciences of the United States of America *103*,
10660-10665. 10.1073/pnas.0600447103.
53.   Peretti, D., Maraschio, P., Lambiase, S., Lo Curto, F., and Zuffardi, O.
(1986). Indirect immunofluorescence of inactive centromeres as indicator
of centromeric function. Human genetics *73*, 12-16.
54.   Cherry, L.M., and Johnston, D.A. (1987). Size variation in kinetochores of
human chromosomes. Human genetics *75*, 155-158.
55.   Cherry, L.M., Faulkner, A.J., Grossberg, L.A., and Balczon, R. (1989).
Kinetochore size variation in mammalian chromosomes: an image
analysis study with evolutionary implications. Journal of cell science *92 (*
Pt 2), 281-289.
56.   Sanchez, L., Martinez, P., and Goyanes, V. (1991). Analysis of
centromere size in human chromosomes 1, 9, 15, and 16 by electron
microscopy. Genome *34*, 710-713.
57.   Tomkiel, J., Cooke, C.A., Saitoh, H., Bernat, R.L., and Earnshaw, W.C.
(1994). CENP-C is required for maintaining proper kinetochore size and
for a timely transition to anaphase. The Journal of cell biology *125*,
531-545.
58.   Nixon, F.M., Honnor, T.R., Clarke, N.I., Starling, G.P., Beckett, A.J.,
Johansen, A.M., Brettschneider, J.A., Prior, I.A., and Royle, S.J. (2017).
Microtubule organization within mitotic spindles revealed by serial block
face scanning EM and image analysis. Journal of cell science.
10.1242/jcs.203877.
59.   Logsdon, G.A., Rozanski, A.N., Ryabov, F., Potapova, T., Shepelev, V.A.,
Catacchio, C.R., Porubsky, D., Mao, Y., Yoo, D., Rautiainen, M., et al.
(2024). The variation and evolution of complete human centromeres.
Nature. 10.1038/s41586-024-07278-3.
60.   Parl, F.F. (2024). Analysis of CENP-B Boxes as Anchor of Kinetochores in
Centromeres of Human Chromosomes. Bioinform Biol Insights *18*,
11779322241248913. 10.1177/11779322241248913.
61.   Kixmoeller, K., Tarasovetc, E.V., Mer, E., Chang, Y.W., and Black, B.E.

62. Pereira, A., Sousa, M., Almeida, A.C., Ferreira, L.T., Costa, A.R., Novais-Cruz, M., Ferras, C., Sousa, M.M., Sampaio, P., Belsley, M., and Maiato, H. (2019). Coherent-hybrid STED: high contrast sub-diffraction imaging using a bi-vortex depletion beam. Optics express *27*, 8092-8111. 10.1364/OE.27.008092.

63. Zaytsev, A.V., Sundin, L.J., DeLuca, K.F., Grishchuk, E.L., and DeLuca, J.G. (2014). Accurate phosphoregulation of kinetochore-microtubule affinity requires unconstrained molecular interactions. The Journal of cell biology *206*, 45-59. 10.1083/jcb.201312107 jcb.201312107 [pii].

64. Volkov, V.A., Huis In 't Veld, P.J., Dogterom, M., and Musacchio, A. (2018). Multivalency of NDC80 in the outer kinetochore is essential to track shortening microtubules and generate forces. Elife *7*. 10.7554/eLife.36764.

65. Stimac, V., Koprivec, I., Manenica, M., Simunic, J., and Tolic, I.M. (2022). Augmin prevents merotelic attachments by promoting proper arrangement of bridging and kinetochore fibers. Elife *11*. 10.7554/eLife.83287.

66. Thompson, S.L., and Compton, D.A. (2010). Proliferation of aneuploid human cells is limited by a p53-dependent mechanism. The Journal of cell biology *188*, 369-381. 10.1083/jcb.200905057.

67. Santaguida, S., Richardson, A., Iyer, D.R., M'Saad, O., Zasadil, L., Knouse, K.A., Wong, Y.L., Rhind, N., Desai, A., and Amon, A. (2017). Chromosome Mis-segregation Generates Cell-Cycle-Arrested Cells with Complex Karyotypes that Are Eliminated by the Immune System. Developmental cell *41*, 638-651 e635. 10.1016/j.devcel.2017.05.022.

68. Hinchcliffe, E.H., Day, C.A., Karanjeet, K.B., Fadness, S., Langfald, A., Vaughan, K.T., and Dong, Z. (2016). Chromosome missegregation during anaphase triggers p53 cell cycle arrest through histone H3.3 Ser31 phosphorylation. Nature cell biology *18*, 668-675. 10.1038/ncb3348.

69. Giam, M., Wong, C.K., Low, J.S., Sinelli, M., Dreesen, O., and Rancati, G. (2020). P53 induces senescence in the unstable progeny of aneuploid cells. Cell Cycle *19*, 3508-3520. 10.1080/15384101.2020.1850968.

70. Phan, T.P., Maryniak, A.L., Boatwright, C.A., Lee, J., Atkins, A., Tijhuis, A., Spierings, D.C., Bazzi, H., Foijer, F., Jordan, P.W., et al. (2021). Centrosome defects cause microcephaly by activating the 53BP1-USP28-TP53 mitotic surveillance pathway. The EMBO journal *40*, e106118. 10.15252/embj.2020106118.

71. Brito, D.A., and Rieder, C.L. (2006). Mitotic checkpoint slippage in humans occurs via cyclin B destruction in the presence of an active checkpoint. Curr Biol *16*, 1194-1200. 10.1016/j.cub.2006.04.043.

72. Gascoigne, K.E., and Taylor, S.S. (2008). Cancer cells display profound intra- and interline variation following prolonged exposure to antimitotic drugs. Cancer Cell *14*, 111-122. 10.1016/j.ccr.2008.07.002.

73. Renda, F., Magidson, V., Tikhonenko, I., Fisher, R., Miles, C., Mogilner,
A., and Khodjakov, A. (2020). Effects of malleable kinetochore
morphology on measurements of intrakinetochore tension. Open biology
10, 200101. 10.1098/rsob.200101.
74. Sacristan, C., Samejima, K., Ruiz, L.A., Deb, M., Lambers, M.L.A.,
Buckle, A., Brackley, C.A., Robertson, D., Hori, T., Webb, S., et al. (2024).
Vertebrate centromeres in mitosis are functionally bipartite structures
stabilized by cohesin. Cell. 10.1016/j.cell.2024.04.014.
75. Altemose, N., Logsdon, G.A., Bzikadze, A.V., Sidhwani, P., Langley, S.A.,
Caldas, G.V., Hoyt, S.J., Uralsky, L., Ryabov, F.D., Shew, C.J., et al.
(2022). Complete genomic and epigenetic maps of human centromeres.
Science (New York, N.Y *376*, eabl4178. doi:10.1126/science.abl4178.
76. Zou, Y., Yi, X., Wright, W.E., and Shay, J.W. (2002). Human telomerase
can immortalize Indian muntjac cells. Exp Cell Res *281*, 63-76.
77. Zhang, G., Kruse, T., Guasch Boldu, C., Garvanska, D.H., Coscia, F.,
Mann, M., Barisic, M., and Nilsson, J. (2019). Efficient mitotic checkpoint
signaling depends on integrated activities of Bub1 and the RZZ complex.
The EMBO journal *38*. 10.15252/embj.2018100977.
78. Lukinavicius, G., Reymond, L., D'Este, E., Masharina, A., Gottfert, F., Ta,
H., Guther, A., Fournier, M., Rizzo, S., Waldmann, H., et al. (2014).
Fluorogenic probes for live-cell imaging of the cytoskeleton. Nat Methods
11, 731-733. 10.1038/nmeth.2972
nmeth.2972 [pii].
79. Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M.,
Pietzsch, T., Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., et al.
(2012). Fiji: an open-source platform for biological-image analysis. Nat
Methods *9*, 676-682. 10.1038/nmeth.2019.
80. Pereira, A.J., Matos, I., Lince-Faria, M., and Maiato, H. (2009). Dissecting
mitosis with laser microsurgery and RNAi in Drosophila cells. Methods
Mol Biol *545*, 145-164. 10.1007/978-1-60327-993-2_9.
81. Tinevez, J.Y., Perry, N., Schindelin, J., Hoopes, G.M., Reynolds, G.D.,
Laplantine, E., Bednarek, S.Y., Shorte, S.L., and Eliceiri, K.W. (2017).
TrackMate: An open and extensible platform for single-particle tracking.
Methods *115*, 80-90. 10.1016/j.ymeth.2016.09.016.
82. Kletter, T., Munoz, O., Reusch, S., Biswas, A., Halavatyi, A., Neumann,
B., Kuropka, B., Zaburdaev, V., and Reber, S. (2025). Cell state-specific
cytoplasmic density controls spindle architecture and scaling. Nature cell
biology *27*, 959-971. 10.1038/s41556-025-01678-x.
83. Jeff Reback, W.M., jbrockmendel, Joris Van den Bossche, Tom
Augspurger, Phillip Cloud, gfyoung, Sinhrks, Simon Hawkins, Adam Klein,
Matthew Roeschke, Jeff Tratner, Terji Petersen, Chang She, William Ayd,
MomIsBestFriend, Marc Garcia, Jeremy Schendel, Andy Hayden, …
Martin Winkel (2020). pandas-dev/pandas: Pandas 1.1.3 (v1.1.3).
Zenodo. .
84. VanderPlas, J., Granger, B., Heer, J., Moritz, D., Wongsuphasawat, K.,
Satyanarayan, A., Lees, E., Timofeev, I., Welsh, B., Sievert, S. (2018).
Altair: Interactive Statistical Visualizations for Python. Journal of Open

Source Software *3*, 1057. https://doi.org/10.21105/joss.01057.
1613

Figure 2

Figure 3

A Merge | Microtubules | ACA | Venus-MAD1

B Merge | 3D rendering

C Longitudinal KT line scanning

Relative signal intensity vs Distance along KT(μm)
— MAD1
— MTs

D Individual kinetochores vs Pearson correlation coefficient

E Synthetic data

Strong positive correlation
r ≈ 1

No correlation
r ≈ 0

Strong negative correlation
r ≈ -1

Z_{MAD1} / Z_{MTs}

Norm. intensity profile

F Experimental data

r = -0.29
(CI -0.24/-0.33)
p < 1e-10

Z_{MAD1} / Z_{MTs}

Figure 4

A Partial laser ablation of the KT   Longitudinal KT line scanning

● MAD1

B Pre-Surgery   After Surgery   Nocodazole

-01:40 min   00:00 min   08:00 min   15:20 min

□ 2xGFP-CENP-A; GFP-CENTRIN   ◧ mScarlet-MAD1

C

0.0015

Time after surgery (min)

Mean ± SD
□ 3.6 ± 2.8
◼ 8.2 ± 3.8

KT Bending   MAD1 Recruitment

B'

— NOC   — Surgery

MAD1/CENP-A relative signal intensity

Normalized distance along KT
0.0   0.5 (Ablated sister KT)   1.0

D

— NOC   — Surgery

MAD1/CENP-A relative signal intensity

Normalized distance along KT
0.0   0.5 (Ablated sister KT)   1.0

E

<0.0001

Coefficient of variation (%)

NOC   Surgery

F Live   Pre-Surgery   After Surgery

-01:00 min   12:00 min

□ 2xGFP-CENP-A; GFP-CENTRIN   ◧ mScarlet-MAD1

F' Fixed (CH-STED)   After Surgery

□ 2xGFP-CENP-A; GFP-CENTRIN   ▨ Microtubules

F''

— MAD1/CENP-A   — MTs/CENP-A

Relative signal intensity

Normalized distance along KT
0.0   0.5   1.0

Figure 5

A

Mock

2x: 0 s   600 s   1200 s   1800 s

siHAUS6

2x: 0 s   600 s   1200 s   1800 s

☐ SiR-Tubulin   ☐ Venus-MAD1

B

Relative Venus-MAD1 levels vs Time (s)

☐ Mock
☐ siHAUS6

C

	Half reduction time (s)	p-value	95 % CI	R²
Mock	142		128 - 158	0.53
siHAUS6	222	<0.0001	186 - 272	0.29

D

% Cells w/ MAD1(+) KTs

p<0.0001 (Prometaphase)
p=0.0008 (Metaphase)

☐ Mock
☐ siHaus6

E

Venus-Mad1 levels (KT/cyto)

p<0.0001 (Prometaphase)
p<0.0001 (Metaphase)

☐ Mock
☐ siHAUS6

F

+MG132 1h

Merge   ■ Microtubules   ☐ ACA   ☐ Venus-MAD1

Control

siHAUS6

Figure 6

Figure 7

Augmin-mediated K-fiber maturation

NEBD — Time — Anaphase

APC/C Activity

SAC Strength — Cyclin B1 / CDK1 — Threshold value

Initial occupancy — Partial occupancy — Full occupancy

Kinetochore — Microtubule — MAD1/MAD2

Accurate mitosis
Daughters proliferate

Inefficient K-fiber maturation (e.g. Augmin depletion)

NEBD — Time — Anaphase

APC/C Activity

SAC Strength — Cyclin B1 / CDK1 — Threshold value

Initial occupancy — Partial occupancy — Partial occupancy — Partial occupancy

Full occupancy

Kinetochore — Microtubule — MAD1/MAD2

STOP

Increased errors
Daughters arrest