Disordered but Different: The Unique Characteristics of Intrinsically Disordered Regions in Human Transcription Factors

Susie Song[1] and Joshua M. Akey[1*]

[1] Lewis-Sigler Institute for Integrative Genomics, Princeton University, Princeton, NJ. 08540, USA

[*] Correspondence: jakey@princeton.edu (JMA)

Abstract

Intrinsically disordered regions (IDRs) of a protein have fundamentally important biological functions, including enabling multivalent interactions and driving biomolecular condensate formation. Although IDRs are widespread in the human proteome, they are significantly enriched in transcription factor (TF) activation domains. However, the evolutionary dynamics of protein disorder and whether TF IDRs systematically differ from those in other proteins remain poorly understood. Here, we comprehensively identify and analyze IDRs across the proteome and show that TF and non-TF IDRs exhibit widespread functional, phenotypic, and evolutionary differences. Strikingly, in contrast to the broader proteome, TFs have evolved to become more disordered over time. Additionally, highly disordered TFs are more likely to regulate developmental processes, govern larger regulatory networks, and be subject to stronger regulatory constraints. Finally, TF IDRs are enriched for pathogenic mutations compared to non-TF IDRs, and disorder content significantly predicts the mode of disease inheritance. Our results provide novel insights into how the evolution of gene regulation has uniquely shaped the molecular function and disease burden of TF IDRs.

Introduction

Transcription factors (TFs) are modular proteins that make information encoded in the genome actionable by binding DNA and recruiting cofactors to activate or repress transcription[1]. TFs bind to regulatory elements, such as promoters and enhancers, through their structured DNA-binding domains (DBDs) and recruit cofactors through their intrinsically disordered activation domains. Unlike well-folded domains, intrinsically disordered regions (IDRs) of a protein lack a stable three-dimensional structure, and their biological functions are governed by the biochemical properties of their underlying amino acid sequences. Although TF activation domains are significantly enriched for IDRs[2], disorder is a pervasive feature across the human proteome that contributes to diverse cellular processes, including transcription, signal transduction, and stress response[3–5]. A hallmark of IDRs is their ability to adopt dynamic conformational ensembles, allowing them to form transient, multivalent interactions.

IDRs also promote liquid-liquid phase separation—a process that drives the formation of biomolecular condensates, which are membraneless compartments that regulate diverse cellular processes[6]. Recent studies suggest that gene expression can be mediated by the assembly of transcriptional condensates, molecular hubs enriched for TFs and other components of the transcriptional machinery[7–9]. Experimental studies further implicate these assemblies in enhancer activation, super-enhancer function, and the rapid transcriptional response to environmental perturbations[7,9–11]. Mechanistically, these condensates have been proposed to act by locally enriching regulatory factors at target loci and altering TF binding kinetics, with downstream effects on transcriptional bursting dynamics[12,13]. Furthermore, aberrant formation of transcriptional condensates has been linked to neurodevelopmental disorders and cancer[14].

Substantial progress has been made in delineating the functional roles that TF IDRs play in gene regulation and condensate dynamics[10,12]. Nonetheless, fundamental questions about the functional, phenotypic, and evolutionary significance of TF IDRs remain unresolved. In particular, TFs are strikingly enriched for IDRs relative to the rest of the proteome, but the evolutionary origins of this enrichment are poorly understood. Additionally, evolutionary and population-genetic analyses provide a powerful lens for addressing these questions, as patterns of variation, selective constraints, and sequence changes over time reveal which IDR

features are required and which can vary without disrupting function. Unlike structured domains, whose evolutionary trajectories are dominated by stability and folding constraints, IDRs evolve under a balance of relaxed purifying selection and episodic positive selection[15–20]. These properties make TF IDRs uniquely informative for understanding how regulatory complexity has evolved and how genetic variation in disordered regions contributes to human phenotypic diversity and disease susceptibility.

Population genetic analyses of protein-coding variation have yielded particularly clear insights for structured protein domains, where mutations can often be more directly associated with stability, binding, or catalytic activity and thus more interpretable in terms of selective pressure. In contrast, the unique mutational characteristics of IDRs and their roles in transient, context-dependent, and multi-protein interactions have been characterized in fewer studies. Existing work has revealed both positive and negative selection acting on these regions[15,16], but many questions remain. Here, we build on this work by focusing on the TF-specific properties of IDRs and leveraging recent large-scale variation and functional genomic datasets. In particular, the mechanisms by which genetic variation in IDRs influences TF function and the evolution of gene regulation remain unresolved. Studying TF IDRs in the context of human population differences is crucial because variations in these regions can influence GRNs, drive local adaptation, and influence phenotypic diversity.

Our study addresses this gap by integrating population genomic data with functional and structural annotations to investigate the evolution of IDRs in TFs. We specifically focus on the evolution of disorder within the human species and compare IDRs in TFs and non-TFs. Additionally, we explore how IDR variation in TFs occurs within human populations. By characterizing the population-level dynamics of genetic variation in these regions, we aim to provide new insights into the evolutionary forces shaping IDR diversity and its implications for human regulatory evolution, phenotypic diversity, and disease susceptibility.

Results

Proteome-wide identification and analysis of IDRs reveals distinct properties of TF disorder

Transcriptional condensates are emerging as a key component of gene regulation[7–9], and delineating the properties of TF IDRs is an essential step towards understanding how they form and regulate expression (Fig. 1a). To better understand the characteristics of TF IDRs and how they compare to disordered regions in other classes of proteins, we comprehensively identified IDRs in the human proteome (Fig. 1b). To this end, we downloaded the human proteome (n=20,659 proteins) from UniProt[21] and filtered out low-confidence proteins (see Methods), resulting in a final set of 19,736 proteins. In total, 8.2% (n=1,612) of proteins were TFs (Fig. 1b).

Next, we identified IDRs in these 19,736 proteins using MetaPredict[22,23], a deep-learning-based consensus predictor shown to achieve high accuracy in independent benchmarking studies[24]. In total, we identified 24,638 IDRs (Fig. 1c), and 80.3% of all proteins (13,832 of 19,736) contained at least one IDR segment. Among proteins containing both ordered and disordered regions (Extended Data Fig. 1a), the median disorder content was significantly higher in TFs compared to non-TFs (56% vs 25%, respectively; Mann-Whitney U test, $P<10^{-30}$; Fig. 1d). Pooling all amino acid residues across the primary sequences of all TF proteins, 56.5% fall within predicted IDRs and 43.5% within ordered regions; for non-TF proteins, the corresponding values are 31.1% and 68.9% (Fig. 1e). Additionally, almost all TFs had at least one IDR segment (Extended Data Fig. 1b), and compared to non-TFs, they are significantly more likely to have two or more IDR segments, even after adjusting for gene length (Mann-Whitney U test, $P<10^{-30}$; Fig. 1f). To confirm this, we stratified proteins by length and observed elevated segment density across length quantiles (Extended Data Fig. 1c). We fit a linear model with protein length as a covariate and the effect size of TF status was 1.41 and was significant ($P=1.7\times10^{-23}$; Extended Data Fig. 1c). When examining the length of individual IDR segments, the median TF IDR length was greater than that of non-TFs (Mann-Whitney U test, $P<10^{-30}$; Fig. 1g). Finally, we examined the location of IDRs along the protein and found a significant terminal bias at both terminals for both TFs and non-TFs (permutation $P<10^{-30}$; Fig. 1h). Our work extends the observation that TFs are enriched for disorder, by showing that TFs are also likelier to have more and longer IDR segments.

To ensure that our inferences were robust to the choice of IDR prediction method, we also identified disordered regions with AIUPred[25], which has also been shown to perform well in benchmarking studies[24] (see Methods). IDR predictions were highly concordant between MetaPredict and AIUPred (Extended Data Fig. 1d-f). For example, the number of amino acids predicted to be disordered per protein and the proportion of a protein predicted to be disordered were significantly correlated between the two methods (Pearson r=0.94 and 0.78, respectively; $P<10^{-30}$ for each correlation; Extended Data Fig. 1d-e). Similarly, the Hamming distance, which quantifies the fraction of residues with discordant predictions of disorder (see Methods), was small (Extended Data Fig. 1f). Perhaps most importantly, all of the key analyses described below resulted in the same qualitative and quantitative inferences when using either the MetaPredict or AIUPred callsets (Extended Data Fig. 1g-h). Thus, these data demonstrate that our results are robust to the specific method used to identify IDRs.

TFs evolved to become more disordered

The disorder-to-order model of protein evolution posits that newly emerged proteins are initially disordered and progressively acquire folded domains with time[26–28] (Fig. 2a). To assess whether this paradigm applies to TFs, we tested whether there was a relationship between the disorder content of TFs and non-TFs as a function of protein age. We obtained estimates of protein age from the GenOrigin[29] catalog, which infers gene age from ortholog mapping across 565 species divergence estimates. Under the disorder-to-order model, we would expect the oldest proteins to have the least disorder and the youngest proteins to have the most. The distribution of disorder content for non-TFs decreases as protein age increases (Kruskal-Wallis, $P<10^{-30}$; Fig. 2b), consistent with the disorder-to-order model. In contrast, TFs follow the opposite pattern, with the disorder content becoming higher as protein age increases (Kruskal-Wallis $P<10^{-30}$; Fig. 2b). Thus, the oldest TFs (>1000 Mya) are significantly enriched for disorder, whereas the oldest non-TFs are significantly depleted for disorder. These data suggest that TF IDRs have experienced a unique evolutionary trajectory compared to non-TF IDRs, in which disorder is not lost but instead maintained or expanded over evolutionary time (Fig. 2c).

Despite the striking differences in IDR content and gene age between TFs and non-TFs, several factors can confound age estimates[28,30–32]. Therefore, we performed additional

analyses to confirm that these patterns were robust. First, we repeated our analysis of disorder content as a function of age using an independent catalog of gene ages[33], which corrects for biases caused by failures in homology detection (Extended Data Fig. 2). Across phylostrata, we observed consistent trends in both TFs and non-TFs (Extended Data Fig. 2a). TFs that arose in Gnathostomata or earlier consistently exhibited higher disorder content than younger TFs (Mann-Whitney U test, P=8.7×10^{-10}; Extended Data Fig. 2b), whereas non-TFs showed the opposite pattern (Mann-Whitney U test, P<10^{-30}; Extended Data Fig. 2b). Second, patterns of IDR content and age for TFs and non-TFs remained the same when stratifying the GenOrigin-based gene age estimates by sequence conservation, protein length, GC content, expression level, and tissue specificity τ (Extended Data Fig. 2c–g), demonstrating that these potential confounding factors do not explain the elevated disorder in ancient TFs. Third, we used multiple linear regression to test whether the relationship between gene age and disorder content differed significantly between TFs and non-TFs, after controlling for the aforementioned covariates (protein length, GC content, mean expression level, tissue specificity; see Methods). The interaction term between disorder content and TF status was strongly positive (β=8.08, P<10^{-6}), indicating that the slope of the age–disorder relationship is significantly steeper in TFs than in non-TFs. In contrast, disorder content was negatively associated with gene age across the entire proteome (β=–5.86, P<10^{-15}), consistent with the canonical trend from disorder to order. Together, these results are consistent with the hypothesis that TFs deviate from the disorder-to-order model, maintaining or expanding their disorder content over time.

Because the relationship between disorder content and gene age is not attributable to technical confounders, we next investigated the evolutionary changes that may have contributed to this pattern. Specifically, we examined how the amino acid composition of IDRs changes over time, as such shifts can reveal selective pressures acting on IDR biochemistry. We tested 5 categories of amino acid residues: polar, aromatic, aliphatic, positively charged, and negatively charged. Of these, we found significant trends for positively-charged and aromatic residues. The youngest TF IDRs are significantly enriched for positively charged (lysine, arginine, histidine; Kruskal-Wallis, P<10^{-30}; Fig. 2d) and aromatic residues (phenylalanine, tyrosine, tryptophan; Kruskal-Wallis, P<10^{-30}; Fig. 2e), consistent with the hypothesis that aromatic residues arose later in the evolution of the genetic code[34–36]. Older TF

IDRs, in contrast, are relatively depleted of these residues (Fig. 2d-e). Non-TF IDRs show the opposite charge trajectory—an increase in basic residues with age—and only modest changes in aromatic content (Fig. 2d-e). The distinct charge trajectories imply that the physicochemical pressures shaping IDRs have diverged between TFs and non-TFs. In summary, these data show that TFs deviate from the classical disorder-to-order evolutionary trajectory, and that the biochemical grammar of their IDRs has itself evolved, potentially reflecting changing requirements for regulatory complexity and interaction capacity.

TF Families Exhibit Marked Heterogeneity in Disorder

TFs are organized into families based on their DNA-binding domain (DBD)[37,38], providing a natural framework to compare disorder content across functionally distinct TFs. We observed that TF families have strikingly different levels of disorder (Fig. 2f). Evolutionarily ancient families, such as HMG and homeodomain TFs, had the highest levels of disorder (median percentage of disordered residues per protein of 81% and 78%, respectively; Fig. 2f). In contrast, the nuclear receptor and KRAB family of TFs exhibit the lowest amount of disorder (median percentage of disordered residues per protein of 32% and 33%, respectively; Fig. 2f). Additionally, there is variation in IDR length, number, and magnitude of terminal bias among TF families (Extended Data Fig. 3a-e).

We next investigated whether highly ordered and disordered TFs were associated with specific biological processes. For example, many developmental TFs, such as *HOXA2* (86%), *HOXA3* (88%), *FOXO1* (90%), and *FOXO3* (90%), are highly disordered (>75% of their sequences are disordered). Indeed, GO enrichment analysis revealed that highly disordered TFs are significantly enriched (FDR ≤ 5%) for the term "developmental process" (GO:0032502) and many additional GO terms related to development and metabolism (Fig. 2g; Extended Data Fig. 4a). As expected, the average disorder of developmentally related TFs is significantly higher than TFs unrelated to development (Mann-Whitney U-test, $P<10^{-30}$; Extended Data Fig. 4b). Non-TFs annotated as developmental also have a significantly higher average disorder content compared to other non-TF proteins (Mann-Whitney U-test, $P=3.4\times10^{-9}$; Extended Data Fig. 4b), but magnitude of the difference was much smaller compared to TFs. Furthermore, pioneer TFs[39–41], which initiate transcription by binding to nucleosomes and promoting chromatin accessibility, are significantly enriched for disorder relative to other TFs

(Mann-Whitney U-test, $P=2.7\times10^{-4}$; Extended Data Fig. 4c). Overall, these data suggest that highly disordered TFs play a crucial role in core biological processes (Fig. 2g, Extended Data Fig. 4a). In contrast, ordered TFs (<45% of protein is disordered) tend to span many different GO terms and are enriched for a smaller number of GO terms. The categories they are enriched for tend to be signaling-related (Fig. 2g). Examples of highly ordered TFs include the STAT family of signaling TFs (*STAT1, STAT3, STAT4*) and the IRF family of immune TFs (*IRF4, IRF8*).

The regulatory characteristics of TFs are shaped by their disorder content

To evaluate whether the amount of disorder in TFs is associated with functional characteristics, we examined protein-protein interaction (PPI) networks and gene regulatory networks (GRNs). We hypothesized that highly disordered TFs would interact with a larger protein network. Using the STRING PPI database, we found a significant positive correlation between TF disorder content and the number of proteins in the TF's functional association network (indirect interactions) (Spearman's $\rho=0.36$, $P<10^{-30}$; Fig. 3a). In contrast, non-TFs exhibit a significantly negative correlation (Spearman's $\rho=-0.14$, $P<10^{-30}$; Fig. 3a). This finding is consistent with the hypothesis that TF IDRs in their effector domains facilitate multiple interactions, enabling flexible coordination of transcriptional cofactors and machinery. Notably, this trend is entirely driven by the largest family of TFs, C2H2 zinc finger TFs (Spearman's $\rho=0.55$, $P<10^{-30}$; Fig. 3a).

We confirmed that the positive relationship between disorder content and PPI connectivity in TFs was robust across multiple definitions and confidence thresholds (Extended Data Fig. 5). For example, when quantified as the total number of disordered amino acids rather than as a percentage, disorder content remained significantly correlated with the number of PPIs (Spearman's $\rho=0.22$, $P=2.5\times10^{-19}$), which was again largely driven by C2H2 zinc-finger TFs (Spearman's $\rho=0.40$, $P=5.7\times10^{-30}$; Extended Data Fig. 5a). We also found that the relationship between disorder content and PPIs for TFs and non-TFs was robust to the confidence threshold used to define an interaction (see Methods), for both all interactions and when restricted to physical interactions (Extended Data Fig. 5b). Furthermore, the analysis was repeated with two additional PPI databases: the HPA[42] and BioGRID[43], and we saw the same trends with these data (Extended Data Fig. 5c-d). These data suggest that the relationship

between disorder and the number of interacting proteins is not an artifact of database coverage or interaction type.

Next, we examined GRNs to determine whether there is a relationship between TF disorder content and the GRN it regulates. We used previously published GTEx GRNs. Interestingly, we observed a significant positive correlation between TF disorder content and the number of target genes it is associated with in GRNs (Spearman's ρ=0.28, P=1.5×10^{-23}; Fig. 3b), across a broad range of GTEx tissues (Spearman's ρ=0.32, P<10^{-30}; Fig. 3c). We also confirmed that these correlations were also observed in an independent dataset, The Cancer Genome Atlas (TCGA)[44], instead of GTEx (Extended Data Fig. 6a-b).

In summary, the degree of intrinsic disorder in TFs is strongly associated with their functional characteristics. As shown in Fig. 3d, TFs with higher levels of disorder tend to participate in more protein–protein interactions, regulate larger gene regulatory networks, and are regulators across a broader range of tissues. Collectively, these findings are consistent with the observation that ancient TFs are more highly disordered and that intrinsic disorder contributes to their broad regulatory roles.

Highly disordered TFs are under strong regulatory constraint

We next explored the genetic architecture underlying the regulation of TFs themselves. The functional and network characteristics of highly disordered TFs suggest that their expression may be more tightly regulated. Using the Human Protein Atlas (HPA)[42], we found that TFs detected in fewer tissues had higher disorder (Kruskal-Wallis, P<10^{-30}; Extended Data Fig. 4c). Beyond transcript detection, we examined expression patterns unique to certain tissues (tissue-specific). TFs expressed in a tissue-specific manner were enriched for disorder (Mann-Whitney U test, P<10^{-30}; Fig. 4a). This was also consistent with cell type-specific expression (Extended Data Fig. 4d). Additionally, we found that TFs with low gene expression variance are enriched for disorder, suggesting that the transcription of these genes is tightly regulated due to their essential roles (Mann-Whitney U test, P=5.0×10^{-11}, Fig. 4b). Since purifying selection can purge genetic variants that affect gene expression, we hypothesized that highly disordered TFs would have fewer expression quantitative trait loci (eQTLs). We examined the fine-mapped sets from GTEx[45] v10 and found that genes that are eGenes in many tissues are depleted for disorder (Fig. 4c). Additionally, we found that genes regulated by

many independent credible sets, indicating multiple causal genetic variants, are depleted for disorder (Extended Data Fig. 6e). Interestingly, the same trend was also seen with fine-mapped splicing QTLs (sQTLs) (Fig. 4e, Extended Data Fig. 6f-g). This finding suggests that isoform selection, not only gene expression, is tightly regulated in these highly disordered TFs, and that population variance is poorly tolerated. Fewer eQTLs may indicate that regulatory variation is not tolerated and requires tighter transcriptional control, thereby reducing gene expression variance. To further investigate this, we investigated translation efficiency, which is affected by codon optimization − a process by which genes evolve to utilize advantageous codons over millions of years. We found that in the highest translational efficiency group, TFs are enriched for disorder while non-TFs exhibited a depletion of disorder, mirroring the trend seen with evolutionary age (Spearman's ρ=0.16, P=4.1×10^{-7}, Fig. 4d). We found that these correlations were significant across most of the 78 cell types, and TFs showed a strong positive correlation between disorder and translation efficiency, whereas Non-TFs had a negative correlation (Fig. 4e). Together, these findings suggest that many highly disordered TFs are indeed older and have thus evolved robust, redundant regulatory architectures that buffer the impact of individual variants or transcriptional noise, consistent with their essential roles in gene regulatory networks (Fig. 4f).

Population genetics characteristics of TF IDRs

Because TF IDRs exhibit distinct evolutionary histories and biochemical properties, we next asked whether these differences are reflected in patterns of human genomic variation. To this end, we leveraged the gnomAD[46] database to compare characteristics of genetic variation in ordered and disordered protein domains between TFs and non-TFs. The gnomAD database is a powerful resource for population-genetic inference, as it aggregates sequence data from up to 1 million individuals.

For both TFs and non-TFs, SNVs and indels are significantly enriched in IDRs relative to non-IDRs (Binomial exact test, P<10^{-30}; Fig. 5a). Specifically, in TFs, 60% of SNVs and 62% of indels lie in IDRs, significantly exceeding the 57% of disordered TF coding sequence (Fig. 5a). For TFs, these numbers translate to a variant density of 355.1 SNVs/kb in IDRs and 307.2 SNVs/kb in non-IDRs. Non-TFs exhibit a similar enrichment, with 32% of SNVs and 36% of indels falling in IDRs, despite IDRs constituting only 31% of their coding genome (Binomial

exact test, $P<10^{-30}$; Fig. 5a). For non-TFs, these numbers translate to a variant density of 339.0 SNVs/kb in IDRs and 322.8 SNVs/kb in non-IDRs. Collectively, these data extend the observation that disordered regions tolerate more mutations than ordered domains by showing that although TFs are subjected to higher overall levels of purifying selection than non-TFs, TF IDRs are still highly tolerant of variation.

Interestingly, although indels exhibit the same qualitative patterns as SNVs, differences in the density of TF indels between IDR and non-IDR regions (Binomial exact test, $P<10^{-30}$; 21.7 vs 17.1 indels/kb, respectively) are substantially smaller compared to non-TFs (Binomial exact test, $P<10^{-30}$; 22.8 vs 18.4 indels/kb in IDRs and non-IDRs, respectively). Therefore, TF IDRs are less tolerant to indels than non-TF IDRs, suggesting that changes in activation-domain length are deleterious despite their propensity to be disordered.

To more quantitatively assess levels of genetic diversity, we calculated nucleotide diversity θ_W and π in IDRs and non-IDRs across the proteome (Fig. 5c-f). θ_W measures the number of segregating sites normalized by sequence length, whereas π measures the average number of pairwise nucleotide differences between chromosomes. Under the standard neutral model, both of these statistics estimate the scaled population mutation rate $\theta = 4N_e\mu$, where N_e is the effective population size and μ is the mutation rate per bp per generation. In 74.4% of TFs, θ_W was higher in IDRs compared to non-IDRs (Wilcoxon, $P<10^{-30}$; Fig.5c). A similar pattern was observed for non-TFs, where 72.4% of genes had elevated θ_W values in IDRs (Wilcoxon, $P<10^{-30}$). The overall difference in θ_W between all IDR and all non-IDR regions is larger for TFs compared to non-TFs (Fig. 5d).

Nucleotide diversity π also varied between ordered and disordered regions of TFs (Fig. 5e-f). Specifically, 65.7% of TFs had higher π in their IDRs compared to their non-IDR regions (Wilcoxon, $P=9.3\times10^{-28}$). In contrast, a higher π in non-IDRs was observed for 50.2% of non-TFs (Wilcoxon, $P=2.9\times10^{-4}$; Fig. 5e-f). The overall difference in π between IDR and non-IDR regions, similar to θ_W differences, was larger for TFs compared to non-TFs (Fig. 5f).

To better understand the different patterns of π and θ_W, we calculated Tajima's D, which is the normalized difference between π and θ_W. Tajima's D is expected to be zero under the standard neutral model. Tajima's D was negative across all groups, consistent with an excess of rare

variants due to human population growth, but IDRs again showed a distinct bias (Fig. 5g-h). For TFs, 59.1% of genes had more negative values in IDRs compared to non-IDRs. Non-TFs showed bias in the other direction, with 73.8% of genes having elevated D in non-IDRs compared to non-IDRs. Across the genome, IDR regions showed more negative D values than non-IDR regions for both TFs and non-TFs (Fig. 5h). We confirmed that these trends were similar using variants from the 1000 Genomes project (Extended Data Fig. 7a-h). Together, these findings indicate that IDRs have a higher density of rare variants than non-IDRs, consistent with their increased mutational tolerance, and that these trends were stronger in TFs than in non-TFs.

Finally, we compared population-structure patterns between IDRs and non-IDRs, as differences could reflect heterogeneity in selective pressures. To quantify population structure, we calculated locus-specific branch lengths (LSBL) in African (AFR), European (EUR), and East Asian (EAS) individuals from the 1000 Genomes Project. LSBL is a simple function of pairwise F_{ST} values among the three populations, yielding a branch-specific FST value for each population, with higher values indicating greater allele-frequency divergence. In the EAS population, we compared per-gene F_{ST} values between IDR and non-IDRs ($\Delta F_{ST} = F_{ST, IDR} - F_{ST, Non-IDR}$), after filtering to proteins with at least 20% disorder and no more than 80% disorder. ΔF_{ST} values were skewed towards positive values as 55.2% of TFs had elevated $F_{ST, IDR}$ compared to $F_{ST, Non-IDR}$ (Wilcoxon, $P=2.6\times10^{-4}$; Fig. 5i, left). However, among non-TFs, nearly equal proportions of genes favored IDRs or non-IDRs (Wilcoxon, $P>0.05$; Fig. 5i, right). This bias was also seen in AFR and EUR populations (Extended Data Fig. 7i-j).

Disorder content shapes the disease landscape of TFs and non-TFs

We next investigated whether the disorder content of TFs and non-TFs was related to disease characteristics. We first analyzed genes associated with lethal phenotypes (see Methods). Lethal TFs had a significantly higher disorder content than non-lethal TFs (Mann-Whitney U test, $P=2.1\times10^{-3}$; Fig. 6a), consistent with the previous observation that highly disordered TFs are enriched for core biological functions (Fig. 2g). In contrast, non-TFs showed the opposite trend with lethal genes being depleted for disorder compared to non-lethal genes (Mann-Whitney U test, $P=6.6\times10^{-15}$; Fig. 6a). Among lethal TFs, the median disorder content was significantly higher for proteins that exhibited prenatal or neonatal lethality compared to

adolescent or adult lethality (Kruskal-Wallis: P=3.5×10^{-3}; Fig. 6b). Again, non-TFs exhibited the opposite pattern with adolescent or adult lethality associated with the highest disorder (Kruskal-Wallis: P=1.9×10^{-6}; Fig. 6b). One contributor to these differences may be that developmental TFs are significantly more disordered than non-developmental TFs (Extended Data Fig. 4b). More broadly, these opposing trends are consistent with the view that TF IDRs often encode interaction capacity that can have far-reaching regulatory effects, whereas lethal non-TFs may be enriched for proteins whose essential function depends on structured catalytic or structural domains.

We also found that the mode of inheritance for pathogenic mutations was significantly related to disorder content (Fig. 6c). Specifically, genes with autosomal dominant (AD) pathogenic mutations were more disordered than genes with autosomal recessive (AR) mutations for both TFs (Mann-Whitney U test, P>0.05; Fig. 6c) and non-TFs (Mann-Whitney U test, P=2.1×10^{-16}; Fig. 6c). Although these results are not significant within TFs, this likely reflects the propensity of IDRs to mediate flexible, multivalent interactions. Thus, mutations are more likely to be dominant-negative, gain-of-function, haploinsufficient, or disrupt the formation of phase-separated condensates in proteins with more disorder, resulting in AD patterns of inheritance[47–49]. In contrast, mutations in proteins with less disorder are more likely to have an AR mode of inheritance as they primarily cause loss-of-function effects that do not disrupt the activity of the wild-type allele[50–52].

We next examined where disease variants fall according to IDR status. High-confidence pathogenic SNVs from Human Gene Mutation Database (HGMD)[53] were overall enriched in ordered regions, consistent with the stronger structural constraints in these regions (Fig. 6d). However, this depletion was weaker for TFs than for non-TFs, indicating that, relative to other proteins, pathogenic SNVs in TFs are more evenly distributed between IDRs and non-IDRs. Thus, although disease variants still preferentially affect non-IDRs, TF IDRs carry a higher burden of pathogenic variants than IDRs in the rest of the proteome. This suggests that pathogenic variants in TF IDRs are more consequential than previously appreciated, potentially because disruption of IDR-mediated interactions or condensate assembly can have broad regulatory consequences.

Finally, we asked whether the balance of IDR versus non-IDR pathogenic variants differs by disease type (Fig. 6e). Grouping HGMD variants by annotated tissue or disease category, we computed odds ratios for IDR versus non-IDR localization separately for TFs and non-TFs. The four disease categories that had significantly different odds ratios between TFs and non-TFs were endocrine/metabolic, neuropsychiatric, blood, and neoplasms. Neuropsychiatric phenotypes showed a relative shift toward IDR localization of pathogenic variants in TFs compared to non-TFs, whereas endocrine or metabolic and blood disorders showed stronger enrichment of variants in non-IDR regions, particularly for non-TF genes. These tissue-specific patterns suggest that the contribution of TF IDRs to disease is not uniform across phenotypes, and that disruption of TF IDRs is especially relevant for certain disorders.

Discussion

TFs are fundamentally essential proteins that regulate the timing, cell-type specificity, and magnitude of gene expression[1,37]. Through their ability to integrate diverse regulatory inputs and coordinate complex gene regulatory networks, TFs influence nearly all biological processes, and many diseases result from aberrant gene expression[37]. Despite decades of work elucidating how TFs bind DNA, recruit cofactors, and modulate chromatin and transcriptional output, our understanding of the mechanistic principles underlying TF function remains incomplete. In particular, accumulating evidence suggests that gene regulation is often organized through the assembly of transcriptional condensates, dynamic, multicomponent hubs that concentrate TFs, coactivators, and the transcriptional machinery at regulatory elements[7,8]. A defining feature of TFs that enables such assemblies is the presence of IDRs, which mediate multivalent interactions and are thought to play a central role in condensate formation and function. These observations raise the possibility that TF IDRs are subject to distinct functional and evolutionary constraints relative to disordered regions in other proteins. Motivated by this hypothesis, we comprehensively compared the properties of IDRs in TFs and non-TFs, finding that distinct evolutionary pressures and functional demands shape TF disorder.

In contrast to most proteins that evolve towards greater order over time as they acquire more energetically favorable secondary structures, we find that TFs systematically deviate from this paradigm. Strikingly, TFs have evolved towards greater disorder, suggesting that unique

evolutionary forces shape their IDRs, as they often serve as effector domains. One interpretation of these results is that TFs are selected to maintain or expand their disorder as their GRNs become increasingly complex, context-specific, and entrenched over time. Ancient TFs are more deeply integrated into regulatory pathways, and selection may favor IDRs that enable multivalency (cofactor recruitment) and more tissue-specific regulatory function. However, it is also possible that neutral drift or mutational bias could act as alternative drivers of rising disorder, independent of selective pressures for multivalency. IDRs in TFs can evolve under constraints that deprioritize fold stability in favor of more IDRs, thereby conferring a more complex cofactor interaction grammar, such as motif spacing, charge patterning, and the density of aromatic residues. Over long evolutionary timescales, constraints on TFs and their GRNs can lead to stable or increasing disorder, even as overall gene age increases. In conclusion, TFs, like most of the proteome, are disordered but different in their characteristics and evolution, so proteome-wide generalizations about disorder should not be assumed. These findings also motivate increasing attention to IDR-mediated transcriptional condensates as potential contributors to regulatory innovation and human phenotypic evolution.

Across interaction and regulatory networks, TF IDRs are uniquely correlated with regulatory scope and complexity. For example, the disorder content of TFs, but not non-TFs, scales with the size of the physical PPI network, consistent with increased interaction capacity for cofactor recruitment during transcription. Furthermore, TF IDRs also scale with increasing regulatory complexity, as demonstrated by positive correlations with the size of the functional association PPI network, the number of target genes, and the number of tissues in which a TF governs a tissue-specific expression pattern. Interestingly, these trends emerge from IDRs rather than from structured DBDs, suggesting that IDR differences, rather than DBD differences, better capture variation between TF GRNs.

These results further suggest an interaction between IDRs and tighter dosage control of TFs themselves. Disorder content in TFs is associated with tighter transcript abundance and isoform regulation, reflected in reduced inter-individual expression and splicing variability and a depletion of fine-mapped eQTL and sQTL signals. One explanation is that IDR-mediated multivalency increases a TF's regulatory influence, so fluctuations in dosage or isoform composition propagate across more targets and contexts, increasing fitness costs and selecting for regulatory mechanisms that stabilize TF abundance.

Human variation in IDRs also differs between TFs and non-TFs. IDRs show higher nucleotide diversity (π and θ_w) compared to non-IDRs, with a larger contrast in TFs rather than non-TFs. Similarly, population differentiation is elevated in IDRs compared to non-IDRs in TFs, but not in non-TFs. Extending to human disease, TFs associated with lethal Mendelian phenotypes tend to have higher disorder and earlier developmental lethality, suggesting that coding variation in TF IDRs can contribute to severe phenotypes even though the overall pathogenic burden remains enriched in non-IDRs. Current variant-effect predictors rely heavily on sequence conservation, which is often weak in IDRs even when these regions are functionally constrained. Our results show that TF IDR variation and pathogenic burden are structured by TF regulatory context, such as network connectivity and tolerated expression variation. Thus, incorporating these gene- and network-level features alongside local sequence properties may improve variant-impact prediction in IDRs beyond conservation alone.

There are several limitations of this work that are worth noting. For instance, although the methods we used to predict disorder have been shown to perform well[22,25,54], future studies will ideally leverage experimentally validated IDRs. Furthermore, the binary classification of a protein segment as being disordered or ordered does not reflect the complexity of IDRs, which can undergo context-dependent order-to-disorder transitions and vice versa. Finally, isoform-aware analyses are needed to resolve cell-type-specific regulatory activity and to avoid conflating protein- and transcript-level effects. Future work integrating improved catalogs of structural variation, isoform-resolved annotations, and analyses of conservation of biochemical grammar would strengthen mechanistic inference. In summary, TF IDRs have unique functional, phenotypic, and evolutionary characteristics, and our results provide novel insights into how protein disorder shapes the evolution of complex gene regulatory systems. More broadly, mapping how variation in TF IDRs uniquely rewires regulatory interactions provides a foundation for interpreting mutations in these regions and for understanding how regulatory dysfunction arises in human disease.

Methods

Obtaining and Curating the Human Proteome

The UniProt reference human proteome was downloaded from the UniProt Knowledgebase. This dataset (Proteome ID: UP000005640; Taxonomy: 9606; Organism: Homo sapiens; Release: 2025_04; Date: 15-Oct-2025) initially comprised 20,659 proteins with canonical sequences. Protein annotations were obtained from the UniProt API[55]. We applied a series of filters to retain high-confidence protein-coding genes. Specifically, entries not included in the manually curated Swiss-Prot (SP) database were excluded. Additional exclusions were made for proteins lacking gene name annotations, those shorter than 20 amino acids, and proteins with the lowest UniProt protein existence score (score=5). Post-filtering, a total of 19,736 proteins were retained.

To identify the subset of proteins with genomic coordinates from the GRCh38/hg38 reference, we linked UniProt accession IDs to CDS-level annotation via the Proteins REST API[56] (https://www.ebi.ac.uk/proteins/api/coordinates). Proteins without genomic coordinates and those that did not map to chromosomes 1–22, X, and Y, were discarded. This process yielded a set of 19,226 proteins. To confirm the accuracy of these coordinates, we retrieved the GRCh38 sequences corresponding to each protein's CDS coordinates using the Ensembl REST API (https://rest.ensembl.org/sequence/region/human), resulting in a final set of 18,899 proteins (18,077 autosomal, 785 X-chromosomal, and 37 Y-chromosomal proteins). To connect UniProt entries to entries in other genomic databases, UniProt ID-to-Ensembl Gene Stable ID mappings were used.

Annotation of Transcription Factors

A total of 1,639 TFs and their corresponding DNA-binding domain (DBD) motifs were identified from the manually curated Human TFs catalog (2018, v1.01; http://humantfs.ccbr.utoronto.ca)[37]. An intersection of the 1,639 TFs with the aforementioned 19,736 UniProt proteins, using Ensembl IDs, resulted in a refined set of 1,613 TFs for subsequent analyses. The set of pioneer TFs was obtained from a previously published list[39]. Our TF set included 32 out of 34 pioneer TFs listed in this resource (*PAX7, PBX1, GRHL2, GATA4, FOXO1, GATA5, SPI1, GRHL1, CEBPB, SOX9, SOX2, FOXA1, GATA1, GATA2,*

FOXM1, RUNX3, FLI1, FOXD3, TFAP2C, GATA3, GATA6, PBX3, POU5F1, ASCL1, TP63, RUNX1, MYB, FOXE1, TP53, OTX2, FOXA2, KLF4).

Identifying IDRs

A consensus disorder prediction tool, MetaPredict (v3.1), was used to identify IDRs, which has been shown to be highly accurate in benchmarking studies[24]. The UniProt FASTA file of primary sequences was input into the MetaPredict Colab notebook (https://colab.research.google.com/github/holehouse-lab/ALBATROSS-colab/blob/main/idrome _constructor/idrome_constructor.ipynb), which assigned disorder scores between 0 and 1 for each residue and used the default score threshold of ≥ 0.5 to classify residues as disordered. In defining IDRs, we used a minimum size threshold of 20 contiguous amino acids to focus on functionally relevant segments and avoid short linker regions. These criteria yielded 28,565 IDR segments.

We also called proteome-wide IDRs on the same UniProt FASTA file using the AIUPred software with the default settings (--analysis-type disorder --no-smoothing False). We selected AIUPred because it exhibits high accuracy in independent benchmarking studies[24] and is methodologically distinct from MetaPredict. Briefly, AIUPred uses a deep learning algorithm trained on an explicit biophysical model to identify IDRs, whereas MetaPredict obtains consensus calls by integrating many disorder prediction methods. In defining IDRs, we once again used a score threshold of ≥ 0.5 and a minimum size threshold of 20 contiguous amino acids. These criteria yielded 28,991 IDR segments.

Gene age estimates

We obtained gene age estimations from GenOrigin[29], a catalog of gene ages inferred across 565 species. Out of the 19,736 proteins described above, 18,123 had TimeTree[57] age estimates and Ensembl IDs that mapped to a UniProt accession number. Although gene intervals ranged from '0–7 Mya' to '>4290 Mya', we aggregated them into four age bins (0–100, 100–500, 500–1000, >1000 Mya) to mitigate potential estimation errors. To confirm the robustness of our results, we used another human gene age catalog[33] that used GenEra[58], which accounts for biases introduced by homology detection failure[30]. We mapped Ensembl protein IDs to UniProt accession numbers and filtered out proteins with incomplete phylostratum information and those with phylostrata more recent than Eutheria.

Furthermore, we also repeated analyses by stratifying the data by conservation, protein length, GC content, expression levels, and tissue specificity. We defined conservation using PhyloP 100-way scores downloaded from UCSC (May 2015 version). We intersected this with the aforementioned set of 18,899 proteins with precise genomic coordinates and calculated the per-gene mean phyloP score. These scores were stratified into tertiles in order to test the robustness of the disorder-evolution relationship in TFs and non-TFs. Expression levels and tissue specificity were obtained from GTEx v10 bulk tissue expression data. Specifically, we first obtained the median gene-level TPM by tissue (GTEx_Analysis_v10_RNASeQCv2.4.2_gene_median_tpm.gct.gz). Next, to bin proteins by gene expression level, we calculated the median TPM across all tissues and stratified these values into tertiles. The tissue specificity index, τ, was calculated across all tissues according to the original method[59].

We modeled gene age (from GenOrigin estimates) using ordinary least squares (OLS) regression to test whether the association between gene age and disorder content differs between TFs and non-TFs. TF status was encoded as a binary indicator variable $\mathbb{1}[TF]$, (1 for TFs, 0 otherwise). Disorder content D (percent disordered residues; 0-100) was mean-centered ($D_C = D - \overline{D}$). The interaction term of these two variables was included. The model also included covariates X_k for protein length, GC content, mean expression level, and tissue specificity (τ). We modeled gene age as follows:

$$GeneAge \sim \beta_0 + \beta_1\,\mathbb{1}[TF] + \beta_2\,D_C + \beta_3\,(\mathbb{1}[TF] \times D_C) + \sum_k \beta_k X_k + \epsilon$$

GO Analysis

Gene Ontology (GO)[60] enrichment of biological process (BP) terms was performed using GOATOOLS[61] with reference annotations from the GO Association File (GAF) (https://geneontology.org/docs/download-go-annotations, release from 2025-07-22). Developmental TFs and non-TFs were labeled based on GO term annotation for "developmental process" (GO:0032502). The full GO term ontology OBO file was obtained from the Gene Ontology Knowledgebase (https://geneontology.org/docs/download-ontology, downloaded 2025-07-25). A custom GO slim set was constructed by subsetting to terms at Levels 3 and 5 to avoid overly broad and overly narrow terms. The default GOATOOLS

settings were used, with Fisher's exact test and multiple-testing correction by the Benjamini–Hochberg method to control the false discovery rate.

PPI network analysis

Protein–protein interaction (PPI) data were obtained from the STRING[62] database (v12.0) after filtering to Homo sapiens-specific data and removing redundant interaction pairs (ex. AB-BA). Analyses were performed using both the full network and the physical subnetwork. To assess robustness, we repeated analyses using multiple interaction confidence thresholds based on the interaction score, which ranges from 0 to 1000. We assessed networks after thresholding to interactions with scores greater than 150, 400, and 700, corresponding to low, medium, and high confidence, respectively. After intersecting this catalog with our protein set using STRING protein IDs and subsetting to proteins with at least 1 interaction, 19,058 proteins were used for further analysis.

We also assessed PPI using the Human Protein Atlas (HPA)[42] Interaction resource, which integrates data from multiple databases to construct a consensus network of interactions present in at least two datasets. We subsetted to proteins with at least 1 interaction in the catalog and removed proteins with a reliability score of "Uncertain", resulting in a set of 15,095 proteins. Lastly, we used physical protein interaction counts from BioGRID[43], resulting in 18,325 proteins with at least 1 interaction.

GRN analysis

TF–target gene regulatory interactions were obtained from GRNdb[63], subsetted to TF-gene interactions with "High" confidence. We used regulons predicted from both GTEx (27 tissues) and TCGA (43 tissues) databases. To obtain the total number of targeted genes per TF, we summed the number of unique genes across all tissue regulons. Tissue breadth was defined as the number of distinct tissues in which a TF was connected to at least one target gene.

Expression Analysis

We obtained gene expression specificity information from the HPA[42] (version 25.0) and removed proteins with a reliability score of "Uncertain".

Inter-individual gene expression variance data were obtained from a previously published dataset[64], which quantified inter-individual expression variance data from 24 GTEx tissues, 11 TCGA (The Cancer Genome Atlas) tissues, and 22 other individual studies. We intersected genes using Ensembl gene IDs and compared the top 50% "High" and "Low" variance TFs and non-TFs. To show the robustness of the variance-disorder relationship in TFs and non-TFs, we also used the per-study mean gene expression variance to calculate Spearman correlation within each of the 58 tissues and used Bonferroni P value correction.

QTL analysis

We obtained fine-mapped expression QTL (eQTL) and splicing QTL (sQTL) data from GTEx[45] v10 (November 2024) found across 50 GTEx tissues. Although QTL credible set (CS) overlap was extremely rare (occurring in only 17 instances), we combined any overlapping CSs by at least one variant to prevent redundancy. 17,085 out of 19,736 genes intersected in GTEx genes, based on HGNC ID and ENSG. For each gene, we counted the number of tissues where it was an eGene (linked to at least 1 CS). For each gene, we also counted the number of distinct CSs per tissue to calculate the per-tissue Spearman correlations between eGene disorder and the number of independent CSs. Bonferroni P value correction was used. As a summary measure, for each eGene, we identified the tissue with the most credible sets and correlated this maximum CS number with the eGene's disorder content.

Translational Efficiency

Translational efficiency (TE) data were obtained from a previously published resource[65], which predicted TE for 10,037 transcripts across 78 studies, trained on 1,076 human ribosome profiling datasets. We intersected this gene catalog using gene symbols, resulting in 9,887 genes. First, we calculated the per-study Spearman correlation between predicted TE and disorder content, and we used Bonferroni P value correction. We then used the mean predicted TE across all samples to summarize the correlation between TE and disorder content.

Human Population Genetics Analyses

Human genetic variation data were obtained from gnomAD v4.1.0 and the 1000 Genomes Project (1KGP; Phase 3, accessed March 12, 2019). We used only high-quality variants

(FILTER=PASS). From gnomAD, we only used polymorphisms within the non-Finnish European (NFE) subpopulation (AC_nfe>0), in order to minimize population structure effects. From the 1KGP, we used the phased biallelic SNP+INDEL call set for unrelated samples ($n=$ 5,248) based on the GRCh38 assembly. To calculate the SNV and indel rates per kilobase, we annotated VCFs by TF status and IDR status (TF IDRs, Non-TF IDRs, TF non-IDRs, and non-TF non-IDRs). We summed variant counts genome-wide within each of the four regions and normalized by the corresponding total genomic region.

To calculate per-base nucleotide diversity metrics, we subsetted to biallelic SNVs. We used the following method to calculate Watterson's theta (θ_W), where S is the number of segregating sites, and N is the maximum number of chromosomes in the sample:

$$\theta_W = \frac{S}{a_1}, \text{ where } a_1 = \sum_{j=1}^{N-1} 1/j \ .$$

Unlike θ_W, nucleotide diversity (π) is influenced by allele frequencies, and since the number of chromosomes sampled is not consistent across all genomic loci, we weighted the contribution of a single nucleotide by the single-locus number of chromosomes sampled. We obtained π and the standard error using the following formula, where p_i is the frequency of the major allele at locus i, q_i is the frequency of the minor allele at locus i, n_i is the total count of chromosomes at locus i, and L is the total region length:

$$\pi = \sum_{i=1}^{S} \left(p_i q_i \times \frac{n_i}{N} \right) \text{ and } SE(\pi) = \sqrt{\frac{\theta_W}{3L + 2\theta_W^2/9}} /N \ .$$

To calculate the per-base π and θ_W, we divided these values by L.

We calculated Tajima's D as follows:

$$Tajima's\ D = \frac{\pi - \theta_W}{\sqrt{e_1 S + e_2 S(S-1)}}$$

with constants e_1 and e_2 defined as in the original method[66].

To obtain per-gene metrics and compare the properties of the IDR-coding sequence and the non-IDR-coding sequence within the same gene, we subsetted to proteins with a disorder content between 10–90%. For gnomAD, we used proteins with at least 10 SNVs in each of the protein's IDR and non-IDR regions; for 1KGP, we used a threshold of 1 SNV.

Using biallelic SNVs from the 1KGP, we estimated unbiased pairwise F_{ST} values between European (EUR), East Asian (EAS), and African (AFR) populations as described by Weir and Cockerham[67], using scikit-allel[68]. We used genes with disorder content between 20–80% and compared per-gene IDR F_{ST} values and per-gene non-IDR F_{ST} values for each population pair. We used these pairwise F_{ST} values to estimate locus-specific branch lengths (*LSBLs*) for each population as previously described[69]. For example, the *LSBL* of the EAS population is:

$$LSBL_{EAS} = \left(F_{ST, EAS:EUR} + F_{ST, EAS:AFR} - F_{ST, EUR:AFR} \right)/2$$

Functional Variation and Disease Analyses

To obtain genes associated with lethality, we also used the Lethal Phenotypes Portal[70] (updated November 2024), a catalog of OMIM genes related to lethality. We grouped the Human Phenotype Ontology (HPO)[71] lethality category annotations to assign the following developmental categories: prenatal/neonatal (L1/L2), infancy/childhood (L3/L4), and adolescence/adulthood (L5/L6). We combined mode-of-inheritance (MOI) annotations from these lethal genes with previously published MOI catalogs[72,73]. We used only autosomal dominant (AD) and autosomal recessive (AR) genes and excluded those that were categorized as both AD and AR.

As an orthogonal source of disease-associated mutations, we also used the Human Gene Mutation Database (HGMD) Professional[53] (accessed May 2025). We filtered for SNVs with the highest confidence disease-causing mutation associations (tag=DM). To analyze disease-type biases, we used HGMD disease variants linked to ICD-10[74] disease codes in the HGMD phenbase. ICD-10 chapters were grouped into the following larger disease categories: Neoplasms (C), Blood (D), Endocrine/Metabolic (E), Neuropsychiatric (F, G), Eye/Ear (H), Circulatory (I), Congenital Abnormalities (Q), and Other (J, K, L, M, N).

Data Availability

The UniProt proteome is available at
https://ftp.uniprot.org/pub/databases/uniprot/current_release/knowledgebase/reference_proteo
mes/Eukaryota/UP000005640/UP000005640_9606.fasta.gz. UniProt ID mappings are
available at
https://ftp.uniprot.org/pub/databases/uniprot/current_release/knowledgebase/reference_proteo
mes/Eukaryota/UP000005640/UP000005640_9606.idmapping.gz. The TF list is available at
https://humantfs.ccbr.utoronto.ca/download/v_1.01/DatabaseExtract_v_1.01.csv. The
GenOrigin human gene age catalog can be downloaded at
http://chenzxlab.hzau.edu.cn/GenOrigin/#!/download. The GenEra-produced gene age catalog
can be downloaded at
https://github.com/LotharukpongJS/phylomapr/blob/main/data-raw/Homo_sapiens_gene_ages.
tsv. PhyloP scores were obtained from
https://hgdownload.cse.ucsc.edu/goldenpath/hg38/phyloP100way/hg38.phyloP100way.bw.
The GO association file can be found at
https://current.geneontology.org/annotations/goa_human.gaf.gz. The GO OBO file can be
found at https://purl.obolibrary.org/obo/go/go-basic.obo. The STRING PPI database is at
https://string-db.org/cgi/download. The GRNdb is available at http://www.grndb.com. The
Human Protein Atlas (v25.0) is available at https://www.proteinatlas.org/. GTEx v10 QTLs can
be downloaded from the GTEx Portal at https://gtexportal.org/home/downloads/adult-gtex/qtl.
Translation efficiency data can be found at Zheng et al. (2025) in Supplementary Table 1 at
https://doi.org/10.1038/s41587-025-02712-x . VCF files from gnomAD v4.1 were downloaded
using Google Cloud Public from https://gnomad.broadinstitute.org/data. 1000 Genomes Project
biallelic SNV and indel VCFs (release 2019) were obtained from their FTP site at
/vol1/ftp/data_collections/1000_genomes_project/release/20190312_biallelic_SNP_and_INDE
L. Lethal OMIM genes data are available at the Lethal Phenotypes Portal
https://lethalphenotypes.research.its.qmul.ac.uk/. HGMD® Professional data were obtained
from https://www.hgmd.cf.ac.uk/ac/index.php.

Code Availability

Code to reproduce the results is available at https://github.com/AkeyLab/tf_idr_paper.

Acknowledgements

We would like to thank the members of the Akey laboratory for their helpful comments and feedback on this work, especially Rob Bierman, Cara Weisman, and Aaron Pfennig. We would also like to thank Thouis Ray Jones, Jap Chae, and Jjam Ppong. This work was supported in part by National Institutes of Health grants T32HG003284 (SS) and R01GM110068 (JMA).

References

1. Badia-i-Mompel, P. *et al.* Gene regulatory network inference in the era of single-cell multi-omics. *Nat. Rev. Genet.* **24**, 739–754 (2023).

2. Liu, J. *et al.* Intrinsic Disorder in Transcription Factors. *Biochemistry* **45**, 6873–6888 (2006).

3. Dyson, H. J. & Wright, P. E. Intrinsically unstructured proteins and their functions. *Nat. Rev. Mol. Cell Biol.* **6**, 197–208 (2005).

4. Babu, M. M. The contribution of intrinsically disordered regions to protein function, cellular complexity, and human disease. *Biochem. Soc. Trans.* **44**, 1185–1200 (2016).

5. Babu, M. M., van der Lee, R., de Groot, N. S. & Gsponer, J. Intrinsically disordered proteins: regulation and disease. *Curr. Opin. Struct. Biol.* **21**, 432–440 (2011).

6. Brangwynne, C. P. *et al.* Germline P Granules Are Liquid Droplets That Localize by Controlled Dissolution/Condensation. *Science* **324**, 1729–1732 (2009).

7. Sabari, B. R. *et al.* Coactivator condensation at super-enhancers links phase separation and gene control. *Science* **361**, eaar3958 (2018).

8. Pei, G., Lyons, H., Li, P. & Sabari, B. R. Transcription regulation by biomolecular condensates. *Nat. Rev. Mol. Cell Biol.* **26**, 213–236 (2025).

9. Hnisz, D., Shrinivas, K., Young, R. A., Chakraborty, A. K. & Sharp, P. A. A Phase Separation Model for Transcriptional Control. *Cell* **169**, 13–23 (2017).

10. Li, X., Liu, C., Lei, Z., Chen, H. & Wang, L. Phase-separated chromatin compartments: Orchestrating gene expression through condensation. *Cell Insight* **3**, 100213 (2024).

11. Cho, W.-K. *et al.* Mediator and RNA polymerase II clusters associate in transcription-dependent condensates. *Science* **361**, 412–415 (2018).

12. Basu, S. *et al.* Unblending of Transcriptional Condensates in Human Repeat Expansion Disease. *Cell* **181**, 1062-1079.e30 (2020).

13. Peng, L., Li, E.-M. & Xu, L.-Y. From start to end: Phase separation and transcriptional regulation. *Biochim. Biophys. Acta BBA - Gene Regul. Mech.* **1863**, 194641 (2020).

14. Tsang, B., Pritišanac, I., Scherer, S. W., Moses, A. M. & Forman-Kay, J. D. Phase Separation as a Missing Mechanism for Interpretation of Disease Mutations. *Cell* **183**, 1742–1756 (2020).

15. Afanasyeva, A., Bockwoldt, M., Cooney, C. R., Heiland, I. & Gossmann, T. I. Human long intrinsically disordered protein regions are frequent targets of positive selection. *Genome Res.* **28**, 975–982 (2018).

16. Ahmed, S. S. *et al.* Characterization of intrinsically disordered regions in proteins informed by human genetic diversity. *PLOS Comput. Biol.* **18**, e1009911 (2022).

17. Brown, C. J., Johnson, A. K., Dunker, A. K. & Daughdrill, G. W. Evolution and disorder. *Curr. Opin. Struct. Biol.* **21**, 441–446 (2011).

18. Singleton, M. D. & Eisen, M. B. Evolutionary analyses of intrinsically disordered regions reveal widespread signals of conservation. *PLOS Comput. Biol.* **20**, e1012028 (2024).

19. Khan, T., Douglas, G. M., Patel, P., Nguyen Ba, A. N. & Moses, A. M. Polymorphism Analysis Reveals Reduced Negative Selection and Elevated Rate of Insertions and Deletions in Intrinsically Disordered Protein Regions. *Genome Biol. Evol.* **7**, 1815–1826 (2015).

20. Nilsson, J., Grahn, M. & Wright, A. P. Proteome-wide evidence for enhanced positive Darwinian selection within intrinsically disordered regions in proteins. *Genome Biol.* **12**, R65 (2011).

21. UniProt Consortium. UniProt: a worldwide hub of protein knowledge. *Nucleic Acids Res.* **47**, D506–D515 (2019).

22. Emenecker, R. J., Griffith, D. & Holehouse, A. S. Metapredict: a fast, accurate, and easy-to-use predictor of consensus disorder and structure. *Biophys. J.* **120**, 4312–4319 (2021).

23. Emenecker, R. J., Griffith, D. & Holehouse, A. S. Metapredict V2: An update to metapredict, a fast, accurate, and easy-to-use predictor of consensus disorder and structure. 2022.06.06.494887 Preprint at https://doi.org/10.1101/2022.06.06.494887 (2022).

24. Mehdiabadi, M. *et al.* Critical Assessment of Protein Intrinsic Disorder Round 3 - Predicting Disorder in the Era of Protein Language Models. *Proteins Struct. Funct. Bioinforma.* **n/a**, (2025).

25. Erdős, G. & Dosztányi, Z. AIUPred: combining energy estimation with deep learning for the enhanced prediction of protein disorder. *Nucleic Acids Res.* **52**, W176–W181 (2024).

26. Carvunis, A.-R. *et al.* Proto-genes and de novo gene birth. *Nature* **487**, 370–374 (2012).

27. Wilson, B. A., Foy, S. G., Neme, R. & Masel, J. Young genes are highly disordered as predicted by the preadaptation hypothesis of de novo gene birth. *Nat. Ecol. Evol.* **1**, 1–6 (2017).

28. James, J. E. *et al.* Universal and taxon-specific trends in protein sequences as a function of age. *eLife* **10**, e57347 (2021).

29. Tong, Y.-B. *et al.* GenOrigin: a comprehensive protein-coding gene origination database on the evolutionary timescale of life. *J. Genet. Genomics* **48**, 1122–1129 (2021).

30. Weisman, C. M., Murray, A. W. & Eddy, S. R. Many, but not all, lineage-specific genes can be explained by homology detection failure. *PLOS Biol.* **18**, e3000862 (2020).

31. Moyers, B. A. & Zhang, J. Phylostratigraphic Bias Creates Spurious Patterns of Genome Evolution. *Mol. Biol. Evol.* **32**, 258–267 (2015).

32. Andreatta, M. E. *et al.* The Recent De Novo Origin of Protein C-Termini. *Genome Biol. Evol.* **7**, 1686–1701 (2015).

33. Lotharukpong, J. S. Phylomap of Homo sapiens. Unpublished (2023).

34. Wong, E. T. C. *et al.* Protein–Protein Interactions Mediated by Intrinsically Disordered Protein Regions Are Enriched in Missense Mutations. *Biomolecules* **10**, 1097 (2020).

35. Trifonov, E. N. The Triplet Code From First Principles. *J. Biomol. Struct. Dyn.* **22**, 1–11 (2004).

36. Higgs, P. G. & Pudritz, R. E. A Thermodynamic Basis for Prebiotic Amino Acid Synthesis and the Nature of the First Genetic Code. *Astrobiology* **9**, 483–490 (2009).

37. Lambert, S. A. *et al.* The Human Transcription Factors. *Cell* **172**, 650–665 (2018).

38. Jolma, A. *et al.* DNA-binding specificities of human transcription factors. *Cell* **152**, 327–339 (2013).

39. Lemma, R. B. *et al.* Pioneer transcription factors are associated with the modulation of DNA methylation patterns across cancers. *Epigenetics Chromatin* **15**, 13 (2022).

40. Barral, A. & Zaret, K. S. Pioneer factors: roles and their regulation in development. *Trends Genet.* **40**, 134–148 (2024).

41. Bulyk, M. L., Drouin, J., Harrison, M. M., Taipale, J. & Zaret, K. S. Pioneer factors — key regulators of chromatin and gene expression. *Nat. Rev. Genet.* **24**, 809–815 (2023).

42. Uhlén, M. *et al.* Tissue-based map of the human proteome. *Science* **347**, 1260419 (2015).

43. Oughtred, R. *et al.* The BioGRID database: A comprehensive biomedical resource of curated protein, genetic, and chemical interactions. *Protein Sci.* **30**, 187–200 (2021).

44. Cancer Genome Atlas Research Network *et al.* The Cancer Genome Atlas Pan-Cancer analysis project. *Nat. Genet.* **45**, 1113–1120 (2013).

45. THE GTEX CONSORTIUM. The GTEx Consortium atlas of genetic regulatory effects across human tissues. *Science* **369**, 1318–1330 (2020).

46. Karczewski, K. J. *et al.* The mutational constraint spectrum quantified from variation in 141,456 humans. *Nature* **581**, 434–443 (2020).

47. Patel, A. *et al.* A Liquid-to-Solid Phase Transition of the ALS Protein FUS Accelerated by Disease Mutation. *Cell* **162**, 1066–1077 (2015).

48. Boeynaems, S. *et al.* Protein Phase Separation: A New Phase in Cell Biology. *Trends Cell Biol.* **28**, 420–435 (2018).

49. Murakami, T. *et al.* ALS/FTD Mutation-Induced Phase Transition of FUS Liquid Droplets and Reversible Hydrogels into Irreversible Hydrogels Impairs RNP Granule Function. *Neuron* **88**, 678–690 (2015).

50. Shendure, J. & Akey, J. M. The origins, determinants, and consequences of human mutations. *Science* **349**, 1478–1483 (2015).

51. Balch, W. E., Morimoto, R. I., Dillin, A. & Kelly, J. W. Adapting Proteostasis for Disease Intervention. *Science* **319**, 916–919 (2008).

52. Gerasimavicius, L., Livesey, B. J. & Marsh, J. A. Loss-of-function, gain-of-function and dominant-negative mutations have profoundly different effects on protein structure. *Nat. Commun.* **13**, 3895 (2022).

53. Stenson, P. D. *et al.* The Human Gene Mutation Database (HGMD®): optimizing its use in a clinical diagnostic or research setting. *Hum. Genet.* **139**, 1197–1207 (2020).

54. Conte, A. D. *et al.* Critical assessment of protein intrinsic disorder prediction (CAID) - Results of round 2. *Proteins Struct. Funct. Bioinforma.* **91**, 1925–1934 (2023).

55. Ahmad, S. *et al.* The UniProt website API: facilitating programmatic access to protein knowledge. *Nucleic Acids Res.* **53**, W547–W553 (2025).

56. Nightingale, A. *et al.* The Proteins API: accessing key integrated protein and genome information. *Nucleic Acids Res.* **45**, W539–W544 (2017).

57. Kumar, S. *et al.* TimeTree 5: An Expanded Resource for Species Divergence Times. *Mol. Biol. Evol.* **39**, msac174 (2022).

58. Barrera-Redondo, J., Lotharukpong, J. S., Drost, H.-G. & Coelho, S. M. Uncovering gene-family founder events during major evolutionary transitions in animals, plants and fungi using GenEra. *Genome Biol.* **24**, 54 (2023).

59. Yanai, I. *et al.* Genome-wide midrange transcription profiles reveal expression level relationships in human tissue specification. *Bioinformatics* **21**, 650–659 (2005).

60. Ashburner, M. *et al.* Gene Ontology: tool for the unification of biology. *Nat. Genet.* **25**, 25–29 (2000).

61. Klopfenstein, D. V. *et al.* GOATOOLS: A Python library for Gene Ontology analyses. *Sci. Rep.* **8**, 10872 (2018).

62. Szklarczyk, D. *et al.* The STRING database in 2023: protein-protein association networks and functional enrichment analyses for any sequenced genome of interest. *Nucleic Acids Res.* **51**, D638–D646 (2023).

63. Fang, L. *et al.* GRNdb: decoding the gene regulatory networks in diverse human and mouse conditions. *Nucleic Acids Res.* **49**, D97–D103 (2021).

64. Wolf, S. *et al.* Characterizing the landscape of gene expression variance in humans. *PLOS Genet.* **19**, e1010833 (2023).

65. Zheng, D. *et al.* Predicting the translation efficiency of messenger RNA in mammalian cells. *Nat. Biotechnol.* 1–14 (2025) doi:10.1038/s41587-025-02712-x.

66. Tajima, F. Statistical method for testing the neutral mutation hypothesis by DNA polymorphism. *Genetics* **123**, 585–595 (1989).

67. Weir, B. S. & Cockerham, C. C. Estimating F-Statistics for the Analysis of Population Structure. *Evolution* **38**, 1358–1370 (1984).

68. Miles, A. *et al.* cggh/scikit-allel: v1.3.13. Zenodo https://doi.org/10.5281/zenodo.13772087 (2024).

69. Shriver, M. D. *et al.* The genomic distribution of population substructure in four populations using 8,525 autosomal SNPs. *Hum. Genomics* **1**, 274 (2004).

70. Cacheiro, P. *et al.* Lethal phenotypes in Mendelian disorders. *Genet. Med.* **26**, 101141 (2024).

71. Robinson, P. N. *et al.* The Human Phenotype Ontology: A Tool for Annotating and Analyzing Human Hereditary Disease. *Am. J. Hum. Genet.* **83**, 610–615 (2008).

72. Blekhman, R. *et al.* Natural selection on genes that underlie human disease susceptibility. *Curr. Biol.* **18**, 883–889 (2008).

73. Berg, J. S. *et al.* An informatics approach to analyzing the incidentalome. *Genet. Med.* **15**, 36–44 (2013).

74. World Health Organization. *International Statistical Classification of Diseases and Related Health Problems (ICD-10).* vol. 2 (Geneva, 2019).

Figures

a Transcriptional Condensate
Activation Domain
DNA-binding Domain
TF
Target Gene

b Proteome Annotation
TFs (n=1,612)
92% 8%
UniProt Proteome (n=19,736)
Non-TFs (n=18,124)

Disorder Prediction
IDR   Non-IDR
MetaPredict Disorder Score
1
0.5   Threshold
N   C
(Disorder Content = 54%)

IDR comparison between TFs and non-TFs
Regulatory Networks
Genetic Variation
Disease

c
% of all proteins
TFs 8.2%   Non-TFs 91.8%
19.7%   80.3%
Proteins with 0 IDRs   Proteins with ≥1 IDR

d
MWU $P < 10^{-30}$
TFs
Non-TFs
25%   56%
Density
0.02
0.01
0.00
0   20   40   60   80   100
Disorder Content (%)

e
IDR   Non-IDR
TFs Σ residues = 0.95 M   56.5% (0.54 M)   43.5% (0.41 M)
NonTFs Σ residues = 10.30 M   31.1% (3.21 M)   68.9% (7.10 M)
0   20   40   60   80   100
% of all residues

f
MWU $P = 9.2 \times 10^{-18}$
IDRs per kb
3.7   3.1
TFs   NonTFs

g
MWU $P < 10^{-30}$
TF IDRs
Non-TF IDRs
59 aa   114 aa
Density
1.0
0.5
0.0
10^1 10^2 10^3
IDR segment length (AA)

h IDR Density Along the Protein
Permutation $P < 10^{-30}$
TFs
Non-TFs
Proportion Disordered
1.00
0.75
0.50
0.25
0.00
N-terminus   C-terminus
0   20   40   60   80   100
Relative Position (% of Protein Length)

Fig. 1 | Identifying and characterizing TF and non-TF IDRs. a, Schematic of a TF with a structured DNA-binding domain (DBD) and a disordered activation domain (AD) in a transcriptional condensate (purple circle). The disordered AD recruits multiple regulatory proteins (grey) to the target gene. **b**, Overview of the study workflow. UniProt human proteins were annotated as TFs or non-TFs. Residue-level disorder was predicted using MetaPredict, and IDR features were compared between TFs and non-TFs. **c,** Sankey diagram of the proportion of TFs (red) and non-TFs (black) with at least one predicted IDR (blue) versus no predicted IDRs (grey). **d**, Distribution of the per-protein disorder content among TFs (n=1,581) and non-TFs (n=13,180). Note, this analysis excludes proteins that are either completely ordered (IDR content=0%) or completely disordered (IDR content=100%). Vertical lines indicate group medians. The P value is from a two-sided Mann–Whitney U test. **e**, Fraction of all residues classified as an IDR or non-IDR, shown separately for TFs and non-TFs. M: million. **f**, Number of IDR segments per kb in TF and non-TF sequences. The median TF segment densities are labelled within the violin plots. The two-sided Mann–Whitney U test P value is shown. **g**, Length distribution of individual IDR segments in TFs (n=3,248 IDRs) and non-TFs (n=25,317 IDRs). The P value is from a two-sided Mann–Whitney U test. **h**, Position-dependent IDR density across normalized protein length, for TFs and non-TFs, showing the fraction of proteins disordered at each relative position. The P value reflects a permutation test of whether the disorder content between TFs and non-TFs across the normalized protein is different.

a

Disorder-First Model of Protein Evolution

Young → → Old

b

TFs   Non-TFs

K-W $P < 10^{-30}$ K-W $P < 10^{-30}$

Disorder Content (%)

<100 100–500 500–1000 >1000   <100 100–500 500–1000 >1000

Young ——— Gene Age (Mya) ——— Old

c

Disorder Content

TFs

Non-TFs

Evolutionary Time

d

Positive Residues in IDRs

TFs   Non-TFs

K-W $P < 10^{-30}$ K-W $P < 10^{-30}$

Fraction of IDR residues

<100 100–500 500–1000 >1000   <100 100–500 500–1000 >1000

Young ——— Gene Age (Mya) ——— Old

e

Aromatic Residues in IDRs

TFs   Non-TFs

K-W $P < 10^{-30}$ K-W $P < 10^{-30}$

Fraction of IDR residues

<100 100–500 500–1000 >1000   <100 100–500 500–1000 >1000

Young ——— Gene Age (Mya) ——— Old

f

Pre-metazoa

HMG $n = 30$ 81%
Homeodomain $n = 192$ 78%
Forkhead $n = 49$ 78%
bZIP $n = 53$ 75%
bHLH $n = 105$ 68%

Post-metazoa

Ets $n = 27$ 64%
C2H2 ZF $n = 394$ 60%
KRAB $n = 342$ 33%
Nuclear receptor $n = 46$ 32%

0   20   40   60   80   100
Disorder Content (%)

g

High disorder TFs (>75% IDR)

cell differentiation
embryonic morphogenesis
neuron differentiation
Wnt signaling pathway
proximal/distal pattern formation
digestive tract development

0.0   0.5   1.0
Gene Ratio

TF Gene Count
● 5
● 20
● 50
● 100

Low disorder TFs (<15% IDR)

transport
defense response
cell surface receptor signaling pathway via STAT
cell surface receptor signaling pathway via JAK-STAT
interleukin-9-mediated signaling pathway
host-mediated perturbation of symbiont process
host-mediated perturbation of viral process

0.0   0.5   1.0
Gene Ratio

3.0
2.5
2.0
1.5
$-\log_{10}$(adjusted P)

Fig. 2 | Evolutionarily ancient TFs are more disordered. a, The "disorder first" hypothesis of protein evolution, in which a peptide begins disordered (blue) and acquires secondary structure (grey) over time. **b,** Disorder content stratified by inferred gene age for TFs and non-TFs. Evolutionarily ancient TFs are enriched for disorder, whereas the opposite trend (with low effect size) is seen in non-TFs. *P* values are from the Kruskal–Wallis test within each protein group. White dots denote medians, and lines denote the interquartile range (IQR). **c,** Summary schematic of the observed divergence between TFs and non-TFs: TF disorder content is maintained or ever expanded over time, whereas non-TFs show the opposite trend, consistent with distinct selective pressures on IDRs in regulatory versus non-regulatory proteins. **d–e,** Composition of IDRs across gene ages, showing that the fraction of (**d**) positively charged residues (K, R, H) and (**e**) aromatic residues (F, W, Y) within IDRs varies with age in both TFs and non-TFs (Kruskal–Wallis tests), suggesting that not only the amount of disorder but also the biochemical character of IDRs shifts across evolutionary time. **f,** Disorder content split by the largest TF DNA-binding domain families, organized into pre- and post-metazoan families. Vertical lines denote medians. **g,** GO enrichment of biological processes in TFs at the extremes of disorder content compared to all TFs (*n*=1,008). Highly disordered TFs (*n*=351) are enriched for developmental and differentiation-related processes, whereas low-disorder TFs (*n*=31) are enriched for immune and signaling pathways. Dot size reflects gene count, and color indicates enrichment significance (Benjamini–Hochberg adjusted *P* values).

a

TFs

r = 0.36, P < 10⁻³⁰

Non-TFs

r = −0.14, P < 10⁻³⁰

C2H2 ZF TFs

r = 0.55, P < 10⁻³⁰

Other TFs

r = −0.00, P > 0.05

Disorder Content (%) / TF Disorder Content (%)

Protein Interactors

b

K-W P = 4.0 × 10⁻²⁶

TF Disorder Content (%)

Number of Target Genes

<50 50-250 >250

c

K-W P < 10⁻³⁰

TF Disorder Content (%)

Number of Regulated Tissues

<5 5-15 >15

d

Disordered

TF Disorder Content (%)

Ordered

PPI   GRN   Tissues

Fig. 3 | TF disorder content is associated with regulatory breadth and protein network connectivity. a, Disorder content as a function of protein–protein interaction (PPI) degree for TFs and non-TFs. Interactions include direct interactions and indirect associations from STRING[62]. TF disorder content increases with the number of interactors in its functional association network, whereas non-TFs show a weak negative association. Lines indicate linear fits with 95% confidence bands. Stratifying TFs by DNA-binding domain family shows that this relationship is strongly driven by C2H2 zinc finger (ZF) TFs, while other TF families show no association. **b,** TF disorder content increases with the number of target genes assigned to each TF in GRNs. The Kruskal–Wallis test P value compares bins of target gene size. Per box TF counts from left to right are n=420, 384, 255. **c,** TF disorder content also increases with the number of tissues in which a TF is inferred to be an active regulator (controlling a GRN regulon). The Kruskal–Wallis test P value compares bins of the number of regulated GTEx tissues. Per box TF counts from left to right are n=655, 484, 356. **d,** Conceptual summary: higher TF disorder content is associated with larger protein functional association networks, broader regulatory targeting within GRNs, and activity across more tissues, consistent with TF IDRs supporting regulatory versatility through expanded interaction capability.

a

TFs | Non-TFs

MWU $P < 10^{-30}$ MWU $P > 0.05$

Disorder Content (%)

100 | 75 | 50 | 25 | 0

Specific Broad Specific Broad

Tissue-Specific Expression

b

TFs | Non-TFs

MWU $P = 5.0 \times 10^{-11}$ MWU $P > 0.05$

Disorder Content (%)

100 | 75 | 50 | 25 | 0

Low High Low High

Inter-Individual Expression Variance

c

TFs | Non-TFs

$\rho = -0.32, P < 10^{-30}$ $\rho = -0.08, P = 8.2 \times 10^{-25}$

Disorder Content (%)

100 | 75 | 50 | 25 | 0

1-10 11-20 21-30 31-40 41-50 1-10 11-20 21-30 31-40 41-50

 # GTEx Tissues as eGene

d

TFs | Non-TFs

$\rho = 0.16, P = 4.1 \times 10^{-7}$ $\rho = -0.23, P < 10^{-30}$

Disorder Content (%)

100 | 75 | 50 | 25 | 0

1 2 3 4 5 1 2 3 4 5

Low ← → High

Translational Efficiency Quintiles

e

Per-Tissue/Study

Spearman's ρ

0.4 | 0.2 | 0.0 | -0.2 | -0.4

TFs
Non-TFs

Increases with more disorder

Decreases with more disorder

Expression Variance | eQTLs per tissue | sQTLs per tissue | Translation Efficiency

f

Disordered

TF Disorder Content (%)

Ordered

Expression Variance | Quantitative Trait Loci (QTLs) | Translational Efficiency

Fig. 4 | Constrained regulatory architecture correlates with disorder in TFs but not in non-TFs. a, Disorder content stratified by HPA tissue specificity for TFs (specific TFs n=714; broad TFs n=835) and non-TFs (specific non-TFs n=9,676; broad non-TFs n=7,026). TFs with broader expression show higher disorder content, whereas non-TFs show no significant difference (two-sided Mann–Whitney U tests). **b,** Disorder content stratified by inter-individual expression variance in GTEx[64] for TFs (high-variance TFs n=510; low-variance TFs n=421) and non-TFs (high-variance non-TFs n=5,270; low-variance non-TFs n=5,194); high-variance TFs exhibit increased disorder content, whereas non-TFs show no significant difference (two-sided Mann–Whitney U tests). **c,** Disorder content as a function of the number of GTEx tissues in which a gene is detected as an eGene. Disorder content decreases with increasing eGene breadth for both TFs and non-TFs. Per box sample sizes for TFs from left to right are n=879, 239, 132, 110, and 101; non-TFs: n=7,537, 3736, 2,143, 1,404, and 1,225. **d,** Disorder content across translational efficiency[65] quintiles; TF disorder content increases with translational efficiency, whereas non-TFs show the opposite trend. n=190 per TF quintile and n=1,849 per non-TF quintile. **e,** Per-study association summary. For each study/tissue, Spearman correlations were computed between disorder content and each metric (expression variance, eQTLs per tissue, sQTLs per tissue, and translational efficiency); boxplots summarize the distribution of per-tissue correlation coefficients for TFs and non-TFs, with the dashed line indicating no association. The sample sizes (number of tissues or studies) for each group, from left to right, are n=57, 50, 50, and 78. **f,** Summary of the relationship between TF disorder content and features of the regulation of the TF itself, such as translation efficiency, expression variance, and QTL abundance.

a

b

SNVs

Indels

c

Nucleotide Diversity θ_w

Per-gene

d

Genome-wide

e

Nucleotide Diversity π

Per-gene

f

Genome-wide

i

g

Tajima's D

Per-gene

h

Genome-wide

Fig. 5 | TF IDRs are enriched for polymorphisms. a, TF IDRs are enriched for SNVs and Indels in gnomAD v4.1. Total genomic region, SNV, and indel counts are shown, split between TF Non-IDRs (grey) and IDRs (blue). *P* values were obtained from the binomial exact test. **** indicates *P*<10^{-30}. **b**, SNV and indel rates. kb: kilobase. **c,** Nucleotide diversity θ_W, in TFs (*n*=1,483; left) and non-TFs (*n*=10,182; right) with between 10% and 90% disorder content. θ_W was calculated separately for the protein's IDR and non-IDR regions. The density plot shows that most TFs and most non-TFs have elevated θ_W in their IDRs. **d**, Nucleotide diversity θ_W, calculated for the TF genome and non-TF genome. **e**, Nucleotide diversity π, in TFs (*n*=1,483; left) and non-TFs (*n*=10,182; right), with between 10% and 90% disorder content. π was calculated separately for the protein's IDR and non-IDR regions. The density plot shows that most TFs have increased π in their IDRs. For non-TFs, this bias is small, showing higher π in their IDR regions as well. **f**, Nucleotide diversity π, calculated for the TF genome and non-TF genome. **g**, Tajima's D in TFs (*n*=1,483; left) and non-TFs (*n*=10,182; right) with between 10% and 90% disorder content. Tajima's D was calculated separately for the protein's IDR and non-IDR regions. **h,** Tajima's D calculated for the TF genome and non-TF genome. **i,** EAS-specific LSBL-derived F_{ST} values in TFs (*n*=1,092; left) and non-TFs (*n*=5,992; right). There is a significant skew towards higher F_{ST} values in a TF's IDR region compared to its non-IDR region, which is not observed in non-TFs.

Fig. 6 | Disease associations differ between TFs and non-TFs by disorder content. a, Disorder content of TFs and non-TFs stratified by OMIM lethality annotation. Lethal TFs show higher disorder content than non-lethal TFs, whereas lethal non-TFs show reduced disorder content relative to non-lethal non-TFs (two-sided Mann–Whitney U tests; TF lethal n=110, TF non-lethal n=1,502; non-TF lethal n=1,529, non-TF non-lethal n=16,595). **b**, Among OMIM-lethal genes, disorder content stratified by the earliest reported age of death. TF disorder content is highest for prenatal/neonatal lethality and decreases for later lethality, while non-TFs show the opposite pattern (Kruskal–Wallis tests; TF n=52, 54, 19; non-TF n=514, 810, 412 for prenatal/neonatal, infancy/childhood, and adolescence/adulthood, respectively). **c**, Disorder content stratified by mode of inheritance (autosomal dominant (AD) versus autosomal recessive (AR)). Disorder content does not differ between AD and AR TFs, whereas AD non-TFs show higher disorder content than AR non-TFs (two-sided Mann–Whitney U tests; TF AD n=113, TF AR n=69; non-TF AD n=621, non-TF AR n=1,593). **d**, Enrichment of HGMD disease variants in IDR versus non-IDR coding sequence for TFs and non-TFs. OR<1 (below the dashed line) indicates relative depletion of disease variants in IDRs (TFs: OR=0.298, 95% CI=0.286–0.310; non-TFs: OR=0.495, 95% CI=0.487–0.504). **e**, Disease-category stratified enrichment of HGMD variants in IDRs within TFs and non-TFs. Points indicate ORs with 95% CIs, and filled markers indicate Bonferroni-significant enrichment. The heatmap summarizes ORs across tissue categories (outlined tiles denote Bonferroni significance), ordered by the ΔOR between TFs and non-TFs.

Extended Data Figures

Extended Data Fig. 1 | Additional IDR characteristics and robustness across disorder predictors. a, Distribution of disorder categories for TFs and non-TFs based on the fraction of residues classified as disordered (fully ordered, 0% IDR; mostly ordered, <50%; mostly disordered, >50%; fully disordered, 100%). Protein-length distributions for all proteins (bottom left) and for partially disordered proteins only (bottom right). Vertical lines denote group medians (all proteins: non-TFs 413 aa, TFs 504 aa; partially disordered: non-TFs 496 aa, TFs 508 aa). P values from two-sided Mann–Whitney U tests (all proteins: $P<10^{-30}$; partially disordered: $P>0.05$). **b**, Distribution of the number of IDR segments per protein (0, 1, 2, or ≥3) in TFs and non-TFs. **c**, Mean number of IDR segments across protein-length quantiles (x axis shows quantile medians, aa). Effect estimate for TF status is shown (β=1.39 [1.31–1.48], P=2.1×10^{-24}; model details in Methods). Across almost all quantiles, TFs have a higher mean number of IDR segments despite having similar protein lengths. **d,e,** Agreement between MetaPredict and AIUPred, a disorder predictor that doesn't rely on conservation information, but solely energy estimation, for predicted disorder length (aa; **d**) and disorder content (%; **e**) across proteins; red line indicates y=x. **f,** Per-protein Hamming distance between binary residue-level disorder calls from MetaPredict and AIUPred (fraction of mismatched residues; median=0.04; n=19,736 proteins). **g,** AIUPred disorder content (%) versus gene age (Mya) for TFs and non-TFs. Violins show distributions within age bins; black points and lines show the median and IQR, respectively. Kruskal–Wallis tests are shown. **h,** AIUPred disorder content (%) stratified by regulatory complexity among TFs with available annotations, shown by the number of target genes and the number of tissues with regulatory activity (box plots: center line, median; box, IQR).

Extended Data Fig. 2 | Robustness of the gene age-disorder relationship. a, Disorder content in TFs stratified by gene ages estimated by a previous gene catalog[33] using genEra[58], a homology-detection failure-aware estimator. Predicted phylostrata, ordered from youngest to oldest clade, are shown separately for TFs (left) and non-TFs (right). **b**, Summary comparison of disorder content for "young" (grouping the youngest 10 phylostrata, Eutheria to Vertebrata, approximately <500 Mya) versus "old" (grouping the oldest 8 phylostrata, Cordata to cellular organisms, >500 Mya) genes within TFs and non-TFs. **c–g,** Stratified analyses of the gene age-disorder relationship, using gene ages from GenOrigin (Fig. 2b), controlling for potential confounders. For TFs (left) and non-TFs (right), their disorder content distribution is shown across gene-age bins within tertiles of **(c)** conservation, **(d)** protein length, **(e)** GC content, **(f)** expression level, and **(g)** tissue specificity.

Extended Data Fig. 3 | TF IDR features by DNA-binding domain family. a, Ridgeline distribution of total protein length (outline) and cumulative IDR length per protein (filled) for major TF DBD families (HMG, Homeodomain, Forkhead, bZIP, bHLH, ETS, C2H2 ZF, KRAB, and Nuclear receptor). This shows that the trends in TF disorder content are not just driven by protein length variation. Vertical lines indicate family medians for IDR length (solid line) and total length (dashed line). **b**, Fraction of proteins in each DBD family classified by disorder content: fully ordered (IDR content=0%), mostly ordered (0%<IDR content<50%), mostly disordered (50%<IDR content<100%), and fully disordered (IDR content=100%). **c**, Distribution of the number of distinct IDR segments per protein (0, 1, 2, or ≥3 segments) within each DBD family. **d**, Positional distribution of disorder along TF sequences by DBD family. Proteins were scaled to relative length (0–100%, N-terminus to C-terminus), and the y axis shows the proportion of proteins that are disordered at each position (based on residue-level disorder calls). **e**, IDR lengths by DBD family, shown as the percentage of segments in short (20–100 aa), medium (100–200 aa), and long (>200 aa) categories.

a

High disorder TFs (>75%)

cellular process
developmental process
pos. reg. of biological process
pos. reg. of cellular process
pos. reg. of metabolic process
pos. reg. of macromolecule biosynthetic process
pos. reg. of macromolecule metabolic process
pos. reg. of biosynthetic process
pos. reg. of nucleobase-containing compd. metab. process
pos. reg. of RNA metabolic process
pos. reg. of DNA-templated transcription
pos. reg. of RNA biosynthetic process
cellular developmental process
pos. reg. of transcription by RNA polymerase II
anatomical structure development
cell differentiation
multicellular organismal process
reg. of developmental process
anatomical structure morphogenesis
reg. of multicellular organismal process
animal organ development
response to stimulus
reg. of cell differentiation
pos. reg. of developmental process
response to chemical
system development
reg. of cell population proliferation
reg. of multicellular organismal development
pos. reg. of multicellular organismal process
cell development
pos. reg. of cell differentiation
pattern specification process
animal organ morphogenesis
embryonic morphogenesis
neuron differentiation
regionalization
cell population proliferation
cell surface receptor signaling pathway
response to endogenous stimulus
gland development
reg. of neuron differentiation
embryonic organ morphogenesis
skeletal system morphogenesis
skeletal system development
central nervous system neuron differentiation
embryonic skeletal system morphogenesis
pos. reg. of neuron differentiation
Wnt signaling pathway
proximal/distal pattern formation
digestive tract development

GeneRatio

Low disorder TFs (<15%)

cell surface receptor signaling pathway via STAT
cell surface receptor signaling pathway via JAK-STAT
interleukin-9-mediated signaling pathway

0.0   0.5   1.0
GeneRatio

TF Gene Count
• 5
• 20
• 50
• 100

5   10   15   20   25   30
$-\log_{10}$(adjusted P)

b

TFs   Non-TFs
MWU $P < 10^{-30}$ MWU $P = 3.4 \times 10^{-9}$

Disorder Content (%)

Developmental Other   Developmental Other
($n = 748$) ($n = 864$)   ($n = 4,659$) ($n = 13,465$)

c

MWU $P = 2.7 \times 10^{-4}$

Disorder Content (%)

Pioneer TFs   Non-Pioneer TFs

Extended Data Fig. 4 | Functional specialization of high- versus low-disorder TFs. a, GO Biological Process enrichment for TFs with high disorder (IDR content>75%; left) and low disorder (IDR content <15%; right). Dot plots show enriched GO terms (*y* axis) versus gene ratios (*x* axis). The full TF list was used as the background gene set. Dot color indicates the −$\log_{10}$ Benjamini-Hochberg-adjusted *P* value, and dot size indicates the number of TFs annotated to each term. **b**, Disorder content (%) for TFs (left) and non-TFs (right) stratified by annotation to the GO term "developmental process" (GO:0032502). **c**, Disorder content (%) for pioneer versus non-pioneer TFs. Black boxes indicate the median and interquartile range. The two–sided Mann–Whitney U test *P* value is shown. Pioneer TFs were obtained from a previously published dataset[39].

Extended Data Fig. 5 | Robustness of the relationship between disorder content and PPI network size. a, The PPI-disorder relationship persists when disorder content is measured as the total number of disordered residues per protein rather than as a percentage. We compare TFs (top left; red) to non-TFs (top right; black) and C2H2 zinc finger TFs (bottom left) to all other TFs (bottom right), indicating that the TF disorder-PPI trend is driven by C2H2 zinc finger TFs. **b,** The PPI-disorder relationship remains when different confidence level thresholds are applied to STRING networks. This shows that the relationship is not purely driven by low-confidence functional links. Disorder content (%) across quartiles of per-protein functional association partner size, shown for TFs and non-TFs, and repeated across STRING confidence thresholds (medium>400, high>700, highest>900). Disorder content (%) across quartiles of PPI size constructed using STRING physical (binding) interactions only. **c–d,** The PPI-disorder relationship remains when using different PPI databases. **(c)** Disorder content and quartiles of protein physical interactions estimated from the HPA database. **(d)** Disorder content and quartiles of protein physical interactions estimated from the BioGRID[43] database.

Extended Data Fig. 6 | Robustness of associations between disorder content and regulatory breadth across independent datasets and specificity metrics. a, Disorder content (%) versus the number of target genes in TCGA-derived GRNs. TFs are stratified by the number of inferred target genes per TF (≤50, 51–250, >250). **b,** Disorder content (%) versus regulatory breadth across tissues (as in Fig. 3b) but in TCGA GRNs instead of GTEx GRNs. TFs are stratified by the number of tissues in which a TF is inferred to regulate tissue-specific expression programs (≤5, 6–15, >15). **c,** Disorder content (%) for TFs and non-TFs stratified by expression breadth across tissues, as annotated by HPA. **d,** Disorder content (%) for TFs and non-TFs stratified by specific versus broadly expressed genes (as in Fig. 4b), but in cell-type specificity instead of tissue specificity, as annotated by HPA. **e,** Disorder content (%) stratified by eQTL burden, defined as the maximum number of eQTL credible sets assigned to a gene within any single tissue, shown for TFs (left) and non-TFs (right). **f,** Disorder content (%) stratified by the number of GTEx tissues in which a gene is identified as an sGene (gene regulated by a splicing-QTL), shown for TFs (left) and non-TFs (right). **g,** Disorder content (%) stratified by sQTL (splicing QTL) burden, defined as the maximum number of sQTL credible sets assigned to a gene within any single tissue, shown for TFs (left) and non-TFs (right).

Extended Data Fig. 7 | Nucleotide diversity and population differentiation in IDRs versus non-IDRs using 1000 Genomes. a, TF IDRs are enriched for SNVs and Indels in the 1000 Genomes Project. Total genomic region, SNV, and indel counts are shown, split between TF Non-IDRs (grey) and IDRs (blue). **b,** SNV and indel rates. **c–d,** Nucleotide diversity (θ_W) for IDRs and non-IDRs. **c,** Per-gene θ_W estimates for TFs (red) and non-TFs (black) with at least 10% IDR and less than 90% IDR. Density plots compare IDR versus non-IDR values; percentages indicate the fraction of genes with higher diversity in IDRs versus non-IDRs. P values are from two-sided Wilcoxon signed-rank tests comparing paired IDR and non-IDR estimates within genes. **d,** Genome-wide θ_W in IDRs and non-IDRs for TFs and non-TFs (top), with the corresponding genome-wide difference ($\Delta\theta_{W=}\theta_{W, IDR} - \theta_{W, Non-IDR}$; bottom). **e–f,** Nucleotide diversity (π) for IDRs and non-IDRs. **(e)** Per-gene π comparisons for TFs and non-TFs with paired Wilcoxon tests and proportions of genes with higher π in IDRs. **(e)** Genome-wide π in IDRs and non-IDRs (top) and the genome-wide difference ($\Delta\pi$; bottom) for TFs and non-TFs. **g–h,** Tajima's D in IDRs and non-IDRs. **(g)** Per-gene comparisons of Tajima's D in IDRs versus non-IDRs for TFs and non-TFs with paired Wilcoxon tests and proportions of genes with higher values in IDRs. **(h)** Genome-wide Tajima's D in IDRs and non-IDRs (top) and the genome-wide difference (ΔD; bottom) for TFs and non-TFs. **i–j,** Distributions of per-gene differences in population differentiation between IDRs and non-IDRs ($\Delta F_{ST} = F_{ST, IDR} - F_{ST, Non-IDR}$) for TFs ($n$=1,092) and non-TFs ($n$=5,992) comparing **(i)** Europeans (EUR) and **(j)** Africans (AFR). Vertical lines denote zero difference; percentages and Wilcoxon P values indicate the skew towards IDRs or non-IDRs.