1 **Nanobody MET CAR-T cells show efficacy in solid tumors**
2

Po-Han Chen[1], Qin Li[1], Sam Deveraux[2], Danielle K. Sohai[1], Pei-Chun Cha[1], Rianna

Raghunandan[3], Nancy Chen[3], Yao Li[1], Mileena Nguyen[1], Michael C. Stankewich[1], Jon S.

Morrow[1,4], Arnaud Augert[1,4], Qin Yan[1,4], and Samuel G. Katz[1,4,*]

Affiliations:

1. Department of Pathology, Yale School of Medicine, New Haven CT, USA

2. Division of Surgery and Interventional Science, University College London, London, UK

3. Yale College, Yale University, New Haven CT, USA

4. Yale Cancer Center, Yale School of Medicine, New Haven, CT, USA

Key words: MET, chimeric antigen receptor, immunotherapy, nanobody, mRNA

*Correspondence: Samuel.Katz@yale.edu

Samuel G. Katz, M.D., Ph.D.

Yale University School of Medicine

310 Cedar Street

Lauder Hall 315C

New Haven, CT 06520

Office: 203-785-2757

Fax: 203-785-3583

Abstract

Background: MET overexpression is associated with poor prognosis in many solid tumors due to its central role in tumor survival, invasion, metastasis, and chemoresistance. While targeting MET with antibody-drug conjugates has shown promising results, engineered cellular immunotherapeutic approaches have not been extensively explored. Compared to conventional single-chain variable fragments (scFv), naturally occurring single-domain antibodies consisting of variable heavy chains only (VHH or nanobodies) are smaller, retain high specificity, and exhibit remarkable biochemical stability. In this study, we tested the efficacy of MET-targeting VHH-CAR-T (chimeric antigen receptor T cells).

Methods: We generated a panel of VHH-CAR-Ts using mRNA electroporation. VHH-CAR-T cells were evaluated in functional assays including cell binding avidity, cytokine production profiles, hydrogel microwell-based cellular kinetics, and in vitro cytotoxicity. We also assessed the therapeutic efficacy of VHH-CAR-T in an in vivo mouse model of metastatic triple negative breast cancer (TNBC).

Results: Among the tested VHH, we identified those with intermediate avidity as most effective for in vitro tumor killing. VHH-CAR-Ts with CD28 costimulatory domains demonstrated augmented cytotoxicity with favorable selectivity, requiring a minimum antigen density threshold for activation. Mechanistically, VHH-CAR-Ts demonstrated low tonic signaling, high avidity, potent cytokine production, and rapid tumor killing kinetics. When administered in an mRNA format, VHH-CAR-Ts exhibited potent and prolonged control of tumor growth in an in vivo metastatic model of TNBC.

Conclusion: Taken together, these results demonstrate that VHH-CAR-Ts exhibit robust MET specificity and potent therapeutic efficacy both in vitro and in vivo. Thus, VHH-CAR-T cell therapy represents a promising immunotherapeutic strategy for targeting MET-overexpressing solid tumors.

55 **What is already known on this topic:** MET signaling is an important contributor to the
56 aggressiveness of many solid tumors, and targeting MET by antibody-drug conjugates has
57 shown efficacy and safety. Targeting MET by CAR-T cells has been under study, though with
58 limited potency.
59
60 **What this study adds:** This study is the first to demonstrate effectiveness of anti-MET VHH-
61 CAR-T cells. Compared with other antigen binding domains, VHH-incorporated CAR-T cells
62 show low tonic signaling, a favorable cytokine profile, and potent tumor killing.
63
64 **How this study might affect research, practice or policy:** With the multiple advantages of
65 VHHs including small size, stability, and low potential for tonic signaling, VHH-CAR-T cells
66 represent a promising approach for CAR-T design against solid tumors.
67
68

INTRODUCTION

While chimeric antigen receptor (CAR) engineered T cells have shown impressive efficacy in hematologic malignancies, solid tumors pose many challenges. Target selection remains a challenge for solid tumors due to the lack of tumor-specific antigens[1]. Innovative CAR designs are also needed to promote tumor infiltration and persistence in the hostile tumor microenvironment, to overcome heterogeneous antigen expression, and to avoid on-target/off-tumor cytotoxicity.

The *MET* oncogene encodes the MET receptor tyrosine kinase, which is overexpressed in many solid tumors and is associated with poor prognosis due to its central role in mediating drug resistance and pro-survival signaling[2]. While tyrosine kinase inhibitors are available, their efficacy is limited to a select number of MET-addicted tumors with MET amplification or mutations[2]. In contrast, targeting MET expression with surface antibodies such as antibody-drug conjugates (ADCs) has shown safety and copy number/mutation-independent efficacy[3]. As CARs leverage the antigen recognition ability of antibodies with the additional benefits of adaptive T cell responses, targeting MET overexpression using CAR-T cells holds great potential.

The antigen recognition domain affects the target sensitivity, durability, and tonic signaling of CAR-T cells[4,5]. Most CARs use a single-chain variable fragment (scFv) as the antigen binding domain, which commonly exhibits self-clustering and tonic signaling leading to T cell exhaustion[6,7].

A variable region of a heavy-chain-only antibody (VHH or nanobody) is emerging as an alternative to scFv[8–10]. VHH-based CARs have been successfully used in BCMA-CAR-T cell therapy ciltacabtagene autoleucel (cilta-cel) against myeloma with approval from the U.S. Food and Drug Administration (FDA) [11]. Of particular importance for solid tumors, the stability of VHH helps avoid receptor clustering and tonic signaling; its uniquely long complementarity-determining region 3 (CDR3) enables targeting of cryptic epitopes and isoforms that may overcome tumor heterogeneity; its compact size allows flexible integration with other moieties (such as tandem VHH in cilta-cel) to facilitate innovative CAR designs with

enhanced functionality; finally, its high degree and ease of humanization reduce

immunogenicity and ensure safety in humans[8,11–15].

In this study, we constructed a panel of mRNA VHH-CAR-T cells targeting MET, and

demonstrate that VHH-CAR-T cells exhibit high avidity, potent cytokine production upon

stimulation, a potential therapeutic window for CAR activation, low tonic signaling, and

effective control of metastatic tumors in vivo – features that warrant further exploration as a

novel form of T cell immunotherapy for solid cancers.

MATERIALS AND METHODS

Cells cultures: Breast cancer cell line MDA-MB-231 was purchased from ATCC (HTB-26),

and its metastatic sub-population LM2 was a kind gift from J. Massagué. The non-small cell

lung cancer cell line A549 cell line was obtained from Peter Glazer, and the head and neck

squamous cell carcinoma cell lines FaDu and CAL27 cell lines were obtained from Barbara

Burtness. These cells were cultured in DMEM+10% FBS+1% penicillin/streptomycin. The

renal cancer cell line 786-O was obtained from Harriet Kluger and cultured in RPMI1640+10%

FBS+1% penicillin/streptomycin. The diffuse large B-cell lymphoma cell lines Karpas-422 and

OCI-LY3 were obtained from Markus Müschen and cultured in RPMI1640+20% FBS+1%

penicillin/streptomycin. Small-cell lung cancer cell lines NCI-H146 (ATCC HB173) and NCI-

H1341(ATCC CRL-5864) were cultured in DMEM-F12 supplemented with 10% FBS, 1X

Insulin-Transferrin-Selenium and 1% penicillin/streptomycin.

Construction of CAR and _in vitro_ transcription: The CAR construct consists of a T7

promoter, 5'UTR (human beta-globin), CAR, P2A self-cleavable sequence, GFP, and 3'UTR

(human beta-globin) (see **Supplemental Figure 1A**). Binding domains for the nanobodies

(VHH1-5), 1E4 (scFv1) and 5D5 (scFv2) sequences were adapted from patents

WO2012/042026[16], US10116622B2[17] and US20190085063A1[18], respectively. The

scFv3 and NK1-CAR sequences were adapted from previous publications [19–21]. The CAR

sequences were synthesized as gene blocks containing T7 promoter and 5' and 3' UTR

sequences. They were cloned into standard pUC57 plasmid using the InFusion cloning kit (Takara).

For *in vitro* transcription, a PCR product encompassing the T7 promoter and 3' UTR, where the reverse primer also added a 100 nucleotide poly-A tail[22], was used in the T7 ARCA HighScribe mRNA Kit with Tailing Reaction (NEB, Cat. #E6020S). Final mRNA was purified using lithium chloride precipitation and concentrated to 1-2 μg/μL in water.

Isolation and activation of primary human T cells: Human peripheral blood mononuclear cells were collected from healthy donors (either local draws under IRB protocol HIC2000033275 or from the Gulf Coast Regional Blood Center). Bulk T cells were isolated using the EasySep Human T Cell Enrichment Kit (STEMCELL Technologies, Cat. #17951), activated with ImmunoCult Human CD3/CD2/CD28 T cell activator (STEMCELL, Cat. #10990) at the recommended concentration, and cultured at 1 x 10^6 cells per mL in RPMI1640 + 10% FBS in the presence of 100 IU/mL recombinant human IL-2 (Miltenyi Biotec, Cat. #130-097743).

Production of mRNA CAR-T cells: On days 7-10 post T cell activation, T cells were pelleted, washed, and resuspended in 20 μL MaxCyte buffer per 2 million cells. Electroporation was performed using the MaxCyte ATX with the Expanded T cell Protocol 1 at 1 μg mRNA/1 million T cells. Following electroporation, cells were rested in a 37^0C incubator with 5% CO_2 for 10 minutes and resuspended in complete T cell culture media at 2x10^6 cells/mL with 100 IU/mL IL-2. Cells were either frozen after 4 hours or used for experiments after overnight incubation. Successful mRNA electroporation was confirmed by GFP expression using either fluorescence microscopy or flow cytometry.

Flow cytometry: Immunophenotypes of all samples were analyzed using CytoFLEX S (Beckman Coulter, CA, USA) flow cytometers. FlowJo software (Tree Star, OR, USA) was used for data analysis. The following antibody-conjugated fluorophores were used: CD3

AlexaFlour 700 (1:100, clone OKT3, BioLegend 317340); CD8 BUV711 (1:100, clone RPA-T8, BioLegend 301044); CD4 BUV605 (1:100, clone OKT4, BioLegend 317438); CD69 PE/Cy5 (1:100, clone FN50, BioLegend 310908); TNF-α PE (1:50, clone MAb11,BioLegend 502909); IFN-γ PerCP/Cy5.5 (1:100, clone XMG1.2, BioLegend 505822); IL-2 BV785 (1:50, clone MQ1-17H12, BioLegend 500348); IL-4 PE/Dazzle 594 (1:100, clone MP4-25D2, BioLegend 500832); PD-1 PE/Cy7 (clone A17188B, BioLegend 621616); TIM-3 APC (clone A18087E, BioLegend 364804); LAG3 APC/Fire750 (clone 7H2C65, BioLegend 369214). MET expression was confirmed by flow cytometry with a monoclonal anti-MET antibody (1:10, Roche, clone SP44) compared to an IgG isotype control (BioLegend 400120).

Gene expression database analysis: Gene expression data for MET across human small cell lung cancer (SCLC) cell lines were obtained from the SCLC-CellMiner database using the SCLC Global response cell line set[23]. Z-scores corresponding to MET expression values were extracted and visualized as a heatmap using GraphPad Prism to compare relative expression levels across cell lines.

Western blot: Protein samples were resolved on 4–20% Mini-PROTEAN TGX precast polyacrylamide gels (15-well, 15 µl; Bio-Rad, #4561096) and transferred to nitrocellulose membranes. Membranes were blocked in 5% non-fat milk in Tris-Buffered Saline with 0.1% Tween-20 (TBST) and incubated with primary antibodies overnight at 4°C (Beta-Actin, 1:5,000, GTX629630; MET, 1:1,000, Cell Signaling Technology #4560). After washing, membranes were incubated with HRP-conjugated secondary antibodies (1:3,000; Cell Signaling Technology; anti-rabbit IgG #7074, anti-mouse IgG #7076) in 5% non-fat milk in TBST for 2 hours at room temperature. Blots were developed using an ultra-sensitive enhanced chemiluminescent (Thermo Scientific, #34095) and imaged on Bio-Rad ChemiDoc imaging system.

Immunohistochemistry (IHC): 4-μm sections were obtained from formalin-fixed, paraffin-embedded (FFPE) tissue. Anti-MET antibody (clone SP44, AbCam, ab227637) was manually applied at 1:50 dilution after antigen retrieval using Tris-EDTA buffer at pH 9.0. Anti-human CD3 antibody (Biocare Medical, CP215) was applied at 1:75 dilution manually following vendor protocol. IHC quantification was performed by visual examination of multiple high power fields for % tumor positive for MET (all intensity).

CD69 activation assay for CAR-Jurkat: The protocol was similar to that previously described [24]. Briefly, 30,000 target cells per well were plated in a 96-well plate. Jurkat cells were electroporated with CAR mRNA and incubated at 37^0C for 4 hours before adding to target cells at various E:T ratios (no target, 100:1, 10:1, 1:1). After overnight incubation, Jurkat cell activation was measured by surface expression of CD69 using flow cytometry.

In vitro tumor killing assays: The flow-based killing assay was similar to that previously described[25] with the following modification. Tumor cells labeled with 1 μM CellTrace Far Red (Invitrogen C34572) were seeded in 24-well plates at a concentration of 1×10^5 cells/well. Mock or CAR-T cells were added to the culture at different ratios (E:T of 0.5:1, 1:1, 2:1, 4:1) without adding exogenous cytokines. Cells were analyzed 16 hours later to measure residual tumor cells by flow cytometry. Dead cells were identified by Live/Dead Violet staining (Invitrogen L34955). Tumor killing was calculated using the formula: % specific lysis = (1-# residual target cells in treated group/# target cells cultured alone without T cells) x100%.

The luciferase-based killing assay was performed using cells expressing firefly luciferase (MDA-MB-231, FaDu, and CAL27). Tumor cells were plated with CAR-T cells at the desired E:T ratio in 384-well plates in 30 μL media overnight. Luminescence was detected using the ONE-Glo Luciferase Assay System (Promega E6120) and analyzed using a plate reader (BioTek, SYNERGYMx). Cell lysis (%) = (1 - (experimental readout - spontaneous readout)/spontaneous readout) x 100.

CellCyte tumor proliferation assay: A lentiviral plasmid encoding mKate2 with puromycin selection was obtained from Dr. David Fox at the University of Michigan. A549 tumor cells expressing mKate2 were co-cultured with T cells in 96-well optically clear flat-bottomed plates at an E:T ratio of 3:1. Plates were imaged using the CellCyte system (ECHO) every 4 hours, and the number of mkate2-positive tumor cells was counted by the provided software.

Mouse tumor xenograft: Six-week-old female NOD.Cg-Prkdc^{scid} Il2rg^{tm1Wjl}/SzJ (NSG) mice were purchased from the Jackson Laboratory (Strain #005557). Tumor inoculation was performed via tail vein injection of $2x10^5$ cells in 100 μL PBS when the mice reached 9-10 weeks of age. Once injected tumors reached a total flux between 10^7 to 10^8 photons/sec by bioluminescence imaging (BLI) (~7-10 days), mice were randomized into respective groups and given weekly injections of PBS or T cells as specified in the text. To avoid graft-versus-host disease, the day before each subsequent T cell injection, 60 mg/kg cyclophosphamide was injected intraperitoneally to deplete the previously injected T cells [26]. Tumor burden was monitored by weekly BLI using the Xenogen *In Vivo* Imaging System (IVIS) (Caliper Life Sciences) and analyzed with Living Image software. Animals were euthanized once luminescence signals reached $1x10^9$ photons/sec or when animals developed significant stress.

Cell force binding assay (avidity): CAR-tumor binding force was measured using the z-Movi device (LUMICKS, Amsterdam, The Netherlands). Target cells were seeded onto microfluidic chips at a concentration of $5x10^7$ cells per mL to obtain a confluent monolayer. CAR-T cells were labeled with 1 μM CellTrace Far Red to enable T cell tracking. After co-incubation of CAR-T cells with the tumor monolayer for 2.5-5 minutes, a force ramp ranging from 0 to 1000 pN was applied for 2.5 minutes. T cell detachment was detected by fluorescence imaging. The binding force was analyzed using the Oceon software (V.1.0, LUMICKS).

Cytokine release assay: For intracellular cytokine staining assays, T cells and target cells were co-cultured at a 1:1 E:T ratio for 5-12 hours in the presence of 2 μM brefeldin A (BD GolgiPlug, cat # 51-2301KZ). For the bulk cytokine assay, the supernatants from overnight co-culture were harvested and stored in -20^0C before shipping to Eve Technologies.

For the multiplexed secretome assay, after stimulation with target cells, approximately 30,000 magnetically enriched CD8+ CAR-T cells were processed for membrane staining (AF647-CD8, STAIN-1002-1). Subsequently, these cells were loaded onto the IsoCode chip (IsoPlexis, ISOCODE-1001-4), comprising 12,000 chambers prepatterned with an array of 32 cytokine capture antibodies. The chip was then incubated in the IsoLight machine for 16 hours at 37°C with 5% CO_2 supplementation. A cocktail of detection antibodies was then applied to detect the secreted cytokines, followed by fluorescence labeling. The resulting fluorescence signals were analyzed using IsoSpeak v.2.8.1.0 (IsoPlexis) to determine the number of specific cytokine-secreting cells and the intensity level of each cytokine.

Structural modeling: The structures of the VHH sequences, scFv1 (1E4) and scFv2 (5D5), and their respective complexes with the human MET extracellular domain were predicted using AlphaFold3[27]. Antibody-MET affinity was estimated using the HADDOCK PRODIGY server [28,29]. Structural modeling and analysis were performed using PyMOL (Schrödinger, LLC).

Microwell co-culture system: We used the TROVO system (Enrich Biosystems) for the microwell assays. For the tumor killing and T cell proliferation assay, T cells (stained with CD4/CD8-Alexa647) and MDA-MB-231 cells were plated at a 2:1 effector to target ratio in a six-well culture plate, with each well containing 3,200 printed hydrogel microwells. Each microwell was indexed by proprietary software and imaged every 12 hours over a four-day period for downstream analysis. Tumor killing was defined as a ≥15% reduction in tumor area compared to day 0. T cell proliferation was defined as a ≥2-fold increase in stained T cell area

compared to day 0. For the single cell killing assay, CAR-T cells and tumor cells were plated

at a 1:10 E:T ratio, and only microwells containing a single T cell were selected for further

imaging analysis. The percentages of microwells with tumor killing/T cell proliferation, and rate

of killing were calculated by the proprietary software built into the TROVO system.

In detail, percentage of microwells with tumor killing or T cell proliferation is calculated as

number of microwells meeting the killing/proliferation threshold stated above divided by total

number of microwells in this well. Bar graph is average +/- standard deviation of 3 technical

replicate wells.

Single cell killing rate was estimated by fitting the paired T cell and tumor signals (area or

counts) at each time point to a mass-action interaction model. For each CAR-T construct (and

matched controls), we ran three independent 3,200-well microwell plates (9,600 wells/clone

total). Dilute seeding targeted ~1 T cell and ~10 tumor cells per well. Plates were imaged from

0–96 h; fluorescence at time zero was used to calibrate area-to-cell conversions. Tumor-only

wells on the same plates were fit to an exponential model $C(t) = C_0 e^{[rC \cdot t]}$ to estimate the

plate-level tumor growth rate rC. From the full set, we retained ~800–900 co-culture wells per

sample that met pre-specified QC (confirmed single T at time zero, adequate tumor baseline,

stable focus/tracking, and finite parameter estimates). An effective killing rate (k_eff) was then

obtained per well using a simplified Lotka–Volterra–type predator–prey coupling $dC/dt = rC \cdot C$

$- k\_eff \cdot T \cdot C$ (classical LV prey equation) and the same mass-action kill term described in

similar CAR-T models[30], i.e., tumor loss regressed against cumulative T cell exposure. Fits

were performed by nonlinear least squares with plate-specific rC; Fitting results with

goodness-of-fit $R^2 > 0.95$ are included. For each CAR-T construct, the distribution of k_eff was

summarized per plate, and the construct-level average k_eff and standard deviation was

calculated across the triplicated experiments.

Statistics: All statistical analyses were conducted using Prism v.10 (GraphPad). Unless

otherwise specified, data were presented as mean ± SD. Statistical significance was

determined by paired t-test, one-way analysis of variance (ANOVA), or two-way ANOVA as

indicated in the figure legends. The log-rank (Mantel-Cox) test was used to determine

statistical significance for overall survival in mouse experiments.

RESULTS

VHH-CAR-T with intermediate avidity showed the most potent tumor killing

We curated five camelid-derived VHH nanobody sequences and incorporated them into a

second-generation mRNA CAR design with a CD8 hinge and transmembrane domain,

followed by 4-1BB and CD3ζ signaling domains (**Supplemental Figure 1 and Figure 1A**).

Since avidity represents the strength of the immune synapse and has been shown to

predict clinical response[31], we used the z-Movi system to measure the avidity of different

VHH-CAR-T cells. To ensure that avidity measurement was not confounded by different CAR

expression levels, the amount of electroporated mRNA was adjusted so that all constructs

had >85% expression as determined by GFP; however, VHH5-CAR was consistently

expressed at ~75-80% regardless of amount of mRNA input (**Figure 1B**).

All VHH-CAR-Ts showed varying percentages of bound T cells with increasing rupture

force (**Figure 1C**). VHH1, VHH2, and VHH3 showed relatively low avidity compared to VHH4

and VHH5. Despite lower expression, VHH5 surprisingly showed the highest avidity. The

mean percentage of remaining T cells bound to targets at 1000 pN, from the lowest to highest,

was VHH3 (7%), VHH1 (12%), VHH2(19%), VHH4 (31%), and VHH5 (36%) (**Figure 1D**).

To gain structural insight into functional avidity, we modeled the binding of the different

VHHs to MET using AlphaFold3 (**Figure 1E**). VHH1, VHH2, and VHH5 are predicted to

overlap with the high affinity binding site for the HGF-α chain. VHH3 is predicted to bind to a

more membrane-distal epitope that overlaps with HGF-β chain. VHH4 appears to bind to a

distinct epitope separate from the other VHHs, slightly away from either HGF binding site.

These predictions are consistent with the reported antagonistic effect of these VHHs on the

HGF-MET interaction. The dissociation constant K_D was estimated by PRODIGY for the top-

ranked structures. The affinity for VHH3 ranged from ~10-50 nM; VHH1 and VHH2 showed

intermediate affinity, averaging 5-10 nM; VHH4 and VHH5 had the highest affinity, ranging

from 0.4-1.2 and 2-3 nM, respectively. The estimated affinities correlated with the observed

avidity, suggesting a direct relationship between affinity and immune synapse strength.

We then investigated whether different avidities correlate with the in vitro cytotoxic

functions of effector cells, expecting higher avidity to achieve better killing of MDA-MB-231

cells. Surprisingly, we found that lower avidity CARs mediated stronger tumor killing, with

VHH1 and VHH2 achieving the highest level of cytotoxicity (**Figure 1F**). VHH3 showed the

lowest avidity, but achieved slightly better killing than VHH4 and VHH5. The superior tumor

killing by VHH1, VHH2, and VHH3 was most notable at low E:T ratios. There was no significant

difference in cytotoxicity between VHH1 and VHH2. Consistently, we showed that both VHH1

and VHH2 strongly inhibit proliferation of A549 lung cancer cells over a 70-hr period using the

CellCyte system (**Figure 1G**).

Although high-avidity CARs are generally favored and have been shown to correlate with

clinical response, our findings are consistent with emerging evidence documenting potential

advantages of lower-avidity CARs (see Discussion). Given that there was no significant

difference between VHH1 and VHH2 in terms of avidity and in vitro cytotoxicity, we used VHH2

for further characterization.

CD28 costimulatory domain conferred stronger VHH-CAR-T activation than 4-1BB

Using VHH2, we next investigated how different cytoplasmic costimulatory domains,

namely,   CD28   versus   4-1BB,   affect   VHH-CAR-T   function.

Although not significantly different, VHH2-28z trends towards higher avidity than VHH2-

BBz (**Figures 2A and B**). VHH2-28z also achieved higher in vitro tumor killing and more

immune cytokine (IFN-γ, TNF-α, IL-2, and IL-4) and cytotoxic molecule Granzyme B

production, as well as higher polyfunctionality (**Figures 2C-E, and Supplemental Figure 2**).

In addition to these fixed-time point assays, we also used a microwell-based co-culture

platform to gain insights into the kinetics of CAR-T cell killing. In microwells where we seeded

1 effector T cell per 10 tumor cells (1:10 E:T), we found that VHH2-28z demonstrated a higher

single-cell killing rate than VHH2-BBz (**Figure 2F**). At an experimental setup of a 20:10 E:T

ratio, we observed more tumor killing and higher T cell proliferation associated with VHH2-28z

compared to VHH2-BBz over a four-day period (**Figures 2G-I**). Although CD28 costimulatory

domain confers immediate potency, it is reported to be more prone to exhaustion than 4-

1BB[32]. However, most likely due to the transient nature of mRNA CAR expression, we did

not observe significant increase in exhaustion markers PD-1, LAG3, or PD-1+/TIM3+ for

VHH2-28z (**Supplemental Figure 3**).Our data is consistent with higher and more rapid

cytotoxic function generally reported for CD28 compared to 4-1BB[32,33]. Beyond four days,

most likely due to diminishing mRNA, there was no significant killing or proliferation observed

with either construct (data not shown). In subsequent studies, we used VHH2-28z unless

otherwise specified.

VHH2-CAR-T eliminated MET+ tumor cells in vitro and in vivo

Since MET overexpression is associated with poor prognosis in many solid tumors, we

tested the efficacy of VHH2-CAR-T against different subtypes of solid tumors with known MET

overexpression. VHH2-CAR-T demonstrated cytotoxicity against tumor cell lines from TNBC

(MDA-MB-231 and LM2), clear cell renal cell carcinoma (786-O), and head and neck

squamous cell carcinoma (CAL27 and FaDu) (**Figures 3A and B**).

Recent clinical trials have shown that repeated CAR-T cell injections exert better tumor

control than single CAR-T administration in solid tumors[34,35]; presumably because fresh

supplies of T cells help overcome T cell exhaustion in the tumor microenvironment [34]. We

tested the in vivo efficacy through weekly injections of fresh mRNA VHH2-CAR-T cells (**Figure**

3C). We focused on the aggressive lung metastatic TNBC model LM2. Luciferase-expressing

LM2 tumor cells were injected intravenously (IV) via the tail vein into NSG mice. After 7-10

days, we confirmed tumor growth in bilateral lungs by bioluminescence imaging (BLI). We

then injected PBS, 20×10^6 mRNA VHH2-CAR-T cells or an equal number of mock T cells via

the tail vein, followed by three additional weekly injections of 5×10^6 effector cells or PBS.

Significant reduction in tumor growth was observed in the VHH2-CAR-T group compared

to the mock T cell group (**Figure 3D**). Mice were euthanized on day 25 for necropsy.

Immunohistochemical (IHC) staining with anti-MET revealed heterogeneous MET expression

within the tumor bed in the mock T cell and PBS groups (**Figure 3E**). Although some residual

tumor cells were seen in the VHH2-CAR-T group, they were mostly MET negative by IHC

(**Figure 3E**); the tumor growth curve also suggested slow growth of these residual tumors

(**Figure 3D**). In the VHH2-CAR-T group, IHC for human CD3 showed clear infiltration and

aggregation of T cells in the residual tumors, whereas the mock T cell group showed only

limited numbers of single scattered T cells (**Figure 3F**). Given the aggressive nature of tumors

generally associated with MET overexpression, these findings indicate that VHH2-CAR-T

could efficiently control tumor growth by eliminating MET-positive tumor cells.

VHH2-CAR-T activity is dependent on MET expression level

Given the necropsy findings that MET can be expressed at heterogenous levels in different

tumor cells, we next tested the antigen threshold for VHH2-CAR-T activation. First, in a panel

of small cell lung cancer cell lines with varying levels of MET expression[23] (**Figure 4A**), we

observed that increased MET expression resulted in increased cytotoxicity by VHH2-CAR-T,

indicating that targeted tumor lysis was dependent on MET expression level (**Figure 4B-E**).

To further investigate antigen-dependent cytotoxicity, we compiled and analyzed a series

of cell lines with varying levels of MET expression and tested cytotoxicity at a high E:T ratio of

5:1 (**Figure 4F-G**). As expected, co-culture of VHH2-CAR-T with cell lines with high levels of

MET expression (786-O, NCIH1341, LM2, and Karpas-422) led to significant tumor cell death

ranging from 80-100%. Although the intermediate MET-expressing cell line, NCIH146, still

showed significant cell death, the killing by VHH2-CAR-T dropped to approximately 45-50%

(against a background of ~10% mock T cell killing). For the low-MET cell line, OCI-LY3, and

MET-negative cell line, K562, VHH2-CAR-T showed cytotoxicity not significantly different from

mock T cells.

Taken together, our findings suggest the existence of an antigen threshold level for VHH2-

CAR-T activity. In normal tissues, MET is expressed at the highest level in hepatocytes, whose

MET expression level is predicted to fall between OCI-LY3 and K562 cell lines, for which we

observed limited cytotoxicity and a potential therapeutic window for VHH2-CAR-T cells.

VHH-CAR-T cells showed low tonic signaling

Since nanobodies generally show high stability[8,12], we tested how incorporation of VHH

influences CAR-T cell function compared to MET-CAR-T cells engineered with different

sources of antigen binding domains (ABDs).

To facilitate comparison, we adapted a previously reported CAR-Jurkat (CAR-J) assay [24]

(**Figure 5A**). We electroporated mRNA encoding CARs with various ABDs into Jurkat cells,

co-cultured them with tumor cell lines at various effector to target ratios for 16 hours, and

assessed expression of CD69 activation marker by flow cytometry. The diffuse large B-cell

lymphoma (DLBCL) cell line, Karpas-422, served as a convenient target for this assay as it

expresses high levels of CD19 in addition to MET, thus allowing comparison with CD19-CAR

(FMC63) as a positive control (**Supplemental Figure 3**).

The selected ABD constructs included three different scFvs, one natural ligand, and VHH1-

4 (**Figure 5B**). ScFv1 and scFv3-CAR-J were constructed similarly to previously described

cell lines [19,20]; scFv2-CAR-J was derived from the monoclonal antibody Onartuzumab

(5D5)[36]. Finally, a natural ligand CAR previously shown to mediate cytotoxicity against

mesothelioma was derived from the NK1 domain of HGF[21].

As expected, CD19-CAR-J cells showed an antigen-dependent increase in CD69

expression, whereas mock Jurkat cells showed only basal levels of expression at all E:T ratios

(**Figure 5B**). The tested scFv-CAR-J cells, however, demonstrated a wide range of antigen-

dependent responses. Notably, scFv1-CARJ showed high basal levels of CD69 expression in

the absence of Karpas-422, indicating tonic signaling. Indeed, increasing the target cell ratio

did not lead to a further increase in CD69 expression, suggesting T cell dysfunction. In contrast,

scFv2-CAR-J showed low basal levels and an antigen-dependent increase in CD69. scFv3-

CAR-J showed low basal CD69 levels, but for an unknown reason did not show an antigen-

dependent response. The ligand-based NK1-CAR-J also failed to show significant CD69

expression upon stimulation. In contrast to the varied behaviors of the tested scFv- and ligand-

CAR-J cells, all VHH-CAR-J cells demonstrated low basal levels and an antigen-dependent

increase in CD69 expression. The various tonic signaling profiles seen in Jurkat cells could

be replicated in co-cultures of primary human T cells and MDA-MB-231 target cells (**Figure**

5C).

To further show that tonic signaling leads to T cell dysfunction, we chose the tonic scFv1

and the low-tonic scFv2 and VHH2 for comparison. As expected, scFv1-CAR-T cells showed

only limited tumor killing compared to scFv2- and VHH2-CAR-T cells, both of which showed

equivalent tumor killing at various E:T ratios (**Figure 5D and 5E**).

The predisposition of different ABDs for tonic signaling is intriguing. Because previous

reports have correlated positive surface charges of ABDs with tonic signaling[37], we modeled

the electrostatic surfaces for scFv1, scFv2, and VHH2 (**Figure 5F**). However, structural

modeling showed no obvious differences in net surface charges, suggesting that other

mechanisms might contribute to tonic signaling (see Discussion).

Different binders endowed varying functional profiles to CAR

We further tested whether low tonic signaling of VHH affects avidity, cytokine secretion,

and tumor killing kinetics. Again, we compared VHH2 with scFv1 and scFv2. All three appear

to share similar predicted affinities (5-10 nM). However, structural modeling predicts different

binding epitopes on MET; while VHH2 and scFv1 occupy a more membrane-proximal epitope,

scFv2 binds to a more membrane-distal one (**Figure 6A**).

We first compared how the three ABDs influence avidity, expecting roughly equal avidity

given their equivalent affinities. Interestingly, despite similar expression levels of all constructs

(**Figure 6B**), VHH2 consistently showed stronger avidity than scFv2, which was greater than

scFv1 for two different tumor cell lines (**Figures 6C-D**). The low avidity of scFv1 could be

related to factors that also contributed to its tonic signaling. The stronger avidity of VHH2 over

scFv2 was unclear, but one possible explanation could be related to VHH2's binding to a more membrane-proximal epitope, as targeting membrane-proximal epitopes has been shown to stabilize the immune synapse[38,39].

In terms of cytokine production, both scFv2 and VHH2 showed higher levels of IFN-γ, TNF-α, and IL-2 after target stimulation than scFv1 (**Figure 6E** and **Supplemental Figure 4**). Among the three constructs, scFv2 was noted to have the highest baseline cytokine secretion in the absence of antigens (especially IFN-γ), suggesting some tonic signaling in scFv2. Using IsoPlexis, VHH2 demonstrated the highest polyfunctional strength index (PSI) in vitro (**Figure 6F**), including the highest effector, stimulatory, and chemoattractive and lower regulatory cytokine levels. Although there were increased inflammatory cytokines detected by IsoPlexis for VHH2, by ELISA VHH2 produced lower levels of proinflammatory cytokines such as IL-6, IL-8, IL-10, IL-1β, and granzyme A (**Supplemental Figure 4**). The reason for this difference is unclear but might be due to different assay conditions.

We also examined the kinetics of tumor cell killing using a microwell platform monitored over four days (as in Figure 2F-I). Although not statistically significant, VHH2-CAR-T cells trended to have the highest single-cell killing rate (**Figure 6G**), the highest percentage of microwells with tumor killing (**Figure 6H**), as well as the highest T cell proliferation (**Figure 6I**). Taken together, these data suggest diverse behaviors of different MET-CAR-T cells, and that VHH2-CAR-T cells have high avidity, potent immune cytokine production including polyfunctionality, and rapid tumor killing kinetics.

VHH2-CAR-T showed potent in vivo efficacy in comparison with scFv2-CAR-T

Emerging evidence suggests that avidity correlates with in vivo and clinical efficacy more so than in vitro cytotoxicity[31]. We tested this hypothesis by comparing VHH2 and scFv2 in vivo given their overall similar in vitro cytotoxicity but differing avidities. Since multiple high doses of VHH2-CAR-T cells appeared to provide potent tumor control (**Figure 3**), we

performed a CAR-T cell stress test by reducing the weekly dose to 2×10^6 CAR-T cells per IV

injection and monitored tumor growth over a longer time period (**Figure 7A**).

Both VHH2-CAR-T and scFv2-CAR-T groups showed significantly better survival than

mock T cells and PBS, and VHH2-CAR-T conferred slightly longer survival than scFv2-CAR-

T (close to statistical significance) (**Figures 7B-D**). There was a significant reduction in tumor

growth rate in the VHH2 and scFv2 groups compared to saline or mock T cells (**Figures 7C-**

D). Interestingly, after the last CAR-T cell injection at day 21, many animals in the VHH2 group

showed lower tumor burdens than the scFv2 group. Three animals in the VHH2 group showed

continuous tumor control until approximately day 67. Overall, there were also fewer

extrapulmonary metastases in the surviving mice in the VHH2 group compared to the scFv2

group (**Figure 7D**). Collectively, our in vivo model indicates that VHH2-CAR-T cells are as

effective as scFv2-CAR-T cells with modestly enhanced tumor control, likely due to more

complete elimination of MET-positive tumor cells by the stronger avidity VHH2.

Although there was no overall survival difference between VHH2- and scFv2-CAR-T cells,

some of the mice appeared to die due to factors other than tumor burden thus complicating

survival curve analysis. On-target/off-tumor cytotoxicity (OTOT) was unlikely as the VHH2

does not bind murine MET[40], and histology showed no obvious evidence of tissue damage.

Other etiologies including cytokine storm or development of early GVHD were also possible.

DISCUSSION

We used an array of functional assays to comprehensively characterize nanobody-based

CAR-T against MET-positive solid tumors. Among a panel of VHH, we found that VHH1 and

VHH2 with intermediate avidity mediated the best killing especially at low E:T ratios. The CD28

costimulatory domain enhanced VHH2-CAR-T cell in vitro cytotoxicity and effectively

controlled tumor growth in vivo. Despite their potency, VHH2-CAR-T cells displayed an

antigen threshold that could potentially discriminate between tumor and normal tissues.

Compared to the more variable behaviors of scFv-CAR-T cells, VHH-CAR-T cells showed low

tonic signaling, high avidity, a favorable cytokine profile, and rapid tumor killing kinetics -

features that likely contributed to prolonged tumor control in vivo despite the transient nature

of the injected mRNA CAR-T cells.

This study highlights construct design as a crucial step in optimizing CARs. We examined

the fundamental aspects of changing the ABD with VHH as a comparison to scFv or ligand-

based approaches. Nanobodies are known for their small size and high stability [8–10]. We

demonstrated that all our tested VHHs have low tonic signaling, whereas the selected scFv

were more variable, with scFv1 being the most prominent example of a tonic signaling CAR.

Interestingly, structural modeling of scFv1 did not indicate excessive surface positive charges

that were previously reported to increase tonic signaling potential[37]. However, other

mechanisms, such as hydrophobic interactions and light-chain-heavy-chain mispairing are

also known to occur in scFv and might contribute to the tonic signaling of scFv1[5]. Although

not without limitations (such as the need for humanization), VHHs might generally offer an

alternative solution for some tonic signaling CARs [4].

We found that the two intermediate avidity VHH1 and VHH2 were more efficacious at

eliminating tumors than either the higher avidity VHH4/VHH5 or the lower avidity VHH3. There

is a complicated relationship between avidity and cytotoxicity—some prior publications

suggest that increased avidity leads to better CAR-T cell performance[41,42], while others

indicate that decreased avidity is preferable[38,43]. One potential explanation is that CAR-T

cells with high avidity have low *Koff* and therefore prolonged engagement with tumor cells,

which can lead to activation-induced cell death and compromised tumor killing[38,43]; on the

other hand, excessively low avidity translates to an unstable immune synapse. In another

context, VHH2-28z showed a trend toward higher avidity and cytotoxic function than VHH2-

BBz, likely due to the predisposition of CD28 to dimerize and thus promote multivalent

interactions[32,44,45]. Finally, the higher avidity of VHH2 compared to scFv2 could potentially

be related to VHH2's binding to a more membrane-proximal epitope and correlated with

slightly better tumor killing kinetics and tumor control in vivo. These findings reflect the

multifactorial nature of avidity, which depends on CAR-target antigen affinity, binding valency,

and CAR-antigen orientation, among other factors. Future studies will benefit from a better

understanding of the factors that influence the avidity-CAR function relationship.

Although the CD28 costimulatory domain led to higher and faster killing as shown here,

other studies have demonstrated earlier T cell exhaustion than 4-1BB[32]. However, we did

not observe notable differences in PD-1 and LAG3 expression between VHH2-28z versus

VHH2-BBz, perhaps due to short duration of mRNA CAR expression. For targeting solid

tumors, it might be advantageous to couple CARs with the CD28 costimulatory domain in an

mRNA format to achieve immediate and maximal tumor killing without the development of T

cell exhaustion, as outlined in a recent review[34].

VHH2-CAR-T cells appeared to have an antigen threshold for activation and thus a

possible therapeutic window. Encouragingly, a phase 1 trial of a MET CAR using the same

scFv2 (5D5) demonstrated safety with mostly clinically manageable grade 1-2 cytokine

release syndrome (CRS)[46]. Given that CAR-T cells generally require a high number of target

antigens to be present to become activated, therapeutic windows are likely present between

normal tissues and MET-overexpressed tumors, as shown here and illustrated elsewhere[47].

One future direction includes exploring logic-gated CAR design to further minimize on-

target/off-tumor toxicity.

Despite the short duration of our mRNA CARs, some VHH2-treated mice displayed

remarkably prolonged tumor control until day 67, even after cessation of CAR-T cell injections

on day 21. This prolonged tumor control was likely due to the ability of the high avidity and

polyfunctional VHH2-CAR-T cells to kill most of the MET-positive cells, which were

presumably more proliferative than the MET-negative cells within the same tumor. One can

imagine a future strategy against solid tumors that would deploy potent effector like VHH2-

CAR-T cells to target a surface antigen related to tumor aggressiveness (e.g. MET) to achieve

prolonged tumor control before bridging to other therapies.

The use of mRNA in the current study offered many advantages but also some limitations.

The main advantages included uniform CAR expression and high T cell viability, which

facilitated construct comparisons in multiple assays. Looking forward, mRNA represents a

particularly promising platform for CAR-T cell reprogramming given the current progress in

developing in vivo nanoparticle delivery systems[48]. However, in some experiments, such as

the microwell co-culture and xenograft mouse models, it would have been beneficial to use a

persistent CAR system so that one could monitor long-term tumor control and T cell functions.

   In summary, we have shown the efficacy of a novel class of VHH-CAR-T cells against MET

with promising in vitro attributes as well as strong tumor control in vivo. The VHH-CAR-T cell

system warrants further preclinical study and development, including VHH humanization and

further construct optimization.

Declarations:

Data availability statement: All data relevant to the study are included in the article or

uploaded as supplementary information.

Ethics statements: Patient consent for publication: Not applicable.

Ethics approval: This study did not include human subjects. All animal studies were

approved by the Yale Institutional Animal Care and Use Committee.

Funding: This work was supported by National Institutes of Health (NIH) grants

R21CA263437 and R21CA282629, Department of the Army award ME220220, a

Developmental Research Program Grant from the Yale Head and Neck SPORE, P50

DE030707, Public Health Service grant number P50CA302572 from the University of Iowa

Neuroendocrine Tumor Specialized Program of Research Excellence and the National Cancer

Institute, and the CT Breast Health Initiative grant to S.G.K., Lion Heart Breast Cancer Pilot

Grant from Yale Cancer Center and Yale Cancer Center Immuno-Oncology K12 training grant

K12CA215110 to P-H.C., and NIH grant R01CA237586, Deparatment of Defense Breast

Cancer Research Program Award HT9425-23-1-0602 and CT Breast Health Initiative grant to

Q.Y. Work carried out by S.D. was supported by the UKRI Medical Research Council

[MR/W006774/1]. D.K.S was supported by the Yale School of Medicine immunohematology

T32 training grant 5T32HL007974-24. Funders had no involvement in study design, data

collection, data analysis, data interpretation, manuscript writing, or the decision to submit.

Conflict of Interest: P-H.C. and S.G.K. have a patent pending on the MET CAR.

Author contributions: P-H.C. and S.G.K. designed all the experiments and wrote the

manuscript. P-H.C. designed and performed all the experiments. Q.L., S.D., D.K.S., and P-

C.C. contributed to data figures and experiments. R.R. and N.C. contributed to molecular

biology and cloning of different CAR constructs. M.N. and A.A. contributed cell lines and

performed western blot and mRNA analysis. Y.L. and Q.Y. helped design and perform the *in*

vivo experiment. All authors reviewed the manuscript.

Acknowledgments: The research was performed as part of the Yale Pathology Physician-

Scientist training track. We also appreciate the support from the K12 Yale Immuno-Oncology

Training Program at Yale Cancer Center. We thank the Yale Animal Research Facility, for

excellent technical support. We also thank the Yale Flow Cytometry Core Facility for

maintaining and making available the cytometers used in the research. The Core is supported

in part by an NCI Cancer Center Support Grant # NIH P30 CA016359. We also thank Keck

Microarray Shared Resource (KMSR) at Yale University for providing the necessary services

using the IsoSpark system, which is funded in part by the National Institutes of Health

instrument grant 1S10OD034429-01.

REFERENCES:

1. Kembuan GJ, Kim JY, Maus MV, Jan M. Targeting solid tumor antigens with chimeric
receptors: cancer biology meets synthetic immunology. Trends Cancer. 2024;10:312–31.
https://doi.org/10.1016/j.trecan.2024.01.003

2. Comoglio PM, Trusolino L, Boccaccio C. Known and novel roles of the MET oncogene in
cancer: a coherent approach to targeted therapy. Nat Rev Cancer. Nature Publishing Group;
2018;18:341–58. https://doi.org/10.1038/s41568-018-0002-y

3. Wang J, Anderson MG, Oleksijew A, Vaidya KS, Boghaert ER, Tucker L, et al. ABBV-
399, a c-Met Antibody-Drug Conjugate that Targets Both MET-Amplified and c-Met-
Overexpressing Tumors, Irrespective of MET Pathway Dependence. Clin Cancer Res Off J
Am Assoc Cancer Res. 2017;23:992–1000. https://doi.org/10.1158/1078-0432.CCR-16-1568

4. Shukla D, Gabunia K, McGettigan SE, Patel PR, Christensen S, Fan T-J, et al. CAR
Binders Affect CAR T-cell Tonic Signaling, Durability, and Sensitivity to Target. CANCER
Immunol Res.

5. Ajina A, Maher J. Strategies to Address Chimeric Antigen Receptor Tonic Signaling. Mol
Cancer Ther. 2018;17:1795–815. https://doi.org/10.1158/1535-7163.MCT-17-1097

6. Frigault MJ, Lee J, Basil MC, Carpenito C, Motohashi S, Scholler J, et al. Identification of
Chimeric Antigen Receptors That Mediate Constitutive or Inducible Proliferation of T Cells.
Cancer Immunol Res. American Association for Cancer Research; 2015;3:356–67.
https://doi.org/10.1158/2326-6066.CIR-14-0186

7. Minguet S, Maus MV, Schamel WW. From TCR fundamental research to innovative
chimeric antigen receptor design. Nat Rev Immunol. 2025;25:212–24.
https://doi.org/10.1038/s41577-024-01093-7

8. De Meyer T, Muyldermans S, Depicker A. Nanobody-based products as research and
diagnostic tools. Trends Biotechnol. 2014;32:263–70.
https://doi.org/10.1016/j.tibtech.2014.03.001

9. Vincke C, Loris R, Saerens D, Martinez-Rodriguez S, Muyldermans S, Conrath K. General
strategy to humanize a camelid single-domain antibody and identification of a universal
humanized nanobody scaffold. J Biol Chem. 2009;284:3273–84.
https://doi.org/10.1074/jbc.M806889200

10. Ditlev SB, Florea R, Nielsen MA, Theander TG, Magez S, Boeuf P, et al. Utilizing
nanobody technology to target non-immunodominant domains of VAR2CSA. PloS One.
2014;9:e84981. https://doi.org/10.1371/journal.pone.0084981

11. Sun Y, Yang X-N, Yang S-S, Lyu Y-Z, Zhang B, Liu K-W, et al. Antigen-induced
chimeric antigen receptor multimerization amplifies on-tumor cytotoxicity. Signal Transduct
Target Ther. 2023;8:445. https://doi.org/10.1038/s41392-023-01686-z

12. Eisa AR, Tavassoli Razavi F, Hoseinzadeh A, Haghmorad D, Baharlou R. Nanobody
CAR-T cells in cancer: From molecular design to clinical translation. Biomed Pharmacother.
2025;193:118819. https://doi.org/10.1016/j.biopha.2025.118819

13. Blueggel M, Gül D, Stauber RH, Knauer SK. Small but mighty: the versatility of
nanobodies in modern medicine. Nanoscale Horiz. 2025;10:2158–71.
https://doi.org/10.1039/D5NH00283D

14. Lou H, Cao X. Antibody variable region engineering for improving cancer
immunotherapy. Cancer Commun. 2022;42:804–27. https://doi.org/10.1002/cac2.12330

15. Zhang XG, Wang L, Yang J, Hu XN, Wang H, Zhang LN, et al. Efficacy and safety of
BCMA nanobody CAR T-cell therapy in relapsed or refractory plasma cell myeloma. Blood
Adv. 2025;9:4543–52. https://doi.org/10.1182/bloodadvances.2025016322

16. Beste G, Denayer T, Hermans G, Szyroki A, Steffensen S, Ververken CJN. Biological
Materials Related to C-Met [Internet]. 2012 [cited 2026 Jan 4].
https://patentscope.wipo.int/search/en/WO2012042026. Accessed 4 Jan 2026

17. Yoo J, Nam JR, Yoo JS, Kim S-W, Lee WS, Shim SR, et al. Human antibody specific to
c-Met and preparation method thereof [Internet]. 2018 [cited 2026 Jan 4].
https://patents.google.com/patent/US10106622B2/en. Accessed 4 Jan 2026

18. Frigault MJ, Zhao Y, Scholler J, June CH. Compositions and Methods for Generating a
Persisting Population of T Cells Useful for the Treatment of Cancer [Internet]. 2019 [cited
2026 Jan 4].
https://patents.google.com/patent/US20190085063A1/en?oq=US20190085063A1. Accessed
4 Jan 2026

19. Kang CH, Kim Y, Lee DY, Choi SU, Lee HK, Park CH. c-Met-Specific Chimeric
Antigen Receptor T Cells Demonstrate Anti-Tumor Effect in c-Met Positive Gastric Cancer.
Cancers. Multidisciplinary Digital Publishing Institute; 2021;13:5738.
https://doi.org/10.3390/cancers13225738

20. Jiang W, Li T, Guo J, Wang J, Jia L, shi X, et al. Bispecific c-Met/PD-L1 CAR-T Cells
Have Enhanced Therapeutic Effects on Hepatocellular Carcinoma. Front Oncol [Internet].
2021 [cited 2022 June 28];11. https://www.frontiersin.org/article/10.3389/fonc.2021.546586.
Accessed 28 June 2022

21. Thayaparan T, Petrovic RM, Achkova DY, Zabinski T, Davies DM, Klampatsa A, et al.
CAR T-cell immunotherapy of MET-expressing malignant mesothelioma. OncoImmunology.
2017;6:e1363137. https://doi.org/10.1080/2162402X.2017.1363137

22. Rabinovich PM, Komarovskaya ME, Wrzesinski SH, Alderman JL, Budak-Alpdogan T,
Karpikov A, et al. Chimeric receptor mRNA transfection as a tool to generate antineoplastic
lymphocytes. Hum Gene Ther. 2009;20:51–61. https://doi.org/10.1089/hum.2008.068

23. Tlemsani C, Pongor L, Elloumi F, Girard L, Huffman KE, Roper N, et al. SCLC-
CellMiner: A Resource for Small Cell Lung Cancer Cell Line Genomics and Pharmacology
Based on Genomic Signatures. Cell Rep. 2020;33:108296.
https://doi.org/10.1016/j.celrep.2020.108296

24. Bloemberg D, Nguyen T, MacLean S, Zafer A, Gadoury C, Gurnani K, et al. A High-
Throughput Method for Characterizing Novel Chimeric Antigen Receptors in Jurkat Cells.
Mol Ther - Methods Clin Dev. 2020;16:238–54. https://doi.org/10.1016/j.omtm.2020.01.012

25. Rabinovich PM, Zhang J, Kerr SR, Cheng B-H, Komarovskaya M, Bersenev A, et al. A
versatile flow-based assay for immunocyte-mediated cytotoxicity. J Immunol Methods.
2019;474:112668. https://doi.org/10.1016/j.jim.2019.112668

26. Tchou J, Zhao Y, Levine BL, Zhang PJ, Davis MM, Melenhorst JJ, et al. Safety and
Efficacy of Intratumoral Injections of Chimeric Antigen Receptor (CAR) T Cells in
Metastatic Breast Cancer. Cancer Immunol Res. 2017;5:1152–61.
https://doi.org/10.1158/2326-6066.CIR-17-0189

27. Abramson J, Adler J, Dunger J, Evans R, Green T, Pritzel A, et al. Accurate structure
prediction of biomolecular interactions with AlphaFold 3. Nature. 2024;630:493–500.
https://doi.org/10.1038/s41586-024-07487-w

28. Vangone A, Bonvin AM. Contacts-based prediction of binding affinity in protein–protein
complexes. eLife. 2015;4:e07454. https://doi.org/10.7554/eLife.07454

29. Xue LC, Rodrigues JP, Kastritis PL, Bonvin AM, Vangone A. PRODIGY: a web server for predicting the binding affinity of protein–protein complexes. Bioinformatics. 2016;32:3676–8. https://doi.org/10.1093/bioinformatics/btw514

30. Sahoo P, Yang X, Abler D, Maestrini D, Adhikarla V, Frankhouser D, et al. Mathematical deconvolution of CAR T-cell proliferation and exhaustion from real-time killing assay data. J R Soc Interface. 2020;17:20190734. https://doi.org/10.1098/rsif.2019.0734

31. Lee L, Lim WC, Galas-Filipowicz D, Fung K, Taylor J, Patel D, et al. Limited efficacy of APRIL CAR in patients with multiple myeloma indicate challenges in the use of natural ligands for CAR T-cell therapy. J Immunother Cancer. 2023;11:e006699. https://doi.org/10.1136/jitc-2023-006699

32. Lotze MT, Olejniczak SH, Skokos D. CD28 co-stimulation: novel insights and applications in cancer immunotherapy. Nat Rev Immunol. 2024;24:878–95. https://doi.org/10.1038/s41577-024-01061-1

33. Salter AI, Ivey RG, Kennedy JJ, Voillet V, Rajan A, Alderman EJ, et al. Phosphoproteomic analysis of chimeric antigen receptor signaling reveals kinetic and quantitative differences that affect cell function. Sci Signal. 2018;11:eaat6753. https://doi.org/10.1126/scisignal.aat6753

34. Albelda SM. CAR T cell therapy for patients with solid tumours: key lessons to learn and unlearn. Nat Rev Clin Oncol. 2024;21:47–66. https://doi.org/10.1038/s41571-023-00832-4

35. Monje M, Mahdi J, Majzner R, Yeom KW, Schultz LM, Richards RM, et al. Intravenous and intracranial GD2-CAR T cells for H3K27M+ diffuse midline gliomas. Nature. 2025;637:708–15. https://doi.org/10.1038/s41586-024-08171-9

36. Shah PD, Huang AC, Xu X, Orlowski R, Amaravadi RK, Schuchter LM, et al. Phase I Trial of Autologous RNA-electroporated cMET-directed CAR T Cells Administered Intravenously in Patients with Melanoma and Breast Carcinoma. Cancer Res Commun. 2023;3:821–9. https://doi.org/10.1158/2767-9764.CRC-22-0486

37. Chen J, Qiu S, Li W, Wang K, Zhang Y, Yang H, et al. Tuning charge density of chimeric antigen receptor optimizes tonic signaling and CAR-T cell fitness. Cell Res. 2023;33:341–54. https://doi.org/10.1038/s41422-023-00789-0

38. Wang Y, Buck A, Piel B, Zerefa L, Murugan N, Coherd CD, et al. Affinity fine-tuning anti-CAIX CAR-T cells mitigate on-target off-tumor side effects. Mol Cancer. 2024;23:56. https://doi.org/10.1186/s12943-024-01952-w

39. Freeman R, Shahid S, Khan AG, Mathew SC, Souness S, Burns ER, et al. Developing a membrane-proximal CD33-targeting CAR T cell. J Immunother Cancer. 2024;12:e009013. https://doi.org/10.1136/jitc-2024-009013

40. Slørdahl TS, Denayer T, Moen SH, Standal T, Børset M, Ververken C, et al. Anti-c-MET Nanobody - a new potential drug in multiple myeloma treatment. Eur J Haematol. 2013;91:399–410. https://doi.org/10.1111/ejh.12185

41. Porazzi P, Nason S, Yang Z, Carturan A, Ghilardi G, Guruprasad P, et al. EZH1/EZH2
inhibition enhances adoptive T cell immunotherapy against multiple cancer models. Cancer
Cell. 2025;43:537-551.e7. https://doi.org/10.1016/j.ccell.2025.01.013

42. Barisa M, Zappa E, Muller H, Shah R, Buhl J, Draper B, et al. Functional avidity of anti-
B7H3 CAR-T constructs predicts antigen density thresholds for triggering effector function
[Internet]. 2024 [cited 2025 July 4]. https://doi.org/10.1101/2024.02.19.580939

43. Zhang Y, Patel RP, Kim KH, Cho H, Jo J-C, Jeong SH, et al. Safety and efficacy of a
novel anti-CD19 chimeric antigen receptor T cell product targeting a membrane-proximal
domain of CD19 with fast on- and off-rates against non-Hodgkin lymphoma: a first-in-human
study. Mol Cancer. 2023;22:200. https://doi.org/10.1186/s12943-023-01886-9

44. Alabanza L, Pegues M, Geldres C, Shi V, Wiltzius JJW, Sievers SA, et al. Function of
Novel Anti-CD19 Chimeric Antigen Receptors with Human Variable Regions Is Affected by
Hinge and Transmembrane Domains. Mol Ther J Am Soc Gene Ther. 2017;25:2452–65.
https://doi.org/10.1016/j.ymthe.2017.07.013

45. Hamieh M, Mansilla-Soto J, Rivière I, Sadelain M. Programming CAR T Cell Tumor
Recognition: Tuned Antigen Sensing and Logic Gating. Cancer Discov. 2023;13:829–43.
https://doi.org/10.1158/2159-8290.CD-23-0101

46. Shah PD, Huang ACC, Xu X, Zhang PJ, Orlowski R, Matlawski T, et al. Phase I trial of
autologous cMET-directed CAR-t cells administered intravenously in patients with
melanoma & breast carcinoma. J Clin Oncol. Wolters Kluwer; 2020;38:10035–10035.
https://doi.org/10.1200/JCO.2020.38.15_suppl.10035

47. Chiriaco C, Donini C, Cortese M, Ughetto S, Modica C, Martinelli I, et al. Efficacy of
CAR-T immunotherapy in MET overexpressing tumors not eligible for anti-MET targeted
therapy. J Exp Clin Cancer Res. 2022;41:309. https://doi.org/10.1186/s13046-022-02479-y

48. Kong B, Kim Y, Kim EH, Suk JS, Yang Y. mRNA: A promising platform for cancer
immunotherapy. Adv Drug Deliv Rev. 2023;199:114993.
https://doi.org/10.1016/j.addr.2023.114993

766
767

 FIGURE LEGENDS

 Figure 1 VHH-CAR-T with intermediate avidity showed the most potent killing.

 (A) Schematic diagram of VHH-CAR construct. (B) Expression level of VHH-CAR by GFP.

 Red peak = mock electroporated T cells. (C) Avidity curve of VHH-CAR-T against MDA-MB-

 231, with the % T cells bound to target cells at 1000 pN shown in (D). N = 4 technical

 replicates; *P* values comparing avidity at 1000 pN were calculated using one-way ANOVA).

 (E) AlphaFold3 modeling of VHH binding sites on MET. MET = cyan; HGFα chain = light

 gray; HGFβ chain = dark gray; VHH1 = red; VHH2 = blue; VHH3 = orange; VHH4 = green;

 VHH5 = brown. (F) VHH-CAR-T cytotoxicity against MDA-MB-231 (n=4 technical replicates;

 P values by two-way ANOVA). (G) VHH1 and VHH2-CAR-T against A549, with tumor

 proliferation monitored in CellCyte (n=6 technical replicates; *P* values by two-way ANOVA).

 Data shown are mean$\pm$SD; *P<0.05; **P<0.01; ***P<0.001; ****P<0.0001*.

 Figure 2: CD28 costimulatory domain conferred stronger VHH2-CAR activation. (A)

 Expression level of mRNA VHH2-BBz and VHH2-28z in primary human T cells. (B) Avidity

 assay using Lumicks z-Movi, comparison between VHH2-BBz and VHH2-28z (n = 3

 technical replicates; two-way ANOVA). (C) Comparison of MDA-MB-231 killing between

 VHH2-BBz and VHH2-28z (n=8 technical replicates, two-way ANOVA). (D) Intracellular

 cytokine production analyzed by flow cytometry; cells harvested 16 hours after co-culture

 with MDA-MB-231 in the presence of 2μM of the Golgi inhibitor monensin (n = 4 technical

 replicates, two-way ANOVA). (E) Polyfunctioinal strength index (PSI) using IsoPlexis,

 comparison between CD8+ VHH2-BBz and CD8+ VHH2-28z 16 hours after co-culture with

 Karpas-422. Effector cytokines (Granzyme B, IFN-γ, MIP-1a, perforin, TNF-α, TNF-β);

 Stimulatory cytokines (GM-CSF, IL-12, IL-15, IL-2, IL-21, IL-5, IL-7, IL-8, IL-9);

 Chemoattractive cytokines (CCL-11, IP-10, MIP-1b, RANTES); Regulatory cytokines (IL-10,

 IL-13, IL-22, IL-4, sCD137, sCD40L, TGF-b1); Inflammatory cytokines (IL-17A, IL-17F, IL-1b,

 IL-6, MCP-1, MCP-4). (F-I) Microwell assay using TROVO, comparing single cell killing rate

(F) (n=3 technical replicates; two-tailed Student's *t*-test), T cell proliferation (G) (n=2

donorsx3 replicates; two-tailed Student's *t*-test), and tumor killing (H) (n=2 donors x 3

replicates each, two-tailed Student's *t*-test). (I) Representative images used to calculate

data shown in (G) and (H). Each row represents one microwell monitored over 4 days. Each

microwell is set up with E:T =20:10. T cell = red. *P<0.05; **P<0.01; ***P:<0.001;*

*****P:<0.0001.*

Figure 3: In vitro and in vivo efficacy of VHH2-CAR. (A) MET expression in tumor cell

lines. Red = IgG isotype control. (B) Cytotoxicity of VHH2-CAR-T cells at different E:T ratios

(n=3 technical replicates; two-way ANOVA). (C) Scheme of in vivo experiment using the

LM2 tumor model. (D) BLI of tumor growth in vivo. PBS = 3 animals; Mock T = 2 animals;

VHH2-CAR-T = 3 animals. *P* values were calculated using two-way ANOVA. (E) MET IHC,

10x objective on the left and 20x objective on the right. The boxed regions were magnified

and shown on the right. The bar graph represents quantifications of MET expression. *P-*

value was calculated by two-tailed Student's *t*-test. (F) CD3 IHC, 20x objective. *P<0.05;*

***P<0.01; ***P<0.001; ****P<0.0001.*

Figure 4: Antigen threshold for VHH2-CAR. (A) mRNA and western blot of MET

expression levels in a series of small cell lung cancer cell lines. The mRNA data comes from

CellMiner[23]. (B and C) MET expression by flow cytometry and cytotoxicity in NCI-H1341.

N= 3 technical replicates in the cytotoxicity assay; *P* values from two-way ANOVA. (D-E)

MET expression by flow cytometry and cytotoxicity in NCI-H146. N= 3 technical replicates; *P*

values calculated using two-way ANOVA. (F) Histogram of MET expression levels in

different cell lines. (G) Cytotoxicity of VHH2-CARs against the same cell lines as in (E). N= 3

technical replicates; *P* values calculated using two-tailed Student's *t*-test. *P<0.05; **P<0.01;*

****P<0.001; ****P<0.0001.*

Figure 5: VHH-CARs showed low tonic signaling. (A) Experimental design of CAR-Jurkat assay. (B) Comparison of CD69 expression in CAR-Jurkat cells with varying levels of antigen stimulation. (C) Comparison of CD69 expression in primary human T cells electroporated to express CARs. E:T ratio was 1:1, and co-culture was incubated for 16 hours before staining for flow cytometry. (D-E) Cytotoxicity comparison between VHH2, scFv1, and scFv2 using two different cell lines with different MET expression levels. (F) Electrostatic surface charge of MET binders. Modeling was performed using PyMol. Blue = more basic/positive charge; red = more acidic/negative charge; white = neutral. In experiments shown in B-E, n=3 technical replicates, and P values were calculated with two-way ANOVA. *$P<0.05$; **$P<0.01$; ***$P<0.001$; ****$P<0.0001$.

Figure 6: Avidity and kinetics comparisons of CARs. (A) Different binding epitopes predicted by AlphaFold3 for VHH2, scFv1, and scFv2. (B) Expression of compared CARs. (C) Avidity measurement by Lumicks z-Movi using MDA-MB-231 as target cells. N = 3 technical replicates, with P values from two-tailed Student's t-test. (D) Avidity measurement against the 786-O cell line. N = 4-5 technical replicates; P values from two-tailed Student's t-test. (E) Intracellular cytokine production by flow cytometry. Cells were stained before and 16-hours after stimulation by Karpas-422. N=3 technical replicates; P values from two-way ANOVA. (F) Polyfunctional strength index by IsoPlexis. Effector cytokines (Granzyme B, IFN-γ, MIP-1a, perforin, TNF-α, TNF-β); Stimulatory cytokines (GM-CSF, IL-12, IL-15, IL-2, IL-21, IL-5, IL-7, IL-8, IL-9); Chemoattractive cytokines (CCL-11, IP-10, MIP-1b, RANTES); Regulatory cytokines (IL-10, IL-13, IL-22, IL-4, sCD137, sCD40L, TGF-b1); Inflammatory cytokines (IL-17A, IL-17F, IL-1b, IL-6, MCP-1, MCP-4). (G-I) Hydrogel-based microwell co-culture assay by TROVO, showing single cell killing rate (G, 1:10 E:T), killing (H, 20:10 E:T), and T cell proliferation (I, 20:10 E:T). N=3 technical replicates; P values using two-tailed Student's t-test. (J) Representative images of microwell co-culture at 20:10 E:T. Each row represents one microwell monitored over time. *$P<0.05$; **$P<0.01$; ***$P<0.001$; ****$P<0.0001$.

Figure 7: In vivo comparison between scFv2 and VHH2-CAR. (A) Experimental design of

in vivo efficacy. (B) Kaplan-Meier survival curve using a predefined tumor burden threshold

of 1×10^9 photons/sec as the survival endpoint for analysis purposes. *P*-value shown for

VHH2-CAR versus scFv2-CAR comparison. (C) Tumor growth curve by BLI imaging. Each

group = 7 mice; each line = 1 mouse. (D) Representative images of BLI.

SUPPLEMENTAL FIGURES:

Supplemental figure 1: mRNA CAR manufacturing through electroporation. (A) mRNA

CAR design. (B) RNA gel showing examples of successful mRNA synthesis and poly(A) tail

incorporation. (C) CAR expression was dose-dependent as shown by GFP. (D) Time course

of GFP decay after mRNA electroporation in T cells.

Supplemental figure 2: Intracellular cytokine assessment. (A) Gating strategy. (B)

Example flow cytometry plot of cytokine production.

Supplemental figure 3: Lack of significant exhaustion marker expression in VHH2-28z.

(A) Representative flow cytometry plot of PD-1 versus TIM3 and PD-1 versus LAG3

expression. CAR-T cells were co-cultured with Karpas-422 for 48 hours before harvesting for

flow cytometry. (B) Comparison of PD-1 and LAG3 expression, and PD-1/TIM3

coexpression between mock T, VHH2-28z, and VHH2-BBz T cells. N=3 technical replicates;

P values from two-way ANOVA. **P<0.05; **P<0.01; ***P<0.001; ****P:<0.0001.*

Supplemental Figure 4: Karpas-422 cell line with MET and CD19 expression. (A) Flow

cytometry demonstrating expression of both antigens. Red curve = IgG isotype control. (B)

Confirmation of cytotoxicity of different MET CAR-T cells and CD19-CAR-T cells against

Karpas-422. N = 3 technical replicates, with *P* values calculated from two-tailed Student's *t*-

test. **P<0.05; **P<0.01; ***P<0.001; ****P:<0.0001.*

880 **Supplemental Figure 5: Bulk cytokine secretion by ELISA between various CAR**

881 **constructs against MDA-MB-231.** Panel A shows selected effector cytokines, while panel

882 B shows selected cytokines known to mediate CAR T cell related toxicity and inflammatory

883 responses.

Chen, et al., Figure 1

Chen, et al., Figure 2

Chen, et al., Figure 3

Chen, et al., Figure 4

Chen, et al., Figure 5

Chen, et al., Figure 6

Chen, et al., Figure 7