Dendritic delay lines shape the computation of sound location in neurons of the

gerbil medial superior olive

Authors/affiliations

Jared Casarez[1]*, Rebecca I. Voglewede[1]*, Bradley D. Winters[2], Ken Ledford[1] and Nace

L. Golding[1]

* Equal contributions

[1] The University of Texas at Austin, Department of Neuroscience and Center for Learning

and Memory, 1 University Station C7000, Austin TX 78712-0248, USA.

[2] Current address: Department of Biological Sciences and University Hospitals –

NEOMED Hearing Research Center, Northeast Ohio Medical University, Rootstown, OH,

44272, United States and Brain Health Research Institute, Kent State University, Kent,

OH, 44242, United States

Corresponding author and lead contact

Nace L. Golding: The University of Texas at Austin, Department of Neuroscience and

Center for Learning and Memory, 1 University Station C7000, Austin TX 78712-0248,

golding@austin.utexas.edu.

Summary

In mammals, neurons of the medial superior olive encode information about sounds arising from discrete spatial locations along the horizon. This tuning requires that an internal delay in the brain must offset acoustic disparities to ensure coincident arrival of excitatory inputs driven from the two ears. The source of this optimal internal delay, originally assumed to arise from axonal delay lines, is currently controversial in mammals. Here we use 2-photon guided paired dendritic and somatic recordings together with compartmental modeling of 40 complete MSO neuron morphologies to demonstrate that the dendrites themselves serve as a significant source of internal delay. We show that most MSO neurons exhibit morphological asymmetries that impose different EPSP delays across dendrites and shifts in optimal interaural time differences. Dendrite-based delays in the mammalian MSO are heterogeneous within each isofrequency laminae and provide a stable, structural mechanism to help tune individual neurons to sounds from different azimuthal locations.

Keywords

dendrite; dendritic filtering; coincidence detection; auditory; sound localization.

INTRODUCTION

Spatial information in the auditory system is critical for humans and other mammals to assemble a 3-dimensional view of the world as well as to understand speech and communication signals, especially in environments with many sounds overlapping in time and in frequency. The auditory system is distinct from vision or touch in that there is no explicit peripheral representation of space, in this case in the cochlea, only a representation of frequency. Accordingly, the central auditory system must compute horizontal sound location indirectly by combining a series of monaural and binaural cues [1,2]. Spatial coding in the central auditory system is a prime example of a centrally synthesized or computational map.

Interaural time difference cues are used by birds and mammals to compute horizontal sound location at low frequencies [3]. In mammals this computation takes place in the medial superior olive (MSO), whose neurons are the first to receive converging excitatory and inhibitory inputs driven from the two ears. Intriguingly, in individual neurons, binaural inputs are integrated within a bipolar dendritic architecture, with ipsilateral and contralateral excitatory inputs segregated onto lateral and medial dendritic branches respectively [4,5]. Ipsilateral and contralateral inhibition converges onto the soma and proximal dendrites [4–8]. MSO neurons detect the optimal combination of these inputs, converting temporal coincidence into a rate code [6,9,10].

The concept of internal delay is central to coincidence detection in the MSO, where acoustic delays to the two ears are countered in time by a mechanism of delay inside the brain. This internal temporal asymmetry was postulated in early models to arise from

axonal delay lines formed by ladder-like combinations of bilateral excitatory inputs (the

"Jeffress model") [11] . While the Jeffress model enjoys strong support in birds [12–14] ,

comparable studies in mammals have failed to reveal systematic axonal input delay lines

nor any mapping of delay along the nucleus [6,15,16] . More recently, there is evidence that

fast inhibition mediates internal delay by shifting the peaks of EPSPs [17–19] . However, this

mechanism is subject to vigorous debate [20–22] , and it is not clear that this mechanism

alone can explain the range of receptive field locations in the MSO or their insensitivity to

sound level. Other mechanisms have been proposed to mediate internal binaural

asymmetry, including asymmetrical axon position, medio-lateral differences in the rising

slopes of EPSPs [23], and differences in tuning of binaural inputs ("stereausis") [24,25].

All models of coincidence detection must address the issue of the temporal distortions

imposed by the dendrites themselves. However, to date models of MSO neurons have

been represented either as point neurons or as simplified, symmetrical "ball-and-stick"

models [17,26–30] . Previous measurements of dendritic propagation have revealed that

EPSPs encounter delays in their propagation to the soma up to ~150 µs, despite some

degree of temporal compensation by activation of low voltage-activated K channels in the

soma and proximal dendrites [27] . However, these measurements have largely been

restricted to the proximal half of the dendritic arbor and have relied on averaging highly

variable populations of neurons. Given that the dendrites of MSO neurons alter their

branching pattern and diameters during early hearing and following disruptions of normal

hearing [31–33], they might serve as a potential mechanism for adjusting the timing

requirements for sounds from the two ears to arrive coincidently at the soma and axon.

Here we demonstrate that the dendritic arbors of MSO neurons are highly heterogeneous

and show a striking continuum of asymmetry across the medial and lateral branches that

receive segregated inputs driven respectively from the two ears. We show with dual 2-

photon fluorescence-guided patch recordings that inputs to smaller and/or more distal

dendrites show more substantial propagation delays than previously recognized, and

computational modeling of a large population of intact neuronal morphologies shows that

these dendritic asymmetries translate into stable internal delays that, as a population,

could account for the full range of naturally occurring horizontal locations. Finally, we

show that inhibition preserves the effects of dendritic asymmetry on interaural time

differences (ITDs) and helps bring the region of maximum slope of firing changes within

the ecologically relevant range. Thus dendrite-based delay lines likely work synergistically

with other integrative mechanisms purported to shape the perception of horizontal

position in mammals.

RESULTS

To understand how MSO neurons process interaural time differences in the dendrites, we made fluorescence-guided dendritic and somatic patch-clamp recordings visualized under 2-photon microscopy. Using this approach, we were able to target the distal third of the dendritic arbor, including secondary and tertiary branches, regions that have largely escaped investigation in these neurons. We then injected simulated EPSCs (τ_{rise} and τ_{decay}, 0.2 ms) into the dendritic pipette and recorded potentials simultaneously at both dendrites and soma. We observed strong variability in both the percent amplitude attenuation at the soma $\left(\left(\frac{V_{soma}}{V_{dend}}\right) \times 100\right)$ as well as the propagation delay of the EPSP,

Fig. 1: Heterogeneity in attenuation and timing of simulated excitatory synaptic potentials following propagation from distal dendritic locations to the soma.
a,b, 2-photon fluorescence-guided dual dendritic and somatic patch recordings from two MSO neurons of comparable distances (78 and 73 µm, respectively). Example neuron in **a₁** shows stronger attenuation of somatic propagation of a simulated EPSP (**a₂** vs. **b₂**) and longer peak delay (**a₃** vs. **b₃**). EPSC amplitudes: cell **a**, 600 pA; cell **b**, 400 pA. **c,d,** Group data for attenuation and peak delay for all paired MSO neuron recordings. There is a poor correlation of both measures as a function of distance.

measured as the difference in EPSP peak time at the soma relative to the dendrite (Fig. 1). Notably, the delays measured in these recordings ranged between 100 and 300 µs, exceeding the physiological range of ITDs measured in gerbils. In these recordings,

111 there was little correlation between local morphological features of recorded dendrites

112 and propagation delay of the EPSP, measured as the difference in the peak timing of

113 dendritic and somatic EPSPs (Fig. 2a-c). However, there was a stronger correlation

114 between the local dendritic membrane time constant and EPSP propagation delay (Fig.

115 2d). In addition, there was a weak trend of increasing dendritic membrane time constant

Fig. 2: EPSP propagation delays are not correlated with local morphology but rather more global physiological properties.
a, Closeups of distal dendritic morphology during paired dendritic and somatic recordings. b,c, Delay of peak EPSPs during propagation from dendrite to soma shows no significant correlation with local diameter or tapering rate of local dendrite. (dendritic EPSCs: 400-1000 pA). d, Peak EPSP delay from dendrite to soma was weakly correlated with the membrane time constant recorded at the dendritic recording site. e, Dendritic membrane time constant increased with distance of the recording site from the soma.

with recording distance (Fig. 2e), possibly because of proximity to the dendritic

terminations. These results may reflect that an understanding of the integrative properties

of the dendrites requires an analysis of the larger morphological and electrotonic structure

of the neuron.

To understand how overall dendritic structure impacts binaural coincidence detection, we

analyzed a database of biocytin-filled MSO neurons that exhibited complete dendritic

structures (40 of 270 neurons from Bondy et al., 2021 [34] ; Fig. 3). We quantified dendritic

length, surface area and branching, with special emphasis on comparisons of medial and

lateral dendritic arbors. Higher-order (2° and higher) dendritic branches were prominent

in these neurons, comprising 70% of the membrane surface area (Fig. S1).

A striking feature of most neurons in this data set was an overall asymmetry between the

lateral and medial dendritic arbors, reflecting differences in length, branching, and/or

diameter of the dendrites (Fig. 3a-c). We quantified this asymmetry according to two

Fig. 3: MSO neuron morphology is inhomogeneous and asymmetric, regardless of tonotopy.
a, Reconstructed morphologies rank ordered by tonotopy (left) and degree of asymmetry (right; average path length). Orderings span from low frequency (top) to high frequency (bottom) and medial (top) and lateral (bottom) asymmetry, respectively. Colored gradients added to visualize ordering. Uniquely colored dots placed beneath each morphology for differentiation. **b**, Fold difference between average path length of medial and lateral dendrites for all reconstructed morphologies (n=40) with vertical axes ordered by tonotopic location (top) or degree of asymmetry (bottom). Gradients added as in **a**. Colored points correspond to **a**. **c**, Same as **b**, except comparing total surface area of medial and lateral branches. Average medio-lateral path length and surface area fold differences ($\pm$ SEM) were 1.080 ± 0.057 and 1.112 ± 0.133 (biased laterally) and tonotopic R^2 values were -0.42 and -0.23.

parameters, the average path length (average length from each dendritic end to the

soma) and membrane surface area, with the latter parameter reflecting the collective

influence of dendritic length, diameter and branching pattern. We then defined the degree

of bilateral asymmetry as fold differences between the larger vs. smaller dendritic

measure for each parameter. We observed no obvious correlation between dendritic

asymmetry and tonotopic location (Fig. 3a, *left* and 3b, *top*). However, we found that

dendritic asymmetry formed a finely graded continuum, with a slight bias toward lateral

dendrites (Fig. 3b, *bottom*). Interestingly, there were no significant differences in

population averages of the medio-lateral path length and surface area (1.080 ± 0.057 and

1.112 ± 0.133 fold differences; p=0.562), though there was a slight lateral bias. These

results indicate that the averaging of dendritic morphological parameters across neurons

obscures what is likely a key structural and computational feature.

To understand how morphological asymmetry impacts synaptic integration, we

constructed compartmental models using the same 40 reconstructions analyzed in Fig. 3.

We introduced an excitatory synaptic conductance (alpha function: τ = 290 µs) into each

dendritic compartment and then measured the voltage attenuation and propagation delay

from the dendritic input location to the soma (Figs. 4-5). At comparable distances from

Fig. 4: MSO morphology nonlinearly filters and attenuates propagating EPSPs through its passive cable properties.
a, Visualization EPSP filtering at different dendritic sites. Cell morphology (top) with similarly distant example synaptic sites indicated with green circles and corresponding numbers. Red triangle indicates the somatic compartment. Synaptic conductance indicated as g_{syn}. Four sets of paired dendritic (green) and somatic (red) traces (bottom) for each example synaptic site. Numbering of traces corresponds to labels on morphology above. Dendritic distance from soma for each site indicated below traces. **b**: Same procedure as in **a**, with an adjusted unitary synaptic conductance (see Methods), with measurements from each computational compartment. Model morphology and orientation same as in **a**. Comparison between active channel (left) and passive channel model (right). Recorded EPSP amplitude at synaptic (top, green) and somatic (bottom, red) compartment as a function of distance from soma. **c**, Actual (top) and electrotonically transformed (bottom) morphologies. Transformation visualizes degree of EPSP attenuation from each compartment (see Methods). **d**, Group data (n=40) of average somatic depolarization asymmetry between mediolateral sides. Gradient as in Fig 3. Dotted line indicates equal medial and lateral average somatic depolarizations. Cell from **a-c** indicated with a triangle.

the soma, EPSP amplitude at the site of synaptic input and following propagation to the

soma varied considerably, like the physiological data from paired recordings in Figure 1

(Fig. 4a). Plots of local dendritic and propagated somatic EPSP amplitudes across all

synapse locations further revealed striking asymmetries in propagation efficacy of EPSPs

to the soma (Fig. 4b, "Active cell"). In MSO neurons, the timing and attenuation of EPSPs

along dendrites are influenced both by passive cable filtering as well as interactions with

low voltage-activated K channels, which are active in the subthreshold voltage range in

these neurons [27] . To determine the degree to which asymmetries in low voltage-activated

K channels impact EPSP attenuation at the soma we converted all voltage-activated

conductances to leak channels (Fig. 4b, "Passive cell"; see Methods). While EPSPs

propagated slightly more effectively to the soma from all dendritic locations in the passive

model, it did not fundamentally disrupt mediolateral asymmetry, indicating that these

differences arose primarily from the passive electrotonic architecture of the cell. The

branching structure had a significant effect on dendritic propagation asymmetry. This

could be better visualized in inward morphoelectrotonic transforms of cells, which exhibits

dendritic compartments according to their electrotonic, versus physical lengths. In the

example neuron in Fig. 4c, large primary dendrites are deemphasized, while thinner

secondary and tertiary branches appear as electrotonically elongated. Differences

between the medial and lateral dendritic arbors in this neuron and many others gave rise

to asymmetries in the magnitude of the average EPSP arriving at the soma.

Further, we observed that our population of cells formed a continuum in the degree of

somatic depolarization asymmetry, in line with the asymmetry in dendritic structure (Fig.

4d). Dendritic asymmetry also translated into differences in average EPSP timing across

the medial and lateral dendrites. Figure 5a shows for two example neurons the spatial

profile of delays of EPSPs propagating from each dendritic compartment to the soma.

Fig. 5: EPSP timing significantly varies within and between cells, with morphological asymmetry informing overall cell timing. a, EPSP delay visualization in two exemplar MSO neurons. Coloring of each cell corresponds to the propagation delay of EPSPs to the soma from its respective compartment. Legend of propagation delay coloring in top right. Cells are oriented mediolaterally according to the scale bar. Outline added to increase gradient readability. **b,** Same data as in **a**, except as a function of distance from the soma. Orientation of cells same as in **a. c,** Group data (n=40) of the difference in average EPSP propagation delay between medial and lateral dendrites. Cells 1 & 2 are indicated with same color as in **b**. Dotted line at 0 ms indicates an equal average EPSP delay between sides. Blue gradient along horizontal axis (as in Fig. 3) to indicate spectrum of asymmetry. **d,** Morphological transforms according to EPSP delays. Morphological centroids of actual cell morphologies (top) and transformed morphologies (bottom). Length corresponds to each sections EPSP delay contribution. Coloring of cell sections is consistent between actual and transformed morphologies for easier differentiation and comparison. Dotted line added to mark the midline.

The asymmetry in dendritic timing can be more quantitatively visualized in plots of peak

EPSP delay ($t_{dend} - t_{soma}$) as a function of dendritic distance from the soma (Fig. 5b).

Dendritic branch points often introduced substantial jumps in propagation delays, and this

is reflected in these plots as increases in slope. To quantify the influence of dendritic

asymmetry on the optimal summation of bilateral EPSPs at the soma, we measured the

electrotonic delay at the soma for every 2 μm compartment of lateral and medial dendritic

subtrees. In our population of 40 neuron models, there were differences in the average

relative time of arrival of EPSPs to the soma from compartments in the medial and lateral

dendrites. Further, these differences formed a smoothly graded continuum of delays

(Fig. 5c).

MSO neurons exhibit a range of intrinsic electrical properties, and so we examined a

subset of neuron morphologies to understand the functional implications of this

physiological diversity. Using a neuron with an asymmetric dendritic arbor, we

systematically slowed the membrane time constant by reducing the density of both low

voltage-activated K channels and HCN channels (Fig. S2). In both cases, the timing of

EPSPs reaching the soma was linearly related to the value of the membrane time

constant. Thus, the effects of dendritic asymmetry on EPSP timing and amplitude at the

soma are expected to be substantially higher in MSO neurons with slower intrinsic

membrane properties. To understand how the differential timing of EPSPs propagating

from medial and lateral dendrites is translated into ITDs, we simulated binaural excitatory

input using a spiking model of MSO neurons (see Methods). This model featured one

axonal input per terminal branch segment as well as a centripetal (dendrite-to-soma)

pattern of dendritic innervation, running from the terminal compartments to the soma, in

accordance with results from anatomical studies (Fig. 6a) [15,16,28]. The conduction velocity

of these inputs was set to 1 ms^{-1}, approximating values of unmyelinated terminal axonal

arborizations of similar caliber [13,35] . The timing of ipsilateral and contralateral inputs was

varied systematically, triggering spiking only when the two inputs were in close

coincidence (Fig. 6b). The location of these excitatory ITD functions were dependent on

medio-lateral asymmetry. Functions shifted toward the side of the electrically longer

dendrite, reflecting the need for earlier activation of that side to compensate for a longer

dendritic propagation delay (Fig. 6c). Our population of 40 neurons showed peak firing at

Fig. 6: Simulated ITDs demonstrate conservation of morphological impact on coincident detection.
a, Schematic visualization of ITD innervation. Somatic compartment and dendrites are shown in gray and black, respectively. Green bulbs represent excitatory synapses, with corresponding arrows to illustrate axonal input velocity (ν_{input}). Dotted lines illustrate point at which there is considered to be no axonal input delay (see Methods). Axon attached to soma, with initial segment, nodes, and internodes. **b**, Example somatic (top) and axonal (bottom) traces from ITD simulations. ITDs of -0.2 ms (left), 0.1 ms (middle, optimal coincidence), 0.3 ms (right) shown. Traces colored only for differentiation. Dotted line ($V_{threshold}$) indicates depolarization threshold for action potential consideration. Morphology of model cell indicated in upper right. **c**, Three example ITD firing curves from varied morphologies, medially biased (left), symmetrical (middle), and laterally biased (right). Corresponding morphologies imaged above. Light orange span represents range of ecologically possible ITDs for model species. Dashed line placed at ITD of 0 ms. Dot indicates centroid ITD value. **d**, Sorted group data (n=39) of centroid ITDs, with ν_{input} = 1 ms^{-1}. Cells from **c** indicated with filled colored points. **e,f**, Sorted group data (n=39) of absolute centroid ITDs, varying axonal input velocity (**e**) and peak firing probability (**f**). Cells from **c** indicated with correspondingly filled colored points.

delays that traversed a continuum of values (-0.11 to 0.08 ms, median -0.02; Fig. 6d; Fig.

S3). The precise range of values depended in part on the assumptions on the velocity of

excitatory axons. The largest range of ITDs was achieved when there was closer

matching between the velocity of axonal input velocity and postsynaptic EPSPs

propagating from the dendrites to the soma, which maximized temporal coincidence at

the soma of spatially disparate inputs in the dendrites (Fig. 6e). Finally, ITDs shifted less

than 3% in the face of input amplitude changes that resulted in firing probabilities between

0.2 and 0.8, indicating that dendritic asymmetry is highly stable across changing stimulus

conditions.

Inhibition is a critical component of ITD encoding, although its precise role is subject to

vigorous debate. To understand how dendritic asymmetry impacts the effects of inhibition,

we added bilateral inhibitory conductances to the somatic compartment of the bilateral

excitatory model in Fig. 6, consistent with the known somatically biased innervation of

MSO neurons from the lateral and medial nuclei of the trapezoid body (LNTB and

MNTB; [8,36]). We set the timing of inhibitory conductances to be 500 μs in advance of

excitatory inputs from the same side, in accordance with a previously published

estimate [21]. We then adjusted the amplitude of inhibitory conductances to produce

summed IPSPs of between ~1-3 mV, which reduced firing probability by 20, 40, 60 and

100% (Fig. 7a). With increasing amplitude of IPSPs, resultant ITD functions decreased in

halfwidth by 57%, but the location of functions with increasing levels of inhibition were not

significantly different from the model with only excitatory inputs (0.070-0.078 ms; centroid,

p=0.984; group data). However, the region of maximum slope consistently shifted towards

zero, in accordance with a sinking iceberg receptive field scenario (Fig. 7d; -0.27 to 0.07

ms). We then tested how the relative timing of excitatory and inhibitory inputs altered ITD

functions. Varying the timing of IPSPs relative to EPSPs from –0.5 ms to 0.25 ms altered

Fig. 7: Inhibition may serve to modulate a pre-established morphological baseline ITD response.
a, Schematic of innervation pattern with excitatory (green) and inhibitory (red) inputs. **b,** Somatic (top) and axonal (bottom) traces from ITD stimulation (ITD: 0.1 ms; IPSP timing: -0.5 ms) for each tested IPSP amplitude. Tested amplitude values and their unique colors above corresponding traces. Morphology of the exemplar neuron shown in the upper right. **c,** Exemplar cell's ITD response curves colored corresponding to IPSP amplitude in **b**. Centroid (square) and maximum slope (circle) are indicated for each curve. Dotted line at an ITD of zero. **d,** ITD centroid and maximum slope as a function of IPSP amplitude (colors corresponding to **b**). **e,f,** Group data (n=39) for ITD centroids (**e**) and steepest sections (**f**) for each IPSP amplitude. Data in **e** are largely overlapping. Shifting of population data in **f** corresponds to trend seen in the exemplar cell (**d**). **g-k,** Same as **b-f**, instead varying the timing of the inhibition (IPSP amplitude: 3 mV). Absolute firing probability, in **h**, is lower than **c**, resulting in lower signal to noise ratio. Centroids in **i** show more variation than **d**, though still minimal. Shift of exemplar cell's maximum slope ITD section (**i**) corresponds to degree of reduction in absolute firing probability (**h**).

the location of ITD functions by 0.04 ms, from 0.07 to 0.11 ms. Over the population of

MSO neuron morphologies, the presence of inhibition reduced firing probability without

fundamentally shifting the location of ITD functions. However, the region of the function

where there is maximal change in firing rate is necessarily impacted by the decrease in

ITD width (Fig. S4), suggesting that dendritic asymmetry and inhibition provide separate

contributions to MSO neurons' receptive field locations.

DISCUSSION

The passive cable properties of neuronal dendrites are known to impart both attenuation and delay on EPSPs as they propagate to the soma, commensurate with the distance traversed and the anatomical features of the dendrite itself. However, the functional significance of dendritic geometry and its impact on action potential firing is often difficult to assess. Here we have newly identified a critical function for dendritic structure in the processing of sound localization cues in MSO neurons of the auditory brainstem. We have shown that the bipolar dendritic arbor of MSO neurons is typically asymmetrical, imposing a corresponding asymmetry in the relative timing required for excitatory inputs to drive optimal patterns of action potential output. Our compartmental models show that variation in dendritic morphology across cells exerts a continuum of temporal influences sufficient to shift spatial receptive fields across the full biological range of ITDs available to the gerbil. These shifts are insensitive to differences in intensity and likely work together with other sources of internal delay to enable MSO neurons to stably encode spatial locations along the horizontal plane.

Role of dendritic-based delays in the detection of input sequences

The idea that dendritic structure could dictate detection of temporal sequences of input was first explored in the pioneering computer models of Wilfred Rall. These studies showed that attenuation and propagation delays of EPSPs traveling from the dendrites to the soma could impart greater sensitivity to somatically vs distally directed activation sequences of excitatory inputs, where the longer propagation time of more distal EPSPs is compensated by their earlier activation [37]. Preferential firing to centripetal (distal to

proximal) temporal sequences of uncaged glutamate at spines has been demonstrated

in cortical neurons in slices, but it is not clear what natural conditions would generate such

sequences [38]. By contrast, single excitatory axonal inputs to MSO dendrites mediate a

natural centripetal sequence of synaptic activity [15,28,39], and both current and previous

modeling results highlight the importance of matching of input sequence conduction

velocity with that of MSO dendrites, which conduct EPSPs at an average of ~0.3 ms^{-1}. In

the present study, with the aid of 2-photon fluorescence-guided recordings we have been

able to sample smaller diameter secondary and tertiary dendrites at longer distances from

the soma. Our results reveal that the attenuation and delays of EPSPs propagating from

these processes to the soma are more severe than previously estimated [40] . In our sample

of modeled MSO neurons, these slower-conducting non-primary dendrites make up 70%

of the surface area of the dendritic arbor in our sample of MSO neurons and thus play an

outsized role in exerting temporal shifts in ITD curves in neurons with asymmetrical

dendritic trees. These results emphasize that both branching and dendritic diameter are

critical determinants of dendritic delay, as seen reflected in the morphoelectrotonic

transforms (Fig. 5).

In the visual system, asymmetrical length and branching of the dendrites drives temporal

disparities in the pattern of inhibition of ganglion cells, providing directional tuning to

object movement [41–43]. The computation in the auditory system differs from vision in

several fundamental ways. In the retina, spatial information is directly represented at the

level of the photoreceptors, and tuning to spatial movement relies on the relative arrival

times of excitatory and inhibitory inputs to the dendrites and soma. By contrast, in the

cochlea space is not explicitly represented on the basilar membrane, which decomposes

complex sounds spatially into different frequency channels. Accordingly, horizontal spatial

receptive fields must be constructed *de novo* in the MSO through the detection of synaptic

binaural coincidence. In this way, ITDs serve as a proxy for horizontal position.

Other sources of asymmetry in binaural coincidence detection

Although our data reveal that the dendrites may contribute significant interaural delays in

the binaural circuitry in the MSO, they likely work together with other mechanisms that

have been proposed to influence binaural coincidence detection. In ~25% of MSO

neurons in gerbils, the axon emerges from either the lateral or medial dendrites [44,45], and

these results appear to extend to other mammalian species, including guinea pigs and

cats [46,47]. Results from computer models suggest these offset axons, like asymmetrical

dendrites, introduce strong asymmetries in synaptic integration and ITD tuning [48].

Inhibition has received considerable attention as a mechanism for ITD tuning based on

shifts of tuning curves measured in single-unit recordings *in vivo* upon iontophoretic block

of glycinergic synapses [18,19]. However, these results are controversial, as strong

inhibition has not been observed in both juxtacellular and intracellular patch recordings *in*

vivo [20,21,49], and there have been concerns about iontophoretic strychnine having off-

target effects that that alter both firing probability and input resistance [20]. We found that

inhibition did not shift the overall location of ITD curves across the functional range of

IPSP magnitudes *(*Fig. 7), consistent with pressure-applied strychnine data *in vivo* [20].

Inhibition shifted the location of maximal change in firing rate, consistent with a

conventional "iceberg" model of inhibitory action on neuronal receptive fields. In

mammals, the maximum rate of change, or slope of ITDs may be more important for

population encoding of ITDs. Given that the maximal rising slope of firing invariably shifts

toward 0 µs ITD when inhibition narrows ITD curves, both inhibition and dendritic

asymmetry would both play key roles in setting the location of individual MSO neurons'

receptive fields.

Comparison with avian binaural coincidence detectors.

Binaural coincidence detection has been extensively studied in birds, and thus the

comparison to mammals is noteworthy. Unlike mammals, dendritic morphology in bird

MSO (nucleus laminaris) shows striking dependence on tonotopic location, exhibiting

longer, more highly branched dendrites in neurons from low frequency regions [33,50]. While

these tonotopic differences in dendritic length have garnered considerable attention, a

close examination of dendritic parameters shows that nucleus laminaris neurons, like

those in mammalian MSO, exhibit a continuum of dendritic asymmetry in the length of

dendrites receiving ipsilateral and contralateral inputs (Smith and Rubel 1979, their Fig.

15 [33]). In these low frequency regions, inputs are restricted to distal, higher order

branches [51]. Based on both our electrophysiological and modeling data, these higher

order branches incur the largest delays as compared to the large-caliber primary

dendrites and thus would be the primary determinants of dendritic propagation delay on

each side. In the more highly specialized barn owls, the dendrites of laminaris neurons

are more limited, and thus internal delay compensation must lie squarely on those

provided by unmyelinated afferents [13] .

Dendritic morphology and the scaling problem across species.

The biological range of ITDs varies with head size [52,53] , and thus any mechanism of

internal delay must be scalable across species, accounting for ITDs differing by hundreds

of microseconds (e.g., gerbil: ±130 µs [54] , human: ± 600-700 µs [55]). A dendrite-based

source of internal delay would be easily scalable, as the length of MSO dendrites

increases with the size of different species. Although quantitative data are not available,

previous morphological studies in guinea pig, ferret, cat and humans show that the

dendritic arbors show considerable asymmetries [46,47,56,57] .

METHODS

All procedures were conducted in agreement with the Institutional Animal Care and Use

Committee at The University of Texas at Austin (UT-Austin), which followed the guidelines

of the National Institutes of Health. Mongolian gerbils (*Meriones unguiculatus*) were

housed and raised in a colony at the UT-Austin Animal Resource Center. Animals

experienced a 50/50 day/night cycle and had continuous access to food and water.

Brain slice preparation: both male and female gerbils (17-40 days old) were anesthetized

with isoflurane and decapitated upon loss of the choroid reflex. The brain was removed

from the cranium under warm (34-35°C) artificial cerebrospinal fluid (ACSF), the

cerebellum was removed and the isolated hindbrain was glued to the stage of an

oscillating tissue slicer (Leica VT-1200S). Horizontal sections were prepared at 200 µm

thickness and stored at 35°C in a holding chamber for 45-60 min., and then gradually

cooled to room temperature (24°C). All solutions were continuously bubbled with

95% O_2 / 5% CO_2. ACSF contained (in mM): 125 mM NaCl, 25 mM D-glucose, 2.5 mM

KCl, 25 mM NaHCO$_3$, 1.25 mM NaH$_2$PO$_4$, 1.5mM MgSO$_4$, 1.5 mM CaCl$_2$, pH adjusted to

7.45 with NaOH, and final osmolarity at 315 mOsm.

Imaging and electrophysiological acquisition and analyses: neurons in the MSO were

visualized under gradient contrast optics on a Leica TCS SP5 System equipped with a

Coherent Chameleon Ultra II Ti:sapphire laser and an 8 kHz resonant scanner under

control of Leica LAS AF Imaging Software. We found that gradient contrast images

acquired via the photomultiplier tubes were of poor contrast in the heavily myelinated

regions of the auditory brainstem, and so cell selection and seal formation was carried

out under gradient contrast imaging through the epifluorescence port of the microscope

and detected on a high-resolution monochrome digital camera (Zeiss Axiocam 503)

running Zen software. The soma was recorded in whole-cell current clamp mode with a

pipette (4 to 8 MΩ resistance) containing a potassium gluconate-based solution with 40-

80 µM Alexa 594 added for visualizing cell morphology. After waiting ~10 min. for the dye

to fill the dendritic arbor, a dendritic recording was made under 2-photon fluorescence

with a second pipette (8 to 15 MΩ resistance) containing the same K Gluconate solution

with 40-80 µM Alexa 488. Pipette solutions had a composition of 115 mM K-gluconate,

4.42 mM KCl, 0.5 mM EGTA, 10 mM HEPES, 10 mM, Na$_2$Phosphocreatine, 4 mM MgATP,

and 0.3 mM NaGTP. Osmolality was adjusted to 300 mOsm/L with sucrose, and pH was

adjusted to 7.30 with KOH. Seal resistances exceeded 1 GΩ, and recordings were

discontinued if series resistances exceeded 20 or 50 MΩ for somatic and dendritic

recordings, respectively.

Electrophysiological recordings were under the control of custom routines written in

IGOR-Pro (Wavemetrics). Pipettes were prepared on a Sutter P-1000 puller using

borosilicate glass (1.5 mm O.D.). Data was acquired using a pair of Dagan BVC-700,

amplifiers in conjunction with an ITC-18 computer interface (Heka Instruments). Data was

low-pass filtered at 5 kHz, sampled at 50 kHz, and stored on a Macintosh computer for

further analyses using custom-written routines in IGOR-Pro.

Compartment model construction:

A subpopulation of 40 fully intact reconstructed MSO neurons was selected from those

gathered in Bondy et al. (2021) [34] . The reconstructed morphologies were imported into

the NEURON simulation environment, using its Neurolucida importing tool.

Computational compartments were created every 2 µm (Fig. S5a). Somatic and dendritic

compartments were applied channel kinetics differentially, consisting of low and high-

voltage-activated K channels [27,58] , HCN channels[59] , and leak channels, with sodium

channels[60] exclusively in the somatic compartments. In simulations modeling a passive

cell, high and low-voltage potassium and HCN channels were replaced with passive leak

channels with conductances set to the average resting conductance of the preexisting

active channels. In simulations with an axon, a model axon, with morphology as in

Lehnert (2014) [61] , was attached at the center of the model soma. The axon channel

kinetics implemented consisted of sodium channels, low-threshold potassium channels

and leak channels in the nodes and initial segments, with solely leak channels in the

internodes. Measurements from the axon were all recorded from the fifth node. Simulated

synapses were conductance-based and shaped by an alpha function (τ = 290 µs).

Channel	Dendrite	Soma	TAIS	CAIS	Internode	Node
KLT (Mathews, 2010)	20	40	155	155		155
KHT (Nabel, 2019)	0.55	0.55				
HCN (Khurana, 2011)	1	1	2	2		
Na (Scott, 2010)	0	30	250	250		250
Leak	0.05	0.05	0.05	0.05	0.02	5

Table 1: Maximal channel conductances ($\mu S/cm^2$). Tapered axon initial segment (TAIS) and constant disameter axon initial segment (CAIS).

Simulation variations: When measuring EPSP propagation, synapses were placed every

2 µm, with a unitary conductance that would maximally depolarize the soma 1 mV (unless

noted otherwise). Each synapse was then stimulated and recorded separately. In ITD

coincident detection procedures, dendrites were innervated with synapses (release

probability=0.45) placed every 2 µm, totaling ~125 synapses for each side [62] . To mimic

axonal activation, an artificial axonal input delay (Δt_{axon}) was calculated for each synapse

as a function of conduction velocity (ν_{input}) and axonal travel distance. The travel distance

was considered zero at each side's furthest point from the soma. Otherwise, the axonal

travel distance consists of two components: the mediolateral distance from the furthest

point (Δt_{axon}=0) to the synapse's closest terminal segment; the path length from that

terminal segment to the synapse (Fig. S5b). Artificial ITDs could then be added onto the

calculated axonal delay. Each artificial ITD value (-0.5 to 0.5 ms, incremented by 0.01),

was tested with 400 trials, and each trial consisted of a single stimulation of all synapses.

To provide a more binary output, a model axon was attached to the cell's soma, and a

spike was reported if a 25 mV depolarization from rest was recorded at the fifth and final

axonal node. The synapses could then be adjusted with a unitary conductance to achieve

an acceptable maximum firing probability. When implementing IPSP inputs to the ITD

procedure, two inhibitory synapses were added at the soma, one corresponding to each

mediolateral side. Each synapse generated an IPSP shaped by an alpha function with a

time constant of 1.5 ms [62]. The timing of the synapses was adjusted relative to the

artificial excitatory ITD signals. The amplitude of the IPSPs was adjusted by modifying

the inhibitory synapses' conductances to achieve a combined IPSP hyperpolarization of

approximately 1, 2, and 3 mV.

Data analyses

Measures of dendritic distance, both in slice experiments and in compartment models,

were taken as the length along the dendritic arbor to the center of the somatic

compartment. To address the variable size of the MSO within the slice plane, a normalized

tonotopic location was calculated as the cell's distance from the most dorsal point of the

MSO, relative to the entire MSO dorsoventral length. To provide a quantitative value of

asymmetry between each cell's morphological measures, a unit of fold difference

$\left(\log_2 \left(\frac{\max\{\text{medial, lateral}\}}{\min\{\text{medial, lateral}\}}\right) + 1\right)$ was used, with a value of 1 and 2 representing equal and

double values, respectively. Average path length measurements were taken from the

distances along the dendrite to the soma from each terminal section of the corresponding

mediolateral arbor. Measurements for propagation attenuation and timing were taken at

both the synaptic and somatic compartments of the model cell. For attenuation, the

somatic depolarization for mediolateral sides was calculated by averaging each synaptic

compartments propagated EPSP amplitude at the soma. Morpho-electrotonic transforms

were created by scaling each sections length according to the natural logarithm of its

attenuation $\left(\text{section length} = \ln\left(\frac{V_{\text{dend}}}{V_{\text{soma}}}\right)\right)$. EPSP delay values were calculated by

measuring the time between peak depolarization in the synaptic and somatic

compartments. The corresponding delay transforms are scaled according to the delay

contribution of each section. Mediolateral comparisons of delay were calculated by

subtracting each sides average EPSP delay for all compartments. To minimize noise and

data filtering for the ITD response, the centroid of the curve was measured, where half of

the total area under the curve exists on either side. Comparisons were made using the

absolute ITD centroid values to highlight the population of neurons' collective change.

The ITD firing probabilities were smoothed with a Savitzky–Golay filter for the purpose of

reasonably identifying the region of max slope for the curves.

REFERENCES

1. Grothe, B., and Pecka, M. (2014). The natural history of sound localization in mammals--a story of neuronal inhibition. Front Neural Circuits *8*, 116. https://doi.org/10.3389/fncir.2014.00116.

2. Joris, P., and Yin, T.C. (2007). A matter of time: internal delays in binaural processing. Trends Neurosci *30*, 70–78. https://doi.org/10.1016/j.tins.2006.12.004.

3. Carr, C.E., Soares, D., Parameshwaran, S., and Perney, T. (2001). Evolution and development of time coding systems. Current Opinion in Neurobiology *11*, 727–733. https://doi.org/10.1016/s0959-4388(01)00276-8.

4. Stotler, W.A. (1953). An experimental study of the cells and connections of the superior olivary complex of the cat. J Comp Neurol *98*, 401–431.

5. Lindsey, B.G. (1975). Fine structure and distribution of axon terminals from the cochlear nucleus on neurons in the medial superior olivary nucleus of the cat. J Comp Neurol *160*, 81–103. https://doi.org/10.1002/cne.901600106.

6. Yin, T.C., and Chan, J.C. (1990). Interaural time sensitivity in medial superior olive of cat. J Neurophysiol *64*, 465–488.

7. Magnusson, A.K., Kapfer, C., Grothe, B., and Koch, U. (2005). Maturation of glycinergic inhibition in the gerbil medial superior olive after hearing onset. J Physiol *568*, 497–512. https://doi.org/10.1113/jphysiol.2005.094763.

8. Kapfer, C., Seidl, A.H., Schweizer, H., and Grothe, B. (2002). Experience-dependent refinement of inhibitory inputs to auditory coincidence-detector neurons. Nat Neurosci *5*, 247–253. https://doi.org/10.1038/nn810.

9. Spitzer, M.W., and Semple, M.N. (1995). Neurons sensitive to interaural phase disparity in gerbil superior olive: diverse monaural and temporal response properties. J Neurophysiol *73*, 1668–1690.

10. Goldberg, J.M., and Brown, P.B. (1969). Response of binaural neurons of dog superior olivary complex to dichotic tonal stimuli: some physiological mechanisms of sound localization. J Neurophysiol *32*, 613–636.

11. Jeffress, L.A. (1948). A place theory of sound localization. J Comp Physiol Psychol *41*, 35–39.

12. Carr, C.E., and Konishi, M. (1988). Axonal delay lines for time measurement in the owl's brainstem. Proc Natl Acad Sci U S A *85*, 8311–8315.

13. Carr, C., and Konishi, M. (1990). A circuit for detection of interaural time differences
in the brain stem of the barn owl.

14. Moiseff, A., and Konishi, M. (1981). Neuronal and behavioral sensitivity to binaural
time differences in the owl. J Neurosci *1*, 40–48. https://doi.org/10.1523/jneurosci.01-01-
00040.1981.

15. Beckius, G.E., Batra, R., and Oliver, D.L. (1999). Axons from anteroventral cochlear
nucleus that terminate in medial superior olive of cat: observations related to delay
lines. J Neurosci *19*, 3146–3161.

16. Smith, P.H., Joris, P.X., Carney, L.H., and Yin, T.C. (1991). Projections of
physiologically characterized globular bushy cell axons from the cochlear nucleus of the
cat. J Comp Neurol *304*, 387–407. https://doi.org/10.1002/cne.903040305.

17. Myoga, M.H., Lehnert, S., Leibold, C., Felmy, F., and Grothe, B. (2014). Glycinergic
inhibition tunes coincidence detection in the auditory brainstem. Nat Commun *5*, 3790.
https://doi.org/10.1038/ncomms4790.

18. Pecka, M., Brand, A., Behrend, O., and Grothe, B. (2008). Interaural time difference
processing in the mammalian medial superior olive: the role of glycinergic inhibition. J
Neurosci *28*, 6914–6925. https://doi.org/10.1523/jneurosci.1660-08.2008.

19. Brand, A., Behrend, O., Marquardt, T., McAlpine, D., and Grothe, B. (2002). Precise
inhibition is essential for microsecond interaural time difference coding. Nature *417*,
543–547. https://doi.org/10.1038/417543a.

20. Franken, T.P., Roberts, M.T., Wei, L., Golding, N.L., and Joris, P.X. (2015). In vivo
coincidence detection in mammalian sound localization generates phase delays. Nat
Neurosci *18*, 444–452. https://doi.org/10.1038/nn.3948.

21. Roberts, M.T., Seeman, S.C., and Golding, N.L. (2013). A mechanistic
understanding of the role of feedforward inhibition in the mammalian sound localization
circuitry. Neuron *78*, 923–935. https://doi.org/10.1016/j.neuron.2013.04.022.

22. Heijden, M. van der, Lorteije, J.A., Plauska, A., Roberts, M.T., Golding, N.L., and
Borst, J.G. (2013). Directional hearing by linear summation of binaural inputs at the
medial superior olive. Neuron *78*, 936–948.
https://doi.org/10.1016/j.neuron.2013.04.028.

23. Jercog, P.E., Svirskis, G., Kotak, V.C., Sanes, D.H., and Rinzel, J. (2010).
Asymmetric excitatory synaptic dynamics underlie interaural time difference processing
in the auditory system. PLoS Biol *8*, e1000406.
https://doi.org/10.1371/journal.pbio.1000406.

24. Shamma, S.A., Shen, N., and Gopalaswamy, P. (1989). Stereausis: binaural
processing without neural delays. The Journal of the Acoustical Society of America *86*,
989–1006. https://doi.org/10.1121/1.398734.

25. Plauška, A., Heijden, M.V.D., and Borst, J.G.G. (2017). A test of the stereausis
hypothesis for sound localization in mammals. Journal of Neuroscience *37*, 7278–7289.
https://doi.org/10.1523/jneurosci.0233-17.2017.

26. Agmon-Snir, H., Carr, C.E., and Rinzel, J. (1998). The role of dendrites in auditory
coincidence detection. Nature *393*, 268–272. https://doi.org/10.1038/30505.

27. Mathews, P.J., Jercog, P.E., Rinzel, J., Scott, L.L., and Golding, N.L. (2010). Control
of submillisecond synaptic timing in binaural coincidence detectors by K(v)1 channels.
Nat Neurosci *13*, 601–609. https://doi.org/10.1038/nn.2530.

28. Callan, A.R., Hess, M., Felmy, F., and Leibold, C. (2021). Arrangement of Excitatory
Synaptic Inputs on Dendrites of the Medial Superior Olive. J Neurosci *41*, 269–283.
https://doi.org/10.1523/jneurosci.1055-20.2020.

29. Franken, T.P., Bremen, P., and Joris, P.X. (2014). Coincidence detection in the
medial superior olive: Mechanistic implications of an analysis of input spiking patterns.
Frontiers in Neural Circuits *8*, 42. https://doi.org/10.3389/fncir.2014.00042.

30. Dasika, V.K., White, J.A., and Colburn, H.S. (2007). Simple models show the
general advantages of dendrites in coincidence detection. J Neurophysiol *97*, 3449–
3459. https://doi.org/10.1152/jn.00669.2005.

31. Feng, A.S., and Rogowski, B.A. (1980). Effects of monaural and binaural occlusion
on the morphology of neurons in the medial superior olivary nucleus of the rat. Brain
Research *189*, 530–534. https://doi.org/10.1016/0006-8993(80)90112-2.

32. Russell, F.A., and Moore, D.R. (1999). Effects of unilateral cochlear removal on
dendrites in the gerbil medial superior olivary nucleus. Eur J Neurosci *11*, 1379–1390.

33. Smith, D.J., and Rubel, E.W. (1979). Organization and development of brain stem
auditory nuclei of the chicken: dendritic gradients in nucleus laminaris. J Comp Neurol
186, 213–239. https://doi.org/10.1002/cne.901860207.

34. Bondy, B.J., Haimes, D.B., and Golding, N.L. (2021). Physiological diversity
influences detection of stimulus envelope and fine structure in neurons of the medial
superior olive. J Neurosci. https://doi.org/10.1523/jneurosci.2354-20.2021.

35. Hoffmeister, B., Jänig, W., and Lisney, S.J.W. (1991). A proposed relationship
between circumference and conduction velocity of unmyelinated axons from normal and
regenerated cat hindlimb cutaneous nerves. Neuroscience *42*, 603–611.
https://doi.org/10.1016/0306-4522(91)90402-a.

36. Werthat, F., Alexandrova, O., Grothe, B., and Koch, U. (2008). Experience-
dependent refinement of the inhibitory axons projecting to the medial superior olive. Dev
Neurobiol *68*, 1454–1462. https://doi.org/10.1002/dneu.20660.

37. Rinzel, J., and Rall, W. (1974). Transient response in a dendritic neuron model for
current injected at one branch. Biophys J *14*, 759–790. https://doi.org/10.1016/s0006-
3495(74)85948-5.

38. Branco, T., Clark, B.A., and Häusser, M. (2010). Dendritic discrimination of temporal
input sequences in cortical neurons. Science *329*, 1671–1675.
https://doi.org/10.1126/science.1189664.

39. Smith, P.H., Joris, P.X., and Yin, T.C. (1993). Projections of physiologically
characterized spherical bushy cell axons from the cochlear nucleus of the cat: evidence
for delay lines to the medial superior olive. J Comp Neurol *331*, 245–260.
https://doi.org/10.1002/cne.903310208.

40. Winters, B.D., Jin, S.X., Ledford, K.R., and Golding, N.L. (2017). Amplitude
Normalization of Dendritic EPSPs at the Soma of Binaural Coincidence Detector
Neurons of the Medial Superior Olive. J Neurosci *37*, 3138–3149.
https://doi.org/10.1523/jneurosci.3110-16.2017.

41. Bloomfield, S.A. (1994). Orientation-sensitive amacrine and ganglion cells in the
rabbit retina. Journal of Neurophysiology *71*, 1672–1691.
https://doi.org/10.1152/jn.1994.71.5.1672.

42. Trenholm, S., Johnson, K., Li, X., Smith, R.G., and Awatramani, G.B. (2011).
Parallel mechanisms encode direction in the retina. Neuron *71*, 683–694.
https://doi.org/10.1016/j.neuron.2011.06.020.

43. Bloomfield, S.A. (1991). Two types of orientation-sensitive responses of amacrine
cells in the mammalian retina. Nature *350*, 347–350. https://doi.org/10.1038/350347a0.

44. Scott, L.L., Mathews, P.J., and Golding, N.L. (2005). Posthearing developmental
refinement of temporal processing in principal neurons of the medial superior olive. J
Neurosci *25*, 7887–7895. https://doi.org/10.1523/jneurosci.1016-05.2005.

45. Rautenberg, P.L., Grothe, B., and Felmy, F. (2009). Quantification of the three-
dimensional morphology of coincidence detector neurons in the medial superior olive of
gerbils during late postnatal development. J Comp Neurol *517*, 385–396.
https://doi.org/10.1002/cne.22166.

46. Cajal., S.R.Y. (1907). Structure et connexions des neurones. Nordiskt Medicinskt
Arkiv *40*, 1–28. https://doi.org/10.1111/j.0954-6820.1907.tb00034.x.

47. Smith, P.H. (1995). Structural and functional differences distinguish principal from
nonprincipal cells in the guinea pig MSO slice. J. Neurophysiol. *73*, 1653–1667.
https://doi.org/10.1152/jn.1995.73.4.1653.

48. Zhou, Y., Carney, L.H., and Colburn, H.S. (2005). A model for interaural time
difference sensitivity in the medial superior olive: interaction of excitatory and inhibitory
synaptic inputs, channel dynamics, and cellular morphology. J Neurosci *25*, 3046–3058.
https://doi.org/10.1523/jneurosci.3064-04.2005.

49. Plauska, A., Borst, J.G., and Heijden, M. van der (2016). Predicting binaural
responses from monaural responses in the gerbil medial superior olive. J Neurophysiol
115, jn 01146 2015. https://doi.org/10.1152/jn.01146.2015.

50. Wang, Y., and Rubel, E.W. (2008). Rapid regulation of microtubule-associated
protein 2 in dendrites of nucleus laminaris of the chick following deprivation of afferent
activity. Neuroscience *154*, 381–389.
https://doi.org/10.1016/j.neuroscience.2008.02.032.

51. Yamada, R., and Kuba, H. (2021). Dendritic synapse geometry optimizes binaural
computation in a sound localization circuit. Science Advances *7*.
https://doi.org/10.1126/sciadv.abh0024.

52. Jones, H.G., Koka, K., Thornton, J.L., and Tollin, D.J. (2011). Concurrent
development of the head and pinnae and the acoustical cues to sound location in a
precocious species, the chinchilla (Chinchilla lanigera). J Assoc Res Otolaryngol *12*,
127–140. https://doi.org/10.1007/s10162-010-0242-3.

53. Koka, K., Jones, H.G., Thornton, J.L., Lupo, J.E., and Tollin, D.J. (2011). Sound
pressure transformations by the head and pinnae of the adult Chinchilla (Chinchilla
lanigera). Hear Res *272*, 135–147. https://doi.org/10.1016/j.heares.2010.10.007.

54. Maier, J.K., and Klump, G.M. (2006). Resolution in azimuth sound localization in the
Mongolian gerbil (Meriones unguiculatus) . The Journal of the Acoustical Society of
America *119*, 1029–1036. https://doi.org/10.1121/1.2159429.

55. George, K. (1977). Model for the intramural time differences in the azimuthal plane.
J. Acoust. Soc. Am *62*, 157–167.

56. Henkel, C.K., and Brunso-Bechtold, J.K. (1990). Dendritic morphology and
development in the ferret medial superior olivary nucleus. J Comp Neurol *294*, 377–388.
https://doi.org/10.1002/cne.902940307.

57. Kulesza, R. (2008). Cytoarchitecture of the human superior olivary complex: nuclei
of the trapezoid body and posterior tier. Hear Res *241*, 52–63.
https://doi.org/10.1016/j.heares.2008.04.010.

58. Nabel, A.L., Callan, A.R., Gleiss, S.A., Kladisios, N., Leibold, C., and Felmy, F.
(2019). Distinct distribution patterns of potassium channel sub-units in somato-dendritic
compartments of neurons of the medial superior olive. Frontiers in Cellular
Neuroscience *13*, 409471. https://doi.org/10.3389/fncel.2019.00038/bibtex.

59. Khurana, S., Liu, Z., Lewis, A.S., Rosa, K., Chetkovich, D., and Golding, N.L.
(2012). An essential role for modulation of hyperpolarization-activated current in the
development of binaural temporal precision. J Neurosci *32*, 2814–2823.
https://doi.org/10.1523/jneurosci.3882-11.2012.

60. Scott, L.L., Mathews, P.J., and Golding, N.L. (2010). Perisomatic voltage-gated
sodium channels actively maintain linear synaptic integration in principal neurons of the
medial superior olive. J Neurosci *30*, 2039–2050.
https://doi.org/10.1523/jneurosci.2385-09.2010.

61. Lehnert, S., Ford, M.C., Alexandrova, O., Hellmundt, F., Felmy, F., Grothe, B., and
Leibold, C. (2014). Action potential generation in an anatomically constrained model of
medial superior olive axons. J Neurosci *34*, 5370–5384.
https://doi.org/10.1523/jneurosci.4038-13.2014.

62. Couchman, K., Grothe, B., and Felmy, F. (2010). Medial superior olivary neurons
receive surprisingly few excitatory and inhibitory inputs with balanced strength and
short-term dynamics. J Neurosci *30*, 17111–17121.
https://doi.org/10.1523/jneurosci.1760-10.2010.