Dendritic delay lines shape the computation of sound location in neurons of the

gerbil medial superior olive

Authors/affiliations

Jared Casarez[1]*, Rebecca I. Voglewede[1]*, Bradley D. Winters[2], Ken Ledford[1] and Nace

L. Golding[1]

* Equal contributions

[1] The University of Texas at Austin, Department of Neuroscience and Center for Learning

and Memory, 1 University Station C7000, Austin TX 78712-0248, USA.

[2] Current address: Department of Biological Sciences and University Hospitals –

NEOMED Hearing Research Center, Northeast Ohio Medical University, Rootstown, OH,

44272, United States and Brain Health Research Institute, Kent State University, Kent,

OH, 44242, United States

Corresponding author and lead contact

Nace L. Golding: The University of Texas at Austin, Department of Neuroscience and

Center for Learning and Memory, 1 University Station C7000, Austin TX 78712-0248,

golding@austin.utexas.edu.

ABSTRACT

In mammals, neurons of the medial superior olive encode information about sounds arising from discrete spatial locations along the horizon. This tuning requires that an internal delay in the brain must offset acoustic disparities to ensure coincident arrival of excitatory inputs driven from the two ears. The source of this optimal internal delay, originally assumed to arise from axonal delay lines, is currently controversial in mammals. Here we use 2-photon guided paired dendritic and somatic recordings together with compartmental modeling of 40 complete MSO neuron morphologies to demonstrate that the dendrites themselves serve as a significant source of internal delay. We show that most MSO neurons exhibit morphological asymmetries that impose different EPSP delays across dendrites and shifts in optimal interaural time differences. Dendrite-based delays in the mammalian MSO are heterogeneous within each isofrequency laminae and provide a stable, structural mechanism to help tune individual neurons to sounds from different azimuthal locations.

KEYWORDS

dendrite; dendritic filtering; coincidence detection; auditory; sound localization.

INTRODUCTION

Spatial information in the auditory system is critical for humans and other mammals to assemble a 3-dimensional view of the world as well as to understand speech and communication signals, especially in environments with many sounds overlapping in time and in frequency. The auditory system is distinct from vision or touch in that there is no explicit peripheral representation of space, in this case in the cochlea, only a representation of frequency. Accordingly, the central auditory system must compute horizontal sound location indirectly by combining a series of monaural and binaural cues (Grothe and Pecka, 2014; Joris and Yin, 2007) . Spatial coding in the central auditory system is a prime example of a centrally synthesized or computational map.

Interaural time difference cues are used by birds and mammals to compute horizontal sound location at low frequencies (Carr et al., 2001) . In mammals this computation takes place in the medial superior olive (MSO), whose neurons are the first to receive converging excitatory and inhibitory inputs driven from the two ears. Intriguingly, in individual neurons, binaural inputs are integrated within a bipolar dendritic architecture, with ipsilateral and contralateral excitatory inputs segregated onto lateral and medial dendritic branches respectively (Lindsey, 1975; Stotler, 1953) . Ipsilateral and contralateral inhibition converges onto the soma and proximal dendrites (Kapfer et al., 2002; Lindsey, 1975; Magnusson et al., 2005; Stotler, 1953; Yin and Chan, 1990) . MSO neurons detect the optimal combination of these inputs, converting temporal coincidence into a rate code (Goldberg and Brown, 1969; Spitzer and Semple, 1995; Yin and Chan, 1990) .

The concept of internal delay is central to coincidence detection in the MSO, where

acoustic delays to the two ears are countered in time by a mechanism of delay inside the

brain. This internal temporal asymmetry was postulated in early models to arise from

axonal delay lines formed by ladder-like combinations of bilateral excitatory inputs (the

"Jeffress model") (Jeffress, 1948) . While the Jeffress model enjoys strong support in

birds (Carr and Konishi, 1990, 1988; Moiseff and Konishi, 1981) , comparable studies in

mammals have failed to reveal systematic axonal input delay lines nor any mapping of

delay along the nucleus (Beckius et al., 1999; Smith et al., 1991; Yin and Chan, 1990) .

More recently, there is evidence that fast inhibition mediates internal delay by shifting the

peaks of EPSPs (Brand et al., 2002; Myoga et al., 2014; Pecka et al., 2008) . However,

this mechanism is subject to vigorous debate (Franken et al., 2015; Heijden et al., 2013;

Roberts et al., 2013) , and it is not clear that this mechanism alone can explain the range

of receptive field locations in the MSO or their insensitivity to sound level. Other

mechanisms have been proposed to mediate internal binaural asymmetry, including

asymmetrical axon position, medio-lateral differences in the rising slopes of

EPSPs (Jercog et al., 2010), and differences in tuning of binaural inputs

("stereausis") (Plauška et al., 2017; Shamma et al., 1989).

All models of coincidence detection must address the issue of the temporal distortions

imposed by the dendrites themselves. However, to date models of MSO neurons have

been represented either as point neurons or as simplified, symmetrical "ball-and-stick"

models (Agmon-Snir et al., 1998; Callan et al., 2021; Dasika et al., 2007; Franken et al.,

2014; Mathews et al., 2010; Myoga et al., 2014) . Previous measurements of dendritic

propagation have revealed that EPSPs encounter delays in their propagation to the soma

up to ~150 μs, despite some degree of temporal compensation by activation of low

voltage-activated K channels in the soma and proximal dendrites (Mathews et al., 2010) .

However, these measurements have largely been restricted to the proximal half of the

dendritic arbor and have relied on averaging highly variable populations of neurons. Given

that the dendrites of MSO neurons alter their branching pattern and diameters during

early hearing and following disruptions of normal hearing (Feng and Rogowski, 1980;

Russell and Moore, 1999; Smith and Rubel, 1979), they might serve as a potential

mechanism for adjusting the timing requirements for sounds from the two ears to arrive

coincidently at the soma and axon.

Here we demonstrate that the dendritic arbors of MSO neurons are highly heterogeneous

and show a striking continuum of asymmetry across the medial and lateral branches that

receive segregated inputs driven respectively from the two ears. We show with dual 2-

photon fluorescence-guided patch recordings that inputs to smaller and/or more distal

dendrites show more substantial propagation delays than previously recognized, and

computational modeling of a large population of intact neuronal morphologies shows that

these dendritic asymmetries translate into stable internal delays that, as a population,

could account for the full range of naturally occurring horizontal locations. Finally, we

show that inhibition preserves the effects of dendritic asymmetry on interaural time

differences (ITDs) and helps bring the region of maximum slope of firing changes within

the ecologically relevant range. Thus dendrite-based delay lines likely work synergistically

with other integrative mechanisms purported to shape the perception of horizontal

position in mammals.

RESULTS

To understand how MSO neurons process interaural time differences in the dendrites, we made fluorescence-guided dendritic and somatic patch-clamp recordings visualized under 2-photon microscopy. Using this approach, we were able to target the distal third of the dendritic arbor, including secondary and tertiary branches, regions that have largely escaped investigation in these neurons. We then injected simulated EPSCs (τ_{rise} and τ_{decay}, 0.2 ms) into the dendritic pipette and recorded potentials simultaneously at both dendrites and soma. We observed strong variability in both the percent amplitude attenuation at the soma $\left(\left(\frac{V_{soma}}{V_{dend}}\right) \times 100\right)$ as well as the propagation delay of the EPSP,

Fig. 1: Heterogeneity in attenuation and timing of simulated excitatory synaptic potentials following propagation from distal dendritic locations to the soma.
a,b, 2-photon fluorescence-guided dual dendritic and somatic patch recordings from two MSO neurons of comparable distances (78 and 73 μm, respectively). Example neuron in **a₁** shows stronger attenuation of somatic propagation of a simulated EPSP (**a₂** vs. **b₂**) and longer peak delay (**a₃** vs. **b₃**). EPSC amplitudes: cell **a**, 600 pA; cell **b**, 400 pA. **c,d,** Group data for attenuation and peak delay for all paired MSO neuron recordings. There is a poor correlation of both measures as a function of distance.

measured as the difference in EPSP peak time at the soma relative to the dendrite (Fig. 1). Notably, the delays measured in these recordings ranged between 100 and 300 μs, exceeding the physiological range of ITDs measured in gerbils. In these recordings,

121 there was little correlation between local morphological features of recorded dendrites

122 and propagation delay of the EPSP, measured as the difference in the peak timing of

123 dendritic and somatic EPSPs (Fig. 2a-c). However, there was a stronger correlation

124 between the local dendritic membrane time constant and EPSP propagation delay (Fig.

125 2d). In addition, there was a weak trend of increasing dendritic membrane time constant

Fig. 2: EPSP propagation delays are not correlated with local morphology but rather more global physiological properties.
a, Closeups of distal dendritic morphology during paired dendritic and somatic recordings. **b,c**, Delay of peak EPSPs during propagation from dendrite to soma shows no significant correlation with local diameter or tapering rate of local dendrite. (dendritic EPSCs: 400-1000 pA). **d**, Peak EPSP delay from dendrite to soma was weakly correlated with the membrane time constant recorded at the dendritic recording site. **e**, Dendritic membrane time constant increased with distance of the recording site from the soma.

with recording distance (Fig. 2e), possibly because of proximity to the dendritic

terminations. These results may reflect that an understanding of the integrative properties

of the dendrites requires an analysis of the larger morphological and electrotonic structure

of the neuron.

To understand how overall dendritic structure impacts binaural coincidence detection, we

analyzed a database of biocytin-filled MSO neurons that exhibited complete dendritic

structures (40 of 270 neurons from Bondy et al., 2021 (Bondy et al., 2021) ; Fig. 3). We

quantified dendritic length, surface area and branching, with special emphasis on

comparisons of medial and lateral dendritic arbors. Higher-order (2° and higher) dendritic

branches were prominent in these neurons, comprising 70% of the membrane surface

area (Fig. S1).

A striking feature of most neurons in this data set was an overall asymmetry between the

lateral and medial dendritic arbors, reflecting differences in length, branching, and/or

diameter of the dendrites (Fig. 3a-c). We quantified this asymmetry according to two

Fig. 3: MSO neuron morphology is inhomogeneous and asymmetric, regardless of tonotopy.
a, Reconstructed morphologies rank ordered by tonotopy (left) and degree of asymmetry (right; average path length). Orderings span from low frequency (top) to high frequency (bottom) and medial (top) and lateral (bottom) asymmetry, respectively. Colored gradients added to visualize ordering. Uniquely colored dots placed beneath each morphology for differentiation. **b**, Fold difference between average path length of medial and lateral dendrites for all reconstructed morphologies (n=40) with vertical axes ordered by tonotopic location (top) or degree of asymmetry (bottom). Gradients added as in **a**. Colored points correspond to **a**. **c**, Same as **b**, except comparing total surface area of medial and lateral branches. Average medio-lateral path length and surface area fold differences (± SEM) were 1.080 ± 0.057 and 1.112 ± 0.133 (biased laterally) and tonotopic R^2 values were -0.42 and -0.23.

parameters, the average path length (average length from each dendritic end to the

soma) and membrane surface area, with the latter parameter reflecting the collective

influence of dendritic length, diameter and branching pattern. We then defined the degree

of bilateral asymmetry as fold differences between the larger vs. smaller dendritic

measure for each parameter. We observed no obvious correlation between dendritic

asymmetry and tonotopic location (Fig. 3a, *left* and 3b, *top*). However, we found that

dendritic asymmetry formed a finely graded continuum, with a slight bias toward lateral

dendrites (Fig. 3b, *bottom*). Interestingly, there were no significant differences in

population averages of the medio-lateral path length and surface area (1.080 ± 0.057 and

1.112 ± 0.133 fold differences; p=0.562), though there was a slight lateral bias. These

results indicate that the averaging of dendritic morphological parameters across neurons

obscures what is likely a key structural and computational feature.

To understand how morphological asymmetry impacts synaptic integration, we

constructed compartmental models using the same 40 reconstructions analyzed in Fig. 3.

We introduced an excitatory synaptic conductance (alpha function: τ = 290 µs) into each

dendritic compartment and then measured the voltage attenuation and propagation delay

Fig. 4: MSO morphology nonlinearly filters and attenuates propagating EPSPs through its passive cable properties.
a, Visualization EPSP filtering at different dendritic sites. Cell morphology (top) with similarly distant example synaptic sites indicated with green circles and corresponding numbers. Red triangle indicates the somatic compartment. Synaptic conductance indicated as g_{syn}. Four sets of paired dendritic (green) and somatic (red) traces (bottom) for each example synaptic site. Numbering of traces corresponds to labels on morphology above. Dendritic distance from soma for each site indicated below traces. **b**: Same procedure as in **a**, with an adjusted unitary synaptic conductance (see Methods), with measurements from each computational compartment. Model morphology and orientation same as in **a**. Comparison between active channel (left) and passive channel model (right). Recorded EPSP amplitude at synaptic (top, green) and somatic (bottom, red) compartment as a function of distance from soma. **c**, Actual (top) and electrotonically transformed (bottom) morphologies. Transformation visualizes degree of EPSP attenuation from each compartment (see Methods). **d**, Group data (n=40) of average somatic depolarization asymmetry between mediolateral sides. Gradient as in Fig 3. Dotted line indicates equal medial and lateral average somatic depolarizations. Cell from **a-c** indicated with a triangle.

from the dendritic input location to the soma (Figs. 4-5). At comparable distances from

the soma, EPSP amplitude at the site of synaptic input and following propagation to the

soma varied considerably, like the physiological data from paired recordings in Figure 1

(Fig. 4a). Plots of local dendritic and propagated somatic EPSP amplitudes across all

synapse locations further revealed striking asymmetries in propagation efficacy of EPSPs

to the soma (Fig. 4b, "Active cell"). In MSO neurons, the timing and attenuation of EPSPs

along dendrites are influenced both by passive cable filtering as well as interactions with

low voltage-activated K channels, which are active in the subthreshold voltage range in

these neurons (Mathews et al., 2010) . To determine the degree to which asymmetries in

low voltage-activated K channels impact EPSP attenuation at the soma we converted all

voltage-activated conductances to leak channels (Fig. 4b, "Passive cell"; see Methods).

While EPSPs propagated slightly more effectively to the soma from all dendritic locations

in the passive model, it did not fundamentally disrupt mediolateral asymmetry, indicating

that these differences arose primarily from the passive electrotonic architecture of the cell.

The branching structure had a significant effect on dendritic propagation asymmetry. This

could be better visualized in inward morphoelectrotonic transforms of cells, which exhibits

dendritic compartments according to their electrotonic, versus physical lengths. In the

example neuron in Fig. 4c, large primary dendrites are deemphasized, while thinner

secondary and tertiary branches appear as electrotonically elongated. Differences

between the medial and lateral dendritic arbors in this neuron and many others gave rise

to asymmetries in the magnitude of the average EPSP arriving at the soma.

Further, we observed that our population of cells formed a continuum in the degree of

somatic depolarization asymmetry, in line with the asymmetry in dendritic structure (Fig.

Fig. 5: EPSP timing significantly varies within and between cells, with morphological asymmetry informing overall cell timing.
a, EPSP delay visualization in two exemplar MSO neurons. Coloring of each cell corresponds to the propagation delay of EPSPs to the soma from its respective compartment. Legend of propagation delay coloring in top right. Cells are oriented mediolaterally according to the scale bar. Outline added to increase gradient readability. **b,** Same data as in **a**, except as a function of distance from the soma. Orientation of cells same as in **a**. **c,** Group data (n=40) of the difference in average EPSP propagation delay between medial and lateral dendrites. Cells 1 & 2 are indicated with same color as in **b**. Dotted line at 0 ms indicates an equal average EPSP delay between sides. Blue gradient along horizontal axis (as in Fig. 3) to indicate spectrum of asymmetry. **d,** Morphological transforms according to EPSP delays. Morphological centroids of actual cell morphologies (top) and transformed morphologies (bottom). Length corresponds to each sections EPSP delay contribution. Coloring of cell sections is consistent between actual and transformed morphologies for easier differentiation and comparison. Dotted line added to mark the midline.

179 4d). Dendritic asymmetry also translated into differences in average EPSP timing across

180 the medial and lateral dendrites. Figure 5a shows for two example neurons the spatial

181 profile of delays of EPSPs propagating from each dendritic compartment to the soma.

182 The asymmetry in dendritic timing can be more quantitatively visualized in plots of peak

183 EPSP delay ($t_{dend} - t_{soma}$) as a function of dendritic distance from the soma (Fig. 5b).

184 Dendritic branch points often introduced substantial jumps in propagation delays, and this

185 is reflected in these plots as increases in slope. To quantify the influence of dendritic

186 asymmetry on the optimal summation of bilateral EPSPs at the soma, we measured the

187 electrotonic delay at the soma for every 2 µm compartment of lateral and medial dendritic

188 subtrees. In our population of 40 neuron models, there were differences in the average

189 relative time of arrival of EPSPs to the soma from compartments in the medial and lateral

dendrites. Further, these differences formed a smoothly graded continuum of delays

(Fig. 5c).

MSO neurons exhibit a range of intrinsic electrical properties, and so we examined a

subset of neuron morphologies to understand the functional implications of this

physiological diversity. Using a neuron with an asymmetric dendritic arbor, we

systematically slowed the membrane time constant by reducing the density of both low

voltage-activated K channels and HCN channels (Fig. S2). In both cases, the timing of

EPSPs reaching the soma was linearly related to the value of the membrane time

constant. Thus, the effects of dendritic asymmetry on EPSP timing and amplitude at the

soma are expected to be substantially higher in MSO neurons with slower intrinsic

membrane properties. To understand how the differential timing of EPSPs propagating

from medial and lateral dendrites is translated into ITDs, we simulated binaural excitatory

input using a spiking model of MSO neurons (see Methods). This model featured one

axonal input per terminal branch segment as well as a centripetal (dendrite-to-soma)

pattern of dendritic innervation, running from the terminal compartments to the soma, in

accordance with results from anatomical studies (Fig. 6a) (Beckius et al., 1999; Callan et

al., 2021; Smith et al., 1991). The conduction velocity of these inputs was set to 1 ms^{-1},

approximating values of unmyelinated terminal axonal arborizations of similar

caliber (Carr and Konishi, 1990; Hoffmeister et al., 1991) . The timing of ipsilateral and

contralateral inputs was varied systematically, triggering spiking only when the two inputs

were in close coincidence (Fig. 6b). The location of these excitatory ITD functions were

dependent on medio-lateral asymmetry. Functions shifted toward the side of the

electrically longer dendrite, reflecting the need for earlier activation of that side to

compense for a longer dendritic propagation delay (Fig. 6c). Our population of 40

Fig. 6: Simulated ITDs demonstrate conservation of morphological impact on coincident detection.
a, Schematic visualization of ITD innervation. Somatic compartment and dendrites are shown in gray and black, respectively. Green bulbs represent excitatory synapses, with corresponding arrows to illustrate axonal input velocity (ν_{input}). Dotted lines illustrate point at which there is considered to be no axonal input delay (see Methods). Axon attached to soma, with initial segment, nodes, and internodes. **b**, Example somatic (top) and axonal (bottom) traces from ITD simulations. ITDs of -0.2 ms (left), 0.1 ms (middle, optimal coincidence), 0.3 ms (right) shown. Traces colored only for differentiation. Dotted line ($V_{threshold}$) indicates depolarization threshold for action potential consideration. Morphology of model cell indicated in upper right. **c**, Three example ITD firing curves from varied morphologies, medially biased (left), symmetrical (middle), and laterally biased (right). Corresponding morphologies imaged above. Light orange span represents range of ecologically possible ITDs for model species. Dashed line placed at ITD of 0 ms. Dot indicates centroid ITD value. **d,** Sorted group data (n=39) of centroid ITDs, with ν_{input} = 1 ms^{-1}. Cells from **c** indicated with filled colored points. **e,f**, Sorted group data (n=39) of absolute centroid ITDs, varying axonal input velocity (**e**) and peak firing probability (**f**). Cells from **c** indicated with correspondingly filled colored points.

neurons showed peak firing at delays that traversed a continuum of values (-0.11 to 0.08

ms, median -0.02; Fig. 6d; Fig. S3). The precise range of values depended in part on the

assumptions on the velocity of excitatory axons. The largest range of ITDs was achieved

when there was closer matching between the velocity of axonal input velocity and

postsynaptic EPSPs propagating from the dendrites to the soma, which maximized

temporal coincidence at the soma of spatially disparate inputs in the dendrites (Fig. 6e).

Finally, ITDs shifted less than 3% in the face of input amplitude changes that resulted in

firing probabilities between 0.2 and 0.8, indicating that dendritic asymmetry is highly

stable across changing stimulus conditions.

Inhibition is a critical component of ITD encoding, although its precise role is subject to

vigorous debate. To understand how dendritic asymmetry impacts the effects of inhibition,

we added bilateral inhibitory conductances to the somatic compartment of the bilateral

excitatory model in Fig. 6, consistent with the known somatically biased innervation of

MSO neurons from the lateral and medial nuclei of the trapezoid body (LNTB and

MNTB; (Kapfer et al., 2002; Werthat et al., 2008)). We set the timing of inhibitory

conductances to be 500 µs in advance of excitatory inputs from the same side, in

accordance with a previously published estimate (Roberts et al., 2013) . We then adjusted

the amplitude of inhibitory conductances to produce summed IPSPs of between ~1-3 mV,

which reduced firing probability by 20, 40, 60 and 100% (Fig. 7a). With increasing

amplitude of IPSPs, resultant ITD functions decreased in halfwidth by 57%, but the

location of functions with increasing levels of inhibition were not significantly different from

the model with only excitatory inputs (0.070-0.078 ms; centroid, p=0.984; group data).

However, the region of maximum slope consistently shifted towards zero, in accordance

with a sinking iceberg receptive field scenario (Fig. 7d; -0.27 to 0.07 ms). We then tested

how the relative timing of excitatory and inhibitory inputs altered ITD functions. Varying

the timing of IPSPs relative to EPSPs from –0.5 ms to 0.25 ms altered the location of ITD

functions by 0.04 ms, from 0.07 to 0.11 ms. Over the population of MSO neuron

morphologies, the presence of inhibition reduced firing probability without fundamentally

shifting the location of ITD functions. However, the region of the function where there is

maximal change in firing rate is necessarily impacted by the decrease in ITD width (Fig.

S4), suggesting that dendritic asymmetry and inhibition provide separate contributions to

MSO neurons' receptive field locations.

Fig. 7: Inhibition may serve to modulate a pre-established morphological baseline ITD response.
a, Schematic of innervation pattern with excitatory (green) and inhibitory (red) inputs. **b,** Somatic (top) and axonal (bottom) traces from ITD stimulation (ITD: 0.1 ms; IPSP timing: -0.5 ms) for each tested IPSP amplitude. Tested amplitude values and their unique colors above corresponding traces. Morphology of the exemplar neuron shown in the upper right. **c,** Exemplar cell's ITD response curves colored corresponding to IPSP amplitude in **b**. Centroid (square) and maximum slope (circle) are indicated for each curve. Dotted line at an ITD of zero. **d,** ITD centroid and maximum slope as a function of IPSP amplitude (colors corresponding to **b**). **e,f,** Group data (n=39) for ITD centroids (**e**) and steepest sections (**f**) for each IPSP amplitude. Data in **e** are largely overlapping. Shifting of population data in **f** corresponds to trend seen in the exemplar cell (**d**). **g-k,** Same as **b-f**, instead varying the timing of the inhibition (IPSP amplitude: 3 mV). Absolute firing probability, in **h**, is lower than **c**, resulting in lower signal to noise ratio. Centroids in **i** show more variation than **d**, though still minimal. Shift of exemplar cell's maximum slope ITD section (**i**) corresponds to degree of reduction in absolute firing probability (**h**).

DISCUSSION

The passive cable properties of neuronal dendrites are known to impart both attenuation

and delay on EPSPs as they propagate to the soma, commensurate with the distance

traversed and the anatomical features of the dendrite itself. However, the functional

significance of dendritic geometry and its impact on action potential firing is often difficult

to assess. Here we have newly identified a critical function for dendritic structure in the

processing of sound localization cues in MSO neurons of the auditory brainstem. We have

shown that the bipolar dendritic arbor of MSO neurons is typically asymmetrical, imposing

a corresponding asymmetry in the relative timing required for excitatory inputs to drive

optimal patterns of action potential output. Our compartmental models show that variation

in dendritic morphology across cells exerts a continuum of temporal influences sufficient

to shift spatial receptive fields across the full biological range of ITDs available to the

gerbil. These shifts are insensitive to differences in intensity and likely work together with

other sources of internal delay to enable MSO neurons to stably encode spatial locations

along the horizontal plane.

Role of dendritic-based delays in the detection of input sequences

The idea that dendritic structure could dictate detection of temporal sequences of input

was first explored in the pioneering computer models of Wilfred Rall. These studies

showed that attenuation and propagation delays of EPSPs traveling from the dendrites to

the soma could impart greater sensitivity to somatically vs distally directed activation

sequences of excitatory inputs, where the longer propagation time of more distal EPSPs

is compensated by their earlier activation (Rinzel and Rall, 1974) . Preferential firing to

centripetal (distal to proximal) temporal sequences of uncaged glutamate at spines has

been demonstrated in cortical neurons in slices, but it is not clear what natural conditions

would generate such sequences (Branco et al., 2010). By contrast, single excitatory

axonal inputs to MSO dendrites mediate a natural centripetal sequence of synaptic

activity (Beckius et al., 1999; Callan et al., 2021; Smith et al., 1993) , and both current

and previous modeling results highlight the importance of matching of input sequence

conduction velocity with that of MSO dendrites, which conduct EPSPs at an average of

~0.3 ms^{-1}. In the present study, with the aid of 2-photon fluorescence-guided recordings

we have been able to sample smaller diameter secondary and tertiary dendrites at longer

distances from the soma. Our results reveal that the attenuation and delays of EPSPs

propagating from these processes to the soma are more severe than previously

estimated (Winters et al., 2017) . In our sample of modeled MSO neurons, these slower-

conducting non-primary dendrites make up 70% of the surface area of the dendritic arbor

in our sample of MSO neurons and thus play an outsized role in exerting temporal shifts

in ITD curves in neurons with asymmetrical dendritic trees. These results emphasize that

both branching and dendritic diameter are critical determinants of dendritic delay, as seen

reflected in the morphoelectrotonic transforms (Fig. 5). Earlier work reported systematic

bilateral asymmetries in excitatory input dynamics and apparent synaptic reversal

potentials in MSO neurons (Jercog, 2008; Jercog et al., 2010), which were interpreted as

reflecting differences in electrotonic length and synaptic location between medial and

lateral dendrites. These functional asymmetries are consistent with the dendritic

propagation delays measured here, suggesting that dendritic filtering may underlie

previously observed excitatory timing differences.

In the visual system, asymmetrical length and branching of the dendrites drives temporal

disparities in the pattern of inhibition of ganglion cells, providing directional tuning to

object movement (Bloomfield, 1994, 1991; Trenholm et al., 2011). The computation in the

auditory system differs from vision in several fundamental ways. In the retina, spatial

information is directly represented at the level of the photoreceptors, and tuning to spatial

movement relies on the relative arrival times of excitatory and inhibitory inputs to the

dendrites and soma. By contrast, in the cochlea space is not explicitly represented on the

basilar membrane, which decomposes complex sounds spatially into different frequency

channels. Accordingly, horizontal spatial receptive fields must be constructed *de novo* in

the MSO through the detection of synaptic binaural coincidence. In this way, ITDs serve

as a proxy for horizontal position.

Other sources of asymmetry in binaural coincidence detection

Although our data reveal that the dendrites may contribute significant interaural delays in

the binaural circuitry in the MSO, they likely work together with other mechanisms that

have been proposed to influence binaural coincidence detection. In ~25% of MSO

neurons in gerbils, the axon emerges from either the lateral or medial

dendrites (Rautenberg et al., 2009; Scott et al., 2005) , and these results appear to extend

to other mammalian species, including guinea pigs and cats (Cajal., 1907; Smith, 1995) .

Results from computer models suggest these offset axons, like asymmetrical dendrites,

introduce strong asymmetries in synaptic integration and ITD tuning (Zhou et al., 2005) .

Inhibition has received considerable attention as a mechanism for ITD tuning based on

shifts of tuning curves measured in single-unit recordings *in vivo* upon iontophoretic block

of glycinergic synapses (Brand et al., 2002; Pecka et al., 2008) . However, these results

are controversial, as strong inhibition has not been observed in both juxtacellular and

intracellular patch recordings *in vivo* (Franken et al., 2015; Plauska et al., 2016; Roberts

et al., 2013), and there have been concerns about iontophoretic strychnine having off-

target effects that that alter both firing probability and input resistance (Franken et al.,

2015). We found that inhibition did not shift the overall location of ITD curves across the

functional range of IPSP magnitudes *(*Fig. 7), consistent with pressure-applied strychnine

data *in vivo* (Franken et al., 2015). Inhibition shifted the location of maximal change in

firing rate, consistent with a conventional "iceberg" model of inhibitory action on neuronal

receptive fields. In mammals, the maximum rate of change, or slope of ITDs may be more

important for population encoding of ITDs. Given that the maximal rising slope of firing

invariably shifts toward 0 µs ITD when inhibition narrows ITD curves, both inhibition and

dendritic asymmetry would both play key roles in setting the location of individual MSO

neurons' receptive fields.

Comparison with avian binaural coincidence detectors.

Binaural coincidence detection has been extensively studied in birds, and thus the

comparison to mammals is noteworthy. Unlike mammals, dendritic morphology in bird

MSO (nucleus laminaris) shows striking dependence on tonotopic location, exhibiting

longer, more highly branched dendrites in neurons from low frequency regions (Smith and

Rubel, 1979; Wang and Rubel, 2008). While these tonotopic differences in dendritic

length have garnered considerable attention, a close examination of dendritic parameters

shows that nucleus laminaris neurons, like those in mammalian MSO, exhibit a continuum

of dendritic asymmetry in the length of dendrites receiving ipsilateral and contralateral

inputs (Smith and Rubel 1979, their Fig. 15 [33]). In these low frequency regions, inputs are

restricted to distal, higher order branches (Yamada and Kuba, 2021). Based on both our

electrophysiological and modeling data, these higher order branches incur the largest

delays as compared to the large-caliber primary dendrites and thus would be the primary

determinants of dendritic propagation delay on each side. In the more highly specialized

barn owls, the dendrites of laminaris neurons are more limited, and thus internal delay

compensation must lie squarely on those provided by unmyelinated afferents (Carr and

Konishi, 1990) .

Dendritic morphology and the scaling problem across species.

The biological range of ITDs varies with head size (Jones et al., 2011; Koka et al., 2011) ,

and thus any mechanism of internal delay must be scalable across species, accounting

for ITDs differing by hundreds of microseconds (e.g., gerbil: ±130 µs (Maier and Klump,

2006) , human: ± 600-700 µs (George, 1977)). A dendrite-based source of internal delay

would be easily scalable, as the length of MSO dendrites increases with the size of

different species. Although quantitative data are not available, previous morphological

studies in guinea pig, ferret, cat and humans show that the dendritic arbors show

considerable asymmetries (Cajal., 1907; Henkel and Brunso-Bechtold, 1990; Kulesza,

2008; Smith, 1995) .

METHODS

All procedures were conducted in agreement with the Institutional Animal Care and Use

Committee at The University of Texas at Austin (UT-Austin), which followed the guidelines

of the National Institutes of Health. Mongolian gerbils (*Meriones unguiculatus*) were

housed and raised in a colony at the UT-Austin Animal Resource Center. Animals

experienced a 50/50 day/night cycle and had continuous access to food and water.

Brain slice preparation: both male and female gerbils (17-40 days old) were anesthetized

with isoflurane and decapitated upon loss of the choroid reflex. The brain was removed

from the cranium under warm (34-35°C) artificial cerebrospinal fluid (ACSF), the

cerebellum was removed and the isolated hindbrain was glued to the stage of an

oscillating tissue slicer (Leica VT-1200S). Horizontal sections were prepared at 200 µm

thickness and stored at 35°C in a holding chamber for 45-60 min., and then gradually

cooled to room temperature (24°C). All solutions were continuously bubbled with

95% O_2 / 5% CO_2. ACSF contained (in mM): 125 mM NaCl, 25 mM D-glucose, 2.5 mM

KCl, 25 mM $NaHCO_3$, 1.25 mM NaH_2PO_4, 1.5mM $MgSO_4$, 1.5 mM $CaCl_2$, pH adjusted to

7.45 with NaOH, and final osmolarity at 315 mOsm.

Imaging and electrophysiological acquisition and analyses: neurons in the MSO were

visualized under gradient contrast optics on a Leica TCS SP5 System equipped with a

Coherent Chameleon Ultra II Ti:sapphire laser and an 8 kHz resonant scanner under

control of Leica LAS AF Imaging Software. We found that gradient contrast images

acquired via the photomultiplier tubes were of poor contrast in the heavily myelinated

regions of the auditory brainstem, and so cell selection and seal formation was carried

out under gradient contrast imaging through the epifluorescence port of the microscope

and detected on a high-resolution monochrome digital camera (Zeiss Axiocam 503)

running Zen software. The soma was recorded in whole-cell current clamp mode with a

pipette (4 to 8 MΩ resistance) containing a potassium gluconate-based solution with 40-

80 μM Alexa 594 added for visualizing cell morphology. After waiting ~10 min. for the dye

to fill the dendritic arbor, a dendritic recording was made under 2-photon fluorescence

with a second pipette (8 to 15 MΩ resistance) containing the same K Gluconate solution

with 40-80 μM Alexa 488. Pipette solutions had a composition of 115 mM K-gluconate,

4.42 mM KCl, 0.5 mM EGTA, 10 mM HEPES, 10 mM, Na_2Phosphocreatine, 4 mM MgATP,

and 0.3 mM NaGTP. Osmolality was adjusted to 300 mOsm/L with sucrose, and pH was

adjusted to 7.30 with KOH. Seal resistances exceeded 1 GΩ, and recordings were

discontinued if series resistances exceeded 20 or 50 MΩ for somatic and dendritic

recordings, respectively.

Electrophysiological recordings were under the control of custom routines written in

IGOR-Pro (Wavemetrics). Pipettes were prepared on a Sutter P-1000 puller using

borosilicate glass (1.5 mm O.D.). Data was acquired using a pair of Dagan BVC-700,

amplifiers in conjunction with an ITC-18 computer interface (Heka Instruments). Data was

low-pass filtered at 5 kHz, sampled at 50 kHz, and stored on a Macintosh computer for

further analyses using custom-written routines in IGOR-Pro.

Compartment model construction:

A subpopulation of 40 fully intact reconstructed MSO neurons was selected from those

gathered in Bondy et al. (2021) (Bondy et al., 2021) . The reconstructed morphologies

were imported into the NEURON simulation environment, using its Neurolucida importing

tool. Computational compartments were created every 2 μm (Fig. S5a). Somatic and

dendritic compartments were applied channel kinetics differentially, consisting of low and

high-voltage-activated K channels (Mathews et al., 2010; Nabel et al., 2019) , HCN

channels(Khurana et al., 2012) , and leak channels, with sodium channels(Scott et al.,

2010) exclusively in the somatic compartments. In simulations modeling a passive cell,

high and low-voltage potassium and HCN channels were replaced with passive leak

channels with conductances set to the average resting conductance of the preexisting

active channels. In simulations with an axon, a model axon, with morphology as in

Lehnert (2014) (Lehnert et al., 2014) , was attached at the center of the model soma. The

axon channel kinetics implemented consisted of sodium channels, low-threshold

potassium channels and leak channels in the nodes and initial segments, with solely leak

channels in the internodes. Measurements from the axon were all recorded from the fifth

node. Simulated synapses were conductance-based and shaped by an alpha function (τ

= 290 µs).

Channel	Dendrite	Soma	TAIS	CAIS	Internode	Node
KLT (Mathews, 2010)	20	40	155	155		155
KHT (Nabel, 2019)	0.55	0.55				
HCN (Khurana, 2011)	1	1	2	2		
Na (Scott, 2010)	0	30	250	250		250
Leak	0.05	0.05	0.05	0.05	0.02	5

Table 1: Maximal channel conductances ($\mu S/cm^2$). Tapered axon initial segment (TAIS) and constant disameter axon initial segment (CAIS).

Simulation variations: When measuring EPSP propagation, synapses were placed every

2 µm, with a unitary conductance that would maximally depolarize the soma 1 mV (unless

noted otherwise). Each synapse was then stimulated and recorded separately. In ITD

coincident detection procedures, dendrites were innervated with synapses (release

probability=0.45) placed every 2 μm, totaling ~125 synapses for each side (Couchman et

al., 2010) . To mimic axonal activation, an artificial axonal input delay (Δt_{axon}) was

calculated for each synapse as a function of conduction velocity (v_{input}) and axonal travel

distance. The travel distance was considered zero at each side's furthest point from the

soma. Otherwise, the axonal travel distance consists of two components: the mediolateral

distance from the furthest point (Δt_{axon}=0) to the synapse's closest terminal segment; the

path length from that terminal segment to the synapse (Fig. S5b). Artificial ITDs could

then be added onto the calculated axonal delay. Each artificial ITD value (-0.5 to 0.5 ms,

incremented by 0.01), was tested with 400 trials, and each trial consisted of a single

stimulation of all synapses. To provide a more binary output, a model axon was attached

to the cell's soma, and a spike was reported if a 25 mV depolarization from rest was

recorded at the fifth and final axonal node. The synapses could then be adjusted with a

unitary conductance to achieve an acceptable maximum firing probability. When

implementing IPSP inputs to the ITD procedure, two inhibitory synapses were added at

the soma, one corresponding to each mediolateral side. Each synapse generated an

IPSP shaped by an alpha function with a time constant of 1.5 ms (Couchman et al.,

2010) . The timing of the synapses was adjusted relative to the artificial excitatory ITD

signals. The amplitude of the IPSPs was adjusted by modifying the inhibitory synapses'

conductances to achieve a combined IPSP hyperpolarization of approximately 1, 2, and

3 mV.

Data analyses

Measures of dendritic distance, both in slice experiments and in compartment models,

were taken as the length along the dendritic arbor to the center of the somatic

compartment. To address the variable size of the MSO within the slice plane, a normalized

tonotopic location was calculated as the cell's distance from the most dorsal point of the

MSO, relative to the entire MSO dorsoventral length. To provide a quantitative value of

asymmetry between each cell's morphological measures, a unit of fold difference

$\left(\log_2 \left(\frac{\max\{\text{medial, lateral}\}}{\min\{\text{medial, lateral}\}}\right) + 1\right)$ was used, with a value of 1 and 2 representing equal and

double values, respectively. Average path length measurements were taken from the

distances along the dendrite to the soma from each terminal section of the corresponding

mediolateral arbor. Measurements for propagation attenuation and timing were taken at

both the synaptic and somatic compartments of the model cell. For attenuation, the

somatic depolarization for mediolateral sides was calculated by averaging each synaptic

compartments propagated EPSP amplitude at the soma. Morpho-electrotonic transforms

were created by scaling each sections length according to the natural logarithm of its

attenuation $\left(\text{section length} = \ln\left(\frac{V_{\text{dend}}}{V_{\text{soma}}}\right)\right)$. EPSP delay values were calculated by

measuring the time between peak depolarization in the synaptic and somatic

compartments. The corresponding delay transforms are scaled according to the delay

contribution of each section. Mediolateral comparisons of delay were calculated by

subtracting each sides average EPSP delay for all compartments. To minimize noise and

data filtering for the ITD response, the centroid of the curve was measured, where half of

the total area under the curve exists on either side. Comparisons were made using the

absolute ITD centroid values to highlight the population of neurons' collective change.

The ITD firing probabilities were smoothed with a Savitzky–Golay filter for the purpose of

reasonably identifying the region of max slope for the curves.

REFERENCES

468
469

Agmon-Snir H, Carr CE, Rinzel J. 1998. The role of dendrites in auditory coincidence detection.
Nature **393**:268–272. DOI: https://doi.org/10.1038/30505, PMID: 9607764

Beckius GE, Batra R, Oliver DL. 1999. Axons from anteroventral cochlear nucleus that
terminate in medial superior olive of cat: observations related to delay lines. J Neurosci
19:3146–3161. PMID: 10191328

Bloomfield SA. 1994. Orientation-sensitive amacrine and ganglion cells in the rabbit retina.
Journal of Neurophysiology **71**:1672–1691. DOI: https://doi.org/10.1152/jn.1994.71.5.1672,
PMID: 8064341

Bloomfield SA. 1991. Two types of orientation-sensitive responses of amacrine cells in the
mammalian retina. Nature **350**:347–350. DOI: https://doi.org/10.1038/350347a0, PMID:
1706822

Bondy BJ, Haimes DB, Golding NL. 2021. Physiological diversity influences detection of
stimulus envelope and fine structure in neurons of the medial superior olive. J Neurosci. DOI:
https://doi.org/10.1523/jneurosci.2354-20.2021, PMID: 34083255

Branco T, Clark BA, Häusser M. 2010. Dendritic discrimination of temporal input sequences in
cortical neurons. Science (New York, N.Y.) **329**:1671–1675. DOI:
https://doi.org/10.1126/science.1189664

Brand A, Behrend O, Marquardt T, McAlpine D, Grothe B. 2002. Precise inhibition is essential
for microsecond interaural time difference coding. Nature **417**:543–547. DOI:
https://doi.org/10.1038/417543a, PMID: 12037566

Cajal. SRY. 1907. Structure et connexions des neurones. Nordiskt Medicinskt Arkiv **40**:1–28.
DOI: https://doi.org/10.1111/j.0954-6820.1907.tb00034.x

Callan AR, Hess M, Felmy F, Leibold C. 2021. Arrangement of Excitatory Synaptic Inputs on
Dendrites of the Medial Superior Olive. J Neurosci **41**:269–283. DOI:
https://doi.org/10.1523/jneurosci.1055-20.2020, PMID: 33208467

Carr C, Konishi M. 1990. A circuit for detection of interaural time differences in the brain stem of
the barn owl. The Journal of Neuroscience.

Carr CE, Konishi M. 1988. Axonal delay lines for time measurement in the owl's brainstem.
Proceedings of the National Academy of Sciences of the United States of America **85**:8311–
8315.

Carr CE, Soares D, Parameshwaran S, Perney T. 2001. Evolution and development of time
coding systems. Current Opinion in Neurobiology **11**:727–733. DOI:
https://doi.org/10.1016/s0959-4388(01)00276-8

Couchman K, Grothe B, Felmy F. 2010. Medial superior olivary neurons receive surprisingly few
excitatory and inhibitory inputs with balanced strength and short-term dynamics. J Neurosci
30:17111–17121. DOI: https://doi.org/10.1523/jneurosci.1760-10.2010, PMID: 21159981

Dasika VK, White JA, Colburn HS. 2007. Simple models show the general advantages of
dendrites in coincidence detection. J Neurophysiol **97**:3449–3459. DOI:
https://doi.org/10.1152/jn.00669.2005, PMID: 16914612

Feng AS, Rogowski BA. 1980. Effects of monaural and binaural occlusion on the morphology of
neurons in the medial superior olivary nucleus of the rat. Brain Research **189**:530–534. DOI:
https://doi.org/10.1016/0006-8993(80)90112-2, PMID: 7370789

Franken TP, Bremen P, Joris PX. 2014. Coincidence detection in the medial superior olive:
Mechanistic implications of an analysis of input spiking patterns. Frontiers in Neural Circuits
8:42. DOI: https://doi.org/10.3389/fncir.2014.00042, PMID: 24822037

Franken TP, Roberts MT, Wei L, Golding NL, Joris PX. 2015. In vivo coincidence detection in
mammalian sound localization generates phase delays. Nat Neurosci **18**:444–452. DOI:
https://doi.org/10.1038/nn.3948, PMID: 25664914

George K. 1977. Model for the intramural time differences in the azimuthal plane. J. Acoust.
Soc. Am **62**:157–167.

Goldberg JM, Brown PB. 1969. Response of binaural neurons of dog superior olivary complex
to dichotic tonal stimuli: some physiological mechanisms of sound localization. J
Neurophysiol **32**:613–636.

Grothe B, Pecka M. 2014. The natural history of sound localization in mammals--a story of
neuronal inhibition. Front Neural Circuits **8**:116. DOI:
https://doi.org/10.3389/fncir.2014.00116, PMID: 25324726

Heijden M van der, Lorteije JA, Plauska A, Roberts MT, Golding NL, Borst JG. 2013. Directional
hearing by linear summation of binaural inputs at the medial superior olive. Neuron **78**:936–
948. DOI: https://doi.org/10.1016/j.neuron.2013.04.028, PMID: 23764292

Henkel CK, Brunso-Bechtold JK. 1990. Dendritic morphology and development in the ferret
medial superior olivary nucleus. J Comp Neurol **294**:377–388. DOI:
https://doi.org/10.1002/cne.902940307, PMID: 2341617

Hoffmeister B, Jänig W, Lisney SJW. 1991. A proposed relationship between circumference and
conduction velocity of unmyelinated axons from normal and regenerated cat hindlimb
cutaneous nerves. Neuroscience **42**:603–611. DOI: https://doi.org/10.1016/0306-
4522(91)90402-a

Jeffress LA. 1948. A place theory of sound localization. J Comp Physiol Psychol **41**:35–39.
PMID: 18904764

Jercog P. 2008. Study of different physiological components that underlie interaural time delay
detection in the auditory system. NYU Department of Physics Dissertation.

Jercog PE, Svirskis G, Kotak VC, Sanes DH, Rinzel J. 2010. Asymmetric excitatory synaptic
dynamics underlie interaural time difference processing in the auditory system. PLoS Biol
8:e1000406. DOI: https://doi.org/10.1371/journal.pbio.1000406, PMID: 20613857

Jones HG, Koka K, Thornton JL, Tollin DJ. 2011. Concurrent development of the head and
pinnae and the acoustical cues to sound location in a precocious species, the chinchilla
(Chinchilla lanigera). J Assoc Res Otolaryngol **12**:127–140. DOI:
https://doi.org/10.1007/s10162-010-0242-3, PMID: 20957506

Joris P, Yin TC. 2007. A matter of time: internal delays in binaural processing. Trends Neurosci
30:70–78. DOI: https://doi.org/10.1016/j.tins.2006.12.004, PMID: 17188761

Kapfer C, Seidl AH, Schweizer H, Grothe B. 2002. Experience-dependent refinement of
inhibitory inputs to auditory coincidence-detector neurons. Nat Neurosci **5**:247–253. DOI:
https://doi.org/10.1038/nn810, PMID: 11850629

Khurana S, Liu Z, Lewis AS, Rosa K, Chetkovich D, Golding NL. 2012. An essential role for
modulation of hyperpolarization-activated current in the development of binaural temporal
precision. J Neurosci **32**:2814–2823. DOI: https://doi.org/10.1523/jneurosci.3882-11.2012,
PMID: 22357864

Koka K, Jones HG, Thornton JL, Lupo JE, Tollin DJ. 2011. Sound pressure transformations by
the head and pinnae of the adult Chinchilla (Chinchilla lanigera). Hear Res **272**:135–147.
DOI: https://doi.org/10.1016/j.heares.2010.10.007, PMID: 20971180

Kulesza R. 2008. Cytoarchitecture of the human superior olivary complex: nuclei of the
trapezoid body and posterior tier. Hear Res **241**:52–63. DOI:
https://doi.org/10.1016/j.heares.2008.04.010, PMID: 18547760

Lehnert S, Ford MC, Alexandrova O, Hellmundt F, Felmy F, Grothe B, Leibold C. 2014. Action
potential generation in an anatomically constrained model of medial superior olive axons. J
Neurosci **34**:5370–5384. DOI: https://doi.org/10.1523/jneurosci.4038-13.2014, PMID:
24719114

Lindsey BG. 1975. Fine structure and distribution of axon terminals from the cochlear nucleus
on neurons in the medial superior olivary nucleus of the cat. J Comp Neurol **160**:81–103.
DOI: https://doi.org/10.1002/cne.901600106, PMID: 1112923

Magnusson AK, Kapfer C, Grothe B, Koch U. 2005. Maturation of glycinergic inhibition in the
gerbil medial superior olive after hearing onset. J Physiol **568**:497–512. DOI:
https://doi.org/10.1113/jphysiol.2005.094763, PMID: 16096336

Maier JK, Klump GM. 2006. Resolution in azimuth sound localization in the Mongolian gerbil (
Meriones unguiculatus) . The Journal of the Acoustical Society of America **119**:1029–1036.
DOI: https://doi.org/10.1121/1.2159429, PMID: 16521765

Mathews PJ, Jercog PE, Rinzel J, Scott LL, Golding NL. 2010. Control of submillisecond
synaptic timing in binaural coincidence detectors by K(v)1 channels. Nat Neurosci **13**:601–
609. DOI: https://doi.org/10.1038/nn.2530, PMID: 20364143

Moiseff A, Konishi M. 1981. Neuronal and behavioral sensitivity to binaural time differences in
the owl. The Journal of Neuroscience: The Official Journal of the Society for Neuroscience
1:40–48. DOI: https://doi.org/10.1523/jneurosci.01-01-00040.1981

Myoga MH, Lehnert S, Leibold C, Felmy F, Grothe B. 2014. Glycinergic inhibition tunes
coincidence detection in the auditory brainstem. Nat Commun **5**:3790. DOI:
https://doi.org/10.1038/ncomms4790, PMID: 24804642

Nabel AL, Callan AR, Gleiss SA, Kladisios N, Leibold C, Felmy F. 2019. Distinct distribution
patterns of potassium channel sub-units in somato-dendritic compartments of neurons of the
medial superior olive. Frontiers in Cellular Neuroscience **13**:409471. DOI:
https://doi.org/10.3389/fncel.2019.00038/bibtex

Pecka M, Brand A, Behrend O, Grothe B. 2008. Interaural time difference processing in the
mammalian medial superior olive: the role of glycinergic inhibition. J Neurosci **28**:6914–6925.
DOI: https://doi.org/10.1523/jneurosci.1660-08.2008, PMID: 18596166

Plauska A, Borst JG, Heijden M van der. 2016. Predicting binaural responses from monaural
responses in the gerbil medial superior olive. J Neurophysiol **115**:jn 01146 2015. DOI:
https://doi.org/10.1152/jn.01146.2015, PMID: 27009164

Plauška A, Heijden MVD, Borst JGG. 2017. A test of the stereausis hypothesis for sound
localization in mammals. Journal of Neuroscience **37**:7278–7289. DOI:
https://doi.org/10.1523/jneurosci.0233-17.2017, PMID: 28659280

Rautenberg PL, Grothe B, Felmy F. 2009. Quantification of the three-dimensional morphology of
coincidence detector neurons in the medial superior olive of gerbils during late postnatal
development. J Comp Neurol **517**:385–396. DOI: https://doi.org/10.1002/cne.22166, PMID:
19760600

Rinzel J, Rall W. 1974. Transient response in a dendritic neuron model for current injected at
one branch. Biophysical Journal **14**:759–790. DOI: https://doi.org/10.1016/s0006-
3495(74)85948-5

Roberts MT, Seeman SC, Golding NL. 2013. A mechanistic understanding of the role of
feedforward inhibition in the mammalian sound localization circuitry. Neuron **78**:923–935.
DOI: https://doi.org/10.1016/j.neuron.2013.04.022, PMID: 23764291

Russell FA, Moore DR. 1999. Effects of unilateral cochlear removal on dendrites in the gerbil
medial superior olivary nucleus. Eur J Neurosci **11**:1379–1390.

Scott LL, Mathews PJ, Golding NL. 2010. Perisomatic voltage-gated sodium channels actively
maintain linear synaptic integration in principal neurons of the medial superior olive. J
Neurosci **30**:2039–2050. DOI: https://doi.org/10.1523/jneurosci.2385-09.2010, PMID:
20147532

Scott LL, Mathews PJ, Golding NL. 2005. Posthearing developmental refinement of temporal
processing in principal neurons of the medial superior olive. J Neurosci **25**:7887–7895. DOI:
https://doi.org/10.1523/jneurosci.1016-05.2005, PMID: 16135745

Shamma SA, Shen N, Gopalaswamy P. 1989. Stereausis: binaural processing without neural
delays. The Journal of the Acoustical Society of America **86**:989–1006. DOI:
https://doi.org/10.1121/1.398734, PMID: 2794252

Smith DJ, Rubel EW. 1979. Organization and development of brain stem auditory nuclei of the
chicken: dendritic gradients in nucleus laminaris. J Comp Neurol **186**:213–239. DOI:
https://doi.org/10.1002/cne.901860207, PMID: 447882

Smith PH. 1995. Structural and functional differences distinguish principal from nonprincipal
cells in the guinea pig MSO slice. Journal of Neurophysiology **73**:1653–1667. DOI:
https://doi.org/10.1152/jn.1995.73.4.1653, PMID: 7643173

Smith PH, Joris PX, Carney LH, Yin TC. 1991. Projections of physiologically characterized
globular bushy cell axons from the cochlear nucleus of the cat. J Comp Neurol **304**:387–407.
DOI: https://doi.org/10.1002/cne.903040305, PMID: 2022755

Smith PH, Joris PX, Yin TC. 1993. Projections of physiologically characterized spherical bushy
cell axons from the cochlear nucleus of the cat: evidence for delay lines to the medial
superior olive. J Comp Neurol **331**:245–260. DOI: https://doi.org/10.1002/cne.903310208,
PMID: 8509501

Spitzer MW, Semple MN. 1995. Neurons sensitive to interaural phase disparity in gerbil superior
olive: diverse monaural and temporal response properties. J Neurophysiol **73**:1668–1690.

Stotler WA. 1953. An experimental study of the cells and connections of the superior olivary
complex of the cat. J Comp Neurol **98**:401–431.

Trenholm S, Johnson K, Li X, Smith RG, Awatramani GB. 2011. Parallel mechanisms encode
direction in the retina. Neuron **71**:683–694. DOI:
https://doi.org/10.1016/j.neuron.2011.06.020, PMID: 21867884

Wang Y, Rubel EW. 2008. Rapid regulation of microtubule-associated protein 2 in dendrites of
nucleus laminaris of the chick following deprivation of afferent activity. Neuroscience
154:381–389. DOI: https://doi.org/10.1016/j.neuroscience.2008.02.032

Werthat F, Alexandrova O, Grothe B, Koch U. 2008. Experience-dependent refinement of the
inhibitory axons projecting to the medial superior olive. Dev Neurobiol **68**:1454–1462. DOI:
https://doi.org/10.1002/dneu.20660, PMID: 18777566

Winters BD, Jin SX, Ledford KR, Golding NL. 2017. Amplitude Normalization of Dendritic
EPSPs at the Soma of Binaural Coincidence Detector Neurons of the Medial Superior Olive.
J Neurosci **37**:3138–3149. DOI: https://doi.org/10.1523/jneurosci.3110-16.2017, PMID:
28213442

Yamada R, Kuba H. 2021. Dendritic synapse geometry optimizes binaural computation in a
sound localization circuit. Science Advances **7**. DOI: https://doi.org/10.1126/sciadv.abh0024,
PMID: 34818046

Yin TC, Chan JC. 1990. Interaural time sensitivity in medial superior olive of cat. J Neurophysiol
64:465–488. PMID: 2213127

Zhou Y, Carney LH, Colburn HS. 2005. A model for interaural time difference sensitivity in the
medial superior olive: interaction of excitatory and inhibitory synaptic inputs, channel
dynamics, and cellular morphology. J Neurosci **25**:3046–3058. DOI:
https://doi.org/10.1523/jneurosci.3064-04.2005, PMID: 15788761

658