Structural mechanism of Necrocide 1 activation of human TRPM4 that triggers necrosis by sodium overload

Celso M. Teixeira-Duarte[1,2,#], Wan Fu[3,#], Weizhong Zeng[1,2], Jianghuang Wang[3], Xinzhe Jiang[3], Ziye Zhao[3], Qing Zhong[3,*] & Youxing Jiang[1,2,*]

Affiliations:

[1] Howard Hughes Medical Institute and Department of Physiology, University of Texas Southwestern Medical Center, Dallas, Texas, USA

[2] Department of Biophysics, University of Texas Southwestern Medical Center, Dallas, Texas, USA

[3] Institute for Translational Medicine on Cell Fate and Disease, Shanghai Ninth People's Hospital, Key Laboratory of Cell Differentiation and Apoptosis of National Ministry of Education, Department of Pathophysiology, Shanghai Jiao Tong University School of Medicine, Shanghai, China

[#] These authors contribute equally.

[*]Correspondence to:

Youxing Jiang, Ph.D., Department of Physiology, UT Southwestern Medical Center, 5323 Harry Hines Blvd., Dallas, Texas 75390-9040, Tel. 214 645-6027; Fax. 214 645-6042; E-Mail: youxing.jiang@utsouthwestern.edu

Qing Zhong, Ph.D., Department of Pathophysiology, Shanghai Jiao Tong University School of Medicine; Tel. 021-63846590-776429; E-Mail: qingzhong@shsmu.edu.cn

Abstract

The small molecule Necrocide 1 (NC1) constitutively activates human TRPM4, triggering Na^+ influx and leading to necrotic cell death, a process termed *Necrosis by Sodium Overload* (NECSO). NC1 activation is specific to human TRPM4 and does not affect most of the other mammalian TRPM4 orthologs. Here, we elucidate the molecular mechanism underlying NC1 activation and its species-specific selectivity for human TRPM4 using a combination of single-particle cryo-EM, electrophysiology, and cell death assays. We demonstrate that NC1 functions as a non-competitive surrogate of the endogenous Ca^{2+} ligand – it binds to a pocket within the S1-S4 domain adjacent to the Ca^{2+} site and induces the same conformational changes as those triggered by Ca^{2+}. Like Ca^{2}-mediated activation, NC1-induced channel opening also requires membrane $PI(4,5)P_2$ to stabilize the open state. Through comparative mutagenesis and structural analysis of human and mouse TRPM4, we identify the molecular determinants of NC1 specificity. Our results reveal that the insensitivity of mouse TRPM4 to NC1 arises not from a lack of binding, but from drug-induced conformational changes that destabilize the selectivity filter and inactivate the channel. We identify three critical residues that confer NC1 sensitivity, and their substitution renders mouse TRPM4 responsive to NC1, akin to the human channel. Given the upregulation of TRPM4 in various human cancers, our mechanistic insights into NC1 activation and specificity provide a framework for the potential development of cancer therapeutics targeting TRPM4-mediated necrosis.

Introduction

TRPM4 is a Ca^{2+}-activated, monovalent-selective cation channel belonging to the transient receptor potential melastatin (TRPM) subfamily [1-3]. It is ubiquitously expressed [4] and plays a key role in regulating various physiological processes, including cardiac conduction [5-8], immune response [9], and cell death [10-14]. Notably, TRPM4 is upregulated in several cancer types [15], such as prostate [16-18], breast [19,20], diffuse large B-cell lymphoma [21], and acute myeloid leukemia [22], making it an attractive target for therapeutic intervention. Despite this potential, commonly used TRPM4 modulators suffer from low specificity [5,23-26], prompting extensive efforts to develop monoclonal antibodies [27-30] and small molecules [31,32] specific for this channel. Among the latter, anthranilic acid derivatives, CBA, NBA, and IBA, have emerged as promising inhibitors [31,33,34]. Recent structural studies revealed that NBA and IBA bind in the vanilloid binding pocket, adjacent to the PI(4,5)P₂ binding site, to exert their inhibitory effects [34,35].

TRPM4 is activated by intracellular calcium and PI(4,5)P₂, promoting membrane depolarization via monovalent cation influx [1,23]. Interestingly, TRPM4 channel opening has been implicated in necrotic cell death triggered by anticancer treatments in human breast cancer cells [12]. Necrocide 1 (NC1) ((S')-3-cycloheptyl-3-(4-hydroxyphenyl)-6-methoxy-7-methyl-1,3-dihydroindole-2-one) is a small molecule derived from the optimization of a compound identified in a screen for anti-tumorigenic activity [36,37]. NC1 induces necrotic cell death in several cancer cell lines, including PC-3 (prostate) and MCF-7 (breast), and has been shown to suppress tumor growth in human cancer xenografts [38]. Importantly, NC1 does not trigger necrosis in non-tumorigenic cells such as fibroblasts and umbilical vein endothelial cells [38].

Recent findings demonstrate that NC1 induces necrotic cell death by directly activating human TRPM4, triggering Na^+ influx that leads to osmotic swelling and membrane rupture, a process termed Necrosis by Sodium Overload (NECSO) [39]. Interestingly, NC1 exhibits a high degree of species specificity: it activates human TRPM4 but not most other mammalian TRPM4 orthologues or other human TRPM family members. Since evading apoptosis is a hallmark of cancer cells [40,41], targeting alternative, non-apoptotic pathways such as TRPM4-mediated necrosis may offer novel therapeutic avenues. Understanding the molecular mechanism of NC1-induced human TRPM4 activation and its species specificity is therefore critical for the rational development of TRPM4-targeted anticancer therapies.

In this study, we combine single-particle cryo-EM, electrophysiology, and cell death assays to elucidate the structural mechanism by which NC1 activates human TRPM4. We identify the NC1 binding site and key molecular determinants responsible for channel activation. In addition, we explain the insensitivity of mouse TRPM4 to NC1 and pinpoint specific residues that define NC1 specificity for human TRPM4, offering insights into the species selectivity of NC1 and paving the way for future therapeutic development.

Results

NC1 activates hTRPM4 and promotes cell death

The small molecule Necrocide 1 (NC1) was recently shown to induce necrotic cell death by activating the human TRPM4 (hTRPM4) channel [39]. The key electrophysiological features of NC1-activated hTRPM4 were characterized using whole-cell patch-clamp recordings of hTRPM4-expressing HEK293 cells (Methods). NC1 activates TRPM4 slowly, requiring about two minutes for the current to reach its maximum, indicating a slow on-rate of NC1 binding (Fig. 1a and 1b). Most channels remain activated even after several minutes of washout, suggesting high-affinity binding and an extremely slow off-rate. With the presence of 1 mM EGTA in the pipette solution (cytosolic side), our recording conditions eliminate the potential contribution to channel activation from cytosolic Ca^{2+}, the endogenous ligand of TRPM4, confirming the Ca^{2+}-independence of NC1 activation.

To complement electrophysiological data, we conducted cell death and viability assays to assess hTRPM4-dependent necrosis in HeLa cells (Methods). While HeLa cells are insensitive to NC1 due to a lack of detectable endogenous TRPM4, they become susceptible to NC1-induced necrosis upon transient or stable expression of hTRPM4. In the cell death assay, EGFP-tagged hTRPM4 was transiently expressed in HeLa cells, and necrotic morphology, characterized by 'balloon-like' cell swelling, was observed 24 hours after NC1 treatment (Fig. 1c). No necrosis was observed in the NC1-treated control cells transiently expressing EGFP or EGFP-tagged mouse TRPM4. This assay provides a qualitative readout of hTRPM4-dependent necrosis, enabling the screening of mutational effects on the NC1 sensitivity of the channel, as discussed later. In the cell viability assay, hTRPM4 was stably expressed in HeLa cells. Following 24-hour NC1 treatment

at varying concentrations, cytotoxicity was assessed by measuring intracellular ATP levels,

indicative of cell viability, using the CellTiter-Glo (CTG) assay (Fig. 1d, Methods). This assay

provides a quantitative measure of NC1's dose-dependent toxicity in hTRPM4-expressing cells.

Structure of NC1-bound TRPM4

To elucidate the structural basis of NC1 binding and its activation mechanism, we purified

detergent-solubilized hTRPM4 and determined its structure in complex with NC1 at 2.7 Å

resolution using single-particle cryo-EM (Fig. 2a, Fig. S1a, Table S1, and Methods). The NC1

ligand is unambiguously resolved in the EM density map (Fig. 2b). Occupying a pocket within the

S1–S4 domain just above the Ca^{2+} activation site, NC1 forms extensive hydrophobic and

hydrogen-bonding interactions with surrounding residues (Fig. 2b).

To validate NC1 binding, we performed mutagenesis of residues interacting with the ligand.

Due to the extensive contact surface between NC1 and its binding pocket, single-alanine

substitutions were insufficient to disrupt NC1 binding and channel activation (Fig. S2a). We

therefore introduced bulkier tryptophan substitutions to block NC1 access sterically. Among all

tested, three hTRPM4 single-tryptophan mutants - Y790W, L824W, and I909W – completely lost

sensitivity to NC1 (Fig. 2c and 2d). These mutants remained responsive to Ca^{2+}/PI(4,5)P$_2$

activation, particularly I909W (Fig. S2b), confirming that these mutations prevent NC1 binding

while retaining native activation. The loss of NC1 activation in these mutants was further

supported by cell viability assays using stable HeLa cell lines expressing either wild-type or mutant

hTRPM4. While NC1 treatment induced a dose-dependent loss of viability in WT-expressing cells,

the mutant-expressing cells were resistant to NC1-induced necrosis (Fig. 2e).

Despite robust NC1 activation observed in functional assays, the NC1-bound TRPM4

structure adopts a conformation intermediate between the closed and open states, when compared

to previously determined apo (closed) and Ca^{2+}/PI(4,5)P$_2$-bound (open) structures (Fig. 3a and Fig.

S3a). NC1 binding induces local conformational changes in the S1-S4 domain resembling those

triggered by Ca^{2+} (Fig. 3b and Fig. S3b); however, the channel pore and the cytosolic domains

remain in a closed conformation (Fig. 3a, 3c, 3d, and Fig. S3a). Within S1-S4, NC1 binding causes

the S3 helix to shift inward toward the NC1-binding pocket, moving the W864 side chain closer

to the C-terminal end of S4 (Fig. 3b). Through tight packing with H908 on S4, W864 motion

displaces the C-terminal part of S4 toward S5 in the adjacent pore domain (Fig. 3b). In the apo

TRPM4 structure, F910 at the C-terminal part of S4 forms a key contact with F935 near the

cytosolic end of S5 - a coupling essential for transducing Ca^{2+}-induced movements into pore

opening, as shown in the $Ca^{2+}/PI(4,5)P_2$-bound structure [35] (Fig. 3e). In contrast, the NC1-bound

structure shows disordered side chains for both F910 and F935, indicating a loss of this critical

contact (Fig. 3e). These structural features of NC1-bound hTRPM4, including an activated S1-S4

domain, a closed pore, and disrupted F910–F935 coupling, mirror the Ca^{2+}-bound hTRPM4

structure in the absence of $PI(4,5)P_2$, previously proposed to represent a desensitized state [35] (Fig

3e and Fig. S3c). Since $PI(4,5)P_2$ binding is required to stabilize the open state upon Ca^{2+} activation

[35], the resemblance between NC1- and Ca^{2+}-bound structures suggests that NC1 triggers local

conformational changes similar to those induced by Ca^{2+} and also relies on $PI(4,5)P_2$ binding to

stabilize the open channel.

$PI(4,5)P_2$-dependent NC1 activation of hTRPM4

To test whether $PI(4,5)P_2$ is indeed required for NC1 activation, we performed patch-clamp

recordings of hTRPM4-expressing HEK293 cells in the inside-out configuration. Following initial

activation by cytosolic Ca^{2+}, TRPM4 undergoes desensitization to a steady-state level with reduced

open probability due to depletion of membrane $PI(4,5)P_2$, caused by Ca^{2+}-induced phospholipase

C activation [42,43] (Fig. 4a). This loss of $PI(4,5)P_2$ after Ca^{2+} activation creates a condition suitable

for assessing NC1 activation in the absence of $PI(4,5)P_2$. Indeed, hTRPM4 becomes unresponsive

to NC1 following $PI(4,5)P_2$ depletion (Fig. 4a). While $PI(4,5)P_2$ alone cannot activate TRPM4, its

addition in the presence of NC1 robustly activates the channel, restoring current to the pre-

desensitization level (Fig. 4a). Consistent with whole-cell recordings, NC1 activates TRPM4

slowly in the presence of $PI(4,5)P_2$, and the ligand remains tightly bound, making it difficult to

wash out (Fig. 4b). These results indicate that $PI(4,5)P_2$ is a prerequisite for NC1-induced

activation of hTRPM4.

To further support this conclusion, we mutated key positively charged $PI(4,5)P_2$-interacting

residues, R664A, K925A, and K928A, previously identified in the $Ca^{2+}/PI(4,5)P_2$-activated

TRPM4 structure [35]. These mutations significantly impaired NC1 activation in whole-cell

recordings, with much stronger effects on inward currents at negative potentials (Fig. 4c and 4d).

Additionally, HeLa cells overexpressing these mutants were no longer sensitive to NC1-induced

cell death, further validating the requirement of PI(4,5)P$_2$ for NC1-mediated TRPM4 activation

(Fig. 4e and Fig. S4).

The PI(4,5)P$_2$ dependence of NC1-induced hTRPM4 activation was further confirmed by

determining the structure of the channel bound to both PI(4,5)P$_2$ and NC1 at 2.8 Å resolution (Fig.

5a and 5b, Fig. S1b, Table S1, and Methods). In this structure, hTRPM4 adopts an open

conformation virtually identical to that of the Ca^{2+}/PI(4,5)P$_2$-activated channel (Fig. 5a). Compared

to the structure with NC1 alone, the presence of PI(4,5)P$_2$ enables the NC1-induced conformational

changes within the S1-S4 domain to be effectively transmitted to the pore through the reformation

of the critical F910/F935 contact, thereby leading to pore opening (Fig. 3c and 3d and Fig. 5c and

5d). This pore-opening movement is further coupled to large-scale rotational and upward swinging

motions of the cytosolic domains via the TRP domain, as also observed in the Ca^{2+}/PI(4,5)P$_2$-

activated structure [35] (Fig. 5c and 5e). Interestingly, despite no exogenous Ca^{2+} being added, the

canonical Ca^{2+}-binding site within the S1-S4 domain is occupied by a Ca^{2+} ion in the

NC1/PI(4,5)P$_2$-bound structure (Fig. 5b). The stabilized, active conformation of S1-S4 induced by

NC1 and PI(4,5)P$_2$ likely increases the apparent Ca^{2+} affinity, allowing residual micromolar Ca^{2+}

in nominally Ca^{2+}-free conditions to bind. To confirm that Ca^{2+} is not required for channel opening

in the NC1/PI(4,5)P$_2$-bound structure, we also determined the structure of the complex prepared

in the presence of EGTA (Fig. S1c, Table S1, and Methods). In this case, no bound Ca^{2+} is observed,

yet the channel remains in the same open conformation, confirming that NC1 and PI(4,5)P$_2$ are

sufficient to activate the channel independently of Ca^{2+} (Fig. S5a and S5b).

Together, these structural and functional results reveal several key insights into PI(4,5)P$_2$-

dependent NC1 activation: NC1 induces conformational changes in the S1-S4 domain similar to

those triggered by Ca^{2+} without interfering with Ca^{2+} binding; Like Ca^{2+}, NC1 alone stabilizes the

channel in a desensitized state, requiring PI(4,5)P$_2$ to promote channel opening; In the presence of

PI(4,5)P$_2$, both NC1 and Ca^{2+} can independently activate hTRPM4.

Molecular determinants of NC1 specificity for human TRPM4

NC1 activates human but not mouse TRPM4 [39], despite the high sequence and structural homology between the two orthologs. This species-specific activation is particularly intriguing given the identical amino acid composition and conserved structure of the NC1-binding pocket. To identify the molecular determinants underlying this specificity, we used the cell death assay to screen various mutations in both human and mouse TRPM4 expressed in HeLa cells. Through a combination of chimeric constructs and point mutations (Fig. S6, Fig. S7, and associated legends), we identified three residues that potentially contribute to the selective activation of hTRPM4 by NC1: V904 on the S4 helix (corresponding to L900 in mTRPM4), P997 in the pore domain (R993 in mTRPM4), and R1064 on the TRP helix (S1060 in mTRPM4) (Fig. 6a).

We next tested whether mouse TRPM4 could be converted into an NC1-sensitive channel by introducing single, double, and triple mutations in which the three residues (L900, R993, and S1060) were replaced with their human TRPM4 counterparts. Figure 6b and 6c show the electrophysiological recordings of NC1 activation in these mutants. Strikingly, the single S1060R mutation in mTRPM4 was sufficient to confer NC1 sensitivity; however, the resulting current was outwardly rectifying, with greater conductance at depolarized membrane potentials. In contrast, mTRPM4 mutants bearing L900V and/or R993P substitutions remained insensitive to NC1 when tested individually or in combination. Notably, when these mutations were combined with S1060R, NC1 sensitivity was significantly enhanced. The mTRPM4 triple mutant (L900V/R993P/S1060R) displayed an NC1 activation profile nearly identical to that of hTRPM4. Complementing these gain-of-function experiments, we also tested loss-of-function mutations in hTRPM4 (Fig. S8a and S8b). Consistent with the mTRPM4 mutagenesis findings, mutations at V904 and P997 did not affect NC1 activation of hTRPM4. In contrast, both the single R1064S mutation and the V904L/P997R/R1064S triple mutation significantly reduced NC1 activation, though not completely abolishing it.

In addition, we generated the HeLa stable cell lines expressing those mTRPM4 gain-of-function mutants and measured their viability in response to NC1 treatment (Fig. 6d). Consistent with our electrophysiology result, expressing mTRPM4 with L900V and R993P mutations has little effect on cell viability similar to control cells expressing WT mTRPM4 or empty vector; however, the expression of S1060R single mutant or S1060R-containing double or triple mutants

progressively decreases the cell viability in response to NC1; cells expressing the

L900V/R993P/S1060R triple mutant of mTRPM4 exhibit similar sensitivity to NC1-induced

necrosis as hTRPM4-expressing cells.

The structural basis of NC1 specificity

To gain structural insight into the NC1 specificity of TRPM4, we determined the cryo-EM

structures of both the NC1-insensitive WT mTRPM4 and its NC1-activated

L900V/R993P/S1060R triple mutant, each in the presence of NC1 (Fig. S9, Table S2, and

Methods). As $PI(4,5)P_2$ is required for NC1 activation in hTRPM4, it was also included during

mTRPM4 protein preparation (Methods).

The structure of WT mTRPM4 in the presence of NC1 and $PI(4,5)P_2$ was resolved at 2.9

Å resolution (Fig. 7a). Surprisingly, NC1 still binds to mTRPM4 at the same pocket as in hTRPM4

with identical surrounding residues (Fig. 7b), despite failing to activate the channel. Relative to

the apo mTRPM4 structure, NC1 binding induces similar local conformational changes within the

S1-S4 domain, as observed in hTRPM4 (Fig. S10a). However, unlike NC1/$PI(4,5)P_2$-activated

hTRPM4, where large swinging and rotational movements of the cytosolic domains are coupled

to pore opening, the cytosolic domains of NC1/$PI(4,5)P_2$-bound mTRPM4 closely resemble the

apo closed conformation, with only a subtle rigid-body shift (Fig. 7a). Notably, three distinct

conformational changes in the transmembrane region of NC1/$PI(4,5)P_2$-bound mTRPM4

distinguish it from the NC1-sensitive hTRPM4 and likely underlie its lack of activation (Fig. 7c,

7d and 7e). Firstly, the entire S1-S4 domain moves laterally toward the center of the channel (Fig.

7c). Secondly, the central filter and surrounding pore helices exhibit no clear density, suggesting

the filter becomes disordered upon NC1 binding (Fig. 7d) - a surprising finding, as this tightly

packed central filter region remains steady during channel gating and is typically the best-resolved

in TRPM4 structures. Thirdly, the S5 helix shifts inward toward the filter, while the pore-lining

S6 helix moves outward, widening the intracellular gate (Fig. 7c and 7e).

The latter two structural changes appear to originate from the lateral movement of S1-S4.

Due to tight S5-mediated packing between the S1-S4 domain and the neighboring subunit's pore

domain, this lateral shift of S1-S4 drives S5 inward, compressing the adjacent pore helix and filter

toward the central axis (Fig. 7c). Unable to accommodate this compression, the filter likely
undergoes a promiscuous movement and adopts a disordered, non-conductive conformation,
leading to the collapse of the ion conduction pathway. The accompanying outward motion of S6
likely serves to relieve the spatial strain caused by the rearrangement of the central pore.
Importantly, these conformational changes are solely induced by NC1 binding, as an NC1-bound
mTRPM4 structure prepared without PI(4,5)P$_2$ is identical to the NC1/PI(4,5)P$_2$-bound form (Fig.
S9b, Fig. S10b, Table S2, and Methods). Thus, the insensitivity of mTRPM4 to NC1 activation is
not due to a lack of drug binding, but rather to NC1-induced collapse of the selectivity filter,
rendering the channel non-conductive, analogous to C-type inactivation observed in certain K$^+$
channels [44].

We next determined the structure of the NC1-sensitive mTRPM4 triple mutant
(L900V/R993P/S1060R) in complex with NC1 and PI(4,5)P$_2$ (Fig. 7f, Fig. S9c, Table S2, and
Methods). Unlike WT mTRPM4, the mutant's pore domain is well structured and adopts an open
conformation (Fig. 7e), and its S1-S4 domain no longer undergoes a lateral movement. The overall
structure of the mutant closely resembles that of the NC1/PI(4,5)P$_2$-bound open hTRPM4 (Fig.
S10c), faithfully recapitulating the activation mechanism observed in the human channel. Given
that the S1060R mutation in the TRP helix is the key determinant for conferring NC1 sensitivity,
we focused our structural comparison on this region between NC1/PI(4,5)P$_2$-bound WT and
mutant mTRPM4. In the mutant, R1060 forms a network of inter-domain interactions absent in
the WT channel (Fig. 7g). Specifically, R1060 mediates a cluster of salt bridges and hydrogen
bonds involving D858 and W860 at the N-terminus of S3, as well as the glycerol phosphate head
group of PI(4,5)P$_2$. These interactions serve to stabilize the S1-S4 domain, preventing the inward
lateral movement observed in NC1-bound WT mTRPM4 that disrupts the pore. Notably, this
arginine residue, crucial for NC1 sensitivity, is conserved in human and primate TRPM4 orthologs
but is replaced by a serine in most other mammalian species.

Discussion

The small molecule NC1 activates hTRPM4, triggering Na$^+$ influx and leading to necrotic
cell death via sodium overload. Through structural and functional analyses, we elucidated the
molecular mechanism underlying NC1-mediated activation of hTRPM4. NC1 binds near the Ca^{2+}

activation site within the S1-S4 domain and induces local conformational changes that mirror those
caused by Ca^{2+}. Like Ca^{2+}, NC1 requires membrane PI(4,5)P₂ to stabilize the open channel
conformation. Thus, NC1 acts as a non-competitive surrogate of the native ligand. NC1 engages
in extensive hydrophobic and polar interactions with surrounding residues, resulting in high-
affinity binding. Its binding site, deeply buried within the protein, likely contributes to its slow
activation kinetics and poor washout. While distinct from the binding sites of other known TRPM4
modulators, the NC1 site is similar to some modulator-binding pockets in TRPM3 (e.g., primidone
and CIM 0216) [45,46] and TRPM8 (e.g., N-(3-aminopropyl)-2-[(3-methylphenyl)methoxy]-N-
[(thiophen-2-yl)methyl]benzamide (AMTB), cryosim-3 (C3), icilin, and the menthol analog WS-
12) [47-49].

NC1 selectively activates human TRPM4 but not most other mammalian TRPM4 orthologs.
Through comparative studies of human and mouse TRPM4, we determined that the species
specificity of NC1 activation is not due to a failure in drug binding. Instead, in mTRPM4, NC1
binding destabilizes the selectivity filter, rendering the channel non-conductive, akin to C-type
inactivation observed in certain K^+ channels. We identified R1064, located on the TRP helix of
hTRPM4, as a key determinant of NC1 specificity. This arginine mediates a network of
interdomain interactions that stabilize the S1-S4 domain's lateral position within the membrane.
In TRPM4s from mice and most other mammals, a serine occupies this position. The absence of
arginine-mediated stabilization permits a lateral shift of the S1-S4 domain upon NC1 binding,
which collapses the selectivity filter and inactivates the channel. Interestingly, akin to NC1, the
TRPM4 modulator CBA was shown to inhibit the human but not the mouse channel [33] and does
not exert any effect on other TRPM channel subfamily members [31]. However, the mechanistic
details that confer TRPM4 species-specificity to this compound have not been thoroughly explored.

The mechanistic insights into NC1 activation of hTRPM4 and its remarkable specificity
revealed in this study establish a structural framework for understanding how small molecules can
selectively modulate TRPM4 function. Our structural analyses identify a previously unrecognized
drug-binding pocket that is distinct from those of other known TRPM4 modulators, thereby
expanding the current understanding of ligand recognition and channel activation in the TRPM
family. The defined NC1-binding mode provides an essential molecular basis for rational drug

design aimed at exploiting TRPM4-mediated necrosis as a potential therapeutic avenue for cancer
treatment.

The discovery that NC1 binds to an identical site in both human and mouse TRPM4
orthologs but induces markedly different functional outcomes emphasizes the structural subtleties
governing allosteric coupling and species-dependent channel gating. Given the widespread
reliance on murine and other mammalian models in preclinical studies, our findings underscore
the necessity of validating compound efficacy and mechanism in human-derived systems to ensure
accurate interpretation of pharmacological responses. This cross-species discrepancy also serves
as a cautionary example for drug development pipelines targeting ion channels, where even small
structural variations can result in divergent functional consequences. Collectively, our results
provide a comprehensive molecular foundation for future high-throughput screening, structure-
guided optimization, and in vivo evaluation of TRPM4 modulators.

Methods

Protein expression and purification

Full-length wild-type human TRPM4 (UniProtKB: Q8TD43) containing an N-terminal FLAG-tag (DYKDDDDK) and full-length wild-type or L900V;R993P;S1060R mouse TRPM4 (UniprotKB: Q7TN37) containing a C-terminal FLAG-tag were cloned into a pEZT-BM plasmid [50]. *E.coli* DH10Bac cells were used to synthesize the bacmid that was employed in baculovirus production in Sf9 cells (Thermo Fisher Scientific) using Cellfectin II reagent (Thermo Fisher Scientific). For protein expression, HEK293S GnTI⁻ cells (American Type Culture Collection (ATCC), CRL-3022) grown in suspension to a density of 3×10^6 cells/mL were infected with P3 viruses at a ratio of 1:40 (virus:cell, v/v) and supplemented with 10 mM sodium butyrate to boost protein expression. The cells were then incubated at 37 °C for 48 h before being harvested by centrifugation (5000 g, 15 min, 4 °C). The cell pellet was resuspended in lysis buffer (50 mM Tris-HCl pH 7.5, 150 mM NaCl) supplemented with protease inhibitors (0.5 μg/ml pepstatin, 2 μg/ml leupeptin, 1 μg/ml aprotinin, and 1 mM PMSF) and homogenized by sonication. TRPM4 was then extracted with 1 % (w/v) lauryl maltose neopentyl glycol (LMNG, Anatrace) in gentle agitation for 2 h at 4 °C. The supernatant was subsequently collected by centrifugation (40000 g, 40 min, 4 °C) and incubated with Anti-DYKDDDDK G1 Affinity Resin (GenScript) for 1 h and 30 min at 4 °C in gentle agitation. The resin was then washed with Buffer A (50 mM Tris-HCl pH 7.5, 150 mM NaCl, 0.01 % LMNG) and elution was performed by incubating for 45 min at room temperature in gentle agitation with Buffer B (50 mM Tris-HCl pH 7.5, 150 mM NaCl, 0.01 % LMNG, 0.2 mg/mL FLAG peptide). The eluate was then concentrated and further purified by size-exclusion chromatography in Buffer C (50 mM Tris-HCl pH 7.5, 150 mM NaCl, 0.0035 % LMNG) on a Superose 6 Increase 10/300 GL (GE Healthcare). For the Ca^{2+} free samples, 2 mM EGTA was also included in Buffer C and for the samples requiring PI(4,5)P$_2$, 50 μM Brain PI(4,5)P$_2$ (L-α-phosphatidylinositol-4,5-bisphosphate (Brain, Porcine) (ammonium salt), Avanti) was added to the buffer B during the elution step and further used to supplement the concentrated eluate before size-exclusion chromatography.

Cryo-EM sample preparation and data acquisition

hTRPM4 and mTRPM4 samples were concentrated to ~7-8 mg/mL and supplemented with 0.3

mM   Necrocide   1   ((S')-3-cycloheptyl-3-(4-hydroxyphenyl)-6-methoxy-7-methyl-1,3-

dihydroindole-2-one) (NC1). 1 mM EGTA was also added when required. Supplemented samples

were incubated for 1 h on ice before vitrification. 4 µL was then applied to a glow-discharged

Quantifoil R1.2/1.3 300-mesh gold holey carbon grid (Quantifoil, Micro Tools GmbH, Germany),

blotted for 3.5 s under 100 % humidity at 12 °C and plunged into liquid ethane using a Mark IV

Vitrobot (FEI).

All data collections were performed using SerialEM with a nominal defocus range set from -0.9

to -2.2 µm and each movie was recorded with a total dose of 60 e$^-$/Å^2.

For the dataset of hTRPM4 containing NC1 and EGTA, movies were acquired on a Titan Krios

microscope (FEI) operated at 300 kV with a Falcon4i direct electron detector camera (Thermo

Fisher Scientific) at 0.738 Å pixel size.

For the dataset of hTRPM4 containing NC1 and PI(4,5)P$_2$, movies were acquired on a Titan Krios

microscope (FEI) operated at 300 kV with a K3 direct electron detector camera (Gatan) using the

CDS (Correlated Double Sampling) mode with a super-resolution pixel size of 0.4172 Å.

For the datasets of hTRPM4 containing NC1, PI(4,5)P$_2$ and EGTA and mTRPM4 wild-type

containing NC1 and PI(4,5)P$_2$, movies were acquired on a Titan Krios microscope (FEI) operated

at 300 kV with a K3 direct electron detector camera (Gatan) using the CDS (Correlated Double

Sampling) mode with a super-resolution pixel size of 0.4285 Å.

For the datasets of mTRPM4 wild-type containing NC1 and mTRPM4 L900V/R993P/S1060R

triple mutant containing NC1 and PI(4,5)P$_2$, movies were acquired on a Titan Krios microscope

(FEI) operated at 300 kV with a Falcon4i direct electron detector camera (Thermo Fisher Scientific)

at 0.735 Å pixel size.

Cryo-EM data processing

The workflow for data processing is summarized in Fig. S1 and Fig. S9. Data processing was

performed using cryoSPARC (versions 4.3.1 or 4.4.1) [51] following the general scheme described

below with some modifications between the datasets. For the dataset of hTRPM4 containing NC1

and PI(4,5)P$_2$, RELION (version 3.1) [52,53] was also employed in the processing workflow.

For the dataset of hTRPM4 containing NC1 and PI(4,5)P$_2$, movies were motion-corrected and

dose-weighted using MotionCor2 [54] and CTF estimation was performed using GCTF [55].

Micrographs were manually inspected to remove those with bad defocus values and ice

contamination. Particles were then picked using Laplacian-of-Gaussian (LoG) and extracted in

RELION. Extracted particles were subjected to multiple rounds of 2D classification and classes

displaying clear features of the TRPM4 channel were used for subsequent 3D classification. The

particles from the best-resolving 3D classes along with the respective 3D volumes were then

imported into cryoSPARC and used for 3D heterogeneous refinement and 3D classification

without alignment to remove junk particles and differentiate channel conformations. The best-

resolving 3D classes were refined using Non-Uniform refinement with imposed C4 symmetry [56].

For all other datasets, movies were subjected to patch motion correction and subsequent patch CTF

estimation in cryoSPARC. The resulting micrographs were manually curated and those exhibiting

drift, poor CTF estimation and outliers in defocus values and ice thickness were excluded from

further processing (individually assessed for each parameter relative to the overall distribution).

An initial round of particle picking was carried out with blob picker. Particles were then extracted

and subjected to one round of 2D classification. Classes displaying clear features of the TRPM4

channel were selected and used either directly for subsequent processing or to re-pick particles

with template picker. Additional rounds of 2D classification were further performed and particles

from selected classes were used to obtain an initial 3D reconstruction with Ab-Initio. 3D

heterogeneous refinement was then employed to remove junk particles, and the resulting particles

were subjected to 3D classification without alignment to differentiate channel conformations. The

best-resolving 3D classes were refined using Non-Uniform refinement with imposed C4 symmetry

[56].

Map resolutions were reported according to the gold-standard Fourier shell correlation (FSC) using

the 0.143 criterion [57]. Local resolutions and angular distributions were estimated in cryoSPARC.

Model building

Initial models were obtained using a combination of ModelAngelo [58] and previously available

hTRPM4 (PDBs: 9MTA and 9MRT) and mTRPM4 (PDB:6BCJ) structures. The models were then

manually adjusted in Coot [59] and refined against the respective maps in PHENIX [60]. Ligand

restraints CIF files were generated using the Grade2 Web Server [61]. The geometry statistics of the

models were obtained using MolProbity [62]. All the structural figures were prepared using UCSF

ChimeraX [63,64].

Electrophysiology

Wild-type human and mouse TRPM4, along with the respective mutants, were cloned into a

pEGFP-N1 plasmid (Clontech). Mutants were generated by site-directed mutagenesis using the

QuickChange method and verified by sequencing. 1 µg of plasmid was transfected into HEK293

cells using Lipofectamine 2000 (Life Technology). 48 h after transfection, cells were dissociated

by trypsin treatment and kept in a complete serum-containing medium before being re-plated onto

35-mm tissue culture dishes and incubated in a tissue culture incubator until recording.

Channel currents were recorded in whole cell or excised inside-out configuration. The long-chain

native Brain $PI(4,5)P_2$ employed in the structural studies is insoluble in water and forms liposomes

in the recording solutions, making it difficult to fuse into the patch membrane. Therefore, the

water-soluble short-chain synthetic $PI(4,5)P_2$ diC8 (Phosphatidylinositol 4,5-bisphosphate diC8,

Echelon Bioscience) was used in the functional experiments as needed. For inside-out patches, the

standard bath solution (cytosolic side) contained (in mM): 145 caesium methanesulfonate (Cs–

MS), 5 NaCl, 1 $MgCl_2$, 0.3 $CaCl_2$, 10 HEPES buffered with Tris, pH 7.4. 10 µM NC1 and/or 10

µM $PI(4,5)P_2$ diC8 were added to the bath solution when required. For the calcium-free condition,

0.5 mM EGTA was added to the bath solution without $CaCl_2$. The pipette solution (extracellular

side) contained (in mM): 140 Na–MS, 1 $MgCl_2$, 5 $CaCl_2$, 10 HEPES buffered with Tris, pH 7.4.

For whole cell current recordings, the pipette solution contained (in mM): Cs-MS 140, CsCl 4,

$MgCl_2$ 1, EGTA 1 or 10, HEPES 10, buffered with Tris, pH 7.4. Bath solution contained (in mM):

Na-MS 140, NaCl 5, $MgCl_2$ 1, $CaCl_2$ 1, HEPES 10, buffered with Tris, pH=7.4. 10 µM NC1 was

added to the bath solution as needed. Patch pipettes were pulled from borosilicate glass (Harvard

Apparatus) and heat polished to a resistance of 3–5 MΩ. After the patch pipette was attached to

the cell membrane, a giga seal (> 10 GΩ) was formed by gentle suction. The inside-out

configuration was formed by pulling the pipette away from the cell and the pipette tip was exposed

to air for a short time in some cases. The holding potential was set to 0 mV. The current and voltage

relationship (I–V curve) was obtained using voltage pulses ramped from – 100 to + 100 mV over

an 800 ms duration every 2 seconds. Data was acquired using an AxoPatch 200B amplifier

(Molecular Devices) and a low-pass analog filter set to 1 kHz. The current signal was sampled at

a rate of 20 kHz using a Digidata 1322A digitizer (Molecular Devices) and further analyzed with

pClamp 11 software (Molecular Devices). Sample traces for the I–V curves of macroscopic

currents shown were obtained from recordings on the same patch. Bars represent mean ± SEM of

five measurements from different patches (n = 5 independent biological replicates).

Cell death assay

Cell death was assessed in HeLa cells transiently expressing EGFP-tagged human or mouse

TRPM4 by monitoring the morphological change of transfected cells using microscopy, as

described previously [39]. Briefly, HeLa cells were seeded onto glass coverslips and grown to 30–

40% confluency before transfection. 24 hours after TRPM4-EGFP transfection, cells were

subjected to 1 µM NC1 treatment. Fluorescence images were taken before and after the indicated

period of treatment. The morphological changes of cells with green fluorescence were monitored

using microscopy, and over 50 cells expressing TRPM4 were analyzed in each group.

Cell viability assay

Cell viability was assessed in HeLa cells stably expressing human or mouse TRPM4, using the

CellTiter-Glo (CTG) assay as described previously [39]. Briefly, cells were seeded in a 96-well plate

at a density of 1×10^4 cells per well and allowed to adhere overnight. After treatment with various

experimental conditions, the intracellular ATP was measured using the CTG luminescent assay kit

(Promega, G7570) according to the manufacturer's instructions. Briefly, 100 µL of CTG reagent

was added to each well, and the plate was gently shaken for 10 minutes to induce cell lysis and

release of intracellular ATP. The luminescence signal, proportional to the amount of ATP present

and indicative of cell viability, was measured using a multi-well plate reader. All experiments were

repeated three times, and the results were presented as mean ± SD.

Data availability

The cryo-EM density maps of the human TRPM4 have been deposited in the Electron Microscopy

Data Bank (EMDB) under accession numbers EMD-73526 (NC1-bound in the presence of EGTA),

EMD-73527 (NC1/PI(4,5)P₂-bound), and EMD-73528 (NC1/PI(4,5)P₂-bound in the presence of

EGTA). Atomic coordinates have been deposited in the Protein Data Bank (PDB) under accession

numbers 9YVK (NC1-bound in the presence of EGTA), 9YVL (NC1/PI(4,5)P$_2$-bound), and

9YVM (NC1/PI(4,5)P$_2$-bound in the presence of EGTA).

The cryo-EM density maps of the mouse TRPM4 have been deposited in the Electron Microscopy

Data Bank (EMDB) under accession numbers EMD-73530 (NC1/PI(4,5)P$_2$-bound wild-type

mTRPM4), EMD-73529 (NC1-bound wild-type mTRPM4), and EMD-73531 (NC1/PI(4,5)P$_2$-

bound L900V/R993P/S1060R triple mutant mTRPM4). Atomic coordinates have been deposited

in the Protein Data Bank (PDB) under accession numbers 9YVO (NC1/PI(4,5)P$_2$-bound wild-type

mTRPM4), 9YVN (NC1-bound wild-type mTRPM4), and 9YVP (NC1/PI(4,5)P$_2$-bound

L900V/R993P/S1060R triple mutant mTRPM4).

Acknowledgements

Single particle cryo-EM data were collected at the University of Texas Southwestern Medical

Center Cryo-EM Facility (CEMF) funded by the CPRIT Core Facility Support Award RP170644

and at Howard Hughes Medical Institute Janelia Cryo-EM Facility. Cryo-EM sample grids were

prepared at the Structural Biology Laboratory at UT Southwestern Medical Center partially

supported by grant RP170644 from CPRIT. This work was supported in part by the Howard

Hughes Medical Institute (to Y.J.) and by grant from the National Institute of Health

(R35GM140892 to Y.J.), and by grants from MOST (Ministry of Science and Technology of the

People's Republic of China) (2023YFA0914900 to Q.Z.), NSFC (National Natural Science

Foundation of China) (92254307 to Q.Z.).

Competing interests

490   The authors declare no competing interests.

References

Launay, P. *et al.* TRPM4 is a Ca2+-activated nonselective cation channel mediating cell membrane depolarization. *Cell* **109**, 397-407 (2002). https://doi.org/10.1016/s0092-8674(02)00719-5

Montell, C. *et al.* A Unified Nomenclature for the Superfamily of TRP Cation Channels. *Molecular Cell* **9**, 229-231 (2002). https://doi.org/https://doi.org/10.1016/S1097-2765(02)00448-3

Nilius, B. *et al.* Voltage dependence of the Ca2+-activated cation channel TRPM4. *J Biol Chem* **278**, 30813-30820 (2003). https://doi.org/10.1074/jbc.M305127200

Fonfria, E. *et al.* Tissue Distribution Profiles of the Human TRPM Cation Channel Family. *Journal of Receptors and Signal Transduction* **26**, 159-178 (2006). https://doi.org/10.1080/10799890600637506

Demion, M., Bois, P., Launay, P. & Guinamard, R. TRPM4, a Ca2+-activated nonselective cation channel in mouse sino-atrial node cells. *Cardiovasc Res* **73**, 531-538 (2007). https://doi.org/10.1016/j.cardiores.2006.11.023

Guinamard, R., Demion, M., Chatelier, A. & Bois, P. Calcium-Activated Nonselective Cation Channels in Mammalian Cardiomyocytes. *Trends in Cardiovascular Medicine* **16**, 245-250 (2006). https://doi.org/https://doi.org/10.1016/j.tcm.2006.04.007

Hof, T. *et al.* TRPM4 non-selective cation channel variants in long QT syndrome. *BMC Med Genet* **18**, 31 (2017). https://doi.org/10.1186/s12881-017-0397-4

Stallmeyer, B. *et al.* Mutational spectrum in the Ca2+-activated cation channel gene TRPM4 in patients with cardiac conductance disturbances. *Human Mutation* **33**, 109-117 (2012). https://doi.org/https://doi.org/10.1002/humu.21599

Launay, P. *et al.* TRPM4 regulates calcium oscillations after T cell activation. *Science* **306**, 1374-1377 (2004). https://doi.org/10.1126/science.1098845

Becerra, A. *et al.* Transient receptor potential melastatin 4 inhibition prevents lipopolysaccharide-induced endothelial cell death. *Cardiovascular Research* **91**, 677-684 (2011). https://doi.org/10.1093/cvr/cvr135

Gerzanich, V. *et al.* De novo expression of Trpm4 initiates secondary hemorrhage in spinal cord injury. *Nature Medicine* **15**, 185-191 (2009). https://doi.org/10.1038/nm.1899

Ghosh, S. *et al.* Plasma Membrane Channel TRPM4 Mediates Immunogenic Therapy-Induced Necrosis. *Cancer Res* **83**, 3115-3130 (2023). https://doi.org/10.1158/0008-5472.Can-23-0157

Schattling, B. *et al.* TRPM4 cation channel mediates axonal and neuronal degeneration in experimental autoimmune encephalomyelitis and multiple sclerosis. *Nature Medicine* **18**, 1805-1811 (2012). https://doi.org/10.1038/nm.3015

Simon, F. *et al.* Hydrogen peroxide removes TRPM4 current desensitization conferring increased vulnerability to necrotic cell death. *J Biol Chem* **285**, 37150-37158 (2010). https://doi.org/10.1074/jbc.M110.155390

Borgström, A., Peinelt, C. & Stokłosa, P. TRPM4 in Cancer-A New Potential Drug Target. *Biomolecules* **11** (2021). https://doi.org/10.3390/biom11020229

Berg, K. D. *et al.* TRPM4 protein expression in prostate cancer: a novel tissue biomarker associated with risk of biochemical recurrence following radical prostatectomy. *Virchows Archiv* **468**, 345-355 (2016). https://doi.org/10.1007/s00428-015-1880-y

Holzmann, C. *et al.* Transient receptor potential melastatin 4 channel contributes to migration of androgen-insensitive prostate cancer cells. *Oncotarget* **6** (2015).

Sagredo, A. I. *et al.* TRPM4 channel is involved in regulating epithelial to mesenchymal transition, migration, and invasion of prostate cancer cell lines. *Journal of Cellular Physiology* **234**, 2037-2050 (2019). https://doi.org/https://doi.org/10.1002/jcp.27371

19 Guo, J. *et al.* TRPM4 channels contribute to Adriamycin chemoresistance in breast cancer cells.
European Journal of Pharmacology **998**, 177637 (2025).
https://doi.org/https://doi.org/10.1016/j.ejphar.2025.177637
20 Wong, K. K. & Hussain, F. A. TRPM4 is overexpressed in breast cancer associated with estrogen
response and epithelial-mesenchymal transition gene sets. *PLOS ONE* **15**, e0233884 (2020).
https://doi.org/10.1371/journal.pone.0233884
21 Loo, S. K. *et al.* TRPM4 expression is associated with activated B cell subtype and poor survival in
diffuse large B cell lymphoma. *Histopathology* **71**, 98-111 (2017).
https://doi.org/10.1111/his.13204
22 Wang, F. *et al.* Aberrant TRPM4 expression in MLL-rearranged acute myeloid leukemia and its
blockade induces cell cycle arrest via AKT/GLI1/Cyclin D1 pathway. *Cellular Signalling* **72**, 109643
(2020). https://doi.org/https://doi.org/10.1016/j.cellsig.2020.109643
23 Amarouch, M.-Y., Syam, N. & Abriel, H. Biochemical, single-channel, whole-cell patch clamp, and
pharmacological analyses of endogenous TRPM4 channels in HEK293 cells. *Neuroscience Letters*
541, 105-110 (2013). https://doi.org/https://doi.org/10.1016/j.neulet.2013.02.011
24 Burris, S. K., Wang, Q., Bulley, S., Neeb, Z. P. & Jaggar, J. H. 9-Phenanthrol inhibits recombinant
and arterial myocyte TMEM16A channels. *Br J Pharmacol* **172**, 2459-2468 (2015).
https://doi.org/10.1111/bph.13077
25 Gardam, K. E., Geiger, J. E., Hickey, C. M., Hung, A. Y. & Magoski, N. S. Flufenamic acid affects
multiple currents and causes intracellular Ca2+ release in Aplysia bag cell neurons. *J
Neurophysiol* **100**, 38-49 (2008). https://doi.org/10.1152/jn.90265.2008
26 Guinamard, R. *et al.* Functional characterization of a Ca2+-activated non-selective cation channel
in human atrial cardiomyocytes. *The Journal of Physiology* **558**, 75-83 (2004).
https://doi.org/https://doi.org/10.1113/jphysiol.2004.063974
27 Chen, B. *et al.* TRPM4-specific blocking antibody attenuates reperfusion injury in a rat model of
stroke. *Pflugers Arch* **471**, 1455-1466 (2019). https://doi.org/10.1007/s00424-019-02326-8
28 Low, S. W. *et al.* Development and characterization of a monoclonal antibody blocking human
TRPM4 channel. *Sci Rep* **11**, 10411 (2021). https://doi.org/10.1038/s41598-021-89935-5
29 Wei, S. *et al.* Binding epitope for recognition of human TRPM4 channel by monoclonal antibody
M4M. *Scientific Reports* **12**, 19562 (2022). https://doi.org/10.1038/s41598-022-22077-4
30 Wei, S. *et al.* Development and in vitro characterization of humanized antibodies for blocking
human TRPM4 channel. *Sci Rep* **15**, 19769 (2025). https://doi.org/10.1038/s41598-025-05256-x
31 Ozhathil, L. C. *et al.* Identification of potent and selective small molecule inhibitors of the cation
channel TRPM4. *Br J Pharmacol* **175**, 2504-2519 (2018). https://doi.org/10.1111/bph.14220
32 Binkle-Ladisch, L. *et al.* Identification and development of TRPM4 antagonists to counteract
neuronal excitotoxicity. *iScience* **27**, 111425 (2024). https://doi.org/10.1016/j.isci.2024.111425
33 Arullampalam, P. *et al.* Species-Specific Effects of Cation Channel TRPM4 Small-Molecule
Inhibitors. *Front Pharmacol* **12**, 712354 (2021). https://doi.org/10.3389/fphar.2021.712354
34 Ekundayo, B. *et al.* Identification of a binding site for small molecule inhibitors targeting human
TRPM4. *Nat Commun* **16**, 833 (2025). https://doi.org/10.1038/s41467-025-56131-2
35 Teixeira-Duarte, C. M., Zeng, W. & Jiang, Y. Structural landscape of activation, desensitization
and inhibition in the human TRPM4 channel. *Nature Structural & Molecular Biology* (2025).
https://doi.org/10.1038/s41594-025-01705-3
36 Christensen, M. K. *et al.* Synthesis and antitumor effect in vitro and in vivo of substituted 1,3-
dihydroindole-2-ones. *J Med Chem* **53**, 7140-7145 (2010). https://doi.org/10.1021/jm100763j
37 Trojel-Hansen, C. *et al.* Novel small molecule drugs inhibit tumor cell metabolism and show
potent anti-tumorigenic potential. *Cancer Chemother Pharmacol* **68**, 127-138 (2011).
https://doi.org/10.1007/s00280-010-1453-3

38 Zhang, J. *et al.* Necrocide 1 mediates necrotic cell death and immunogenic response in human
cancer cells. *Cell Death Dis* **14**, 238 (2023). https://doi.org/10.1038/s41419-023-05740-0
39 Fu, W. *et al.* Persistent activation of TRPM4 triggers necrotic cell death characterized by sodium
overload. *Nature Chemical Biology* (2025). https://doi.org/10.1038/s41589-025-01841-3
40 Fulda, S. & Debatin, K. M. Extrinsic versus intrinsic apoptosis pathways in anticancer
chemotherapy. *Oncogene* **25**, 4798-4811 (2006). https://doi.org/10.1038/sj.onc.1209608
41 Singh, R., Letai, A. & Sarosiek, K. Regulation of apoptosis in health and disease: the balancing act
of BCL-2 family proteins. *Nature Reviews Molecular Cell Biology* **20**, 175-193 (2019).
https://doi.org/10.1038/s41580-018-0089-8
42 Nilius, B. *et al.* The Ca2+-activated cation channel TRPM4 is regulated by phosphatidylinositol
4,5-biphosphate. *Embo j* **25**, 467-478 (2006). https://doi.org/10.1038/sj.emboj.7600963
43 Zhang, Z., Okawa, H., Wang, Y. & Liman, E. R. Phosphatidylinositol 4,5-bisphosphate rescues
TRPM4 channels from desensitization. *J Biol Chem* **280**, 39185-39192 (2005).
https://doi.org/10.1074/jbc.M506965200
44 Ong, S. T., Tyagi, A., Chandy, K. G. & Bhushan, S. Mechanisms Underlying C-type Inactivation in
Kv Channels: Lessons From Structures of Human Kv1.3 and Fly Shaker-IR Channels. *Frontiers in*
Pharmacology **Volume 13 - 2022** (2022). https://doi.org/10.3389/fphar.2022.924289
45 Yin, Y. *et al.* Molecular basis of neurosteroid and anticonvulsant regulation of TRPM3. *Nature*
Structural & Molecular Biology **32**, 828-840 (2025). https://doi.org/10.1038/s41594-024-01463-
8
46 Kumar, S. *et al.* Structural basis for agonist and heat activation of nociceptor TRPM3. *Nature*
Structural & Molecular Biology (2025). https://doi.org/10.1038/s41594-025-01692-5
47 Yin, Y. *et al.* Structural basis of cooling agent and lipid sensing by the cold-activated TRPM8
channel. *Science* **363**, eaav9334 (2019). https://doi.org/doi:10.1126/science.aav9334
48 Yin, Y. *et al.* Mechanisms of sensory adaptation and inhibition of the cold and menthol receptor
TRPM8. *Science Advances* **10**, eadp2211 (2024). https://doi.org/doi:10.1126/sciadv.adp2211
49 Zhao, C. *et al.* Structures of a mammalian TRPM8 in closed state. *Nature Communications* **13**,
3113 (2022). https://doi.org/10.1038/s41467-022-30919-y
50 Morales-Perez, C. L., Noviello, C. M. & Hibbs, R. E. Manipulation of Subunit Stoichiometry in
Heteromeric Membrane Proteins. *Structure* **24**, 797-805 (2016).
https://doi.org/10.1016/j.str.2016.03.004
51 Punjani, A., Rubinstein, J. L., Fleet, D. J. & Brubaker, M. A. cryoSPARC: algorithms for rapid
unsupervised cryo-EM structure determination. *Nature Methods* **14**, 290-296 (2017).
https://doi.org/10.1038/nmeth.4169
52 Scheres, S. H. W. RELION: Implementation of a Bayesian approach to cryo-EM structure
determination. *Journal of Structural Biology* **180**, 519-530 (2012).
https://doi.org/https://doi.org/10.1016/j.jsb.2012.09.006
53 Zivanov, J. *et al.* New tools for automated high-resolution cryo-EM structure determination in
RELION-3. *Elife* **7** (2018). https://doi.org/10.7554/eLife.42166
54 Zheng, S. Q. *et al.* MotionCor2: anisotropic correction of beam-induced motion for improved
cryo-electron microscopy. *Nature Methods* **14**, 331-332 (2017).
https://doi.org/10.1038/nmeth.4193
55 Zhang, K. Gctf: Real-time CTF determination and correction. *Journal of Structural Biology* **193**, 1-
12 (2016). https://doi.org/https://doi.org/10.1016/j.jsb.2015.11.003
56 Punjani, A., Zhang, H. & Fleet, D. J. Non-uniform refinement: adaptive regularization improves
single-particle cryo-EM reconstruction. *Nature Methods* **17**, 1214-1221 (2020).
https://doi.org/10.1038/s41592-020-00990-8

57   Henderson, R. *et al.* Outcome of the first electron microscopy validation task force meeting.
Structure **20**, 205-214 (2012). https://doi.org/10.1016/j.str.2011.12.014
58   Jamali, K. *et al.* Automated model building and protein identification in cryo-EM maps. *Nature*
628, 450-457 (2024). https://doi.org/10.1038/s41586-024-07215-4
59   Emsley, P., Lohkamp, B., Scott, W. G. & Cowtan, K. Features and development of Coot. *Acta*
Crystallographica Section D **66**, 486-501 (2010).
https://doi.org/doi:10.1107/S0907444910007493
60   Liebschner, D. *et al.* Macromolecular structure determination using X-rays, neutrons and
electrons: recent developments in Phenix. *Acta Crystallographica Section D* **75**, 861-877 (2019).
https://doi.org/doi:10.1107/S2059798319011471
61   Smart, O. S., Sharff A., Holstein, J., Womack, T.O., Flensburg, C., Keller, P., Paciorek, W.,
Vonrhein, C. and Bricogne G. Grade2 version 1.6.0. *Cambridge, United Kingdom:Global Phasing*
Ltd. (2021).
62   Chen, V. B. *et al.* MolProbity: all-atom structure validation for macromolecular crystallography.
Acta Crystallogr D Biol Crystallogr **66**, 12-21 (2010).
https://doi.org/10.1107/s0907444909042073
63   Pettersen, E. F. *et al.* UCSF ChimeraX: Structure visualization for researchers, educators, and
developers. *Protein Sci* **30**, 70-82 (2021). https://doi.org/10.1002/pro.3943
64   Meng, E. C. *et al.* UCSF ChimeraX: Tools for structure building and analysis. *Protein Science* **32**,
e4792 (2023). https://doi.org/https://doi.org/10.1002/pro.4792

Fig. 1: NC1 activation of hTRPM4. a Macroscopic currents of hTRPM4 overexpressed in HEK293 cells at ±100 mV in a whole-cell patch with the presence or absence of 10 μM NC1 in the bath (extracellular). Triangles mark the time points of three NC1-mediated hTRPM4 activation states (left to right): Closed channel in the absence of NC1; Fully activated channel 2 min after NC1 addition; Channel current retained 5 min after NC1 washout. **b** Sample I-V curves measured at the time points marked by the triangles shown in **a**. **c** NC1-induced cell death assay. Top panel: schematic illustration of the assay. Bottom panels: morphologies of EGFP, EGFP-tagged mTRPM4, or EGFP-tagged hTRPM4-expressing HeLa cells 24 hours after DMSO (NC1 solvent) or NC1 (1 μM) treatment. **d** Dose-dependent NC1 cytotoxicity in HeLa cells stably expressing wild-type hTRPM4. Data points represent mean ± SD of n = 3 independent replicates.

Fig. 2: NC1 binding in hTRPM4. a Structure of NC1-bound hTRPM4. The front subunit is depicted in a cartoon representation, with each domain individually colored. **b** Zoomed-in view of the NC1 binding site. NC1 and its surrounding residues are shown in stick representation. The light green sphere marks the calcium activation site. NC1 density (grey surface) is contoured at 0.2 in ChimeraX. Residues sensitive to tryptophan substitution are labeled in red. **c** Sample I-V curves of wild-type hTRPM4 and its tryptophan substitutions at the NC1-binding site recorded in whole-cell patches with 10 µM NC1 in the bath (extracellular). **d** Outward current density of NC1-activated wild-type hTRPM4 and its tryptophan mutations recorded at +100 mV in whole-cell patches with 10 µM NC1 in the bath (extracellular). Bars represent mean ± SEM of n=5 independent replicates. **e** Dose-dependent NC1 cytotoxicity in HeLa cells stably expressing wild-type hTRPM4 or its tryptophan mutations. Data points represent mean ± SD of n = 3 independent replicates.

Fig. 3: hTRPM4 conformational changes induced by NC1 binding. a Superposition of the hTRPM4 structures in the NC1-bound and apo closed (PDB:9MTA) states with the front subunits highlighted in cyan and orange, respectively. **b** Local conformational changes at the S1-S4 domain between NC1-bound (cyan) and apo closed (orange) hTRPM4. Only S3-S4 and the neighboring S5 (marked with a single quotation mark) are displayed for clarity. Red arrows mark the major movements upon NC1 binding. Key residues are shown in stick representation. **c** Structural comparison of the hTRPM4 ion conduction pore in the apo closed (orange), NC1-bound (cyan), and NC1/PI(4,5)P$_2$-bound (navy blue) states. Gating residues I1040 and S1044 are shown in stick representation. The front and back subunits are removed for clarity. **d** Pore radius of hTRPM4 along the central axis in the apo closed, NC1-bound, and NC1/PI(4,5)P$_2$-bound structures. **e** F935'-F910 contact interface in the NC1-bound, apo closed, Ca^{2+}-bound desensitized (PDB:9MT8), and Ca^{2+}/PI(4,5)P$_2$-bound open (PDB:9MRT) states. The EM density maps for F910 and F935' are shown in grey surface contoured at 0.23 in ChimeraX.

Fig. 4 PI(4,5)P$_2$-dependent NC1 activation of hTRPM4. a Macroscopic currents of hTRPM4-expressing HEK293 cells at ±100 mV in an inside-out patch with the presence or absence of Ca^{2+}, NC1, and PI(4,5)P$_2$ diC8 in the bath (cytosolic). NC1 and PI(4,5)P$_2$ diC8 were introduced after Ca^{2+} activation and desensitization due to Ca^{2+}-induced membrane PI(4,5)P$_2$ depletion. A-E mark the time points of hTRPM4 in various states: A, apo closed state; B, initial Ca^{2+}-activated state before desensitization; C, Ca^{2+}-bound desensitized state due to PI(4,5)P$_2$ depletion; D, closed state in the presence of NC1; E, NC1/PI(4,5)P$_2$-activated state. The panel on the right depicts sample I-V curves corresponding to A-E time points. **b** Similar recordings as **a** except that PI(4,5)P$_2$ diC8 was introduced before NC1, and the time point D marks the closed state in the presence of PI(4,5)P$_2$ diC8. The panel on the right depicts sample I-V curves corresponding to A-E time points. **c** Sample I-V curves of wild-type hTRPM4 and its alanine substitutions at the PI(4,5)P$_2$-binding site recorded in whole-cell patches with 10 µM NC1 in the bath (extracellular). **d** Outward and inward current densities of NC1-activated wild-type hTRPM4 and its mutants recorded at ±100 mV in whole-cell patches with 10 µM NC1 in the bath (extracellular). Bars represent mean ± SEM of n=5 independent replicates. **e** Responsiveness of HeLa cells to NC1-induced cell death when expressing wild-type hTRPM4 or its PI(4,5)P$_2$ site mutants. Data points represent mean ± SD of n = 3 independent replicates. Over 50 cells were counted in each replicate.

Fig. 5: Structure of NC1/PI(4,5)P$_2$-bound hTRPM4. a Superposition of the NC1/PI(4,5)P$_2$-bound and Ca^{2+}/PI(4,5)P$_2$-bound (PDB:9MRT) hTRPM4 open structures with the front subunits highlighted in navy blue and pink, respectively. NC1 and PI(4,5)P$_2$ are depicted in stick representation, and Ca^{2+} is shown as a green sphere. **b** Zoomed-in view of the ligand binding sites in the NC1/PI(4,5)P$_2$-bound structure. Ca^{2+} is represented as a green sphere. NC1, PI(4,5)P$_2$, and ligand-interacting residues are shown in stick representation. Density (grey surface) is contoured at 0.5 for NC1 and Ca^{2+} and at 0.45 for PI(4,5)P$_2$ in ChimeraX. **c** Superposition of the NC1/PI(4,5)P$_2$-bound open and NC1-bound hTRPM4 structures with the front subunits highlighted in navy blue and cyan, respectively. **d** Enlarged view of the conformational changes at the transmembrane region between the NC1/PI(4,5)P$_2$-bound open (navy blue) and NC1-bound (cyan) structures. The S4 helix from the neighboring subunit (S4') is also shown. The inset shows a close-up highlighting the coupling between F935 and neighboring F910' in the NC1/PI(4,5)P$_2$-bound structure that leads to the movement (red arrow) at the S4-S5 linker and S5 and channel opening. Density for F935 and F910' is shown in a grey surface and contoured at 0.55 in ChimeraX. **e** Conformational changes at the cytosolic MHR domains between the NC1/PI(4,5)P$_2$-bound (navy blue) and NC1-bound (cyan) structures. Red arrows mark the upward swing of the cytosolic MHR domains and the rotation of MHRs 1 and 2.

a

Pre-S1, S1, S2, S3, S4, V904, S4-S5 linker, Pore domain, P997, S5, S6, R1064, TRP helix 1, TRP helix 2

S1-S4, P997, Pore helix, TRP helix 2, V904, S5, R1064, S6, TRP helix 1

b

mTRPM4 WT, mTRPM4 L900V, mTRPM4 R993P, mTRPM4 L900V/R993P, mTRPM4 S1060R, mTRPM4 L900V/S1060R, mTRPM4 R993P/S1060R, mTRPM4 L900V/R993P/S1060R, hTRPM4 WT

Current (pA), Vm (mV), 0 µM NC1, 10 µM NC1

c

Current density (pA/pF)

mTRPM4, mTRPM4

hTRPM4_WT, mTRPM4_WT, L900V, R993P, S1060R, R993P/S1060R, L900V/S1060R, L900V/R993P/S1060R

At 100 mV, At -100 mV

d

HeLa NC1 24h

Cell viability (%), log[NC1] (M)

Vector
hTRPM4 WT
mTRPM4 WT
mTRPM4 L900V/R993P/S1060R
mTRPM4 L900V/R993P
mTRPM4 L900V/S1060R
mTRPM4 R993P/S1060R
mTRPM4 S1060R

Fig. 6: Molecular determinants of NC1 specificity for human TRPM4. a Topology (left) and structure (right) of the hTRPM4 TM domain with the three key residues important for NC1 specificity marked. **b** Sample I-V curves of wild-type mTRPM4 and its single, double, and triple mutants at the three key residues. Currents were recorded in whole-cell patches with 10 µM NC1 in the bath (extracellular). Current from NC1-activated wild-type hTRPM4 is shown in the last panel for comparison. **c** Outward and inward current densities of the wild-type mTRPM4 and its mutants recorded at ±100 mV in whole-cell patches with 10 µM NC1 in the bath (extracellular). Bars represent mean ± SEM of n=5 independent replicates. Current densities of hTRPM4 are also graphed for comparison. **d** Dose-dependent NC1 cytotoxicity in stable HeLa cell lines expressing wild-type hTRPM4, wild-type mTRPM4, or mTRPM4 mutants. Data points represent mean ± SD of n = 3 independent replicates.

Fig. 7: NC1 binding in mTRPM4. a Superposition of the NC1/PI(4,5)P$_2$-bound (brown) and apo closed (PDB:6BCJ) (green) wild-type mTRPM4 structures along the pore axis, with the front subunits highlighted. NC1 (blue) and PI(4,5)P$_2$ (yellow) are shown in stick representation. **b** Zoomed-in view of the NC1 binding site in mTRPM4. NC1 and its interacting residues are shown in stick representation. NC1 density (grey surface) is contoured at 0.23 in ChimeraX. **c** Conformational changes at S1-S4 and its neighboring pore domain upon NC1 binding to mTRPM4. The red line and red diamond mark the central 4-fold axis. Red arrows indicate the major movements in this region, namely the lateral movement of S1-S4 towards the central axis, the inward movement of the neighboring subunit S5 (S5'), and the outward movement of the neighboring subunit S6 (S6'). **d** Comparison of the density maps at the pore domain between the apo closed (PDB:6bcj) (green) and NC1/PI(4,5)P$_2$-bound (brown) mTRPM4 structures. The disordered region in the pore domain of the NC1/PI(4,5)P$_2$-bound structure is depicted as a grey semi-transparent cartoon. **e** Structural comparison of the ion conduction pore in the apo closed (green), NC1/PI(4,5)P$_2$-bound inactivated (brown) wild-type mTRPM4 and in the NC1/PI(4,5)P$_2$-bound open (purple) triple mutant (L900V/R993P/S1060R). Gating residues I1036 and S1040 are shown in stick representation. The front and back subunits were removed for clarity. The panel on the right depicts the pore radius along the central axis in these structures. The dotted line illustrates the disordered filter region in the NC1/PI(4,5)P$_2$-bound wild-type mTRPM4. **f** Structure of the NC1/PI(4,5)P$_2$-bound mTRPM4 triple mutant (L900V/R993P/S1060R) with the front subunit highlighted in purple. NC1 and PI(4,5)P$_2$ are shown in stick representation and Ca^{2+} as a green sphere. **g** Structural comparison at S1060 region between NC1/PI(4,5)P$_2$-bound wild-type mTRPM4 (light brown) and its triple mutant (L900V/R993P/S1060R, light purple). PI(4,5)P$_2$ and key residues important for stabilizing S1-S4 in the mutant are shown in stick representation. The wild-type S1060 side chain is also shown for comparison.